Assessing genomic diversity and signatures of selection in Original Braunvieh cattle using whole-genome sequencing data

Meenu Bhati[1, *], Naveen Kumar Kadri[1], Danang Crysnanto[1], Hubert Pausch[1]

[1]Animal Genomics, ETH Zürich, Zürich, Switzerland

Authors and email addresses:

    Meenu Bhati: meenu.bhati@usys.ethz.ch

    Naveen Kumar Kadri: naveen.kadri@usys.ethz.ch

    Danang Crysnanto: danang.crysnanto@usys.ethz.ch

    Hubert Pausch: hubert.pausch@usys.ethz.ch

    [*]Corresponding author

Abstract

Background: Autochthonous cattle breeds represent an important source of genetic variation because they might carry alleles that enable them to adapt to local environment and food conditions. Original Braunvieh (OB) is a local cattle breed of Switzerland used for beef and milk production in alpine areas. Using whole-genome sequencing (WGS) data of 49 key ancestors, we characterize genomic diversity, genomic inbreeding, and signatures of selection in Swiss OB cattle at nucleotide resolution.

Results: We annotated 15,722,811 million SNPs and 1,580,878 million Indels including 10,738 and 2,763 missense deleterious and high impact variants, respectively, that were discovered in 49 OB key ancestors. Six Mendelian trait-associated variants that were previously detected in breeds other than OB, segregated in the sequenced key ancestors including variants causal for recessive xanthinuria and albinism. The average nucleotide diversity (1.6×10^{-3}) was higher in OB than many mainstream European cattle breeds. Accordingly, the average genomic inbreeding quantified using runs of homozygosity (ROH) was relatively low ($F_{ROH}=0.14$) in the 49 OB key ancestor animals. However, genomic inbreeding was higher in more recent generations of OB cattle ($F_{ROH}=0.16$) due to a higher number of long (> 1 Mb) runs of homozygosity. Using two complementary approaches, composite likelihood ratio test and integrated haplotype score, we identified 95 and 162 genomic regions encompassing 136 and 157 protein-coding genes, respectively, that showed evidence ($P < 0.005$) of past and ongoing selection. These selection signals were enriched for quantitative trait loci related to beef traits including meat quality, feed efficiency and body weight and pathways related to blood coagulation, nervous and sensory stimulus.

Conclusions: We provide a comprehensive overview of sequence variation in Swiss OB cattle genomes. With WGS data, we observe higher genomic diversity and less inbreeding in OB than many European mainstream cattle breeds. Footprints of selection were detected in genomic regions that are possibly relevant for meat quality and adaptation to local

1 environmental conditions. Considering that the population size is low and genomic

2 inbreeding increased in the past generations, the implementation and adoption of optimal

3 mating strategies seems warranted to maintain genetic diversity in the Swiss OB cattle

4 population.

5
6

Introduction

Following the domestication of cattle, both natural and artificial selection led to the formation

of breeds with distinct phenotypic characteristics including morphological, physiological and

adaptability traits [1]. With an increasing demand for animal-based food products, few

breeds were intensively selected for high milk (e.g., Holstein, Brown Swiss) and beef (e.g.,

Angus) production. The predominant selection of cattle from specialized breeds caused a

sharp decline in the population size of local cattle breeds [2, 3]. Although less productive

under intensive production conditions, local breeds of cattle might carry alleles that enable

them to adapt to local conditions. Therefore, local breeds represent an important genetic

resource to facilitate animal breeding in the future under challenging and changing

production conditions [4, 5]. Characterizing the genetic diversity of local cattle breeds is

important to optimally manage these genetic resources.

The Swiss Original Braunvieh (OB) cattle breed is a dual purpose taurine cattle breed that is

used for beef and milk production in alpine areas [6, 7]. In transhumance farming, the cattle

graze at alpine pastures (between 1000 and 2400 meter above sea level) during the

summer months and return to the stables for the winter months [7]. Mainly due to their

strong and firm legs and claws, they are well adapted to the alpine terrain. Under extensive

farming conditions, OB cattle may outperform specialized dairy breeds in terms of fertility,

longevity and health status [8]. However, in the early 1960s, Swiss cattle breeders began

inseminating OB cows with semen from US Brown Swiss sires to increase milk yield, reduce

calving difficulties and improve mammary gland morphology of the Swiss OB cattle

population [9]. The extensive cross-breeding of OB cows with Brown Swiss sires decreased

the number of female OB calves entering the herdbook to less than 2,000 by mid 1990's [9]

(Additional file 1). Since then, the size of the Swiss OB population increased steadily,

facilitated by governmental subsidies.

A number of studies investigated the genomic diversity and population structure of the Swiss

OB cattle breed using either pedigree or microarray data [9, 10]. In spite of the small

population size, genetic diversity is higher in OB than many commercial breeds likely due to

the use of many sires in natural mating and lower proportion of artificial insemination [9, 10].

Genomic inbreeding and footprints of selection have been compared between OB and other

Swiss cattle breeds using SNP microarray-derived genotypes [10]. Because the SNP

microarrays were designed in a way that they interrogate genetic markers that are common

in the mainstream breeds of cattle, they might be less informative for breeds of cattle that

are diverged from the mainstream breeds [11]. Ascertainment bias is inherent in the

resulting genotype data because rare, breed-specific, and less-accessible genetic variants

are underrepresented among the microarray-derived genotypes [12]. This limitation causes

observed allele frequency distributions to deviate from expectations which can distort

population genetics estimates [13].

With the availability of whole genome sequencing (WGS), it has become possible to

discover sequence variant genotypes at population scale [14]. While sequence variant

genotypes might be biased toward the reference allele, this reference bias is less of a

concern when the sequencing coverage is high [15]. According to Boitard et al. 2016 [16],

WGS data facilitate detecting selection signatures at higher resolution than SNP microarray

data. Moreover, the WGS-based detection of runs of homozygosity (ROH) is more sensitive

for short ROH that are typically missed using SNP microarray-derived genotypes.

In the present study, we analyze more than 17 million WGS variants of 49 key ancestors of

the Swiss OB cattle breed that were sequenced to an average fold-coverage of 12.75 per

animal [17]. These data enabled us to assess genomic diversity and detect signatures of

past or ongoing selection in the breed at nucleotide resolution. Moreover, we estimate

genomic inbreeding in the population using runs of homozygosity.

2 Material and Methods

Sequence variant genotyping

We considered genotypes at 17,303,689 biallelic variants (15,722,811 SNPs and 1,580,878

Indels) that were discovered and genotyped previously [17] in the autosomes of 49 key

ancestors of the OB population using a genome graph-based sequence variant genotyping

approach [18]. In brief, 49 OB cattle were sequenced at between 6 and 38-fold genome

coverage using either Illumina HiSeq 2500 (30 animals) or Illumina HiSeq 4000 (19 animals)

instruments. The sequencing reads were filtered and subsequently aligned to the UMD3.1

assembly of the bovine genome [19] using the mem-algorithm of the Burrows Wheeler

Aligner (BWA) software package [20]. Single nucleotide and short insertion and deletion

polymorphisms were discovered and genotyped using the Graphtyper software [18].

Following recommended filtration criteria (see Crysnanto et al. [17] for more details),

15,722,811 SNPs and 1,580,878 Indels were retained for subsequent analyses. Beagle [21]

phasing and imputation was applied to improve the primary genotype calls from Graphtyper

and infer missing genotypes. Unless stated otherwise, imputed genotypes were considered

for subsequent analyses, because Beagle imputation considerably improved the primary

genotype calls particularly in samples that had been sequenced at low coverage [17].

Variant annotation and evaluation

Functional consequences of 15,722,811 SNPs and 1,580,878 Indels were predicted

according to the Ensembl (release 91) annotation of the bovine genome assembly UMD3.1

using the Variant Effect Predictor tool (VEP v.91.3) [22] with default parameter settings. The

impacts of amino acid substitutions on protein function were predicted using the sorting

intolerant from tolerant (SIFT) (version 5.2.2) [23] algorithm that has been implemented in

the VEP tool. Variants with SIFT scores less than 0.05 were considered to be likely

deleterious to protein function. In order to assess if known Mendelian trait-associated

variants segregate among 49 sequenced OB cattle, we downloaded genomic coordinates of

155 trait-associated variants that are curated in the Online Mendelian Inheritance in Animals

(OMIA) database [24, 25].

Population genetic analysis

Nucleotide diversity (π) quantifies the average number of nucleotide differences per site

between two DNA sequences that originated from the studied population [26]. We estimated

π of the OB population over the entire autosomal genome using VCFtools v0.1.15 (in

windows of 10 kb) [27].

Detection of runs of homozygosity

Runs of homozygosity were identified using a Hidden Markov Model (HMM)-based approach

implemented in the BCFtools/RoH software [28, 29]. The recombination rate was assumed

to be constant along the genome at 10^{-8} per base pair (1 cM/Mb). For the HMM-based

detection of ROH, we considered phred-scaled likelihoods (PL) and allele frequencies of

15,722,811 filtered SNPs before Beagle imputation. Because samples that are sequenced at

low coverage are enriched for ROH [30], we considered only 33 samples with average

sequencing coverage greater than 10-fold for the detection of ROH (Additional file 2). We

only considered ROH longer than 50 kb because they were less likely to contain false-

positives (Phred-score > 67 in our data, Additional file 2). Genomic inbreeding (F_{ROH}) was

calculated for each animal as $F_{ROH}=\sum L_{ROH}/L_{GENOME}$, where $\sum L_{ROH}$ is the length of all ROH

longer than 50 kb and L_{GENOME} is the length of the genome covered by SNPs [31], which is

2,512,054,768 bp in our data.

Further, ROH were classified into short (50 - 100 kb), medium (0.1 - 2 Mb) and long ROH

(>2 Mb) reflecting ancient, historical, and recent inbreeding, respectively [32]. The

contribution of each ROH category to F_{ROH} was calculated for each animal. Average

genomic inbreeding was compared between animals born before and after 1989 using the

two samples t-test.

27
Detection of signatures of selection

To avoid potential bias arising from extended relationships among the sequenced animals,

we did not consider nine sons from sire-son pairs for the detection of signatures of selection.

For the remaining 40 cattle, we considered genotypes at 9,051,833 SNPs for which the

ancestral allele provided by Rocha et al. [33] was detected in at least two species other than

cattle and where it agreed with either the reference or alternate allele in our data.

Haplotypes were phased using the Eagle2 software [34] with default parameter settings and

assuming a constant recombination rate along the chromosome.

Integrated haplotype score (iHS)
9

To identify signatures of ongoing selection, integrated haplotype scores (iHS) were

calculated for 8,465,912 variants with minor allele frequency (MAF) greater than 0.01 using

the R package rehh v.2 [35]. We obtained iHS that ranged from -6.6 to 6.4. Subsequently,

|iHS| were averaged for non-overlapping windows of 40 kb over the whole genome.

Windows with either less than 10 SNPs were removed. To test if variants with similar |iHS|

properties were pooled in 40 kb windows, we followed the approach of Granka et al. [36].

Specifically, we randomly selected the same number of SNPs that were pooled in 40 kb

windows and calculated the mean variance of |iHS| in the true and permuted 40 kb windows

for each chromosome. This procedure was repeated for 10,000 randomly selected 40 kb

windows. The variance of |iHS| in the non-overlapping 40 kb windows (0.24) was

significantly ($P < 0.01$) less than in windows of randomly selected SNPs (0.37) indicating that

SNPs that were grouped in 40 kb windows had |iHS| values that were more alike than

random SNPs.

Composite likelihood ratio (CLR)

Composite likelihood ratio (CLR) tests were carried out to identify alleles that are either close

to fixation or already reached fixation due to past selection. Following the recommendation

of Huber et al. [37], we removed 118,124 SNPs from the data which were fixed for the

ancestral alleles because such sites are not informative for CLR tests. Using a pre-

computed empirical allele frequency spectrum of 8,933,709 SNPs for which ancestral and

derived alleles were assigned (see above), we calculated CLR statistics in non-overlapping

40 kb windows using SweepFinder2 [38, 39]. A window size of 40 kb was chosen to allow

comparison and alignment between |iHS| and CLR values.

Empirical *P* values were calculated for CLR and |iHS| windows [40] and the top 0.5% of

windows of each statistic were considered as candidate signatures of selection. Adjacent top

0.5% windows were merged separately for each statistic using BEDTools v2.27.1 [41]. For

each merged candidate signature of selection, the lowest *P* value among the merged

windows was retained.

Characterization of signatures of selection

Genes within candidate signatures of selection were determined based on the Ensembl

(release 91) annotation of the UMD3.1 assembly of the bovine genome. Gene-set

enrichment analysis of genes within candidate signatures of selection was performed using

PANTHER v.14.1 [42]. Specifically, we investigated if these genes were enriched in the

functional categories of GO-slim Biological Process and PANTHER pathways using $P \leq 0.05$

as significance level.

To determine the overlap between QTL and candidate signatures of selection, we

downloaded genomic coordinates for 122,893 QTL from the Animal QTL Database [43, 44].

We classified 85,722 unique QTL that were located on the 29 autosomes into six trait

categories: exterior, health, milk, meat and carcass, production and reproduction (Additional

file 3). QTL with identical genomic coordinates in associated trait categories were

considered as one QTL. We used the intersect module of BEDTools v2.27.1 [41] to identify

QTL that overlapped with CLR and |iHS| candidate regions for each of the six trait categories

separately. To test if QTL were enriched in candidate signatures of selection, we used a

permutation test with 10,000 permutations. In each permutation, we randomly sampled the

same number of regions of the same size as the candidate signatures of selection from CLR

and |iHS| for each chromosome separately, and overlapped them with QTL of the respective

trait categories using BEDTools (see above). The number of QTL that overlapped permuted

regions was used as the empirical null distribution to calculate P values. P values less than

0.05 were considered as indicators for a significant enrichment of QTL in candidate

signatures of selection.

7 Results

8 Overview of genomic diversity in OB cattle

We annotated 15,722,811 biallelic SNPs and 1,580,878 Indels that were discovered in 49

OB cattle [17]. The average genome wide nucleotide diversity within the OB breed was

0.001637/bp. Among the detected variants, 546,419 (3.5%) SNPs and 307,847 (19.5%)

Indels were found novel when compared to the 102,090,847 polymorphic sites of the NCBI

bovine dbSNP database version 150.

Functional annotation of the polymorphic sites revealed that the vast majority of SNPs were

located in either intergenic (73.8%) or intronic regions (25.2%). Only 1% of SNPs (160,707)

were located in the exonic regions (Table 1). In protein-coding sequences, we detected

58,387, 47,249 and 1,264 synonymous, missense, and high impact SNPs, respectively.

According to the SIFT scoring, 10,738 missense SNPs were classified as likely deleterious

to protein function (SIFT score < 0.05). Among the high impact variants, we detected 580,

33, 106, 273 and 272 stop gain, stop lost, start lost, splice donor and splice acceptor

variants, respectively. Deleterious and high impact variants were more frequent in the low

than high allele frequency classes (Additional file 4).

The majority of 1,580,878 Indels were in either intergenic (72.7%) or intronic (26.7%)

regions. Only 2,213 (0.14%) Indels affected coding sequences. Among these, 1,499 were

classified as high impact variants including 1,324, 16, 4, 71 and 84 frameshift, stop gain,

start lost, splice donor and splice acceptor variants, respectively. Similar to previous studies

in cattle [14, 45], coding regions were enriched for Indels with lengths in multiples of three

indicating that they are less likely to be deleterious to protein function than frameshift

variants (Additional File 5).

3
4   **Table 1**: Number of SNPs and Indels in sequence ontology classes annotated using the

5   VEP software

Sequence ontology class	SNP	Indel
splice_acceptor_variant	272	84
splice_donor_variant	273	71
stop_gained	580	16
frameshift_variant	0	1324
stop_lost	33	0
start_lost	106	4
inframe_insertion	0	290
inframe_deletion	0	440
missense_variant	47429	0
protein_altering_variant	0	12
splice_region_variant	9553	1059
stop_retained_variant	45	2
synonymous_variant	58387	0
coding_sequence_variant	166	125
mature_miRNA_variant	83	23
5_prime_UTR_variant	6744	600
3_prime_UTR_variant	30716	4074
non_coding_transcript_exon_variant	6296	434
intron_variant	3960673	422764
non_coding_transcript_variant	24	25
upstream_gene_variant	526000	56715
downstream_gene_variant	454753	51672

intergenic_variant	10620678	1041144
Total	15722811	1580878

1
2
OMIA variants segregating in the OB population

We obtained genomic coordinates of 155 variants that are associated with Mendelian traits

in cattle from the OMIA database to analyze if they segregate among the 49 OB cattle. It

turned out that six OMIA variants were also detected in the 49 OB cattle including two

variants in the *MOCOS* and *SLC45A2* genes that are associated with severe recessive

disorders (Additional file 6). Two OB key ancestor bulls born in 1967 and 1974 (ENA SRA

sample accession numbers SAMEA4827662 and SAMEA4827664) were heterozygous

carriers of a single base pair deletion (BTA24:g.21222030delC) in the *MOCOS* gene (OMIA

001819-9913) that causes xanthinuria in the homozygous state in Tyrolean grey cattle [46].

Another two OB key ancestor bulls (sire and son; ENA SRA sample accession numbers

SAMEA4827659 and SAMEA4827645) that were born in 1967 and 1973 were heterozygous

carriers of two missense variants in *SLC45A2* (BTA20:g.39829806G>A and

BTA20:g.39864148C>T) that are associated with oculocutaneous albinism (OMIA 001821-

9913) in Braunvieh cattle [47].

Runs of homozygosity and genomic inbreeding

Runs of homozygosity were analyzed in 33 OB animals that had an average sequencing

depth greater than 10-fold. We found an average number of 2,044 ± 79 autosomal ROH per

individual with an average length of 179 kb ± 17.6 kb. The length of the ROH ranged from 50

kb (minimum size considered, see methods) to 5,025,959 bp. On average, 14.58% of the

genome (excluding sex chromosome) was in ROH (Additional file 7). Average genomic

inbreeding for the 29 chromosomes ranged from 11.5% (BTA29) to 18.6% (BTA26) (Figure

1a).

In order to study the demography of the OB population, we calculated the contributions of

short, medium and long ROH to the total genomic inbreeding (Additional file 7). The

medium-sized ROH were the most frequent class (50.46%), and contributed most (75.01%)

to the total genomic inbreeding. While short ROH occurred almost as frequent (49.17%) as

medium-sized ROH, they contributed only 19.52% to total genomic inbreeding (Figure 1b&c;

Additional file 7). Long ROH were rarely (0.36%) observed among the OB key ancestors and

contributed little (5.47%) to total genomic inbreeding. The number of long ROH was

correlated (r=0.77) with genomic inbreeding.

Figure 1 - ROH in 33 OB cattle with average sequencing depth greater than 10-fold. (a)

Average genomic inbreeding and corresponding standard error for the 29 autosomes. (b)

Average genomic inbreeding (F_{ROH}) calculated from short (50-100 kb), medium (0.1 - 2 Mb)

and long (>2 Mb) ROH. (c) Average number of short, medium and long ROH.

Genomic inbreeding (F_{ROH}) was significantly (P =0.0002) higher in 20 animals born between

1990 and 2012 than in 13 animals born between 1965 and 1989 (0.16 vs. 0.14) (Additional

file 8). The high F_{ROH} in animals born in more recent generations was mainly due to more

long (>2 Mb; P =0.00004) and medium-sized ROH (0.1-1 Mb; P =0.001) (Figure 2).

Figure 2 - Cumulative genomic inbreeding (%) in animals born between 1965-1989 (blue

lines) and 1990-2012 (red lines) from ROH sorted on length and binned in windows of 10 kb.

Thin dashed lines represent individuals and thick solid lines represent the average

cumulative genomic inbreeding of the two groups of animals.

Signatures of selection

We identified candidate signatures of selection using two complementary methods: the

composite likelihood ratio (CLR) test and the integrated haplotype score (iHS) (Figure 3a &

3b). The CLR test detects 'hard sweeps' where genomic regions encompass beneficial

adaptive alleles that recently reached fixation [48]. The iHS detects 'soft sweeps' at genomic

regions where selection for beneficial alleles is still ongoing [49, 50]. We detected 95 and

162 candidate regions of signatures of selection (*P* < 0.005) using CLR and iHS,

respectively, encompassing 12.56 Mb and 12.48 Mb (Additional file 9; Additional file 10).

These candidate signatures of selection were not evenly distributed over the genome

(Figure 3c). Functional annotation revealed that 136 and 157 protein-coding genes

overlapped with 50 and 86 candidate regions from CLR and iHS analyses, respectively. All

other candidate signatures of selection were located in intergenic regions. Closer inspection

of the top selection regions of both analyses revealed that 16 CLR candidate regions

overlapped with 25 iHS candidate regions on chromosomes 5, 7, 11, 14, 15, 17 and 26

(Figure 3c) encompassing 35 coding genes (Additional File 11).

12
13   **Figure 3 -** Genome wide distribution of top 0.5% signatures of selection from CLR (a) and

14   iHS (b) analyses and their overlap (c). Each point represents a non-overlapping window of

15   40 kb along the autosomes.

16
17   **Top candidate signatures of selection**
18

On chromosome 11, we identified 12 and 36 candidate regions of selection using CLR and

iHS analyses, respectively. The top CLR candidate region (P_{CLR} = 3.1 x 10^{-5}) was located on

chromosome 11 between 66 Mb and 68.5 Mb (Figure 4a) and it encompassed 24 protein-

coding genes (Additional file 9). The same region was also in ROH in 77% of 33 animals that

were sequenced at high coverage. The peak of this top CLR region was located between

67.5 and 68.2 Mb and it contained several adjacent windows with CLR values higher than

5,000 (P_{CLR} < 0.003). The top region encompassed 5 genes (Figure 4a & 4e). The variant

density in the top region was low and SNP allele frequency was skewed which is typical for

the presence of a hard sweep (Figure 4c). The top iHS candidate region was located on

chromosome 11 between 68.4 and 69.2 Mb (P_{iHS} = 3.2 x 10^{-5}) encompassing 7 genes

(Figure 4b & 4f). The allele frequencies of the SNPs within the top iHS region are

approaching fixation indicating ongoing selection possibly due to hitchhiking with the

neighboring hard sweep (Figure 4d).

Another striking CLR signal (P_{CLR} = 0.0012) was detected on chromosome 6 between 38.5

and 39.4 Mb. This genomic region encompasses the *DCAF16, FAM184B, LAP3, LCORL,*

MED28 and *NCAPG* genes, and the window with the highest CLR value overlapped the

NCAPG gene (Figure 5a & 5c). This signature of selection coincides with a QTL that is

associated with stature, feed efficiency and fetal growth [51–53]. Most SNPs detected within

this region were fixed for the alternate allele in the OB key ancestor animals of our study.

For instance, all 49 sequenced OB cattle were homozygous for the Chr6:38777311 G-allele

which results in a likely deleterious (SIFT score 0.01) amino acid substitution (p.I442M) in

the *NCAPG* gene that is associated with increased pre- and postnatal growth and calving

difficulties [51].

Figure 4 - Detailed view of a top candidate selection region on chromosome 11 in OB that was detected using CLR tests (a) and iHS (b). Each point represents a non-overlapping window of 40 kb. The dotted horizontal lines indicate the cutoff values (top 0.5%) for CLR (210) and iHS (2.13) statistics. The allele frequencies of the derived (red) or alternate alleles (black) (c and d) and genes (e and f) in the peak region (67.5-68.2 Mb) of the top CLR (66 to 68.5 Mb) and iHS (68.4-69.2 Mb) regions. Green and black colour indicates genes on the forward and reverse strand of DNA, respectively.

Figure 5 - Top CLR candidate region on chromosome 6 (a). Each point represents a non-

overlapping window of 40 kb. The frequencies of the derived (red) or alternate alleles (black)

(c) and genes (e) annotated between 38.5 and 39.4 Mb. Green and black colour indicates

genes on the forward and reverse strand of DNA, respectively.

GO enrichment analysis

Genes within candidate signatures of selection from CLR and iHS analyses were enriched

(after correcting for multiple testing) in the panther pathway (P00011) related to "Blood

coagulation". Genes within candidate signatures of selection from CLR tests were also

enriched in the pathway "P53 pathway feedback loops 1" (Additional file 12). Although we

did not find any enrichment of GO-slim biological processes after correcting for multiple

testing, 21 GO-slim biological processes including cellular catabolic processes, oxygen

transport and different splicing pathways were nominally enriched for genes within CLR

candidate signatures of selection and 14 GO-slim biological processes including nervous

system, sensory perception (olfactory receptors) and multicellular processes were nominally

enriched for genes within iHS candidate signatures of selection (Additional file 12).

QTL enrichment analysis

We investigated if candidate selection regions overlapped with trait-associated genomic

regions using QTL information curated at the Animal QTL Database (Animal QTLdb). We

found that 74.7% and 83.9% of CLR and iHS candidate signatures of selection, respectively,

were overlapping at least one QTL (Additional file 13). We tested for enrichment of these

signatures of selection in QTL for six trait classes: exterior, health, milk, meat, production,

and reproduction using permutation. It turned out that QTL associated with meat quality

(P_{CLR} = 0.0004, P_{iHS} = 0.0003) and production traits (P_{CLR} = 0.0027, P_{iHS} = 0.0039) were

significantly enriched in both CLR and iHS candidate signatures of selection. We did not

detect any enrichment of QTL associated with milk, reproduction, health, and exterior traits

neither in CLR nor in iHS candidate signatures of selection.

16
Discussion

We discovered 107,291 variants in coding sequences of 49 sequenced OB cattle. In

agreement with previous studies in cattle [14, 54], missense deleterious and high impact

variants occurred predominantly at low allele frequency likely indicating that variants which

disrupt physiological protein functions are removed from the population through purifying

selection [55]. However, deleterious variants may reach high frequency in livestock

populations due to the frequent use of individual carrier animals in artificial insemination [56],

hitchhiking with favorable alleles under artificial selection [57, 58], or demography effects

such as population bottlenecks [59]. Because we predicted functional consequences of

missense variants using computational inference, they have to be treated with caution in the

absence of experimental validation [60]. High impact variants that segregated among the 49

sequenced OB key ancestors were also listed as Mendelian trait-associated variants in the

OMIA database. For instance, we detected frameshift and missense variants in *MOCOS* and

SLC45A2 that are associated with recessive xanthinuria [46] and oculocutaneous albinism

[47], respectively. To the best of our knowledge, calves neither with xanthinuria nor

oculocutaneous albinism have been reported in the Swiss OB cattle population. The

absence of affected calves is likely due to the low frequencies of the deleterious alleles and

avoidance of matings between closely related heterozygous carriers. Among 49 sequenced

cattle, we detected only two bulls that carried the disease-associated *MOCOS* and *SLC45A2*

alleles in the heterozygous state. However, the frequent use of individual carrier bulls in

artificial insemination might result in an accumulation of diseased animals within short time

even when the frequency of the deleterious allele is low in the population [61]. Because the

deleterious alleles were detected in sequenced key ancestor animals that were born

decades ago, we cannot preclude that they were lost due to genetic drift or during the recent

population bottleneck in OB (Additional file1). A frameshift variant in *SLC2A2*

(NM_001103222:c.771_778delTTGAAAAGinsCATC, rs379675307, OMIA 000366-9913)

causes a recessive disorder in cattle that resembles human Fanconi-Bickel syndrome [62–

64]. Recently, the disease-causing allele was detected in the homozygous state in an OB

calf with retarded growth due to liver and kidney disease [65]. We did not detect the disease-

associated allele in our study. This may be because it is located on a rare haplotype that

does not segregate in the 49 sequenced cattle. Most of the sequenced animals of the

present study were selected for sequencing using the key ancestor approach, as their genes

contributed significantly to the current population [17, 66]. More sophisticated methods to

select animals for sequencing might prioritize rare haplotypes, thus increasing the likelihood

to detect rarer alleles when the sequencing budget is constrained [67–69].

24
Genomic diversity and genomic inbreeding

Original Braunvieh is a local cattle breed with approximately 10,000 cows registered in the

breeding population and 4,500 calves entering the herd book every year (Additional file 1). In

spite of the small population size, the nucleotide diversity ($\pi = 1.6 \times 10^{-3}$) is higher in OB

than many taurine cattle breeds with considerably more breeding animals including Holstein,

Jersey and Fleckvieh ($\sim$1.2-1.4 × 10^{-3}) [70, 71]. However, nucleotide diversity is lower in OB

than African indigenous cattle breeds (2.0-4.0 × 10^{-3}), New Danish Red (1.7 × 10^{-3}) and

Yakutian cattle breeds (1.7 × 10^{-3}) [71–73]. The average F_{ROH} estimated from WGS data was

0.14 in OB. This is lower than WGS-based F_{ROH} in Holstein (0.18), Jersey (0.24), Old Danish

Red (0.23) and Belgian Blue (0.3) cattle [74, 75]. However, the genomic inbreeding is slightly

higher in OB than New Red Danish cattle (0.11), an admixed breed that contains genes from

old Danish and other red breeds [74]. The relatively high genomic diversity of OB cattle is

assumed to be the result of many different sires contributing to the gene pool due to frequent

use of natural mating [10]. Our WGS based estimate of F_{ROH} (0.14) is substantially higher

than previous estimates obtained using 50K SNP microarray data (F_{ROH}= 0.029, [10]) for the

same population. Genotype data obtained using SNP microarrays with medium density (e.g.,

BovineSNP50) facilitate to detect long ROH (>1 Mb). However, due to low SNP density (~1

SNP per 50 kb) detecting short ROH is not possible using microarray-derived genotype data.

In our data, short and medium-sized ROH accounted for 80.48% of total inbreeding. Most

short and medium-sized ROH are not reliably detectable with the SNP microarrays that were

used to quantify F_{ROH} in Signer-Hasler et al. [10], resulting in an underestimation of genomic

inbreeding. Our estimate of the genomic inbreeding using WGS variants also includes short

and medium-sized ROH that were previously missed using SNP array data, thus

representing a realistic estimate of total genomic inbreeding in OB cattle.

Apart from genomic inbreeding, ROH also provide information about population and

individual demography based on length and number of ROH [76–78]. Our findings show that

medium-sized ROH that reflect historical inbreeding contribute most to the genomic

inbreeding of the current OB population. The minor contribution of long ROH to the genomic

inbreeding indicates that recent inbreeding is relatively low in OB possibly due to use of

many sires in natural matings as suggested by Hagger [9] and Signer-Hasler et al. [10]. Our

results based on ROH inferred from WGS variants corroborate that genomic inbreeding is

lower in OB than most mainstream breeds [10]. However, comparing the number and size

distribution of ROH across studies is subject to bias because misplaced genomic segments

might break ROH into multiple small- and medium-sized ROHs and different ROH-detection

approaches yield results that are not readily comparable [21, 27, 76, 79, 80]. Genomic

inbreeding is increasing in the OB population in recent years mainly due to an increase in

occurrence of long ROH. The recent population bottleneck in the OB population (Additional

file 1) might promote matings between closely related animals that caused inbreeding to

increase in recent generations. In this regard, genome-based mating strategies seem to be

warranted to achieve sufficient genetic gain while maintaining genetic diversity and avoiding

matings between carriers of disease-associated alleles [81, 82].

12   Signatures of selection

With WGS data, we were able to identify more signatures of selection compared to SNP

array data [10, 83], even though we used only 9 million SNP for which we could readily

assign ancestral and derived alleles [33]. Using two complementary approaches, we found

several new and known candidate regions that seem to be targets of recent or ongoing

selection in OB. Many signatures of selection were located in non-coding regions

corroborating that selection frequently acts on regulatory sites [16]. However, it is possible

that an improved annotation of the bovine genome might place these regions in yet to be

annotated coding regions. We applied methods to detect signatures of selection that depend

on frequency changes of alleles (CLR) and haplotypes (iHS). The detected signatures of

selection may be confounded by other evolutionary forces including genetic drift and

background selection [84–86].

We detected candidate selection regions in OB cattle that harbor genes associated with

stature or milk production (*NCAPG, LCORL, LAP3*), feed efficiency or lipid metabolism

(*R3HDM1, AOX1*), and unknown functions (*SLC25A33, TMEM201*) that were previously

reported to be targets of selection in taurine and indicine cattle breeds [16, 87–89]. The

presence of signatures of selection that are common in several breeds indicates that

selection at these regions has happened either before the breeds diverged or independently

after the formation of breeds [16, 89]. A number of genes that are targets of selection in

various cattle breeds are associated with either coat colour (*MC1R*, *KIT*), milk production

(*DGAT1*, *ABCG2*, *GHR*) or stature (*PLAG1*) [40, 71, 89, 90]. These genes were not detected

within the top 0.5% CLR and iHS windows in OB cattle possibly either due to absence of

trait-associated genomic variation in our data or these genes are not under selection in OB

cattle. While some cattle breeds including Holstein and Fleckvieh are selected for particular

coat colour patterns [40, 91], animals with variation in coat colour are rarely observed in the

OB cattle breed [92]. Moreover, due to the use of OB cattle for both milk and beef production

under extensive conditions, the milk production-associated variants that are under strong

artificial selection in many dairy breeds seem to be less important in OB cattle [88].

Some of the genes (*PLAG1*, *DGAT1*, *ABCG2*, *GHR*) that have been reported to be targets

of selection in specialized breeds contain well-known variants that contribute to the genetic

variation of economically important traits. We investigated if these variants segregate in our

data although they were not detected in our selection signature analysis. A number of

variants in high linkage disequilibrium stimulate the expression of *PLAG1*, thus increasing

pre- and postnatal growth in cattle [52, 53, 93]. Among 14 candidate causal variants for the

PLAG1 QTL, six were fixed for the stature-decreasing alleles in our study (Additional File

14). The other candidate causal variants were either fixed for the stature-increasing allele or

segregated at low allele frequency. This pattern indicates that a recombinant haplotype

might segregate in Swiss OB cattle that could facilitate fine-mapping of this region. Among

known mutations affecting milk production traits, a mutation in *ABCG2* (p.Y581S,

rs43702337 at 38,027,010 bp) [94] did not segregate in our population of Swiss OB cattle

which corroborates previous findings in Brown Swiss cattle [95]. A variant (BTA14,

g.1802265_1802266GC>AA, p.A232K) of the *DGAT1* gene is associated with milk

production traits in cattle [96, 97]. The milk fat-enhancing and milk yield-lowering lysine-

allele segregates in OB cattle at low frequency (0.03). A missense variant (BTA20,

g.31909478A>T, p.Y279F) in the *GHR* gene is associated with milk protein percentage [98].

The protein fat percentage-lowering T-allele segregates at low frequency (0.06) in OB cattle.

We observed a striking signature of selection on chromosome 11 that has previously been

detected in the Swiss Fleckvieh, Simmental, Eringer and Evolèner breeds using microarray-

called genotypes [10, 83]. Our results in OB cattle indicate that this region harbors a rapid

sweep which seems to act on alleles with selection advantage [49]. While large sweeps are

easy to detect using dense sequencing data, pinpointing causal alleles underpinning such

regions remains challenging. Most of the variants in such regions are either fixed or

segregate at very low frequency [16] which we also observed for the signature of selection

on chromosome 11. In our study, the signature of selection on chromosome 11

encompassed millions of nucleotides (between 66 and 72 Mb) and many genes, rendering

the identification of underpinning genes and variants a difficult task. The windows with the

highest CLR and |iHS| values did not encompass *PROKR1*, which was previously suggested

to be the target of selection at this region due to its association with fertility [10, 83].

However, closer inspection of the sequence variants detected in our study revealed that a

stop-gained variant in *PROKR1* (g.66998234C>A, rs476744845, p.Y293*) segregates at

high frequency in the 49 sequenced OB key ancestors. Yet, it remains to be elucidated, if

the presence of a high-impact variant in immediate proximity to a massive selective sweep is

causal or just due to hitchhiking. The window with the largest |iHS| value on chromosome 11

was right next to the *CAPN13* gene which is associated with meat tenderness and was also

suggested as a potential target of selection by Signer-Hasler et al. [10].

Genes within the top 0.5% CLR and iHS windows were enriched in pathways related to

reactive oxygen species, metabolic process, blood coagulation and nervous system,

indicating that the identified regions under selection might harbor genomic variants that

confer adaptive advantage to harsh environments. Moreover, QTL associated with meat

quality and production traits including feed efficiency and body weight were enriched in

selection signatures possibly indicating that OB cattle harbor variants that enabled them to

adapt to particular feed conditions. Combining results from selection signature and

association analyses might reveal phenotypic characteristics associated with genomic

regions that showed evidence of past or ongoing selection [40], thus providing additional

hints why particular genomic regions are under selection in OB cattle.

Conclusions

We provide a comprehensive overview of genomic variation segregating in the Swiss OB

cattle population using sequencing data of 49 key ancestor bulls. In spite of the small

population size, genetic diversity is higher and genomic inbreeding is lower in OB than many

other mainstream cattle breeds. However, genomic inbreeding is increasing in recent

generations mainly due to large ROH which should be considered in future management of

this breed. Finally, this study highlights regions that show evidence of past and ongoing

selection in OB which are enriched for QTL related to meat quality and production traits and

pathways related to blood coagulation, cellular metabolic process, and nervous system.

16
List of abbreviations

WGS- Whole-genome sequencing, CLR – Composite likelihood ratio; iHS – Integrated

haplotype score; OB – Original Braunvieh; OMIA – Online Mendelian Inheritance in Animals;

QTL – Quantitative trait loci; ROH – runs of homozygosity; SNP – Single nucleotide

polymorphism;

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Raw sequencing read data of all animals are available from the European Nucleotide Archive (ENA) (http://www.ebi.ac.uk/ena) under primary accession PRJEB28191 (SAMEA4827645-SAMEA4827674 and SAMEA5059741-SAMEA5059759).

Funding

This study was financially supported by the Federal Office for Agriculture (FOAG), Bern. The funding body was not involved in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Author's contributions

Analysis of whole-genome sequencing data: MB NKK CD HP; Conceived and designed the experiments: MB NKK HP; Wrote the paper: MB HP; Read and approved the final version of the manuscript: all authors

Acknowledgements

We thank Braunvieh Schweiz for providing pedigree and genotype data of Original Braunvieh cattle.

References

1. Food and Agriculture Organization. The Second Report on the State of the World's Animal Genetic Resources for Food and Agriculture. Rome: FAO Commission on Genetic Resources for Food and Agriculture Assessments. 2015. www.fao.org/publications.

2. Diversity ECG, Consortium. Marker-assisted conservation of European cattle breeds: an evaluation. Animal Genetics. 2006;37:475–81.

3. Medugorac I, Medugorac A, Russ I, Veit-Kensch CE, Taberlet P, Luntz B, et al. Genetic diversity of European cattle breeds highlights the conservation value of traditional unselected breeds with high effective population size. Molecular Ecology. 2009;18:3394–410.

4. Boettcher PJ, Hoffmann I, Baumung R, Drucker AG, McManus C, Berg P, et al. Genetic resources and genomics for adaptation of livestock to climate change. Frontiers in Genetics. 2014;5:461.

5. Biscarini F, Nicolazzi E, Alessandra S, Boettcher P, Gandini G. Challenges and opportunities in genetic improvement of local livestock breeds. Frontiers in Genetics. 2015;6:33.

6. Jurt C, Häberli I, Rossier R. Transhumance Farming in Swiss Mountains: Adaptation to a Changing Environment. Mountain Research and Development. 2015;35:57–65.

7. Herzog F, Seidl I. Swiss alpine summer farming: Current status and future development under climate change. Rangeland Journal. 2018;40:501–11.

8. Bieber A, Wallenbeck A, Leiber F, Fuerst-Waltl B, Winckler C, Gullstrand P, et al. Production level, fertility, health traits, and longevity in local and commercial dairy breeds under organic production conditions in Austria, Switzerland, Poland, and Sweden. Journal of Dairy Science. 2019;102:5330–41.

9. Hagger C. Estimates of genetic diversity in the brown cattle population of Switzerland obtained from pedigree information. Journal of Animal Breeding and Genetics. 2005;122:405–13.

10. Signer-Hasler H, Burren A, Neuditschko M, Frischknecht M, Garrick D, Stricker C, et al. Population structure and genomic inbreeding in nine Swiss dairy cattle populations. Genetics Selection Evolution. 2017;49:83.

11. Matukumalli LK, Lawley CT, Schnabel RD, Taylor JF, Allan MF, Heaton MP, et al. Development and Characterization of a High Density SNP Genotyping Assay for Cattle.

PLoS ONE. 2009;4:e5350.

12. Rowan TN, Hoff JL, Crum TE, Taylor JF, Schnabel RD, Decker JE. A Multi-Breed Reference Panel and Additional Rare Variation Maximizes Imputation Accuracy in Cattle. Preprint at the bioRxiv. 2019;517144. doi:https://doi.org/10.1101/517144. Accessed 27 November 2019.

13. Albrechtsen A, Nielsen FC, Nielsen R. Ascertainment biases in SNP chips affect measures of population divergence. Molecular Biology and Evolution. 2010;27:2534–47.

14. Daetwyler HD, Capitan A, Pausch H, Stothard P, van Binsbergen R, Brøndum RF, et al. Whole-genome sequencing of 234 bulls facilitates mapping of monogenic and complex traits in cattle. Nature Genetics. 2014;46:858–65.

15. Günther T, Nettelblad C. The presence and impact of reference bias on population genomic studies of prehistoric human populations. PLoS Genetics. 2019;15:e1008302.

16. Boitard S, Boussaha M, Capitan A, Rocha D, Servin B. Uncovering adaptation from sequence data: Lessons from genome resequencing of four cattle breeds. Genetics. 2016;203:433–50.

17. Crysnanto D, Wurmser C, Pausch H. Accurate sequence variant genotyping in cattle using variation-aware genome graphs. Genetics Selection Evolution. 2019;51:21.

18. Eggertsson HP, Jonsson H, Kristmundsdottir S, Hjartarson E, Kehr B, Masson G, et al. Graphtyper enables population-scale genotyping using pangenome graphs. Nature Genetics. 2017;49:1654–60.

19. Zimin A V, Delcher AL, Florea L, Kelley DR, Schatz MC, Puiu D. A whole-genome assembly of the domestic cow, Bos taurus. Genome Biology. 2009;10:R42.

20. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. Preprint at the arXiv:13033997v2. http://arxiv.org/abs/1303.3997. Accessed 27 November 2019.

21. Browning SR, Browning BL. Rapid and Accurate Haplotype Phasing and Missing-Data Inference for Whole-Genome Association Studies By Use of Localized Haplotype Clustering. The American Journal of Human Genetics. 2007;81:1084–97.

22. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The Ensembl

Variant Effect Predictor. Genome Biology. 2016;17:122.

23. Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants

on protein function using the SIFT algorithm. Nature Protocols. 2009;4:1073–82.

24. Online Mendelian Inheritance in Animals database (OMIA). https://omia.org/home.

Accessed 27 November 2019.

25. Nicholas FW, Hobbs M. Mutation discovery for Mendelian traits in non-laboratory

animals: A review of achievements up to 2012. Animal Genetics. 2014;45:157–70.

26. Nei M, Li WH. Mathematical model for studying genetic variation in terms of restriction

endonucleases. Proceedings of the National Academy of Sciences of the United States of

America. 1979;76:5269–73.

27. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant

call format and VCFtools. Bioinformatics. 2011;27:2156–8.

28. Narasimhan V, Danecek P, Scally A, Xue Y, Tyler-Smith C, Durbin R. BCFtools/RoH: A

hidden Markov model approach for detecting autozygosity from next-generation sequencing

data. Bioinformatics. 2016;32:1749–51.

29. Xue Y, Prado-Martinez J, Sudmant PH, Narasimhan V, Ayub Q, Szpak M, et al.

Mountain gorilla genomes reveal the impact of long-term population decline and inbreeding.

Science. 2015;348:242–5.

30. van der Valk T, Díez-del-Molino D, Marques-Bonet T, Guschanski K, Dalén L. Historical

Genomes Reveal the Genomic Consequences of Recent Population Decline in Eastern

Gorillas. Current Biology. 2019;29:165–70.

31. McQuillan R, Leutenegger AL, Abdel-Rahman R, Franklin CS, Pericic M, Barac-Lauc L,

et al. Runs of Homozygosity in European Populations. The American Journal of Human

Genetics. 2008;83:359–72.

32. Kirin M, McQuillan R, Franklin CS, Campbell H, McKeigue PM, Wilson JF. Genomic runs

of homozygosity record population history and consanguinity. PloS ONE. 2010;5:e13996.

33. Rocha D, Billerey C, Samson F, Boichard D, Boussaha M. Identification of the putative

ancestral allele of bovine single-nucleotide polymorphisms. Journal of Animal Breeding and Genetics. 2014;131:483–6.

34. Loh PR, Danecek P, Palamara PF, Fuchsberger C, Reshef YA, Finucane HK, et al. Reference-based phasing using the Haplotype Reference Consortium panel. Nature Genetics. 2016;48:1443–8.

35. Gautier M, Klassmann A, Vitalis R. rehh 2.0: a reimplementation of the R package rehh to detect positive selection from haplotype structure. Molecular Ecology Resources. 2017;17:78–90.

36. Granka JM, Henn BM, Gignoux CR, Kidd JM, Bustamante CD, Feldman MW. Limited evidence for classic selective sweeps in African populations. Genetics. 2012;192:1049–64.

37. Huber CD, DeGiorgio M, Hellmann I, Nielsen R. Detecting recent selective sweeps while controlling for mutation rate and background selection. Molecular Ecology. 2015;25:142–56.

38. Nielsen R. Molecular Signatures of Natural Selection. Annual Review of Genetics. 2005;39:197–218.

39. Degiorgio M, Huber CD, Hubisz MJ, Hellmann I, Nielsen R. SweepFinder2: Increased sensitivity, robustness and flexibility. Bioinformatics. 2016;32:1895–7.

40. Qanbari S, Pausch H, Jansen S, Somel M, Strom TM, Fries R, et al. Classic Selective Sweeps Revealed by Massive Sequencing in Cattle. PLoS Genetics. 2014;10:e1004148.

41. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics. 2010;26:841–2.

42. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: More genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic Acids Research. 2019;47:D419–26.

43. Hu ZL, Park CA, Reecy JM. Building a livestock genetic and genomic information knowledgebase through integrative developments of Animal QTLdb and CorrDB. Nucleic Acids Research. 2019;47:D701–10.

44. The Animal Quantitative Trait Loci (QTL) Database. https://www.animalgenome.org/QTLdb/cattle. Accessed 27 November 2019.

45. Das A, Panitz F, Gregersen VR, Bendixen C, Holm LE. Deep sequencing of Danish Holstein dairy cattle for variant detection and insight into potential loss-of-function variants in protein coding genes. BMC Genomics. 2015;16:1043.

46. Murgiano L, Jagannathan V, Piffer C, Diez-Prieto I, Bolcato M, Gentile A, et al. A frameshift mutation in MOCOS is associated with familial renal syndrome (xanthinuria) in Tyrolean Grey cattle. BMC Veterinary Research. 2016;12:276.

47. Rothammer S, Kunz E, Seichter D, Krebs S, Wassertheurer M, Fries R, et al. Detection of two non-synonymous SNPs in SLC45A2 on BTA20 as candidate causal mutations for oculocutaneous albinism in Braunvieh cattle. Genetics Selection Evolution. 2017;49:73.

48. Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C. Genomic scans for selective sweeps using SNP data. Genome Research. 2005;15:1566–75.

49. Sabeti PC, Schaffner SF, Fry B, Lohmueller J, Varilly P, Shamovsky O, et al. Positive natural selection in the human lineage. Science. 2006;312:1614–20.

50. Voight BF, Kudaravalli S, Wen X, Pritchard JK. A map of recent positive selection in the human genome. PLOS Biology. 2006;4:e72.

51. Eberlein A, Takasuga A, Setoguchi K, Pfuhl R, Flisikowski K, Fries R, et al. Dissection of genetic factors modulating fetal growth in cattle indicates a substantial role of the non-SMC condensin I complex, subunit G (NCAPG) gene. Genetics. 2009;183:951–64.

52. Pausch H, Flisikowski K, Jung S, Emmerling R, Edel C, Götz KU, et al. Genome-wide association study identifies two major loci affecting calving ease and growth-related traits in cattle. Genetics. 2011;187:289–97.

53. Bouwman AC, Daetwyler HD, Chamberlain AJ, Ponce CH, Sargolzaei M, Schenkel FS, et al. Meta-analysis of genome-wide association studies for cattle stature identifies common genes that regulate body size in mammals. Nature Genetics. 2018;50:362–7.

54. Jansen S, Aigner B, Pausch H, Wysocki M, Eck S, Benet-Pagès A, et al. Assessment of the genomic variation in a cattle population by re-sequencing of key animals at low to medium coverage. BMC Genomics. 2013;14:446.

55. Miosge LA, Field MA, Sontani Y, Cho V, Johnson S, Palkova A, et al. Comparison of

predicted and actual consequences of missense mutations. Proceedings of the National Academy of Sciences of the United States of America. 2015;112:E5189–98.

56. Pausch H, Kölle S, Wurmser C, Schwarzenbacher H, Emmerling R, Jansen S, et al. A Nonsense Mutation in TMEM95 Encoding a Nondescript Transmembrane Protein Causes Idiopathic Male Subfertility in Cattle. PLoS Genetics. 2014;10:e1004044.

57. Kadri NK, Sahana G, Charlier C, Iso-Touru T, Guldbrandtsen B, Karim L, et al. A 660-Kb Deletion with Antagonistic Effects on Fertility and Milk Production Segregates at High Frequency in Nordic Red Cattle: Additional Evidence for the Common Occurrence of Balancing Selection in Livestock. PLoS Genetics. 2014;10:e1004049.

58. Pausch H, Ammermüller S, Wurmser C, Hamann H, Tetens J, Drögemüller C, et al. A nonsense mutation in the COL7A1 gene causes epidermolysis bullosa in Vorderwald cattle. BMC Genetics. 2016;17:149.

59. Bosse M, Megens HJ, Derks MFL, de Cara ÁMR, Groenen MAM. Deleterious alleles in the context of domestication, inbreeding, and selection. Evolutionary Applications. 2019;12:6–17.

60. Dong C, Wei P, Jian X, Gibbs R, Boerwinkle E, Wang K, et al. Comparison and integration of deleteriousness prediction methods for nonsynonymous SNVs in whole exome sequencing studies. Human Molecular Genetics. 2015;24:2125–37.

61. Schwarzenbacher H, Wurmser C, Flisikowski K, Misurova L, Jung S, Langenmayer MC, et al. A frameshift mutation in GON4L is associated with proportionate dwarfism in Fleckvieh cattle. Genetics Selection Evolution. 2016;48:25.

62. Santer R, Schneppenheim R, Dombrowski A, Gotze H, Steinmann B. Mutations in GLUT2, the gene for the liver-type glucose transporter, in patients with Fanconi-Bickel syndrome. Nature Genetics. 1997;17:324–6.

63. Pausch H, Schwarzenbacher H, Burgstaller J, Flisikowski K, Wurmser C, Jansen S, et al. Homozygous haplotype deficiency reveals deleterious mutations compromising reproductive and rearing success in cattle. BMC Genomics. 2015;16:312.

64. Burgstaller J, Url A, Pausch H, Schwarzenbacher H, Egerbacher M, Wittek T. Clinical

and biochemical signs in Fleckvieh cattle with genetically confirmed Fanconi-Bickel syndrome (cattle homozygous for Fleckvieh haplotype 2). Berliner und Münchener tierärztliche Wochenschrift. 2016;129:132–7.

65. Joller S, Stettler M, Locher I, Dettwiler M, Seefried F, Meylan M, et al. Fanconi-Bickel-Syndrom□: eine bislang unerkannte Erbkrankheit beim Braunvieh. Schweizer Archiv Tierheilkunde. 2018;160:179–84.

66. Goddard ME, Hayes BJ. Genomic selection based on dense genotypes inferred from sparse genotypes. Proceedings of the 18th Conference of the Association for the Advancement of Animal Breeding and Genetics. 2009;18:26–9.

67. Gusev A, Shah MJ, Kenny EE, Ramachandran A, Lowe JK, Salit J, et al. Low-pass genome-wide sequencing and variant inference using identity-by-descent in an isolated human population. Genetics. 2012;190:679–89.

68. Bickhart DM, Hutchison JL, Null DJ, VanRaden PM, Cole JB. Reducing animal sequencing redundancy by preferentially selecting animals with low-frequency haplotypes. Journal of Dairy Science. 2016;99:5526–34.

69. Ros-Freixedes R, Gonen S, Gorjanc G, Hickey JM. A method for allocating low-coverage sequencing resources by targeting haplotypes rather than individuals. Genetics Selection Evolution. 2017;49:78.

70. Charlier C, Li W, Harland C, Littlejohn M, Coppieters W, Creagh F, et al. NGS-based reverse genetic screen for common embryonic lethal mutations compromising fertility in livestock. Genome Research. 2016;26:1333–41.

71. Mei C, Wang H, Liao Q, Wang L, Cheng G, Wang H, et al. Genetic architecture and selection of Chinese cattle revealed by whole genome resequencing. Molecular Biology and Evolution. 2018;35:688–99.

72. Kim J, Hanotte O, Mwai OA, Dessie T, Salim B, Diallo B, et al. The genome landscape of indigenous African cattle. Genome Biology. 2017;18:34.

73. Weldenegodguad M, Popov R, Pokharel K, Ammosov I, Ming Y, Ivanova Z, et al. Whole-Genome Sequencing of Three Native Cattle Breeds Originating From the Northernmost

Cattle Farming Regions. Frontiers in Genetics. 2019;9:728.

74. Zhang Q, Guldbrandtsen B, Bosse M, Lund MS, Sahana G. Runs of homozygosity and distribution of functional variants in the cattle genome. BMC Genomics. 2015;16:542.

75. Solé M, Gori AS, Faux P, Bertrand A, Farnir F, Gautier M, et al. Age-based partitioning of individual genomic inbreeding levels in Belgian Blue cattle. Genetics Selection Evolution. 2017;49:92.

76. Bosse M, Megens H, Madsen O, Paudel Y, Frantz LAF, Schook LB, et al. Regions of Homozygosity in the Porcine Genome□: Consequence of Demography and the Recombination Landscape. PLoS Genetics. 2012;8:e1003100.

77. Purfield DC, Berry DP, McParland S, Bradley DG. Runs of homozygosity and population history in cattle. BMC Genetics. 2012;13:70.

78. Ceballos FC, Joshi PK, Clark DW, Ramsay M, Wilson JF. Runs of homozygosity: windows into population history and trait architecture. Nature Reviews Genetics. 2018;19:220–34.

79. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. American journal of human genetics. 2007;81:559–75.

80. Bertrand AR, Kadri NK, Flori L, Gautier M, Druet T. RZooRoH: an R package to characterize individual genomic autozygosity and identify homozygous-by-descent segments. Methods in Ecology and Evolution. 2019;10:860–6.

81. Wellmann R. Optimum contribution selection for animal breeding and conservation: The R package optiSel. BMC Bioinformatics. 2019;20:25.

82. Cole JB. A simple strategy for managing many recessive disorders in a dairy cattle breeding program. Genetics Selection Evolution. 2015;47:94.

83. Rothammer S, Seichter D, Förster M, Medugorac I. A genome-wide scan for signatures of differential artificial selection in ten cattle breeds. BMC Genomics. 2013;14:908.

84. Biswas S, Akey JM. Genomic insights into positive selection. Trends in Genetics. 2006;22:437–46.

85. Akey JM. Constructing genomic maps of positive selection in humans: Where do we go from here? Genome Research. 2009;19:711–22.

86. Charlesworth B. The role of background selection in shaping patterns of molecular evolution and variation: Evidence from variability on the Drosophila X chromosome. Genetics. 2012;191:233–46.

87. Gutiérrez-Gil B, Arranz JJ, Wiener P. An interpretive review of selective sweep studies in Bos taurus cattle populations: Identification of unique and shared selection signals across breeds. Frontiers in Genetics. 2015;6:167.

88. Zhao F, McParland S, Kearney F, Du L, Berry DP. Detection of selection signatures in dairy and beef cattle using high-density genomic information. Genetics Selection Evolution. 2015;47:49.

89. Xu L, Bickhart DM, Cole JB, Schroeder SG, Song J, Van Tassell CP, et al. Genomic signatures reveal new evidences for selection of important traits in domestic cattle. Molecular Biology and Evolution. 2015;32:711–25.

90. Randhawa IAS, Khatkar MS, Thomson PC, Raadsma HW. Composite Selection Signals for Complex Traits Exemplified Through Bovine Stature Using Multibreed Cohorts of European and African Bos taurus. G3 (Bethesda, Md). 2015;5:1391–401.

91. Joerg H, Fries HR, Meijerink E, Stranzinger GF. Red coat color in Holstein cattle is associated with a deletion in the MSHR gene. Mammalian Genome. 1996;7:317–8.

92. Durkin K, Coppieters W, Drögüller C, Ahariz N, Cambisano N, Druet T, et al. Serial translocation by means of circular intermediates underlies colour sidedness in cattle. Nature. 2012;482:81–4.

93. Karim L, Takeda H, Lin L, Druet T, Arias JAC, Baurain D, et al. Variants modulating the expression of a chromosome domain encompassing PLAG1 influence bovine stature. Nature Genetics. 2011;43:405–15.

94. Cohen-Zinder M, Seroussi E, Larkin DM, Loor JJ, Everts-Van Der Wind A, Lee JH, et al. Identification of a missense mutation in the bovine ABCG2 gene with a major effect on the QTL on chromosome 6 affecting milk yield and composition in Holstein cattle. Genome

Research. 2005;15:936–44.

95. Pausch H, Emmerling R, Gredler-Grandl B, Fries R, Daetwyler HD, Goddard ME. Meta-analysis of sequence-based association studies across three cattle breeds reveals 25 QTL for fat and protein percentages in milk at nucleotide resolution. BMC Genomics. 2017;18:853.

96. Grisart B, Coppieters W, Farnir F, Karim L, Ford C, Berzi P, et al. Positional candidate cloning of a QTL in dairy cattle: Identification of a missense mutation in the bovine DGAT1 gene with major effect on milk yield and composition. Genome Research. 2002;12:222–31.

97. Winter A, Kramer W, Werner FAO, Kollers S, Kata S, Durstewitz G, et al. Association of a lysine-232/alanine polymorphism in a bovine gene encoding acyl-CoA:diacylglycerol acyltransferase (DGAT1) with variation at a quantitative trait locus for milk fat content. Proceedings of the National Academy of Sciences of the United States of America. 2002;99:9300–5.

98. Blott S, Kim JJ, Moisio S, Schmidt-Küntzel A, Cornet A, Berzi P. Molecular dissection of a quantitative trait locus: a phenylalanine-to-tyrosine substitution in the transmembrane domain of the bovine growth hormone receptor is associated with a major effect on milk yield and composition. Genetics. 2003;163:253–66.

Supplementary Information

Additional file 1 File S1

File format: .docx

Title: Original Braunvieh herdbook population

Description: Number of female calves entering the OB herdbook between 1980 and 2016.

Additional file 2 File S2

File format: .pdf

Title: Runs of homozygosity in 49 OB cattle

Description: a) Total genome fraction in ROH in 49 cattle with high (<10x) and low

(<10x) coverage (b) Phred confidence score for ROH in 33 cattle sequenced at

average sequencing depth higher than 10-fold. Red dots indicate mean confidence

scores for ROH.

Additional file 3 File S3

File format: .xlsx

Title: Bovine QTL information downloaded from the AnimalQTL database.

Description: Number of QTL for each trait. Traits are grouped in six trait categories.

Additional file 4 File S4

File format: .pdf

Title: Allele frequency distribution in different functional annotations

Description: Allele frequency of SNPs with different consequences according to VEP

prediction, like high impact, deleterious (missense SNP with SIFT score < 0.05).

tolerated (missense SNPs with SIFT score > 0.05) and synonymous SNPs.

Additional file 5 File S5

File format: .pdf

Title: Distribution of length of Indels

Description: a) Number of Indels (x1000) with size less than 12 bp detected

according to the number of affected bases. b) number of Indels detected in coding

sequences.

Additional file 6 File S6

File format: .xlsx

Title: OMIA variants detected in OB

Description: Six OMIA variants detected in 49 sequenced OB animals with their

respective frequency and information.

Additional file 7 File S7

File format: .xlsx

Title: ROH statistics of each animal (high coverage)

Description: Number, length, average length and genomic fraction of all ROH in each

animal. Also categorized in short, medium and long ROH category.

Additional file 8 File S8

File format: .pdf

Title: Genomic inbreeding in OB cattle stratified by birth year

Description: Genomic inbreeding in two groups of animals born either between 1960

and 1989 or between 1990 and 2012.

Additional file 9 File S9

File format: .xlsx

Title: Candidate selection signatures detected using CLR

Description: Genomic coordinates, CLR values, p values and encompassed genes

for 95 candidate selection signatures.

Additional file 10 File S10

File format: .xlsx

Title: Candidate selection signatures detected using iHS

Description: Genomic coordinates, |iHS| values, p values and encompassed genes

for 162 candidate selection signatures.

Additional file 11 File S11

File format: .xlsx

Title: Overlap between the top CLR and iHS selection signatures

Description: Overlapped regions and genes between CLR and iHS candidate

selection regions

Additional file 12 File S12

File format: .xlsx

Title: Summary of Gene Ontology enrichment analysis

Description: PANTHER and GO-Slim pathways enriched using genes encompassed

in signatures of selection from CLR and iHS analyses.

Additional file 13 File S13

File format: .xlsx

Title: Overlap between QTL and signatures of selection

Description: QTL (from all 6 categories) that overlapped with CLR and iHS selection signatures.

Additional file 14 File S14

File format: .xlsx

Title: Frequency of candidate causal variants for a stature QTL on BTA14

Description: Genomic coordinates and allele frequencies of 14 variants nearby bovine *PLAG1* that were reported as candidate causal variants for a stature QTL in cattle by Karim et al. [95].