Coupling Day Length Data and Genomic Prediction tools for Predicting Time-Related Traits under Complex Scenarios

Diego Jarquin[¥,*], Hiromi Kajiya-Kanegae[†], Chen Taishen[†], Shiori Yabe[‡], Reyna Persa[¥], Jianming Yu[**], Hiroshi Nakagawa[a], Masanori Yamasaki[§], Hiroyoshi Iwata[†,*]

[¥]Department of Agronomy and Horticulture, University of Nebraska – Lincoln, Lincoln NE, 68583, USA.

[†]Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo, Tokyo 113-8657, Japan.

[‡]Institute of Crop Sciences, National Agriculture and Food Research Organization (NARO), Tsukuba, Ibaraki 305-8518, Japan

[**]Department of Agronomy, Iowa State University, Ames, USA.

[§]Food Resources Education and Research Center, Graduate School of Agricultural Science, Kobe University, Kasai, Hyogo 675-2103, Japan

[a]Institute for Agro-Environmental Sciences, National Agriculture and Food Research Organization (NARO), Tsukuba Ibaraki 305-8604

[*]Corresponding authors: University of Nebraska – Lincoln, Lincoln NE, 68583, USA. E-mail: diego.jarquin@gmail.com ORCID ID 0000-0002-5098-2060 (DJ). Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo, Tokyo 113-8657, Japan. E-mail: iwata@ut-biomet.org, ORCID 0000-0002-6747-7036 (HI).

Abstract

Genomic selection (GS) has proven to be an efficient tool for predicting crop-rank performance
of untested genotypes; however, when the traits have intermediate optima (phenology stages)
this implementation might not be the most convenient. GS might deliver high rank correlations
but incurring in serious bias. Days to heading (DTH) is a crucial development stage in rice for
regional adaptability with significant impact in yield potential. The objectives of this research
consisted in the development of a method that accurately predicts time related traits like DTH in
unobserved environments. For this, we developed an implementation that incorporate day length
information (DL) in the prediction process for two relevant scenarios: CV0, predicting tested
genotypes in unobserved environments (C method); and CV00, predicting untested genotypes in
unobserved environments (CB method). The use of DL has advantages over weather data since it
can be determined in advance just by knowing the location and planting date. The proposed
methods showed that DL information significantly helps to improve the predictive ability of
DTH in unobserved environments. Under CV0, the C method returned an root-mean-square-
error (RMSE) of 3.9 days, a Pearson Correlation (PC) of 0.98 and the differences between the
predicted and observed environmental means (EMD) ranged between -4.95 and 4.67 days. For
CV00, the CB method returned an RMSE of 7.3 days, a PC of 0.93 and the EMD ranged
between -6.4 and 4.1 days while the conventional GS implementation produced an RMSE of
18.1 days, a PC of 0.41 and the EMD ranged between -31.5 and 28.7 days.

Introduction

To satisfy the daily demand of crop consumption for a growing population, it is necessary to improve the production systems to allow a sustainable increase of the yield potential and nutritional value of the new and up-coming varieties[1]. In plant breeding there are different tools and methodologies to increase the rate of genetic gain[2]. Traditional plant breeding methods use phenotypic and/or pedigree information to perform selection of experimental lines for developing superior breeding lines. However, phenotypic selection may not be permissible in many situations due to the high phenotyping costs; also, pedigree selection may not be accurate since it assumes a constant rate of recombination (50% of the genetic material coming from mother and the remaining 50% coming from father) and this might not be a feasible premise. In this regard, the use of molecular markers can help to detect the deviations from the expected genetic material of the parents and these can be used to determine which parental haplotypes were inherited.

Genomic selection[3] (GS) is an emerging tool that allows screening genotypes from a very large population without having to observe them in fields[4,5]. This method only requires phenotypic and genomic information for calibrating models, then other genotyped candidates are selected based on the predicted values obtained via their marker profiles[6]. In this context, the implementation of genomic tools and resources translates our understanding of the relationship between genotype and phenotype into predicted genetic merits for direct selection. This is highly desirable dealing with those traits that are controlled by a large number of genes with small individual effects, also known as complex traits[5,7]. GS has shown to be an effective tool for plant and animal breeding applications, especially dealing with complex traits[8]. The successfulness of GS predicting phenotypes from genotypic information has been reported in various crop species[9].

In GS, the genomic estimated breeding values (GEBVs) of complex traits of untested genotypes are computed based on genome-wide marker information. The GEBVs can be considered as deviations around a fixed mean (e.g., zero). Once the GEBVs are obtained, the breeders may conduct selections based on the ranked values by selecting those individuals in the tails. Depending on the trait, the individuals are selected either on the upper or lower tails of the ranked GEBVs. For example, the upper side is selected for increasing genetic gains of traits like grain yield, stress resistance, or the content of essential nutrients. On the other hand, genotypes in the lower tail are selected for reducing lodging, shattering, the content of anti-nutrients, etc. In these cases, the EBVs provide sufficient information when the correlation between the EBVs and the direct genetic values (or phenotypic values) is high. However, the rank-based selection criteria might not be the most accurate when the target trait has an intermediate optima. This is the case of those traits related to phenological stages of crop plants. For example, Onogi *et al*[10]., the prediction of heading date of untested genotypes within and across environments by integrating genomic prediction with crop/ecophysiological modelling. These authors predicted DTH for untested genotypes within environments and for untested genotypes in unobserved environments among other cross-validations schemes. The corresponding root-mean-square-error (RMSE) predicting untested genotypes in unobserved environments was almost twice than for

103   the within environments prediction schemes. Moreover, in both cases high correlations between
104   predicted and observed values were obtained. For the within environments the correlation was of
105   0.96 while for predicting untested genotypes in unobserved environments it was of 0.87. These
106   results highlight the importance of improving the accuracy of predicting untested genotypes in
107   unobserved environments despite the fact of obtaining high correlations.

108   In rice, days to heading (DTH; typically defined as the number of days after planting
when more than 50% of individuals show the heading stage) is one of the critical traits of
ideotypes with an increased yield potential. Hence it is important to optimize DTH for high yield
potential in different environments[11]. Usually multi environment trials (METs) are performed for
developing stable cultivars across a wide range of environmental conditions, and the breeders
may be interested in estimating DTH for a large set of environments instead of a single one.
However, when the GEBVs are computed for an unobserved environment these might contain a
large bias if the phenotypic information of genotypes in the training set do not resemble the
target environment. The GEBVs of one environment may not be valid for another environment
due to variations in the environmental *stimuli* causing inconsistencies in the response patters as
result of the interaction between genotypes to environments (G×E interaction). These differences
remain also for same genotypes observed in same locations but tested at different times (planting
date, years). As consequence of the aforementioned issues, the problem of correctly estimating
DTH for a set of untested genotypes in a wide range of unobserved environments becomes
harder to solve than for non-time related traits. As pointed before, the advantage of GS
techniques over conventional phenotypic methods is its ability for performing accurate rank
predictions of untested genotypes based exclusively on their marker profiles without having to
observe these in fields. Thus, a method capable of delivering highly accurate predictions of
untested genotypes in unobserved environments without incurring in a large bias is desirable.

127   As was pointed before, DTH in rice is important for regional adaptability and has usually
great impact on yield potential[11], genetic systems controlling it have been studied for a long time
by integrating Mendelian genetics and genomics[12]. The potential of GS in DTH has been
evaluated in rice[13,14] and sorghum[15]. Onogi *et al.*[10] focused on the genomic prediction of DTH in
multiple environments and demonstrated the potential of a novel method that integrates GS with
a crop growth model. Li *et al.*[15] also focused on the genomic prediction of DTH in multiple
environments and demonstrated the potential of models combining GS with joint regression
analysis on photothermal time. To our knowledge, no studies combining DL at the day when
genotypes reach DTH with GS tools have been done before. The objectives of this research
consisted in the development of a method that accurately predicts time related traits like DTH in
unobserved environments by leveraging information from day length and molecular markers.

138   Two cross-validation schemes of interest for designing and planning experiments were
considered for evaluating the performance of the proposed implementation for predicting DTH in
unobserved environments: CV0, predicting already tested genotypes in unobserved environments
(targeting environments); and CV00, predicting untested genotypes in unobserved environments
(targeting genotypes and environments). For assessing the proposed implementation the root-

mean-square-error (RMSE), the Pearson correlation (PC) and the difference between predicted
and observed environmental means (EMD) values were computed.

In this study, we propose a novel methods to predict DTH in a precise and accurate way
(i.e., small bias and high correlation between predicted and observed values) of tested (C
method) and untested (CB method) genotypes in unobserved environments.

Under the hardest cross-validation scheme (i.e., CV00), the results of the introduced CB
method are compared with those obtained using the conventional GS implementation (G-BLUP).
For studying the realistic scenario of predicting already tested genotypes in unobserved
environments, the cross validation scheme CV0 was considered. These results are used for
assessing the levels of predictability that can be reached when phenotypic information of a
genotype is available in other environments. Here the environments are considered "target" in a
strict sense because these can be selected even before conducting field experiments using the
theoretical DL values of locations for a given planting date.

Results

Descriptive statistics. The box-plot of DTH is depicted in Figure S1. The environments were
ordered based on the environmental medians. In general, those environments with late planting
date appear first. The medians ranged between 59 ("Fukuyama 2010 Late") and 116 ("Akita
2015") days. The SDs of the environments varied between 5.07 and 16.83 days. Also, the
smallest and the largest DTH values were 42 and 148 days and these were observed in
"Tsukubamirai 2016 Late" and "Akita 2015" environments, respectively.

Evaluating the bias and accuracy of the methods. Table 1 contains the observed and the
predicted environmental means (E-mean) for two different cross-validation schemes (CV00 and
CV0) as well as the corresponding RMSE and the PC between predicted and observed values.
For the CV00 scheme, the CB and GS methods were implemented. The objective of using this
cross-validation scheme was to assess the model performance of the proposed method with
respect to the conventional GS implementation for predicting untested genotypes in unobserved
environments. Since no phenotypic information is available for connecting yet to be predicted
genotypes in unobserved environments with genotypes in training sets, the CV00 scheme is
considered the hardest and the most interesting prediction problem. Across environments, the
conventional GS method returned an RMSE of 18.1 days and a PC of 0.41; within environments,
the average RMSE and PC were of 17.5 days and 0.69. The EMD ranged between -31.5 and 28.7
days. On the other hand, across environments the proposed CB method returned an RMSE of 7.3
days and a PC of 0.93; within environments, the average RMSE and PC were of 7.2 days and
0.86. The EMD ranged between -6.4 and 4.1 days.

The C method was implemented under the CV0 scheme and their results were used for
assessing the predictive ability of predicting an already tested genotype in an untested
environments considering only the phenotypic and DL information of the same genotype. Across
environments, this method returned an RMSE of 3.9 days and a PC of 0.98; within

environments, the average RMSE and PC were of 3.7 days and 0.98. The EMD ranged between -
4.95 and 4.67 days.

Table 1. Environmental means, Root Means Squared Error (RMSE) and the Pearson correlation between predicted
and observed days to heading (DTH) for 112 rice genotypes tested in 51 environments in Japan. Two cross-
validation schemes were implemented for mimicking realistic prediction scenarios: CV0 represents the scenario of
predicting tested genotypes in unobserved environments; CV00 considers the prediction of untested genotypes in
unobserved environments. For CV0, the C method was used combining phenotypic and day length information (DL)
of tested genotypes (one at a time) in other environments. For CV00, two methods were considered: (*i*) the
conventional GS implementation; and (*ii*) the CB method, which combines phenotypic and DL information from the
tested genotypes (training set) and genomic BLUP values ($\hat{g}$) of the untested genotypes. In both cases, the
prediction procedure was conducted by leaving one genotype out across environments and deleting any phenotypic
information from the target environment as well.

	CV00							CV0		
	Mean			RMSE		Corr		Mean	RMSE	Corr
	Real	CB	GS	CB	GS	CB	GS	C	C	C
Tsukubamirai 2004 Late	79.4	76.7	85.9	7.0	12.6	0.86	0.70	76.3	4.6	0.99
Tsukubamirai 2004 Early	87.5	93.2	84.1	10.0	12.9	0.84	0.55	92.2	6.4	0.97
Tsukubamirai 2005 Early	98.5	102.1	87.1	6.9	14.0	0.87	0.75	101.0	3.5	0.98
Tsukubamirai 2005 Late	76.7	74.8	84.8	5.7	12.4	0.86	0.68	73.7	3.4	0.99
Kasai 2006	87.4	83.8	88.3	7.9	8.8	0.85	0.75	82.8	6.8	0.93
Fukuyama 2006	80.6	83.0	82.5	5.6	13.4	0.87	0.29	82.0	2.0	0.99
Fukuyama 2007	84.1	83.4	86.6	5.0	6.1	0.87	0.84	82.5	2.9	0.98
Tsukuba 2008	104.5	107.7	87.2	9.1	19.2	0.87	0.85	107.3	4.0	0.98
Kasai1st 2008	102.2	99.9	88.7	9.0	17.8	0.89	0.71	99.2	4.3	0.99
Fukuyama 2008	77.8	82.2	85.6	7.8	11.1	0.86	0.77	81.3	4.5	0.98
Tsukuba 2009	103.9	107.7	87.1	8.6	19.0	0.89	0.81	108.0	5.2	0.98
Kasai1st 2009	101.9	100.6	89.4	7.6	15.1	0.89	0.83	100.0	2.7	0.99
Fukuyama 2009	78.7	78.7	89.6	5.2	13.2	0.86	0.83	77.8	1.5	0.99
Tsukuba 2010 Late	83.5	89.1	88.9	8.4	11.1	0.86	0.66	88.4	5.9	0.97
Kasai1st 2010	100.6	100.7	89.5	7.1	14.3	0.88	0.77	100.1	1.9	0.99
Fukuyama 2010	76.2	79.5	88.2	5.9	15.3	0.86	0.50	78.1	2.6	0.99
Fukuyama 2010 Late	59.0	65.4	87.7	9.2	29.7	0.84	0.82	62.5	4.2	0.96
Tsukuba 2011 Early	105.9	109.9	85.5	9.9	23.9	0.87	0.66	109.9	5.3	0.98
Tsukuba 2011 Late	82.8	80.8	90.0	5.9	11.9	0.86	0.78	80.4	4.1	0.95
Tsukuba 2011 Middle	94.7	95.7	90.9	6.5	11.1	0.88	0.67	95.4	3.0	0.98
Kasai 2011	101.4	100.6	89.3	7.8	15.7	0.89	0.76	100.0	2.4	0.99
Fukuyama 2011	75.7	79.4	91.1	6.4	16.9	0.87	0.83	78.6	3.1	0.99
Fukuyama 2011 Late	66.4	72.7	97.9	8.9	32.4	0.68	0.58	66.2	2.9	0.95
Tsukuba 2012 Early	97.1	96.0	86.9	5.7	12.4	0.81	0.74	101.1	5.8	0.97
Tsukuba 2012 Late	84.3	81.6	88.2	6.1	7.5	0.86	0.80	81.0	4.3	0.96
Tsukuba 2012 Middle	92.6	95.4	88.4	6.3	8.3	0.87	0.82	94.8	3.7	0.97
Kasai 2012	103.8	102.7	86.4	7.9	21.3	0.89	0.60	102.1	2.7	0.99
Fukuyama 2012	75.2	78.8	89.9	6.2	16.3	0.87	0.83	78.0	3.1	0.99
Fukuyama 2012 Late	71.5	77.0	103.1	7.5	32.4	0.70	0.66	68.6	3.4	0.98
Fukuoka 2012	79.6	76.0	89.1	6.7	13.0	0.84	0.69	75.4	5.2	0.96
Akita 2012	113.1	115.9	90.1	8.3	26.1	0.86	0.54	114.5	2.8	0.98
Kasai 2013	104.8	104.1	87.3	8.0	20.1	0.90	0.80	103.5	3.2	0.98
Fukuyama 2013	74.9	80.1	89.4	7.4	16.3	0.87	0.82	79.3	4.6	0.99
Fukuyama 2013 Late	62.1	64.8	85.7	6.3	27.1	0.83	0.36	64.1	3.6	0.93
Fukuoka 2013	77.4	77.3	93.3	5.7	17.5	0.86	0.77	76.7	1.9	0.99
Akita 2013	116.6	115.8	91.1	8.5	28.1	0.88	0.66	115.1	2.4	0.99
Kasai 2014	102.4	105.1	88.9	8.4	18.0	0.90	0.66	104.4	2.8	0.99
Fukuyama 2014	79.6	82.1	87.9	5.9	12.1	0.88	0.68	81.3	2.1	0.99
Fukuyama 2014 Late	67.6	64.5	88.1	6.7	21.4	0.83	0.79	62.9	5.1	0.98
Fukuoka 2014	81.8	77.7	80.4	6.7	11.5	0.87	0.45	77.1	5.4	0.97
Akita 2014	115.2	116.2	89.6	9.2	28.5	0.88	0.68	115.6	2.2	0.99
Fukuyama 2015	76.3	72.6	88.1	6.5	14.4	0.86	0.63	71.9	4.6	0.99
Akita 2015	118.1	119.8	89.4	7.7	29.9	0.81	0.76	117.6	2.5	0.99
Kasai 2016	102.3	101.2	85.6	7.2	20.4	0.90	0.61	100.7	3.1	0.98
Fukuyama 2016	79.1	81.1	85.2	6.0	8.9	0.86	0.82	80.8	2.4	0.99
Fukuyama 2016 Late	67.5	65.2	89.6	6.8	26.1	0.79	0.63	64.6	4.5	0.95
Tsukubamirai 2016 Late	73.6	77.9	84.6	7.0	13.6	0.90	0.80	77.2	4.9	0.98
Kasai 2017	108.6	106.2	85.7	8.3	26.3	0.89	0.56	105.5	4.3	0.98
Fukuyama 2017	77.4	80.2	87.7	6.3	16.3	0.88	0.42	79.6	2.7	0.99
Fukuyama 2017 Late	68.4	67.7	88.8	6.0	22.9	0.82	0.69	67.0	3.0	0.97
Tsukubamirai 2017 Late	69.5	73.1	86.9	6.8	19.8	0.87	0.65	72.2	4.3	0.98
Mean				**7.2**	**17.5**	**0.86**	**0.69**		**3.7**	**0.98**

The results of CB and GS methods are directly comparable between them (CV00
scheme). Figure 1 depicts the scatter plot between predicted and observed DTH values for all
genotypes across environments. The top panel contains the results obtained with the proposed
CB method, while the bottom panel shows the results derived from the conventional GS
implementation. The blue and black lines correspond to the fitted and the on target predictions
with respect to the observed DTH values. In general, across environments (Figure 1) and within
environments (Table 1) the results obtained with the CB method showed a better fit than those
derived from the GS implementation.

209
1. Figure 1. Scatter plot of predicted (*x-axis*) and observed (*y-axis*) DTH values under CB (top) and GS (bottom)
methods for the CV00 scheme (predicting untested genotypes in unobserved environments) for a rice data set
comprising 112 genotypes tested in 51 environment in Japan between 2005 and 2017. The CB method combines the

predicted environmental means (E-mean) and the genomic BLUPs ($\hat{g}$) derived from the conventional GS
implementation while the GS implementation considers a common mean across environments plus the genomic
BLUPs. The blue line represents the fitted line between predicted and observed DTH values, and the black line show
the exact match (on target) of the predicted values on the observed values.
217

218
The advantages of CB method for predicting DTH are clearly appreciated considering the
predicted environmental means (E-means). Figure 2 displays the predicted (*x-axis*) and observed
(*y-axis*) E-means under the CB (orange) and GS (blue) methods. As expected, when no
information from the target environment was available the conventional GS implementation
predicted the DTH values to be centered on the overall mean (~88 days) of the environments
observed in training set. On the other hand, the CB method consistently returned closer values to
the diagonal black line, which represents the exact match.
226

227
2. Figure 2. Scatter plot of predicted and observed environmental means (E-means) using the CB method (orange)
and the conventional GS implementation (blue) for the CV00 scheme (predicting untested genotypes in unobserved
environments) for a rice data set comprising 112 genotypes tested in 51 environment in Japan between 2005 and
2017. The vertical pink line indicates the position of the overall phenotypic mean across environments (88 days).
The diagonal black line shows the exact match between predicted and observed means. ARMSE and ACP stand for
the Average Root Mean Square Error and the Average Person Correlation between predicted and observed DTH
values across environments. The RMSE and PC were first computed within environments and the obtained values
were averaged.

Discussion

The boxplot (Figure S1) showed a high heterogeneity among the environments in the study with
those with "late" planting date exhibiting smaller phenotypic variability and shorter occurrence

time for DTH compared with the "early" planting environments. Among other considerations,
the conventional GS models assume homogeneity of variances between environments. However,
the violation of this assumption may induce serious inconsistencies in the results. Especially
when genotype-by-environment interaction parameters are not explicitly included or accounted
in models.

The results from GS model provide an overview of the possible problems that we might
incur predicting time related traits when no phenotypic information from the target environment
is available (CV00). This cross-validation scheme represents among the four prediction scenarios
that breeders face in field (CV2: predicting tested genotypes in observed environments; CV1:
predicting untested genotypes in observed environments; CV0: predicting tested genotypes in
unobserved environments; and CV00: predicting untested genotypes in unobserved
environments) the most interesting, useful and challenging. Dealing with traits where the
selection process is based in ranks rather than in the *per se* estimated values, the conventional
implementation (GS) usually provides good results (moderate to high correlations) under the four
schemes.

However, in traits related with important growth stages like DTH where the *per se*
estimated values are crucial for designing, planning and establishing future experiments (i.e.,
CV0 and CV00 situations) the implementation of the conventional implementations (GS) might
not be the best choice. When no phenotypic information of the target environment is available,
the E-mean of the target environment is determined as the weighted average of those
environments in the training set forcing it to be centered towards a general mean corresponding
to the weighted mean of those environments in the training set. Hence, the final prediction is
composed of this general weighted mean plus deviations corresponding to the
estimated/predicted genomic effects. Since these deviations are conceptualized to be centered
round zero, under CV0 and CV00 schemes the GS method would produce similar environmental
means (i.e., centered on an overall mean determined by the environments in the calibration set).

The results from the C method showed that this implementation provides good
estimations of DTH for the CV0 scheme. Across environments, the RMSE was of 3.9 days and
for a PC of 0.98. These results were very similar to those obtained using the conventional GS
models when complete information of the genotypes across environments was available (i.e.,
CV2 scheme). Under CV2, the SMSE and Pearson correlation across environment were of 3.8
days and 0.98, respectively (data not shown).

As mentioned before, it is expected to obtain similar environmental means using either
CV0 or CV00 schemes under the conventional GS implementations. Under the CV00 scheme,
the GS strategy returned a RMSE of 18.1 and a PC of 0.41 across environments. The EMD
between predicted and observed values ranged between -31.5 and 28.7 days. On the other hand,
the C method returned an EMD ranging between -4.95 and 4.67 days under the CV0 scheme. In
this case, only the phenotypic information of the genotype to be predicted in an observed
environment but tested in others was used. Since under the CV0 and CV00 schemes the
conventional GS implementation would produce similar E-means when no information of the

target environment is available then it is also expected that the proposed C method will produce
better results than those derived from the GS implementation under the CV0 scheme.

As was pointed before, one of the main goals of this research was to develop a method
able to deliver small RMSE between predicted and observed values when predicting DTH of
untested genotypes in unobserved environments (CV00 scheme). Under this scenario the CB
method produced encouraging results. The RMSE across environments was of 7.3 days and a PC
of 0.86. A significant improvement was achieved compared with those results obtained using the
conventional GS implementation (RMSE of 18.1 days and PC of 0.41). In this case, the CB
method doubled the accuracy of the conventional GS method and reduced the SMSE by around
60%. However, the advantages of using this method under CV00 scheme can be appreciated
better when comparing the predicted environmental means. Under the conventional GS
implementation, the differences between predicted and observed environmental means ranged
between -31.5 and 28.7 days while with the CB method these only ranged between -6.4 and 4.1
days. The CB method reduced the extremes (left and right) of the range obtained with the GS
implementation in between 80 and 85%, respectively. The advantages of the CB method are
evident in Figure 2 were their environmental means (orange dots) were always closer to the
diagonal line that represents the exact match with true means compared with those means from
the conventional GS implementation (blue dots).

At first, the implementation of the C and CB methods requires enough information for
accurately characterize the genotype's DL curves. For this, sets of genotypes tested in several
environments are necessary. However, once a good characterization of the DL curves is
accomplished potentially any untested genotypes may be accurately predicted in any unobserved
environment. Thus, the obtained results should be accurate not only for the set of genotypes and
environments analyzed in this study but also for others.

The methods here introduced (C and CB) showed that the use of DL information
significantly improves the predictive ability of the conventional GS implementations with a
significant reduction of the RSME between the predicted and observed DTH values. An
advantage of the C method is that if phenotypic information of the same genotype is available for
a set of environments, it is easy to estimate the expected DTH in an unobserved environment
without needing molecular marker information or phenotypic information from other genotypes.
The CB method improved the prediction of DTH of untested genotypes in unobserved
environments by coupling the marker profiles of the untested genotypes with the phenotypic and
DL information from other genotypes.

Both of the introduced methods require the day length (DL) information of the
environments (training and testing) in their design. The convenience of these methods resides in
the fact that the theoretical DL information can be obtained in advance before establishing and
designing the experiments thus the day length data can be help breeders to schedule the crop
season better. The proposed methods capture the genotype-by-environment interaction (G×E)
component via the fitted curves thus the explicitly inclusion of this term is not needed. The
conventional implementation (GS) returns the same estimated genomic effect across
environments for a particular genotype when the G×E component is absent. The curves precisely

capture the sensitivity experienced by genotypes under different environments driving specific
responses to each environment. Thus, although the CB method is composed of the genomic
component computed without explicitly considering the G×E term the final prediction accounts
for this interaction term via the curves. In addition, the formal modeling of heterogeneity of
variances is not necessary because these patterns are also accounted with the current method via
the curves.

Materials and Methods

Phenotypic data. The phenotypic data set comprises the seeding and heading date of 112
Japanese rice cultivars with different origin from different regions and these correspond to
improved lines and landraces (Table S1). These genotypes were tested in 51 locations (North to
South of Japan) from 2004 to 2017 (Table S2). This multi-environmental trials (METs)
experiment was conducted for investigating heading date related genes. The materials had
polymorphisms known heading date related genes (e.g., Hd1, Hd2, Hd5 and Hd6).

The number of DTH was calculated as the difference in days between the heading date and the
seeding date. The heading date was determined when more than 50% of the individuals in the
plot reached the heading stage.

Marker genotype data. Details procedures of DNA extraction and library construction for
whole genome shotgun sequence are described by Yabe *et al*[17].. For each variety, total DNA was
extracted using the CTAB method[18]. The library was constructed using the Illumina TruSeq
DNA Sample preparation kit according to manufacturer's instructions (Illumina, Inc., San Diego,
CA, USA). Next-generation sequencing data was obtained using the Illumina HiSeq 2000, HiSeq
4000 and HiseqX systems via paired-end sequencing (Illumina, Inc., San Diego, CA, USA). The
sequence data have been deposited in the DNA Data Bank of Japan (DDBJ) Sequence Read
Archive, under Submission DRA008071. Data sets deposited in the DDBJ Sequence Read
Archive (SRA106223, ERA358140, DRA000158, DRA000307, DRA000897, DRA000927,
DRA007273, DRA007256) were reanalyzed. Adapters and low-quality bases were removed
from paired reads using the Trimmomatic v0.36 program[19]. The preprocessed reads were mapped
to the Os-Nipponbare-Reference-IRGSP-1.0[20] by using the bwa-0.7.12 mem algorithm with the
default options[21]. The mean depth of the reads against the Os-Nipponbare-Reference-IRGSP-1.0
ranged from 7.4x to 72.4x with average of 16.9x.

SNP calling was based on alignment using the Genome Analysis Toolkit[22] (GATK,
3.7-0-gcfedb67) and Picard package V2.5.0 (http://broadinstitute.github.io/picard). The mapped
reads were realigned using RealignerTargetCreator and indelRealigner of GATK software. SNPs
and InDel were called at the population level using the UnifiedGenotyper of GATK with the -
glm BOTH option. Heterozygous loci were treated as missing, owing to the assumption that
almost all markers were well fixed in rice cultivars after several generations of self-pollination.
We used only biallelic non-missing sites over all cultivars with a minor allele frequency (MAF)

≥ 0.025. Initially, 629,602 SNP markers were considered in the study. After applying discarding
those with more than 50% of missing values 408,372 SNPs remaining in the study for analysis.

Day length data. Although there are several definitions for day length (DL) according to the
Smithsonian Institution[23] roughly speaking it can be defined as the amount of time in a day with
sunlight. The theoretical DL values were computed by the default method[16] included in the
geosphere (v1.5-5) R package[24]. DL information can be known in advance just by knowing the
location and date and it is not affected by the weather conditions.

Cross-Validation Schemes. Two cross-validation schemes were considered in the study and
these mimic different situations that breeders might face in fields. Figure S2 shows the graphical
representation of CV0 (predicting tested genotypes in unobserved environments) and CV00
(predicting untested genotypes in unobserved environments). For CV0, the cross-validation is
conducted by deleting the phenotypic information for all genotypes at each environment (one at a
time) and using the remaining environments as training set. Using the information from Figure
S2 (top panel), this procedure is sequentially repeated six times (one for each environment).

For implementing the CV00 scheme, for each environment in the testing set all the
phenotypic information is deleted as in the previous cross-validation scenario; however, also
sequentially the phenotypic information from each one of the genotypes in the testing
environment is deleted from all environments (one at a time). In the bottom panel of Figure S2,
we have that the goal was to predict genotype G3 in environment E4. For this environment, the
same procedure is repeated for each one of the genotypes (G1-G5). Such that with six
environments and five genotypes we have that this procedure is implemented 30 (5×6) times.

GBLUP model. The Genomic Best Linear Unbiased Predictor model (Genomic BLUP or
GBLUP) is a main effects model used in GS applications[25,8]. It attempts to explain the DTH
value of the i^{th} genotype (i=1, 2,…, I) observed in the j^{th} environment (j=1, 2, …, J) as the sum of
a common mean, random environmental and genomic effects, and an error term and it is
represented as follows:

$$Y_{ij} = \mu + E_j + g_i + \varepsilon_{ij} \qquad\qquad [1]$$

where μ is a common effect to all genotypes across all environments; E_j are IID (independent
and identically distributed) outcomes from a $N(0, \sigma_E^2)$ representing the effect of the j^{th}
environment; $g = \{g_i\} \sim N\left(0, G\sigma_g^2\right)$ is the vector of genomic effects, with $G = \frac{XX'}{p}$ as the genomic
relationship matrix whose entries describe the genetic similarities between pairs of individuals, X
is the standardized (by columns) matrix and p is the number of molecular markers; ε_{ij} are IID
outcomes from a $N(0, \sigma_\varepsilon^2)$ representing measure errors; and σ_E^2, σ_g^2 and σ_ε^2 are the corresponding
associated variance components of the described random effects.

Usually, when METs are analyzed the environmental component explains the larger
proportion of the phenotypic variability across environments[26,27]. Hence, from [1] is concluded

that the predicted value of the i^{th} genotype in the j^{th} environment would strongly depend on the
environmental mean, which is composed by a common mean across environments plus an
environmental deviation that corresponds to the effect of j^{th} environment.

Conceptually, the common mean effect can be viewed as the average performance of all
genotypes across environments. The environmental deviation corresponds to the mean of the
specific affectations of the current environmental *stimuli* in the set of the tested genotypes
beyond the common mean. Thus, these environmental effects can be considered deviations
around the common mean. Hence, by construction the contribution of the genomic component to
the idealized phenotype is much smaller in comparison with the overall mean and the
environmental effects.

If phenotypic information (DTH) of the target environment is available, good
approximations of the environmental effects can be computed. Otherwise, the environmental
effects are estimated using the phenotypic information of the genotypes observed in other
environments. Thus, the environmental component is estimated as the weighted average of the
environmental effects in the training set. Since these random effects are conceptualized to be
centered on zero, the weighted average will be close to zero. Hence, the predicted DTH value
will be compose mainly of a common mean and a small genomic effect shrinking the predicted
values towards the common mean. Although this method usually delivers high correlations
between predicted and observed values, it also might induce a large bias (RMSE) negatively
affecting the accuracy of the prediction of time related traits.

For alleviate this, the two introduced methods (C and CB) incorporate DL information in
the prediction process. The idea for using DL information in the prediction process was
motivated when examining the relationship between the Cumulative Day Length (CDL) and
DTH of all genotypes across environments (Figure S3). A simple linear regression of DTH on
CDL returned an R-squared value of 0.99 exhibiting a strong relationship between these two
variables. From there we can conclude that if CDL values where to be known then accurate
estimations of DTH may be obtained for any genotype in any environment. However, in order to
compute the CDL the exact number of days between planting date and the heading occurrence
(DTH) should be known at first complicating the estimation process.

Since each genotype was observed in a considerable number of environments (48-50)
these might show response patterns between DTH and DL when analyzing one at a time. In this
study, we observed that across environments the DL values at the DTH occurrence showed
strong relationship response patterns with DTH. For illustrating the conceptualization of this idea
we use as example the data depicted in Figure 3. It corresponds to phenotypic (DTH) and DL
information of genotype 56 observed in 50 environments out of 51 (1-46, and 48-51 but not in
env 47). The goal in this example is to estimate DTH for this genotype in the unobserved
environment (env 47: Tsukubamirai 2016 Late). For each environment, the planting date was
considered as day 1 (*x-axis*) then the daily DL (*y-axis*) values were plotted (pink lines) from day
1 (planting) until genotype 56 reached DTH (blue dots). In Figure 3, a pattern relating DTH on
DL at the occurrence day can be identified. This gradient seems not to be strongly affected by the
initial DL values at the different planting dates across environments. However, it appears to be

linked with the initial DL's changing rate at the planting day. Those environments where DL's

changing rates decreasing at the planting time (late cultivars) showed shortest DTH values and

vice-versa.

For CV00 scheme, the proposed method requires: (*i*) first the estimation of DTH for a set

of tested (training) genotypes (one at a time) yet to be observed in a target/unobserved

environment (CV0); (*ii*) then the computing of the expected environmental mean of the

unobserved environment using the estimated values derived from (*i*) step; (*iii*) and finally to add

to the estimated environmental mean the resulting BLUPs of the untested genotypes derived

from conventional GS methods. Further details of these steps are given below.

C method. The DTH estimation of a tested genotype in an unobserved environment (CV0)

requires phenotypic and DL information across several environments. The first step considers the

characterization of DL at the heading time of occurrence across environments. For this, a third

degree polynomial equation is used for fitting DL as function of DTH (orange curve in Figure 3)

as follows:

453
$$dl_{ij} = b_0 + b_1 t + b_2 t^2 + b_3 t^3 \qquad [1]$$

454   where dl_{ij} represents the observed DL of the i^{th} genotype in the j^{th} environment; t is the time

455   independent variable; b_0, b_1, b_2 and b_3 are the model coefficients; and e_{ij} is the error term and it

456   is assumed to be IID $N(0, \sigma^2)$. This model is referred as C_1.

457   Similarly, the daily DL information of the target environment (green curve) is modeled

458   using a third degree polynomial equation:

459
$$E_j = b_{E0} + b_{E1} t + b_{E2} t^2 + b_{E3} t^3 \qquad [2]$$

with b_{0E}, b_{1E}, b_{2E} and b_{3E} acting as the correspondent model coefficients. The resulting

fitted model is referred as C_2. The solution to this system of two equations (C_1, C_2) will produce

an estimate of DTH of the genotype in the target environment. In order to find the intersection of

these two curves numerical evaluation was implemented. In this example, the solution was found

to be 93 days to heading (Figure 3; vertical dotted gray line) while the observed DTH was of 93

days.

Geno_56 env 47 Tsukubamirai 2016 Late

3. Figure 3. Graphical representation of the C method for estimating days to heading (DTH, *x-axis*) of genotype 56 in environment 47 (env 47 Tsukubamirai 2016 Late). Pink curves represent daily day length in hours (DL, *y-axis*) values from planting day (day 1) until heading time for genotype 56 in 50 environments (1-46, and 48-51). The blue dots indicate the corresponding DL values when genotype 56 reached heading time. The green curve (C_2) represents the target environment (Environment 47) where no genotypes have been observed yet. The orange curve (C_1) represents the fitted line for genotype 56 using a third degree polynomial equation relating DL at the occurrence time (DTH across environments). The vertical dotted gray line shows the intersection (93 days) between the green and the orange curves (C_1, C_2). This value is the estimated DTH for genotype 56 in the unobserved environment "env 47 Tsukubamirai 2016 Late". The vertical dotted black line corresponds to the actual DTH (90 days) value for env 47.

CB Method. Since the ultimate goal is to predict DTH for untested genotypes in unobserved environments, the first step is to accurately estimate the expected DTH mean of the target environment. Then, the final predicted value is composed of the expected DTH mean of the target environment plus the genomic estimated breeding value ($\hat{g}_i$) of the untested genotype obtained via conventional GS implementation. For estimating the DTH mean, the C method was first implemented for predicting DTH on the target environment. Here, only the phenotypic and DL information of those genotypes in training set was considered and the predicted values were averaged.

Figure 4 shows the process for predicting untested genotypes in unobserved environments using a toy example consisting of five genotypes (G_1, G_2, G_3, G_4, and G_5) and four environments (E_1, E_2, E_3, and E_4). In this example we describe the procedure for predicting DTH of the untested genotype G_5 in the unobserved environment E_4. The top panel of Figure 4 contains the observed values of genotypes 1-4 in environments 1-3; the middle panel shows the

predicted values presumably obtained with the C method for genotypes 1-4 (104, 105, 110 and
114) in the unobserved environment 4, also the predicted E-mean (108.3) was computed using as
the mean of the predicted DTH values; and the bottom panel shows the predicted DTH for
genotype G_5 in environment 4 (106). This value (106 = 108.3-2.3) was obtained as sum of the E-
mean and the BLUP value (2.3) obtained for genotype G_5 using the conventional GS
implementation.

		G1	G2	G3	G4	G5	Mean
Observed	**E1**	65	69	71	66		67.8
	E2	80	85	87	90		85.5
	E3	90	95	100	102		96.8
	E4						
Predicted	**E1**						
	E2						
	E3						
	E4	104	105	110	114		108.3
E-Mean + BLUP	**E1**						
	E2						
	E3						
	E4					106	

499
4. Figure 4. Representation of the procedure for predicting DTH for the untested genotype G_5 in the unobserved
environment E4 using information of four genotypes (G_1, G_2, G_3 and G_4) observed in three environments (E1, E2
and E3). The top panel contains the observed values for genotypes (G_1, G_2, G_3 and G_4) in environments (E_1, E_2, and
E_3). The middle panel contains the predicted values that presumably were obtained with the C method for these
genotypes in the unobserved environment E4. Also the E-mean (108.3) of E4 is computed as the mean of the
predicted values for genotypes G_1, G_2, G_3 and G_4. The bottom panel contains the predicted DTH value (106) of
genotype G_5 in environment E4 as the sum of the E-mean (108.3) computed in the previous step and the BLUP (2.3)
obtained with the conventional GS implementation.

Software
The C and CB methods as well all the statistical analyses were performed using the R-software R Core
Team (2018)[28]. The conventional GS model described was fitted using the Bayesian Generalized
Linear Regression (BGLR) R-package[29, 30].

Data availability. Genome data can be accessed from DDBJ under submission DRA008071.

516   **REFERENCES**

1. FAO, IFAD, UNICEF, WFP and WHO. 2018. The State of Food Security and Nutrition in the
World 2018. Building climate resilience for food security and nutrition. Rome, FAO

2. Bernardo R (2002) Breeding for Quantitative Traits in Plants. Stemma Press, USA.

3. Meuwissen TH, Hayes BJ, Goddard ME (2001) Prediction of total genetic value using
genome-wide dense marker maps. Genetics 157:1819–1829.

4. Heffner HL, Sorrells ME, Jannink JL (2009) Genomic selection for crop improvement. Crop
Science 49: 1–2.

5. Jannink JL, Lorenz AJ, and Iwata H (2010) Genomic selection in plant breeding: from theory
to practice. Brief. Funct. Genomics 9: 166–177.

6. Crossa, J., G. de los Campos, P. Pérez, D. Gianola, J. Burgueño, J.L. Araus, D. Makumbi,
R.P. Singh, S. Dreisigacker, J. Yan, V. Arief, M. Banziger, and H.-J. Braun. 2010.
Prediction of genetic values of quantitative traits in plant breeding using pedigree and
molecular markers. Genetics 186:713–724.

7. Perez-de-Castro, A.M., S. Vilanova, J. Canizares, L. Pascual, J.M. Blanca, M.J. Diez, J.
Prohens, and B. Pico. 2012. Application of genomic tools in plant breeding. Curr.
Genomics 13:179–195. doi:10.2174/138920212800543084.

8. de los Campos, G., J. M. Hickey, R. Pong-Wong, H. D. Daetwyler, and M.P. L. Calus, 2013
Whole-genome regression and prediction methods applied to plant and animal breeding.
Genetics 193: 327–345.

9. Voss-Fels KP, Cooper M, Hayes BJ (2019) Accelerating crop genetic gains with genomic
selection. Theor Appl Genet 132:669–686.

10. Onogi A, Watanabe M, Mochizuki T, Hayashi T, Nakagawa H, Hasegawa T, Iwata H (2016)
Toward integration of genomic selection with crop modelling: the development of an
integrated approach to prediction rice heading dates. Theor Appl Genet 129:805–817.

11. Yin X, Kropff MJ, Aggarwal PK, Peng S, Horie T (1997) Optimal preflowering phenology
of irrigated rice for high yield potential in three Asian environments: a simulation study.
Field Crop Res 51:19–27.

12. Hori K, Matsubara K, Yano M (2016) Genetic control of flowering time in rice: integration
of Mendelian genetics and genomics. Theor Appl Genetics 129:2241–2252.

13. Onogi A, Ideta O, Inoshita Y, Ebana K, Yoshioka T, Yamasaki M, Iwata H (2015) Exploring
the areas of applicability of whole-genome prediction methods for Asian rice (Oryza
sativa L.). Theor Appl Genet 128:41–53.

14. Spindel J, Begum H, Akdemir D, Virk P, Collard B, Redoña, Atlin G, Jannink JL, McCouch
SR (2015) Genomic selection and association mapping in rice (Oryza sativa): effect of
trait genetic architecture, training population composition, marker number and statistical
model on accuracy of rice genomic selection in elite, tropical rice breeding lines. PLoS
Genetics 11: e1004982.

15. Li, X., Guo, T., Mu, Q., Li, X. & Yu, J. Genomic and environmental determinants and their
interplay underlying phenotypic plasticity. Proc. Natl Acad. Sci. USA 115, 6679–6684
(2018).

16. Forsythe, William C., Edward J. Rykiel Jr., Rand al S. Stahl, Hsin-i Wu and Robert M.
Schoolfield, 1995. A model comparison for daylength as a function of latitude and day of
the year. Ecological Modeling 80:87-95.

17. Yabe S, Yoshida H, Kajiya-Kanegae H, Yamasaki M, Iwata H, Ebana K, Hayashi T,
Nakagawa H. Description of grain weight distribution leading to genomic selection for
grain-filling characteristics in rice. PLoS One. 2018 Nov 20;13(11):e0207627. doi:
10.1371/journal.pone.0207627. eCollection 2018. PubMed PMID: 30458025; PubMed
Central PMCID: PMC6245794.

18. Murray, M. G., and Thompson, W. F. 1980, Rapid isolation of high molecular weight plant
DNA. Nucleic Acids Res., 8, 4321–5.

19. Bolger, A. M., Lohse, M., and Usadel, B. 2014, Trimmomatic: A flexible trimmer for
Illumina sequence data. Bioinformatics, 30, 2114–20.

20. Kawahara, Y., de la Bastide, M., Hamilton, J. P., et al. 2013, Improvement of the Oryza
sativa Nipponbare reference genome using next generation sequence and optical map
data. Rice, 6, 1–10.

21. Li, H. 2012, Exploring single-sample SNP and INDEL calling with whole-genome de novo
assembly. Bioinformatics, 28, 1838–44.

22. McKenna, A., Hanna, M., Banks, E., et al. 2010, The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. Genome
Res., 20, 1297–303.

23. Smithsonian Institution, 1939. Smithsonian Meteorological Tables. Lord Baltimore Press,
Baltimore, MD, pp. lxxvilxxviii, pp. 211-224.

24. Hijmans, R.J. (2017). geosphere: Spherical Trigonometry. R package version 1.5-5.
https://CRAN.R-project.org/package=geosphere

25. VanRaden, P.M. Efficient methods to compute genomic predictions. J. Dairy Sci. 2008, 91,
4414–4423.

26. Crossa, J., D. Jarquín, J. Franco, P. Pérez-Rodríguez, J. Burgueñoet al., 2016.Genomic
prediction of gene bank wheat landraces. G3 (Bethesda) 6:1819–1834.

27. Jarquín, D., C. Lemes da Silva, R.C. Gaynor, J. Poland, A. Fritz, R. Howard, S. Battenfield,
and J. Crossa. 2017. Increasing Genomic-Enabled Prediction Accuracy by Modeling
Genotype $\times$ Environment Interactions in Kansas Wheat. Plant Genome 10:0.
doi:10.3835/plantgenome2016.12.0130

28. R Core Team, R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing (2018), Vienna, Austria.

29. de los Campos, G., and P. Pérez-Rodríguez. 2013. BGLR: Bayesian generalized linear
regression. R package version 1(3).

30. Pérez-Rodríguez, P., J. Crossa, K. Bondalapati, G. De Meyer, F. Pita, and G.D.L. Campos.
2015. A pedigree-based reaction norm model for prediction of cotton yield in
multienvironment trials. Crop Sci. 55:1143–1151. doi:10.2135/cropsci2014.08.0577

Acknowledgements

598 We are grateful to Dr. Vikas Belamkar for their valuable feedback and intellectual contribution
599 during the elaboration of this manuscript.

Author Contributions

602 D.J. and H.I. conceptualized the study; H.K.K., S.Y., H.N., and M.Y. performed phenotyping;
603 H.K.K. and C.T. generated and processed molecular marker, environmental and day length data
604 sets; D.J., C.T., R.P. and H.I. analyzed the data; D.J., H.I., R.P., C.T., H.K.K and J.Y. prepared the
605 draft M.S. and all authors contributed to finalize; all authors have read and approved the M.S.

606 **Competing Interests:** The authors declare no competing interests.