Coupling Day Length Data and Genomic Prediction tools for Predicting Time-Related
Traits under Complex Scenarios

Diego Jarquin[*,1], Hiromi Kajiya-Kanegae[†], Chen Taishen[†], Shiori Yabe[‡], Reyna Persa[*], Jianming

Yu[**], Hiroshi Nakagawa[‡], Masanori Yamazaki[§], Hiroshi Iwata[†,1]

[*]Department of Agronomy and Horticulture, University of Nebraska – Lincoln, Lincoln NE,

68583, USA.

[†]Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo, Tokyo

113-8657, Japan.

[‡]Institute of Crop Sciences, National Agriculture and Food Research Organization (NARO),

Tsukuba, Ibaraki 305-8604, Japan

[**]Department of Agronomy, Iowa State University, Ames, USA.

[§]Food Resources Education and Research Center, Graduate School of Agricultural Science,

Kobe University, Kasai, Hyogo 675-2103, Japan

[‡]Institute for Agro-Environmental Sciences, National Agriculture and Food Research

Organization (NARO), Tsukuba, Ibaraki 305-8604, Japan

[1]Corresponding authors: University of Nebraska – Lincoln, Lincoln NE, 68583, USA. E-mail:
diego.jarquin@gmail.com ORCID ID 0000-0002-5098-2060 (DJ). Graduate School of
Agricultural and Life Sciences, The University of Tokyo, Bunkyo, Tokyo 113-8657, Japan. E-
mail: iwata@ut-biomet.org, ORCID 0000-0002-6747-7036 (HI).

ABSTRACT

Genomic selection (GS) has proven to be an efficient tool for predicting crop rank performance of untested genotypes via their maker profiles; however, when the trait of interest have intermediate optima, as those traits related to crucial phenology stages of plants, this implementation might not be the most convenient when phenotypic information of the target environment is not available. In these cases, GS might deliver high rank correlations between predicted and observed values but incurring in serious bias. Days to heading (DTH) is a crucial development stage in rice for regional adaptability with a significant impact in yield potential. For this reason, the information about DTH is important for designing and planning rice cultivars experiments. We developed two methods that incorporate day length information (DL) in the prediction process: C method, which models DL and DTH aiding the prediction of tested genotypes in unobserved environments; CB method, which combines the C method plus genomic BLUPs for predicting untested genotypes in unobserved environments. The use of DL has clear advantages over the use of weather data since this information can be determined in advance just by knowing the location and planting date for establishing future cultivars. In this research, we considered three different prediction scenarios of interest for breeders: (*i*) CV2, predicting tested genotypes in observed environments; (*ii*) CV0, predicting tested genotypes in unobserved environments; and (*iii*) CV00, predicting untested genotypes in unobserved environments. Under CV00 scenario, the proposed CB method returned a square root of the mean square error (SMSE) of 7.3 days, a Pearson correlation of 0.93 and the environmental mean differences (EMD) ranged between -6.4 and 4.1 days. The conventional GS method produced a SMSE of 18.1 days for a correlation of 0.41 and EMD ranging between -31.5 and 28.7 days. For CV0, the proposed C method returned a SMSE of 3.9 days for a correlation of 0.98 and EMD ranging between -4.95 and 4.67 days. The results from the CV2 scheme were used for comparison purposes to assess the predictability levels that can be reached when complete information on genotypes and environments are available. Here, the conventional GS implementation produced a SMSE of 3.8 days for a correlation of 0.98 and an EMD ranging between -0.02 and 0.03 days. The results obtained with the proposed methods are promising and showed that the efficient use of DL information aids to improve significantly the predictive ability of time-related traits.

Key words: Time-related traits; days to heading; day length; genome-enabled prediction.

INTRODUCTION

To satisfy the daily demand of crop consumption for a growing population, it is necessary to improve the production systems to allow a sustainable increase of the yield potential and nutritional value of the new and up-coming varieties (FAO et al., 2018). In plant breeding there are different tools and methodologies to increase the rate of genetic gain (Bernardo 2002). Traditional plant breeding methods use phenotypic and/or pedigree information to perform selection of experimental lines for developing superior cultivars. However, phenotypic selection may not be permissible in many situations due to the high phenotyping costs; also, pedigree selection may not be accurate since recombination effects are not accounted for under this method.

Genomic selection (GS, Meuwissen et al., 2001) is an emerging tool that allows screening genotypes from a very large population without having to observe them in fields (Heffner et al., 2009; Jannink et al., 2010). This method only requires phenotypic and genomic information for calibrating models, then other genotyped candidates are selected based on the predicted values using their marker profiles (Crossa et al., 2010). In this context, the implementation of genomic tools and resources translates our understanding of the relationship between genotype and phenotype into predicted genetic merits for direct selection. This is highly desirable dealing with those traits that are controlled by a large number of genes with small individual effects, also known as complex traits (Jannink et al., 2010, Perez-de-Castro et al., 2012). GS has shown to be an effective tool for plant and animal breeding applications, especially when dealing with complex traits (de los Campos et al., 2013). The successfulness of GS predicting phenotypes from genotypic information has been reported in various crop species as reviewed by Voss-Fels et al. (2019).

In GS, the estimated breeding values (EBVs) of complex traits of untested genotypes are computed based on genome-wide marker information. The EBVs can be considered as deviations around a fixed mean (e.g., zero). Once the EBVs are obtained, the breeders may perform selections based on the ranked values by selecting those individuals in the tails. Depending on the trait, the individuals are selected either on the upper or lower tails of the ranked EBVs. For example, the upper side is selected for increasing genetic gains of traits like grain yield, stress resistance, or the content of essential nutrients. On the other hand, genotypes in the lower tail are selected for reducing lodging, shattering, the content of anti-nutrients, etc. In these cases, the EBVs provide sufficient information when the correlation between the EBVs and the direct genetic values (or phenotypic values) is high. However, the rank-based selection criteria might not be the most accurate when the target trait has an intermediate optima. This is the case of those traits related to phenological stages of crop plants. For example, days to heading (DTH) is one of the critical traits of ideotypes with an increased yield potential, and thus it is important to optimize DTH for high yield potential in different environments (Yin et al., 1997).

Usually multi environment trials (METs) are performed for developing stable cultivars
across a wide range of environmental conditions, and the breeders may be interested in
estimating DTH for a large set of environments instead of a single one. However, the EBVs may
contain a large bias if no phenotypic information from a targeted unobserved environment is
available. The EBVs of one environment may not be valid for another environment due to
variations in the environmental *stimuli*. These differences remain also for same genotypes
observed in same locations but at different times (planting date, years). As consequence of the
aforementioned issues, the problem of correctly estimating DTH for a set of untested genotypes
in a wide range of unobserved environments becomes harder to solve than for non-time related
traits. As pointed before, the advantage of GS techniques over conventional phenotypic methods
is its ability for performing accurate rank predictions of untested genotypes based exclusively on
their marker profiles without having to observe these in fields. Thus, a method capable of
delivering highly accurate predictions of untested genotypes in unobserved environments without
incurring in a large bias is desirable.

In this study, we propose a novel method to predict DTH in a precise and accurate way
(i.e., small bias and high correlation between predicted and observed values). We demonstrated
the potential of the method, analyzing a rice DTH dataset comprised of 112 genotypes evaluated
in 51 environments. Because rice DTH is important for regional adaptability and has usually
great impact on yield potential (Yin et al., 1997), genetic systems controlling DTH have been
studied for a long time by integrating Mendelian genetics and genomics (Hori et al., 2016). The
potential of GS in DTH has been evaluated in rice (Spindel et al., 2015 and Onogi et al., 2015)
and sorghum (Li et al., 2018). Onogi et al. (2016) focused on the genomic prediction of DTH in
multiple environments and demonstrated the potential of a novel method that integrates GS with
a crop growth model. Li et al. (2018) also focused on the genomic prediction of DTH in multiple
environments and demonstrated the potential of models combining GS with joint regression
analysis on photothermal time. In this study, we also focus on the accurate prediction of DTH
because its optimization is indispensable for high yield potential in different environments (Yin
et al., 1997). The optimization will become more important in near future to attain high and
stable yield under the changing climate conditions.

Firstly, we explored the usefulness of cumulative day length (CDL) information (planting
date until heading) for predicting DTH. We found that CDL produced accurate predictions with
small to null bias. However, for this method to work it is necessary know first the exact number
of days that DL will accumulate from planting until heading and these values correspond to the
response we want predict complicating the process.

An alternative way of using DL information is considering the DL values of the day
when DTH are reached. In principle, DTH information of genotypes observed in different
environments may be used for finding patterns that relate DTH and DL. Apparently, each
genotype poses a unique fingerprint that relates the DL values at the occurrence day with DTH
across environments. These patterns are characterized via a third degree polynomial equation
that depends exclusively on the time covariate. This fingerprint provides a rough estimation of
the DTH potential for a target candidate environment (location $\times$ planting-date combination).

The ultimate goal of this research was to develop a method for predicting DTH of untested
genotypes in unobserved environments. For this, a third degree polynomial equation was used
for characterizing the DL at the DTH occurrence across environments of tested genotypes (one at
the time). Then, the DL values of a target unobserved environment were fitted via a third degree
polynomial equation. An estimation of DTH of a tested genotype in an unobserved environment
can be obtained by finding the intersection of these two equations.

This procedure was repeated for each genotype (one at a time) yet to be observed in the
candidate environment. As was pointed before, one of the disadvantages of using GS methods
for predicting DTH in unobserved environments is that a large bias between predicted and
observed values may be generated if no phenotypic information of the candidate environment is
available. In this case, the DTH environmental mean of the target environment is determined as
the weighted mean of those environments in the training set. Thus, the genetic components
become just random deviations around this weighted mean. If the real mean of this candidate
environment is far from the weighted mean, a large bias may be expected predicting DTH, yet a
high correlation can be obtained between the random deviations and the observed DTH values.

Thus, the problem of obtaining reliable predictions of untested genotypes in an
unobserved environment may be solved just by finding the accurate DTH mean and then
combine it with the genomic BLUPs obtained with GS methods. The BLUPs may be added just
as deviations around this estimated DTH environmental mean. For assessing the proposed
method, the square root of the mean square error (SMSE) and the Pearson correlation between
predicted and observed values were computed within environments.

Three cases that are of interest for designing and planning experiments for addressing
different problems were considered. CV2, predicting tested genotypes in observed environments
(incomplete field trials); CV0, predicting tested genotypes in unobserved environments
(targeting environments); and CV00, predicting untested genotypes in unobserved environments
(targeting genotypes and environments).

Under the hardest cross-validation scheme (CV00), the results of the proposed method
(CB) were compared with those obtained using the conventional GS model (E+G). For studying
other applications for realistic scenarios, the two remaining cross validation schemes were also
considered (CV2 and CV0). Results from CV2 scheme using the conventional GS model (E+G)
were used to contrast the performance of the proposed method when complete information on
genotypes and environments are available. Thus, these results may provide a better idea of the
model performance of the proposed method when no information of genotypes and environments
are available. Results under the CV0 scheme correspond to the proposed C method. Here the
environments are considered "target" in a strict sense because based on theoretical DL values the
best location and planting date can be determined before conducting field experiments.

Across environments, under the CV2 scheme the conventional GS model (E+G) returned
a SMSE of 3.8 days and a Pearson correlation of 0.98. With CV00, the proposed BC method
gave a SMSE of 7.2 days and a Pearson correlation of 0.93; also, the difference between the
predicted and observed environmental means (DM) ranged between -4.1 and 6.4 days. On the

other hand, the conventional GS model produced a SMSE of 18.1 days and a Pearson correlation
of 0.41; the DM ranged between -28.73 and 31.53 days. Finally, the C method produced accurate
results predicting tested genotypes in untested environments (CV0 scheme). It returned a SMSE
of 3.8 days and a Pearson correlation of 0.98; the DM ranged between -4.67 and 4.95 days. This
method does not require genomic information at all neither phenotypic information from other
genotypes.

193   **MATERIALS AND METHODS**

Phenotypic data

The phenotypic data set comprises the seeding and heading date of 112 Japan rice cultivars
tested in multiple locations from 2004 to 2017. This multi-environmental experiment was
established to investigate heading related genes. During the first 2 years (2004 and 2005), the
experiments were conducted in Tsukubamirai for 2 different cultivation periods. Then gradually
more locations were considered for testing DTH, and these are distributed from North to South
of Japan. Since 2012 at least 7 experiment per year have been conducted until 2018. Originally,
the total number of trials/experiments (location-by-year-by-planting date combinations) is 79
(see Table A1); however, in this research we used information of only 51 of these (see Table 1).

The DTH was calculated as the difference between the heading date and the seeding date.
The heading date was determined when more than 50% of the individuals reached the heading
stage. However, DTH was measured in a different way in Kasai location. There, the heading date
was initially determined as the event when 10% of the individuals started heading. Then,
following the suggestions of an expert rice scientist, the final DTH for the genotypes tested in
this location were obtained by adding 2 days to those cultivars from Hokkaido and 1 day to the
cultivars originally developed in other locations. The replicated number of plants evaluated for
each cultivar is different by experiments, and these ranged from 7 to 30. The cultivars that did
not reach the heading stage at the end of the experiment were removed from the analysis.

Marker genotype data

For each variety, total DNA was extracted using the CTAB method (Murray et al., 1980). Next-
generation sequencing data was obtained using the Illumina HiSeq 2000, HiSeq 4000 and
HiseqX systems via paired-end sequencing (Illumina, Inc., San Diego, CA, USA). Data sets
deposited in the DDBJ Sequence Read Archive (SRA106223, ERA358140, DRA000158,
DRA000307, DRA000897, DRA000927, DRA007273, DRA007256) were reanalyzed. Adapters
and low-quality bases were removed from paired reads using the Trimmomatic v0.36 program
(Bolger et al., 2014). The preprocessed reads were aligned on the Os-Nipponbare-Reference-
IRGSP-1.0 (Kawahara et al., 2013) by using the bwa-0.7.12 mem algorithm with the default
options (Li, 2012).
SNP calling was based on alignment using the Genome Analysis Toolkit (GATK, 3.7-
0-gcfedb67) (McKena et al., 2010) and Picard package V2.5.0
(http://broadinstitute.github.io/picard). The mapped reads were realigned using

RealignerTargetCreator and indelRealigner of GATK software. SNPs and InDel were called at
the population level using the UnifiedGenotyper of GATK with the -glm BOTH option.
Heterozygous loci were treated as missing, owing to the assumption that almost all markers were
well fixed in rice cultivars after several generations of self-pollination. We used only biallelic
non-missing sites over all cultivars with a minor allele frequency (MAF) $\geq$ 0.025. For further
analysis, genotypes were represented as -1 (Homozygous Reference) and 1 (Homozygous
Alternate).

Day length data

The original data set also consider environmental data (ED) and it was downloaded from the
Agro-Meteorological Grid Squared Data (NIAES; https://amu.rd.naro.go.jp). However, in this
study we only day length DL information was considered for developing the proposed model.
Further developments will integrated both types of layers (DL & ED) in the analysis. The
theoretical DL values were computed by the default method (Forsythe et al., 1995) included in
the geosphere (v1.5-5) R package (Hijmans., 2017).

GBLUP model.

The GBLUP (Genomic BLUP) model is a main effects model used in GS applications
(vanRaden et al., 2007, de los Campos et al., 2013). It attempts to explain the DTH value of the
ith genotype (i=1, 2,…, I) observed in the jth environment (j=1, 2, …, J) as the sum of a common
mean, random environmental and genomic effects, and an error term. It can be represented as
follows:

$$Y_{ij} = \mu + E_j + g_i + \varepsilon_{ij} \qquad [1]$$

where μ is a common effect to all genotypes across all environments; E_j are IID (independent
and identically distributed) outcomes from a $N(0, \sigma_E^2)$ representing the effect of the jth
environment; $\boldsymbol{g} = \{g_i\} \sim N\left(\boldsymbol{0}, \mathbf{G}\sigma_g^2\right)$ is the vector of genomic effects, with $\mathbf{G} = \frac{\mathbf{XX'}}{p}$ as the
genomic relationship matrix whose entries describe the genetic similarities between pairs of
individuals, $\mathbf{X}$ is the standardized (by columns) matrix for p molecular markers; ε_{ij} are IID
outcomes from a $N(0, \sigma_\varepsilon^2)$ representing measure errors; and σ_E^2, σ_g^2 and σ_ε^2 are the corresponding
associated variance components of the described random effects.

Usually, when METs are analyzed the environmental component explains the larger
proportion of the phenotypic variability across environments. For example, Crossa et al. 2016
and Jarquin et al., 2017 showed that this component captured between 65% and 95% of the
variability in grain yield for different wheat populations (Mexican, Iranian and hard red winter).
As consequence of this, from [1] we can conclude that the predicted value of the ith genotype in
the jth environment strongly depends on an environmental mean, which is composed by a

common mean across environments plus an environmental deviation that corresponds to the
effect of j^{th} environment.

Conceptually, the common mean effect can be viewed as the average performance of all
genotypes across environments. The environmental deviation corresponds to the mean of the
specific affectations of the current environmental *stimuli* in the set of the tested genotypes
beyond the common mean. Thus, these environmental effects can be considered deviations
around the common mean. Hence, by construction the contribution of the genomic component to
the idealized phenotype is much smaller in comparison with the overall mean and the
environmental effects.

If phenotypic information of the target environment is available, good approximations of
the environmental effects can be computed. When this information is not available, the
environmental effects are estimated using genotypes observed (phenotype) in other
environments. In this case, the environmental component is estimated as the weighted average of
the environmental effects in the training set. Since these random effects are conceptualized to be
centered on zero, the weighted average will be close to zero. Hence, the predicted DTH value
will be compose mainly of a common mean and a small genomic effect shrinking the predicted
values towards the common mean. Dealing with traits where the selection process is based on
rank performance (yield), the selected genotypes would correspond to those with the
largest/smallest values (depending on the desired direction of the trait to improve) without
considering the estimated values *per se*.

However, for those traits linked to important physiological crop growth stages (e.g.,
DTH) the selection based on rakings might not be the best strategy. Although this method
usually delivers high correlations between predicted and observed values, it also might induce a
large bias (MSE) when no information of the target environment is included. For solving this
issue, we developed a method that incorporates DL information in the prediction process.

C method

The idea for using DL information in the prediction process was motivated examining the
relationship between CDL and DTH of all genotypes across environments. A visual assessment
of this relationship can be done with Figure A1. This plot contains 94% of the all possible
combinations between 112 genotypes and 51 environments. Here, a simple linear regression
returned an R-squared value of 0.99 exhibiting a strong relationship between DTH and CDL. It
can be concluded that if CDL values are known then accurate estimations of DTH can be
obtained for any genotype in any environment. However, a recursive issue is faced because in
order to compute CDL the number of days between planting date and the heading occurrence
should be known beforehand. Hence, DTH should be known first before compute CDL leading
this to a recursive problem. However, analyzing the DL values of the day when a single genotype
reaches the heading stage across environments could lead us to recognize clear patterns relating
DTH and DL. Under this assumption, DTH becomes a function that depends on DL; however for
a better understanding of the concept this relationship can be switched (i.e., DL depending on
DTH).

For illustrating this concept, Figure 1 is used as example. It corresponds to genotype 56
observed in 50 environments out of 51 (1-46, and 48-51 but not in environment 47). The goal
here is to estimate DTH for this genotype in the remaining unobserved environment
(environment 47: Tsukubamirai 2016 Late). For each environment, the planting date was
considered as day 1 (*x-axis*). Then for these environments, the daily DL (*y-axis*) values were
plotted (pink lines) from day 1 (planting) until genotype 56's DTH (blue point). Examining the
cloud of blue points, a pattern can be identified where DL at the occurrence day depends at some
grade on DTH. This gradient seems not to be strongly affected by the initial DL values at the
different planting dates across environments. However, it appears to be linked with the initial
DL's changing rate at the planting day. Those environments where DL's changing rates
decreases at the planting time (late cultivars) showed shortest DTH values. On the other hand,
the environments with increasing DL's rates (early cultivars) presented the largest occurrence
times.

CB method

For predicting untested genotypes in unobserved environments (CV00) under the
aforementioned concept (i.e., using DL information), the proposed method requires: (*i*) the
estimation of DTH for a set of tested (training) genotypes (one at the time) yet to be observed in
a target/unobserved environment (CV0); (*ii*) to compute the expected environmental mean of the
unobserved environment using the estimated values from (*i*); (*iii*) to add the resulting blups of
the untested genotypes derived from conventional GS methods to the estimated environmental
mean. Further details of these steps are given next.

The DTH estimation of a tested genotype in an unobserved environment (CV0) requires
the phenotypic information across several environments and their corresponding DL values at the
occurrence time. The first step considers the characterization of DL at the heading occurrence
time across environments. For this, a third degree polynomial equation is used for fitting DL as
function of DTH (orange curve in Figure 1) as follows:
$dl_{ij} = b_0 + b_1 t + b_2 t^2 + b_3 t^3 + e_{ij}$, where dl_{ij} represents the observed DL of the i^{th} genotype in
the j^{th} environment; t is the time independent variable; b_0, b_1, b_2 and b_3 are the model parameters;
and e_{ij} is the error term and it is assumed to be IID $N(0, \sigma^2)$. For facility we will refer to this
model as C_1 curve.

In a similar way, the daily DL information of the target environment (green curve) can be
modeled using a third degree polynomial equation. The resulting fitted model is referred as C_2
curve. The solution to this system of two equations (C_1, C_2) will produce an estimate of DTH of
the tested genotype yet to bet observed in the target environment. In order to find the intersection
of these two curves, deterministic and numerical methods can be used. For facility in this study,
the numerical evaluation method was considered as follows: (*i*) the fitted model C_1 (orange
curve) was evaluated within the range of values of those phenotypes (DTH) of the same
genotype tested in other environments. For example, in Figure 1 the smallest and the largest
DTH values for genotype 56 were 67 and 137 days. Thus, the model was evaluated for the values
in the range [67, 137] in steps of 1 day. Similarly, for the fitted model of the target environment
it was evaluated in the same range in steps of 1 day. Then, the element to element differences
between the two resulting vectors were computed and the label [67-137] of the position with the

closets value to zero was considered as the intersection point (day 93 in Figure 1; dotted gray
line).
348

349
Figure 1. Graphical representation of the curve's C method for estimating days to heading (DTH) of genotype 56 in
environment 47 (Tsukubamirai 2016 Late). Pink curves represent daily day length (DL) values from planting day
(day 1) until heading time for genotype 56 in 50 environments (1-46, and 48-51). The blue points indicate the
corresponding DL values when genotype 56 reached heading time. The green curve (C_2) represents the target
environment (Environment 47) where no genotypes have been observed yet. The orange curve (C_1) depicts the fitted
line for genotype 56 using a third degree polynomial equation relating DL at the occurrence time and DTH across
environments. The vertical dotted gray line shows the intersection (93 days) between the green and the orange
curves (C_1, C_2). This value is the estimated DTH for genotype 56 in the unobserved environment "Tsukubamirai
2016 Late". The vertical dotted black line corresponds to the observed DTH (90 days) in environment 47.
359

Empirical assessment

For testing the accuracy of this method for predicting DTH for a set of tested genotypes yet to be
observed in a target environment, the C method was implemented considering one genotype at
the time. To evaluate the performance of this method across environments, the previous
procedure was implemented on each one of the 51 environment. The described method mimics
the scenario of predict DTH for tested genotypes in unobserved environments. This scenario is
similar to the cross-validation scheme CV0 (Jarquin et al. 2017) which attempts to predict
already tested genotypes in unobserved/target environments by combining phenotypic and
genomic information from the same genotype and from other genotypes observed in other
environments. In this case, only the phenotypic information of the same genotype observed in
other environments is used. Thus, the proposed C method does not considers genomic
information, neither phenotypic information from other genotypes.

Since the ultimate goal is to predict DTH for untested genotypes in unobserved
environments, the first step is to accurately estimate the expected DTH mean of the target
environment. Then, the final predicted value will be composed of the expected DTH mean of the
target environment (computed ignoring any phenotypic record of the untested genotype) plus the
genomic estimated breeding value ($\hat{g}_i$) of the untested genotype obtained via conventional GS
methods (E+G model). For estimating the DTH mean, the C method was applied first to the
genotypes in the training set to be observed in the target environment (one at the time). Then the
environmental DTH mean was computed by averaging these predicted values. So far, we have
not include any information (genomic and phenotype) from the untested genotype in the analysis.

Using a toy example consisting of four genotypes (G_1, G_2, G_3, and G_4) and three
environments (E_1, E_2, and E_3), in Figure 2 we describe the procedure for finding the
environmental mean of the target environment for untested genotypes. The top panel contains the
observed values; the middle panel shows the predicted values presumably were obtained with the
C method; and the bottom panel displays the conditional (given the tested genotypes) estimated
DTH environmental means for each genotype at each environment. Once the DTH means are
computed for each genotype in the candidate environment the prediction will be composed of
these means plus the genomic effects obtained using model [1] under CV00 scheme. These
genomic effects are estimated by deleting from the training set the information of the target
environment and the information from the same genotype observed in other environments.
Hence, within each environment the procedure of leaving out one genotype at the time was
applied.

		G_1	G_2	G_3	G_4	Mean
Observed	E_1	65	69	71	66	67.75
	E_2	80	85	87	90	85.50
	E_3	100	105	108	110	105.75
Predicted	E_1	68	69	73	70	70.00
	E_2	77	83	85	88	83.25
	E_3	104	103	110	114	107.75
E-Mean	E_1	70.7	70.3	69	70	69.98
	E_2	85.3	83.3	82.7	81.7	83.24
	E_3	109	109	107	106	107.65

394
Figure 2. Representation of the procedure for estimating the conditional DTH environmental mean for untested
genotypes in unobserved environments. The top panel contains the observed values for four genotypes (G_1, G_2, G_3
and G_4) observed in three environments (E_1, E_2, and E_3). The middle panel contains the predicted values presumably
obtained with the C method assuming that these genotypes were not tested in a particular environment. This method
attempts predicts DTH of tested genotypes in unobserved environments using the phenotypic information of these
genotypes but observed in other environments (one genotype at the time). For example, the predicted value for G_2 in
E_1 is 69 and it was computed using the information of G_2 tested in E_2 (85) and E_3 (105) via the C method. The

bottom panel contains the estimated DTH means (E-Mean). These are particular to each genotype and were compute
as the mean of the predicted values of those genotypes in the training set for the current environment. For example,
G_2 in E_1 (70.3) was computed as the mean of the predicted values of G_1 (68), G_3 (73) and G_4 (70) in E_1.
405

RESULTS

Descriptive statistics

Box-plot of DTH for each environment are depicted in Figure A2. The environments were
ordered based on the environmental medians. In general, those environments with late planting
date appeared first. The medians ranged between 59 (Fukuyama 2010 Late) and 116 (Akita
2015) days. The SDs of the environments varied between 5.07 and 16.83 days. Also, the smallest
and the largest DTH values were 42 and 148 days and these were observed in "Tsukubamirai
2016 Late" and "Akita 2015" environments, respectively.

Evaluating the bias and accuracy of the methods

Table 1 contains the observed and the predicted environmental means (E-mean) for four
different strategies (cross-validation scheme-method combination) as well as the corresponding
square root of the mean squared error (SMSE) and the Pearson correlation between the predicted
and observed values. These different strategies were used for different purposes. CB and GS
methods were implemented under the CV00 scheme (CV00_CB, CV00_GS) for comparing the
model performance of the proposed model with the conventional implementation (GS) predicting
untested genotypes in unobserved environments. C method was implemented under the CV0
scheme (CV0_C) for assessing the predictive ability of the C method predicting tested genotypes
in untested environments (also this is the intermediate step of the CB method used in the
previous scheme). Lastly, the conventional method (GS) was implemented under the CV2
scheme (CV2_GS) for providing an idea of the levels of predictability that can be reached when
complete information of genotypes and environments are available.

CV2 is the scenario that includes the most complete sets of information on genotypes and
environments for calibrating models. Across environments, the CV2_GS strategy returned a
SMSE of 3.8 days and a Pearson correlation of 0.98; within environments, the average SMSE
and Pearson correlation values were of 3.7 days and 0.97, respectively. The difference of the
environmental means (EMD) between predicted and observed DTH values ranged between -0.02
and 0.03 days. Since this scenario uses the most complete sets of information for calibrating
models, these results can be used as reference for assessing the performance of the proposed
methods under more complex prediction scenarios.

CV0_C results provide information of the ability for predicting tested genotypes in
unobserved environment when considering only phenotypic and DL information of the same
genotype using the C method. Across environments, this method respectively returned a SMSE
of 3.9 days and a Pearson correlation of 0.98; within environments, the average SMSE and
Pearson correlation were of 3.7 days and 0.98. The EMD ranged between -4.95 and 4.67 days.

CV00 corresponds to the scenario of predicting untested genotypes in unobserved
environments and it is considered the hardest and most interesting prediction problem because no
phenotypic information is available for connecting untested genotypes in unobserved
environments with calibration sets. Across environments, the conventional GS method
(CV00_GS) respectively returned a SMSE and a Pearson correlation of 18.1 days and 0.41;
within environments, the average SMSE and Pearson correlation were of 17.5 days and 0.69. The
EMD ranged between -31.5 and 28.7 days. For the same scenario, across environments the
proposed CB method (CV00_BC) respectively returned a SMSE and a Pearson correlation of 7.3
days and 0.93; within environments, the average SMSE and Pearson correlation were 7.2 days
and 0.86. The EMD ranged between -6.4 and 4.1 days.

Table 1. Environmental means, Square root of the Mean Squared Error (SMSE) and the Pearson correlation between
predicted and observed days to heading (DTH) for 112 rice genotypes tested in 51 environments. Three cross-
validation schemes were implemented for mimicking realistic prediction scenarios. CV2 resembles incomplete field
trails were some lines are tested in some environments but not in others. CV0 represents the scenario of predicting
tested genotypes in unobserved environments. CV00 considers the prediction of untested genotypes in unobserved
environments. CV2 was implemented using the conventional implementation (GS, E+G) leaving one phenotype out
at the time. For CV0 the C method was used combining phenotypic and day length information (DL) of each
genotype (one at the time) already tested in other environments. For CV00 two methods were considered: (*i*) the
conventional implementation (GS, E+G) leaving one genotype out across environments and deleting any phenotypic
information from the target environment; and (*ii*) the proposed CB method, which combines phenotypic and DL
information from the tested genotypes (training set) with the genomic blup values ($\hat{g}$) of the untested genotypes
obtained in (*i*).

	Mean					Square root of the Mean Square Error				Correlation			
	Mean	CV00_CB	CV00_GS	CV0_C	CV2_GS	CV00_CB	CV00_GS	CV0_C	CV2_GS	CV00_CB	CV00_GS	CV0_C	CV2_GS
Tsukubamirai 2004 Late	79.4	76.7	85.9	76.3	79.4	7.0	12.6	4.6	2.5	0.86	0.70	0.99	0.98
Tsukubamirai 2004 Early	87.5	93.2	84.1	92.2	87.5	10.0	12.9	6.4	4.2	0.84	0.55	0.97	0.97
Tsukubamirai 2005 Early	98.5	102.1	87.1	101.0	98.5	6.9	14.0	3.5	2.0	0.87	0.75	0.98	0.99
Tsukubamirai 2005 Late	76.7	74.8	84.8	73.7	76.7	5.7	12.4	3.4	2.2	0.86	0.68	0.99	0.99
Kasai 2006	87.4	83.8	88.3	82.8	87.4	7.9	8.8	6.8	4.7	0.85	0.75	0.93	0.94
Fukuyama 2006	80.6	83.0	82.5	82.0	80.6	5.6	13.4	2.0	2.2	0.87	0.29	0.99	0.99
Fukuyama 2007	84.1	83.4	86.6	82.5	84.1	5.0	6.1	2.9	3.0	0.87	0.84	0.98	0.98
Tsukuba 2008	104.5	107.7	87.2	107.3	104.5	9.1	19.2	4.0	4.7	0.87	0.85	0.98	0.98
Kasai1st 2008	102.2	99.9	88.7	99.2	102.2	9.0	17.8	4.3	4.8	0.89	0.71	0.99	0.99
Fukuyama 2008	77.8	82.2	85.6	81.3	77.8	7.8	11.1	4.5	2.8	0.86	0.77	0.98	0.98
Tsukuba 2009	103.9	107.7	87.1	108.0	103.9	8.6	19.0	5.2	4.7	0.89	0.81	0.98	0.97
Kasai1st 2009	101.9	100.6	89.4	100.0	101.9	7.6	15.1	2.7	3.5	0.89	0.83	0.99	0.99
Fukuyama 2009	78.7	78.7	89.6	77.8	78.7	5.2	13.2	1.5	2.2	0.86	0.83	0.99	0.99
Tsukuba 2010 late	83.5	89.1	88.9	88.4	83.5	8.4	11.1	5.9	3.0	0.86	0.66	0.97	0.97
Kasai1st 2010	100.6	100.7	89.5	100.1	100.6	7.1	14.3	1.9	2.9	0.88	0.77	0.99	0.99
Fukuyama 2010	76.2	79.5	88.2	78.1	76.2	5.9	15.3	2.6	2.8	0.86	0.50	0.99	0.99
Fukuyama 2010 Late	59.0	65.4	87.7	62.5	59.0	9.2	29.7	4.2	7.7	0.84	0.82	0.96	0.90
Tsukuba 2010 Early	105.9	109.9	85.5	109.9	105.9	9.9	23.9	5.3	5.6	0.87	0.66	0.98	0.97
Tsukuba 2010 Late	82.8	80.8	90.0	80.4	82.8	5.9	11.9	4.1	3.6	0.86	0.78	0.95	0.96
Tsukuba 2010 Middle	94.7	95.7	90.9	95.4	94.7	6.5	11.1	3.0	3.1	0.88	0.67	0.98	0.98
Kasai 2011	101.4	100.6	89.3	100.0	101.3	7.8	15.7	2.4	3.6	0.89	0.76	0.99	0.99
Fukuyama 2011	75.7	79.4	91.1	78.6	75.7	6.4	16.9	3.1	2.0	0.87	0.83	0.99	0.99
Fukuyama 2011 Late	66.4	72.7	97.9	66.2	66.4	8.9	32.4	2.9	5.7	0.68	0.58	0.95	0.86
Tsukuba 2012 Early	97.1	96.0	86.9	101.1	97.1	5.7	12.4	5.8	3.2	0.81	0.74	0.97	0.94
Tsukuba 2012 Late	84.3	81.6	88.2	81.0	84.3	6.1	7.5	4.3	3.5	0.86	0.80	0.96	0.97
Tsukuba 2012 Middle	92.6	95.4	88.4	94.8	92.6	6.3	8.3	3.7	2.9	0.87	0.82	0.97	0.97
Kasai 2012	103.8	102.7	86.4	102.1	103.8	7.9	21.3	2.7	3.8	0.89	0.60	0.99	0.99
Fukuyama 2012	75.2	78.8	89.9	78.0	75.2	6.2	16.3	3.1	2.0	0.87	0.83	0.99	0.99
Fukuyama 2012 Late	71.5	77.0	103.1	68.6	71.6	7.5	32.4	3.4	4.6	0.70	0.66	0.98	0.92
Fukuoka 2012	79.6	76.0	89.1	75.4	79.6	6.7	13.0	5.2	4.2	0.84	0.69	0.96	0.96
Akita 2012	113.1	115.9	90.1	114.5	113.1	8.3	26.1	2.8	4.4	0.86	0.54	0.98	0.97
Kasai 2013	104.8	104.1	87.3	103.5	104.8	8.0	18.0	3.2	4.7	0.90	0.80	0.98	0.98
Fukuyama 2013	74.9	80.1	89.4	79.3	74.9	7.4	16.3	4.6	1.8	0.87	0.82	0.99	0.99
Fukuyama 2013 Late	62.1	64.8	85.7	64.1	62.1	6.3	27.1	3.6	4.9	0.83	0.36	0.93	0.95
Fukuoka 2013	77.4	77.3	93.3	76.7	77.4	5.7	17.5	1.9	2.6	0.86	0.77	0.99	0.98
Akita 2013	116.6	115.8	91.1	115.1	116.5	8.5	28.1	2.4	4.9	0.88	0.66	0.99	0.98
Kasai 2014	102.4	105.1	88.9	104.4	102.3	8.4	18.0	2.8	4.4	0.90	0.66	0.99	0.99
Fukuyama 2014	79.6	82.1	87.9	81.3	79.6	5.9	12.1	2.1	1.3	0.88	0.68	0.99	0.99
Fukuyama 2014 Late	67.6	64.5	88.1	62.9	67.6	6.7	21.4	5.1	6.0	0.83	0.79	0.98	0.94
Fukuoka 2014	81.8	77.7	80.4	77.1	81.8	6.7	11.5	5.4	3.1	0.87	0.45	0.97	0.97
Akita 2014	115.2	116.2	89.6	115.6	115.2	9.2	28.5	2.2	5.8	0.88	0.68	0.99	0.98
Fukuyama 2015	76.3	72.6	88.1	71.9	76.3	6.5	14.4	4.6	3.3	0.86	0.63	0.99	0.98
Akita 2015	118.1	119.8	89.4	117.6	118.0	7.7	29.9	2.5	3.6	0.81	0.76	0.99	0.98
Kasai 2016	102.3	101.2	85.6	100.7	102.2	7.2	20.4	3.1	3.9	0.90	0.61	0.98	0.98
Fukuyama 2016	79.1	81.1	85.2	80.8	79.1	6.0	8.9	2.4	2.9	0.86	0.82	0.99	0.98
Fukuyama 2016 Late	67.5	65.2	89.6	64.6	67.5	6.8	26.1	4.5	6.3	0.79	0.63	0.95	0.91
Tsukubamirai 2016 Late	73.6	77.9	84.6	77.2	73.6	7.0	13.6	4.9	1.9	0.90	0.80	0.98	0.99
Kasai 2017	108.6	106.2	85.7	105.5	108.6	8.3	26.3	4.3	4.4	0.89	0.56	0.98	0.98
Fukuyama 2017	77.4	80.2	87.7	79.6	77.4	6.3	16.3	2.7	2.0	0.88	0.42	0.99	0.99
Fukuyama 2017 Late	68.4	67.7	88.8	67.0	68.4	6.0	22.9	3.0	4.9	0.82	0.69	0.97	0.95
Tsukubamirai 2017 Late	69.5	73.1	86.9	72.2	69.5	6.8	19.8	4.3	1.6	0.87	0.65	0.98	0.99
Mean						7.2	17.5	3.7	3.7	0.86	0.69	0.98	0.97

Only the results of CB and GS methods used under CV00 scheme are directly comparable between them. Figure 3 depicts the scatter plot between predicted and observed DTH values for all genotypes across environments. The top panel contains the results obtained with the proposed CB method, while the bottom panel shows the results derived from the conventional implementation (GS, E+G). The blue and black lines correspond to the fitted and the on target predictions with respect to the observed DTH values. In general, across and within environments (Table 1) the results obtained with CB method showed a better fit than those derived from the GS model.

472
Figure 3. Scatter plot of predicted (*x-axis*) and observed (*y-axis*) DTH values using two different methods (CB a and
GS) for predicting untested genotypes in unobserved environments (CV00 scheme) for a rice data set comprising
112 genotypes tested in 51 environment in Japan between 2005 and 2017. The top panel presents the results

obtained with the proposed CB method. It combines the predicted environmental means (E-mean) obtained using the
proposed C-method with the genomic blups ($\hat{g}$) derived from the conventional implementation (GS, E+E) under the
CV00 scheme. The bottom panel displays the results obtained with the conventional implementation (GS, E+G GS).
The blue line represents the fitted line between predicted and observed DTH values while the black line show the
exact match (on target) of the predicted values on the observed values.
481
The advantages of considering the proposed CB method for predicting DTH under the
CV00 scheme are more clearly appreciated analyzing the predicted environmental means (E-
means). Figure 4 displays the predicted (*x-axis*) and observed (*y-axis*) E-means. The predicted E-
means obtained with the CB and GS methods are indicated in orange and blue colors,
respectively. As expected, when no information from the target environment was available for
calibrating models the conventional implementation (GS, E+G) regresses the predicted values
towards the overall mean (~87.74) from the training sets. On the other hand, the CB method
consistently returned closer values to the diagonal black line, which represents the exact match.

490
Figure 4. Scatter plot of predicted and observed environmental means (E-means) using the proposed CB method
(orange) and the conventional implementation (GS, E+G) (blue) under the CV00 scheme (predicting untested
genotypes in unobserved environments). The vertical pink line indicates the position of the overall phenotypic mean
across environments (87.74 days). The diagonal black line shows the exact match between predicted and observed
means. SMSE and cor stand for square root of the mean square error and the Person correlation between predicted
and observed DTH values. These were computed within environments; and the mean SMSE and mean cor
correspond to the across environments means.

499   Since the objective of this study is to develop a method for predicting DTH for untested
500 genotypes in unobserved environments, the predicted values obtained with the CB method were
501 used for fitting the DTH curves for each one of the 112 genotypes across environments. Figure
502 A3 shows the fitted curves for the predicted (orange) and observed (gray) DTH values across the
503 51 environments. In principle, these fitted curves can be used for predicting DTH for other
504 environments not included in this study. The information needed for defining a target
505 environment are the location and planting date allowing the computation of the theoretical DL
506 values using different methods (Forsythe et al., 1995). Once these theoretical values are obtained
507 the C method is implemented to find the estimated DTH E-means. In general, the fitted curves
508 (orange) derived from the predicted DTH values were very similar to those derived from the
509 observed DTH values (blue).

511              **DISCUSION**

512 The boxplot from Figure A2 shows a high heterogeneity among the environments in the study
513 with those with "late" planting date showing smaller phenotypic variability and shorter
514 occurrence time for DTH in comparison with the "early" planting environments. Conventional
515 GS models assume homogeneity of variances among environments and the violation of this
516 assumption may induce serious inconsistencies in the results. Especially when no genotype-by-
517 environment parameters are include in models.

518   The results obtained with the CV2_GS strategy can be viewed as the threshold of the
519 predictive ability levels that can be reached with the conventional implementation when
520 complete information about genotypes and environments is available. On the other hand, the
521 results from the CV00_GS strategy provide information of the model performance of the
522 conventional implementation when no phenotypic information is available for connecting
523 genotypes in target environments. This cross-validation scheme is among the three cases (CV2:
524 tested genotypes in observed environments; CV0: tested genotypes in unobserved environments;
525 and CV00: untested genotypes in unobserved environments) the most interesting, useful and
526 challenging for breeders. Dealing with traits where the selection process is based in ranks rather
527 than in the estimated *per se* values, the conventional implementation (GS) usually provides good
528 results (moderate to high correlations) under this scheme (CV00).

529   However, in traits related with important growth stages like DTH where the estimated *per*
530 *se* values are crucial for designing, planning and establishing experiments the implementation of
531 the conventional implementations (GS) might not be best choice. When no phenotypic
532 information of the target environment is available, the E-mean of the target environment is
533 determined as a weighted average of those environments in the training set forcing it towards a
534 general mean. Hence, the final prediction is composed of this general mean plus deviations due
535 to genomic effects. Since these deviations are conceptualized to be centered in zero, under CV0
536 and CV00 schemes the GS methods would produce similar environmental means (i.e., centered
537 on an overall mean determined by the environments in the calibration set).

538   The obtained results showed that the C method provides good estimations of DTH for the
539 CV0 scheme. Across environments, the SMSE was of 3.9 days and a Pearson correlation of 0.98
540 was obtained. These results were very similar to those obtained using the conventional GS
541 models when complete information of the genotypes and environments were available (i.e., CV2
542 scheme). Under CV2, the SMSE and Pearson correlation across environment were of 3.8 days
543 and 0.98, respectively.

544   As mentioned before, it is expected to obtain similar environmental means using either
545 CV0 or CV00 schemes under the conventional GS implementations. The, CV00_GS strategy
546 returned, across environments, a SMSE of 18.1 and a Pearson correlation of 0.41. The EMD
547 between predicted and observed values ranged between -31.5 and 28.7 days while for the CV0_C
548 strategy these ranged between -4.95 and 4.67 days. In this case, the CV0_C strategy used only
549 the phenotypic information of the genotype to be predicted but observed in other environments.
550 Since the CV0 and CV00 schemes produce similar E-means when no information of the target
551 environment is available using the conventional implementations (GS) it is also expected that the
552 proposed C method will produce good results with respect to the CV0_GS strategy not
553 considered in this study.

554   As was pointed before, the ultimate goal of this research is to develop a method able to
555 deliver small SMSE when predicting unobserved genotypes in unobserved environments (CV00
556 scheme). The BC method seems to produce good results under this scenario. The SMSE across
557 environments was 7.3 days with a Pearson correlation of 0.86. These results are very
558 encouraging compared with those obtained using the conventional implementation (GS) under
559 this scheme (SMSE: 18.1 and cor: 0.41). In this case, the BC method doubled the accuracy of the
560 conventional GS method and reduced the SMSE by 60%. However, the advantages of this
561 method were more evident when comparing the range of the differences between the predicted
562 and observed environmental means. Under the conventional implementation (CV00_GS) these
563 differences ranged between -31.5 and 28.7 days while with the proposed CB method (CV00_BC)
564 these ranged between -6.4 and 4.1 days. The proposed CB method reduced the extremes of this
565 range between 80 and 85%, respectively. These improvements can be observed in Figure 4 were
566 the environmental means of the proposed BC method (orange dots) were always closer to the
567 diagonal line, representing the exact match with true means, than those means obtained using the
568 conventional GS implementation (blue dots).

569   At beginning, the implementation of the proposed methods (C and BC) requires enough
570 information for an accurate characterization of the genotype's DL curves. For this, sets of
571 genotypes are tested in several environments. Perhaps it might sound expensive conduct these
572 experiments but this will allow the accurate estimation of the environmental means of the
573 unobserved environments. Once this information is available, potentially any untested genotypes
574 could be accurately predicted in any unobserved environment. Thus, the obtained results should
575 hold not only for the set of analyzed genotypes and environments of this study but also for
576 others.

CONCLUSIONS

GS is an emergent tool successfully implemented for screening candidate genotypes based on
their rank performance via the genomic estimated breeding values (GEBV). However, dealing
with traits linked to important growth development stages like DTH, the punctual prediction of
these traits (*per se* time wise values) becomes more important than the rank accuracy. In
scenarios where partial information of the target environment is available (CV2 and CV1
schemes), good estimations of the environmental means can be obtained. However, when no
information of the target environment is available (CV0 and CV00 schemes) a large bias
between the predicted and estimated environmental means can be expected; even when the rank
correlations may exhibit high values.

The developed methods here presented (C and BC) showed that the use of DL
information significantly improves the predictive ability of the conventional implementations
(GS) also reducing the bias between the predicted and observed DTH values. An advantage of
the C method is that if phenotypic information of the same genotype is available for a set of
environments, it is easy estimate the expected DTH in an unobserved environment without need
molecular marker information or phenotypic information from other genotypes. The BC method
improved the prediction of DHT of untested genotypes in unobserved environments by coupling
the marker profiles of the untested genotypes with the phenotypic and DL information from other
genotypes.

Both of the proposed methods require the day length (DL) information of the
environments (training and testing) in their construction. The convenience of the proposed
method resides in the fact that the theoretical DL information can be obtained in advance even
before establishing the experiments and thus it can be exploited during the planning and
designing stages. The proposed methods capture the genotype-by-environment interaction (G×E)
component via the fitted curves thus the explicitly inclusion of this term is not needed. The
conventional implementation (GS) returns the same estimated genomic effect across
environments for a particular genotype when the G×E component is absent. The curves precisely
capture the sensitivity experienced by genotypes under different environments driving specific
responses to each environment. Thus, although the proposed BC method is composed of the
genomic component computed without explicitly considering the G×E term the final prediction
accounts for this interaction term via the curves. In addition, the formal modeling of
heterogeneity of variances is not necessary because these patterns are also accounted with the
current method via the different curves.

ACKNOWLEDGEMENTS

613 We are grateful to Drs. Jose Crossa and Vikas Belamkar for their valuable feedback and
614 intellectual contribution during the elaboration of this manuscript.

REFERENCES

Bernardo R (2002) Breeding for Quantitative Traits in Plants. Stemma Press, USA.

Bolger, A. M., Lohse, M., and Usadel, B. 2014, Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics, 30, 2114–20.

Crossa, J., G. de los Campos, P. Pérez, D. Gianola, J. Burgueño, J.L. Araus, D. Makumbi, R.P. Singh, S. Dreisigacker, J. Yan, V. Arief, M. Banziger, and H.-J. Braun. 2010. Prediction of genetic values of quantitative traits in plant breeding using pedigree and molecular markers. Genetics 186:713–724

Crossa, J., D. Jarquín, J. Franco, P. Pérez-Rodríguez, J. Burgueñoet al., 2016.Genomic prediction of gene bank wheat landraces. G3 (Bethesda) 6:1819–1834.

de los Campos, G., J. M. Hickey, R. Pong-Wong, H. D. Daetwyler, and M.P. L. Calus, 2013 Whole-genome regression and prediction methods applied to plant and animal breeding. Genetics 193: 327–345.

FAO, IFAD, UNICEF, WFP and WHO. 2018. The State of Food Security and Nutrition in the World 2018. Building climate resilience for food security and nutrition. Rome, FAO

Forsythe, William C., Edward J. Rykiel Jr., Rand al S. Stahl, Hsin-i Wu and Robert M. Schoolfield, 1995. A model comparison for daylength as a function of latitude and day of the year. Ecological Modeling 80:87-95

Heffner HL, Sorrells ME, Jannink JL (2009) Genomic selection for crop improvement. Crop Science 49: 1–2.

Hori K, Matsubara K, Yano M (2016) Genetic control of flowering time in rice: integration of Mendelian genetics and genomics. Theor Appl Genetics 129:2241–2252.

Jannink JL, Lorenz AJ, and Iwata H (2010) Genomic selection in plant breeding: from theory to practice. Brief. Funct. Genomics 9: 166–177.

Jarquín, D., C. Lemes da Silva, R.C. Gaynor, J. Poland, A. Fritz, R. Howard, S. Battenfield, and J. Crossa. 2017. Increasing Genomic-Enabled Prediction Accuracy by Modeling Genotype × Environment Interactions in Kansas Wheat. Plant Genome 10:0. doi:10.3835/plantgenome2016.12.0130

Kawahara, Y., de la Bastide, M., Hamilton, J. P., et al. 2013, Improvement of the Oryza sativa Nipponbare reference genome using next generation sequence and optical map data. Rice, 6, 1–10.

Li, H. 2012, Exploring single-sample SNP and INDEL calling with whole-genome de novo assembly. Bioinformatics, 28, 1838–44.

Li, X., Guo, T., Mu, Q., Li, X. & Yu, J. Genomic and environmental determinants and their interplay underlying phenotypic plasticity. Proc. Natl Acad. Sci. USA 115, 6679–6684 (2018).

McKenna, A., Hanna, M., Banks, E., et al. 2010, The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res., 20, 1297–303.

Meuwissen TH, Hayes BJ, Goddard ME (2001) Prediction of total genetic value using genome-wide dense marker maps. Genetics 157:1819–1829.

Murray, M. G., and Thompson, W. F. 1980, Rapid isolation of high molecular weight plant DNA. Nucleic Acids Res., 8, 4321–5.

Onogi A, Ideta O, Inoshita Y, Ebana K, Yoshioka T, Yamasaki M, Iwata H (2015) Exploring the areas of applicability of whole-genome prediction methods for Asian rice (Oryza sativa L.). Theor Appl Genet 128:41–53.

Onogi A, Watanabe M, Mochizuki T, Hayashi T, Nakagawa H, Hasegawa T, Iwata H (2016) Toward integration of genomic selection with crop modelling: the development of an integrated approach to prediction rice heading dates. Theor Appl Genet 129:805–817.

Perez-de-Castro, A.M., S. Vilanova, J. Canizares, L. Pascual, J.M. Blanca, M.J. Diez, J. Prohens, and B. Pico. 2012. Application of genomic tools in plant breeding. Curr. Genomics 13:179–195. doi:10.2174/138920212800543084.

Robert J. Hijmans (2017). geosphere: Spherical Trigonometry. R package version 1.5-5. https://CRAN.R-project.org/package=geosphere

Spindel J, Begum H, Akdemir D, Virk P, Collard B, Redoña, Atlin G, Jannink JL, McCouch SR (2015) Genomic selection and association mapping in rice (Oryza sativa): effect of trait genetic architecture, training population composition, marker number and statistical model on accuracy of rice genomic selection in elite, tropical rice breeding lines. PLoS Genetics 11: e1004982.

VanRaden, P.M. Efficient methods to compute genomic predictions. J. Dairy Sci. 2008, 91, 4414–4423.

Voss-Fels KP, Cooper M, Hayes BJ (2019) Accelerating crop genetic gains with genomic selection. Theor Appl Genet 132:669–686.

Yin X, Kropff MJ, Aggarwal PK, Peng S, Horie T (1997) Optimal preflowering phenology of irrigated rice for high yield potential in three Asian environments: a simulation study. Field Crop Res 51:19–27.

APPENDIX

684
685 Figure A1. Scatter plot of days to heading (DTH) as function of the cumulative day length (CDL) from planting date
686 until the occurrence event for a rice data set comprising 112 genotypes observed in 51 environments (in Japan
687 between 2005 and 2017). From the total of all possible genotype-environment combinations (112×51=5712), 94%
688 (5365) of these were observed. The black line represents the regression line between DTH and CDL (R-squared of
689 0.99).
690

691
692 Figure A2. Boxplot for days to heading (DTH) for a rice data set comprising 112 genotypes tested in 51
693 environments (in Japan between 2005 and 2017). From the total of all possible genotype-environment combinations
694 (112×51=5712), 94% (5365) of these were observed. Environments were ordered based on the environmental
695 medians.
696

697
698   Figure A3. Fitted curves for observed (gray) and predicted (orange) DTH (Day length-*axis*) values for a rice data set
699   comprising 112 genotypes (Genotype-*axis*) tested in 51 environments between days 1 (planting date) and 180 (Days-
700   *axis*). The orange color curves were fitted using the predicted values obtained with the CB method under the CV00
701   scheme (predicting untested genotypes in unobserved environments) while the gray color curves considered the
702   observed data.

703
704   Table A1. Rice trials for studying days to heading for 112 genotypes tested in 79 environments (site-by-year-by-
705   planting date combinations) in multiple locations in Japan. In locations with more than 1 experiment per year
706   different seeding times were tested.

Location	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	Latitude	Altitude
Fukuoka									1	1	1	1	1	1	33.21	130.49
Fukuyama			1	1	1	1	2	2	2	2	2	2	2	2	34.5	133.39
Kasai			1	1	2	2	2	1	1	1	1	1	1	1	34.88	134.86
Tsukubamirai	2	2										1	2	2	36.01	140.02
Tsukuba					1	1	2	3	3	1	1				36.02	140.11
Toyama												1	1	1	36.62	137.23
Hokuriku												1	1	1	37.11	138.27
Akita									1	1	1	1	1	1	39.48	140.49
Sapporo										1	1	1	1	1	43.01	141.41

707