Gut microbiota structure differs between honey bees in winter and summer

Lucie Kešnerová[*1], Olivier Emery[*1], Michaël Troilo[1], Berra Erkosar[2*], Philipp Engel[1*†]

[1]Department of Fundamental Microbiology, University of Lausanne, CH-1015 Lausanne, Switzerland

[2]Department of Ecology and Evolution, University of Lausanne, CH-1015 Lausanne, Switzerland

[*]equal contribution between adjacent authors

[†]Correspondence:

Prof. Philipp Engel

Department of Fundamental Microbiology

University of Lausanne, CH-1015 Lausanne, Switzerland

Tel.: +41 (0)21 692 56 12

e-mail: philipp.engel@unil.ch

Keywords: *Bartonella, Lactobacillus*, *Apis mellifera*, diutinus bees, pollen diet, seasonality

Running title: Microbiota of summer and winter bees

Abstract

Honey bees harbor a relatively simple and highly conserved gut microbiota. Yet, little is known about the dynamics of the bee gut microbiota across seasons, especially in winter when adult worker bees live much longer and are critical for colony survival. We quantified the major community members of the bee gut microbiota in a single colony over two years and monitored 14 additional colonies during summer and winter. Our data shows that the total abundance of the gut microbiota in nectar/honey-feeding foragers was significantly lower compared to pollen-feeding nurses and winter bees. Strikingly, long-lived winter bees had the highest bacterial loads, with a characteristic shift towards a predominance of Lactobacillus Firm-5 and Bartonella apis. By colonizing microbiota-depleted bees with a community reconstituted from bacterial isolates, we provide experimental evidence that diet is a major contributor to the observed differences in bacterial loads between winter bees, nurses, and foragers. Overall, our study reveals that the gut microbiota of winter bees is remarkably different compared to foragers and nurses. Considering the importance of winter bees for colony survival, future work should focus on the role of the gut microbiota in health and disease in winter bees.

Introduction

The European honey bee, *Apis mellifera,* is an important pollinator species for natural ecosystems and agricultural production [1]. Its health status is threatened by numerous factors including habitat loss, pesticide exposure, and high parasite and pathogen loads [2–4]. Accumulating evidence point that the gut microbiota of adult honey bees plays a critical role for bee health [5]. It converts dietary compounds [6, 7] and produces short chain fatty acids [8] in the gut, enhances sucrose responsiveness of the host [8], and stimulates the immune system [9, 10]. Moreover, disruption of the gut microbiota composition by antibiotic treatment, pesticide exposure, or dietary manipulations was associated with increased pathogen loads resulting in increased mortality [11–14].

A striking feature of the honey bee gut microbiota is its low taxonomic complexity. In female worker bees, the community is dominated by seven to ten phylotypes (i.e. clusters of strains sharing ≥97% sequence identity in the 16S rRNA gene), which typically make up >95% of the bacterial cells in the gut [5, 15–18]. These phylotypes have been consistently detected in honey bees, regardless of geographic location, life stage, or sampling season [16, 19, 20], and are acquired horizontally through contact with nest mates and hive components [21]. They include five core phylotypes (*Gilliamella, Snodgrassella alvi, Lactobacillus* Firm-4 and Firm-5, and *Bifidobacterium*), which are present in all bees, and two non-core phylotypes, i.e. *Frischella perrara* and *Bartonella apis,* which are highly prevalent in colonies, but not in every bee [22]. On the other hand, phylotypes such as Alpha 2.1, Alpha 2.2, or *Lactobacillus kunkeii* account for a relatively small proportion of the gut community [23].

While these phylotypes are consistently present in adult worker bees, their abundance

may change across bees influencing their impact on the host. For example, the type and

amount of nutrients (i.e. pollen and nectar) available during the foraging season may

impact the metabolic activity of the gut microbiota and therefore modulate its

composition and function [24]. Likewise, distinct dietary habits or variation in life span of

worker bees during summer and winter may influence gut microbiota composition. From

spring to autumn, young worker bees (nurses) stay inside the hive to take care of larvae,

and feed on nutrient-rich pollen, whereas older worker bees become foragers that feed

on nectar and honey to fuel their energy-expensive flights [25]. In late autumn, newly

emerged adult worker bees become winter bees (also called 'diutinus') that have an

extended lifespan (~6 months) and that ensure the colony survival during the cold winter

season in the absence of brood [26]. These bees form a tight cluster for thermoregulation

inside the hive, feed strictly on food stores (pollen beebread and honey) and retain their

feces all winter [27], which is likely to impact the ecology of their gut microbiota.

A number of studies have looked at the gut microbiota composition of different worker

bee types or throughout seasons, with the overall conclusion that the community

composition is relatively stable [19, 28–31]. However, previous studies were mostly based

on comparative analyses of relative community member abundance using 16S rRNA

amplicon sequencing. Such analysis cannot provide insights about the extent or

directionality of changes in taxa abundance, especially if microbial loads vary substantially

between samples [32]. In fact, a change in the total abundance of the microbiota could by

itself be an important characteristic of different bee types, season, or environmental

exposure. A striking example is the experimental exposure of bees to antibiotics which

did not result in a strong shift in the relative composition, but in an overall reduction of

bacterial load, rendering bees more susceptible to pathogen invasion [11].

In addition to the limitations of current studies using relative abundance data, almost

nothing is known about the gut microbiota of winter bees as compared to foragers or

nurses. This is surprising, as winter bees are critical for colony health and survival during

the cold season of the year, when resources are limited and most colony losses occur [33,

34].

Characterizing the gut microbiota structure of winter bees and identifying the factors that

shape the gut communities may help us to understand the physiological adaptations that

are necessary to survive the cold season of the year in temperate regions. In this study,

we monitored the gut microbiota of 566 individual worker bees from a single hive over

two years using a qPCR-based approach. We confirmed our initial findings by analyzing

the microbiota of nurses, foragers, and winter bees from 14 hives in two subsequent

years, and performed experiments with gnotobiotic bees to test the influence of diet on

gut microbiota loads.

Materials and Methods

Sampling of honey bees
101

For monitoring the gut microbiota of worker bees over the period of two years, ~24 bees

were collected monthly from a single hive located at the apiary of the University of

Lausanne, Switzerland. During the foraging season, we sampled foragers returning to the

hive entrance with pollen on their legs, while during the cold winter months, we sampled

"winter bees" on top of the frames from inside the hive. Each sampling time point took

place at the middle of each month (+/- 3 days) between April 2015 and April 2017.

Samples from July 2017 were not included in the analysis due to a manipulation error

during DNA extraction.

To compare the gut microbiota of three different worker bee types (nurses, foragers, and

winter bees) across colonies, we sampled bees from 14 hives. Eleven hives were located

on the Dorigny campus of the University of Lausanne and three hives were located in the

village of Yens, about 17 km away. Foragers and nurses were sampled in July 2017 and

August 2018, and winter bees were sampled in January 2018 and January 2019. "Nurses"

and winter bees were collected inside the hive in the center of the colony. Foragers were

collected outside the hive at the same time point as nurses. The bees sampled in July

2017 and January 2018 were used for gut content visualization and correlation of gut

weight and 16S rRNA copy numbers. The guts of the bees sampled in August 2018 and

January 2019 were pooled (20 guts per bee type per hive) and used to monitor the

abundance of individual community members.

To dissect the guts, bees were anesthetized with CO_2 for 10 s, and the guts including crop,

mid- and hindgut and Malpighian tubules were carefully removed using sterile forceps.

For the monthly sampling of the single bee hive, each gut sample was placed in a drop of

PBS, scored for the scab phenotype [35], and placed in a bead beating tube containing

approx. 150 mg of glass beads (0.75-1 mm in diameter, Carl Roth) and 750 µL of CTAB

lysis buffer (0.2 M Tris-HCl, pH 8; 1.4 M NaCl; 0.02 M EDTA, pH 8; 2% CTAB, w/v, dissolved

at 56°C overnight; 0.25% β-mercaptoethanol, v/v). For the sampling of the 14 colonies in

August 2018 and January 2019, the dissected guts were pooled into a single Falcon tube

(14 hives x 3 bee types = 42 pooled samples). Tubes were flash frozen in liquid nitrogen

and stored at -80°C until DNA extraction.

Experimental colonization of honey bees

Microbiota-depleted bees were generated and colonized as described in Kešnerová et al.

[6]. The treatment group of bees fed on pollen and sugar water was the same as in this

previous study. The other treatment group (bees fed on sugar water only) was carried out

in parallel with the same batch of bees. Bees were fed *ad libitum* with sterilized bee

pollen and sterilized sugar water (50% sucrose, w/v) (SW+P treatment), or with only

sterilized sugar water (SW treatment). Bees were sampled 10 days after colonization and

the guts were dissected as described above.

DNA extraction from honey bee gut tissue

A previously established CTAB/phenol-based extraction protocol [6] was used to extract

DNA/RNA from individual guts. At the end of the protocol, the precipitated dried pellet

was resuspended in 200 μl and split into two samples à 100 μl. One sample was

processed with the Nucleospin PCR Clean-up kit (Macherey-Nagel, Germany) according to

the manufacturer's instructions and the resulting DNA was used for qPCR. For the pooled

samples containing 20 guts, 2 ml of glass beads and 15 ml of CTAB lysis buffer were added

to each Falcon tube. Samples were then homogenized in a Fast-Prep24^{TM}5G homogenizer

at 6 m/s for 40 s, briefly centrifuged, and an aliquot of 750 μl corresponding to the

sample volume of one bee gut was transferred to a new 2 ml bead beating tube with glass

beads and homogenized again. All further steps in DNA extractions were performed as

previously described [6].

Quantitative PCR (qPCR) to determine absolute abundance of community members

We used the same qPCR methods and data analysis as described in Kešnerová et al. [6]. In

short, to determine the absolute copy number (i.e. abundance) of each target in each

sample, we performed standard curves on serial dilutions of plasmids containing the

targeted sequence. The number of plasmid copies in these dilutions ranged from 10^1–10^7

copies per µl. The plasmid copy number was calculated from the molecular weight of the

plasmid and the DNA concentration of the purified plasmid measured with NanoDrop.

The slope and intercept of standard curves were calculated based on the Cq values

(quantification cycle [36]) obtained from the dilutions with 10^2 - 10^7 copies. For most

primer pairs, no amplification was obtained for the highest dilution, or the amplification

reached a similar Cq as the negative water control. As the *limit of detection* (LOD) of a

given primer pair, we consider the detectable Cq value of the highest plasmid dilution.

The target was considered absent in the sample when the Cq value was higher than the

LOD Cq. The *Efficiency of primers (E)* was estimated from the slope according to the

equation: $E = 10^{(-1/slope)}$ [37]. Primer characteristics and their performance are summarized

in **Table S1**.

All samples for which the Cq value of actin was >24 were excluded from the analysis, as

the extracted DNA was considered to be not of sufficient quality. For each DNA sample

that passed the initial quality check, we determined the number of bacterial cells per gut

as follows: We first calculated the 'raw' copy number (*n*) of each target in 1 µl of DNA

sample from the Cq value and the standard curve using the formula $n = E^{(\text{intercept} - Cq)}$ [38].

Then, we normalized bacterial 16S rRNA gene copies to the median number of actin gene

copies in a given dataset and corrected this number considering the total volume of the

samples (200µl) to obtain normalized copy numbers per gut (*Normn*): *Normn =*

n_{target}/n_{actin} *x Median($n_{(actin)}$) x 200.* Normalization with the actin gene was carried out to

reduce the effect of gut size variation and extraction efficiency. To infer the number of

microbial cells from the normalized 16S rRNA gene copy number we additionally divided

it by the number of 16S rRNA loci present in the genome of the given species (as listed in

Table S1).

Statistics and diversity analyses of the gut microbiota composition

All statistical analyses were performed using R (version 3.5.1). We measured alpha

diversity based on the absolute quantification of the seven dominant gut microbiota

phylotypes using effective number of species [39] that is calculated by taking the

exponent of Shannon's diversity index ("diversity function", from "Vegan" package in R

[40].

We performed a principal components analysis (PCA) with the *prcomp* function of the R

package "stats" to determine the similarity of the bacterial communities between

foragers and winter bees across the monthly sampling over two years using absolute

abundance measures of the seven gut microbiota phylotypes.

We tested the effect of bee type on bacterial loads, diversity indices and wet gut weight

using Student's t-test (in case of two group comparisons) or general linear models (in case

of three group comparisons). Since the residuals obtained for certain models showed

heteroscedasticity, we used a permutation approach to test the significance of the effects

as described before [41]. Briefly, we randomized the values of the response variable

10'000 times and computed the F-values/t-values for the tested effect for each

randomized dataset. The p-values corresponding to the effects were calculated by the

proportion of 10'000 F-values that were equal or higher than the observed one. Pairwise

comparisons between different factors were performed by Tukey's HSD using

"multcomp" package [42] using *glht* function on the model. P-values were adjusted using

the Bonferroni method.

Results

Bacterial loads and community composition differ between foraging and winter season in a honey bee colony monitored over two years.

To characterize the gut microbiota of worker bees across seasons, we tracked the total abundance of five core (*Gilliamella*, *S. alvi*, *Bifidobacterium*, *Lactobacillus* Firm4, and *Lactobacillus* Firm5) and two non-core members (*F. perrara* and *B. apis*; **Table S1**) in adult worker bees from a single hive over two years. Our analyses included 566 individual bee samples.

The core members *Gilliamella*, *S. alvi*, *Lactobacillus* Firm-5 and *Bifidobacterium* were present in all analyzed bees, and the core member *Lactobacillus* Firm-4 was detectable in 98.4% of all bees (**Supplementary Figure S1A**). Notably, the two designated non-core members *B. apis and F. perrara* were also present at relatively high prevalence with 94.7% and 73.1%, respectively (**Supplementary Figure S1B** & **C**). Consistent with our previous results, *F. perrara* prevalence strongly correlated with the presence of the scab phenotype (**Supplementary Figure S2**), a local melanization response that is induced by *F. perrara* upon colonization [35].

The absolute abundance of the monitored phylotypes varied little among the bees sampled at the same time point, with the exception of the non-core members *F. perrara* and *B. apis* (**Supplementary Figure S3**). However, there were clear differences in bacterial abundances between months (Permutation ANOVA P=1e-4) for all microbiota members. In particular, we observed remarkable differences in the bacterial loads between bees sampled during the foraging and the winter season in both years. This became evident from the abundance of individual phylotypes and from the total bacterial load as based

on the summed abundances of all seven phylotypes (**Supplementary Figure S3, Figure**

1A). Specifically, we found a 10- to 100-fold increase in the levels of the core members

Firm-4, Firm-5, and *Bifidobacterium*, as well as the non-core members *B. apis* when

comparing across all winter bees relative to foragers (**Figure 1B**, Permutation T-Test P=1e-

4). We also observed a small increase of *S. alvi* levels in winter bees (**Figure 1B**,

Permutation T-Test P=6e-4), but no difference in the levels of *Gilliamella* (**Figure 1B**,

Permutation T-Test P=0.7). *F. perrara* was the only member of the community which

displayed the opposite trend, i.e. lower abundance in winter bees (**Figure 1B**,

Permutation T-Test P=1e-4). The overall bacterial load was about 10x larger in winter

bees than in foragers as based on both the summed abundances of all seven phylotypes

(**Figure 1B**, Permutation T-Test P=1e-4) as well as the number of universal 16S rRNA gene

copies, which was determined for a subset of the samples (**Figure 1C**, Permutation T-Test,

P=1e-4).

Considering that the monitored phylotypes comprise the majority of the detected species

in the honey bee gut microbiota, we analyzed the diversity of microbial communities

based on our dataset neglecting the presence of other (likely very low-abundant) non-

core members. In both years, the communities of winter bees were largely dominated by

the phylotypes Firm-5 and *B. apis.* In contrast, forager bees seem to have more even

community compositions (**Figure 1D**). The dominance of Firm-5 and *B. apis* in winter bees

was reflected by the lower effective number of species (calculated from Shannon's

diversity index) in winter bees compared to foragers (**Figure 1E**, Permutation T-Test,

P=1e-4). Moreover, PCA revealed a clear separation between foragers and winter bees

(**Figure 1F,** MANOVA Wilks =0.6 $F_{(7, 392)}$=39.4, P<2.2e-4) along the principal component 1

(PC1). This separation was driven by Firm-4, Firm-5, *B. apis* and *Bifidobacterium*, the four

phylotypes with the largest differences in abundance between the two types of bees

(**Figure 1B**).

Taken together, these results suggest that the gut microbiota of winter bees and foragers

markedly differ from each other in the monitored hive, both in terms of the absolute

bacterial abundance and the relative community composition, with a few phylotypes

dominating the community in winter bees.

Consistent difference in bacterial loads and community composition between foragers,

nurses, and winter bees across colonies

The observed differences in bacterial loads between foragers and winter bees prompted

us to check for similar patterns across 14 different hives in two subsequent years. In

addition to foragers and winter bees, we also monitored the bacterial loads in nurses,

because this would help us understand whether the differences between foragers and

winter bees are linked to different seasons or to behavioral or dietary differences.

Strikingly, we found that the pooled gut samples from the fourteen different hives

showed a similar pattern as the individual samples of the hive monitored over two years.

Winter bees had higher bacterial loads than forager bees based on the summed

abundances of the seven monitored phylotypes (**Figure 2A,** Permutation ANOVA followed

by Tukey P=3.782-10) and the overall bacterial quantification with the universal 16S rRNA

primers (**Figure 2B,** Permutation ANOVA followed by Tukey, P=5.1e-9). As for the

previously monitored hive, *Bifidobacterium*, *Lactobacillus* Firm4, *Lactobacillus* Firm5, and

B. apis had increased levels (Permutation ANOVA followed by Tukey, P=1e-4), while *F.*

perrara had lower levels in winter bees compared to foragers (**Figure 2A,** Permutation ANOVA followed by Tukey P=3.36e10-3). The only two phylotypes which showed abundance patterns inconsistent with the results from the previous hive were *S. alvi* and *Gilliamella*. *S. alvi* did not experience any differences in abundance (**Figure 2A,** Permuation ANOVA P=5.52e-1), while *Gilliamella* had increased levels in winter bees as compared to foragers (**Figure 2A,** Permutation ANOVA followed by Tukey P=4.33e-08). Interestingly, we also found that nurse bees had increased bacterial loads compared to foragers (**Figure 2A,** P=1.33e-5; and **Figure 2B**). The levels of *Gilliamella* and Firm4 in nurse bees were as high as in winter bees, while the levels of *Bifidobacterium, Lactobacillus* Firm5*, and B. apis* were at intermediate levels between foragers and winter bees (see **Supplementary Table S2** for statistics).

The differences of the gut microbiota of foragers, nurses, and winter bees were also evident from the analysis of the relative abundances at the community level. *B. apis* and *Lactobacillus* Firm5 were both more dominant in winter bees as compared to foragers, while in foragers the phylotypes *F. perrara* and *S. alvi* accounted for a larger proportion of the community (**Figure 2C,** see **Supplementary Table S2** for statistics). In the PCA, foragers and winter bees clustered separately along PC1, explaining 57.3% of the variation in the data (**Figure 2D**). The communities of most nurse samples were less dominated by *B. apis* than those of winter bees (**Figure 2C**). However, they still clustered more closely with the winter bees than the foragers (**Fig. 2D,** MANOVA Wilks =0.1, $F_{(14,66)=}7.9$, P=1.7e-9).

In summary, our analysis across 14 hives suggests that the differences in bacterial loads and community structure between winter bees, foragers, and nurses is a general pattern,

and that the microbiota of winter bees resembles more closely the one of nurses in terms

of absolute bacterial abundances.

Pollen diet increases gut community size in gnotobiotic bees

One of the drivers of the observed differences in bacterial load and community structure

between winter bees, foragers, and nurses could be diet. Dietary differences between the

three types of bees were clearly visible from the dissected guts (**Figure 3A-C**). The

rectums of winter bees and nurses appeared yellow indicating the presence of pollen,

while the rectums of forager bees appeared translucent indicating the absence of pollen.

Moreover, the wet weight of the guts was significantly different between the three types

of bees (ANOVA $F_{(2,68)}$ = 24.13, P= 1.205e-8), with foragers having on average two times

lighter guts than nurses (Tukey P=1.14e-6) and winter bees (Tukey P=6.81e-8) (**Figure 3D**).

When plotting normalized 16S rRNA gene copy numbers as a function of gut weight, we

found that gut weight positively correlated with total microbiota abundances across the

three bee types (**Figure 3E**).

In order to demonstrate that pollen diet is directly associated with increased bacterial

loads in honey bees, we experimentally colonized newly emerged bees with a community

of 11 bacterial strains representing the seven major bacterial phylotypes of the bee gut

microbiota [6]. The colonized bees were kept in the laboratory for ten days and fed *ad*

libitum either sterile sugar water and pollen (SW+P treatment), or sugar water only (SW

treatment). We found a significant difference in the gut weights between the two

treatments (**Figure 3F**, Welch's T-Test *t*=9.433, P=1.452e-11). While the gut weights of the

bees of the SW treatment were comparable to those of forager bees, the gut weights of

the bees of the SW+P treatment were markedly higher, exceeding even those of winter

bees (**Figure 3D** & **3F**). Strikingly, we observed a positive correlation between gut weight

and microbiota abundance for both the colonized bees in the laboratory (**Figure 3G**) and

the worker bees sampled from the hive (**Figure 3E**). Moreover, differences in bacterial

loads of individual community members between the two experimental treatments

mirrored, to a large extent, the difference found between winter bees, nurses, and

foragers: most phylotypes were more abundant in bees fed on pollen as compared to

bees fed sugar water only (**Figure 3H**, see **Supplementary Table S2** for statistics). Two

exceptions were *B. apis* and *F. perrara.* While *B. apis* had similar levels between the two

experimental treatments **(Figure 3H)**, its abundance was higher in winter bees and nurses

as compared to foragers **(Figure 2A)**. Interestingly, *B. apis* was able to colonize only 75%

of all bees when pollen was absent. The dependence on pollen for gut colonization was

even more pronounced for *F. perrara.* Less than 50% of the experimentally colonized bees

of the SW treatment had detectable levels of *F. perrara,* and the loads in bees that were

colonized were relatively low. In contrast, bees of the SW+P treatment were all colonized

and had relatively high and consistent loads *of F. perrara* **(Figure 3H)***.*

Taken together, these results show that a pollen diet leads to an increase in gut weight

and overall bacterial load providing a plausible explanation for some of the differences in

the loads observed between foragers, nurses, and winter bees.

Discussion

In contrast to 16S rRNA amplicon sequencing, qPCR provides information about the absolute abundances of bacteria and therefore allows determining whether individual community members increase, decrease, or remain the same in terms of bacterial cell number across samples. For example, in our study, the relative abundance of *S. alvi* went down in winter bees as compared to foragers and nurses (**Figure 2C**). However, this effect was not due to a decrease of the total number of *S. alvi* cells, but rather an increase of other community members as identified by qPCR. In fact, the total abundance of *S. alvi* remained the same in foragers, nurses, and winter bees **(Figure 2A)**. Such quantitative microbiome profiling approaches can reveal important associations between bacterial symbionts and the host, as previously demonstrated for the human microbiota [32] or the microbiota of caterpillars [43]. We argue that absolute abundances should be routinely assessed when analyzing microbial communities, as the absolute number of cell of a bacterium - not its relative abundance – may determine its impact on ecosystem processes.

Noteworthy, qPCR is a targeted approach, i.e. we can only quantify members for which specific primer sets have been designed. This is a limitation for microbial communities that are highly diverse. However, in the current study, we took advantage of the fact that the honey bee gut microbiota consists of a relatively simple community typically dominated by the seven analyzed phylotypes [5, 22, 23, 44–46]. Hence, while our approach did not cover some of the non-core members that are sporadically present in the bee gut, we are confident that the majority of the diversity of the bee gut microbiota has been quantified. This is supported by the fact that the community profiles obtained

from the qPCR data looked very similar to those obtained in previous studies based on

16S rRNA gene amplicon sequencing [21, 28, 47–49].

Almost all members of the gut microbiota had increased levels in winter bees when

compared to foragers, and many also showed increased levels when compared to nurses.

The most pronounced differences were found for *Lactobacillus* Firm-5 and *B. apis,*

resulting in a remarkable shift of the community towards these two phylotypes.

Strikingly, this shift was observed across fourteen different colonies and in three different

years. Hence, the "reconfiguration" of the microbiota in winter bees seems to be a

conserved feature in colonies in Western Switzerland. However, regional differences in

floral diversity [24] or climate may influence these changes. Therefore, additional surveys

of winter bees in other geographic regions are needed to test for the conservation of this

pattern. Interestingly, a recent study carried out in Germany on the effects of winter

supplementation feeds also revealed that the relative abundance of *Lactobacillus* Firm-5

(and in one treatment group also the relative abundance of *B. apis*) markedly increased in

winter bees compared to foragers [19], resulting in a community shift. The results of this

study were also consistent with our findings in that the levels of *F. perrara* were

significantly lower in winter bees as compared to forager bees.

What drives the changes in bacterial loads and community composition in winter bees? A

possible explanation could be dietary differences between the analyzed bee types. It is

well known that foragers mainly feed on nectar and honey, while nurses and winter bees

also consume pollen [50]. These dietary differences were also evident from the three bee

types sampled in our study based on consistent changes in appearance and weight of the

gut (**Figure 3A-C**). Strikingly, our bee colonization experiment with a defined bacterial

community showed that the presence of pollen in the diet substantially increased gut

weight and bacterial loads to levels comparable to those in winter bees, while the levels

in bees fed on sugar water were more similar to those of foragers (**Figure 3F-H**).

Therefore, we conclude that diet is an important factor that can explain many of the

differences observed between worker bee types. Seasonal changes in gut microbiota

composition in wild rodent populations [51] and humans [52, 53] have also been found to

coincide with dietary shifts, which is in agreement with the general notion that dietary

preferences is the main driver of community differences across a wide range of animals

[54–56].

In the case of honey bees, the larger amount of food in the gut is likely to increase the

carrying capacity for the gut microbiota. In addition, pollen is a more nutrient-rich diet

than nectar, honey, or sucrose offering a larger diversity of different metabolic niches for

gut bacteria. Both factors are likely to contribute to the increased bacterial loads in bees

fed on pollen as compared to those fed on sugar water only. The role of pollen as a main

driver for the observed differences is also supported by the fact that the bacteria with the

largest increase in total abundance (*Lactobacillus*, *Bifidobacteria*, *B. apis*) are located in

the rectum, which is the last part of the hindgut where pollen accumulates until bees

defecate. In line with these, a previous report showed that the abundances of total

bacteria, as well as certain individual phylotypes (*Lactobacillus* Firm-5, *Bifidobacterium*)

increase in rectum upon pollen consumption [57]. However, this increase was dependent

on the age factor and it was not significant when sterile pollen (similar to our pollen

treatment) was used instead of stored pollen [57]. In contrast to our study the

experimental bees were not inoculated by bacteria [57], which may greatly impact

community growth and dynamics. Overall, despite certain experimental differences, our

results seem overall to be consistent with the data that has been published before.

However, not all changes observed in winter bees could be recapitulated in our

colonization experiment. For example, the differences observed *in B. apis* levels between

foragers, nurses, and winter bees were not observed in the experimental bees that were

fed with or without pollen. Another example is *F. perrara*. While this bacterium was less

abundant in winter bees than in foragers and nurses, the colonization success of

F. perrara was largely dependent on the presence of pollen in the experimental bees. This

suggests that other factors may contribute to community differences found in winter

bees as compared to foragers or nurses. First, winter bees have an extended lifespan with

an average life expectancy of ~6 months as compared to ~4 weeks in the case of summer

bees (i.e. nurses and foragers) [58]. In contrast, the bees of the colonization experiment

were age-matched and sampled ten days after emergence. It is well known that the

physicochemical state of the gut changes as animals age, modulating their gut microbiota

[59–61]. Second, winter bees feed on pollen that has been stored in the hive for several

weeks to months. It has previously been shown that the consumption of an aged pollen-

diet affects the gut microbiota composition of nurses, causing dysbiosis and higher

mortality [12]. Third, winter bees retain their feces in the gut for extended periods of

time, which likely affects the physico-chemical conditions and the availability of nutrients

in the gut. Forth, the body temperature of bees is at ~21°C during winter and at ~35°C

during summer [62]. This may influence bacterial growth rates. Indeed, in a recent metagenomic study, it was shown that gut bacteria have lower average population replication in old winter bees as compared to young nurse bees, which is indicative of decreased replication rates [18]. Fifth, winter bees show reduced expression of immune genes [63–66], and have an altered protein metabolism [67].

How the different factors influence the dynamics of specific gut microbiota members should be the focus of future studies. In the case of *F. perrara* it is tempting to speculate that the decrease in colonization levels in old winter bees, may be a consequence of the specific immune response elicited by the host towards this bacterium [10], eliminating it from the gut as the bee ages. Another interesting finding is that the levels of *S. alvi* barely changed across worker bee type or the two diet treatments in the experiment. This suggests that the colonization of *S. alvi* is not modulated by the dietary state, the physicochemical conditions in the gut, or the abundance of other community members. A possible explanation could be that the niche of *S. alvi* is dependent on the host rather than the diet, because it selectively colonizes the epithelial lining of the ileum and was suggested to rely on the presence of oxygen for growth released from the host epithelium [8, 44, 68].

The two phylotypes that predominate in winter bees, *Lactobacillus* Firm5 and *B. apis*, seem to profit from the specific gut conditions in winter bees. The primary metabolism of *Lactobacillus* Firm5 is fermentation of sugars or sugar derivatives into organic acids [22, 69]. Genomic analysis revealed that these bacteria harbor a large diversity of glycoside

hydrolases and phosphotransferase transporters for the cleavage and transport of sugars.

Moreover, a recent metabolomics study has found that flavonoids, secondary plant

metabolites that are abundant in pollen, are prominent substrates of these bacteria. In

contrast to *Lactobacillus* Firm5, *B. apis* encodes a complete TCA cycle and harbors a

respiratory chain. Some strains of *B. apis* also possess genes for the degradation of

aromatic compounds. It is possible that metabolic interactions among these two

particular phylotypes or the extended gut-retention time of pollen (resulting in the

availability of additional nutrients) facilitates their increased abundance in winter bees.

Most of the recent colony losses occur during the winter months [33, 34]. Consequently,

winter bees are highly critical for colony survival and a better understanding of the

factors influencing their health status - including the gut microbiota - is needed. In

summary, our analysis revealed that the gut microbiota of winter bees undergo

characteristic shifts. These changes may have important consequences for the host. For

example, they may influence the physiology or metabolic state of the host by optimizing

the energy uptake [70], or increase colonization resistance against pathogens. Therefore,

future studies should specifically focus on the functional role of the gut microbiota in

winter bees, and colony health.

Acknowledgements

We would like to thank Paul Reymond, Clément Etter, Katherine Lane for their help

during the sampling of honey bees. We are also grateful to Andrew Quinn and Fabienne

Wichmann for comments on the manuscript. This study was supported by the European

Research Council (ERC-StG 'MicroBeeOme') and the Swiss National Science Foundation

(grant number 31003A_160345 and 31003A_179487) received by PE. The funder had no

role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript.

Competing Interests

Authors declare to have no conflicts of interest.

Data accessibility

All scripts and datasets can be found on:

https://drive.switch.ch/index.php/s/GsX9TBGcNJ8WEuv

Supplementary Material

Supplementary Figures: Contains Figures S1-S3

Supplementary Table S1: Contains information on the primers that are used in this study.

Supplementary Table S2: Contains the results of the statistical tests used in this study.

References

1. Kleiner A-M, Cane James H, Steffan-Dewenter Ingolf, Cunningham Saul A, Kremen Claire, Tscharntke Teja. Importance of pollinators in changing landscapes for world crops. *Proc R Soc B Biol Sci* 2007; **274**: 303–313.

2. Barron AB. Death of the bee hive: understanding the failure of an insect society. *Curr Opin Insect Sci* 2015; **10**: 45–50.

3. Nazzi F, Pennacchio F. Disentangling multiple interactions in the hive ecosystem. *Trends Parasitol* 2014; **30**: 556–561.

4. Goulson D, Nicholls E, Botías C, Rotheray EL. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* 2015; **347**: 1255957.

5. Raymann K, Moran NA. The role of the gut microbiome in health and disease of adult honey bee workers. *Curr Opin Insect Sci* 2018.

6. Kešnerová L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, Engel P. Disentangling metabolic functions of bacteria in the honey bee gut. *PLOS Biol* 2017; **15**: e2003467.

7. Engel P, Martinson VG, Moran NA, Moran NA. Functional diversity within the simple gut microbiota of the honey bee. *Proc Natl Acad Sci* 2012; **109**: 11002–11007.

8. Zheng H, Powell JE, Steele MI, Dietrich C, Moran NA. Honeybee gut microbiota promotes host weight gain via bacterial metabolism and hormonal signaling. *Proc Natl Acad Sci* 2017; **114**: 4775–4780.

9. Kwong WK, Mancenido AL, Moran NA. Immune system stimulation by the native gut microbiota of honey bees. *Open Sci* 2017; **4**: 170003.

10. Emery O, Schmidt K, Engel P. Immune system stimulation by the gut symbiont Frischella perrara in the honey bee (Apis mellifera). *Mol Ecol* 2017; **26**: 2576–2590.

11. Raymann K, Shaffer Z, Moran NA. Antibiotic exposure perturbs the gut microbiota and elevates mortality in honeybees. *PLOS Biol* 2017; **15**: e2001861.

12.   Maes PW, Rodrigues PAP, Oliver R, Mott BM, Anderson KE. Diet-related gut bacterial

dysbiosis correlates with impaired development, increased mortality and Nosema disease in

the honeybee (Apis mellifera). *Mol Ecol* 2016; **25**: 5439–5450.

13.   Motta EVS, Raymann K, Moran NA. Glyphosate perturbs the gut microbiota of honey bees.

Proc Natl Acad Sci 2018; 201803880.

14.   Blot N, Veillat L, Rouzé R, Delatte H. Glyphosate, but not its metabolite AMPA, alters the

honeybee gut microbiota. *PLOS ONE* 2019; **14**: e0215466.

15.   Kwong WK, Moran NA. Gut microbial communities of social bees. *Nat Rev Microbiol* 2016;

14: 374–384.

16.   Moran NA, Hansen AK, Powell JE, Sabree ZL. Distinctive Gut Microbiota of Honey Bees

Assessed Using Deep Sampling from Individual Worker Bees. *PLoS ONE* 2012; **7**: e36393.

17.   Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, Moran NA. A simple and

distinctive microbiota associated with honey bees and bumble bees. *Mol Ecol* 2011; **20**: 619–

628.

18.   Ellegaard KM, Engel P. Genomic diversity landscape of the honey bee gut microbiota. *Nat*

Commun 2019; **10**: 446.

19.   D'Alvise P, Böhme F, Codrea MC, Seitz A, Nahnsen S, Binzer M, et al. The impact of winter

feed type on intestinal microbiota and parasites in honey bees. *Apidologie* 2018; **49**: 252–264.

20.   Corby-Harris V, Maes P, Anderson KE. The Bacterial Communities Associated with Honey

Bee (Apis mellifera) Foragers. *PLoS ONE* 2014; **9**: e95056.

21.   Powell JE, Martinson VG, Urban-Mead K, Moran NA. Routes of Acquisition of the Gut

Microbiota of the Honey Bee Apis mellifera. *Appl Environ Microbiol* 2014; **80**: 7378–7387.

22.   Kwong WK, Moran NA. Gut microbial communities of social bees. *Nat Rev Microbiol* 2016;

14: nrmicro.2016.43.

23.   Corby-Harris V, Maes P, Anderson KE. The Bacterial Communities Associated with Honey

Bee (Apis mellifera) Foragers. *PLOS ONE* 2014; **9**: e95056.

24. Jones JC, Fruciano C, Hildebrand F, Al Toufalilia H, Balfour NJ, Bork P, et al. Gut

microbiota composition is associated with environmental landscape in honey bees. *Ecol Evol*

2017; **8**: 441–451.

25. Brodschneider R, Crailsheim K. Nutrition and health in honey bees. *Apidologie* . 2010. , 278–

294

26. Winston M. The biology of the honey bee, Harvard University Press. 1987.

27. Pavlovsky EN, Zarin EJ. Memoirs: On the Structure of the Alimentary Canal and its

Ferments in the Bee (Apis mellifera L.). *Q J Microsc Sci* 1922; **66**: 509–556.

28. Corby-Harris V, Maes P, Anderson KE. The Bacterial Communities Associated with Honey

Bee (Apis mellifera) Foragers. *PLoS ONE* 2014; **9**: e95056.

29. Rothman JA, Carroll MJ, Meikle WG, Anderson KE, McFrederick QS. Longitudinal Effects

of Supplemental Forage on the Honey Bee (<Emphasis Type="Italic">Apis

mellifera</Emphasis>) Microbiota and Inter- and Intra-Colony Variability. *Microb Ecol*

2018; 1–11.

30. Kapheim KM, Rao VD, Yeoman CJ, Wilson BA, White BA, Goldenfeld N, et al. Caste-

Specific Differences in Hindgut Microbial Communities of Honey Bees (Apis mellifera).

PLoS ONE 2015; **10**.

31. Ludvigsen J. Seasonal trends in the midgut microbiota of honeybees. *88* 2013.

32. Vandeputte D, Kathagen G, D'hoe K, Vieira-Silva S, Valles-Colomer M, Sabino J, et al.

Quantitative microbiome profiling links gut community variation to microbial load. *Nature*

2017; **551**.

33. Guzmán-Novoa E, Eccles L, Calvete Y, Mcgowan J, Kelly PG, Correa-Benítez A. Varroa

destructor is the main culprit for the death and reduced populations of overwintered honey

bee (Apis mellifera) colonies in Ontario, Canada. *Apidologie* 2010; **41**: 443–450.

34. Genersch E, Ohe W von der, Kaatz H, Schroeder A, Otten C, Büchler R, et al. The German

bee monitoring project: a long term study to understand periodically high winter losses of

honey bee colonies. *Apidologie* 2010; **41**: 332–352.

35.   Engel P, Bartlett KD, Moran NA. The Bacterium Frischella perrara Causes Scab Formation

573   in the Gut of its Honeybee Host. *mBio* 2015; **6**: e00193-15.

574   36.   Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al. The MIQE

575   Guidelines: Minimum Information for Publication of Quantitative Real-Time PCR

576   Experiments. *Clin Chem* 2009; **55**: 611–622.

577   37.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT–PCR.

578   *Nucleic Acids Res* 2001; **29**: e45–e45.

579   38.   Gallup JM. qPCR Inhibition and Amplification of Difficult Templates. *PCR Troubleshooting*

580   *and Optimization: The Essential Guide*. 2011. Horizon Scientific Press, pp 23–65.

581   39.   Lou Jost. Entropy and diversity. *Oikos* 2006; **113**: 363–375.

582   40.   Oksanen J, Blanchett FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:

583   Community Ecology Package. R package version 2.5-3. 2018.

584   41.   Lavanchy G, Strehler M, Llanos Roman MN, Lessard-Therrien M, Humbert J, Dumas Z, et

585   al. Habitat heterogeneity favors asexual reproduction in natural populations of grassthrips.

586   *Evol Int J Org Evol* 2016; **70**: 1780–1790.

587   42.   Hothorn T, Bretz F, Westfall P. Simultaneous Inference in General Parametric Models. *Biom*

588   *J* 2008; **50**: 346–363.

589   43.   Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N. Caterpillars lack a resident gut

590   microbiome. *Proc Natl Acad Sci* 2017; **114**: 9641–9646.

591   44.   Martinson VG, Moy J, Moran NA. Establishment of Characteristic Gut Bacteria during

592   Development of the Honeybee Worker. *Appl Environ Microbiol* 2012; **78**: 2830–2840.

593   45.   Sabree ZL, Hansen AK, Moran NA. Independent Studies Using Deep Sequencing Resolve

594   the Same Set of Core Bacterial Species Dominating Gut Communities of Honey Bees. *PLOS*

595   *ONE* 2012; **7**: e41250.

596   46.   Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, Moran NA. A simple and

597   distinctive microbiota associated with honey bees and bumble bees. *Mol Ecol* 2011; **20**: 619–

598   628.

599   47.   Sabree ZL, Hansen AK, Moran NA. Independent Studies Using Deep Sequencing Resolve

600   the Same Set of Core Bacterial Species Dominating Gut Communities of Honey Bees. *PLoS*

601   *ONE* 2012; **7**: e41250.

602   48.   Moran NA, Hansen AK, Powell JE, Sabree ZL. Distinctive Gut Microbiota of Honey Bees

603   Assessed Using Deep Sampling from Individual Worker Bees. *PLoS ONE* 2012; **7**: e36393.

604   49.   Yun J-H, Jung M-J, Kim PS, Bae J-W. Social status shapes the bacterial and fungal gut

605   communities of the honey bee. *Sci Rep* 2018; **8**.

606   50.   Anderson KE, Carroll MJ, Sheehan T, Mott BM, Maes P, Corby-Harris V. Hive-stored pollen

607   of honey bees: many lines of evidence are consistent with pollen preservation, not nutrient

608   conversion. *Mol Ecol* 2014; **23**: 5904–5917.

609   51.   Maurice CF, CL Knowles S, Ladau J, Pollard KS, Fenton A, Pedersen AB, et al. Marked

610   seasonal variation in the wild mouse gut microbiota. *ISME J* 2015; **9**: 2423–2434.

611   52.   Davenport ER, Mizrahi-Man O, Michelini K, Barreiro LB, Ober C, Gilad Y. Seasonal

612   Variation in Human Gut Microbiome Composition. *PLoS ONE* 2014; **9**: e90731.

613   53.   Smits SA, Leach J, Sonnenburg ED, Gonzalez CG, Lichtman JS, Reid G, et al. Seasonal

614   cycling in the gut microbiome of the Hadza hunter-gatherers of Tanzania. *Science* 2017; **357**:

615   802–806.

616   54.   Sonnenburg ED, Smits SA, Tikhonov M, Higginbottom SK, Wingreen NS, Sonnenburg JL.

617   Diet-induced extinctions in the gut microbiota compound over generations. *Nature* 2016;

618   **529**: 212–215.

619   55.   Martino ME, Joncour P, Leenay R, Gervais H, Shah M, Hughes S, et al. Bacterial Adaptation

620   to the Host's Diet Is a Key Evolutionary Force Shaping Drosophila-Lactobacillus Symbiosis.

621   *Cell Host Microbe* 2018; **24**: 109-119.e6.

622   56.   Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, et al. Diet

623   Drives Convergence in Gut Microbiome Functions Across Mammalian Phylogeny and

624   Within Humans. *Science* 2011; **332**: 970–974.

57. Ricigliano VA, Fitz W, Copeland DC, Mott BM, Maes P, Floyd AS, et al. The impact of

pollen consumption on honey bee (*Apis mellifera*) digestive physiology and carbohydrate

metabolism: RICIGLIANO et al. *Arch Insect Biochem Physiol* 2017; **96**: e21406.

58. Omholt SW, Amdam GV. Epigenetic Regulation of Aging in Honeybee Workers. *Sci Aging*

Knowl Environ 2004; **2004**: pe28.

59. Li H, Qi Y, Jasper H. Preventing Age-Related Decline of Gut Compartmentalization Limits

Microbiota Dysbiosis and Extends Lifespan. *Cell Host Microbe* 2016; **19**: 240–253.

60. Broderick NA, Buchon N, Lemaitre B. Microbiota-Induced Changes in Drosophila

melanogaster Host Gene Expression and Gut Morphology. *mBio* 2014; **5**: e01117-14.

61. Clark RI, Salazar A, Yamada R, Fitz-Gibbon S, Morselli M, Alcaraz J, et al. Distinct Shifts in

Microbiota Composition during Drosophila Aging Impair Intestinal Function and Drive

Mortality. *Cell Rep* 2015; **12**: 1656–1667.

62. Fahrenholz L, Lamprecht I, Schricker B. Thermal investigations of a honey bee colony:

thermoregulation of the hive during summer and winter and heat production of members of

different bee castes. *J Comp Physiol B* 1989; **159**: 551–560.

63. Gätschenberger H, Azzami K, Tautz J, Beier H. Antibacterial Immune Competence of Honey

Bees (Apis mellifera) Is Adapted to Different Life Stages and Environmental Risks. *PLOS*

ONE 2013; **8**: e66415.

64. Hystad EM, Salmela H, Amdam GV, Münch D. Hemocyte-mediated phagocytosis differs

between honey bee (Apis mellifera) worker castes. *PLOS ONE* 2017; **12**: e0184108.

65. Cristian R, Neagu A, Mihalache, Lazar V, Ecovoiu A. Molecular and bioinformatics analysis

of the relative expression profiles of dorsal, Toll-1, Relish and Duox genes in young versus

old diutinus workers of Apis mellifera. 2016.

66. Steinmann N, Corona M, Neumann P, Dainat B. Overwintering Is Associated with Reduced

Expression of Immune Genes and Higher Susceptibility to Virus Infection in Honey Bees.

PLOS ONE 2015; **10**: e0129956.

67. Crailsheim K. Dependence of protein metabolism on age and season in the honeybee (Apis

mellifica carnica Pollm). *J Insect Physiol* 1986; **32**: 629–634.

68. Kwong WK, Moran NA. Cultivation and characterization of the gut symbionts of honey bees

and bumble bees: description of Snodgrassella alvi gen. nov., sp. nov., a member of the

family Neisseriaceae of the Betaproteobacteria, and Gilliamella apicola gen. nov., sp. nov., a

member of Orbaceae fam. nov., Orbales ord. nov., a sister taxon to the order

'Enterobacteriales' of the Gammaproteobacteria. *Int J Syst Evol Microbiol* 2013; **63**: 2008–

2018.

69. Ellegaard KM, Tamarit D, Javelind E, Olofsson TC, Andersson SG, Vásquez A. Extensive

intra-phylotype diversity in lactobacilli and bifidobacteria from the honeybee gut. *BMC*

Genomics 2015; **16**: 284.

70. Sommer F, Ståhlman M, Ilkayeva O, Arnemo JM, Kindberg J, Josefsson J, et al. The Gut

Microbiota Modulates Energy Metabolism in the Hibernating Brown Bear Ursus arctos. *Cell*

Rep 2016; **14**: 1655–1661.

Figures and Figure Legends

Figure 1. Gut bacterial communities differ between foraging and winter season in a

single colony monitored over two years. (A) Monthly changes in the absolute abundance

of the seven phylotypes monitored over a period of two years depicted as mean values

(±SE) of the analyzed bees. The number of bees per month is indicated. Asterisk indicates

missing data for July 2015 due to extraction failure. **(B)** Number of cells per individual

honey bee gut in foragers (F) and winter bees (W) for each bacterial phylotype and their

sum. Mean values are shown as black horizontal lines. Only bees with detectable levels

were plotted (number of bees is given at the bottom of the plot; for prevalence see

Supplementary Figure S1). **(C)** Copy number of the 16S rRNA gene in gut samples of a

subset of the analyzed months (Apr 2015, Aug 2015, Oct 2015, Jan 2016, Apr 2016, Jul

2016, Oct 2016, Jan 2017, Apr 2017). **(D)** Relative community composition of the gut

microbiota of bees sampled in each month. Community composition is based on the

abundances of the seven monitored phylotypes **(E)** Shannon's diversity index calculated

from cell numbers of different bacterial phylotypes in foragers (F) and winter bees (W).

(F) Projection of the abundances of monitored phylotypes into first and second principle

components in all analyzed bees, together with correlation vectors representing variables

driving the separation on both axes. Permutation T-Test was used for pairwise

comparisons. ns, non-significant; ***, P < 0.001.

Figure 2. Gut community size and structure differ between foragers, nurses, and winter

bees across 14 colonies. Data from 20 pooled guts sampled from foragers (F), nurses (N)

and winter bees (W) across multiple hives (n=14). **(A)** Normalized cell numbers (see

methods) of the seven monitored phylotypes and their summed abundances. **(B)** 16S

rRNA gene copy numbers per sample. **(C)** Relative community composition of the gut

microbiota based on the abundances of the seven monitored phylotypes (left y-axis).

Black diamonds indicate the sum of the total abundances of seven monitored phylotypes

(right y-axis). Capital letters below the stacked bars indicate the hive of origin. **(D)**

Projection of the abundances of monitored phylotypes into first and second principle

components in the pooled samples. Each dot represents a different hive. Correlation

vectors represent different variables driving the separation on both axes. Black lines show

mean values. Different letters indicate groups that are significantly different from each

other based on Tukey's post-hoc tests following Permutation ANOVA. See **Supplementary**

Table S2 for full statistics results.

Figure 3. Diet is a major factor shaping the gut communities of honey bees. **(A-C)**

Dissected guts of foragers **(A)**, nurses **(B)**, and winter bees **(C)** consisting of the crop (c),

(missing in some samples), the midgut (m), the ileum (i), and the rectum (r) with attached

stinger and last abdominal tergite. **(D)** Wet gut weight of individual foragers (F), nurses

(N), and winter bees (W). **(E)** Spearman correlation between gut weight and bacterial

loads (assessed with universal bacterial 16S rRNA primers) across all bee types. The grey

area indicates the 95% confidence interval. **(F)** Wet gut weight of experimentally

colonized bees that were fed either sugar water (SW) or sugar water and pollen (SW+P).

(G) Spearman correlation between gut weight and bacterial loads (assessed with

universal bacterial 16S rRNA primers) across the two treatment groups of colonized bees.

(H) Total abundances of the seven monitored phylotypes in the two treatment groups of

colonized bees. Bees with loads below the detection limit of the qPCR method are shown

below the dashed line. Different letters indicate groups that are significantly different

from each other based on Tukey's post-hoc tests following Permutation ANOVA. Two-

group comparisons were done by Permutation T-Test. ns, non-significant; ***, $P < 0.001$.

Supplementary Material

Supplementary Figures

Figure S1. Prevalence of phylotypes that were not detect in every honey bee of the colony sampled over two years. Monthly prevalence (i.e. the proportion of bees in which the bacterial phylotype was detected in a given month) of **(A)** *Lactobacillus* Firm-4, **(B)** *B. apis* and **(C)** *F. perrara*. The targetted bacterium was considerred as absent when the qPCR signal was below the detection limit of the primer set (see Supplementary Table S1). The numbers in the upper and lower part of the bar plots indicate the number of bees in which the target is absent and present, respectively. Asterisk indicates missing data for July 2015 due to a sampling/extraction error.

733 **Figure S2. *F. perrara* loads correlate with scab presence.** *F. perrara* normalized number

734 of cells (see methods) in guts of foragers (F) and winter bees (W) with or without scab

735 phenotype in the pylorus region. Data points under the black dashed line correspond to

736 samples with no detectable level of *F. perrara* as based on the detection limit of our qPCR

737 method. Statistics: permutation T-Test.

Fig. S3 Monthly trends in the abundance of honey bee-specific bacteria.

Absolute abundance of the seven monitored phylotypes in individual foragers and winter bees
sampled over two years: Gilliamella, *S. alvi*, Bifidobacterium, Lactobacillus Firm-5, Lactobacillus
Firm-4, *B. apis*, and *F. perrara*. The last plot shows the summed abundance of each bee. Only bees
with detectable levels were considered. Bees with loads below the detection limit of the qPCR
method are shown below the dashed line. Asterisk indicates missing data for July 2015 due to a
sampling/extraction error.

Supplementary Tables

Target (gene name and accession no.)	Sequence (5' to 3')	Tm (°C)[a]	Amplicon size	# Gene loci per genome[b]	Standard curve Efficiency, R²	Slope, intercept	LOD Cq[c] (# copies)	Reference
Apis mellifera (*actin*, AB023025)	**F:** TGCCAACACTGTCCTTTCTG **R:** AGAATTGACCCACCAATCCA	58.4 56.4	156 bp	-	1.896 = 89.6 %, 1.0	-3.6, 37.579	33.726 (10)	[1]
Gilliamella (*16S rRNA*, JQ936674)	**F:** CTTTGTTGCCATCGGTTAGGCC **R:** CCGCTTGCTCTCGCGAGG	64.2 62.9	160 bp	4	1.86 = 86 %, 0.997	-3.709, 40.807	37.026 (10)	[2]
Frischella perrara (*16S rRNA*, JX8783306)	**F:** GGAAGTTATGTGTGGGATAAGC **R:** CTATTCTCAGGTTGAGCCCG	60.1 60.5	185 bp	4	1.946 = 94.6 %, 0.995	-3.457, 38.042	31.704 (100)	[3]
Snodgrassella alvi (*16S rRNA*, JQ746651)	**F:** CTTAGAGATAGGAGAGTGCCTT **R:** AACTTAATGATGGCAACTAATGACAA	60.1 60.1	132 bp	4	1.966 = 96.6 %, 0.999	-3.406, 38.236	31.256 (100)	[4], modified
Bartonella apis (*16S rRNA*, KP987885)	**F:** GTGGGAATCTACCTATTTCTACG **R:** AACGCGGGCTCATCTATCTC	60.9 60.5	103 bp	2	2.051 = 105.1 %, 0.997	-3.205, 36.771	30.752 (100)	[3]
Bifidobacterium (*16S rRNA*, AB437355)	**F:** ATGCAAGTCGAACGGGATCC **R:** CATCCCATRCCGGTAAACCC	60.5 60.5 – 62.5	174 bp	2	1.948 = 94.8 %, 0.997	-3.453, 38.983	34.906 (10)	[3]
Lactobacillus Firm-4 (*16S rRNA*, DQ837632)	**F:** AGTCGAGCGCGGGAAGTCA **R:** AGCCGTCTTTCAACCAGCACT	61.6 61.2	169 bp	4	1.912 = 91.2 %, 0.999	-3.554, 37.196	29.966 (100)	[3]
Lactobacillus Firm-5 (*16S rRNA*, JX099547)	**F:** GCAACCTGCCCTWTAGCTTG **R:** GCCCATCCTKTAGTGACAGC	60.5 60.5 – 62.5	118 bp	4	2.144 = 114.4 %, 0.998	-3.02, 36.306	29.94 (100)	[3]
Universal bacteria (*16S rRNA*)	**F:** AGGATTAGATACCCTGGTAGTCC **R:** YCGTACTCCCCAGGCGG	62.9 60 – 62	variable	-	1.946 = 94.6 %, 0.993	-3.457, 37.986	33.546 (10)	[3]

Table S1. Primers used in this study and standard curve characteristics. [a]Melting temperatures were calculated with the online tool described in [5]. [b]The number of loci of the target gene per genome of the corresponding organism was assessed based on JGI database annotations. [c]*LOD* refers to the limit of detection of primers sets, expressed here as the lowest number of plasmid copies reliably detected by qPCR when standard curves were performed.

Supplementary References

1. Zufelato MS, Lourenço AP, Simões ZLP, Jorge JA, Bitondi MMG. Phenoloxidase activity in Apis mellifera honey bee pupae, and ecdysteroid-dependent expression of the prophenoloxidase mRNA. *Insect Biochem Mol Biol* 2004; **34**: 1257–1268.

2. Engel P, Bartlett KD, Moran NA. The Bacterium Frischella perrara Causes Scab Formation in the Gut of its Honeybee Host. *mBio* 2015; **6**: e00193-15.

3. Kešnerová L, Mars RAT, Ellegaard KM, Troilo M, Sauer U, Engel P. Disentangling metabolic functions of bacteria in the honey bee gut. *PLOS Biol* 2017; **15**: e2003467.

4. Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, Moran NA. A simple and distinctive microbiota associated with honey bees and bumble bees. *Mol Ecol* 2011; **20**: 619–628.

5. Kibbe WA. OligoCalc: an online oligonucleotide properties calculator. *Nucleic Acids Res* 2007; **35**: W43–W46.

Supplementary Table S2

Statistics for data presented in Fig 1

Permutation ANOVA (Month Effect)

Response Variable	P-val
Universal 16s	1.00E-04
Ba	1.00E-04
Bi	1.00E-04
F4	1.00E-04
F5	1.00E-04
Fp	1.00E-04
Ga	1.00E-04
Sa	1.00E-04

Permutation T-Test (F vs W)

Response Variable	P-val
Sum	1.00E-04
Universal 16s	1.00E-04
Ba	1.00E-04
Bi	1.00E-04
F4	1.00E-04
F5	1.00E-04
Fp	1.00E-04
Ga	0.6572
Sa	6.00E-04

Regression for general trends During foraging season

2015 foraging season	slope	r^2	p-val
Ba	0.02881199	0.0049618	0.3652
Bi	0.02773742	0.0153763	0.0943
F4	0.058836	0.070771	5.00E-04
F5	0.08169238	0.1790406	1.00E-04
Fp	-0.01085094	0.0006541	0.7533
Ga	0.01323004	0.0063768	0.2792
Sa	0.006376515	0.0008233	0.6935
2016 foraging season			
Ba	0.0412078	0.0085632	0.2464
Bi	0.0154952	0.0050785	0.3637
F4	0.0861126	0.0608465	0.0014
F5	0.007147429	0.0006722	0.7411
Fp	-0.08212353	0.0207844	0.1037
Ga	-0.1530585	0.2926288	1.00E-04
Sa	0.01242151	0.0018879	0.5773

Alpha-Diversity: Permutation T-Test on Shannon's Diversity Index

F vs W	1.00E-04

Multivariate Analysis of Variance (MANOVA) Wilk's Lambda

	Df	Wilks	Approx F	Num Df	Den Df	P-Val
Bee type	1	0.58162	57.342	7	558	< 2.2e-16
Residulas	564					

Statistics for data presented in Fig 2

Permutation One-Way ANOVA (F vs W vs N)

Response Variable	P-val
Sum	1.00E-04
Universal 16S	1.00E-04
Ba	1.00E-04
Bi	1.00E-04
F4	1.00E-04
F5	1.00E-04
Fp	2.00E-04
Ga	1.00E-04
Sa	5.52E-01

Tukey's Post Hoc Test

Response Variable	P-Val N vsF	P-Val W vs F	P-Val W vs N
Sum	1.33E-05	3.78E-10	0.00561
Universal 16S	1.66E-06	5.10E-09	0.224
Ba	0.039604	1.02E-07	0.000376
Bi	4.99E-07	2.67E-10	0.0602
F4	1.04E-07	5.42E-07	1.00E+00
F5	8.66E-09	1.08E-12	0.0118
Fp	1.00E+00	3.36E-03	0.00112
Ga	1.05E-07	4.33E-08	1.00E+00
Sa	1.00E+00	1.00E+00	0.957

Multivariate Analysis of Variance (MANOVA) Wilk's Lambda

	Df	Wilks	Approx F	Num Df	Den Df	P-Val
Bee type	2	0.13955	7.9056	14	66	1.70E-09
Residulas	39					

Statistics for data presented in Fig 3

One-Way Anova (F vs N vs W) on gut weight			
	Df	F-value	P-value
Bee type	2	24,128	1.21E-08
residuals	66		
Tukey's Post Hoc Test			
	N vs F	W vs F	W vs N
P-value	1.14E-06	6.81E-08	1

Permutation T-Test Experimental Bees	
Response Variable	P-val
Universal 16s	3.00E-04
Ba	2.00E-02
Bi	2.00E-04
F4	1.00E-04
F5	1.00E-04
Fp	1.00E-04
Ga	2.00E-04
Sa	2.85E-01

Welch's T-test on gut weight of experimental bees	
t	9.43
Df	38.57
P-value	1.45E-11

Statistics for data Presented in Fig S2

Permutation T-Test onF. perrara abundance in the presence and absebce of scab	
P-Value	1.00E-04

Statistics for data Presented in Fig S2

Permutation T-Test onF. perrara abundance in the presence and absebce of scab	
P-Value	1.00E-04