Recovirus NS1-2 has viroporin activity that induces aberrant cellular calcium signaling to facilitate virus replication

Alicia C. Strtak[1], Jacob L. Perry[1], Mark N. Sharp[1,4], Alexandra L. Chang-Graham[1], Tibor Farkas[2,3], and Joseph M. Hyser[1*]

[1]Alkek Center for Metagenomic and Microbiome Research and Department of Molecular Virology and Microbiology, Baylor College of Medicine, Houston, TX 77303. [2]Department of Pathobiological Sciences, Louisiana State University School of Veterinary Medicine, and [3]Louisinan Animal Disease Diagnostic Laboratory, Baton Rouge, LA 70803. [4]Texas Medical Center Summer Research Internship Program, Augustana College, Rock Island, IL 61201.

Correspondence should be sent to:
Joseph M. Hyser
Alkek Center for Metagenomic and Microbiome Research
Department of Molecular Virology and Microbiology
MS: BCM385
Baylor College of Medicine
Houston, TX, 77303
PH: 713-798-4514
FAX: 713-798-3586
Joseph.Hyser@bcm.edu

No conflicts of interest exist.

Abstract

Enteric viruses in the *Caliciviridae* family cause acute gastroenteritis in humans and animals, but the cellular processes needed for virus replication and disease remain unknown. A common strategy among enteric viruses, including rotaviruses and enteroviruses, is to encode a viral ion channel (i.e., viroporin) that is targeted to the endoplasmic reticulum (ER) and disrupts host calcium (Ca^{2+}) homeostasis. Previous reports have demonstrated genetic and functional similarities between the nonstructural proteins of caliciviruses and enteroviruses, including the calicivirus NS1-2 protein and the 2B viroporin of enteroviruses. However, it is unknown whether caliciviruses alter Ca^{2+} homeostasis for virus replication or whether the NS1-2 protein has viroporin activity like its enterovirus counterpart. To address these questions, we used Tulane virus (TV), a rhesus enteric calicivirus, to examine Ca^{2+} signaling during infection and determine whether NS1-2 has viroporin activity that disrupts Ca^{2+} homeostasis. We found that TV disrupts increases Ca^{2+} signaling during infection and increased cytoplasmic Ca^{2+} levels is important for efficient replication. Further, TV NS1-2 localizes to the endoplasmic reticulum (ER), the predominant intracellular Ca^{2+} store and the NS2 region has characteristics of a viroporin domain (VPD). NS1-2 had viroporin activity in a classic bacterial functional assay and caused aberrant Ca^{2+} signaling when expressed in mammalian cells, but truncation of the VPD abrogated these functions. Together, our data provide new mechanistic insights into the function of the NS2 region of NS1-2 and show that like many other enteric viruses, enteric caliciviruses also exploit host Ca^{2+} signaling to facilitate their replication.

Importance

Tulane virus is one of many enteric caliciviruses that cause acute gastroenteritis and diarrheal disease. Globally, enteric caliciviruses affect both humans and animals and result in >65 billion

dollars per year in treatment and healthcare-associated costs, thus imposing an enormous

economic burden. Recent progress has resulted in several cultivation systems (B cell, enteroid

and zebrafish larvae) to study human noroviruses, but mechanistic insights into the viral factors

and host pathways important for enteric calicivirus replication and infection are largely still

lacking. Here we used Tulane virus, a calicivirus that is biologically similar to human

noroviruses and can be cultivated in conventional cell culture, to identify and functionally

validate NS1-2 as an enteric calicivirus viroporin. Viroporin-mediated calcium signaling may be

a broadly utilized pathway for enteric virus replication, and its existence within caliciviruses

provides a novel approach to developing antivirals and comprehensive therapeutics for enteric

calicivirus diarrheal disease outbreaks.

Introduction

The *Caliciviridae* family consists of small, non-enveloped single-stranded RNA viruses with five major genera: *Sapovirus, Lagovirus, Vesivirus, Nebovirus,* and *Norovirus* (1, 2). Caliciviruses infect a wide-array of hosts and have importance in medical, veterinary, and agricultural fields (3). Of particular importance are human noroviruses (HuNoVs) that are the leading cause of acute gastroenteritis (AGE) in every age group, and can cause life-threatening illness in the young, immunocompromised, and elderly subpopulations (4–7). Estimates show that every individual will experience at least five symptomatic norovirus infections in their life (8), which underlines the need for anti-viral drugs, vaccines, or anti-diarrheal therapies for HuNoV infection (9, 10). However, many aspects of calicivirus pathogenesis, including that of HuNoV, remain uncharacterized, which represents a challenge to developing effective therapies (4, 6, 9). One strategy to address this challenge is to study other enteric caliciviruses, such as porcine sapoviruses and rhesus enteric caliciviruses (*Recovirus*). Recoviruses are a new proposed genus of enteric CVs initially identified in the stool from rhesus macaques, of which Tulane virus (TV) is the prototype strain (11, 12). While Recoviruses constitute a separate genus, these viruses are most closely related to HuNoVs and studies of TV show that it retains both biologic and genetic similarities to HuNoVs, including genomic organization, tissue tropism (intestinal epithelia), and clinical presentation (self-limiting vomiting and diarrhea) (1, 11–13). Furthermore, TV robustly replicates in cell culture in monkey kidney cell lines (e.g., LLC-MK2 cells), which facilitates investigation into the host pathways exploited by TV during infection. This makes TV an excellent model system to identify host signaling pathways broadly exploited by caliciviruses for replication and pathogenesis.

Like other caliciviruses, TV has 3 main open reading frames (ORFs), with ORF1

encoding the nonstructural proteins (NS1-7), and ORFs2-3 encoding the capsid proteins VPI

(ORF2) and VP2 (ORF3) (10, 14-15). During replication ORF1 is synthesized into the

polyprotein, which is subsequently cleaved by the viral protease NS6 to produce six

nonstructural proteins that orchestrate viral replication (4, 12, 14–16). In particular, the roles of

the NS1-2 protein (N-terminal protein) plays in viral replication and pathogenesis are not well-

characterized. Recent reports have identified that murine norovirus NS1, the N-terminal portion

of NS1-2, antagonizes the interferon pathway, but studies of full-length NS1-2 or the NS2

domain remain limited (17–19). Recombinant expression of NS1-2 from feline calicivirus

(FCV), murine norovirus (MNV), and HuNoV GII.4 shows that the protein traffics to the

endoplasmic reticulum (ER), concentrates perinuclearly, colocalizes with the ER-resident protein

calnexin, and contains C-terminal hydrophobic sequences (20–22). In contrast, Norwalk virus

(GI.1) NS1-2 (p48) was primarily found in the Golgi apparatus and implicated in disrupting ER-

to-Golgi trafficking (23, 24). The similarities in ER/Golgi membrane association and domain

organization of NS1-2 from different viruses suggest that NS1-2 may have a conserved function

among caliciviruses.

The ER, and to a lesser extent Golgi apparatus, are important intracellular calcium (Ca^{2+})

storage organelles, with the ER Ca^{2+} concentration being up to 1 mM (25, 26). As a ubiquitous

secondary messenger, Ca^{2+} is at the epicenter of many cellular processes and host machinery

tightly regulates Ca^{2+} levels to ensure low (nM) cytoplasmic Ca^{2+} concentrations at cellular rest

(26–32). Importantly, Ca^{2+} signaling regulates several aspects of viral life cycles including entry,

genome replication, and release (30, 33–35). To exploit Ca^{2+} signaling, many viruses express an

ion channel (i.e., viroporin) to dysregulate Ca^{2+} homeostasis in order to usurp Ca^{2+}-dependent

host processes (30, 36–39). The best characterized Ca^{2+} disrupting viroporins are the

nonstructural protein 4 (NSP4) from rotavirus (RV) and the 2B nonstructural protein of

enteroviruses (EV) and some other picornaviruses (36, 37, 40–44). Like all *bona fide* viroporins,

NSP4 and 2B have canonical biophysical motifs, including being oligomeric, having an

amphipathic α-helix that forms the pore, and a cluster of basic residues that facilitate membrane

insertion (37–39, 42–44). While no study has specifically looked at whether caliciviruses

dysregulate Ca^{2+} signaling or have a viroporin, they belong to the picornavirus-like superfamily

of positive-sense RNA viruses, among which there is considerable positional homology of the

cognate proteins of the nonstructural polyprotein (23, 45, 46). Within this rubric, the

picornavirus 2AB region constitutes the positional homolog of the calicivirus NS1-2 protein, and

several sequence motifs in the NS1 are conserved in the 2A protein of some picronaviruses (23).

While no functional homology between EV 2B and the NS2 region of NS1-2 has been yet been

identified, it is tempting to speculate that NS1-2 may have viroporin activity and dysregulate

host Ca^{2+} signaling analogous to that of EV 2B proteins.

In this study, we investigated the role of Ca^{2+} signaling in TV replication and whether TV

NS1-2 has viroporin activity that can dysregulate Ca^{2+} homeostasis. Using long-term live-cell

Ca^{2+} imaging, we sought to determine whether TV infection causes aberrant Ca^{2+} signaling

during infection and identify the cellular Ca^{2+} pools critical for the TV-induced Ca^{2+} signaling.

Finally, we tested TV NS1-2 for viroporin activity and determined whether the putative NS1-2

viroporin domain caused aberrant Ca^{2+} signaling similar to TV infection.

Methods

Cell lines, GECI lentiviruses, and viruses: All experiments were performed in LLC-MK2

cells. Lentivirus packaging and recombinant protein expression for western blot lysate

production was performed in HEK293FT cells (ATCC CRL-3216). Cell lines were grown in

high-glucose DMEM (Sigma: D6429) containing 10% fetal bovine serum (FBS) (Corning lot no.

35010167) and antibiotic/antimycotic (Invitrogen), and maintained at 37ºC with 5% CO_2.

Lentivirus packaging in HEK293FT cells was performed as previously described (41). Briefly,

LLC-MK2 cells were transduced with a lentivirus vector encoding GCaMP6s 1-day post-seeding

(~85% confluency). We confirmed positive expression of GCaMP6s at 48-72 hours post-

transduction, then passaged cells 1:2 and added hygromycin (100 μg/mL) for selection of the

LLC-MK2-GCaMP6s cell lines, henceforth referred to as MK2- G6s. We determined GCaMP6s

activity and dynamic range using thapsigargin (0.5μM). Tulane virus (TV) stocks were made in-

house by infecting cells with MOI 0.01 and harvesting at ~95% cytopathic effect (CPE). Virus

titer was determined by plaque assay. Irradiated virus controls were made by gamma-irradiating

TV stocks for 19 hours.

Replication assays: LLC-MK2 cells were seeded at 125,000 cells/well in 24 well plates (Costar

3524, Corning) and inoculated next day with TV at MOI 1 for 1 hour. Inoculum was removed,

and cell media was replaced containing different extracellular Ca^{2+} conditions (0 mM Ca^{2+}, 4

mM Ca^{2+}), intracellular Ca^{2+} chelator 50 μM BAPTA-AM, or the sarco/endoplasmic reticulum

calcium ATPase (SERCA) blocker thapsigargin (TG). Ca^{2+}-free DMEM was purchased from

Gibco (Cat #21068-028). Standard high-glucose DMEM (Sigma) has 1.8 mM $CaCl_2$, which we

refer to as "2 mM Ca^{2+}, and media with 4 mM Ca^{2+} was made by adding 2 mM $CaCl_2$ to the

standard high-glucose DMEM (Sigma). We maintained TV-infected cells under these conditions

until the positive control (normal media) had ~90% CPE. Progeny virus was harvested by 3

freeze/thaw cycles, the virus yield was determined by plaque assay. For plaque assays, cells are

seeded at 75,000 cells/well in 24 well plates and 2-days post-seeding, the cells were inoculated

for 1 hour with 10-fold serial dilutions of the sample. Then, we removed the inoculum and added

the overlay. Overlays for plaque assays was made mixing equal parts 1.2% Avicel (FMC

Corporation) and 2X DMEM (Gibco). Plaque assays were harvested at 72 hours; fixed and

stained with crystal violet (3% solution) to visualize plaques. Titer is represented as plaque

forming units per milliliter (PFU/mL).

One-step growth curves: One step growth curves for TV were performed using a modified

protocol from previous reports (11, 15). Briefly, LLC-MK2 cells were inoculated with TV at

MOI 1 in serum-free DMEM (0% FBS DMEM). At 1-hour post-infection (HPI), the inoculum

was removed and replaced with 0% FBS DMEM. Cells were harvested at 0, 4, 6, 8, 10, 12, 16,

20, 24, and 28 HPI and virus yield determined by plaque assay. Each biological replicate was

performed in duplicate.

Long-term Ca^{2+} imaging experiments: Calcium imaging experiments were set up by adapting

a protocol detailed in previous reports (47). MK2-G6s cells were seeded at 78,500 cells/well in

15 μ-slide 8 well chambers (Ibidi, Germany) and infected the next day with TV at the indicated

MOI. After one hour, the inoculum was removed and replaced with FluoroBrite DMEM (Gibco).

For studies involving pharmacological compounds, the FluoroBrite was mixed with DMSO

(0.1%, vehicle control) or the indicated pharmacological compounds dissolved in DMSO. Then

the slide was mounted into a stage-top environmental chamber (Okolab H301-Mini) that

maintained 37°C with humidity control and 5% CO_2. Time lapse live-cell Ca^{2+} imaging

experiments were conducted for from ~2 HPI until ~18-24 HPI on a Nikon TiE epifluorescence

microscope using a SPECTRAX LED light source (Lumencor) and a 20X Plan Apo (NA 0.75)

objective. Images were acquired at 1-2 images/minute. Images were acquired and analyzed using

the NIS elements advanced research software package (Nikon). Prior to image analysis,

background camera noise was subtracted from the images using an averaged file of 10 no-light

camera images. Cells that underwent division during the imaging run were excluded from

analysis. Intracellular Ca^{2+} signaling over time was quantified by calculating the number of Ca^{2+}

spikes per cell. This was determined as follows: Raw fluorescence intensity values were

measured in Nikon software, then exported to Microsoft Excel to normalize the fluorescence to

the first image (F/F_0). The Ca^{2+} spikes were calculated by subtracting each normalized

fluorescence measurement from the previous measurement to determine the change in GCaMP6s

fluorescence (ΔF) between each timepoint. Ca^{2+} signals with a ΔF magnitude of >5% were

counted as Ca^{2+} spikes. For each condition testes, Ca^{2+} spikes in ≥30 cells were determined.

Heatmap generation: To generate heatmaps of the normalized GCaMP6s fluorescence over

time for long-term Ca^{2+} imaging experiments we used the TidyR (48) and ggplot2 (49) packages

available through R studio. Normalized GCaMP6s data from Excel was used to create a R-

compatible file (.csv) containing the normalized fluorescence and the acquisition time data for

the dataset, and imported the file into R. We used the TidyR package to organize data into a

format accessible by ggplot2. We then used ggplot2 to generate heatmaps.

Prediction of viroporin motifs *in silico*: We used the Hydropathy Analysis program at the

Transporter Classification Database to generate Kyte & Doolittle Hydropathy and Amphipathic

moment plots to identify putative viroporin motifs within full-length TV NS1-2 (50). Secondary

structure, membrane topology, and membrane integration predictions were performed using

PSIPred prediction analysis suite (website: http://bioinf.cs.ucl.ac.uk/introduction/) (51). Helical

wheel plots to identify clustered basic residues within the putative viroporin domain were

generated using the PepWheel analysis program at Heliquest (website:

http://heliquest.ipmc.cnrs.fr/) (52).

Expression vectors: *E. coli* expression constructs for the lysis assay were generated via ligation-

independent cloning (LIC) using the pET46-Ek/LIC kit (MilliporeSigma, Darmstadt, Germany).

The pET46-Ek/LIC constructs all have an N-terminal 6x histidine tag. Mammalian expression

vectors were generated by inserting c-myc tag and mRuby3 red fluorescent protein upstream of

full length NS1-2 and then subcloning this into the pTagRFP-N vector in place of TagRFP

(Epoch Life Sciences, Missouri City, TX). This construct will be referred to as RFP-NS1-2. The

NS1-2(Δ176) and NS1-2(Δ157) truncation mutations in both bacterial and mammalian

expression vectors were generated using the NEB Q5 site-directed mutagenesis kit (New

England Biolabs, Ipswich, MA). All constructs were sequence-verified using universal primers

specific to the construct backbone (GENEWIZ, South Plainfield, NJ). The mammalian

expression vector for EV 2B was generated by cloning the 2B from enterovirus 71 upstream into

pTagRFP-N and the construction of the NSP4-TagRFP expression vector was previously

described (53).

Transfection experiments: MK2-G6s cells were seeded in 15 μ-slide 8 well chambers (Ibidi,

Germany) and at 85% confluency transfected with mammalian expression constructs in Opti-

MEM (ThermoFisher) and Lipofectamine 2000 (Invitrogen). Transfection was optimized so cells

received 400 ng plasmid DNA and 0.5 μL of Lipofectamine 2000 per well. 10 μM trichostatin A

(TSA) was added from 1-3 hours post-transfection. TSA is a histone deacetylase (HDAC)

inhibitor used to increase expression from the vectors (54–56). Time-lapse Ca^{2+} imaging was

performed beginning 8 hours post-transfection to capture expression kinetics, and up to 24 hours

post-transfection to measure changes in Ca^{2+} signaling during expression of the RFP-tagged

proteins.

Deconvolution microscopy: LLC-MK2 cells were seeded in 15 μ-slide 8 well chambers (Ibidi,

Germany) and transfected one day prior to imaging. Cells were transfected with intracellular

markers for the plasma membrane (LCK-GFP; Addgene plasmid #61099), endoplasmic

reticulum (pLV-ER GFP; Addgene plasmid #80069), Golgi apparatus (pLV-Golgi GFP;

Addgene plasmid #79809), and mitochondria (HyPer-dMito, Envrogen). Control wells received

TagRFP (Envrogen), while experimental wells received either full length RFP-NS1-2, RFP-NS1-

2(Δ157) or RFP-NS1-2(Δ176). Cells were imaged 24 hours post-transfection on the DeltaVision

LIVE high resolution deconvolution microscope (GE Healthcare) using the 60X/1.4 Plan-Apo

NA oil objective (Olympus), and acquired using a pco.edge sCMOS_5.5 camera. Images were

acquired and deconvolved in SoftWoRx software. Upon deconvolving, images were further

processed in FIJI (ImageJ) to adjust for brightness/contrast and pseudo-coloring (57).

E. coli **Lysis Assay**: *E. coli* Lysis assays were performed as previously described (41). Briefly,

pET46-Ek/LIC constructs of the TV NS1-2 full-length and truncation mutants were transformed

into BL21(DE3)pLysS cells. Transformations were plated on LB + 1% glucose + 100μg/mL

ampicillin, + 35μg/mL chloramphenicol and grown at 37°C overnight. Isolated colonies were

picked next day and cultured overnight in liquid LB + 1% glucose, + 100μg/mL ampicillin, +

35μg/mL chloramphenicol at 37°C in an orbital shaker at 250 rpm. The next day, overnight

cultures were subcultured by 1:100 dilution into 200 mL LB + 1% glucose, + 100μg/mL

ampicillin, + 35μg/mL chloramphenicol. Subcultures were grown at 37°C in an orbital shaker at

250 rpm for ~3 hours to an OD_{630} between 0.3-0.5, then induced with 1mM IPTG. Absorbance

measurements at 630 nm (OD_{630}) were taken every 10 minutes for 90 minutes and normalized to

the induction OD_{630} to determine the percent growth or lysis over time after induction. Each

experiment was performed ≥3 times. Protein expression was determined by SDS-PAGE using a

4-20% Tris-glycine gel (Bio-RAD, Hercules, CA) and western blot for the 6x histidine tag. An

uninduced culture served as the negative control for viroporin activity and NS1-2 synthesis.

Membrane association experiment: Membrane association experiments were performed using

a modified protocol from previously reported experiments (36, 41). Lysed membranes from a

200 mL culture were centrifugated at 21,000 x g for 20 minutes and supernatants decanted.

Pellets were resuspended in PBS and sonicated 3 times for 1 minute on ice. Total lysate was

collected after sonication. Then membranes were pelleted by ultracentrifugation at 49,000 x g for

1 hour using a TLA-100.3 rotor in an Optima TL Ultracentrifuge (Beckman Coulter,

Indianapolis, IN), and the supernatant was collected for the soluble fraction. Finally, the

membrane fraction pellet was resuspended in PBS + 1% SDS to solubilize membrane proteins.

Samples from the total lysate, soluble fraction, and membrane fraction analyzed by western blot.

Production of TV and Vpg antisera: For the anti-TV antisera to detect VP1, adult male and

female CD-1 mice (purchased from Center for Comparative Medicine, Baylor College of

Medicine) were immunized 5 times with $CsCl_2$-gradient-purified TV at 10 μg/dose in AddaVax

adjuvant (InvivoGen). Immunizations were given at 3-week intervals. For the anti-Vpg antisera,

adult Balb/C were immunized 3 times with 10-20 μg/dose purified Vpg expressed in *E. coli*. The

priming dose was given in Freund's complete adjuvant and the subsequent boosts were given in

Freund's incomplete adjuvant. All experiments were performed in accordance with the

recommendations in the Guide for the Care and Use of Laboratory Animals of the National

Institutes of Health.

Immunoblot analysis: Samples were prepared using procedures adapted from (36). Briefly,

samples were mixed with 5X sample buffer containing 2-mercaptonethanol and boiled for 10

minutes at 100℃. Samples were then run on a 4-20% Tris-glycine gel (Bio-RAD, Hercules CA)

and transferred onto a nitrocellulose membrane using the Transblot Turbo transfer system (BIO-

RAD, Hercules, CA). To detect the bacterial constructs of NS1-2 and NSP4, we used the mouse

α-HIS Tag monoclonal antibody at 1:1000 (Genscript, Piscataway, NJ). To detect mammalian

expression constructs of NS1-2, we used the mouse α-c-Myc monoclonal (clone 9E10) antibody

at 1:1000 (R&D Systems, MN). To detect TV structural protein VP1, we used the mouse α-TV

polyclonal we made in-house by hyperimmunizing CD1 mice with purified TV particles. To

detect TV nonstructural protein Vpg, we used the mouse α-Vpg polyclonal antibody made by

hyperimmunizing mice with bacterially expressed and purified Vpg. For loading control of

mammalian cell lysates, we used the mouse α-GAPDH at 1:3000 (Novus Biologicals, CO). We

used alkaline phosphatase-conjugated goat α-mouse IgG at 1:2000 (Southern Biotech,

Birmingham, AL), and visualized using alkaline phosphatase substrate [Tris-base, nitro blue

tetrazolium (NBT), 5-bromo-4-chloro-3-indolyl phosphate (BCIP)].

Statistical analysis: Statistical analyses were completed using GraphPad Prism (ver. 7.03). Data

in this manuscript is presented as mean ± standard deviation. We performed columns statistics to

collect descriptive statistics and to determine the normality of the datasets. We then used the

unpaired Student's t-test for datasets with a parametric distribution, or a Mann-Whitney test for

datasets with a nonparametric distribution. Differences were determined statistically significant

if the $p < 0.05$. Authors had access to the data for this manuscript, and all authors approved the

final manuscript.

Data availability: RConsole code for the heatmaps generated in this paper is available upon

request.

Results

TV infection disrupts host calcium signaling kinetics in LLC-MK2 cells. Ca^{2+} is a ubiquitous

secondary messenger and many enteric viruses (*e.g.*, RVs and EVs) require elevated cytosolic

Ca^{2+} to facilitate replication (30, 36–39, 42, 43). To determine if TV causes aberrant Ca^{2+}

signaling like other enteric viruses, we examined whether Ca^{2+} signaling dynamics changed

during TV infection. We infected LLC-MK2 cells stably expressing GCaMP6s (MK2-G6s) with

different infectious doses [multiplicity of infection (MOI) 1, 5, 10] or γ-irradiated inactivated TV

and performed live-cell fluorescent microscopy during the infection. GCaMP6s is a GFP-based

genetically encoded Ca^{2+} indicator that reports changes in cytosolic Ca^{2+} as an increase in

fluorescence (58). TV-infected MK2-G6s cells show increased cytoplasmic Ca^{2+} levels (Fig. 1A)

beginning at roughly 8 HPI (MOI 10) and progressing for the remainder of the infection

(Supplemental Video S1). The observed increase in Ca^{2+} signaling coincides with the synthesis

of TV nonstructural proteins, assessed by western blot using an α-Vpg antisera (Fig. 1B) and TV

structural proteins, assessed by western blot using an α-TV antisera to detect VP1 (Fig. 1C),

which show increased TV protein production between 8-12 HPI. Further, based on a one-step

growth curve, the increased cytosolic Ca^{2+} also coincides with the onset of progeny virus

production, which occurs between 6-8 HPI (Fig. 1D). The increases in cytosolic Ca^{2+} were

dynamic during TV infection (Supplemental Video S2). In infected cells, we noted that changes

in cytosolic Ca^{2+} occurred through an increased number of discrete Ca^{2+} signals, much like what

we recently observed in RV-infected cells (Fig. 1E) (47). We refer to these high-amplitude,

transient Ca^{2+} signals as "Ca^{2+} spikes" and quantitated the number of Ca^{2+} spikes per cell during

infection. Compared to uninfected controls, TV-infected cells have significantly more Ca^{2+}

spikes/cell, but cells inoculated with γ-irradiated TV did not exhibit increased Ca^{2+} signaling

(Fig. 1F). Together, these data indicate that increased Ca^{2+} signaling requires replication

competent virus and occurs at later during infection, well after entry has occurred. Additionally,

Ca^{2+} signaling in infected cells increases in an infectious dose-dependent manner, saturating at

MOI=5 (Fig. 1F). To visualize the aberrant Ca^{2+} signaling induced by TV, we generated

heatmaps plotting normalized GCaMP6s fluorescence over time (Fig. 1G). Heatmap data shows

an increased number and magnitude of Ca^{2+} signals and that cytosolic Ca^{2+} levels change earlier

and more frequently throughout infection as infectious dose increases (Fig. 1G). The heatmaps

also show that MK2-G6s cells inoculated with γ-irradiated TV do not have increased Ca^{2+}

signaling compared to mock-inoculated cells (Fig. 1G), consistent with the lack of increased

Ca^{2+} spikes (Fig. 1F). Taken together, these data suggest that, like other enteric viruses, TV

disrupts host Ca^{2+} signaling kinetics during infection.

Intracellular Ca^{2+} is critical for TV replication. Since we observed aberrant Ca^{2+} signaling

during TV infection, we sought to determine whether Ca^{2+} was involved in TV replication. To

test this, we manipulated extracellular and intracellular Ca^{2+} levels and determined their effect

this had on TV yield. Doubling the extracellular Ca^{2+} concentration (~4 mM) did not affect TV

yield (Fig. 2A, right). In contrast, TV propagated in Ca^{2+}-free media significantly reduced total

yield (Fig. 2A, middle). Interestingly, plaques of TV propagated in Ca^{2+}-free media were

substantially smaller than that propagated in normal media, even though the plaque assay

titrations were performed in normal media (Fig. 2C). Next, to investigate the role of intracellular

Ca^{2+} during infection, we treated LLC-MK2 cells with BAPTA-AM, which chelates cytosolic

Ca^{2+} and therefore buffers cytosolic Ca^{2+} (59, 60). TV replication in Ca^{2+}-free media

supplemented with BAPTA-AM ($0Ca^{2+}$/BAPTA) was reduced up to 4-log (Fig. 2B), which was

a greater inhibition than Ca^{2+}-free media alone (Fig. 2A versus 2B). We next sought to determine

whether intracellular Ca^{2+} stores are important for TV replication by testing the effect of

thapsigargin (TG) on TV replication. TG is an inhibitor of sarco/endoplasmic reticulum Ca^{2+}

ATPase (SERCA), which pumps cytosolic Ca^{2+} into the ER to help maintain ER Ca^{2+} stores. We

treated TV-infected cells with TG and measured TV yield as above and found that TV

replication is ~3-log less in TG-treated than in DMSO-treated cells (Fig. 2B). Finally, we tested

these different manipulations of extracellular or intracellular Ca^{2+} on TV yield a different time

points during infection (8, 16, 24 HPI) (Fig. 2D). These studies confirmed that reduction of

extracellular Ca^{2+} or treatment with TG significantly inhibited total virus replication; however,

the rate of progeny virus production was not substantially reduced. Together, the replication

assays demonstrate that intracellular Ca^{2+} levels facilitate TV replication, and that the ER Ca^{2+}

store is particularly important for robust virus production.

TV-induced Ca^{2+} signaling requires ER Ca^{2+} stores. We next sought to determine the effect

that the manipulations to extracellular and intracellular Ca^{2+} had on the TV-induced Ca^{2+}

signaling exhibited in Figure 1. We altered extracellular and intracellular Ca^{2+} concentrations as

above and performed live Ca^{2+} imaging of mock and TV-infected MK2-G6s cells. TV-infected

cells in 2 mM Ca^{2+} (normal media) exhibited increased Ca^{2+} signaling, as observed above (Fig.

3A). Supplementing media with additional extracellular Ca^{2+} (4 mM Ca^{2+} total) did not further

increase the Ca^{2+} spikes but removing extracellular Ca^{2+} abolished the TV-induced Ca^{2+} spikes

(Fig. 3A). Using heatmaps, we plotted the relative change in GCaMP6s fluorescence over time

and observed increased signaling starting at ~8 HPI in both the 2 mM Ca^{2+} and 4 mM Ca^{2+}

conditions (Fig. 3B). Further, the heatmaps show that infected cells in Ca^{2+}-free media have a

signaling profile that phenotypically mimics uninfected controls (Fig. 3B). Like the results

obtained in replication assays, buffering cytoplasmic Ca^{2+} using BAPTA-AM reduced the

number of Ca^{2+} spikes per cell to a level comparable to that of mock-infected cells (Fig. 3C)

(Supplemental Video S2). Similarly, blocking the ER SERCA pump with TG significantly

reduces TV-induced Ca^{2+} signaling (Fig. 3D), supporting replication data and demonstrating that

ER Ca^{2+} stores are a critical source of Ca^{2+} for enhancing replication.

Tulane virus NS1-2 is targeted to the ER membrane. Our data indicates that TV activates

aberrant Ca^{2+} signaling involving the ER Ca^{2+} store, much like the dysregulation of Ca^{2+}

homeostasis by other enteric viruses observed in RV and EV infections. Both RV and EV encode

a viroporin, or viral ion channel, that targets the ER Ca^{2+} store to activate aberrant Ca^{2+} signaling

pathways that are critical for virus replication (35, 36, 38, 39, 42, 43). Viroporins are integral

membrane proteins that have some common characteristics, including being oligomeric, having

an amphipathic α-helix that serves as the channel lumen through the membrane, and a cluster of

basic amino acid residues that facilitate insertion into the membrane (24, 34, 35, 39, 61).

Previous work with NS1-2 from several different caliciviruses shows it is membrane-associated

and primarily to the ER (17–20, 24) and/or Golgi (17, 20, 21, 23). Thus, we hypothesized that

calicivirus NS1-2 could be a viroporin involved in the aberrant Ca^{2+} signaling we observed

during TV infection. Notably, the calicivirus NS2 domain is the positional homolog of the EV

2B viroporin (Fig. S1). This is potentially significant because previous studies have found

conserved functional characteristics between the positional homologs of the other nonstructural

proteins (20, 21, 23, 46, 61–64), and functional homology between EV 2AB and human

norovirus (HuNoV) GII.4 NS1-2 (20, 23). Additionally, when performing multiple sequence

alignments of other calicivirus NS1-2s, we found that the C-terminal domain (CTD) of this

protein is highly conserved (Fig. S2). To determine whether TV NS1-2 has viroporin-like

characteristics, we examined TV NS1-2 for viroporin motifs. First, we performed a Kyte-

Doolittle plot to detect hydrophobic regions of NS1-2 and an amphipathicity plot to identify

potential amphipathic domains (Fig. 4A). We found that aa195-215 (Fig. 4A, dark green box) in

the CTD of NS1-2 has a high amphipathic moment. We then used PSIPred (51) to model NS1-2

predicted secondary structure (Fig. 4B). Output from this analysis suggested that the NS1-2 CTD

was predominantly comprised of alpha-helices (Fig. 4B, pink residues), and accompanying

confidence scores for prediction of these C-terminal helices were ≥75% (Fig. S3). Interestingly,

the region of peak amphipathicity (Fig. 4A) was located within one of the PSIPred helix

predictions of the CTD (Fig. 4B, dark green bar) and contained clustered basic residues (blue

asterisks), two key features of viroporins. Additionally, NS1-2 topology modeling identified two

putative transmembrane domains (TMD): the first (TMD1) from aa164-179, and the second

(TMD2) from aa202-225 (Fig. 4C, top). The membrane topology schematic indicated that both

TMD1 and TMD2 had predicted pore-lining regions within their helices (Fig. 4B bottom left).

To explore this, we used HeliQuest (52) to generate a helical wheel diagram for TMD2 (aa198-

215), since TMD2 had the clustered basic residues common among viroporins. The helical wheel

shows that TMD2 is highly amphipathic with clear polar and non-polar faces to the helix (Fig.

4D). The calculated hydrophobic moment for TMD2 is 0.522, supporting the above

amphipathicity predictions (Fig. 4A). Given the results of these computational studies, we

predicted that NS1-2 TMD2 (aa195-215) is a viroporin domain and set out to test this prediction

experimentally.

First, we tested whether TV NS1-2 was an integral membrane protein and whether it

localized to the ER similar to NS1-2 from other caliciviruses. To do so, we generated bacterial

and mammalian expression vectors of full-length NS1-2. For mammalian expression vectors, we

N-terminally fused full-length NS1-2 to mRuby3 (henceforth referred to as RFP-NS1-2). From

these constructs, we generated two truncation mutants of WT NS1-2 in both mammalian and

bacterial expression vectors: the first, NS1-2 Δ176, was predicted to have TMD1 but lack the

viroporin domain, and the second, NS1-2 Δ157, was predicted to lack both TMD1 and the VPD.

We then transfected wild-type, full-length (WT) RFP-NS1-2, RFP-NS1-2 Δ157, and RFP-NS1-2

Δ176 into HEK 293FT cells and harvested cell suspensions next day. Samples following cell

lysis, sonication, and fractionation were collected for SDS/PAGE western blots. We found both

Δ176 and WT TV NS1-2 in the total fraction (T) and membrane pellets (M), but not in the

supernatant (S), suggesting that TMD1 mediates membrane association (Fig. 4E). Additionally,

in the non-reducing, unboiled conditions used, oligomers of both both Δ176 and WT RFP-NS1-2

were detected by western blot (Fig 4E, arrowheads). Similar results were obtained from

membrane fractionation of analogous bacterially expressed NS1-2 constructs (Fig. S4). Using the

mammalian expression vectors of RFP-NS1-2, we performed colocalization assays with

fluorescent markers of the ER, Golgi apparatus, and mitochondria. RFP-NS1-2 showed no

colocalization with the Golgi or mitochondria (Fig. 4F). In contrast, RFP-NS1-2 strongly

colocalized with the ER-GFP marker (Fig. 4F), indicating that, like NS1-2 from other

caliciviruses and EV 2B and RV NSP4, TV NS1-2 traffics to the ER membrane.

TV NS1-2 has viroporin activity that disrupts Ca^{2+} signaling. Since our predictive modeling

suggested that NS1-2 met the biophysical requirements for a viroporin, and our live-cell Ca^{2+}

imaging data exhibited large changes in cytosolic Ca^{2+} during TV infection, we tested whether

NS1-2 has viroporin activity. We performed the *E. coli* lysis assay, which is a classical viroporin

functional assay, wherein viroporin expression by BL21(DE3)pLysS *E. coli* results in

permeabilization of the inner membrane, resulting in T7 lysozyme-mediated cell lysis (41). This

assay has been used to identify and initially characterize many viroporins (36, 65, 66). We

expressed full-length HisNS1-2 in BL21(DE3)pLysS cells and measured optical density (OD)

over time following protein induction with IPTG. For the lysis assay, strong viroporin activity is

characterized by large decreases in OD over time, whereas no viroporin activity is characterized

by increases in OD over time. Our results show that induced NS1-2 has strong viroporin activity,

similar to that of RV NSP4, our positive control for viroporin activity (Fig. 5A). We see no

changes in OD over time for uninduced NS1-2, indicating that HisNS1-2 viroporin activity

correlated with protein expression, detected by immunoblot for the 6xHis-tag (Fig. 5B). We then

asked whether recombinant expression of RFP-NS1-2 alone increases Ca^{2+} signaling in MK2-

G6s cells. To test this, we transfected MK2-G6s cells with mammalian expression vectors for

RFP-NS1-2 as well as RFP-NSP4 and RFP-EV 2B, our positive controls for viroporin-mediated

Ca^{2+} signaling. Expressing RFP-tagged viroporins in MK2-G6s cells significantly increases both

the number and amplitude of Ca^{2+} spikes. However, this was not observed in cells expressing

RFP alone, as illustrated by the representative single-cell traces (Fig. 5C) (Supplemental Video

S3). As above, we quantitated the number of Ca^{2+} spikes and confirmed that recombinant

expression of RFP-NS1-2 increased the number of Ca^{2+} spikes per cell approximately 2-fold,

similar to that of EV 2B and RV NSP4 (Fig. 5D). Taken together, our results demonstrate that

TV NS1-2 has viroporin activity in the lysis assay, similar to *bona fide* viroporins, and causes

aberrant host Ca^{2+} signaling when expressed in mammalian cells.

NS1-2 viroporin activity maps to the putative viroporin domain. Our computational studies

above identified a putative TV NS1-2 VPD from aa195-212. To determine whether the NS1-2

viroporin activity maps to this putative VPD, we generated C-terminal truncation mutants in

bacterial expression vectors with deletions after aa212 (A212-Δ), after aa194 (W194-Δ), or after

aa176 (D176-Δ) and characterized them in the lysis assay (Fig. 6A). We found that the A212-Δ

truncation (red) had strong lysis activity comparable to full-length NS1-2 (black) (Fig. 6B). In

contrast the D176-Δ truncation (blue) exhibited no lysis activity, comparable to uninduced NS1-

2 (grey) (Fig. 6B). Immunoblot analysis confirm that protein expression correlated with

viroporin activity, and that the impaired activity of W194-Δ was not due to lower expression

levels, since the expression was comparable to that of full-length and A212-Δ (Fig 6C). Since the

W194-Δ truncation (green) had impaired viroporin activity, this suggests that the VPD

functionally extends to aa177-212.

Next, we characterized truncation mutants for their activation of aberrant Ca^{2+} signaling

in MK2-G6s cells. Since recombinant expression of full-length RFP-NS1-2 induced aberrant

Ca^{2+} signaling (Fig. 5D), we tested whether truncating the putative viroporin domain alone

(Δ176) or both TMDs (Δ157) would compromise NS1-2-induced Ca^{2+} signaling (Fig. 6D). First,

we examined the subcellular distribution and expression levels of the constructs. While the full-

length and Δ176 truncation both appeared reticular, the Δ157 truncation had cytoplasmic

distribution, consistent with it lacking both TMDs (Fig. 6E). Immunoblot analysis shows that the

expression of both truncations was much greater than that of full-length NS1-2 (Fig. 6F, left

blots) and by loading less lysate we can better resolve the 2 kDa size difference in the Δ157 and

Δ176 truncations (Fig. 6F, right blots). Next, we examined whether these truncations could

induce Ca^{2+} signaling by long-term live Ca^{2+} imaging in MK2-G6s cells. Individual cell traces

illustrate that neither the $\Delta157$ nor the $\Delta176$ truncation dramatically increased Ca^{2+} signaling

similar to full-length RFP-NS1-2 (Fig. 6G). Quantitation of the Ca^{2+} spikes per cell showed that

while both truncations exhibited higher Ca^{2+} signaling than RFP alone (Fig 6G), the amplitude of

these spikes was significantly reduced compared to full-length RFP-NS1-2 (Fig. 6H). The

significant reduction in the number and amplitude of Ca^{2+} spikes/cell for both mutants highlights

the critical importance of an intact VPD for disrupting host Ca^{2+} signaling. Together this work

demonstrates that TV NS1-2 is an ER-targeted viroporin that induces aberrant Ca^{2+} signaling.

Discussion

As obligate intracellular pathogens, viruses are adept at exploiting host pathways to

facilitate replication. Viruses from many different taxonomic families activate aberrant Ca^{2+}

signaling because Ca^{2+} signals are used by all cells to regulate a vast array of cellular functions.

Therefore, this represents a powerful strategy to reconfigure host cell physiology via targeted

disruption of host Ca^{2+} homeostasis. The overarching goal of this study was to determine

whether dysregulation of Ca^{2+} signaling is a characteristic of caliciviruses and if this is due to the

production of a viroporin protein similar to picornaviruses. To address these questions, we

studied TV, as a model calicivirus, using a combination of live-cell Ca^{2+} imaging and other

classical techniques. The major new findings of this study are as follows: (i) TV infection causes

aberrant Ca^{2+} signaling that coincides with viral protein synthesis and replication, (ii) cellular

Ca^{2+} is critical for TV replication and buffering of cytosolic Ca^{2+} severely reduced viral yield,

(iii) TV NS1-2 has viroporin activity and dysregulates Ca^{2+} signaling in mammalian cells similar

to TV infection, and (iv) NS1-2 viroporin activity maps to a C-terminal integral membrane

viroporin domain and truncation of this domain abrogates the NS1-2-induced activation of Ca^{2+}

signaling. To our knowledge, these results are the first to show exploitation of Ca^{2+} signaling by

a calicivirus and identification of NS1-2 as a Ca^{2+}-disrupting viroporin. These findings further

extend the functional homology between the calicivirus nonstructural proteins and their

picornavirus positional homologs.

The exploitation of host Ca^{2+} signaling to facilitate virus replication is a common feature

of many viruses (30). Our finding that TV coopts Ca^{2+} signaling is consistent with previous

studies showing that elevated Ca^{2+} levels are important for picornavirus replication, especially

since caliciviruses and picornaviruses utilize a similar replication strategy (43). Similar to other

Ca^{2+}-disrupting viruses, TV also induces aberrant Ca^{2+} signaling peak of virus replication, many

hours after cell entry. This is consistent with the reduced virus yield in media with reduced

extracellular Ca^{2+} or treatments to buffer cytosolic Ca^{2+} (BAPTA-AM) or block refilling of ER

Ca^{2+} stores (TG). Further, as we recently reported for RV infection, the TV-induced increase in

cytosolic Ca^{2+} manifests as many discrete Ca^{2+} signals rather than a monophasic increase in Ca^{2+}

over the infection (47). This raises the question of what cellular pathways are activated by this

Ca^{2+} signaling and how they benefit TV replication. Both RV and EV have been shown to

exploit Ca^{2+} signaling to activate the biosynthetic early stages of autophagy, which facilitates

virus replication through rearrangement of cellular membranes to form replication complexes

(67). MNV infection of primary macrophages or RAW264.7 cell line activates autophagy, but, in

contrast to RV and EV, autophagy limits MNV replication (68). Thus, it remains to be

determined whether autophagy plays a role in calicivirus replication complex assembly or

whether Ca^{2+} signaling regulates autophagy activation during calicivirus infection. Further,

elevated Ca^{2+} signaling may serve to modulate cellular apoptotic responses. Strong monophasic

increases in cytosolic Ca^{2+} activate apoptosis through mitochondrial Ca^{2+} overload, but transient

and oscillatory Ca^{2+} fluxes serve as pro-survival signals (69). Activation of apoptosis has been

seen in norovirus and feline calicivirus-infected cells, and caspase activation is critical for

cleavage and release of MNV NS1 from NS1-2, which in turn modulates cellular innate immune

responses (19, 22). Thus, increased transient Ca^{2+} signaling may serve to counteract apoptosis

activation and help prolong cell viability to maximize virus replication.

Within the superfamily of picornavirus-like positive sense RNA viruses, there is

positional homology between the ORF1 nonstructural proteins of caliciviruses (and likely

astroviruses) and the P2-P3 nonstructural proteins of picronaviruses (23, 45, 46). We used this

framework to determine whether TV NS1-2 exhibited viroporin activity, since the positional

homolog, the picornavirus 2B protein, is a well-established Ca^{2+}-conducting viroporin (34, 44).

We found that TV NS1-2 has viroporin activity, similar to 2B and RV NSP4, and the viroporin

activity mapped to the integral membrane NS2 domains. Since both the N- and C-termini are

likely oriented in the cytoplasm, NS1-2 is classified as a Class II2 viroporin, similar to the

picornavirus 2B proteins (34, 44). This topology is supported by the cytosolic accessibility of the

NS1 domain and the need for the C-terminus to also be localized in the cytosol to enable

cleavage by the NS6 protease. This raises the question of whether NS1-2 viroporin activity is

conserved throughout the *Caliciviridae* family. Though 2B and NS1-2 lack appreciable primary

sequence homology, this is not surprising because viroporins, even from the same virus family,

often only share the common viroporin motifs (*i.e.*, having [i] an amphipathic α-helix, [ii] a

cluster of basic residues, and [iii] being oligomeric) (33, 38, 39). Among NS1-2 from different

caliciviruses, we found that these characteristic features are conserved, so we predict viroporin

activity of NS1-2 is a common function. Furthermore, since blunting cytosolic Ca^{2+} signaling

with BAPTA-AM reduced TV replication, blocking NS1-2 viroporin activity with mutations or

drugs should also reduce replication. This is supported by a previous study showing that

recombinant coxsackie B3 virus with mutations of the 2B viroporin exhibited significantly

impaired replication or were completely replication deficient (70). Analogous studies can be

done using the TV reverse genetics system once residues critical for viroporin activity can be

identified through mutagenesis screens of the TV NS1-2 viroporin domain we mapped in this

study.

The increased Ca^{2+} signaling observed in TV-infected cells is phenotypically similar to

that induced by recombinant expression of full-length NS1-2, but the Ca^{2+} signaling is abrogated

by truncation of the viroporin domain. Further, NS1-2 primarily localized to the ER, which is a

major intracellular Ca^{2+} storage organelle. Thus, our model predicts that NS1-2 directly releases

Ca^{2+} from the ER; however, it is likely that both NS1-2 and activation of host Ca^{2+} signaling

pathways contribute to the observed Ca^{2+} signals. Ca^{2+} signals from NS1-2 require it to directly

conduct Ca^{2+} and have a high enough conductance that the ER Ca^{2+} release event can be detected

by a fluorescent Ca^{2+} indicator, yet these unitary events are challenging to detect even for large

channels like the IP3R (71). Future studies using patch clamp electrophysiology are needed to

confirm that NS1-2 conducts Ca^{2+} and determine its conductivity. Nevertheless, based on the

similarities between NS1-2 and other Ca^{2+}-conducting viroporins, EV 2B and RV NSP4, NS1-2

viroporin activity would reduce ER Ca^{2+} levels and this in turn will activate host Ca^{2+} signaling

pathways. First, the moderately increased steady-state cytosolic Ca^{2+} levels could foster more ER

Ca^{2+} release by potentiating the IP3R Ca^{2+} release channel (72). Second, reduced ER Ca^{2+} levels

activate the store-operated Ca^{2+} entry (SOCE) pathway, wherein decreased ER Ca^{2+} levels

activate the ER Ca^{2+} sensing protein stromal interaction molecule 1 (STIM1). Activated STIM1

translocates to ER microdomains adjacent to the plasma membrane and opens Ca^{2+} influx

channels, like Orai1, to elevate cytosolic Ca^{2+} (31, 32). This Ca^{2+} influx, in concert with SERCA,

helps to refill ER stores for continued signaling.

HuNoV and human sapoviruses cause outbreaks of AGE and are a major cause of

foodborne illnesses. However, the molecular mechanisms of how these caliciviruses cause

vomiting and diarrhea, the chief symptoms of AGE, have not been characterized. The

dysregulation of Ca^{2+} signaling by TV may provide insights into the pathophysiology of enteric

caliciviruses. Both IP_3-mediated ER Ca^{2+} release and SOCE have been shown to activate

chloride secretion from epithelial cells (73, 74). Additionally, in studies of other viroporins, the

viroporin-induced elevated cytosolic Ca^{2+} induces cytoskeleton rearrangement, leading to

disassembly of tight junctions and loss of barrier integrity (39). Hyperactivation of chloride

secretion and loss of tight junctions would contribute to excess fluid secretion and diarrhea. In

our study we have shown that dysregulated Ca^{2+} signaling is a feature of calicivirus infection

using TV. Thus, future studies can further examine the role of aberrant Ca^{2+} signaling in

calicivirus pathophysiology using human intestinal enteroid cultures that support the replication

of many HuNoV strains (4).

In summary, we have shown that TV activates aberrant Ca^{2+} singling during infection,

and cellular Ca^{2+} is critical for robust TV replication. Further, we found that the NS2 domain of

the NS1-2 nonstructural protein is a viroporin that alone induces Ca^{2+} signaling similar to TV

infection. Together, these results indicate that NS1-2 is functionally analogous to EV 2B and RV

NSP4. While little is known about the function(s) of NS1-2, and particularly the NS2 domain of

NS1-2, the similarity with other Ca^{2+} conducting viroporins may provide a broader insight for

understanding NS1-2 functions. Finally, antiviral drugs against viroporins have been developed

for influenza M2 and HIV Vpu (34). Thus, the NS1-2 viroporin may be a viable antiviral drug

target against caliciviruses.

Primers:

Primer name	Sequence
pET-46 EK/LIC NS1-2 Fwd	GACGACGACAAGATGGATACGTCCATAGATTCTGTGTTATCTGACACCAGC
pET-46 EK/LIC NS1-2 Rev	GAGGAGAAGCCCGGTTACTGTGGGATCAGTGATGAGGTAAGAAATGTGATCAAACTGG
pET-46 EK/LIC NS1-2 D176 Rev	GACGACAAGCCCGGTTAATCTTTGACAATAAGGTACACGATCTCGAG
pET-46 EK/LIC NS1-2 W194 Rev	GAGGAGAAGCCCGGTTACCAATTGTGCCAAAGGTTC
pET-46 EK/LIC NS1-2 A212 Rev	GAGGAGAAGCCCGGTTAAGCAACCCTTGTCAGGAATG
pET-46 EK/LIC NS1-2 Truncations Fwd	GACGACGACAAGATGGATACGTCCATAGATTCTGTGTTATC
pET-46 EK/LIC NS1-2 Δ157 Rev	GAGGAGAAGCCCGGTTTATTTGGTCCCTCCGACG
pET-46 EK/LIC NS1-2 Δ176 Rev	GAGGAGAAGCCCGGTTTAATTGACATCTTTGACAATGAGG
pTagRFP-N mRuby3-NS1-2 Δ157 Q5 Mut. Fwd	CGGAGGGACCTAAAGCGGATGGAC
pTagRFP-N mRuby3-NS1-2 Δ157 Q5 Mut. Rev	GTCGTACCTATTTTCATTCCAATCTTGGACATGTGTG
pTagRFP-N mRuby3-NS1-2 Δ176 Q5 Mut. Fwd	CAAAGATGTCTAATGGGCAAAGATTTG
pTagRFP-N mRuby3-NS1-2 Δ176 Q5 Mut. Rev	ACAATGAGGTACACGATCTCGAG

Figure 1 TV infection disrupts host calcium signaling kinetics in LLC-MK2 cells.
A. Representative images at early (4 HPI), onset (8 HPI), and late (12 HPI) stages of mock (top)
and TV-infected (bottom) LLC-MK2 GCaMP6s cells. **B, C.** Western blots for nonstructural
protein Vpg (B) and structural protein VP1 (C) confirm that aberrant Ca^{2+} signaling in infected
cells coincides with both structural and nonstructural protein synthesis. **D**. One-step growth curve
for TV at a low MOI (1) shows that virus replication is concomitant with viral protein synthesis
(1B, C) and with changes in Ca^{2+} signaling (1A). **E.** Still from overlay of α-Vpg staining (red) onto

short (10 minute) continuous imaging runs of TV-infected cells (MOI=5) 12 HPI. Accompanying
Ca^{2+} cell traces (right) shows the dynamic increases in cytosolic Ca^{2+} in infected cells **F.** Compared
to mock, TV-infected cells have an increased number of Ca^{2+} spikes per cell that increases in an
infectious dose-dependent manner, saturating at MOI=5. **G.** Heatmap data suggests that Ca^{2+}
signaling increases with infectious dose, and that a higher MOI disrupts host Ca^{2+} signaling earlier
in infection and sustains this aberrant Ca^{2+} signaling throughout. Mock and irradiated have similar
heatmap profiles, suggesting that replication competent virus is required to drive these changes in
Ca^{2+} signaling. Data shown as mean $\pm$ SD. **p<0.001, ****p<0.0001.
621

Figure 2: Intracellular calcium is critical for TV replication. A. Buffering out extracellular
calcium hinders TV replication, significantly reducing the total plaque forming units (PFU). In
contrast, excess extracellular Ca^{2+} (4 mM Ca^{2+}, right) does not impact replication. **B.** Buffering
intracellular calcium reduces replication. Depleting ER calcium stores with the SERCA-inhibitor
thapsigargin and reducing cytoplasmic Ca^{2+} with BAPTA-AM significantly reduce TV infectious
yield (PFU/mL). **C.** Representative images of plaques under normal (2mM) Ca^{2+} conditions (top,
DMSO) and reduced Ca^{2+} (0 mM Ca^{2+}/Bapta-AM, TG), conditions (bottom). **D.** Partial one-step
growth curve data altering free intracellular (IC) and extracellular (EC) Ca^{2+}. TV replication is
stunted in Ca^{2+} free IC and EC conditions (0 mM Ca^{2+}, 0 mM Ca^{2+}/BAPTA-AM). Inhibiting ER
Ca^{2+} replenishment with thapsigargin also blunts replication, suggesting that IC Ca^{2+} stores are
critical for TV replication. Data shown as mean ± SD. *$p<0.01$, **$p<0.001$.

Figure 3: TV-induced Ca^{2+} signaling requires ER Ca^{2+} stores. A. Ca^{2+}-free media reduces
Ca^{2+} signaling in TV-infected cells, suggesting Ca^{2+} signaling is activated during infection. **B.**
TV infection in 0 mM Ca^{2+} phenocopies mock Ca^{2+} traces in heatmap data, suggesting that EC
Ca^{2+} facilitates TV infection. **C.** Intracellular Ca^{2+} chelator BAPTA-AM abrogated TV-induced
Ca^{2+} signaling. BAPTA-AM-treated TV-infected cells (light green) returns Ca^{2+} signaling to
uninfected levels (grey). **D.** Depleting ER Ca^{2+} with SERCA blocker thapsigargin (TG)
significantly reduces TV-induced Ca^{2+} signaling (pink), suggesting that ER Ca^{2+} stores are a key
source of Ca^{2+} leveraged during infection. Data shown as mean ± SD. ****p<0.0001.

A — Putative VPD; Hydropathy; Amphipathicity; Value; Residue Number

B — Helical wheel plot: aa198-215; Hydrophobic moment: 0.522; Basic resiudes; Polar resiudes; Nonpolar resiudes

B — Strand; Helix; Coil; *Basic residue; Putative VPD

C — Transmembrane Helix; Extracellular; Transmembrane Helix; Pore-lining Helix; Extracellular; ER Lumen; 163-178: Pore-lining; 201-225: Pore-lining; Membrane; Cytoplasmic; N-terminus; C-terminus

E — UI NS1-2; Δ157 NS1-2; Δ176 NS1-2; WT NS1-2; L T S M; moαc-myc

F — Mito; Golgi; ER; RFP; GFP; MERGE

Figure 4: Tulane virus NS1-2 is targeted to the ER membrane. **A.** Predictive modeling of TV NS1-2 reveals that is has essential features of *bona fide* viroporins. Kyte-Doolittle-hydropathy plots predict an amphipathic moment from aa195-212 (dark green bar), consistent with alpha-helical structure required for channel formation. **B.** PSIPred secondary structure algorithms predict the C-terminus of NS1-2 is helical in nature, with the putative viroporin domain (VPD) contained to helices. **C.** PSIPred membrane topology predictions suggest that NS1-2 has two transmembrane helices (grey cubes, top figure). PSIPred algorithms predicting transmembrane helices suggest NS1-2 transmembrane domains are pore-lining (bottom left) and propose a model of membrane insertion and orientation where the putative VPD (aa195-212) comprises the pore-lining helix (bottom right). **D.** Helical wheel plot generated from the NS1-2 amphipathic segment (dark green bar) show clustered basic residues (green asterisks) and a hydrophobic moment of 0.522 from aa198-215, coinciding with the putative VPD **E.** Mammalian expressed full-length RFP-NS1-2 and RFP NS1-2 Δ176 is membrane-associated, but RFP NS1-2 Δ157 is not. Both the total fraction (T) and membrane pellets (M) extracted with 1% SDS contain RFP-NS1-2 and Δ176, but centrifuged supernatant (S) does not, suggesting that RFP-NS1-2 and Δ176 are membrane-associated proteins. In contrast, the supernatant contains RFP-NS1-2 Δ157. Further, immunoblot assays run under non-reducing conditions show that full-length RFP-NS1-2 and Δ176 oligomerize (black arrowheads) **F.** Co-transfection experiments using intracellular markers for predominant intracellular Ca^{2+} stores mitochondria (mito), Golgi, and endoplasmic reticulum (ER) to determine whether TV NS1-2 associated with any intracellular organelle(s). Based on deconvolution microscopy data, RFP-NS1-2 localized to the ER (right), but not with the Golgi (middle). RFP-NS1-2 did not localize to the mitochondria (left) (N≥2).

Figure 5: V NS1-2 has viroporin activity that disrupts Ca^{2+} signaling in mammalian cells. A.
Inducing TV NS1-2 in the lysis assay strongly reduces optical density similar to rotavirus NSP4,
the positive control for viroporin activity. B. Western blot data to verify protein expression
during the lysis assay for TV NS1-2 (bottom) and RV NSP4 (top). **C, D.** Mammalian
recombinant RFP-NS1-2 increases the number (B) and amplitude (C, top row, right) of Ca^{2+}
spikes when transfected into cells similar to RV NSP4 and EV 2B, the viroporin controls for
these experiments. Data shown as mean ± SD from ≥8 fields-of-view. *p<0.01, ****p<0.0001.
675

Figure 6: NS1-2 viroporins mutants do not increase cytoplasmic Ca²⁺ A. Schematic of TV
NS1-2 C-terminal truncation mutants to functionally map the viroporin domain. **B.** In the lysis
assay, truncating the C-terminal domain to amino acid 212 (red) results in wild-type activity

(black), but truncating to W194 (green) impairs activity. Truncating to D176 (blue) abrogates
viroporin activity, suggesting the viroporin domain functionally spans from aa177-212. **C.**
Western blots verifying protein expression in the lysis assay. **D.** Schematic for the mammalian
C-terminal truncation mutant constructs. **E.** IF data for truncation mutants. The $\Delta157$ is
cytoplasmic (far right), whereas the $\Delta176$, which retains one transmembrane segment, is
membrane localized (middle). **F.** Western blot data confirm the $\Delta157$ and $\Delta176$ mutant
constructs. The left blot is run with 20 μL/well to visualize FL NS1-2, whereas the right blot is
run with 5 μL/well to resolve the size difference between the $\Delta157$ and 176 NS1-2 mutants. **G.**
Both the $\Delta157$ and $\Delta176$ truncation mutants have significantly less Ca^{2+} spikes/cell compared to
wild-type full length RFP-NS1-2. **H.** Compared to full length RFP-NS1-2, both the $\Delta157$ and
$\Delta176$ truncation mutants have a significant reduced Ca^{2+} spike amplitudes, resulting in a change
in cytosolic fluorescence (ΔF) that phenotypically mimics RFP alone. Data shown as mean $\pm$ SD
from ≥ 8 fields-of-view. **p<0.001, ****p<0.0001.
693

References

1. Wobus CE, Cunha JB, Elftman M, Kolawole AO. 2016. Animal Models of Norovirus
Infection, p. 397–415. *In* Svensson, L, Desselberger, U, Greenberg, HB, Estes, MK (eds.),
Viral Gastroenteritis: Molecular Epidemiology and Pathogenesis. Academic Press,
Cambridge, MA.

2. Kniel KE. 2014. The makings of a good human norovirus surrogate. Curr Opin Virol
4:85–90.

3. Desselberger U. 2019. Caliciviridae Other Than Noroviruses. Viruses 11:286.

4. Karst SM, Wobus CE, Goodfellow IG, Green KY, Virgin HW. 2014. Advances in
norovirus biology. Cell Host Microbe 15:668–680.

5. Li J, Predmore A, Divers E, Lou F. 2012. New Interventions Against Human Norovirus:
Progress, Opportunities, and Challenges. Annu Rev Food Sci Technol 3:331–352.

6. Pendu J Le, Rydell GE, Nasir W, Larson G. 2016. Chapter 3.3 - Human Norovirus
Receptors, p. 379–396. *In* Svensson, L, Desselberger, U, Greenberg, HB, Estes, MK
(eds.), Viral Gastroenteritis. Academic Press, Cambridge, MA.

7. Ettayebi K, Crawford SE, Murakami K, Broughman JR, Karandikar U, Tenge VR, Neill
FH, Blutt SE, Zeng X-L, Qu L, Kou B, Opekun AR, Burrin D, Graham DY, Ramani S,
Atmar RL, Estes MK. 2016. Replication of human noroviruses in stem cell-derived human
enteroids. Science (80-) 353:1387–1393.

8. Bartnicki E, Cunha JB, Kolawole AO, Wobus CE. 2017. Recent advances in
understanding noroviruses. F1000Research 6:79.

9. Richardson C, Bargatze RF, Goodwin R, Mendelman PM. 2013. Norovirus virus-like
particle vaccines for the prevention of acute gastroenteritis. Expert Rev Vaccines 12:155–
167.

10. Ramani S, Atmar RL, Estes MK. 2014. Epidemiology of human noroviruses and updates
on vaccine development. Curr Opin Gastroenterol 30:25–33.

11. Farkas T, Sestak K, Wei C, Jiang X. 2008. Characterization of a Rhesus Monkey
Calicivirus Representing a New Genus of Caliciviridae. J Virol 82:5408–5416.

12. Wei C, Farkas T, Sestak K, Jiang X. 2008. Recovery of Infectious Virus by Transfection
of In Vitro-Generated RNA from Tulane Calicivirus cDNA. J Virol 82:11429–11436.

13. Smits SL, Rahman M, Schapendonk CME, van Leeuwen M, Faruque ASG, Haagmans
BL, Endtz HP, Osterhaus ADME. 2012. Calicivirus from novel recovirus genogroup in
human diarrhea, Bangladesh. Emerg Infect Dis 18:1192–1195.

14. Goodfellow IG, Taube S. 2016. Calicivirus Replication and Reverse Genetics, p. 355–378.
In Svensson, L, Desselberger, U, Greenberg, HB, Estes, MK (eds.), Viral Gastroenteritis:
Molecular Epidemiology and Pathogenesis. Academic Press, Cambridge, MA.

15. Farkas T. 2015. Rhesus enteric calicivirus surrogate model for human norovirus
gastroenteritis. J Gen Virol 96:1504–1514.

16. Green KY. 2013. The Noroviruses, p. 949–979. *In* Knipe, D, Howley, P (eds.), Field's
Virology, 6th ed. Wolter Kluwer Health/Lippincott Williams & Wilkins, Philadelphia,
PA.

17. Bailey D, Kaiser WJ, Hollinshead M, Moffat K, Chaudhry Y, Wileman T, Sosnovtsev S
V., Goodfellow IG. 2010. Feline calicivirus p32, p39 and p30 proteins localize to the
endoplasmic reticulum to initiate replication complex formation. J Gen Virol 91:739–749.

18. Hyde JL, Mackenzie JM. 2010. Subcellular localization of the MNV-1 ORF1 proteins and
their potential roles in the formation of the MNV-1 replication complex. Virology
406:138–148.

19. Lee S, Liu H, Wilen CB, Sychev ZE, Desai C, Hykes BL, Orchard RC, McCune BT, Kim
K-W, Nice TJ, Handley SA, Baldridge MT, Amarasinghe GK, Virgin HW. 2019. A
Secreted Viral Nonstructural Protein Determines Intestinal Norovirus Pathogenesis. Cell
Host Microbe 1–13.

20. Urakova N, Frese M, Hall RN, Liu J, Matthaei M, Strive T. 2015. Expression and partial
characterisation of rabbit haemorrhagic disease virus non-structural proteins. Virology
484:69–79.

21. Doerflinger SY, Cortese M, Romero-Brey I, Menne Z, Tubiana T, Schenk C, White PA,
Bartenschlager R, Bressanelli S, Hansman GS, Lohmann V. 2017. Membrane alterations
induced by nonstructural proteins of human norovirusPLoS Pathogens.

22. Lee S, Wilen CB, Orvedahl A, McCune BT, Kim KW, Orchard RC, Peterson ST, Nice TJ,
Baldridge MT, Virgin HW. 2017. Norovirus Cell Tropism Is Determined by
Combinatorial Action of a Viral Non-structural Protein and Host Cytokine. Cell Host
Microbe 22:449-459.e4.

23. Fernandez-Vega V, Sosnovtsev S V, Belliot G, King AD, Mitra T, Gorbalenya A, Green
KY. 2004. Norwalk virus N-terminal nonstructural protein is associated with disassembly
of the Golgi complex in transfected cells. J Virol 78:4827–37.

24. Ettayebi K, Hardy ME. 2003. Norwalk virus nonstructural protein p48 forms a complex
with the SNARE regulator VAP-A and prevents cell surface expression of vesicular
stomatitis virus G protein. J Virol 77:11790–7.

25. Jing J, He L, Sun A, Quintana A, Ding Y, Ma G, Tan P, Liang X, Zheng X, Chen L, Shi
X, Zhang SL, Zhong L, Huang Y, Dong M-Q, Walker CL, Hogan PG, Wang Y, Zhou Y.
2015. Proteomic mapping of ER–PM junctions identifies STIMATE as a regulator of
Ca^{2+} influx SUPPLEMENTARY. Nat Cell Biol 17:1339–1347.

26. Raffaello A, Mammucari C, Gherardi G, Rizzuto R. 2016. Calcium at the Center of Cell
Signaling: Interplay between Endoplasmic Reticulum, Mitochondria, and Lysosomes.
Trends Biochem Sci 41:1035–1049.

27. Cui C, Merritt R, Fu L, Pan Z. 2017. Targeting calcium signaling in cancer therapy. Acta
Pharm Sin B 7:3–17.

28. Carafoli E, Krebs J. 2016. Why calcium? How calcium became the best communicator. J
Biol Chem 291:20849–20857.

29. Gudermann T, Mederos y Schnitzler M, Dietrich A. 2004. Receptor-Operated Cation
Entry--More Than Esoteric Terminology? Sci Signal 2004:pe35–pe35.

30. Zhou Y, Frey TK, Yang JJ. 2009. Viral Calciomics: Interplays between Ca2+ and virus.
Cell Calcium 46:1–17.

31. Bagur R, Hajnóczky G. 2017. Intracellular Ca2+ Sensing: Its Role in Calcium
Homeostasis and Signaling. Mol Cell 66:780–788.

32. Dickson EJ, Jensen JB, Hille B. 2016. Regulation of calcium and phosphoinositides at
endoplasmic reticulum–membrane junctions. Biochem Soc Trans 44:467–476.

33. Nieto-Torres JL, Verdiá-Báguena C, Castaño-Rodriguez C, Aguilella VM, Enjuanes L.
2015. Relevance of viroporin ion channel activity on viral replication and pathogenesis.
Viruses 7:3552–3573.

34. Nieva JL, Madan V, Carrasco L. 2012. Viroporins: Structure and biological functions. Nat
Rev Microbiol 10:563–574.

35. Sze CW, Tan YJ. 2015. Viral membrane channels: Role and function in the virus life
cycle. Viruses 7:3261–3284.

36. Hyser JM, Collinson-Pautz MR, Utama B, Estes MK. 2010. Rotavirus disrupts calcium
homeostasis by NSP4 viroporin activity. MBio 1:1–12.

37. Van Kuppeveld FJM, De Jong AS, Melchers WJG, Willems PHGM. 2005. Enterovirus
protein 2B po(u)res out the calcium: A viral strategy to survive? Trends Microbiol 13:41–
44.

38. Wang K, Xie S, Sun B. 2011. Viral proteins function as ion channels. Biochim Biophys
Acta - Biomembr 1808:510–515.

39. Hyser JM, Estes MK. 2015. Pathophysiological Consequences of Calcium-Conducting
Viroporins. Annu Rev Virol 2:473–496.

40. Hyser JM, Estes MK. 2009. Rotavirus Vaccines and Pathogenesis: 2008. Curr Opin
Gastroenterol 25:36–43.

41. Pham T, Perry JL, Dosey TL, Delcour AH, Hyser JM. 2017. The Rotavirus NSP4
Viroporin Domain is a Calcium-conducting Ion Channel. Sci Rep 7:1–11.

42. van Kuppeveld FJ, Galama JM, Zoll J, Melchers WJ. 1995. Genetic analysis of a
hydrophobic domain of coxsackie B3 virus protein 2B: a moderate degree of
hydrophobicity is required for a cis-acting function in viral RNA synthesis. J Virol
69:7782–90.

43. de Jong AS, de Mattia F, Van Dommelen MM, Lanke K, Melchers WJG, Willems
PHGM, van Kuppeveld FJM. 2008. Functional Analysis of Picornavirus 2B Proteins:
Effects on Calcium Homeostasis and Intracellular Protein Trafficking. J Virol 82:3782–
3790.

44. Martinez-Gil L, Bano-Polo M, Redondo N, Sanchez-Martinez S, Nieva JL, Carrasco L,
Mingarro I. 2011. Membrane Integration of Poliovirus 2B Viroporin. J Virol 85:11315–

11324.

45.   Koonin E V, Dolja V V. 1993. Evolution and taxonomy of positive-strand viruses:
Implications and analysis of amino acid sequences. Crit Rev Biochem Mol Biol 28:375–
430.

46.   Zell R. 2017. Picornaviridae—the ever-growing virus family. Arch Virol 163:1–19.

47.   Chang-Graham AL, Perry JL, Strtak AC, Ramachandran NK, Criglar JM, Philip AA,
Patton JT, Estes MK, Hyser JM. 2019. Rotavirus Calcium Dysregulation Manifests as
Dynamic Calcium Signaling in the Cytoplasm and Endoplasmic Reticulum. BioRxiv 1–
19.

48.   Hadley A, Henry L. 2019. Package "tidyr": Easily Tidy Data with "spread()" and
"gather()" Functions.

49.   Wickham H. 2009. ggplot2: Elegant Graphics for Data Analysis, p. 1–16. *In* Valero-Mora,
PM (ed.), UseR! Springer-Verlag, New York.

50.   Saier MH. 2005. TCDB: the Transporter Classification Database for membrane transport
protein analyses and information. Nucleic Acids Res 34:D181–D186.

51.   Buchan DWA, Jones DT. 2019. The PSIPRED Protein Analysis Workbench: 20 years on.
Nucleic Acids Res 1–6.

52.   Gautier R, Douguet D, Antonny B, Drin G. 2008. HELIQUEST: A web server to screen
sequences with specific α-helical properties. Bioinformatics 24:2101–2102.

53.   Hyser JM, Utama B, Crawford SE, Broughman JR, Estes MK. 2013. Activation of the
Endoplasmic Reticulum Calcium Sensor STIM1 and Store-Operated Calcium Entry by
Rotavirus Requires NSP4 Viroporin Activity 87:13579–13588.

54.   Nan X, Hyndman L, Agbi N, Porteous DJ, Boyd AC. 2004. Potent stimulation of gene
expression by histone deacetylase inhibitors on transiently transfected DNA. Biochem
Biophys Res Commun 324:348–354.

55.   Lu HK, Gray LR, Wightman F, Ellenberg P, Khoury G, Cheng WJ, Mota TM, Wesselingh
S, Gorry PR, Cameron PU, Churchill MJ, Lewin SR. 2014. Ex vivo response to histone
deacetylase (HDAC) inhibitors of the HIV long terminal repeat (LTR) derived from HIV-
infected patients on antiretroviral therapy. PLoS One 9.

56.   Paszkiet BJ, Zhang J, Matukonis M, Kaleko M, Luo T. 2016. Histone Deacetylation
Inhibitors Enhance Lentiviral Vector Production and Infectivity. Mol Ther 5:S308.

57.   Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image
analysis. Nat Methods 9:671–5.

58.   Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr
RA, Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS. 2013. Ultrasensitive
fluorescent proteins for imaging neuronal activity. Nature 499:295–300.

59.   Billman GE. 1993. Intracellular calcium chelator, BAPTA-AM, prevents cocaine-induced
ventricular fibrillation. Am J Physiol Circ Physiol H1529–H1535.

60. Horvath B, Szentandrassy N, Veress R, Baranyai D, Kistamas K, Almassy J, Toth A,
Magyar J, Banyasz T, Nanasi PP. 2018. Effect of the intracellular calcium concentration
chelator BAPTA acetoxy-methylester on action potential duration in canine ventricular
myocytes. J Physiol Pharmacol 69:99–107.

61. Gonzalez ME, Carrasco L. 2003. Viroporins. FEBS Lett 552:28–34.

62. Neil JD. 1990. Nucleotide sequence of a region of the feline calicivirus genome which
encodes picornavirus-like RNA-dependent RNA polymerase , cysteine protease and 2C
polypeptides. Virus Res 17:145–160.

63. Kuyumcu-Martinez M, Belliot G, Sosnovtsev S V., Chang K-O, Green KY, Lloyd RE.
2004. Calicivirus 3C-Like Proteinase Inhibits Cellular Translation by Cleavage of
Poly(A)-Binding Protein. J Virol 78:8172–8182.

64. Venkataraman S, Prasad BVLS, Selvarajan R. 2018. RNA dependent RNA polymerases:
Insights from structure, function and evolution. Viruses 10:1–23.

65. Lama J, Carrasco L. 1992. Expression of poliovirus nonstructural proteins in Escherichia
coli cells. J Biol Chem 267:15932–15937.

66. Guinea R, Carrasco L. 1994. Influenza virus M2 protein modifies membrane permeability
in E. coli cells. FEBS Lett 343:242–246.

67. Crawford SE, Estes MK. 2013. Viroporin-mediated calcium-activated autophagy 797–
798.

68. Furlong K, Hwang S. 2019. Autophagy and Noroviruses. Viruses 11:244.

69. Krebs J, Agellon LB, Michalak M. 2015. Ca2+ homeostasis and endoplasmic reticulum
(ER) stress: An integrated view of calcium signaling. Biochem Biophys Res Commun
460:114–121.

70. van Kuppeveld FJ, Galama JM, Zoll J, van den Hurk PJ, Melchers WJ. 1996. Coxsackie
B3 virus protein 2B contains cationic amphipathic helix that is required for viral RNA
replication. J Virol 70:3876–3886.

71. Lock JT, Alzayady KJ, Yule DI, Parker I. 2018. All three IP 3 receptor isoforms generate
Ca 2+ puffs that display similar characteristics . Sci Signal 11:eaau0344.

72. Taylor CW, Tovey SC. 2010. IP3 Receptors: Toward Understanding Their Activation.
Cold Spring Harb Perspect Biol 1–23.

73. Jin X, Shah S, Du X, Zhang H, Gamper N. 2016. Activation of Ca2+-activated Cl-
channel ANO1 by localized Ca2+ signals. J Physiol 594:19–30.

74. Concepcion AR, Vaeth M, Wagner LE, Eckstein M, Hecht L, Jun Y, Crottes D, Seidl M,
Shin HP, Weidinger C, Cameron S, Turvey SE, Issekutz T, Meyts I, Lacruz RS, Cuk M,
Yule DI, Feske S. 2016. Store-operated Ca2+ entry regulates Ca2+-activated chloride
channels and eccrine sweat gland function. J Clin Invest 126:4303–4318.

884