The *C. elegans* 3'UTRome V2: an updated genomic resource to study 3'UTR biology

Steber HS[1,2], Gallante C[3], O'Brien S[2], Chiu P.-L[4], Mangone M*[1,2].

[1]Molecular and Cellular Biology Graduate Program, School of Life Sciences 427 East Tyler Mall Tempe, AZ 85287 4501.

[2]Virginia G. Piper Center for Personalized Diagnostics, The Biodesign Institute at Arizona State University, 1001 S McAllister Ave, Tempe, AZ, USA

[3]Barrett, The Honors College, Arizona State University, 751 E Lemon Mall, Tempe, AZ 85281

[4]Center for Applied Structural Discovery, The Biodesign Institute at Arizona State University, 1001 S McAllister Ave, Tempe, AZ 85287, USA

* To whom correspondence should be addressed. Tel: +1(480) 965-7957; Fax: +1(480) 965-3051;

Email: mangone@asu.edu

Present Address: Marco Mangone, Arizona State University, Biodesign Institute Building A, 1001 S McAllister Ave, Tempe, AZ 85281 USA

Keywords: *C. elegans*, 3'Untranslated Regions, Alternative Polyadenylation, Cleavage and Polyadenylation Complex, miRNAs, 3'UTRome

ABSTRACT

3'Untranslated Regions (3'UTRs) of mRNAs emerged as central regulators of cellular function as they contain important but poorly-characterized *cis*-regulatory elements targeted by a multitude of regulatory factors. The soil nematode *C. elegans* is an ideal model to study these interactions since it possesses a well-defined 3'UTRome. In order to improve its annotation, we have used a genomics approach to download raw transcriptome data for ~1,500 transcriptome datasets from the Sequence Read Archive at the NCBI and then extract and map high-quality 3'UTR data at ultra-deep coverage. Here we describe and release to the Community the updated version of the worm 3'UTRome, which we named 3'UTRome v2. This resource contains high-quality 3'UTR data mapped at single base ultra-resolution for 23,159 3'UTR isoforms variants corresponding to 14,808 protein-coding genes and is updated to the latest release of WormBase. We used this dataset to study and probe principles of RNA cleavage and polyadenylation in *C. elegans*. The worm 3'UTRome v2 represents the most comprehensive and high-resolution 3'UTR dataset available in *C. elegans*, and it allows a better understanding of this important process *in vivo*.

BACKGROUND

3'Untranslated Regions (3'UTRs) are the portions of mRNA located between the

end of the coding sequence and the polyA tail of RNA polymerase II-transcribed genes.

They contain *cis*-regulatory elements targeted by miRNAs and RNA binding proteins and

modulate mRNA stability, localization, and overall translational efficiency (Bartel 2018).

Because multiple 3'UTR isoforms of a particular mRNA can exist, differential regulation of

3'UTRs has been implicated in numerous diseases, and its discriminative processing

influences development and metabolism (Mayr and Bartel 2009; Zhu et al. 2018). 3'UTRs

are processed to full maturity through cleavage of the nascent mRNA and subsequent

polyA tail addition to its 3' end by the nuclear polyA polymerase enzyme (PABPN1) (Kuhn

and Wahle 2004). The mRNA cleavage step is a dynamic regulatory process directly

involved in the control of gene expression in Eukaryotes. The reaction depends on the

presence of a series of sequence elements located within the end of the 3'UTRs. The most

well-characterized sequence is the PolyA Signal (PAS) element, a hexameric motif located

at ~19nt from the polyadenylation site in the 3'UTR of mature mRNAs. In metazoans, the

PAS element is commonly 'AAUAAA', which accounts for more than half of all 3' end

processing in eukaryotes (Mangone et al. 2010; Tian and Graber 2012) although

alternative forms of the PAS elements exist (Sheets et al. 1990; Mangone et al. 2010).

Previous studies have shown that single base substitutions in this sequence reduce the

effectiveness of the cleavage and the polyadenylation of the mRNA transcript (Sheets et

al. 1990; Chen et al. 1995). However, this canonical sequence is necessary and sufficient

for efficient 3' end polyadenylation *in vitro* (Clerici et al. 2018; Sun et al. 2018). A less

defined 'GU rich' element is also known to be present downstream of the cleavage site to

facilitate the cleavage and polyadenylation steps (Chen et al. 1995). Recently, studies in

human cells identified an additional upstream 'UGUA' sequence that is not required for the

cleavage process, but acts as a cleavage enhancer in the context of Alternative

Polyadenylation (APA) (Zhu et al. 2018).

APA is a poorly understood mRNA maturation step that produces mRNAs with

different 3'UTR lengths due to the presence of multiple PAS elements within the same

3'UTR. The usage of the most upstream element, termed the proximal PAS element, leads

to the formation of shorter 3'UTR isoforms while the use of the distal PAS element results

in a longer isoform. Importantly, these changes in size may include or exclude regions to

which regulatory molecules such as microRNAs (miRNAs) and RNA-binding proteins

(RBPs) can bind, substantially impacting gene expression (Matlin et al. 2005; Bartel 2009).

While its function in eukaryotes is still not fully understood, a recent study revealed that

APA may occur in a tissue-specific manner and, at least in the soil nematode *C. elegans*,

is used in specific cellular contexts to evade miRNA-based regulatory networks in a tissue-

specific manner (Blazie et al. 2015; Blazie et al. 2017).

The cleavage and polyadenylation reaction is still poorly characterized in

metazoans. Although it involves a multitude of proteins and is considered to be very

dynamic, the order in which this process is executed and the role of each member of the

complex is still not fully understood.

In humans, the Cleavage and Polyadenylation Complex (CPC) is composed of at

least 17 members (**Figure 1A**) which immunoprecipitate into at least four large sub-

complexes: the Cleavage and Polyadenylation Specificity Factor (CPSF), the Cleavage

Stimulation Factor (CstF), the Cleavage Factor Im (CFIm) and the Cleavage Factor IIm

(CFIIIm) sub-complexes (**Figure 1A**). CPSF forms the minimal core complex necessary

and sufficient to recognize and bind the PAS element of the nascent mRNA *in vitro* (Tian

and Manley 2017) (**Figure 1A**). In humans, the CPSF sub-complex is composed of

CPSF160 (Clerici et al. 2017; Sun et al. 2018), CPSF100 (Mandel et al. 2006), CPSF73

(Mandel et al. 2006), CPSF30 (Clerici et al. 2017; Sun et al. 2018), Fip1 (Kaufmann et al.

2004) and Wdr33 (Clerici et al. 2017; Sun et al. 2018). Initial experiments assigned

CPSF160 with the role of binding the PAS element, but it is now clear that Wdr33 and

CPSF30 are the proteins that instead contact the PAS directly. CPSF160 has a scaffolding

role in this process and keeps this sub-complex structured (Chan et al. 2014). The

interaction between members of the CPSF core complex (Wdr33, CPSF30, and CPSF160)

and the PAS element was recently revealed using single-particle cryo-EM (Clerici et al.

2017; Sun et al. 2018), showing a unique conformation where the PAS element twists to

form an s-shaped structure with a non-canonical pairing between the U3 and the A6 in the

PAS element (Sun et al. 2018).

CPSF73 is the endonuclease that performs the cleavage of the nascent mRNAs

(Ryan et al. 2004; Mandel et al. 2006) (**Figure 1A**). CPSF73 possesses a Metallo-β-

lactamase domain and a β-CASP domain used to recognize and cleave nucleic acids.

Purified recombinant CPSF73 retains RNA endonuclease activity, and mutations that

disrupt the zinc binding in the active site of the enzyme abolish this activity (Mandel et al.

2006), suggesting that this protein's role is to perform mRNA cleavage. Importantly,

CPSF73 is also required in the cleavage of pre-histone mRNAs and is recruited on their

cleavage site by the U7 SNP (Yang et al. 2009).

Fip1 is another member of the CPSF sub-complex. Fip1 interacts with PABPN1,

which is the enzyme that performs the polyadenylation reaction on the cleavage site. Fip1

preferentially binds U-rich elements in the nascent mRNA and stabilizes the cleavage

complex using its arginine-rich RNA-binding domain (Kaufmann et al. 2004). Together with

CPSF160 and PABPN1, Fip1 forms a ternary complex *in vitro* (Kaufmann et al. 2004)

capable of inhibiting endogenous PABPN1activity (Zhelkovsky et al. 1998; Helmling et al.

2001), suggesting a bridging role for this protein in the complex.

The CstF sub-complex is the second most well-characterized sub-complex involved

in the cleavage and polyadenylation reaction (**Figure 1A**). CstF binds to GU rich elements

located downstream of the cleavage site in the nascent mRNA and directly contacts the

CPSF sub-complex using its conserved HAT-C domain (Bai et al. 2007; Yang et al. 2018)

(**Figure 1A**). The CstF sub-complex is a dimer of heterotrimers composed of CstF77,

CstF64 and CstF50 (Yang et al. 2018). CstF77 holds the complex together through its Pro-

rich domain located on its C terminal region (Takagaki and Manley 2000) **(Figure 1A)**.

CstF64 recognizes GU rich sequences through its N-terminal RRM domain (Perez

Canadillas and Varani 2003; Yang et al. 2018) and interacts with the scaffolding protein

Symplekin and CstF77 using its N-terminal hinge domain (**Figure 1A**) (Takagaki and

Manley 2000).

The CFIm and CFIIm sub-complexes are unfortunately less characterized (**Figure**

1A). The CFIm sub-complex is composed of the CFIm68, CFIm59 and CFIm25 subunits,

and it was recently shown to contribute to APA by influencing PAS selection (Martin et al.

2012; Hwang et al. 2016). CFIm25 binds a 'UGUA' RNA element upstream of the cleavage

site and contributes to 3'processing by recruiting CFIm59 and CFIm68 (Yang et al. 2010;

Yang et al. 2011; Zhu et al. 2018).

The Cleavage Factor IIm sub-complex is composed of two factors named Pcf11

and hClp1 (Schafer et al. 2018). Pcf11 binds RNA unspecifically through two zinc fingers

in its C-terminal region and stimulates the RNA 5' kinase activity of hClp1, which is not

required for the cleavage reaction (Schafer et al. 2018). It has been suggested that hClp1

binds CPSF, although the exact interaction has not been determined (de Vries et al. 2000).

Despite the importance of this complex, the CPC remains poorly characterized in

most species, including humans, and most of the research in this field is performed *in vitro*.

The round nematode *C. elegans* represents an attractive, novel system to study the

cleavage and polyadenylation process *in vivo*. Most of the CPC is conserved between

humans and nematodes, including known functional domains and protein interactions

(**Figure 1B and Supplemental Figure S1**). *C. elegans* possess the most well-annotated

3'UTRome available so far in metazoans, with mapped 3'UTR boundaries for ~26,000

distinct *C. elegans* protein-coding genes (Mangone et al. 2010; Jan et al. 2011).

The *C. elegans* 3'UTRome was originally developed in 2011 within the

modENCODE project (Mangone et al. 2008; Gerstein et al. 2010; Mangone et al. 2010)

and represented a milestone in 3'UTR biology since it allowed the Community to study and

identify important regulatory elements such as miRNAs and RBPs targets with great

precision. A second 3'UTRome was later published using a different mapping pipeline (Jan

et al. 2011), confirming most of the previous data such as isoforms numbers, PAS usage,

etc. Other datasets were made available later, mostly focusing on tissue-specific 3'UTRs

and alternative polyadenylation (Haenni et al. 2012; Blazie et al. 2015; Blazie et al. 2017;

Chen et al. 2017; Diag et al. 2018; West et al. 2018).

Although refined and based on several available datasets, only a subset of *C.*

elegans 3'UTRs in protein-coding genes are sufficiently annotated today, and the existing

mapping tools do not yet reach the single-base resolution necessary to execute

downstream analysis and study the cleavage and polyadenylation process in detail. Most

of these 3'UTR datasets were developed using a gene model now considered obsolete

(WS190), and the 3'UTR coordinates often do not match the new gene coordinates.

To address these and other issues, we developed a novel pipeline to

bioinformatically extract 3'UTR data from almost all *C. elegans* transcriptome datasets

stored in the public repository SRA trace archive. This blind approach produced a new

saturated dataset we named 3'UTRome v2. This updated 3'UTRome contains 3'UTR data

for 23,159 3'UTR isoforms variants corresponding to 14,808 protein-coding genes and is

available to the Community as an additional gBrowse track in the *C. elegans* database

WormBase (www.WormBase.org) (Stein et al. 2001) and in the 3'UTR-centric database

3'UTRome (www.UTRome.org) (Mangone et al. 2008; Mangone et al. 2010).

We have also used this dataset to study the PAS sequence requirement and the

cleavage location of the CPC *in vivo* using transgenic *C. elegans* animals. We found that

the canonical CPC can in principle bind different PAS sequences and that elements

downstream of the PAS site can in turn influence the location of the cleavage.

RESULTS

Functional elements of the human cleavage and polyadenylation complex are

conserved in nematodes.

To initially gain structural and functional information for the *C. elegans* CPC, we

downloaded the protein sequences of the orthologs of the *C. elegans* CPC and aligned

them to their human counterparts (**Figure 1B and Supplemental Figure S1**). Based on

sequence similarity, *C. elegans* possess orthologs to all the known members of the human

CPC, with many peaks of conservation interspersed in the subunits within known

interaction domains. The amino acids that make direct contact with PAS elements are also

conserved in *C. elegans*; 11 out of the 12 amino acids that form hydrogen bonds and salt

bridges with the PAS element (Clerici et al. 2017) are present in both the CPSF30 and

WDR33 worm orthologs *cpsf-4* and *pfs-2* ($V67^{CPSF30}$ with $V81^{cpsf-4}$; $K69^{CPSF30}$ with $K83^{cpsf-4}$;

$R73^{CPSF30}$ with $R87^{cpsf-4}$; $E95^{CPSF30}$ with $E109^{cpsf-4}$; $K77^{CPSF30}$ with $K91^{cpsf-4}$; $S106^{CPSF30}$ with

$S120^{cpsf-4}$; $N107^{CPSF30}$ with $N121^{cpsf-4}$; $R54^{WRD33}$ with $R80^{pfs-2}$; $R47^{WRD33}$ with $R71^{pfs-2}$;

$R49^{WRD33}$ with $R73^{pfs-2}$) (**Figure 1B and Supplemental Figure S1**). The only exception is

$Y97^{CPSF30}$, which is substituted with a Phenylalanine residue in the worm ortholog. In

addition, 9 out of the 10 amino acids in CPSF30 and WDR33 that form the π-π stacking

and hydrophobic interactions with the AAUAAA RNA element (Clerici et al. 2017) are also

conserved in the CPSF30 and WDR33 worm orthologs *cpsf-4* and *pfs-2* (A1:K69^{CPSF30}

with K83^{cpsf-4} and F84 ^{CPSF30} with F98^{cpsf-4}; A2: H70^{CPSF30} with H84^{cpsf-4}; U3: I156^{WDR33} with

I181^{pfs-2}; A4: F112^{CPSF30} with F126^{cpsf-4} and F98^{CPSF30} with F112^{cpsf-4}; A5: F98^{CPSF30} with

F112^{cpsf-4}; A6: F153^{WDR33} with F178^{pfs-2}) (**Figure 1B and Supplemental Figure S1**). The

only exception is a F43^{WDR33} substitution to a Glycine residue that interacts with A6 in the

worm ortholog.

CPSF73, the endonuclease that performs the cleavage reaction, has a *C. elegans*

ortholog named *cpsf-3*. Both genes are conserved with an overall 57.61% identity that

increases to 69.52% in the β-lactamase domain, which is the region required to perform

the cleavage reaction (**Figure 1B and Supplemental Figure S1**). Specifically, all eight

amino acids shown previously to form the zinc binding site required for the cleavage

reaction (Mandel et al. 2006) are also conserved (D75^{CPSF73} with D74^{cpsf-3}; H76^{CPSF73} in

H75^{cpsf-3}; H73^{CPSF73} in D72^{cpsf-3}; H396^{CPSF73} with H397^{cpsf-3}; H158^{CPSF73} with H159^{cpsf-3};

D179^{CPSF73} with D180^{cpsf-3}; H418^{CPSF73} with H419^{cpsf-3}; E204^{CPSF73} with E205^{cpsf-3}) (**Figure**

1B and Supplemental Figure S1). This overall similarity is also observed in most of the

other members of the *bona fide C. elegans* CPC complex (**Supplemental Figure S1**),

suggesting similar structure and function.

In addition, when subjected to RNAi analysis, each of the *C. elegans* CPC members

produced a similar strong embryonic lethal phenotype, suggesting that each of these

genes may act as a complex and are required for viability (**Figure 1C and Supplemental**

Figure S2).

An updated 3'end mapping strategy

Next, we used a blind genomic approach to improve the current version of the

3'UTRome. We refined a 3'UTR mapping pipeline we previously developed and used in

the past (Blazie et al. 2015; Blazie et al. 2017). This approach uses raw transcriptome data

as input material to identify and precisely map high-quality 3'UTR end clusters (**Figure 2**

and Supplemental Figure S3).

We wanted to obtain the most accurate, saturated and tissue-independent dataset

possible. To achieve this goal, we downloaded from the Sequence Read Archive (SRA)

the majority of all publicly available transcriptome datasets and processed them through

our 3'UTR mapping pipeline. We reasoned that this blind approach would lead to the

identification of as many 3'UTR isoforms as possible in an unbiased manner since these

downloaded transcriptomes have been sequenced using both *wild-type* and mutant strains

subjected to many different environmental conditions and covering all developmental

stages with many replicates.

We downloaded a total of 1,094 *C. elegans* transcriptome datasets (~2TB of total

raw data), which represent the majority of transcriptomes available for this organism so far

(**Supplemental Table S1**). Most of these datasets have been also used in the past to map

polyadenylation sites in *C. elegans*. Our 3'UTR mapping approach extracted from these

datasets ~5M unique, high-quality polyA reads, which we then used for cluster preparation

and mapping (see Methods). To limit the unavoidable noise produced by using such

diverse datasets as data sources, we implemented very restrictive parameters for cluster

identification and 3'UTR end mapping (**Supplemental Figure S3**), leading to 3'UTR

clusters with ultra-deep coverage of several magnitudes (average cluster coverage ~220X)

(**Figure 2A**).

This approach allowed for the identification of 23,159 3'UTR isoforms

corresponding to 14,808 protein-coding genes. When compared with the previous

3'UTRome v1 dataset (Mangone et al. 2010), we obtained 3'UTR information for an

additional 4,638 new protein-coding genes (6,218 3'UTR isoforms) (**Figure 2B-C**).

The *C. elegans* 3'UTRome v2

Our approach produced high-quality 3'UTR data for 14,808 *C. elegans* protein-

coding genes (**Figure 2B**). The most abundant nucleotide in *C. elegans* 3'UTRs is a

Uridine, which accounts for 40% of all nucleotides in 3'UTRs (**Figure 3A Top Left Panel**).

Adenosine nucleotides are the second most represented nucleotide class with ~30% of

incidence (**Figure 3A Top Left Panel**).

Alternative polyadenylation is common, but occurs at a significantly lesser extent

than what was previously published (Mangone et al. 2010; Jan et al. 2011). The majority of

protein-coding genes (58%) are transcribed with only one 3'UTR isoform (**Figure 3A**

Bottom Left Panel) in contrast with ~61% as it was reported in the past (Mangone et al.

2010; Jan et al. 2011). Genes with two 3'UTR isoforms are notably increased in

occurrence when compared with past studies (32% vs 25%), while the occurrence of

genes with three or more 3'UTRs is comparable with what was previously found (**Figure**

3A Bottom Left Panel) (Mangone et al. 2010; Jan et al. 2011).

Interestingly, in the case of genes with multiple 3'UTRs, the canonical AAUAAA

PAS site is greater than two times more abundant in longer 3'UTR isoforms than in shorter

3'UTR isoforms, suggesting that the preparation of shorter 3'UTR isoforms may be subject

to regulation (**Supplemental Figure S4**).

The average 3'UTR length in the 3'UTRome v2 is 215nt (**Figure 3A Top Right**

Panel), and the occurrence of more 3'UTR isoforms per gene correlates with an overall

extension in length (**Figure 3A Top Right Panel**). We also note a slight correlation

between 3'UTR length and PAS element usage, with longer 3'UTRs more frequently

containing variant PAS elements (**Figure 3A Bottom Right Panel**). The most common

PAS element in *C. elegans* protein-coding genes is consistently the hexamer 'AAUAAA',

which is present in 58.4% of all the 3'UTRs mapped in this study (**Figure 3B Left Panel**).

This element is ~20% more abundant than what was previously identified in past studies

(Mangone et al. 2010; Jan et al. 2011). The PAS sequence is located ~ 18nt from the

cleavage site (**Figure 3B Right Panel**), and a *buffer region* of ~12nt is present between

the PAS element and the cleavage site (**Figure 3C**). The cleavage site occurs almost

invariably at an Adenosine nucleotide, which is often preceded by a Uridine nucleotide

(**Figure 3C**).

An RRYRRR motif in 3'UTRs with variant PAS elements

We could not detect any enrichment for the UGUA motif near the cleavage site

(**Supplemental Figure S5**), but perhaps this element is either not used in *C. elegans* or

the CFIm complex may recognize a variant motif not yet identified in this organism.

Importantly, when we align the 3' ends of 3'UTRs which contain variant PAS elements, we

notice an enrichment of an 'RRYRRR' motif where the canonical PAS element is generally

located; this suggests that in *C. elegans,* an 'RRYRRR' element could be used instead

when the AAUAAA hexamer is absent (**Figure 4A**).

To better understand the molecular details of the interaction between CPSF and the

PAS element, we built a pseudo-atomic homology model of the worm CPSF core complex

containing *cpsf-1* (CPSF160), *pfs-2* (Wdr33), and *cpsf-4* (CPSF30) (**Figure 4B and**

Supplemental Figure S6). Most of this model can be superimposed to the cryo-EM

structure of the human CPSF core complex (**Figure 4B and Supplemental Figure S6**).

The nucleotide-binding pocket can also be fitted into our homology model, which

may implicate a conserved binding region in the *C. elegans* complex (**Figure 4B Right**

Panel). From the structural details of the human CPSF core complex, the interactions

between the RNA nucleotides and CPSF30 or WDR33 are not specific. The nucleotide

binding is mainly established by the π-π ring stacking force between the nucleotide bases

and the residues with aromatic side chains, such as phenylalanine and tyrosine

(**Supplemental Figure S6**). Also, the binding pockets of the Adenine base do not seem to

have a steric hindrance for Guanine base to bind. It is similar for the Uridine base to the

Cytosine base (**Supplemental Figure S6**). Thus, at least in *C. elegans*, the selectivity of

the nucleotide binding may be only at a level to the nucleotide bases, that is, Pyrimidines

or Purines.

An enrichment of Adenosine nucleotide at the cleavage site

We were intrigued by the almost invariable presence of Adenosine nucleotides near

the cleavage site. This enrichment becomes more evident when we sort 3'UTRs with

canonical PAS elements by the length of their respective *buffer regions* (**Figure 5A**). In the

case of the largest group with a *buffer region* of 12-13nt, more than 2,000 3'UTRs

terminate with ~70% occurrence of Adenosine nucleotides at the cleavage site. Since we

bioinformatically removed the polyA sequences from the sequencing reads during our

cluster preparation step, we do not have direct evidence that this last Adenosine

nucleotide is indeed present in the mature transcripts and used as a template for the

polymerization of the polyA tail or that it is attached by PABPN1 during the polymerization

of the polyA tail. Of note, the high abundance of this nucleotide at the cleavage site

suggests that it is somehow important in the cleavage process.

We decided to investigate this issue further and study how precisely the raw reads

produced by our cluster algorithm align to the genome. We noticed that in each gene the

cleavage rarely occurs at a unique position in the transcript. Instead, there are always

slight fluctuations of the exact cleavage site, with a few percentages of reads ending a few

nucleotides upstream and downstream of the most abundant cleavage site for a given

gene (**Figure 5B**). Importantly, almost all the reads in each cluster terminate at an

Adenosine nucleotide (**Figure 5B**). Also, if there are Adenosine nucleotides located within

shorter *buffer regions*, the cleavage rarely occurs at these sites. Perhaps, the large size of

the CPC does not allow for the docking and the cleavage of the pre mRNAs near the PAS

element, which is optimally performed at 12-13nt downstream the PAS (**Figure 5A and**

Figure 5B).

Next, we decided to study the role of the terminal Adenosine nucleotide in the

cleavage process. We reasoned that if this Adenosine nucleotide indeed plays any role in

the cleavage process, we should be able to alter the position of the mRNA cleavage site

by mutating this residue with different Purines or Pyrimidines in the pre mRNAs of selected

test genes.

We selected three test genes; *ges-1*, Y106G6H.9, and M03A1.3. These genes are

processed only with a single 3'UTR isoform, use a single canonical PAS element, have a

buffer region of 12, 13 and 14 nucleotides respectively and possess a terminal Adenosine

nucleotide in their sequence. To capture their entire 3'UTR region, we cloned the genomic

portions of these genes spanning from their translation STOP codons to ~200nt

downstream of their cleavage sites. We then prepared several mutant *C. elegans* strains

replacing their terminal Adenosine nucleotide at their cleavage site with other nucleotides.

In the case of Y106G6H.9 we also prepared a double mutant removing an additional

Adenosine nucleotide upstream of the first one located at the cleavage site (**Figure 5C**

and Supplemental Figure S7-S9).

We cloned these *wt* and mutant 3'UTR regions downstream of a GFP reporter

vector and prepared transgenic *C. elegans* strains that express them in the worm pharynx

using the *myo-2* promoter. We opted to use the pharynx promoter since it is very strong

and produces a robust expression of our constructs (**Supplemental Figure S7-S9**). We

prepared transgenic worm strains expressing these constructs, recovered total RNAs, and

tested using RT-PCR and a sequencing approach if the absence of the terminal

Adenosine nucleotide in our mutants affects the position of the cleavage site (**Figure 5C**

and Supplemental Figure S7-S9).

We observed an overall disruption of the cleavage process, in some case more

pronounced than in others (**Figure 5C and Supplemental Figure S7-S9**). In the case of

M03A1.3, the absence of the terminal Adenosine nucleotide forces the cleavage complex

to backtrack in 40% of the tested clones and cleave the mRNAs 3nt upstream of the

original cleavage site, but still at an Adenosine nucleotide (**Figure 5C and Supplemental**

Figure S7).

In the case of Y106G6H.9, the single mutant does not alter the position of the

cleavage site, but interestingly activates a novel cryptic cleavage site 100 nucleotides

upstream of the canonical cleavage site in 20% of the sequenced clones ~ (**Figure 5C**

and Supplemental Figure S8). This new site also possesses a non-used PAS element

containing the motif YRYRRR, which could still be recognized by the CPSF core complex

in this mutant, and a *buffer region* of 12nt. The Y106G6H.9 double mutant in one case

skips the original cleavage site but still cut at the next Purine residue, which is not an

Adenosine in this case (**Supplemental Figure S8**). In the case of *ges-1*, mutating the

terminal Adenosine does not change the cleavage pattern, although it became more

imprecise (**Supplemental Figure S9**).

Updated miRANDA prediction in *C. elegans*

Next, we used our new UTRome v2 dataset to update MiRanda miRNA target

predictions. We downloaded and locally ran the miRanda prediction software (John et al.

2004) using our new 3'UTRome v2 as a target dataset. We have produced two sets of

predictions; one generic, which contain the entire output produced by the software, and

one more restrictive, in which we only output predictions with high scoring and with low E-

energy scores. These two tracks have been uploaded in both the 3'UTRome database

(Mangone et al. 2008; Mangone et al. 2010) and the WormBase (Stein et al. 2001).

DISCUSSION

Here we have used a blind genome-wide approach to refine and study the

3'UTRome in the nematode *C. elegans*. We have identified 3'UTR data for 14,808 genes,

corresponding to 23,159 3'UTR isoforms, significantly improving their annotation.

Alternative Polyadenylation is widespread in *C. elegans*, with ~42% of genes

possessing at least two 3'UTR isoforms (**Figure 3A**). The PAS usage is still most

commonly the hexamer 'AAUAAA', which is used to process ~58% of all *C. elegans*

3'UTRs (**Figure 3B**). Importantly, we found that the remaining 42% possess a variation of

this canonical PAS element which indeed is very similar in chemical composition and

contain an 'RRYRRR' motif at the same location where the PAS element is expected

(**Figure 4A**). Perhaps this element is sufficient to induce the cleavage reaction in

nematodes.

Our superimposition of the *C. elegans* CPSF ortholog to the human cryo-EM

structure (Clerici et al. 2018; Sun et al. 2018) in **Figure 4B and Supplemental Figure S6**

suggests that in worms, the pocket used by this complex to bind the PAS element may

accommodate other nucleotides as long as they have a similar chemical structure and can

recapitulate the 'RRYRRR' motif. In humans, the second most abundant PAS element is

'AUUAAA' (Sun et al. 2018), which does not follow this guideline, suggesting that perhaps

other factors can contribute to the cleavage of non-canonical PAS elements in other

species.

We also performed an analysis on the cleavage site and found that the Cleavage

and Polyadenylation machinery does not always cleave the same mRNA at the same

position on the 3'UTRs (**Figure 5B**). While a predominant site is often chosen for each

gene, a slight variation of a few nucleotides upstream or downstream of the cleavage site

is also possible. Importantly, this slight variation almost invariably ends at an Adenosine

nucleotide in the Genome, suggesting that this nucleotide is somehow important in the

cleavage process.

The concept of mRNAs terminating with an Adenosine nucleotide is not novel.

Pioneering work using 269 vertebrate cDNA sequences has shown that ~71% of these

genes terminate with a CA nucleotide element (Sheets et al. 1990). These experiments

were biochemically validated a few years later using SV40 Late PolyA signal in

mammalian cells in a more controlled environment (Chen et al. 1995). These experiments

also showed that, at least for the case of this specific 3'UTR, the cleavage could not occur

closer than 11nt from the PAS element, and no farther of 23 nt from it (Chen et al. 1995).

This, in turn, could explain why we do not detect a terminal Adenosine at the cleavage site

with our double mutant Y106G6H.9 (**Supplemental Figure S8**). Of note, in the case of this

gene, the cleavage still occurs at a Purine nucleotide, suggesting that perhaps in the

absence of a terminal Adenosine another Purine can be used.

Overall, our experiments shown in **Figure 5C and Supplemental Figures S7-9**

support and expand both these initial results.

Our study does not have the resolution to probe if this Adenosine nucleotide is

included in the processed mRNAs. More specifically we do not know if this nucleotide is

read by the RNA polymerase II and incorporated in the nascent mRNAs or if the

machinery somehow 'senses' its presence and cleaves the mRNA upstream of it. Another

attractive hypothesis is that CPSF73 cleaves the mRNAs downstream of this terminal

Adenosine nucleotide, and then exonucleases degrade the mRNA molecule until they

reach the first Adenosine in a row. Of note, CPSF73 also cleaves the polyA-lacking

histone mRNAs. In this case, the enzyme is positioned near the cut site by the U7 snRNP,

and interestingly cuts just downstream of an Adenosine nucleotide (Yang et al. 2009),

suggesting that perhaps CPSF73 is capable of either 'sensing' this terminal Adenosine or

is positioned next to it by either other members of the CPC or not yet identified factors.

More experiments need to be performed to answer these questions.

Of note, while we observe a terminal Adenosine nucleotide in most of the mapped

3'UTRs, the Cytosine nucleotide previously identified in humans is replaced with another

Pyrimidine nucleotide in *C. elegans* (Thymidine) (**Figure 3C**).

MiRanda predictions were obsolete and needed to be updated since those present

in the microrna.org database (www.microrna.org) were obtained using a 9-year-old 3'UTR

dataset. Also, prior to this study, WormBase (Stein et al. 2001) did not include miRNA

targeting predictions in its JBrowse software.

The number of predicted miRNA targets is now decreased from 34,186 to 23,160,

mostly because several 3'UTR isoforms in the 3'UTRome V1 were discarded in this new

UTRome V2 release.

In conclusion, this new 3'UTR dataset, which we renamed 3'UTRome v2, has been

uploaded to the WormBase (Stein et al. 2001) and it is shown as a new track in the

JBrowse tool together with updated MiRanda miRNA target predictions. The 3'UTRome v2

expands the old 3'UTRome developed within the modENCODE Consortium and, together

with updated MiRanda predictions, provides the *C. elegans* Community with an important

novel resource to investigate 3'UTR biology and miRNA targeting.

452   **METHODS**

Comparative analysis of *C. elegans* members of the CPC

We have downloaded the protein sequences of each known member of the human CPC

and used BLAT algorithm to identify *C. elegans* genes with high homology to their human

counterparts. We then performed a Protein BLAST analysis using the tools available at the

NCBI website to obtain the amino acid sequences for the fly, rat, and mouse orthologs.

These amino acid sequences were then aligned using Clustal Omega Multiple Sequence

Alignment with standard parameters. At the completion of the analysis, we used the Batch

NCBI Conserved Domain Search (Batch CD-Search) against the database CDD- 52910

PSSMs using standard parameters to identify the conserved domains across the aligned

protein sequences. We then used these results to populate the location of these elements

within the alignment shown in **Supplemental Figure S1**.

3'UTR mapping pipeline

We have use the SRA toolkit from the NCBI to download raw reads from 1,094

transcriptome experiments. The complete list of datasets used in this study is shown in

Supplemental Table S1. We restricted the analysis to sequences produced from *C.*

elegans transcriptomes using the Illumina platform and with reads of at least 150nt in

length. At the completion of the download step, the files were unzipped and stored in our

servers. We then used custom-made Perl scripts to extract reads containing at least 23

consecutive Adenosine nucleotides at their 3'end or 23 consecutive Thymidine nucleotides

at their 5'end. This filter produced 24,973,286 mappable 3'end reads. We then removed

the terminal Adenosine or Thymidine nucleotides from these sequences, converted them

to fastq files using the FASTX-Toolkit (CSHL), and mapped them to the WS250 release of

the *C. elegans* genome using Bowtie2 algorithm with standard parameters (Langmead and

Salzberg 2012). The Bowtie algorithm mapped 7,761,642 reads (31.08%), which were

sorted and separated, based on their respective strand origin (positive or negative).

Cluster Preparations

PolyA clusters were prepared as follow. We stored the ID, genomic coordinates, and the

strand orientation of each mapped read, and used this information throughout the pipeline.

The BAM file produced by the aligners were sorted and converted to BED format using

SAMtools software (Li et al. 2009). Contiguous genomic coordinates were merged using

BEDTools software (Quinlan and Hall 2010) using the following command 'Bedtools

merge -c 1 -o count -I > tmp.cluster'. This new file produced the

characteristic 'shark fin' graph visible in **Figure 2**. We used several stringed filters to

eliminate as much as noise possible. 1) We ignored clusters composed of less than 6

reads. 2) We extracted genomic DNA sequences 20nt downstream the end of each

cluster. If the number of Adenosine nucleotides was more than 65% in the genomic

sequence, we ignored the corresponding cluster and marked it as caused by mispriming

during the second strand synthesis in the RT reaction. 3) We ignored clusters overlapping

with other clusters in the same orientation by 2nt or less were both ignored. 4) We

attached clusters to the closest gene in the same orientation. If no gene could be identified

within 2,000nt the cluster was ignored. 5) In cases with multiple 3'UTR isoforms identified,

we calculated the frequency of occurrence for each isoform and ignored isoforms

occurring at a frequency of less than 1% independently from the number of reads that form

this cluster.

Plasmid DNA isolation, sequencing and visualization

All plasmids used in this study were prepared from cultures grown overnight in LB using

the Wizard Plus SV Minipreps DNA Purification System (Promega) according to the

manufacturer's instructions. DNA samples were sequenced with Sanger sequencing

performed at the DNASU Sequencing Core Facility (The Biodesign Institute, ASU, Tempe,

AZ).

RNAi experiments

RNAi experiments were performed in Standard NGM agar containing 1mM IPTG and 50

µg/ml Ampicillin. These plates were seeded with 75 µl of RNAi clone bacteria and allowed

to induce for a minimum of 16 hours. 5 N2 *C. elegans* at the L1 stage were aliquoted for

each RNAi clone tested. Three days after plating, the progeny was scored for embryonic

lethality. Each RNAi experiment was performed in triplicate. The total number of hatched

and not matched eggs was the following: *cpsf-1(CPSF160)* n=567*; cpsf-2(CPSF100)*

n=557*; cpsf-4(CPSF30)* n=1,251*; cpf-2(CstF64)* n= 652*; cpf-1(CstF50)* n=801*; cfim-*

1(CFIm25) n=644*; cfim-2(CFIm68)* n=739*; lrp-2(CFIm59)* n=1,120*; symk-1(Symplekin)*

n=208*; tag-214(RBBP6)* n=753*; pcf-11(CPF11)* n=428*; clpf-1(CLP1*) n=841*.*

Extraction of 3'UTR regions from the *C. elegans* genome

The 3'UTRs used in the experiments described in **Figure 5C and Supplemental Figure**

S7-9 were initially cloned from N2 *wild type C. elegans* genomic DNA using PCR with

Platinum Taq Polymerase (Invitrogen). Genomic DNA template was prepared as

previously described (Blazie et al. 2017). Forward DNA primers were designed to include

approximately 30 nucleotides upstream of the translation STOP codon and include the

endogenous translation STOP codon. We used the Gateway BP Clonase II Enzyme Mix

(Invitrogen) to clone the 3'UTR region into Gateway entry vectors. The DNA primer was

modified to include the attB Gateway recombination elements required for insertion into

pDONR P2RP3 (Invitrogen). The reverse DNA primers were designed to end between 200

and 250 nucleotides downstream of the RNA cleavage site and to include the reverse

recombination element attB for cloning into pDONR P2RP3 (Invitrogen). At the conclusion

of the recombination step, the entry vectors containing the cloned 3'UTR regions were

transformed into Top10 competent cells (Thermo Fisher Scientific), using agar plates

containing 20mg/μL of Kanamycin. The plasmids were then recovered, and clones were

confirmed using Sanger sequencing with the M13F primer. The list of primers used in this

study is available in **Supplemental Table S2**.

Mutagenesis of 3'UTRs cleavage sites

The mutagenesis reactions to remove the Adenosine nucleotides near the cleavage sites

were carried out using the QuikChange Site-Directed Mutagenesis Kit (Agilent). The

mutagenesis DNA primers for the site mutation reactions are available in **Supplemental**

Table S2. Each mutagenesis reaction was followed by DNA digestion using Dpn-1

enzyme and transformed in Top10 competent cells (Thermo Fisher Scientific) in agar

plates containing 20mg/μL of Kanamycin. We validated the nucleotide mutation using

Sanger sequencing approach. *Wild type* and mutant 3'UTRs cloned in pDONR P2RP3

were then shuttled into destination vectors using the Gateway LR Clonase II Plus Enzyme

Mix (Invitrogen, Carlsbad, CA). The finalized destination vectors contained the *C. elegans*

pharynx promoter (Pmyo-2) in the first position, a GFP sequence with a mutated STOP

codon in the second position, and the *wt* or mutant 3'UTRs used in this study in the third

position. The resultant recombined constructs were then transformed in Top10 competent

cells (Thermo Fisher Scientific) and plated on 10mg/μL Ampicillin plates overnight. The

success of the recombination reaction was confirmed using Sanger sequencing with the

M13F DNA primer.

Preparation of transgenic worm lines

Eg6699 strain worms were kindly provided by Christian Frokjaer-Jensen (Frokjaer-Jensen

et al. 2008). These worm strains were maintained at 18°C on nematode growth media

(NGM) agar plates and propagated on plates seeded with OP50-1 bacteria. To

synchronize worms for injections, Eg6699 worms were bleached with bleaching solution (1

M NaOH) four days before injections. Each construct was mixed with an injection master

mix containing pCFJ601 (25 ng/μl), pgH8 (10 ng/μl), and pCFJ104 (5 ng/μl) vectors.

Injection needles were loaded with the injection mixture and mounted to the Leica

DMI300B microscope. The needle was pressurized with 22 psi through the FemtoJet

(Eppendorf). Young adult Eg6699 worms were picked onto an agarose pad covered with

mineral oil on a glass coverslip. Injected worms were rescued onto an NGM plate and

rinsed with M9 buffer. Two days post-injections, the F1 progeny were screened with a

Leica DMI3000B microscope for both *unc-119* rescues and expression of the red

fluorescence produced by the co-injection marker and then isolated onto individual plates.

These worms were allowed to lay eggs, and then the F2 progeny was screened for

fluorescence. Once 75% of the progeny on a single plate were transgenic, the strains were

used for further experimentation.

Worm genotype validation

Populations obtained from single worms from each of the seven strains were lysed using

worm lysis buffer (EDTA, 0.1 M Tris, 10% Triton-X, Proteinase K, 20% Tween 20). These

worms were subjected to heating in a Bio-Rad T100 Thermal Cycler. To confirm that the

mutated cleavage site was present in the injected strains, we used PCR approach using

Platinum Taq polymerase (Invitrogen) with a forward DNA primer binding the beginning of

the GFP sequence and 3'UTR-specific reverse DNA primers. The PCR product was then

sequenced using Sanger sequencing with a forward DNA primer binding to the GFP

sequence present in the injected construct.

Detection of the 3'UTR cleavage skipping

Total RNA was extracted from transgenic strains using the Direct-zol RNA MiniPrep Plus

kit (RPI) according to the manufacturer's instructions. We tested approximately 10

independent *wt* and mutant clones for each 3'UTR. Approximately 50 μL of worm pellet

was used for extraction. cDNA was synthesized using a reverse transcription reaction

using Superscript II enzyme (Invitrogen). The first strand reaction was performed using a

reverse poly dT DNA primer containing two anchors and the attB Gateway BP

recombination element (Invitrogen). The second strand of the cDNA was synthesized

using a PCR with HiFi taq polymerase (Thermo Fisher Scientific) and the forward DNA

primer containing the pDONR P2RP3 Gateway element (Invitrogen), which binds to GFP

and the same reverse poly dT DNA primer used in the first strand reaction. The BP

Gateway kit (Invitrogen) was once again used to clone the cDNA which contains the polyA

tail into pDONR P2RP3. These constructs were then transfected into Top10 competent

cells (Thermo Fisher Scientific) and plated on agar plates containing 20mg/μL of

Kanamycin. About 8-10 colonies were then sequenced with Sanger sequencing using the

M13F DNA primer to map the location of the cleavage site.

Updated MiRanda Predictions

We downloaded a complete list of *C. elegans* miRNAs from miRBase (Griffiths-Jones et al.

2006) and the miRanda algorithm v3.3a (John et al. 2004) from the microrna.org website.

We queried the 3'UTRome v2 with the miRanda algorithm using both standard and

stringent parameters. The stringent query used was '-strict -sc -1.2'. The standard query

produced 58,330 putative miRNA targets; the stringent query produced 12,136 putative

miRNA targets. Both these predictions are included in WormBase (Stein et al. 2001) as

individual tracks.

Homology model building

Homology modeling was performed using SWISS_MODEL (Waterhouse et al.

2018) with a matched templated of human CPSF160-WDR33-CPSF30 complex

(PDB code: 6DNF) (Sun et al. 2018). The molecular graphics were prepared using

the UCSF ChimeraX software (version 0.8) (Goddard et al. 2018).

Data Availability

Strains and plasmids are available upon request. The authors affirm that all data

necessary for confirming the conclusions of the article are present within the article,

figures and supplemental figures, and tables and supplemental tables. The results

of our analyses are available in the WormBase (www.WormBase.org) (Stein et al.

2001) and in our 3'UTR-centric website www.UTRome.org.

Author Contribution

HSS and MM designed the experiments. MM developed and executed the

bioinformatic analysis and 3'UTR cluster preparation. HSS performed the rescue

experiments in **Figure 5**. PLC performed the homology modelling in **Figure 4 and**

Supplemental Figure S6 and helped writing the manuscript. CG assisted with the

experiments and performed the analysis in **Supplemental Figure S1**. SO

contributed to the experiments in Figure S7-S9. MM uploaded the results to the

WormBase and UTRome.org database. MM and HSS led the analysis and

interpretation of the data, assembled the Figures, and wrote the manuscript. All

authors read and approved the final manuscript.

Funding

This work was supported by the National Institute of Health grant number

1R01GM118796.

Conflict of Interest

The authors declare that they have no competing interests.

Acknowledgements

We thank Heather Hrach for insights and review of the manuscript. We thank

Gabrielle Richardson for maintaining the *C. elegans* strains used in this manuscript.

FIGURE LEGENDS

Figure 1. The *C. elegans* members of the Cleavage and Polyadenylation Complex (CPC).

A) The CPC is composed of at least 4 independent subcomplexes named Cleavage and

Polyadenylation Specificity Complex (Blue), which canonically recognizes the PAS

hexamer 'AAUAAA'; the Cleavage Stimulation Factor Complex (Green), which binds

downstream of the cleavage site to GU rich elements; and the Cleavage Factor CFIm

(Red) and CFIIm (Orange) Complexes. CFIm recognizes the element 'UGUA' located

upstream of the PAS element. Other known required factors are the PolyA Polymerase

enzyme, the scaffolding member Symplekin and RBBP6. The name of the *C. elegans*

orthologs are shown in parenthesis. B) The human and *C. elegans* CPSF subcomplexes

are similar in amino acid composition and structure. 2-species alignments between several

655 members of the human and *C. elegans* CPSF members. Amino acids 100% conserved

656 between these two species are shown in red in the conservation bar. Yellow dotted boxes

657 show the sequence of the proteins that interact with the PAS element. Functional domains

658 are conserved. The two Kyte-Doolittle graphs in each panel indicate the hydrophobic

659 amino acids in human and *C. elegans*. C) We have used RNAi to selectively silence most

660 of the members of the CPC complex in *C. elegans.* We observed a strong embryonic

661 lethality phenotype with all the RNAi experiments performed.

663 **Figure 2.** Cluster preparation and analysis. A) Screenshots showing several

664 mapped 3'UTR clusters for genes with one or two 3'UTR isoforms. miRanda

665 predicted miRNA targets are shown for a particular 3'UTR at the bottom of this

666 Panel. B) Summary of the 3'UTRs in genes identified in this study along with the

667 number of reads mapped and clustered for each 3'UTR. C) Comparison between

668 the 3'UTRs for genes and total isoforms mapped in this study vs the UTRome v1

669 (Mangone et al. 2010) and the dataset from Jan et al., 2001.

671 **Figure 3.** The worm 3'UTRome v2. A) Top Left Panel. Nucleotide composition of

672 3'UTRs in the 3'UTRome v2. Uridine is the most abundant nucleotide within 3'UTRs

673 for *C. elegans*. Bottom Left Panel. The number of 3'UTR isoforms in each gene.

674 42% of the genes in the 3'UTRome v2 possess multiple 3'UTR isoforms. Top Right

675 Panel. 3'UTR length distribution in genes expressed with one, two, or three or more

676 3'UTR isoforms. The median 3'UTR length across these datasets is 122nt. Genes

677 with multiple 3'UTR isoforms are on average longer than genes with one 3'UTR

isoform. Bottom Right Panel. Median 3'UTR length in genes with Canonical (C) or

Variant (V) PAS elements. There is a slight increase in 3'UTR length in genes with

variant PAS elements when compared to those with canonical PAS elements. This

variation is still detected when increasing the stringency of the density of the

clusters (cd) used in this analysis. B) PAS element usage in 3'UTRs. 58.4% of

3'UTRs use the canonical PAS element 'AAUAAA' while the most common variant

PAS element is the hexamer 'AAUGAA', which occurs in 11% of genes. The

distribution of canonical PAS elements within 3'UTRs. The average distance from

the PAS element to the cleavage site is 18nt. C) Alignment of 3'UTRs at the

cleavage site. This alignment in genes with both canonical and variant PAS

elements reveals a region between the PAS element and the cleavage site we

renamed the *buffer region* in which cleavage rarely occurs. The most abundant

nucleotide at the cleavage site is an Adenosine nucleotide preceded by a

Thymidine nucleotide.

Figure 4. The sequence requirements of the *C. elegans* CPSF core complex. A)

PAS element usage of the RRYRRR motif. 3'UTRs from the 3'UTRome v2 aligned

by their cleavage site in genes with canonical or variant PAS element. The motif

RRYRRR is highlighted in yellow, and its spatial conservation is very strong in

single 3'UTR isoforms with canonical PAS elements and is enriched in those with

variant PAS elements. This RRYRRR element is maintained in 3'UTRs that have at

least two isoforms but is not strongly represented in human 3'UTR data due to the

lack of their annotation. R= Purine, Y= Pyrimidine. B) Superimposition of the cryo-

EM structure of the previously published human CPSF core complex (Clerici et al.

2018; Sun et al. 2018) to the worm CPSF core complex: cpsf-1 (CPSF-160) in blue,

pfs-2 (Wdr33) in pink, and cpsf-4 (CPSF30) in green. The PAS element binding

pocket can be fitted into the homology model. The PAS element of the RNA is

represented in yellow. The size and the selectivity of the nucleotide binding pocket

can fit other nucleotides as long as the motif is RRYRRR.

Figure 5. The Adenosine nucleotide is required at the cleavage site for correct

cleavage. A) Sequence Logos produced from 3'UTRs from genes with only 3'UTR

isoforms containing the canonical PAS element 'AAUAAA' and aligned by their

respective *buffer region* length (n=4,374). Two extra nucleotides are included

downstream of each cut site (triangle). The nucleotide distribution of the distance

between the PAS element and the cleavage site is shown in the bar chart below. B)

Example of slight variability in the cleavage site for the gene C09G9.8. While

prevalent forms are observed, the exact cleavage site can vary on several

occasions but predominantly occurs at a different Adenosine nucleotide. C) Test of

the role of the terminal Adenosine nucleotide in the cleavage reaction. The 3'end

regions of several test genes where cloned and used to prepare transgenic *C.*

elegans strains expressing this region with or without mutated terminal Adenosine

nucleotides (Red, see below). The top sequence shows the test 3'end region

(Cyan=ORF, Green=translation STOP signal, Grey=3'UTR, Red=Terminal

Adenosine nucleotide. The PAS element is underscored). The Sanger trace files

show the outcome of the cleavage site location in selected clones. Two genes are

shown (M03A1.3 and Y106G6H.9). In the case of M03A1.3, the loss of the terminal

Adenosine nucleotide sometimes forces the CPC to backtrack and cleave the

mRNAs upstream of the regular cleavage site but still at the closest Adenosine

nucleotide available. In the case of the gene Y106G6H.9, the loss of the terminal

Adenosine nucleotide forces the complex to skip the cleavage site, which

sometimes occurs at the next Purine nucleotide. Additional clones and more test

genes are shown in the **Supplemental Figure S7-S9**.

<div align="center">**REFERENCES**</div>

Bai Y, Auperin TC, Chou CY, Chang GG, Manley JL, Tong L. 2007. Crystal structure of murine CstF-77: dimeric
735   association and implications for polyadenylation of mRNA precursors. *Mol Cell* **25**: 863-875.
736 Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. *Cell* **136**: 215-233.
737 Bartel DP. 2018. Metazoan MicroRNAs. *Cell* **173**: 20-51.
738 Blazie SM, Babb C, Wilky H, Rawls A, Park JG, Mangone M. 2015. Comparative RNA-Seq analysis reveals
739   pervasive tissue-specific alternative polyadenylation in Caenorhabditis elegans intestine and
740   muscles. *BMC Biol* **13**: 4.
741 Blazie SM, Geissel HC, Wilky H, Joshi R, Newbern J, Mangone M. 2017. Alternative Polyadenylation Directs
742   Tissue-Specific miRNA Targeting in Caenorhabditis elegans Somatic Tissues. *Genetics* **206**: 757-774.
743 Chan SL, Huppertz I, Yao C, Weng L, Moresco JJ, Yates JR, 3rd, Ule J, Manley JL, Shi Y. 2014. CPSF30 and
744   Wdr33 directly bind to AAUAAA in mammalian mRNA 3' processing. *Genes Dev* **28**: 2370-2380.
745 Chen F, Chisholm AD, Jin Y. 2017. Tissue-specific regulation of alternative polyadenylation represses
746   expression of a neuronal ankyrin isoform in C. elegans epidermal development. *Development* **144**:
747   698-707.
748 Chen F, MacDonald CC, Wilusz J. 1995. Cleavage site determinants in the mammalian polyadenylation
749   signal. *Nucleic Acids Res* **23**: 2614-2620.
750 Clerici M, Faini M, Aebersold R, Jinek M. 2017. Structural insights into the assembly and polyA signal
751   recognition mechanism of the human CPSF complex. *Elife* **6**.
752 Clerici M, Faini M, Muckenfuss LM, Aebersold R, Jinek M. 2018. Structural basis of AAUAAA polyadenylation
753   signal recognition by the human CPSF complex. *Nat Struct Mol Biol* **25**: 135-138.
754 de Vries H, Ruegsegger U, Hubner W, Friedlein A, Langen H, Keller W. 2000. Human pre-mRNA cleavage
755   factor II(m) contains homologs of yeast proteins and bridges two other cleavage factors. *EMBO J*
756   **19**: 5895-5904.
757 Diag A, Schilling M, Klironomos F, Ayoub S, Rajewsky N. 2018. Spatiotemporal m(i)RNA Architecture and 3'
758   UTR Regulation in the C. elegans Germline. *Dev Cell* **47**: 785-800 e788.
759 Frokjaer-Jensen C, Davis MW, Hopkins CE, Newman BJ, Thummel JM, Olesen SP, Grunnet M, Jorgensen EM.
760   2008. Single-copy insertion of transgenes in Caenorhabditis elegans. *Nat Genet* **40**: 1375-1383.

Gerstein MB Lu ZJ Van Nostrand EL Cheng C Arshinoff BI Liu T Yip KY Robilotto R Rechtsteiner A Ikegami K et
al. 2010. Integrative analysis of the Caenorhabditis elegans genome by the modENCODE project.
Science **330**: 1775-1787.
Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE. 2018. UCSF ChimeraX:
Meeting modern challenges in visualization and analysis. *Protein Sci* **27**: 14-25.
Griffiths-Jones S, Grocock RJ, van Dongen S, Bateman A, Enright AJ. 2006. miRBase: microRNA sequences,
targets and gene nomenclature. *Nucleic Acids Res* **34**: D140-144.
Haenni S, Ji Z, Hoque M, Rust N, Sharpe H, Eberhard R, Browne C, Hengartner MO, Mellor J, Tian B et al.
2012. Analysis of C. elegans intestinal gene expression and polyadenylation by fluorescence-
activated nuclei sorting and 3'-end-seq. *Nucleic Acids Res* **40**: 6304-6318.
Helmling S, Zhelkovsky A, Moore CL. 2001. Fip1 regulates the activity of Poly(A) polymerase through
multiple interactions. *Mol Cell Biol* **21**: 2026-2037.
Hwang HW, Park CY, Goodarzi H, Fak JJ, Mele A, Moore MJ, Saito Y, Darnell RB. 2016. PAPERCLIP Identifies
MicroRNA Targets and a Role of CstF64/64tau in Promoting Non-canonical poly(A) Site Usage. *Cell
Rep* **15**: 423-435.
Jan CH, Friedman RC, Ruby JG, Bartel DP. 2011. Formation, regulation and evolution of Caenorhabditis
elegans 3'UTRs. *Nature* **469**: 97-101.
John B, Enright AJ, Aravin A, Tuschl T, Sander C, Marks DS. 2004. Human MicroRNA targets. *PLoS Biol* **2**:
e363.
Kaufmann I, Martin G, Friedlein A, Langen H, Keller W. 2004. Human Fip1 is a subunit of CPSF that binds to
U-rich RNA elements and stimulates poly(A) polymerase. *EMBO J* **23**: 616-626.
Kuhn U, Wahle E. 2004. Structure and function of poly(A) binding proteins. *Biochim Biophys Acta* **1678**: 67-
84.
Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357-359.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project
Data Processing S. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**:
2078-2079.
Mandel CR, Kaneko S, Zhang H, Gebauer D, Vethantham V, Manley JL, Tong L. 2006. Polyadenylation factor
CPSF-73 is the pre-mRNA 3'-end-processing endonuclease. *Nature* **444**: 953-956.
Mangone M, Macmenamin P, Zegar C, Piano F, Gunsalus KC. 2008. UTRome.org: a platform for 3'UTR
biology in C. elegans. *Nucleic Acids Res* **36**: D57-62.
Mangone M, Manoharan AP, Thierry-Mieg D, Thierry-Mieg J, Han T, Mackowiak SD, Mis E, Zegar C, Gutwein
MR, Khivansara V et al. 2010. The landscape of C. elegans 3'UTRs. *Science* **329**: 432-435.
Martin G, Gruber AR, Keller W, Zavolan M. 2012. Genome-wide analysis of pre-mRNA 3' end processing
reveals a decisive role of human cleavage factor I in the regulation of 3' UTR length. *Cell Rep* **1**: 753-
763.
Matlin AJ, Clark F, Smith CW. 2005. Understanding alternative splicing: towards a cellular code. *Nat Rev Mol
Cell Biol* **6**: 386-398.
Mayr C, Bartel DP. 2009. Widespread shortening of 3'UTRs by alternative cleavage and polyadenylation
activates oncogenes in cancer cells. *Cell* **138**: 673-684.
Perez Canadillas JM, Varani G. 2003. Recognition of GU-rich polyadenylation regulatory elements by human
CstF-64 protein. *EMBO J* **22**: 2821-2830.
Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**: 841-842.
Ryan K, Calvo O, Manley JL. 2004. Evidence that polyadenylation factor CPSF-73 is the mRNA 3' processing
endonuclease. *RNA* **10**: 565-573.

Schafer P, Tuting C, Schonemann L, Kuhn U, Treiber T, Treiber N, Ihling C, Graber A, Keller W, Meister G et
808   al. 2018. Reconstitution of mammalian cleavage factor II involved in 3' processing of mRNA
809   precursors. *RNA* **24**: 1721-1737.
810   Sheets MD, Ogg SC, Wickens MP. 1990. Point mutations in AAUAAA and the poly (A) addition site: effects
811   on the accuracy and efficiency of cleavage and polyadenylation in vitro. *Nucleic Acids Res* **18**: 5799-
812   5805.
813   Stein L, Sternberg P, Durbin R, Thierry-Mieg J, Spieth J. 2001. WormBase: network access to the genome
814   and biology of Caenorhabditis elegans. *Nucleic Acids Res* **29**: 82-86.
815   Sun Y, Zhang Y, Hamilton K, Manley JL, Shi Y, Walz T, Tong L. 2018. Molecular basis for the recognition of
816   the human AAUAAA polyadenylation signal. *Proc Natl Acad Sci U S A* **115**: E1419-E1428.
817   Takagaki Y, Manley JL. 2000. Complex protein interactions within the human polyadenylation machinery
818   identify a novel component. *Mol Cell Biol* **20**: 1515-1525.
819   Tian B, Graber JH. 2012. Signals for pre-mRNA cleavage and polyadenylation. *Wiley Interdiscip Rev RNA* **3**:
820   385-396.
821   Tian B, Manley JL. 2017. Alternative polyadenylation of mRNA precursors. *Nat Rev Mol Cell Biol* **18**: 18-30.
822   Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de Beer TAP, Rempfer C,
823   Bordoli L et al. 2018. SWISS-MODEL: homology modelling of protein structures and complexes.
824   *Nucleic Acids Res* **46**: W296-W303.
825   West SM, Mecenas D, Gutwein M, Aristizabal-Corrales D, Piano F, Gunsalus KC. 2018. Developmental
826   dynamics of gene expression and alternative polyadenylation in the Caenorhabditis elegans
827   germline. *Genome Biol* **19**: 8.
828   Yang Q, Coseno M, Gilmartin GM, Doublie S. 2011. Crystal structure of a human cleavage factor
829   CFI(m)25/CFI(m)68/RNA complex provides an insight into poly(A) site recognition and RNA looping.
830   *Structure* **19**: 368-377.
831   Yang Q, Gilmartin GM, Doublie S. 2010. Structural basis of UGUA recognition by the Nudix protein CFI(m)25
832   and implications for a regulatory role in mRNA 3' processing. *Proc Natl Acad Sci U S A* **107**: 10062-
833   10067.
834   Yang W, Hsu PL, Yang F, Song JE, Varani G. 2018. Reconstitution of the CstF complex unveils a regulatory
835   role for CstF-50 in recognition of 3'-end processing signals. *Nucleic Acids Res* **46**: 493-503.
836   Yang XC, Sullivan KD, Marzluff WF, Dominski Z. 2009. Studies of the 5' exonuclease and endonuclease
837   activities of CPSF-73 in histone pre-mRNA processing. *Mol Cell Biol* **29**: 31-42.
838   Zhelkovsky A, Helmling S, Moore C. 1998. Processivity of the Saccharomyces cerevisiae poly(A) polymerase
839   requires interactions at the carboxyl-terminal RNA binding domain. *Mol Cell Biol* **18**: 5942-5951.
840   Zhu Y, Wang X, Forouzmand E, Jeong J, Qiao F, Sowd GA, Engelman AN, Xie X, Hertel KJ, Shi Y. 2018.
841   Molecular Mechanisms for CFIm-Mediated Regulation of mRNA Alternative Polyadenylation. *Mol*
842   *Cell* **69**: 62-74 e64.

Figure 1

A

B

C

A

B

# 3'UTRs in genes (isoforms)			# reads used	
total	in 3'UTRome v1	new	mapped	in each cluster
14,808 (23,159)	11,377 (16,941)	4,638 (6,218)	4,838,905	220 (median)

clusters >=10%

C

Figure 3

A

1 isoform   ☐: RRYRRR

AAUAAA — variants

2,845 3'UTRs
2,207 3'UTRs

-100   cleavage site   0   -100   cleavage site   0

Median freq.=1
Average freq.=2.5

Median freq.=1
Average freq.=2

>1 isoforms

AAUAAA   variants

4,789 3'UTRs   5,202 3'UTRs

worm 3'UTRome v2

AAUAAA   variants

12,265 3'UTRs   7,409 3'UTRs

human 3'UTRome (hg19)

AAUAAA   variants

14,219 3'UTRs   2,207 3'UTRs

B

Homology model (*C. elegans*)

Cryo-EM structure (*Homo sapiens*)

A

genes w/ unique 3'UTRs with canonical PAS=4,374

B

C