Population genomics and pathotypic evaluation of the bacterial leaf blight

pathogen of rice reveals rapid evolutionary dynamics of a plant pathogen

Jinshui Zheng[a,1], Zhiwei Song[b,1], Dehong Zheng[a], Huifeng Hu[a], Hongxia Liu[c], Yancun

Zhao[b], Ming Sun[a], Lifang Ruan[a,2], Fengquan Liu[b,2]

[a] State Key Laboratory of Agricultural Microbiology, Hubei Key Laboratory of

Agricultural Bioinformatics, Huazhong Agricultural University, Wuhan, 430070, China.

[b] Institute of Plant Protection, Jiangsu Academy of Agricultural Sciences, Nanjing

210014, China.

[c] Department of Plant Pathology, College of Plant Protection, Nanjing Agricultural

University, Nanjing 210095, China

[1] These authors contributed equally to this work.

[2] Corresponding authors：Fengquan Liu (fqliu20011@sina.com) and Lifang Ruan

(ruanlifang@mail.hzau.edu.cn).

Keywords: Plant pathogen, Genomic epidemiology, Large-scale virulence test, Rapid

virulence dynamics

Abstract

Xanthomonas oryzae pv. *oryzae* (*Xoo*) causes bacterial blight disease, which reduces crop yield by up to 50% in rice production. Despite its substantial threat on food production worldwide, knowledge about its population structure, virulence diversity and the relationship between them is limited. We used whole-genome sequencing to explore the diversity and evolution of *Xoo* during the past 30 years in the main rice-planting areas of China. Six separate lineages were revealed by phylogenomic analysis, with CX-5 and CX-6 predominating in the population for decades. The recent sporadic outbreaks were respectively caused by *Xoo* derived from these lineages especially the two major ones. The lineage and sub-lineage distribution of isolates strongly correlated to their geographical origin, which was found to be mainly determined by the planting of the two major rice subspecies, *indica* and *japonica*. Large-scale virulence testing was conducted to evaluate the diversity of pathogenicity for *Xoo.* We found rapid virulence dynamics against rice, and its determinant factors including genetic background of *Xoo*, rice resistance genes and the planting environment of rice. Genetic background was investigated deeply by comparative genomics, which indicates that transposition events contributing the most to evolution of the *Xoo* genome and the rapid diversification of virulence. This study provided a good model to understand the evolution and dynamics of plant pathogens in the context of interaction with their hosts which are influenced by both geographical conditions and farming practices.

Introduction

Population genomics is a powerful tool for understanding the formation and evolution

of bacterial pathogens of human and some important domestic animals [1-6]. It has also

provided new strategies for the detection, prevention and control of important human

pathogens [7,8]. Though plant pathogens are serious threat to food security worldwide,

their evolution and dynamics at population scale in the content of modern agricultural

practice have not been well understood. A few studies have used genome-based

methods to study their population structure, which have provided some new insights

into evolution and transmission of some important crop pathogens [9-13].

Xanthomonas oryzae pv. *oryzae* (*Xoo*) causes bacterial blight (BB) disease of rice

(*Oryza sativa*) and has been considered one of the top 10 plant bacterial pathogens

based on its scientific/economic importance [14]. It is a notorious destructive pathogen

which can cause a considerable reduction in rice production in both temperate and

tropical regions, especially in Asia [15]. On average, BB can lead to 20%~30% overall

yield reduction with some severe cases even causing up to 50% loss [16]. As the world's

largest producer and consumer of rice, China has suffered serious BB outbreaks and

huge food loss since the 1930s, with the most serious damage occurring between 1950s

and 1980s [17,18]. Breeding resistant rice cultivars has been proved to be the most

effective method for controlling this pathogen. Since 1980s, BB-resistant rice varieties

have been widely planted in China and have significantly reduced the loss caused by

BB [19-21]. However, sporadic outbreaks still occurred in different areas of China after

the year of 2000 [22,23]. It is unknown whether the recent outbreaks relate to changes in the genetics and virulence of the past *Xoo* populations in China, allowing it to overcome host defenses. Only few studies focused on population structure of *Xoo* from limited rice-planting areas of China were conducted by DNA fingerprinting methods [24-26]. So very little is known about the evolution, spread and dynamics of *Xoo* involved in different outbreaks during the past decades around the whole country in the context of rapid development of modern agricultural science and technology.

The virulence differentiation among different strains of *Xoo* which was evaluated by defining different races or pathotypes, was frequently studied to infer its virulence dynamics and diversity against rice host. Strains of the same race share a common pathogenic phenotype in a set of tested host cultivars. Rice lines carrying different resistance genes (R genes) determine the race of *Xoo*. Up to now, tens of races of *Xoo* were identified in different rice-planting locations worldwide [22,27-31]. The pathotypes of *Xoo* showed huge diversity among different locations and rapid dynamics along the isolated times. It was predicted that many factors including genetic background of *Xoo* contributing to its virulence diversity [28-30]. However, many crucial details were overlooked due to the intrinsic low sensitivity and stability of the typing method.

Different resistant cultivars carrying bacterial blight R genes have been planted in the different Asian countries for decades, including China [32,33]. Resistance from R genes carried by rice cultivars plays a crucial role of determining *Xoo* evolution during its interaction with rice hosts. Recent study has showed that R gene *Xa4* has largely

affected the race composition of *Xoo* from the Philippines [33]. The pathogenic

diversity of *Xoo* in China has been studied previously [22,23,25,26]. Race compositions

showed huge diversity and rapid dynamics among different areas along different time

points [34,35]. However, it is unknown the genetic basis for the pathogenicity diversity

and dynamics of *Xoo* from China, which means the relationship between genetic

population and the ability *Xoo* to overcome different R genes has not been studied.

Moreover, the evolutionary force behind the genetic and pathogenic diversity and

dynamics is also not clear.

In this study, a comprehensive population genomic study was combined with a large-

scale determination of virulence of *Xoo* that have been isolated in the past 30 years in

China, to obtain a framework of genetic dynamics and virulence diversity over time

and space. Results from this work shed new light on the evolution of this important

plant pathogen.

Results

Population structure of *Xoo* from China

To understand the population structure of *Xoo* in China, genomes of the 247 strains

were sequenced, including 237 isolated from China, 4 and 6 representative ones from

Japan and Philippines, respectively (Table S1). Reads were mapped to the genome of

PXO99[A]; SNPs were called by Samtools. After the prophage region, repeat sequences

and recombinant regions being filtered, the core SNPs contained 5,271 variable sites.

Two methods were used to outline the population structure based on the core SNPs,

Maximum likelihood (ML) phylogenetic analysis and a tree-independent hierarchical

Bayesian clustering (BAPS).

All *Xoo* from China were clustered into 6 lineages (CX-1 to CX-6), with more than 70%

members belonging to CX-5 and CX-6 (Figure 1). The three major rice production areas,

including South China, Yangtze Valley area and North China, were displayed in Figure

2a. We annotated the lineage-specific tree with the time and space information of the

Xoo strains isolated (Figure 1, 2b and 2c). CX-1 and CX-2 were most frequently

represented in South China at the year of 2003, of which most isolates were from

Yunnan province. The *Xoo* belonging to CX-3 were isolated from North China in 1984,

2003 and 2014, with most of which from Northeast China in 2014. CX-5 and CX-6

were nationally distributed during the past 30 years; while most members in CX-5 were

dominant in South China and Yangtze Valley, and CX-6 were more frequently from

North China and Yangtze Valley.

Phylogeography of CX-5 and CX-6

To investigate more details about evolution and dynamics of *Xoo* in China, we focused

on the two major lineages, CX-5 and CX-6. Phylogenetic analysis based on core SNP

alignment (Figures 3a and 3b) and pairwise SNP distance between isolates were

performed for both lineages (Figure 3c). Both CX-5 and CX-6 can be divided into 5

sub-lineages, respectively (Figure 3a and 3b).

CX-5.1, CX-5.2, and CX-5.5 were restricted in South China, while CX-5.3 and CX-5.4

can be found in some places of both South China and Yangtze Valley (Figure 3a). When

time was considered, we found that all these sub-lineages persisted at these places for

at least 30 years. Then we focused on the recent sporadic outbreaks being attributed to this lineage. Besides clustered together on the tree, we considered epidemiologically and genomically linked outbreak with an average SNP pairwise distance less than 5. In 2003, the outbreak in Hunan province of Yangtze Valley was caused by CX-5.4 (HN-2003), while the outbreak in Guangdong province of South China ascribed to two sub-population of CX-5.5 (GD-2003-1 and GD-2003-2). One cluster of CX-5.3 caused outbreak in Guangxi province of South China in 2014 (GX-2014).

Seven out of 10 isolates in CX-6.1 were from different places of South China. This sub-lineage was the most diverse one in CX-6 with long branches. In CX-6.2, *Xoo* from North China and Yangtze Valley were dominant on different branches, respectively. CX-6.3 and CX-6.4 were most frequently found in Yangtze Valley, while *Xoo* from North China predominated in CX-6.5. Similar as CX-5, all these sub-lineages have contributed to BB on rice for a long time in China. These sub-lineages totally caused at less 4 local outbreaks in 2003 and 2014. In 2003, the local outbreaks in two provinces of Yangtze Valley (Henan, and Jiangsu) were caused by CX-6.2 (HeN-2003) and CX-6.4 (JS-2003), respectively, while the one in North China (Jilin province) was ascribed to CX-6.5 (JL-2003). Outbreak in Jiangsu province (Yangtze Valley) in 2014 was caused by one sub-population of CX-6.4 (JS-2014).

Chinese *Xoo* diversity and dynamics in the context of rice host

There are two major domesticated rice subspecies *Oryza sativa japonica* and *indica* existing in different areas of China; and most of rice varieties were derived from these two subspecies [34,35]. Rice planted in high-altitude areas of Yunnan province from

South China and North China belong to subspecies *japonica* (http://www.ricedata.cn).

Xoo isolated from these areas were respectively clustered into 3 lineages, CX-1, CX-2,

CX-3 and one sub-lineage of CX-6 (CX-6.5) (Figure 1 and Figure 3b). Lineage CX-5

mainly contained *Xoo* isolated from rice in South China where subspecies *indica* is

planted (Figure 3a). CX-6 also contained *Xoo* isolated from rice of *japonica* from

Yangtze Valley (CX-6.4), *indica* from Yangtze Valley (CX-6.3) and some *indica* from

South China (CX-6.1), indicating the widest adaptation to different rice varieties of *Xoo*

in this lineage (Figure 3b). In CX-6.2, old *Xoo* were dominant on indica, while recent

isolated ones were almost from *japonica*, suggesting a putative host switch in this sub-

lineage.

Taken together, the distribution of Chinese *Xoo* lineages appeared to be impaired by

biogeography and rapid dynamics over isolated time, which may be mainly due to the

distribution and dynamics of the two major subspecies of rice, *japonica* and *indica*.

Diversity of *Xoo* from different counties of Asia

To place *Xoo* from China into a global context, phylogenomic analysis was performed

on the core genome SNPs of 109 *Xoo* genomes available in Genbank (including 100

from India, 8 from the Philippines and 1 from Japan), and the 247 sequenced ones in

this study. Phylogenetic analysis revealed a similar topological structure of Asian *Xoo*

to that of Chinese *Xoo*, with small differences in some branches (Figure S1). Eleven

genetic lineages were identified for the *Xoo* population. *Xoo* from India and the

Philippines were respectively assigned into PX-A to PX-C and IX-I to IX-V according

to previous studies [36,37]. Eight lineages displayed strict geographic distribution

features, most of them being specific to one or at least a limited amount of rice-planting

areas. Besides the three Chinese lineages (CX-1, CX-2 and CX-3), PX-B and PX-C

were mainly represented in the Philippines, and IX-I to IX-III were from India. CX-4

contained old *Xoo* from China and some recently isolated ones from India belonging to

IX-V. For CX-5 and CX-6, besides both represented the major *Xoo* from China, they

also contained a few *Xoo* from Japan, India and Philippines, with PX-A of Philippines

being nested within the former one and IX-IV from India being encompassed among

the members of the last one, indicating frequently transmission of these 2 lineages not

only among different places of China but also among different Asian countries.

Rapid virulence dynamics of *Xoo* strains isolated in China during the past 30 years

As describe above, the population structure *Xoo* maintains relative stability during the

past 30 years. We next inferred the pathogenic diversity and dynamics of these *Xoo*

strains. Based on the interactions between *Xoo* strains and rice lines, most of the tested

Xoo were classified into 9 pathogenic races (Table S1and Figure 4a). The top two races

with most members, R5 and R8, contain 26% and 18% of the isolates, respectively.

The three major rice-planting regions had different race compositions (Figure 4b).

Yangtze River Valley mainly contained *Xoo* from races R5 and R8, with the respective

percentages of the total population being 50% and 30%, respectively. *Xoo* isolates from

North China mostly belongs to R7 and R5, together accounts for about two third of the

total population. Five of the nine *Xoo* races were well represented in the *Xoo* population

from South China, with each of them account for more than 10% of the total population.

The virulence dynamics of *Xoo* during the last 30 years was inferred by analyzing race

distribution in China at different period (Figure 4c). Strains isolated before the year of

2000 were mainly of the races of R5, R4, R8 and R1, while most *Xoo* isolated in 2003–

2004 were R8, R7, R5 and R6; and about 45% of the isolates in 2014 were R5.

We also found different lineages have different race compositions, indicating genetic

background playing important roles in the interaction between *Xoo* and rice hosts

(Figure 4d). Compared to other lineages, the CX-6 *Xoo* isolates fall mostly within the

races of R5, R8 and R4, while have reduced representation in R1, R2 and R6. Moreover,

the rapid dynamics of pathogenic features can be inferred by focusing on the races of

Xoo isolated from the same places at the same time and belonging to the same lineage

(Figure 1). For example, most *Xoo* from Yunnan, one province of South China, in 2003

were clustered in CX-1 and CX-2. The 19 isolates in CX-1 represented 8 of the 9 races

described above, and the 14 strains in CX-2 belonged to 6 races. Even *Xoo* isolates from

the same outbreak were allocated into different races, with 16 *Xoo* in JL-2003 belonging

to 3 different races and 10 isolates in GD-2003-2 representing 2 races.

Resistance gene of rice contributes to the interaction between *Xoo* and rice

As each of the near-isogenic rice lines used in this study contains one known resistance

(R) gene, we studied the interaction between *Xoo* and rice host by focusing on the

capacity of *Xoo* to overcome the resistances conferred by these genes (Figure 1, Tables

1 and 2). When the time of isolation was considered, we found that *Xoo* from different

times have different capacity to overcome different R genes (Table 1). R gene *xa5*

showed the most effective resistance against *Xoo* all the time. At different periods, more

than 90% *Xoo* could not infect the rice line IRBB5, which contained gene *xa5*. The

resistance mediated by R gene *Xa3* was effective during the period of 1990 to 2004.

However*,* rice containing this R gene only showed resistance against 5% of *Xoo* in 2014.

Rice cultivars carrying R gene *Xa4* are resistant to more than 60% of *Xoo* isolated from

1970s to 2000 and 2014. However, they are only resistant to about 30% of the isolates

recovered in 2003 and 2004.

To determine whether the *Xoo* genetic background contributes to the interaction

between *Xoo* and rice lines, we studied the resistance capacity of R genes against *Xoo*

from different lineages (Table 2). We found that resistance provided by R genes against

Xoo are indeed depended on the genetic background of *Xoo*. R genes *Xa2* and *Xa14*

effectively confer resistance against *Xoo* from Lineages CX-1, CX-2 and CX-3,

preventing infection by more than 70% of *Xoo* strains from these lineages. Similarly, R

gene *Xa3* and *Xa4* showed potential resistance against *Xoo* from CX-1 and CX-2, and

CX-2 and CX-3, respectively. No R gene except *xa5* conferred resistance to more than

50% of *Xoo* strains from CX-5 and CX-6, which is consistent with the situation that

CX-5 and CX-6 had much more extensive distribution across different areas of China

during the past decades, comparing to the other Chinese *Xoo* lineages. Among the sub-

lineages of CX-5 and CX-6, *Xa2* was only effective against *Xoo* of CX-6.5, with

effectiveness of 90%, while *Xa4* could confer the resistance against more than 60% *Xoo*

from these two lineages except for CX-5.1 and CX-6.5; *Xa14* showed potential medium

level of resistance against *Xoo* from CX-6.1 and CX-6.5 (Table S2).

Genome dynamics of *Xoo* is mainly determined by transposition events

To study the genomic evolution contributing to the rapid dynamics of virulence, several

Xoo isolates from different lineages were chosen to complete genome for comparative

genomics. All proteins of the 22 *Xoo* with complete genomes were clustered into 7542

groups, with 3513 belonging to core genes. The pan genes plot showed that Asian *Xoo*

has an open pan genome (**Figure S2**).

It was reported that *Xoo* genome contains numerous insert sequence elements, so we

firstly studied the distribution of transposase genes to understand the dynamics of *Xoo*

genomes. Each *Xoo* genome contained more than 800 transposase genes, taking up

above 16% of the total genes, with most of them from IS5, IS3, IS1595, IS701, IS30

and IS630 (**Figure 5 and Table S3**). In the core gene sets, 8%, (266 of 3543) were

predicted to encode transposases. However, 71% (1251 of 1771) and 62% (942 of 1523)

of the distributed gene and unique gene families were transposases (**Figure S3**). It

suggests that transposon elements contribute the most to the diversity of *Xoo* genomes.

Based on the analysis of gene presence/absence analysis, the core genome of *Xoo* can

be split into about 200 conserved blocks by less conserved genome fragments (Figure

5). Among the blocks, there were genome segments with more sequence diversity, and

most transposases were in these regions.

As a rice pathogen, the host selection was reported to play crucial roles on the evolution

of *Xoo*. Type III effectors (T3Es) were found to be crucial to the pathogenicity of

Xanthomas spp. on plant hosts. Among them, transcription activator-like effectors

(TALEs) (Avirulence protein AvrBs3) are the most important in *Xoo* during the

interaction with rice. TALEs function as transcriptional factors activate expression of

target genes in the host in a sequence specific manner, resulting in enhanced bacterial

growth and development of disease symptom (Bogdanove et al 2010). Most TALEs

from *Xoo* activate the expression of *SWEET* genes in rice to promote susceptibility [32].

Significant variations in the number of TALEs was observed among the 22 *Xoo* with

complete genomes, ranging from 11 to 21 (**Table S4**). Half of the TALE families (21

out of 42) were only predicted in one or two isolates. And some families only distributed

or lost in some special lineages. *Xoo* from the same lineage had some common and

different families of TALEs. Even *Xoo* isolated in the same geographical location at the

same time with the same phylogenetic relationship, JL28 and JL33, contain different

compositions of TALE families. This suggests that interaction among rice and *Xoo* can

determine rapid diversity of TALE effectors. When locations of TALE genes were

identified, we found that almost all of them were in the variable regions where enrich

in transposase genes (Figure 5). Conversely, other T3Es were very conserved, almost

all of them located in the conserved genome blocks with less transposons (Figure 5).

Discussion

Most *Xoo* clusters identified by traditional molecular typing methods from several

Asian countries showed strict geographical distribution [28-30,38]. Previous report

focusing on *Xoo* from India through population genomics defined 5 lineages, with each

of them showing a restricted-region distribution [37]. In this study, based on population

genomics, we found that there is a general correspondence between the areas of *Xoo*

isolated and most of Asian lineages and some sub-lineages in CX-5 and CX-6 (Figures

1, 3 and S2). Moreover, Asian *Xoo* was reported to be very different from those of

Africa and North America regions [39,40]. This suggests that the geographical factors

as affected by climate, soil types, or the choice of different rice cultivars and varieties

may influence on the diversity of *Xoo*. Though thousands of rice cultivars have been

planted in different areas of China, almost all of them were derived from the two

subspecies, *indica* and *japonica*, or hybrid varieties between them

(http://www.ricedata.cn). Information of the *Xoo* host subspecies in China indicates that

variety of rice was one of the major factors forces on the formation and dynamics of

Chinese *Xoo* population. Rice varieties planted in different counties and different areas

of a country are determined by geographical environment, climate factors and local

agricultural policy [41-43]. Taken together, geography-specific rice cultivars in

different countries or different areas in the same country contribute the most to the

genetic diversity of *Xoo* infecting them.

Development resistance for novel cultivars of crops and vegetables, is proved to be the

most effective strategy fighting against bacterial and fungi pathogens [44-47]. But the

durability of plant resistance can be neutralized by evolutionary changes in pathogen

populations. Knowledge about pathogen population evolution and dynamics under the

force from the resistant plant are crucial for further cultivar development [48]. Though

resistance breeding of rice have effectively prevented the unprecedented outbreak of

BB since 1990s, sporadic outbreaks occurred in different areas of China after the year

2000 [49]. Genomic epidemiology analysis indicates all *Xoo* strains causing the recent

epidemic outbreaks except those in CX-1 were derived from the ones leading to heavy

losses in the history. On the other hand, only the minor lineage CX-4 was not

represented by the recent outbreaks, so the diversity of the *Xoo* population was not reduced due to wide planting of BB-resistant rice varieties. For the current practice of rice production, the major BB threat in the three rice-planting areas of China is still mainly from CX-5 and CX-6 as they were present in history, which call for different detection and prevention strategies for different areas and rice varieties.

The virulence diversity of *Xoo* in China against rice was well investigated during the past 30 years, however, the relationships between population genetic structure and pathotype of *Xoo* were not explored well [22-24,26,49]. This study shows that races of *Xoo* from South China were more diverse than those from North China and Yangtze Valley areas. In line with this, phylogenetic analysis revealed that *Xoo* from South China are more diverse compared to other places of China (Figure 1). Some rice growing regions in South China, particularly in the Yunnan province, have three harvests per year and use different rice cultivars for each harvest [50,51]. The greater genetic and virulence diversity of *Xoo* thus could be caused by a greater diversity of rice cultivars in use. Our result also showed that there were diverse races in the same lineage, for example, 85% *Xoo* from lineage CX-6 could be assigned to 8 races (Figure 4d), indicating rapid virulence dynamics in the same lineage. Moreover, the isolates from the same local outbreak even belonged to different races. Although *Xoo* strains isolated in China were consistently assigned to one of the 6 Chinese lineages, the dominating population of *Xoo* races in specific areas displayed rapid dynamics and high level of diversity during the past 30 years (Tables 1 and 2). Rapid virulence dynamics with relative stability of the standing genetic variability of *Xoo* population indicates fast

adaptation to host during interaction for this plant pathogen.

In China, since the 1980s, *indica* hybrid rice (especially Shanyou 63) had been planted

on a large scale [52]. Later, bacterial blight resistant rice varieties, mainly the *india* rice

carrying *Xa4* and *japonica* rice carrying *Xa3*, were grown on a large scale for a long

time [31,52]. Soon after these rice varieties widely used for rice production, the

effectiveness of these R genes conferring resistance against infection causing by *Xoo*

was significantly compromised (Table 1). Actually, the resistance abilities of these two

R genes were also overcome frequently by *Xoo* population in other Asian countries

[30,38,53]. R gene *xa5* showed very good resistance (above 95%) to *Xoo* from all

lineages. Previously, isolates of *Xoo* having compatibility with *xa5* were only reported

from Indian Lineages IX-I and IX-III, and some Philipps and Korea isolates [27,54,55].

The only 2 Chinese *Xoo* from Yunnan province (YN24 and YN04-5) belonging to IX-I

and IX-III showed good infection ability to rice with *xa5*, suggesting putative

transmission of *Xoo* from India to South China. Besides, several isolates from CX-5

and CX-6, especially those obtained recently, can overcome the resistance contributed

by *xa5*, indicating different origins of these resistance compatibilities (Tables 1 and 2).

Since rice variants with gene *xa5* were less used in China, the development of its

compatibility could not be due to a direct selection pressure of this R gene.

It was reported recently that different *Xoo* population from the Philippines evolved

different mechanisms to adapt to the R gene *Xa4* [22]. Chinese *Xoo* Isolates from

different lineages and sub-lineages showed distinct capabilities to fight against the same

R gene. Overall, lineages and sub-lineages with stricter geographical distribution can

be more easily defeated by R genes than the wider distributed ones (Tables 2 and S2).

The underlying genetic determinations of these differences call for advanced

comparative genomics and wet experiments. However, there is no lineage or sub-

lineage with all isolates incompatible with one of the tested R genes, which means that

Xoo with different genetic backgrounds can adapt to different rice hosts with variable

resistances. This suggests that the introduction of new R genes could reduce the loss

causing by *Xoo* but can't change the genetic diversity of the *Xoo* population, as no

lineage or sub-lineage can be eliminated thoroughly. This is partially explained by that

R genes usually restrict the pathogens to the initial infection site, but have no killing

abilities [56].

Most of the R genes cloned up to now were reported to mediate resistance associated

with the TALEs during the interaction between rice and *Xoo* [57,58]. The TALE

proteins are the most import effectors used by *Xoo* to interact with rice, enforcing

virulence, proliferation, and dissemination against rice [59]. All the TALE genes were

found in the several variable regions where enriched in transposase genes belonging to

families IS3, IS5, IS30 and IS701, respectively. We also find that other T3Es didn't

have much diversity. It is predicted that transposons containing these kinds of

transposons contribute to the rapid dynamics of virulence of *Xoo* isolates mainly

through affecting the compositions and structure of TALEs. Similar situation was

reported that Tn3-like transposon was proved to play a major role of the spread of

pathogenicity determinants in *Xanthomonas* spp. [60]. Actually, transposon-rich

genome regions were reported to play crucial roles in rapid evolution of effectors after

host changes in the Irish potato famine pathogen *Phytophthora infestans* [61].

Conclusion

The introduction of several resistance genes into rice tremendously reduced the damage

of this notorious plant pathogen; only sporadic events were reported after 2000s [19,21-

23]. However, this study found that the genome evolution and virulence dynamics of

Xoo are extraordinary rapid, indicating its ability to overcome the resistance conferred

by the current resistance genes and cause the potential for large-scale outbreaks in the

future. It is therefore prudent to continue surveillance of disease outbreaks caused by

this plant pathogen, and to develop novel strategies for its control. The rice varieties

and resistance genes were predicted to be crucial to the evolution and dynamics of *Xoo*.

The subspecies of *Oryza sativa* used for rice production and resistance gene

introduction into the rice planted were mainly decided by human. The results from the

population genomics and large-scale virulence tests provided new insight into how

plant pathogens evolve and spread in the context of human agricultural activities.

Resistance genes from rice can be used to fight against *Xoo*, but *Xoo* can rapid develop

strategies to overcome these resistances. Transposon elements were proved to play a

crucial role in the evolution progress during the interaction between pathogen and it

rice host. The interaction between plant pathogen and host was reported to conform to

zig-zag-zig model [62], this study presented a well-documented example for co-

evolution of a pathogen and its host.

Materials and Methods

Genome sequencing

Xoo strains from 3 countries were used in this study (Table S1). The major strains were isolated from China. And the represented *Xoo* strains for 4 different races from Japan and 7 different races from Philippines were also sequenced.

The bacterial DNA of overnight cultures (30 °C) was extracted and purified using the Easy-DNA kit (Invitrogen, USA) following the manufacturer's protocol. Total DNA was sequenced by with Illumina HiSeq 2000, 2500, or 4000 to produce pair-end reads with lengths of 100, 125 and 150 bp. For each *Xoo*, raw reads were assessed with the FastQC tool (https://github.com/s-andrews/FastQC) and quality filtered using Trimmomatic (https://github.com/timflutre/trimmomatic). The filtered reads were error-corrected by library with Quake to produce clean reads [63]. Most of the analysis were conducted with the PGCGAP pipeline developed by our bioinformatic team [64]; and more details are described as below.

Mapping and SNP calling

To study the population structure of *Xoo* from China, reads from each strain sequenced was mapped onto the reference genome PXO99^{A}, which is used as a model strain for studying the interaction between *Xoo* and rice host and is the first *Xoo* with complete genome sequence [65], using BWA [66]. Variant detection was performed using Samtools mpileup (https://github.com/samtools/samtools) and filtered with a minimum mapping quality of 30. SNP was excluded if its coverage was less than 10% or more than 200% of the average coverage, if it was not supported by at least 5 reads on each

strand. Phage regions and repetitive sequences of the PXO99^A genome were predicted

by PHASTER (http://phaster.ca) and RepeatScout (https://bix.ucsd.edu/repeatscout),

respectively. Snippy (https://github.com/tseemann/snippy) was used to generate the

whole genome alignment of all studied *Xoo*, and then the recombination was detected

by Gubbins (https://sanger-pathogens.github.io/gubbins) on the core genome alignment

after prophage and repeat regions were filtered. SNPs located within phage regions,

repetitive sequences or recombinant regions were excluded.

Assembly and whole genome alignment

To study the population structure of Asian *Xoo*, genomes from different locations were

downloaded from the Genbank as follows: 100 from India, 9 from Philippines, 1 from

South Korea and 1 from Japan [36,37]. We performed genome assembly from all the

strains sequenced in this study. For each strain, the pair-end clean reads were assembled

using Spades [67]. After a few evaluations, different sets of k-mer values were chosen

for Spades according to different read lengths. The final assemblies were obtained by

filtering out contigs with few reads supported or with lengths lower than 200 bp (Table

S1). Whole genome alignment was carried out on all the genomes obtained from public

database and assembled here by Parsnp using the genome of PXO99^A as reference [68].

Core SNPs were extracted from the core genome alignment by snp-sites

(https://github.com/sanger-pathogens/snp-sites). Recombination of the core genome

alignment was inferred by Gubbins. The final SNPs were obtained by filtering out the

prophage region and repeat sequences mentioned above, and the recombined regions

predicted here.

Phylogenetic analysis

The SNPs-based maximum likelihood (ML) phylogenies were built by RAxML with the generalized time-reversible model and a Gamma distribution to model site-specific rate variation [69]. Bootstrap support values were calculated from 500 replicates. *Xoo* population structure of China was defined with the hierBAPS module of the BAPS software, which delineates the population structure by nested clustering [70]. Three independent iterations with upper population sizes of 5, 10, and 15 were used to obtain optimal clustering of the population. Phylogenetic trees were annotated and visualized by iTOL (https://itol.embl.de/).

Comparative genomics of compete genomes

Ten isolates were selected for sequencing by long-read sequencing platform based on their positions on the phylogenetic tree and the races they represented. High quality total DNA of each isolate was sequenced by the PacBio RS II sequencing platform using P6-C4 chemistry to a mean fold coverage of 150. De novo assembly was conducted by HGAP3 [71].

For more comprehensive analysis of the genomic dynamics of *Xoo*, complete *Xoo* genomes available on Genbank were included. All the genomes were annotated by Prokka [72]. The TAL effectors of each genome were identified and classified by AnnoTALE [73]. Pan and core genome analysis was conducted by the tool Roary [74], based the gff format file constructed by Prokka. The inflation value of 2 was used for the MCL clustering.

Virulence evaluation of *Xoo* strains

Virulence of *Xoo* was asseses by inoculating 6 near-isogenic rice lines, each carrying

a specific R gene, IRBB2 (*Xa2*), IRBB3 (*Xa3*), IRBB4 (*Xa4*), IRBB5 (*xa5*), IRBB14

(*Xa14*) and IR24 (*Xa18*). IR24 was used as a susceptible check. Seeds of all lines in

this study were obtained from the China National Rice Research Institute (CNRRI).

Three weeks after sowing on a seedbed, seedlings were uprooted and transplanted into

the field. The management of the rice plants in the field was proceeded as usual.

Bacterial strain was cultivated on NA plates at 28 °C for 48 h. Then the bacterial

colonies were suspended in sterile ddH$_2$O with concentration adjusted to 3×10^8 cfu/ml

before inoculation. The leaf-clipping method was used to test the virulence of all the

Xoo strains [75]. Fifteen top fully expanded leaves per rice plant were inoculated with

each strain. The lesions length was measured after 21 days post inoculation. The ratio

of the lesion length compared to the whole leaf length was calculated as described

previously [76]. When the ratio of the rice line was lower than 1/4, the rice line was

classified as resistant (R); when the ratios were between 1/4 and 1, the rice line was

rated as susceptible (S). The lesion length 14 days following inoculation was also

measured and used to compare with the data obtained 7 days later.

Data access

Sequence data generated for this study have been submitted to the NCBI Sequence Read

Archive under BioProject accession PRJNA350904.

Acknowledgments

This work was supported by National Key Research and Development Program of

Conflict of Interest

The authors declare no conflict of interest.

Reference

1. Didelot X, Walker AS, Peto TE, Crook DW, Wilson DJ (2016) Within-host evolution of bacterial
pathogens. Nat Rev Microbiol 14: 150-162.
2. Baker S, Thomson N, Weill FX, Holt KE (2018) Genomic insights into the emergence and spread of
antimicrobial-resistant bacterial pathogens. Science 360: 733-738.
3. Feasey NA, Hadfield J, Keddy KH, Dallman TJ, Jacobs J, et al. (2017) Distinct *Salmonella Enteritidis*
lineages associated with enterocolitis in high-income settings and invasive disease in low-income settings.
Nat Genet 49: 1211-1217.
4. Okoro CK, Kingsley RA, Connor TR, Harris SR, Parry CM, et al. (2012) Intracontinental spread of
human invasive *Salmonella Typhimurium* pathovariants in sub-Saharan Africa. Nat Genet 44: 1215-1221.
5. Weill FX, Domman D, Njamkepo E, Tarr C, Rauzier J, et al. (2017) Genomic history of the seventh
pandemic of cholera in Africa. Science 358: 785-789.
6. Coll F, Phelan J, Hill-Cawthorne GA, Nair MB, Mallard K, et al. (2018) Genome-wide analysis of
multi- and extensively drug-resistant *Mycobacterium tuberculosis*. Nat Genet 50: 307-316.
7. Quainoo S, Coolen JPM, van Hijum S, Huynen MA, Melchers WJG, et al. (2017) Whole-Genome
Sequencing of Bacterial Pathogens: the Future of Nosocomial Outbreak Analysis. Clin Microbiol Rev
30: 1015-1063.
8. Gardy JL, Loman NJ (2018) Towards a genomics-informed, real-time, global pathogen surveillance
system. Nat Rev Genet 19: 9-20.
9. Zhong Z, Chen M, Lin L, Han Y, Bao J, et al. (2018) Population genomic analysis of the rice blast
fungus reveals specific events associated with expansion of three main clades. ISME J 12: 1867-1878.
10. Baltrus DA, McCann HC, Guttman DS (2017) Evolution, genomics and epidemiology of
Pseudomonas syringae: Challenges in Bacterial Molecular Plant Pathology. Mol Plant Pathol 18: 152-
168.
11. Brader G, Compant S, Vescio K, Mitter B, Trognitz F, et al. (2017) Ecology and Genomic Insights
into Plant-Pathogenic and Plant-Nonpathogenic Endophytes. Annu Rev Phytopathol 55: 61-83.
12. Hartmann FE, Croll D (2017) Distinct Trajectories of Massive Recent Gene Gains and Losses in
Populations of a Microbial Eukaryotic Pathogen. Mol Biol Evol 34: 2808-2822.
13. Hartmann FE, Sanchez-Vallet A, McDonald BA, Croll D (2017) A fungal wheat pathogen evolved

host specialization by extensive chromosomal rearrangements. ISME J 11: 1189-1204.

14. Mansfield J, Genin S, Magori S, Citovsky V, Sriariyanum M, et al. (2012) Top 10 plant pathogenic
bacteria in molecular plant pathology. Mol Plant Pathol 13: 614-629.

15. Mew T, Alvarez A, Leach J, Swings J (1993) Focus on bacterial blight of rice. Plant Dis 77: 5-13.

16. New T Current status and future prospects of research on bacterial blight of rice. Annu Rev
Phytopathol: 359-382.

17. Fang Z (1963) Rice Bacterial Blight: Jiangsu People Press.

18. Zhang Q (2009) Genetics and Improvement of Bacterial Blight Resistance of Hybrid Rice in China.
Rice Science 16: 83-92.

19. Huang N, Angeles ER, Domingo J, Magpantay G, Singh S, et al. (1999) Pyramiding of bacterial
blight resistance genes in rice: marker-assisted selection using RFLP and PCR. Theor Appl Genet 95:
313-321.

20. Kottapalli KR, Narasu ML, Jena KK (2010) Effective strategy for pyramiding three bacterial blight
resistance genes into fine grain rice cultivar, Samba Mahsuri, using sequence tagged site markers.
Biotechnol Lett 32: 989-996.

21. Pradhan SK, Nayak DK, Mohanty S, Behera L, Barik SR, et al. (2015) Pyramiding of three bacterial
blight resistance genes for broad-spectrum resistance in deepwater rice variety, Jalmagna. Rice (N Y) 8:
51.

22. Liu H, Yang W, Hu B, Liu F (2007) Virulence analysis and race classification of *Xanthomonas oryzae*
pv. *oryzae* in China. J Phytopathol 155: 129-134.

23. Chen XL, Yu L, Gao LL, Jiang T, Li QY, et al. (2012) Elevational Variation in Diversity of
Xanthomonas oryzae pv. *oryzae* in South-West China. J Phytopathol 160: 261-268.

24. Hua J, Zhang Y, Qian W, He CZ (2007) Avirulence gene and insertion element-based RFLP as well
as RAPD markers reveal high levels of genomic polymorphism in the rice pathogen *Xanthomonas oryzae*
pv. *oryzae*. Syst Appl Microbiol 30: 587-599.

25. Chen X, Wei S, Yan Q, Huang F, Ma Z, et al. (2019) Virulence and DNA fingerprinting analysis of
Xanthomonas oryzae pv. *oryzae* identify a new pathotype in Guangxi, South China. J Basic Microbiol
59: 1082-1091.

26. Li G, Song CF, Pang XM, Yang Y, Wang JS (2009) Analysis of Pathotypic and Genotypic Diversity
of *Xanthomonas oryzae* pv. *oryzae* in China. J Phytopathol 157: 208-218.

27. Jeung JU, Heu SG, Shin MS, Vera Cruz CM, Jena KK (2006) Dynamics of *Xanthomonas oryzae* pv.
oryzae Populations in Korea and Their Relationship to Known Bacterial Blight Resistance Genes.
Phytopathol 96: 867-875.

28. Adhikari TB, Mew TW, Leach JE (1999) Genotypic and Pathotypic Diversity in *Xanthomonas oryzae*
pv. *oryzae* in Nepal. Phytopathol 89: 687-694.

29. Mishra D, Vishnupriya MR, Anil MG, Konda K, Raj Y, et al. (2013) Pathotype and genetic diversity
amongst Indian isolates of *Xanthomonas oryzae* pv. *oryzae*. PLoS One 8: e81996.

30. Nelson RJ, Baraoidan MR, Cruz CM, Yap IV, Leach JE, et al. (1994) Relationship between Phylogeny
and Pathotype for the Bacterial Blight Pathogen of Rice. Appl Environ Microbiol 60: 3275-3283.

31. Xu Z, Sun Q, Liu F, Chen Z, Hu B, et al. (2004) Race monitoring of rice bacterial blight
(*Xanthomonas oryzae* pv.　*oryzae*) in China. Zhongguo Shuidao Kexue 18: 469-473.

32. Jiang N, Yan J, Liang Y, Shi Y, He Z, et al. (2020) Resistance Genes and their Interactions with
Bacterial Blight/Leaf Streak Pathogens (*Xanthomonas oryzae*) in Rice (*Oryza sativa* L.)—an Updated
Review. Rice 13: 1-12.

33. Quibod IL, Atieza-Grande G, Oreiro EG, Palmos D, Nguyen MH, et al. (2020) The Green Revolution
shaped the population structure of the rice pathogen *Xanthomonas oryzae* pv. *oryzae*. ISME J 14: 492-
505.

34. Huang X, Kurata N, Wei X, Wang ZX, Wang A, et al. (2012) A map of rice genome variation reveals
the origin of cultivated rice. Nature 490: 497-501.

35. Kovach MJ, Sweeney MT, McCouch SR (2007) New insights into the history of rice domestication.
Trends Genet 23: 578-587.

36. Quibod IL, Perez-Quintero A, Booher NJ, Dossa GS, Grande G, et al. (2016) Effector Diversification
Contributes to *Xanthomonas oryzae* pv. *oryzae* Phenotypic Adaptation in a Semi-Isolated Environment.
Sci Rep 6: 34137.

37. Midha S, Bansal K, Kumar S, Girija AM, Mishra D, et al. (2017) Population genomic insights into
variation and evolution of *Xanthomonas oryzae* pv. *oryzae*. Sci Rep 7: 40694.

38. Adhikari TB, Cruz C, Zhang Q, Nelson RJ, Skinner DZ, et al. (1995) Genetic Diversity of
Xanthomonas oryzae pv. *oryzae* in Asia. Appl Environ Microbiol 61: 966-971.

39. Triplett LR, Hamilton JP, Buell CR, Tisserat NA, Verdier V, et al. (2011) Genomic analysis of
Xanthomonas oryzae isolates from rice grown in the United States reveals substantial divergence from
known X. oryzae pathovars. Appl Environ Microbiol 77: 3930-3937.

40. Gonzalez C, Szurek B, Manceau C, Mathieu T, Sere Y, et al. (2007) Molecular and pathotypic
characterization of new *Xanthomonas oryzae* strains from West Africa. Mol Plant Microbe Interact 20:
534-546.

41. Stuecker MF, Tigchelaar M, Kantar MB (2018) Climate variability impacts on rice production in the
Philippines. Plos One 13: e0201426.

42. Lu ZJ, Song Q, Liu KB, Wu WB, Liu YX, et al. (2017) Rice cultivation changes and its relationships
with geographical factors in Heilongjiang Province, China. J Integr Agr 16: 2274-2282.

43. Wang Y, Xue Y, Li J (2005) Towards molecular breeding and improvement of rice in China. Trends
Plant Sci 10: 610-614.

44. Hu J, Xiao C, He YQ (2016) Recent progress on the genetics and molecular breeding of brown
planthopper resistance in rice. Rice (N Y) 9(1): 30.

45. Miedaner T, Korzun V (2012) Marker-Assisted Selection for Disease Resistance in Wheat and Barley
Breeding. Phytopathol 102: 560-566.

46. Mesterhazy A, Lemmens M, Reid LM (2012) Breeding for resistance to ear rots caused by *Fusarium*
spp. in maize - a review. Plant Breeding 131: 1-19.

47. Tiwari JK, Siddappa S, Singh BP, Kaushik SK, Chakrabarti SK, et al. (2013) Molecular markers for
late blight resistance breeding of potato: an update. Plant Breeding 132: 237-245.

48. McDonald BA, Linde C (2002) Pathogen population genetics, evolutionary potential, and durable
resistance. Annu Rev Phytopathol 40: 349-379.

49. Yang SQ, Liu SY, Zhao S, Yu YH, Li RB, et al. (2013) Molecular and pathogenic characterization of
new *Xanthomonas oryzae* pv. *oryzae* strains from the coastline region of Fangchenggang city in China.
World J Microbiol Biotechnol 29: 713-720.

50. Tu M, Lu BR, Zhu Y, Wang Y (2007) Abundant within-varietal genetic diversity in rice germplasm
from Yunnan Province of China revealed by SSR fingerprints. Biochem Genet 45: 789-801.

51. Xiong Z ZS, Wang Y, Ford-Lloyd BV, Tu M, Jin X, Wu Y, Yan H, Yang X, Liu P, Lu BR (2010)
Differentiation and distribution of *indica* and *japonica* rice varieties along the altitude gradients in
Yunnan Province of China as revealed by InDel molecular markers. Genet Resour Crop Evol 57: 891-

902.

52. Chen S, Lin XH, Xu CG, Zhang QF (2000) Improvement of bacterial blight resistance of 'Minghui 63', an elite restorer line of hybrid rice, by molecular marker-assisted selection. Crop Science 40: 239-244.

53. Leach JE, Rhoads ML, Vera Cruz CM, White FF, Mew TW, et al. (1992) Assessment of genetic diversity and population structure of *Xanthomonas oryzae* pv. *oryzae* with a repetitive DNA element. Appl Environ Microbiol 58: 2188-2195.

54. Carpenter SCD, Mishra P, Ghoshal C, Dash PK, Wang L, et al. (2018) A Strain of an Emerging Indian *Xanthomonas oryzae* pv. *oryzae* Pathotype Defeats the Rice Bacterial Blight Resistance Gene xa13 Without Inducing a Clade III SWEET Gene and Is Nearly Identical to a Recent Thai Isolate. Front Microbiol 9: 2703.

55. Huang S, Antony G, Li T, Liu B, Obasa K, et al. (2016) The broadly effective recessive resistance gene xa5 of rice is a virulence effector-dependent quantitative trait for bacterial blight. Plant J 86: 186-194.

56. Shen Y, Ronald P (2002) Molecular determinants of disease and resistance in interactions of *Xanthomonas oryzae* pv. *oryzae* and rice. Microbes Infect 4: 1361-1367.

57. White FF, Yang B (2009) Host and Pathogen Factors Controlling the Rice-*Xanthomonas oryzae* Interaction. Plant Physiology 150: 1677-1686.

58. Ji Z, Wang C, Zhao K (2018) Rice Routes of Countering *Xanthomonas oryzae*. Int J Mol Sci 19: 3008.

59. Boch J, Bonas U (2010) Xanthomonas AvrBs3 family-type III effectors: discovery and function. Annu Rev Phytopathol 48: 419-436.

60. Ferreira RM, de Oliveira AC, Moreira LM, Belasque J, Jr., Gourbeyre E, et al. (2015) A TALE of transposition: Tn3-like transposons play a major role in the spread of pathogenicity determinants of *Xanthomonas citri* and other xanthomonads. mBio 6: e02505-02514.

61. Raffaele S, Farrer RA, Cano LM, Studholme DJ, MacLean D, et al. (2010) Genome evolution following host jumps in the Irish potato famine pathogen lineage. Science 330: 1540-1543.

62. Jones JD, Dangl JL (2006) The plant immune system. Nature 444: 323-329.

63. Kelley DR, Schatz MC, Salzberg SL (2010) Quake: quality-aware detection and correction of sequencing errors. Genome Biol 11: R116.

64. Liu H, Xin B, Zheng J, Zhong H, Yu Y, et al. (2020) Build a Bioinformatics Analysis Platform and Apply it to Routine Analysis of Microbial Genomics and Comparative Genomics , 27 January 2020, PROTOCOL (Version 1) available at Protocol Exchange. [+https://doiorg/1021203/rs221224/v1+].

65. Salzberg SL, Sommer DD, Schatz MC, Phillippy AM, Rabinowicz PD, et al. (2008) Genome sequence and rapid evolution of the rice pathogen *Xanthomonas oryzae* pv. *oryzae* PXO99A. BMC Genomics 9: 204.

66. Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25: 1754-1760.

67. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, et al. (2012) SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19: 455-477.

68. Treangen TJ, Ondov BD, Koren S, Phillippy AM (2014) The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. Genome Biol 15: 524.

69. Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30: 1312-1313.

70. Cheng L, Connor TR, Siren J, Aanensen DM, Corander J (2013) Hierarchical and spatially explicit
clustering of DNA sequences with BAPS software. Mol Biol Evol 30: 1224-1228.
71. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, et al. (2013) Nonhybrid, finished microbial
genome assemblies from long-read SMRT sequencing data. Nat Methods 10: 563-569.
72. Seemann T (2014) Prokka: rapid prokaryotic genome annotation. Bioinformatics 30: 2068-2069.
73. Grau J, Reschke M, Erkes A, Streubel J, Morgan RD, et al. (2016) AnnoTALE: bioinformatics tools
for identification, annotation, and nomenclature of TALEs from Xanthomonas genomic sequences. Sci
Rep 6: 21077.
74. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, et al. (2015) Roary: rapid large-scale
prokaryote pan genome analysis. Bioinformatics 31: 3691-3693.
75. Kauffman H, Reddy A, Hsieh S, Merca S (1973) An improved technique for evaluating resistance of
rice varieties to *Xanthomonas oryzae*. Plant Dis Rep 57: 737-741.
76. Fang C, Xu Z, Guo C, Yin S, Wu S, et al. (1990) Studies on pathotypes of *Xanthomonas campestris*
pv. *oryzae* in China. Acta Phytopathol Sin 20: 81-88.
664
665

Figure 1 Population structure of *X. oryzae* pv. *oryzae*. The maximum likelihood tree was inferred

by RAxML with the generalized time-reversible model and a Gamma distribution to model site-

specific rate variation based on the core SNPs of all the *Xoo* genome sequences. The six lineages

were decorated with different colors. From inner to outer, the first circle refers to major rice planting

areas, the second one describes isolated times of these strains and the third one represents the rice

subspecies which the *Xoo* strain isolates. The outmost 6 squares represent the virulence analysis of

these *Xoo* against 6 different rice near-isogenic lines with one known resistance gene in each line,

with solid square for resistance and hollow one for sensitivity. Bootstrap support values were

calculated from 500 replicates, and only values of > 70% were labelled.

Figure 2 Distribution of the 6 major Chinese *X. oryzae* pv. *oryzae* lineages. Pathotypes of *Xoo*

were determined on the following 13 near-isogenic lines, each carrying a specific R gene, IRBB2

(*Xa2*), IRBB3 (*Xa3*), IRBB4 (*Xa4*), IRBB5 (*xa5*), IRBB14 (*Xa14*). IR24 was used as a susceptible

check. (a) The map of the 3 major rice producing areas in China. (b) Distribution of the 6 *Xoo*

lineages among the 3 rice planting areas. (c) Dynamics of the 6 *Xoo* lineages during the past 30

years.

Figure 3 Phylogeography of CX-5 and CX-6. Maximum likelihood phylogeny of Lineage CX-5

(a) and CX-6 (b). Different sub-lineages were displayed by different branch colors. Bootstrap

support values were calculated from 500 replicates. The recent outbreaks are represented by

different backgrounds. Isolated information of *Xoo* including location, time and rice subspecies

were shown in three colored strips. (c) Mean Pairwise SNP Distances (MPSDs) of *Xoo* from the

outbreak, and between one in an outbreak and the other one out of the outbreak but in the same sub-

lineage were displayed by two different boxes beside the tree.

Figure 4 Virulence analysis and race classification of *X. oryzae* pv. *oryzae* China. Pathogenicity

profile of each race corresponding to was shown in Table S1. (a) Proportions of the 9 *Xoo* races. (b)

Distribution of the 9 races among the 3 major rice planting areas in China. (c) Dynamics of the 9

Xoo races during the past 40 years. (d) Distribution of the 9 races in Lineage CX-6 and others.

Figure 4 Core and pan genome of *Xoo*. The results showed here is from the comparative analysis

of the 22 *Xoo* complete genomes, using OS198 belonging to Lineage CX-6 as reference. From the

inner to the outmost, (1) Circle 1, the black break line, SNPs per 10kbp; (2) Circle 2, the colorized

lines, different families of transposase genes (magenta for IS3, blue for IS5, yellow for IS30, green

for IS630, red for IS701, orange for IS1595, black for others); (3) Circle 3, the red break line,

numbers of genome having that gene family (ranging 1 to 22); (4) Circle 4, the dark blue lines,

positions of core gene; (5) Circle 5, the colorful lines, presences of type III effector genes, with

green for TALEs in OS198, red for TALEs in other Xoo genomes, blue for other T3Es; (6) Circle 6,

scales according to genome of OS198.

Table 1 Resistance of different rice lines to *Xoo* strains isolated at different times.

Shown is the percentage of *Xoo* strains isolated at different periods that are unable to

cause disease.

Time (# of strains)	IRBB2 (*Xa2*)	IRBB3 (*Xa3*)	IRBB4 (*Xa4*)	IRBB5 (*xa5*)	IRBB14 (*Xa14*)	IR24 (*Xa18**)
<1990 (35)	37	0	97	94	25	20
<2000 (69)	25	58	64	100	28	12
2003/2004 (97)	49	35	34	100	56	7
2014 (42)	36	5	79	90	33	5

715   *: Recurrent parent *Xa18*

718 **Table 2 Resistance of different rice lines to *Xoo* strains of different lineages.** Shown

719 is the percentage of *Xoo* strains of different lineages that are unable to cause disease.

Lineage (# of strains)	IRBB2 (*Xa2*)	IRBB3 (*Xa3*)	IRBB4 (*Xa4*)	IRBB5 (*xa5*)	IRBB14 (*Xa14*)	IR24 (*Xa18**)
CX-1 (19)	89	79	42	100	95	21
CX-2 (15)	73	93	80	100	100	7
CX-3 (20)	100	20	70	100	99	10
CX-5 (53)	11	32	47	96	17	11
CX-6 (123)	29	25	62	98	28	6

*: Recurrent parent *Xa18*. CX-4 was not included because of few isolates it contained.

Supplementary Information

Figure S1 Population structure of Asian *Xanthomonas oryzae* pv. *oryzae*. For phylogenetic

analysis, parsnp was used to call SNPs, and RAxML was used to build the ML tree. Background

colors of nodes represent different lineages. The outmost circle refers to different countries where

Xoo were isolated.

Figure S2 Plot of Pan and core genes from the 22 complete *Xoo* genomes.

Figure S3 Function analysis of pan genes. (a) COG function of different types of gene families.
Core, genes shared by all the 22 genomes; distributed, gene shared by 2 ~ 21 genomes; unique,
genes unique to each genome. (b) Classification of transposase genes. COG categories are energy
production and conversion (C), amino acid metabolism and transport (E), carbohydrate metabolism
and transport (G), coenzyme metabolism (H), lipid metabolism (I), translation (J), transcription (K),
cell wall/membrane/envelop biogenesis (M), cell motility (N), posttranslational modification,
protein turnover, and chaperone functions (O), inorganic ion transport and metabolism (P), general
functional prediction only (R), function unknown (S), signal transduction (T), Mobilome: prophages,
transposons (X).

Table S1 *Xoo* isolates used in this study.

Table S2 Resistance of different rice lines to *Xoo* strains of different sub-lineages of CX-5 and CX-6. Shown is the percentage of *Xoo* strains of different lineages that are unable to cause disease.

Lineage (# of strains)	IRBB2 (*Xa2*)	IRBB3 (*Xa3*)	IRBB4 (*Xa4*)	IRBB5 (*xa5*)	IRBB14 (*Xa14*)	IR24 (*Xa18**)
CX-5.1 (5)	0	20	20	100	0	0
CX-5.3 (12)	25	25	67	92	17	17
CX-5.4 (8)	25	50	75	100	28	25
CX-5.5 (18)	0	28	28	100	28	0
CX-6.1(10)	30	30	90	100	50	20
CX-6.2 (47)	13	38	79	98	23	4
CX-6.3 (18)	10	17	68	100	11	0
CX-6.4 (18)	0	0	61	94	0	0
CX-6.5(31)	90	32	29	97	74	10

*: Recurrent parent *Xa18*. CX-5.2 was not included because of few isolates it contained.

Table S3 Transposase content of *Xoo* with complete genomes.

Table S4 Distribution of TAL among different *Xoo* with complete genomes

Figure S1 Population structure of Asian *Xanthomonads oryzae* pv. *oryzae*. For phylogenetic
analysis, parsnp was used to call SNPs, and RAxML was used to build the ML tree. Background
colors of nodes represent different lineages. The outmost circle refers to different countries where
Xoo were isolated.

Table S2 Resistance of different rice lines to *Xoo* strains of different sub-lineages of CX-5 and CX-6. Shown is the percentage of *Xoo* strains of different lineages that are unable to cause disease.

Lineage (# of strains)	IRBB2 (*Xa2*)	IRBB3 (*Xa3*)	IRBB4 (*Xa4*)	IRBB5 (*xa5*)	IRBB14 (*Xa14*)	IR24 (*Xa18**)
CX-5.1 (5)	0	20	20	100	0	0
CX-5.3 (12)	25	25	67	92	17	17
CX-5.4 (8)	25	50	75	100	28	25
CX-5.5 (18)	0	28	28	100	28	0
CX-6.1(10)	30	30	90	100	50	20
CX-6.2 (47)	13	38	79	98	23	4
CX-6.3 (18)	10	17	68	100	11	0
CX-6.4 (18)	0	0	61	94	0	0
CX-6.5(31)	90	32	29	97	74	10

*: Recurrent parent *Xa18*. CX-5.2 was not included because of few isolates it contained.