In silico identification of a molecular circadian system with novel features in the crustacean model organism *Parhyale hawaiensis*

BENJAMIN JAMES HUNT [1], EAMONN MALLON [1] & EZIO ROSATO [1*]

[1] Department of Genetics and Genome Biology, University of Leicester, Leicester, United Kingdom

* Correspondence: Dr. Ezio Rosato, er6@le.ac.uk

Keywords: crustacean, amphipod, parhyale, talitrus, hyalella, circadian, clock, genetics
Running title: "Molecular clock of *Parhyale hawaiensis*"

Word count: 4,265

Abstract

The amphipod *Parhyale hawaiensis* is a model organism of growing importance in the fields of evolutionary development and regeneration. A small, hardy marine crustacean that breeds year-round with a short generation time, it has simple lab culture requirements and an extensive molecular toolkit including the ability to generate targeted genetic mutant lines. Here we identify canonical core and regulatory clock genes using genomic and transcriptomic resources as a first step in establishing this species as a model in the field of chronobiology. The molecular clock of *P. hawaiensis* lacks orthologs of the canonical circadian genes *cryptochrome 1* and *timeless*, in common with the mammalian system but in contrast to many arthropods including *Drosophila melanogaster*. Furthermore the predicted CLOCK peptide is atypical and CRY2 shows an extended 5' region of unknown function. These results appear to be shared by two other amphipod species.

1 Introduction

The subphylum Crustacea has a rich history of chronobiological research. One of the earliest recorded examples of a persistent circadian phenomenon in animals was the colour changes seen in the prawn *Hippolyte varians* (Gamble and Keeble, 1900), while investigations on the circadian and circatidal rhythms of the fiddler crab *Uca pugnax* helped lay the basis for modern chronobiology (Palmer, 1991), with published works on the persistence of chromatophore and locomotor activity rhythms in constant conditions; temperature compensation; persistance of rhythms in isolated tissue; and the phase-shifting effect of light pulses, this latter observation coming a decade before the phase response curve (PRC) was incorporated into circadian concepts (De Coursey, 1960). A variety of endogenously controlled rhythmic phenomena have been documented in wide range of crustaceans, not only circadian rhythms but also tidal, lunar and semi-lunar (Strauss and Dircksen, 2010), under the control of *zeitgebers* such as light, hydrostatic pressure, immersion, wave motion and salinity. Where the clade has historically proved lacking, however, is in providing an organism with a molecular toolkit comparable to that of the fruit fly *Drosophila melanogaster*, in which transgenics, reporter genes and binary expression systems have been used to characterise, locate and dissect the endogenous biological clock at a neurological level (Helfrich-Förster, 2005).

At the centre of the *Drosophila* clock are the bHLH-PAS proteins CLOCK (Allada *et al.*, 1998; Gekakis *et al.*, 1998, CLK) and CYCLE (Rutila *et al.*, 1998, CYC), which form a heterodimer that drives transcription of clock-controlled genes including *period* (Konopka and Benzer, 1971, per) and *timeless* (Sehgal *et al.*, 1994, tim), whose protein products dimerise and interact with CLK:CYC to inhibit its transcription activity and thus their own expression. TIM is degraded by the light-activated CRYPTOCHROME (Emery *et al.*, 1998, CRY), exposing PER to degradation in turn and ending CLK:CYC repression. The transcription-translation feedback loop enacted by these core clock genes is modified to achieve a circadian periodicity of 24 hours through the contributions of numerous regulatory genes such as *doubletime, vrille, shaggy* and *clockwork orange* . In mammals the clock is similarly well understood thanks to work on *Mus musculus* (Reppert and Weaver, 2001), showing that the molecular system is remarkably conserved but not identical. In mammals the CLK:BMAL1 heterodimer (where BMAL1 is an ortholog of CYC) is inhibited by multiple PER and CRY proteins, the latter notably different to the *Drosophila* CRY in lacking photosensitivity. A number of species (Zhu *et al.*, 2005) have been found to have both a *Drosophila*-like CRY (often termed CRY1) and a mammalian-like CRY (CRY2).

In crustaceans, the first core clock gene to be identified was *clock* in the prawn *Macrobrachium rosenbergii* (Yang *et al.*, 2006). The falling cost of next generation sequencing has subsequently brought a wave of such data in crustacean species including the water flea *Daphnia pulex* (Tilden *et al.*, 2011), the copepods *Calanus finmarchicus* (Christie *et al.*, 2013) and *Tigriopus californicus* (Nesbit and Christie, 2014), the lobster *Nephrops norvegicus* (Sbragaglia *et al.*, 2015), the amphipod *Talitrus saltator* (O'Grady *et al.*, 2016) and the Antarctic krill *Euphausia superba* (Hunt *et al.*, 2017), in some cases including identification of rhythmic transcription. The clock of *E. superba* has been further characterised through analysis of the interactions of its protein components (Biscontin *et al.*, 2017), as has that of the isopod *Eurydice pulchra* (Zhang *et al.*, 2013).

Nevertheless much work remains to be done in determining how these genes generate rhythms, and in what tissues and cells they are active and when, and with which genes they interact. More fine-grained investigation into the crustacean clock requires a model organism of similar power to *D. melanogaster*, and the establishment of such a model also promises to significantly advance our understanding of the animal clock through comparative analysis.

The amphipod *Parhyale hawaiensis* was first isolated for lab culture in 1997 from the filtration system of the John G. Shedd Aquarium, "preselected [for] minimal care", in the words of the collectors (Browne *et al.*, 2005). A circumtropical marine detritivore found in shallow water mangrove zones, it is a robust, fecund organism that is cheap to maintain and breeds year-round with a quick generation time, making it well-suited to lab culture. Thanks to a reproductive system that permits the collection and manipulation of embryos from a single cell stage this species has become established as a model organism of some significance in the field of development. It has further proven itself to be amenable to molecular analysis and manipulation, including *in situ* hybridisation, RNAi-mediated knockdown, transposon and integrase-based genetic transformation and CRISPR/Cas9 mutagenesis, and recently the 3.6 Gb genome was published (Kao *et al.*, 2016).

Here we describe the creation of a head transcriptome to identify core and regulatory clock genes

56 and use the genome to gain independent verification of our findings, which include the apparent loss of a
57 number of major clock genes and the identification of a peptide feature that may be unique to amphipods.

Materials and Methods

Animal husbandry

Animals were maintained in polypropylene containers with lids 3/4 on, with a layer of crushed coral approximately 1 cm deep and filled with 1 L artificial seawater prepared using 30g Tetra Marine Sea Salt per litre of distilled tap water to give a specific gravity of 1.022. Cultures were maintained on a 12:12 light:dark cycle with lights on at 9am, illumination coming from fluorescent room lighting at 300 lux, at 20°C. Animals were fed twice weekly, once with shredded carrot and once with a pinch of standard fish flakes.

Head transcriptome

At eight time-points across 24 hours beginning at 9am (lights on, *zeitgeber* time (ZT) 0) and every three hours thereafter, a total of 20 adult males were collected per time-point and snap-frozen on a metal plate atop dry ice, whereupon the heads were dissected and stored at -80°C until use. Total RNA was extracted from five heads per time-point using a micropestle to homogenise the tissue in TRIzol reagent (Invitrogen), cleaned using the PureLink Micro Kit (ThermoFisher Scientific) with on-column DNase treatment to remove genomic DNA contamination and eluted with RNase free water. RNA extractions were quality assessed using Nanodrop and Agilent 2100 Bioanalyzer. Samples were required to meet a minimum quality of > 50 ng/(microlitres symbol) concentration; 260/280 ratio 1.8 − 2.2; 260/230 ratio 2.0 - 2.4 for sequencing use.

One sample per time-point was delivered to Glasgow Polyomics (University of Glasgow) and subject to polyA mRNA enrichment with the TruSeq Stranded mRNA Library Prep Kit (Illumina). The resultant libraries were sequenced using the Illumina NextSeq 500 platform to generate 75 bp paired-end strand-specific reads, which were subject to gentle low quality read removal and Q10 adapter trimming by Glasgow Polyomics. After download of this dataset FastQC 0.11.2 (Andrews, 2010) was used to assess read quality and Cutadapt 1.9.1 (Martin, 2011) used to remove remaining traces of the TruSeq Index and Universal adapters using the sequence "GATCGGAAGAGC" common to both adapter types as criteria. Reads were then filtered using Trimmomatic 0.32 (Bolger *et al.*, 2014) to remove reads with an average quality of less than 20 and/or shorter than 72 bases.

A multi-k-mer, multi-assembler approach was adopted for assembly. Trans-ABySS 1.5.5 (Robertson *et al.*, 2010), Bridger 2014-12-01 (Chang *et al.*, 2015) and Trinity r20140717 (Haas *et al.*, 2013) were used to generate individual assemblies using k-mers ranging from 19 to 71 with a minimum contig length of 200 bp. These were subsequently merged and duplicates and containments were removed at 99% identity using CD-HIT-EST 4.6.6 (Li and Godzik, 2006). TransRate (Smith-Unna *et al.*, 2016) was used to identify well-assembled contigs based on read-mapping evidence. TransDecoder 3.0.0 (Haas *et al.*, 2013) was used to identify predicted peptides from the well-assembled contigs (single_best_orf, -m 90), and CD-HIT used at 90% peptide identity to produce a final list of well-assembled, non-redundant coding contigs that were extracted from the original merged transcriptome assembly and used for all downstream purposes.

Identification of clock genes

The head transcriptome was searched for core and regulatory circadian genes using the tblastn function of BLAST+ with an E-value cutoff of 1.0e-5 (Table 1). The protein sequences inferred from the genome (Kao *et al.*, 2016, see https://doi.org/10.6084/m9.figshare.3498104.v4) were searched in the same manner using blastp. Genes that returned little or no evidence through this approach were searched for in the genome using tblastn and, where evidence was subsequently obtained, the scaffold and coordinates of the alignments were noted. The head transcriptome reads were aligned to the genome using HISAT2 2.1.0 (Kim *et al.*, 2015) and visualised with IGV 2.3.82 (Robinson *et al.*, 2011), and Stringtie 1.3.3b (Pertea *et al.*, 2015) was used to assemble transcripts of interest from these read alignments. Gene sequences identified through BLAST+ searches were extended using these alignments and transcripts where appropriate. Sequence data from two other amphipods, *Hyalella azteca* (GCA_000764305.2) and *Talitrus saltator* (GDUJ00000000) was further used to extend partial transcripts and lend support to atypical results. RNA-seq reads for *H. azteca* (PRJNA312414) were downloaded, aligned and assembled as above. *P. hawaiensis* sequences obtained from from BLAST+ searches and contigs identified as circadian genes in *T. saltator* (O'Grady *et al.*, 2016) were used to search the genomes/transcriptomes of all three

species, with results expanded using read mapping evidence and then these extended sequences used to search all resources again until no further extension was possible. Sequences returned by these processes were translated to peptides and subject to validation using the NCBI BLAST web interface (blastp, Database 'nr', Entrez Query 'all[filter] NOT predicted[title] NOT hypothetical[title] NOT putative[title]') to ascertain their closest match in other species.

Peptide domain analysis was conducted using SMART (Letunic *et al.*, 2015). Peptide sequences were visualised using DOG 2.0 (Ren, et al, 2009). Phylogenetic trees were created using MEGA 7.0.26; peptide sequences were aligned using MUSCLE at default settings, and neighbour-joining trees generated using 1,000 bootstrap replications and complete deletion.

Gene expression analysis

The Trinity utilities 'align_and_estimate_abundance.pl' and 'abundance_estimates_to_matrix.pl' (–est_method RSEM, –aln_method bowtie2) were used to generate a TMM-normalised (Robinson and Oshlack, 2010) expression matrix by aligning the quality-trimmed reads to the head transcriptome. Values for all circadian-related contigs were extracted, along with the values for a number of putative housekeeping genes identified via BLAST search (Table S1). For each contig these were normalised to the highest value (highest = 1 after normalisation), and then the mean normalised value calculated. Finally for each timepoint the departure from the mean normalised value was calculated and plotted using ggplot2 (Wickham, 2016) function in R v3.4.0 (R Core Team, 2018).

Results

Transcriptome assembly and mining

After quality control the RNA-seq library for each head timepoint sample was approximately 22 million read pairs, with a total of 178,028,148 used in a final assembly of 86,686 contigs with an N50 of 2,797 and contig length ranging from 267 – 35,494. This Transcriptome Shotgun Assembly project has been deposited at GenBank under the accession GFVL00000000.

Evidence was found for the existence of the core clock genes *bmal1*, *period* and the mammalian-like *cryptochrome 2* in this assembly. Also identified were 16 regulatory clock genes (Table 1, Table 2). The expression across 24 hours of all clock-related genes present in the head transcriptome is shown in Figure 1, in which a pattern of peak expression at ZT3-6 and ZT18 appears to be common in clock genes, in contrast to the selection of housekeeping genes.

BMAL1/CYCLE

P. hawaiensis bmal1 (Ph-bmal1), found in complete form in the genome-inferred peptides (phaw_30_tra_m.024548) and in fragmentary form in the head transcriptome assembly, shows the typical complement of domains when translated (Figure 2A), with a bHLH domain, two PAS domains, a PAC domain immediately following the second PAS, and a BMAL1 C-terminal Region (BCTR), a transactivation domain characteristic of this gene (except in *D. melanogaster* CYCLE). The peptide returned Bmal1a in *Pacifastacus leniusculus* (accession AFV39705; $1E^{-161}$) from a BLAST confirmation query. *Ph-bmal1* showed very low levels of expression in the head transcriptome (Table S2).

CLOCK

No evidence was found of a *P. hawaiensis clock* gene (*Ph-clock/Ph-clk*) through BLAST mining of the head transcriptome or genome-inferred peptides. Searching the genome using *Eurydice pulchra* CLK 5 (AGV28720) identified orthologous sequences on scaffold LQNS02278184.1 (supplementary data). When translated and combined, these sequences returned *clock* in the American lobster *Homarus americanus* (AWC08577; $3E^{-69}$), while SMART domain analysis identified bHLH, PAS and PAC domains. A small number of reads mapped to these fragments, indicating very low levels of expression in the head.

A putative CLK protein has been reported for the amphipod *T. saltator* (O'Grady *et al.*, 2016). This 1,907 amino acid internal fragment peptide lacks the bHLH domain and the first of the two PAS domains typical to CLK proteins reported in other species, showing only the second PAS domain with an associated PAC near the beginning of the sequence, with the remainder showing no homology to any entry in the NCBI nr database. Searching the *P. hawaiensis* head transcriptome and genome-inferred peptides using this sequence again returned no results, however searching the genome returned further alignments on scaffold LQNS02278184.1, just downstream of the bHLH/PAS/PAC coding sequences described above. Highly similar results are seen in the *H. azteca* genome (supplementary data), with BLAST searches identifying sequences encoding bHLH and PAS domains and sequences downstream of these with strong homology to Tal-CLK. Read mapping evidence further extended this to a final sequence encoding a 2,241 amino acid (aa) peptide (Ha-CLK) with 36% identity to Tal-CLK. Using this to search the *P. hawaiensis* genome extended Ph-CLK to a final sequence encoding a 2,065 aa peptide. Schematics of the predicted peptides for both species are shown in Figure 2B.

Searching the *T. saltator* transcriptome with Ph-CLK and Ha-CLK identified two additional contigs of interest beyond Tal-CLK (Figure 2B): GDUJ01041853 encodes a truncated peptide with a bHLH domain that returns *clock* in *Euphausia superba* (ANW48378; $6E^{-24}$), while GDUJ01061398 shows homology (BLAST alignment 41% identity, $1E^{-40}$) to the downstream sequence identified in Ha-CLK using read mapping evidence. If these two contigs are part of the full *T. saltator clock* gene this would result in a peptide at least 2,455 aa in length.

CRYPTOCHROME 2

P. hawaiensis cryptochrome 2 (*Ph-cryptochrome 2/Ph-cry2*) was identified in complete form in the head transcriptome, returning cryptochrome 2 in *T. saltator* (AFV96168; e-value 0.0) in a BLAST confirmation query. SMART analysis of the translated 931 aa peptide identified a DNA photolyase and FAD

binding domain (Figure 2C), while an extended N-terminus of 357 amino acids had no identifiable functional domain or motif. A neighbour-joining phylogenetic tree confirmed the nature of the peptide as a mammalian-like CRY2 rather than a *Drosophila*-like CRY1 (Figure 3A). This coding sequence was supported by genome search and read mapping evidence: Stringtie assembled an 11-exon transcript (supplementary data) located on scaffold LQNS02278089.1 that shows 99.5% identity to the *de novo* assembled contig, with the first exon incorporating both the N-terminus and a partial DNA photolyase domain in one contiguous sequence, indicating that the *de novo* assembled contig is unlikely to be a chimeric construct. Further to this the genome-inferred peptides, which were generated with reference to an independently assembled *de novo* transcriptome from embryonic and limb tissue (Kao *et al.*, 2016), include a contig (phaw_30_tra_m.017772) covering 189 amino acids of the N-terminus and the entire DNA photolyase domain, and another (phaw_30_tra_m.017773) covering the first 160 amino acids of the N-terminus. Of the core clock genes identified *Ph-cry2* showed the highest levels of expression (Table S2).

Similar results can again be found in the other two amphipod species. In the paper detailing the molecular clock of *T. saltator* (O'Grady *et al.*, 2016) two contigs were identified as representing *cry2*. An assumed complete peptide of 565 amino acids was reported, but one of the *de novo* assembled transcriptome contigs (accession GDUJ01076706.1, reported as comp100937_c0_seq1) includes an N-terminus of 362 amino acids that is not considered part of the CRY2 peptide by the authors. However this sequence shows 55.9% identity and 71.3% similarity to the *P. hawaiensis* N-terminus. In *H. azteca* a predicted *cryptochrome* has been reported (accession XM_018166003), and while somewhat extended at the 5' end it again does not include an N-terminus comparable to that of *P. hawaiensis*. However a BLAST search of the genome using the Ph-CRY2 peptide returned alignments on scaffold KV721600.1 suggesting the existence of such a feature upstream of the reported sequence. Read alignment and assembly generated an 11-exon transcript (supplementary data), the second exon coding for the N-terminus and partial DNA photolyase domain and terminating at the same point as the first exon of the Stringtie *Ph-cry2* transcript. When translated, this transcript encodes a peptide 932 amino acids long (Ha-CRY2) incorporating the extended N-terminus with the previously predicted *H. azteca* CRY.

No evidence was found in any *P. hawaiensis, T. saltator* or *H. azteca* resource for a *Drosophila*-like *cryptochrome*.

PERIOD

A small 3' fragment of *P. hawaiensis period* (*Ph-period/Ph-per*) was found in the head transcriptome, the peptide of which returned *period* in *Macrobrachium nipponense* from a BLAST confirmation query (accession ANN13870; e-value 4E^{-07}). Searching the genome using this sequence and the predicted PER protein of *Hyalella azteca* (accession XP_018012603) returned multiple alignments on scaffold LQNS02276498.1 (supplementary data). A small number of head transcriptome reads mapped to 14 of these these regions. Translated and combined to form a fragmented putative peptide sequence 1,257 amino acids in length, SMART domain analysis revealed two PAS domains, a PAC domain and a PERIOD C domain (Figure 2D). This construct returned *period isoform 1* in *Homarus americanus* (accession AWC08578; 3E^{-99}) from a BLAST confirmation query. *Ph-per* showed very low levels of expression in the head transcriptome (Table S2).

TIMELESS

A mammalian-like TIMELESS was identified in the genome-inferred peptides resource. This is an ortholog of the *D. melanogaster timeless* paralog *timeout* (Figure 3B), a gene crucial to early development that is seemingly ubiquitous in animals (Sandrelli *et al.*, 2008) and may have only peripheral involvement with the endogenous clock. No true TIMELESS could be identified in any *P. hawaiensis, H. azteca* or *T. saltator* resource.

Regulatory and clock-controlled genes

Putative orthologs were identified for the regulatory genes *casein kinase II α* and *β*, *circadian trip, clockwork orange, doubletime, microtubule star, nejire, nemo, par domain protein ϵ, protein phosphatase I, shaggy, supernumerary limbs, twins, vrille* and *widerborst*, and for the clock-controlled genes *pigment*

dispersing hormone (*pdh*) and *pdh receptor*. *Ph-pdh* was found only through the use of the putative
Hyalella azteca preprohormone Hyaaz-prepro-PDH (Christie, 2014) as a query term, and while a BLAST
confirmation query returned only one unrelated result (2) Ph-PDH shares 47.6% identity and 62.1%
similarity with Hyaaz-prepro-PDH. A signal peptide was detected between residues 36 and 37 and through
alignment with Hyaaz-prepro-PDH a 23 amino acid mature peptide was identified with 91.3% identity
and 100% similarity.

Discussion

In silico mining resulted in the identification and characterisation of four putative core clock proteins in *Parhyale hawaiensis*; Ph-BMAL1, Ph-CLK, Ph-PER and Ph-CRY2. Despite searching multiple resources, including the recently published genome, we were unable to identify an ortholog of TIM or a *Drosophila*-like CRY. This suggests that *P. hawaiensis* has evolved what might be termed a mammalian-style molecular clock - a system lacking a TIM or CRY1, as originally reported in *M. musculus* (Reppert and Weaver, 2000). This system is found in vertebrates, hymenopterans and now in crustaceans. The gene reported as *T. saltator timeless* (O'Grady et al., 2016) appears to be a *timeout* (Figure 3B) and no sequence encoding a CRY1 or true TIM could be found in the various transcriptomic and genomic resources available for *T. saltator* and *H. azteca*, indicating that this system may be common in amphipods. This contrasts with other crustaceans studied so far – *Daphnia pulex* possesses the full complement of core clock proteins including a CRY1 and CRY2 (Tilden *et al.*, 2011), as does *Calanus finmarchicus* (Christie *et al.*, 2013), *Trigriopus californicus* (Nesbit and Christie, 2014), *Limulus polyphemus* (Chesmore *et al.*, 2016) and *Euphausia superba* (Hunt *et al.*, 2017). *Eurydice pulchra* lacks only a CRY1 (Zhang *et al.*, 2013), as does *Nephrops norvegicus* (Sbragaglia *et al.*, 2015).

This molecular system is not the only interesting aspect of the *P. hawaiensis* circadian system. We were unable to identify a complete coding sequence for *clock*, relying on alignments and read mapping to generate an incomplete construct that is notably different from the orthologs of other species. The putative peptide products appear to be extremely long (*E. superba* CLK at 1,344 aa is one of the longest complete versions of this gene so far documented, yet is easily surpassed by the partial Tal-CLK at 1,907 aa), lack a second PAS domain and an extensive polyglutamine region that is characteristic of the gene (Saleem *et al.*, 2001), with only very small glutamine-rich regions identified. In *H. azteca* and *T. saltator* there is evidence of fragmentation in this gene, with bHLH-encoding sequences present in separate scaffolds/contigs. This fragmentation was also present in the first draft of the *P. hawaiensis* genome but was resolved with the second draft, and it is likely that the fragmentation is *in silico* rather than *in vivo* for all three species. The default assumption should be that Ph-CLK forms a heterodimer with Ph-BMAL1 to drive clock gene expression as is usual in animals, but the atypical nature of the amphipod CLK peptides encourages speculation about alternative partners for Ph-BMAL1 - in mice BMAL1 is a promiscuous binding protein (Hogenesch *et al.*, 1998) and other bHLH-PAS proteins have substituted for CLK in other species, for example methoprene-tolerent (MET) in the mosquito *Aedes aegypti* (Shin *et al.*, 2012), an ortholog of which can be found in the head transcriptome.

In this context we consider the cryptochrome Ph-CRY2, which shows an extraordinary extension at the N-terminus (and various lines of evidence suggest the same is true in *H. azteca* and *T. saltator*). Whether this serves a function, and what that might be, is again speculative at this point, but an unusual transcriptional heterodimer may require an unusual repressor. Perhaps then Ph-CRY2 serves in the repressive limb of the *P. hawaiensis* circadian feedback loop as does CRY2 in many other species, and it has evolved this N-terminus as a means to interact with, for example, a MET:BMAL1 heterodimer, or an atypical CLK:BMAL1.

A candidate for Ph-PDH proved difficult to identify due to its divergent sequence including at the conserved NSELINS motif. *T. saltator* is reported to possess two PDH peptides, the first (Tal-PDH I) unusual with regards to its long mature peptide form but showing the NSELINS motif and the second (Tal-PDH II) identified using relaxed search criteria. This latter PDH sequence and that of *H. azteca* - itself described as atypical (Christie, 2014) - were the only search terms from dozens used that returned a candidate contig. In insects the homologue of PDH, called Pigment dispersing factor (PDF) is a neuropeptide and neurohormone that is crucial for the synchronisation among the different circadian neurons (Renn *et al.*, 1999). In crustaceans PDH exists in many isoforms and the neuroanatomy of PDH-expressing cells has been described in numerous species, with evidence of PDH involvement in circadian rhythmicity in several of them (Strauss and Dircksen, 2010). The novel nature of the putative Ph-PDH requires further work to confirm its identity and to ascertain whether the sequence divergence has any impact on its functioning in the circadian pathways.

Given a lack of timepoint replicates in our head transcriptome data, and the very low levels of expression seen in a number of clock-related genes (Table S2), we have restricted ourselves to a descriptive portrayal of core and regulatory clock gene expression, wherein a bimodal pattern of rhythmic expression is apparent among clock genes (Figure 1) and which may be of interest to researchers studying circa-tidal rhythms. This transcriptomic data also indicates very low expression levels for the core clock genes

in head tissue, low enough that only short fragments were assembled of *Ph-bmal1* and *Ph-period*, and nothing of *Ph-clock*. This result is consistent with the isopod *E. pulchra* in which it has been determined that Ep-PERIOD is present in only 10 cells in the brain (Zhang *et al.*, 2013).

This *in silico* analysis of these amphipod circadian systems has generated plenty of avenues for future enquiry. These sequences need to be confirmed through further sequencing, and their expression patterns more stringently characterised, both temporally and in terms of location using more extensive RNA-seq and/or RT-PCR. If it is found that Ph-CLK has been accurately described here, its ability to bind with Ph-BMAL1 and drive gene expression needs to be confirmed. The possibility of other partners for Ph-BMAL1 should also be investigated, and the functional significance of the N-terminus of Ph-CRY2 determined. Finally, further to the molecular work required it will be vital to develop behavioural and physiological assays that can be used to study the effects of clock gene manipulation. In *Drosophila melanogaster* a number of rhythmic phenomena have been identified including locomotor activity, eclosion, egg deposition, eye sensitivity, and oxygen consumption (Helfrich-Förster, 1996), while as a model organism *P. hawaiensis* has been largely exploited at a molecular level and often at early developmental stages, and behavioural observations are lacking. We have gathered encouraging preliminary locomotor activity results (Hunt, 2016) that align with the observations of Poovachiranon *et al.* (1986) in indicating a nocturnal peak in activity that persists in constant darkness, but it is a small dataset compromised by issues with temperature control, hunger, age-matching and water evaporation, aeration and fouling. We urge interested parties to work towards optimising such behavioural assays and also to undertake the identification of other relevant characteristics of *P. hawaiensis* that may be experimentally tractable. Only with this information in hand can one begin the task of employing *P. hawaiensis* to bring about new insights in crustacean chronobiology, a field with a rich research history that is ripe for significant advancement through the exploitation of a model with a diverse and powerful genetic toolkit.

Acknowledgements

We are deeply grateful to Ana Patricia Ramos and Michalis Averof for supplying us with animals and husbandry advice and for permitting us to search an unpublished leg and embryo tissue assembly which proved vital in identifying the PER fragment in our own head assembly. Thanks are also due to Damian Kao, Aziz Aboobaker, Anastasios Pavlopoulos and all other authors of Kao et al., (2016) for granting us access to the then unpublished *P. hawaiensis* genome. Parts of this research first appeared in the thesis of BH (Hunt, 2016). This research used the ALICE High Performance Computing Facility at the University of Leicester.

Author contributions

BH performed RNA extraction, *de novo* assembly, genome and transcriptome analysis, and drafted the manuscript. ER and EM conceived the study. All authors approved the final manuscript.

Conflict of interest

None declared.

Funding disclosure

This work was funded by the Natural Environment Research Council (NERC) UK through an Antarctic Funding Initiative grant NE/D008719/1 (ER) and a PhD studentship (BH). EBM was supported by research grant NE/N010019/1 from the Natural Environment Research Council, UK.

Data Accessibility,

The *P. hawaiensis* head transcriptome is archived at the NCBI SRA under project PRJNA399131. Newly identified/expanded putative coding and peptide sequences of core clock genes are detailed in supplementary data.

References

Allada, R., White, N. E., So, W., Hall, J. C., and Rosbash, M. 1998. A Mutant *Drosophila* Homolog of Mammalian *Clock* Disrupts Circadian Rhythms and Transcription of *period* and *timeless*. *Cell*, 93(5): 791–804.

Andrews, S. 2010. FastQC A Quality Control tool for High Throughput Sequence Data.

Biscontin, A., Wallach, T., Sales, G., Grudziecki, A., Janke, L., Sartori, E., Bertolucci, C., Mazzotta, G., Pittà, C. D., Meyer, B., Kramer, A., and Costa, R. 2017. Functional characterization of the circadian clock in the Antarctic krill, *Euphausia superba*. *Scientific Reports*, 7(1): 17742.

Bolger, A. M., Lohse, M., and Usadel, B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15): 2114–2120.

Browne, W. E., Price, A. L., Gerberding, M., and Patel, N. H. 2005. Stages of embryonic development in the amphipod crustacean,*Parhyale hawaiensis*. *Genesis*, 42(3): 124–149.

Chang, Z., Li, G., Liu, J., Zhang, Y., Ashby, C., Liu, D., Cramer, C. L., and Huang, X. 2015. Bridger: a new framework for *de novo* transcriptome assembly using RNA-seq data. *Genome Biology*, 16(1): 30–30.

Chesmore, K. N., Watson, W. H., and Chabot, C. C. 2016. Identification of putative circadian clock genes in the American horseshoe crab, *Limulus polyphemus*. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, 19: 45–61.

Christie, A. E. 2014. Identification of the first neuropeptides from the Amphipoda (Arthropoda, Crustacea). *General and Comparative Endocrinology*, 206: 96–110.

Christie, A. E., Fontanilla, T. M., Nesbit, K. T., and Lenz, P. H. 2013. Prediction of the protein components of a putative *Calanus finmarchicus* (Crustacea, Copepoda) circadian signaling system using a *de novo* assembled transcriptome. *Comparative biochemistry and physiology. Part D, Genomics & proteomics*, 8(3): 165–93.

De Coursey, P. J. 1960. Daily Light Sensitivity Rhythm in a Rodent. *Science*, 131(3392): 33–35.

Emery, P., So, W., Kaneko, M., Hall, J. C., and Rosbash, M. 1998. CRY, a Drosophila Clock and Light-Regulated Cryptochrome, Is a Major Contributor to Circadian Rhythm Resetting and Photosensitivity. *Cell*, 95(5): 669–679.

Gamble, F. W. and Keeble, F. W. 1900. *Hippolyte varians*: a Study in Colour-change. *Quarterly Journal of Microse Science*, 43: 589–703.

Gekakis, N., Staknis, D., Nguyen, H. B., Davis, F. C., Wilsbacher, L. D., King, D. P., Takahashi, J. S., and Weitz, C. J. 1998. Role of the CLOCK protein in the mammalian circadian mechanism. *Science (New York, N.Y.)*, 280(5369): 1564–1569.

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M. B., Eccles, D., Li, B., Lieber, M., Macmanes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C. N., Henschel, R., Leduc, R. D., Friedman, N., and Regev, A. 2013. *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols*, 8(8): 1494–512.

Helfrich-Förster, C. 1996. *Drosophila* rhythms: from brain to behavior. *Seminars in Cell & Developmental Biology*, 7(6): 791–802.

Helfrich-Förster, C. 2005. Neurobiology of the fruit fly's circadian clock. *Genes, Brain and Behavior*, 4(2): 65–76.

Hogenesch, J. B., Gu, Y. Z., Jain, S., and Bradfield, C. A. 1998. The basic-helix-loop-helix-PAS orphan MOP3 forms transcriptionally active complexes with circadian and hypoxia factors. *Proceedings of the National Academy of Sciences*, 95(10): 5474–9.

Hunt, B. J. 2016. *Advancing molecular crustacean chronobiology through the characterisation of the circadian clock in two malacostracan species, Euphausia superba and Parhyale hawaiensis*. Thesis, University of Leicester.

Hunt, B. J., Özkaya, , Davies, N. J., Gaten, E., Seear, P., Kyriacou, C. P., Tarling, G., and Rosato, E. 2017. The *Euphausia superba* transcriptome database, SuperbaSE: An online, open resource for researchers. *Ecology and Evolution*, 7(16): 6060–6077.

Kao, D., Lai, A. G., Stamataki, E., Rosic, S., Konstantinides, N., Jarvis, E., Di Donfrancesco, A., Pouchkina-Stancheva, N., Semon, M., Grillo, M., Bruce, H., Kumar, S., Siwanowicz, I., Le, A., Lemire, A., Eisen, M. B., Extavour, C., Browne, W. E., Wolff, C., Averof, M., Patel, N. H., Sarkies, P., Pavlopoulos, A., and Aboobaker, A. 2016. The genome of the crustacean *Parhyale hawaiensis*, a model for animal development, regeneration, immunity and lignocellulose digestion. *eLife*, 5.

Kim, D., Langmead, B., and Salzberg, S. L. 2015. HISAT: a fast spliced aligner with low memory requirements. *Nature Methods*, 12(4): 357–360.

Konopka, R. J. and Benzer, S. 1971. Clock mutants of *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*, 68(9): 2112–2116.

Letunic, I., Doerks, T., and Bork, P. 2015. SMART: recent updates, new developments and status in 2015. *Nucleic Acids Research*, 43(D1): D257–D260.

Li, W. and Godzik, A. 2006. Cd-hit: A fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, 22(13): 1658–1659.

Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, 17(1): pp. 10–12.

Nesbit, K. T. and Christie, A. E. 2014. Identification of the molecular components of a *Tigriopus californicus* (Crustacea, Copepoda) circadian clock. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, 12: 16–44.

O'Grady, J. F., Hoelters, L. S., Swain, M. T., and Wilcockson, D. C. 2016. Identification and temporal expression of putative circadian clock transcripts in the amphipod crustacean *Talitrus saltator*. *PeerJ*, 4: e2555.

Palmer, J. D. 1991. Contributions Made to Chronobiology by Studies of Fiddler Crab Rhythms. *Chronobiology International*, 8(2): 110–130.

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., and Salzberg, S. L. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology*, 33(3): 290–295.

Poovachiranon, S., Boto, K., and Duke, N. 1986. Food Preference Studies and Ingestion Rate Measurements of the Mangrove Amphipod *Parhyale hawaiensis* (Dana). *Journal of Experimental Marine Biology and Ecology*, 98(316): 129–140.

R Core Team 2018. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Renn, S. C., Park, J. H., Rosbash, M., Hall, J. C., and Taghert, P. H. 1999. A *pdf* Neuropeptide Gene Mutation and Ablation of PDF Neurons Each Cause Severe Abnormalities of Behavioral Circadian Rhythms in *Drosophila*. *Cell*, 99(7): 791–802.

Reppert, S. M. and Weaver, D. R. 2000. Comparing Clockworks: Mouse versus Fly. *Journal of Biological Rhythms*, 15(5): 357–364.

Reppert, S. M. and Weaver, D. R. 2001. Molecular Analysis of Mammalian Circadian Rhythms. *Annual Review of Physiology*, 63(1): 647–676.

Robertson, G., Schein, J., Chiu, R., Corbett, R., Field, M., Jackman, S. D., Mungall, K., Lee, S., Okada, H. M., Qian, J. Q., Griffith, M., Raymond, A., Thiessen, N., Cezard, T., Butterfield, Y. S., Newsome, R., Chan, S. K., She, R., Varhol, R., Kamoh, B., Prabhu, A.-L., Tam, A., Zhao, Y., Moore, R. A., Hirst, M., Marra, M. A., Jones, S. J. M., Hoodless, P. A., and Birol, I. 2010. *De novo* assembly and analysis of RNA-seq data. *Nature Methods*, 7(11): 909–12.

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., and Mesirov, J. P. 2011. Integrative genomics viewer. *Nature Biotechnology*, 29: 24–26.

Robinson, M. D. and Oshlack, A. 2010. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome biology*, 11(3): R25–R25.

Rutila, J. E., Suri, V., Le, M., So, W. V., Rosbash, M., and Hall, J. C. 1998. Cycle is a second bHLH-PAS clock protein essential for circadian rhythmicity and transcription of *Drosophila period* and *timeless*. *Cell*, 93(5): 805–814.

Saleem, Q., Anand, a., Jain, S., and Brahmachari, S. K. 2001. The polyglutamine motif is highly conserved at the *Clock* locus in various organisms and is not polymorphic in humans. *Human Genetics*, 109(2): 136–142.

Sandrelli, F., Costa, R., Kyriacou, C. P., and Rosato, E. 2008. Comparative analysis of circadian clock genes in insects. *Insect Molecular Biology*, 17(5): 447–463.

Sbragaglia, V., Lamanna, F., M. Mat, A., Rotllant, G., Joly, S., Ketmaier, V., de la Iglesia, H. O., and Aguzzi, J. 2015. Identification, Characterization, and Diel Pattern of Expression of Canonical Clock Genes in *Nephrops norvegicus* (Crustacea: Decapoda) Eyestalk. *PLOS ONE*, 10(11): e0141893–e0141893.

Sehgal, A., Price, J. L., Man, B., and Young, M. W. 1994. Loss of circadian behavioral rhythms and per RNA oscillations in the *Drosophila* mutant *timeless*. *Science*, 263(5153): 1603–1606.

Shin, S. W., Zou, Z., Saha, T. T., and Raikhel, A. S. 2012. bHLH-PAS heterodimer of methoprene-tolerant and Cycle mediates circadian expression of juvenile hormone-induced mosquito genes. *Proceedings of the National Academy of Sciences*, 109(41): 16576–81.

Smith-Unna, R., Boursnell, C., Patro, R., Hibberd, J. M., and Kelly, S. 2016. TransRate: reference-free quality assessment of de novo transcriptome assemblies. *Genome research*, 26(8): 1134–44.

Strauss, J. and Dircksen, H. 2010. Circadian clocks in crustaceans: identified neuronal and cellular systems. *Frontiers in Bioscience*, 15: 1040–74.

Tilden, A. R., McCoole, M. D., Harmon, S. M., Baer, K. N., and Christie, A. E. 2011. Genomic identification of a putative circadian system in the cladoceran crustacean *Daphnia pulex*. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, 6(3): 282–309.

Wickham, H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Yang, J. S., Dai, Z. M., Yang, F., and Yang, W. J. 2006. Molecular cloning of Clock cDNA from the prawn, *Macrobrachium rosenbergii*. *Brain Research*, 1067(1): 13–24.

Zhang, L., Hastings, M. H., Green, E. W., Tauber, E., Sladek, M., Webster, S. G., Kyriacou, C. P., and Wilcockson, D. C. 2013. Dissociation of circadian and circatidal timekeeping in the marine crustacean *Eurydice pulchra*. *Current biology*, 23(19): 1863–73.

Zhu, H., Yuan, Q., Froy, O., Casselman, A., and Reppert, S. M. 2005. The two CRYs of the butterfly. *Current Biology*, 15(23): R953–R954.

Figure 1: Expression of putative clock genes and housekeeping genes across 8 timepoints in light:dark 12:12 conditions. ZT = *zeitgeber* time, ZT0 = lights on, ZT12 = lights off.

Figure 2: Schematics of predicted *P. hawaiensis* core clock peptides aligned with examples from other species. Ph = *Parhyale hawaiensis*. Dm = *Drosophila melanogaster*. Es = *Euphausia superba*. Tal = *Talitrus saltator*. Ha = *Hyalella azteca*. QQQ = polyglutamine domain (defined as a run of 10 or more amino acids in which 60% or more are glutamine). [a] - peptide is predicted on the basis of BLAST alignments and read mapping. [b] - peptide is a construct of multiple separate *de novo* assembled contigs. Both Tal-CRY2 and Tal-CLK have been previously reported in a shorter form (O'Grady *et al.*, 2016).

Figure 3: Optimal neighbour-joining trees depicting the evolutionary relationships of (A) CRYP-TOCHROME and (B) TIMELESS/TIMEOUT peptides.

Table 1: *P. hawaiensis* clock gene candidates identified from transcriptome and genome mining. nt = nucleotides. aa = amino acids.

Protein	Query	Contig	E-value	Length (nt)	Length (aa)
BMAL1	ANW48377	phaw_30_tra_m.024548	2.00E-156	-	617
CASEIN KINASE II α	AAN11415	PH.k29.comp2312_seq0	0	4,885	352
CASEIN KINASE II β	AAF48093	PH.k21.comp3428_seq3	5.00E-110	2,357	222
CIRCADIAN TRIP	AAF52092	PH.k31.comp1615_seq0	2.00E-111	12,125	2,996
CLOCKWORK ORANGE	AAF54527	PH.k21.comp80764_seq0	3.00E-26	642	212
CRYPTOCHROME 2	AFV96168	PH.k21.R17911710	0	3,301	931
DOUBLETIME	AAF57110	PH.k27.comp42178_seq0	0	1,599	418
JETLAG	AAF52178	-	-	-	-
MICROTUBULE STAR (PP2A)	AAF52567	PH.k21.comp640_seq0	0	3,538	309
NEJIRE	AAF46516	PH.k51.J3947083	0	5,384	1,794
NEMO	AAF50497	PH.c118708_g1_i2	0	1,416	430
PAR DOMAIN PROTEIN 1 ε	AAF04509	phaw_30_tra_m.019467	2.00E-35	-	591
PERIOD	ALC74274	PH.k25.comp125155_seq0	4.00E-07	280	91
PIGMENT DISPERSING HORMONE	GAJQ01005023	PH.k41.S6600835	2.00E-24	795	103
PDH RECEPTOR	AAF45788	PH.k21.comp140546_seq0	5.00E-29	597	191
PROTEIN PHOSPHATASE 1	CAA39820	PH.k27.comp8511_seq1	0	2,963	328
SHAGGY	AAN09084	PH.k29.comp7576_seq1	0	5,244	457
SUPERNUMERARY LIMBS	AAF55853	PH.k21.comp23826_seq4	0	4,640	642
TIMELESS	ANW48379	phaw_30_tra_m.019341	1.00E-10	-	1,674
TWINS (PP2A)	AAF54498	PH.k21.R18132040	0	2,887	444
VRILLE	AAF52237	PH.k21.comp5330_seq6	3.00E-42	3,178	480
WIDERBORST (PP2A)	AAF56720	PH.k29.comp1930_seq3	0	9,834	458

Table 2: BLAST search results using *P. hawaiensis* clock gene candidates vs. NCBI non-redundant protein database.

Protein	Accession	Name	Species	E-value
Es-BMAL1	AFV39705	Bmal1a	*Pacifastacus leniusculus*	1.00E-161
Es-CASEIN KINASE II α	ARJ31756	Casein kinase II subunit alpha	*Litopenaeus vannamei*	0
Es-CASEIN KINASE II β	ANO53990	CKII beta	*Limulus polyphemus*	1.00E-145
Es-CIRCADIAN TRIP	NP_001084531	E3 ubiquitin-protein ligase TRIP12	*Tribolium castaneum*	0
Es-CLOCKWORK ORANGE	KOC64192	Hairy/enhancer-of-split related with YRPW motif protein 1	*Habropoda laboriosa*	3.00E-34
Es-CRYPTOCHROME 2	AFV96168	cryptochrome 2	*Talitrus saltator*	0
Es-DOUBLETIME	AUI80371	Doubletime	*Euphausia superba*	0
Es-MICROTUBULE STAR (PP2A)	KDR18186	PP2A catalytic subunit alpha	*Zootermopsis nevadensis*	0
Es-NEJIRE	CUT08824	CREB-binding protein	*Blattella germanica*	0
Es-NEMO	KZS07628	Mitogen-activated protein kinase	*Daphnia magna*	0
Es-PAR DOMAIN PROTEIN 1 ε	ABV22507	PAR domain protein 1	*Danaus plexippus*	2.00E-46
Es-PERIOD	ANN13870	period	*Macrobrachium nipponense*	4.00E-07
Es-PDH	WP_082125040	ribonuclease P protein component	*Erythrobacter luteus*	6.8
Es-PDH RECEPTOR	BAH85843	Pigment dispersing hormone receptor	*Marsupenaeus japonicus*	2.00E-103
Es-PROTEIN PHOSPHATASE 1	AQZ36563	Serine/threonine protein phosphatase 1	*Penaeus monodon*	0
Es-SHAGGY	ALK82316	glycogen synthase kinase 3 beta	*Procambarus clarkii*	0
Es-SUPERNUMERARY LIMBS	KDR19729	F-box/WD repeat-containing protein 1A	*Zootermopsis nevadensis*	0
Es-TIMELESS	AUI80375	timeless 2	*Euphausia superba*	0
Es-TWINS (PP2A)	AFK24473	serine/threonine-protein phosphatase PP2A reg. subunit B	*Penaeus monodon*	0
Es-VRILLE	AUI80373	Vrille	*Euphausia superba*	1.00E-57
Es-WIDERBORST (PP2A)	KYB26642	PP2A regulatory subunit epsilon	*Tribolium castaneum*	0