Population structure determined by comparative amplicon gene sequencing in biofilm and planktonic bacterial communities from pesticide-contaminated water

Population structure in biofilm and planktonic bacterial communities from contaminated water

Jhenifer Yonara Lima[1], Cassiano Moreira [1], Gessica Costa[1], Paloma Nathane

Nunes Freitas[1], Luiz Ricardo Olchanheski[2], Sonia Alvim Veiga Pileggi[1], Rafael

Mazer Etto[1], Christopher Staley[3], Michael Jay Sadowski[4] and Marcos Pileggi[1*]

[1] Department of Structural and Molecular Biology and Genetics, Ponta Grossa State

University, Ponta Grossa, Paraná, Brazil

[2] Department of Microbiology, University of São Paulo, Sao Paulo, Brazil

[3] The Biotechnology Institute, University of Minnesota, St. Paul, Minnesota, United

States of America

[4] Department of Soil, Water, and Climate and The Biotechnology Institute, University

of Minnesota, St. Paul, Minnesota, United States of America

[*] Corresponding Author

E-mail: pileggimarcos@gmail.com

Abstract

Molecular and bioinformatics tools for research are very important for biodiversity characterization in contaminated environments due the presence of a high percentage of nonculturable microorganisms. The wide use of pesticides in agriculture exposes microbiomes to stressful and selective conditions that demand survival strategies such as biofilm formation. The purpose of this work was to evaluate the bacterial population structure in planktonic and biofilm communities of water that was used for washing the packaging of herbicides and stored for six months in tanks. This substance is highly contaminated waste. Samples of water and biofilms from tanks and biofilms developed for short times in flasks were used for DNA isolation and 16S rRNA gene sequencing. The physicochemical conditions imposed by water used for washing the containers were inadequate for killing the bacterial genera identified according to water and wastewater standards. The variation in population structure and diversity was lower in the planktonic samples than in the biofilm samples, indicating a possible combination of genetic drift and subsequent selection of individuals surviving under stressful water conditions, such as heating and contact with agrochemicals, over a six-month period. The biofilm formation in water tanks contaminated with pesticides enabled the survival of bacterial genera harboring the essential processes for adaptation to these environments; the presence of these processes was determined according to descriptions obtained from the genomic databases. This study suggests the potential of bacterial genera identified in biofilms obtained from tanks to adapt to contaminated environments through their metabolic complexity. Thus, herbicide biodegradation kinetics can be accessed through a culturable collection obtained from these communities.

Introduction

Environmental microbiology has been gaining ground in academia over the last few years, bringing with it an interest in understanding composition, structure and function in nature [1]. Notably, contaminated environments expose microbiomes to stressful and selective conditions, and once microorganisms exhibit structural and metabolic adaptations, they can overcome these conditions and survive. These adaptive strategies are linked at the cellular level and to specific populations, allowing different responses to compounds such as pesticides [2,3] and enabling their use in biotechnological processes such as agrochemical degradation [4–6]

However, microbial diversity is high and poorly elucidated by culture techniques, even with different types of culture media, since a large percentage of these microorganisms are nonculturable [7,8]. Therefore, it is necessary to create new research tools that allow the characterization of microbiological biodiversity, i.e., molecular microbial ecology, through the use of high-throughput sequencing technologies with the aid of bioinformatics techniques for data analysis [1,9–11].

The knowledge and use of molecular technologies are fundamental to understanding the structure and functions of microbiomes and nonculturable microorganisms. Advances in genomics and metagenomics, marker genes, sequencing of the 16S rRNA gene and other molecular approaches are essential for predicting soil microbiome functions and improving agriculture production [12]. These approaches are fundamental to the creation of genomic libraries that allow us to identify and isolate enzymes with different biocatalytic activities from culturable strains, permitting their

biotechnological exploitation [13,14]. These techniques allowed Krohn-Molt et al.

(2017) to evaluate the genes of an association of algae essential to the carbon cycle with

a unique and specific microbiome. Knowledge of the taxonomic structure of the

microbiome increases the reliable handling of its latent biotechnological potential

[16,17].

Thus, the composition and relative abundance of soil microbiomes in the

Amazon were correlated with different types of land use, such as deforestation, which

leads to carbon cycle disturbances and, consequently, climate change [18]. Therefore,

microbiomes, which play key roles in biogeochemical cycles, are affected by

anthropogenic action, particularly by the use of xenobiotic compounds, leading to

changes in sustainability and environmental quality [18,19]. High-density DNA

microarray and 16S rDNA amplicon next-generation sequencing were used to study the

impact of the insecticide chlorpyrifos, the herbicide isoproturon and the fungicide

tebuconazole on the soil bacterial community, and significant differences have been

found in field experiments but not in soil microcosms exposed to these pesticides [20].

These methods were considered a guide for more specific studies in this area. A

molecular approach was also used to analyze the effects of combinations of glyphosate

and different herbicides on the rhizobacterial community of transgenic plants; this study

indicated that glyphosate alone seems to be the less aggressive [21]. Molecular

approaches also helped Dennis et al., 2018, reach the conclusion that the herbicides

glyphosate, glufosinate, paraquat, and paraquat-diquat did not significantly interfere

with soil microbial diversity and function under laboratory conditions.

In aqueous environments, which are also subject to contamination by

agrochemicals, microbiomes present two possible lifestyles: planktonic and biofilm.

The first term refers to the cells of microorganisms suspended in water (e.g., in an ocean

water column) [22]. However, one of the main adaptive strategies found in microbiomes

in contaminated or extreme environments is the formation of biofilms, in which cells

adhere to a substrate or one another by secreting a gelatinous matrix of extracellular

polymeric substances (EPS), which is configured as an extension of the cell itself

[23,24]. The community can thus capture organic compounds for metabolization by

extracellular enzymes through organized gene expression mediated by cell-cell

communication or quorum sensing. This structure is also responsible for increased

adaptability, stability, and cellular integrity [25–27]. 16S rRNA gene amplicon

sequencing was used to verify the impact of herbicides and fertilizers on planktonic

marine microbial communities at the Great Barrier Reef. Water characteristics, such as

salinity, rainfall, temperature and water quality, have a great influence on the

composition of microbial communities [28].Aquatic environments are complex and

subject to various physicochemical interferences, making it difficult to understand the

impact of agrochemicals on microbiomes.

Thus, the objective of this study was to identify the population structure of

planktonic and biofilm bacterial communities in an environment consisting of tanks

used for storing washing water from the packaging of herbicides, which results in

highly contaminated waste; thus, another objective was to glimpse adaptive strategies

through the composition of genes present in their taxonomic units.

Materials and methods

Study area

108   The object of study was water used to wash the containers (Fig 1) of 30

109   agrochemicals (S1 Table). The water used for the washing of agrochemical packages

came from an artesian well. After washing, the water was kept in two 10,000 L tanks for

6 months at the Capão da Onça Farm School, State University of Ponta Grossa, State of

Paraná, Brazil (25°05'31.3"S; 50°03'28.0"W) (altitude: 1000 m). The first tank (T1) had

water from the washing of agrochemical containers that had lost some volume by an

evaporation process at 100 °C for 190 min and was then stored in a second tank (T2)

until being transported and discarded.

Figure 1: Evaporation of the water from the washing of pesticide containers used for

DNA collection in this study. S: Sanitary sink; TR: Tractor spray washing trough; SH:

Shed where T1 and T2 are; T1: Tank 1; T2: Tank 2; W: Water filters; TI: Timer for

controlling the flow of water; TC: Solar heating temperature control panel; So: Solar

heating system.

Physicochemical analyses of water

The following physicochemical characteristics of water samples collected from

the tanks and the artesian well were analyzed according to the *Standard methods for the*

examination of water and wastewater (1998): chlorine, conductivity, color, fluorine,

pH, total dissolved solids and turbidity, biochemical oxygen demand (BOD) and

chemical oxygen demand (COD).

Collection of samples and extraction of DNA

Four liters of water were collected from T1 and T2 to obtain planktonic samples.

The water was passed through autoclaved glass fiber membranes in 750 mL triplicates:

first, a 5-µm-thick prefilter; then, 1.2 µm; and last, 0.45 µm (Merck Millipore,

Carrigtohill, Ireland). After the last membrane, the samples were stored at -80 °C. The

samples were named M1 (preheating) and M2 (postheating).

The biofilm samples were collected in triplicate from the wall surface of the

tanks in contact with the water line using 3 sterile swabs for each triplicate. The samples

were named B1 (preheating) and B2 (postheating). Likewise, biofilms were collected

from the 4 L flasks in which water samples were collected after being kept immobile for

five days. The samples were named G1 (preheating) and G2 (postheating). The material

collected in triplicate was stored at -80 °C.

Samples of planktonic origin (retained in the membranes) were macerated in

liquid nitrogen with the aid of a pestle. The swabs used for the biofilm collections were

submerged in 1 mL of saline buffer (0.9% NaCl) and homogenized for 5 min to extract

biological material from the cells in the buffer. DNA extractions were performed using

the PowerSoil DNA extraction kit (MO BIO Laboratories Inc., Carlsbad, USA), and the

material obtained was stored at -80 °C.

Sequencing

Sequencing of the samples was performed using 515F/806R primers, which are

specific for the V4 region [29]. Amplification and sequencing were performed using the

dual index method reported by the University of Minnesota Genomics Center

(Minneapolis, MN, USA) [30]. A negative control containing sterile water was included

in each amplified and sequenced sample plate. Sequencing was performed on the

Illumina MiSeq platform using an iSeq 100 System (Illumina, Inc. San Diego, USA)

[29]. Sequence data are available from the Sequence Read Archive of the National

Center for Biotechnology Information under accession number SRA: PRJNA528924.

Bioinformatics

Sequence processing was performed using mothur software (version 1.29.2).

The prokaryotic sequence data were trimmed to the first 160 nucleotides (nt), and end

pairing was carried out using fastq-join software [31]. The sequences were trimmed for

quality following a previous description for V5-V6 data [32]. The overall alignment was

performed in relation to the SILVA database ver. 119 [33], the sequences were

subjected to a 2% preclustering to remove sequence errors [34], and the chimeric

sequences were identified and removed by UCHIME software [35]. Operational

taxonomic units (OTUs) were assigned at ≥ 97% identity by complete linkage

clustering. Taxonomic attributions were made by comparing the Ribosomal Database

Project ver. 14 to a bootstrap limit of 60%, as previously described [36].

Bacterial communities were ordered based on OTU abundance using

multidimensional scaling. The relative abundance of the taxonomic groups and the

functional abundances inferred in PICRUSt were regressed in relation to the ordering

scores to determine significant correlations with the ordering scores of the bacterial

communities [37]. The 16S rDNA data were analyzed as indicated by the PICRUSt

genome prediction software [http://picrust.github.io/picrust/] from raw sequence reads

in the following environment: NumPy (1.7.1), biom-format (1.3.1), PyCogent (1.5.3),

PICRUSt (1.0.0-dev), and PICRUSt script (1.0.0-dev). Functional predictions were

assigned up to KO tier 3 for all genes. Results from the MG-RAST QIIME report were

compared with predictions from PICRUSt using the "compare_biom.py" subroutine

with normalization and observations not in the "expected data" file ignored. Results

were compared using both shotgun metagenomic and PICRUSt predictions as the

expected data. Statistical analyses were performed using the vegan package in R [38].

Results and discussion

This work discusses three levels of results to account for interferences in the
adaptive strategies presented by the different bacterial communities linked to the
contaminated environment in this study: the physicochemical characteristics of the
environment, the taxonomic composition at the time of collection, and the genes
associated with the identified taxonomic units.

Physicochemical characteristics of the tank water

185   The physicochemical data are analyzed in this section because they provide the
186 conditions under which the adaptation of the present microbiome to the environment
187 occurs. The data for the water from T1 and T2 and the artesian well whose water was
188 used in the washing are shown in Table 1.

200   **Table 1:** Physicochemical analyses of tanks 1 (T1) and 2 (T2) and the artesian well

201   (AW). Parameter values are presented as the means of triplicate measurements. WHO:

202   World Health Organization; SD: standard deviation; TDS: total dissolved solids; BOD:

203   biochemical oxygen demand; COD: chemical oxygen demand; ND: not detected. The

204   letters A, B, C, and D represent significant differences within each row.

Parameters	T1	SD	T2	SD	AW	SD	WHO
Chlorine (mg/L)	1.66^A	0.1	1.57^A	0	0^B	0	0.7^C
Conductivity (μS/cm)	418.3^A	2	413^A	0.7	13.57^B	0.564	$10 - 100^B$
Color (Pt/Co)	245^A	1.5	250^B	0	0^C	0	0^C
Fluorine (mg/L)	0.82^A	0	0.81^A	0	0.05^B	0	1.5^C
pH at 25°C	5.45^A	0.1	5.32^A	0	4.75^B	0.045	$6.5 - 8.5^C$
TDS (ppm/NaCl)	186.3^A	1	183.2^A	0.3	0^B	0	$100 - 600^A$
Turbidity (NTU)	204^A	5.8	206.3^A	3.6	0.25^B	0	1^C
BOD	80.3^A	1.5	69^B	4.6	ND	ND	0^C
COD	1352.3^A	29	1105^B	35.5	ND	ND	0^C

According to the parameters shown in Table 1, the water characteristics of tanks

1 and 2 are not significantly different from each other.

The data obtained in this work indicate that the heating process did not interfere

with the physicochemical parameters of the water in the tanks. Nevertheless, there are

significant differences between tank water and the water from the artesian well in every

analyzed item, mainly due to the presence of pesticides that significantly altered the

characteristics (Table 1) and independent of the microbiome of these communities.

The chlorine and fluorine contents of T1 and T2 are noteworthy since, according

to the World Health Organization [39], these specific values are close to the acceptable

concentrations for drinking water, even though the water from the artesian well and that

from the abovementioned tanks had not been added to these samples (Table 1).

Moreover, according to the WHO, the pH is also at inadequate levels for bacterial

growth, since the pH values are close to that of drinking water. In addition to these

parameters, the increase in conductivity and COD indicated important changes in the

physicochemical characteristics of the water of both tanks from those of water from the

artesian well, most likely related to the nonbiological oxidation processes of the

pesticides. The BOD data indicate biological activity, even in an environment

unfavorable to the development of microorganisms. COD values are usually higher than

BOD values due to the greater ease of compound oxidation by the chemical route than

by microorganisms [40].

The physicochemical data obtained in our study were compared to those of

water with different origins and from environments with different characteristics to

determine which water the T1 and T2 waters most resemble. Such environments were

that of drinking water [39], the interior transition zone of the ocean [41], the Red Sea

[42], southwest part of the Gulf of Mexico [43], a hot water spring in Siloam, South

Africa [44], an integrated system of anaerobic-aerobic reactors for the treatment of

wastewater (WW) in Ethiopia [45] and the Yeongsan River Basin of South Korea [46].

The two tanks do not significantly differ in chlorine level, but these levels are

statistically higher than those found for seawater and potable water (Table 2). However,

the levels found for the well water are statistically similar to those of the sea. Fluorine

levels are statistically similar in both tanks; however, they are lower than those of

drinking water and higher than those of the artesian well water (Table 2) [44]. The tanks

have statistically similar pH values but are more acidic than the artesian well, residual

tributaries, ocean, fresh water and drinking water and are statistically significantly

different from these sources in pH (Table 2). In general, when total dissolved solids

(ppm/NaCl, corresponding to salinity) are analyzed, our data present statistical

similarities to the levels in WW, drinking water and a hot water spring in Siloé, South

Africa. The BOD and COD rates are higher in T1 than in T2, indicating that there are

probably more contaminants and microorganisms in the former. For the same reason,

the rates in T1 and T2 are higher than those in the artesian well. The tanks have lower

BOD and higher COD than WW and a wastewater tributary (WT), probably because the

methods used in the treatment of these effluents have microorganisms that increase their

BOD values but not their COD values in relation to their values in the tanks studied in

this work (Table 2).

Table 2: Comparison of physicochemical analyses of tanks 1 (T1) and 2 (T2) and the

artesian well (AW) with different environments. Data were obtained in the literature for

environments such as the oceanic transition zone (OC), drinking water (DW), the Red

Sea (RS), Siloé Hot Spring (SHS), wastewater (WW), wastewater tributary (WT), the

Yeongsan River Basin (YRB), the Gulf of Mexico (GLF) and freshwater aquaculture

systems (FAS). No information (-). The letters A, B, C, D, E, and F represent significant

differences within each column.

Environment	Chlorine	Fluorine	BOD	COD	TDS (ppm/NaCl)	pH
T1	1.66^B	0.82^C	80.33^C	1352.33^A	186.27^D	5.45^E
T2	1.57^B	0.81^C	69.00^D	1105.00^B	183.17^D	5.32^E
AW	0.00^D	0.05^D	0.00^F	0.00^D	0.00^D	4.75^F
OC	0.00^D	-	-	-	-	7.78^C
DW	0.70^C	1.50^B	-	-	600.00^A	7.50^C
RS	0.94^C	-	-	-	39826.47^D	-
SHS	44.35^A	6.11^A	-	-	197.32^D	9.50^B
WW	-	-	4886.26^A	308.91^C	2593.69^D	7.66^C
WT	-	-	12547.50^B	395.00^C	9470.50^C	10.40^A
YRB	-	-	3.28^E	-	-	7.61^C
GLF	-	-	-	-	23602.07^B	-

FAZ	-	-	-	-	-	7.09[D]

In this way, tanks T1 and T2 were characterized as environments with

inadequate levels of chlorine and fluorine for the survival of the genera identified in this

study. However, these genera were probably able to tolerate these conditions, either by

inactivity in planktonic form or by organic and metabolic activity in biofilms [47].

It is important to remember that these physicochemical analyses reflect the

planktonic community more directly than they do biofilms. Biofilms, although

interfacing with the environment, most likely present modifications that improve

metabolism and the adaptability of microorganisms to the studied environment [26], as

will be discussed next.

273   Biodiversity and bacterial taxonomy

The identification of the taxonomic groups present in the different niches

evaluated in this study may indicate changes in bacterial population structures adapting

to the stress produced by the physicochemical conditions and contamination. Sample

coverage data in terms of the Simpson indices from the 16S rRNA gene sequencing and

diversity calculations are shown in Fig 2.

Figure 2: Measurements of alpha diversity in the planktonic triplicate (M1A, M1B,

M1C and M2A, M2B, M2C) and biofilm triplicate (B1A, B1B and B2A, B2B) samples

from the tanks and biofilm samples from the container's triplicates (G1A, G2B, G2C

and G2A, G2B, G2C).

Analyses of the sequences obtained in every community studied in this work

showed that the most representative classes are Gammaproteobacteria,

Alphaproteobacteria and Betaproteobacteria, as shown in Fig 3.

Figure 3: Abundance of bacterial classes in the biofilm samples from the containers and

planktonic samples from tanks 1 and 2. B1 and B2: taxonomic units identified from pre-

and postheating biofilms, respectively; M1 and M2: taxonomic units identified from

pre- and postheating planktonic samples, respectively; G1 and G2: taxonomic units

identified from biofilms formed in containers with pre- and postheating water,

respectively.

The most representative families identified by the sequences obtained in every

community are shown in Fig 4.

Figure 4: Abundance of bacterial families in biofilm samples from tanks 1 and 2. B1

and B2: taxonomic units identified from pre- and postheating biofilms, respectively; M1

and M2: taxonomic units identified from pre- and postheating planktonic samples,

respectively; G1 and G2: taxonomic units identified from biofilms formed in containers

with pre- and postheating water, respectively.

The number of sequences assigned by community and by taxonomic category

and their respective percentages within each community are shown in S1 Fig.

Simpson's diversity index demonstrates the distribution of alpha diversity (Fig

1). The sequences obtained and identified as the genera of the planktonic communities

(M1 and M2) are represented in the biofilm communities (B1 and B2, G1 and G2) but

have different diversity indices that depend on selection and adaptation to the

environment (Figs 1 and S1 Fig).

The data shown in Figs 3 and 4 indicate that there is great variation in the

population structure in different communities in relation to classes and families. The

event of water heating is a nondirectional selective process since none of the taxonomic

groups identified in this study have resistance responses for high temperatures, but it

changes the number of taxonomic units, resulting in decreased diversity in the

postheating samples for every community and less intense changes for the planktonic

communities than for the biofilm communities. The effect of this genetic drift that

began in the planktonic community can be observed in Fig 2 and leads to different

diversity and abundance among the communities that settled in the biofilm structure in

the tanks and in the flasks before and after the heating process. The planktonic

community has a direct influence on biofilm formation [48]. Chronologically, the

communities are formed in the following order: first T1, followed by B1 and then G1.

This ordering is valid for postheating communities, with the proviso that the T2

community is formed after the T1 community.

A genetic drift effect is more conspicuous in the biofilm population structures in

flasks G1 and G2, formed from the planktonic strains, than in the structures in other

communities. This is probably because they had less time to consolidate population

structures that are more adapted to the environment, as observed in B1 and B2.

Population structure variation is lower in the planktonic samples than in the

biofilm samples in tanks and flasks, which may indicate that there was a combination of

the genetic drift of individuals who survived the heating and subsequent selection of

differentiated characteristics to allow individuals to adapt to the tank environment. This

environment for water contaminated with 30 different agrochemicals and with

physicochemical conditions unfavorable to the survival of microorganisms identified in

this work was considered stressful (Table 1 and S1 Table). The organization of bacteria

in biofilms in tanks and containers, with their cooperative or antagonistic interactions,

may lead to a specific combination of species with increased adaptability to the

environment [49]. Thus, the most dominant sequences in B2, corresponding to the

genera *Enterobacter* and *Acinetobacter*, are poorly represented in the other communities

studied, including the biofilms in B1, G1 and G2 (Fig 2).

Theoretically, the transition from planktonic to biofilm life involves changes in

the regulation of different groups of genes and can be seen as a transition phase before

the colonization of a new environment [50]. Therefore, this transition should not present

significant changes in the population structure (Fig 3). Throughout the establishment of

the community, the most abundant genera tend to retain most of the genes related to

basic cellular processes, leading to the same function being shared by individuals of

different species and not only being distributed by the selection of the fittest [51].

Nevertheless, the reduction in diversity after the water heating event possibly led

to genetic drift, altering the fine structure of the M2, G2 and B2 in relation to that of the

M1, G1 and B1 communities and the retention of genes involved in central adaptive

processes. If the genetic drift and the functional diversity of different bacterial

communities were less preponderant, the diversity indices of B1 and B2, B2 and G2,

and G1 and G2 should be closer [48]than what was observed in Fig 2. In addition, the

data shown in S1 Fig indicate that the same genus but with different 16S sequences may

contribute to different communities adapted to the environments studied in this work

[52]. Thus, the genetic drift hypothesis was established from the sequences identified

and not by the taxonomic units.

Aquatic environments are complex [28]; therefore, the analysis of changes in the

structure of communities of microorganisms that are made to adapt to environmental

contaminants is difficult. Restricted environments with a high number of agrochemicals,

such as those presented in this paper, allow basic hypotheses to be tested with increased

safety before being tested at a larger environmental scale.

Functional diversity

The comparative data in Fig 5 cluster genes by the sequences of the 16S rRNA

gene identified in this study. This analysis was carried out as a survey of

biotechnological potential [18,19] mainly related to biodegradation in the storage tanks

of water contaminated with agrochemicals, which are considered a significant

environmental liability.

Figure 5: Ordering the sites based on OTU abundance by multidimensional scaling.

Only the bacterial genus (green), genetic process (red), cellular process (black),

environmental process (blue) and metabolic pathways (yellow) with $p<0.05$ after 999

permutations are displayed. Each vector shows the direction of increase for a given

variable, and its length indicates the strength of the correlation between the variable and

the ordination scores. Legend: M: Motility; GD: Growth and Death; TP: Transportation

in Catabolism; MT: Membrane Transport; SMI: Signaling Molecule Interaction; ST:

Signal Transduction; TL: Translation; TC: Transcription; RR: Replication and Repair;

FSD: Folding, Sorting and Degradation; AA: Amino Acid; SM: Secondary Metabolites;

C: Carbohydrate; E: Energy; EZ: Enzyme; G: Glycan; L: Lipid; CV: Cofactors and

Vitamins; OA: Other Amino Acids; TP: Terpenoids and Polyketides; N: Nucleotide;

XB: Xenobiotic Biodegradation.

Nonmetric multidimensional scaling (NMDS) shows the correlations between

the number and types of 16S rRNA gene sequences grouped by the 6 different

communities. Genes related to bacterial motility have a high correlation with the two

planktonic communities (S2 Table).

On the other hand, genes associated with other cellular processes, environmental

processes, gene information processes and metabolic processes have a strong correlation

with the biofilm community B2 (Table 1), which is predominantly composed of the

genera *Acinetobacter* and *Enterobacter*.

To picture the potential for the degradation and bioremediation of pesticides and

relate the hypotheses about adaptation to the environment described in this study, the

degradation capacity of the pesticides (S1 Table) was related to the genera of the most

prevalent bacteria identified by the 16S rRNA gene sequencing (Fig 2).

Bacteria from the genus *Pseudomonas* (representing 45% of the structure in B1

and 8% of the structure in B2, Fig 4) present in communities identified in Hawaiian soil

can degrade pyraclostrobin, which is present in several fungicides [53]. A strain of

Pseudomonas aeruginosa isolated from rice-cultivated soil has the capacity to degrade

propiconazole [54]. Other strains of the genus *Pseudomonas* isolated from agricultural

soil degrade thiamethoxam [55].

Strains of the genus *Enterobacter* that were identified in Egyptian soil

(representing 63% of the structure in B2, Fig 4) degraded the herbicide Topik [56].

Pseudomonas and *Acinetobacter* (representing 15% of the structure in B2, Fig 4) from

communities where there was tar processing were identified as potential degraders of

mineral oil [57]. A strain of the genus *Burkholderia* was isolated from paddy fields with

pesticide application in Koppal, India, and was a degrader of propiconazole, a

component of the fungicide Tilt [54].

The genera in the literature that are most related to agrochemical biodegradation

are the same as those that dominate B2. Thus, the genera identified in this study have

agrochemical degradation potential and the capacity for structuring biofilm

communities, as shown in Fig 2, suggesting the viability of searching for promising

strains among the culturable collection obtained in this study. The high temperature to

which the planktonic community was subjected is bactericidal for any genus observed

in this study [39]. Therefore, the nondirectional selective effect induced by the high

temperature was independent of the genomes of the identified genera.

Comparative 16S rRNA gene sequencing

In this work, a model was proposed to explain a possible range of characteristics

whose expression could confer adaptive advantages over the stressful environment of

water tanks used in the washing of agrochemicals. The population fluctuations observed

were the result of genetic drift and the capacity for biofilm formation, which involves

membrane transport, interaction of signaling molecules and signal transduction, and

functional bacterial versatility [58]. The biofilm structure depends on the presence of

EPS, which involves the production of glycans and lipid transportation in catabolism.

Biofilms confer genomic plasticity and phenotypic flexibility to the species that

compose them, providing successful reproduction in stressful environments (Fig 5)[23].

Theoretically, these two characteristics, plasticity and flexibility, and community

diversity, which are higher in the planktonic community (Fig 2) than in the biofilm

community, may lead to species fixation over time or even extinction of the less

efficient. Some mutations may be randomly lost by genetic drift; however, significant

advantageous mutations are more likely to fix in the population, as is the case of

motility in a planktonic community, the only feature that has a high correlation with this

community (Fig 5). If the selection pressure in a community is strong, the frequencies

of genotypes that improve competence increase in the population [59]. This model can

be applied for mutations in populations with one species or populations of different

species but with gene flow potential, as might occur in the B2 community.

The environments studied in this work passed through a sequence of

physicochemical changes involving chemical contamination and heat over six months.

During these events, genera that had the capacity to develop useful functions [58] to

consolidate biofilms, such as *Enterobacter* spp. and *Acinetobacter* spp., dominated the

B2 community [48,60], which was formed after the communities M1, M2 and B1 by

selection and genetic drift.

Environmental biofilms have a level of metabolic complexity higher than that of

their individual components [61]. Thus, the study of biofilm formation in environments

contaminated with pesticides, which can be stressful to microorganisms, has the

potential to produce information about response systems for adaptation. This

information provides a possible way to exploit enzymatic potentials, mainly those

related to biodegradation and bioremediation of herbicides [50], in the kind of waste

used in this study. New knowledge can be obtained for the exploitation of the enzymatic

potential of bacterial communities under these environmental conditions, mainly the

potential related to the biodegradation kinetics of herbicides.

Conclusions

The population structure and abovementioned functional profile derived from

amplicon sequencing of bacterial communities from tanks that contained water used for

washing the packaging of herbicides and that had been stored for six months to form a

highly contaminated waste were studied in this work. The communities were characterized as tank biofilms, flask biofilms, and planktonic samples, all from pre- and postheating treatments and formed at different periods of time. Only two genera identified in this study, *Acinetobacter* and *Enterobacter*, were dominant in the tank biofilms after water heating, which probably acted as a genetic drift agent for the planktonic community. A database was accessed after the amplicon sequencing to compare genomic compositions, and this analysis showed that these genera have a metabolic potential for survival under these environmental conditions by possessing genes such as those related to biofilm formation, structure and membrane transport, quorum sensing and xenobiotic degradation. The genetic drift, induced by water heating, decreased the diversity of the bacterial biofilm community, which probably survived as large bacterial populations under the harmful conditions found in the tank due their genomic diversity and adaptation to environmental responses, since it is known that biofilms have a level of metabolic complexity higher than that of their individual components. Thus, the hypotheses about metabolic abilities, generated for a small and controlled environment as that studied in this work, can be tested with communities isolated from the biofilms formed in these tanks. Thus, new knowledge can be obtained for the exploitation of the enzymatic potential of bacterial communities, mainly the potential related to the biodegradation kinetics of herbicides.

Acknowledgments

The authors thank Maria Janina Pinheiro Diniz for assisting in the preparation and execution of experiments.

References

1.   Zhou J, He Z, Yang Y, Deng Y, Tringe SG, Alvarez-cohen L. High-Throughput

Metagenomic Technologies for Complex Microbial Community Analysis: Open

and   Closed   Formats.   CEUR   Workshop   Proc   2015;1542:33–6.

doi:10.1128/mBio.02288-14.Copyright.

2.   Dobrzanski T, Gravina F, Steckling B, Olchanheski LR, Sprenger RF, Espírito

Santo BC, et al. Bacillus megaterium strains derived from water and soil exhibit

differential responses to the herbicide mesotrione. PLoS One 2018;13:1–24.

doi:10.1371/journal.pone.0196166.

3.   Prione LP, Olchanheski LR, Tullio LD, Santo BCE, Reche PM, Martins PF, et al.

GST activity and membrane lipid saturation prevents mesotrione-induced cellular

damage in Pantoea ananatis. AMB Express 2016;6:1–12. doi:10.1186/s13568-

016-0240-x.

4.   Martins PF, Martinez CO, Carvalho G De, Irajara P, Carneiro B, Azevedo RA, et

al.   Selection   of   Microorganisms   Degrading   S-Metolachlor   Herbicide

2007;50:153–9.

5.   Olchanheski LR, Dourado MN, Beltrame FL, Zielinski AAF, Demiate IM,

Pileggi SA V, et al. Mechanisms of tolerance and high degradation capacity of

the herbicide mesotrione by Escherichia coli strain DH5-α. PLoS One 2014;9:1–

8. doi:10.1371/journal.pone.0099960.

6.   Silva TM, Stets MI, Mazzetto AM, D.Andrade F, Pileggi SAV, Fávero PR, et al.

Degradation of 2,4-D herbicide by microorganisms isolated from Brazilian

contaminated soil. Brazilian J Microbiol 2007;38:522–5. doi:10.1590/S1517-

83822007000300026.

7.   Deng Y, Deng C, Yang J, Li B, Wang E. Novel Butane-Oxidizing Bacteria and

Diversity of bmoX Genes in Puguang Gas Field. Front Microbiol 2018;9:1–13.

doi:10.3389/fmicb.2018.01576.

8.   Oueriaghli N, Castro DJ, Llamas I, Béjar V, Martínez-Checa F. Study of bacterial

community composition and correlation of environmental variables in Rambla

Salada, a hypersaline environment in South-Eastern Spain. Front Microbiol

2018;9:1–17. doi:10.3389/fmicb.2018.01377.

9.   Battin TJ, Besemer K, Bengtsson MM, Romani AM, Packmann AI. The ecology

and biogeochemistry of stream biofilms. Nat Rev Microbiol 2016;14:251–63.

doi:10.1038/nrmicro.2016.15.

10.   Gutarowska B, Celikkol-Aydin S, Bonifay V, Otlewska A, Aydin E, Oldham AL,

et al. Metabolomic and high-throughput sequencing analysis-modern approach

for the assessment of biodeterioration of materials from historic buildings. Front

Microbiol 2015;6:1–13. doi:10.3389/fmicb.2015.00979.

11.   Ranjan R, Rani A, Metwally A, McGee HS, Perkins DL. Analysis of the

microbiome: Advantages of whole genome shotgun versus 16S amplicon

sequencing. vol. 469. Elsevier Ltd; 2016. doi:10.1016/j.bbrc.2015.12.083.

12.   Fierer N. Embracing the unknown: disentangling the complexities of the soil

microbiome. Nat Rev Microbiol 2017. doi:10.1038/nrmicro.2017.87.

13.   Al-Amoudi S, Razali R, Essack M, Amini MS, Bougouffa S, Archer JAC, et al.

Metagenomics as a preliminary screen for antimicrobial bioprospecting. Gene

2016;594:248–58. doi:10.1016/j.gene.2016.09.021.

14.   Madhavan A, Sindhu R, Parameswaran B, Sukumaran RK, Pandey A.

517   Metagenome Analysis: a Powerful Tool for Enzyme Bioprospecting. Appl

518   Biochem Biotechnol 2017;183:636–51. doi:10.1007/s12010-017-2568-3.

519   15.   Krohn-Molt I, Alawi M, Förstner KU, Wiegandt A, Burkhardt L, Indenbirken D,

520   et al. Insights into Microalga and bacteria interactions of selected phycosphere

521   biofilms using metagenomic, transcriptomic, and proteomic approaches. Front

522   Microbiol 2017;8:1–14. doi:10.3389/fmicb.2017.01941.

523   16.   Ji P, Rhoads WJ, Edwards MA, Pruden A. Impact of water heater temperature

524   setting and water use frequency on the building plumbing microbiome. ISME J

525   2017;11:1318–30. doi:10.1038/ismej.2017.14.

526   17.   Ofaim S, Ofek-Lalzar M, Sela N, Jinag J, Kashi Y, Minz D, et al. Analysis of

527   microbial functions in the rhizosphere using a metabolic-network based

528   framework for metagenomics interpretation. Front Microbiol 2017;8:1–14.

529   doi:10.3389/fmicb.2017.01606.

530   18.   Kroeger ME, Delmont TO, Eren AM, Meyer KM, Guo J, Khan K, et al. New

531   Biological Insights Into How Deforestation in Amazonia Affects Soil Microbial

532   Communities Using Metagenomics and Metagenome-Assembled Genomes.

533   Front Microbiol 2018;9:1–13. doi:10.3389/fmicb.2018.01635.

534   19.   Imfeld G, Vuilleumier S. Measuring the effects of pesticides on bacterial

535   communities in soil: A critical review. Eur J Soil Biol 2012;49:22–30.

536   doi:10.1016/j.ejsobi.2011.11.010.

537   20.   Storck V, Nikolaki S, Perruchon C, Chabanis C, Sacchi A, Pertile G, et al. Lab to

538   field assessment of the ecotoxicological impact of chlorpyrifos, isoproturon, or

539   tebuconazole on the diversity and composition of the soil bacterial community.

540   Front Microbiol 2018;9:1–13. doi:10.3389/fmicb.2018.01412.

541   21.   Valverde JR, Marín S, Mellado RP. Effect of herbicide combinations on bt-maize

542   rhizobacterial   diversity.   J   Microbiol   Biotechnol   2014;24:1473–83.

543   doi:10.4014/jmb.1405.05054.

544   22.   Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, et al.

545   Structure and function of the global ocean microbiome 2015;348:1–10.

546   23.   Decho AW, Gutierrez T. Microbial extracellular polymeric substances (EPSs) in

547   ocean systems. Front Microbiol 2017;8:1–28. doi:10.3389/fmicb.2017.00922.

548   24.   Maunders E, Welch M. Matrix exopolysaccharides; the sticky side of biofilm

549   formation. FEMS Microbiol Lett 2017;364:1–10. doi:10.1093/femsle/fnx120.

550   25.   Anutrakunchai C, Bolscher JGM, Krom BP, Kanthawong S, Chareonsudjai S,

551   Taweechaisupapong S. Impact of nutritional stress on drug susceptibility and

552   biofilm structures of Burkholderia pseudomallei and Burkholderia thailandensis

553   grown in static and microfluidic systems. PLoS One 2018;13:e0194946.

554   doi:10.1371/journal.pone.0194946.

555   26.   Lerch TZ, Chenu C, Dignac MF, Barriuso E, Mariotti A. Biofilm vs. Planktonic

556   Lifestyle: Consequences for Pesticide 2,4-D Metabolism by Cupriavidus necator

557   JMP134. Front Microbiol 2017;8:1–11. doi:10.3389/fmicb.2017.00904.

558   27.   Zhao K, Li Y, Yue B, Wu M. Genes as early responders regulate quorum-sensing

559   and control bacterial cooperation in Pseudomonas aeruginosa. PLoS One

560   2014;9:1–8. doi:10.1371/journal.pone.0101887.

28.   Angly FE, Heath C, Morgan TC, Tonin H, Rich V, Schaffelke B, et al. Marine

microbial communities of the Great Barrier Reef lagoon are influenced by

riverine floodwaters and seasonal weather events. PeerJ 2016;4:e1511.

doi:10.7717/peerj.1511.

29.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al.

Ultra-high-throughput microbial community analysis on the Illumina HiSeq and

MiSeq platforms. ISME J 2012;6:1621–4. doi:10.1038/ismej.2012.8.

30.   Gohl DM, Vangay P, Garbe J, MacLean A, Hauge A, Becker A, et al. Systematic

improvement of amplicon marker gene methods for increased accuracy in

microbiome studies. Nat Biotechnol 2016;34:942–9. doi:10.1038/nbt.3601.

31.   Aronesty E. Comparison of Sequencing Utility Programs. Open Bioinforma J

2013;7:1–8. doi:10.2174/1875036201307010001.

32.   Staley C, Gould TJ, Wang P, Phillips J, Cotner JB, Sadowsky MJ. Evaluation of

water sampling methodologies for amplicon-based characterization of bacterial

community structure. J Microbiol Methods 2015;114:43–50.

doi:10.1016/j.mimet.2015.05.003.

33.   Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig WG, Peplies J, et al. SILVA:

a comprehensive online resource for quality checked and aligned ribosomal RNA

sequence data compatible with ARB. Nucleic Acids Res 2007;35:7188–96.

34.   Huse SM, Welch DM, Morrison HG, Sogin ML. Ironing out the wrinkles in the

rare biosphere through improved OTU clustering. Environ Microbiol

2010;12:1889–98. doi:10.1111/j.1462-2920.2010.02193.x.

35.   Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. UCHIME improves

584   sensitivity and speed of chimera detection. Bioinformatics 2011;27:2194–200.

585   36.   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid

586   assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ

587   Microbiol 2007;73:5261–7. doi:10.1128/AEM.00062-07.

588   37.   Langille MGI, Clemente JC, Knight R, Burkepile DE, Caporaso JG, McDonald

589   D, et al. Predictive functional profiling of microbial communities using 16S

590   rRNA   marker   gene   sequences.   Nat   Biotechnol   2013;31:814–21.

591   doi:10.1038/nbt.2676.

592   38.   R Foundation for Statistical Computing. R: A language and environment for

593   statistical computing. vol. 2. 2007.

594   39.   World Health Organization. Guidelines for Drinking-water Quality 4th ed.,

595   WHO,   Geneva,   p.   340.   World   Heal   Organ   2011.   doi:10.1016/S1462-

596   0758(00)00006-6.

597   40.   Gholizadeh MH, Melesse AM, Reddi L. A comprehensive review on water

598   quality parameters estimation using remote sensing techniques. Sensors

599   (Switzerland) 2016;16. doi:10.3390/s16081298.

600   41.   Mende DR, Bryant JA, Aylward FO, Eppley JM, Nielsen T, Karl DM, et al.

601   Environmental drivers of a microbial genomic transition zone in the ocean's

602   interior. Nat Microbiol 2017;2:1367–73. doi:10.1038/s41564-017-0008-3.

603   42.   Thompson LR, Williams GJ, Haroon MF, Shibl A, Larsen P, Shorenstein J, et al.

604   Metagenomic covariation along densely sampled environmental gradients in the

Red Sea. ISME J 2017;11:138–51. doi:10.1038/ismej.2016.99.

43. Godoy-Lozano EE, Escobar-Zepeda A, Raggi L, Merino E, Gutierrez-Rios RM,

Juarez K, et al. Bacterial Diversity and the Geochemical Landscape in the

Southwestern Gulf of Mexico. Front Microbiol 2018;9:1–15.

doi:10.3389/fmicb.2018.02528.

44. Tekere M, Lötter A, Olivier J, Jonker N, Venter S. Metagenomic analysis of

bacterial diversity of Siloam hot water spring, Limpopo, South Africa. African J

Biotechnol 2011;10:18005–12. doi:10.5897/AJB11.899.

45. Desta AF, Assefa F, Leta S, Stomeo F, Wamalwa M, Njahira M, et al. Microbial

community structure and diversity in an integrated system of anaerobic-aerobic

reactors and a constructed wetland for the treatment of tannery wastewater in

Modjo, Ethiopia. PLoS One 2014;9:1–22. doi:10.1371/journal.pone.0115576.

46. Jang J, Di DYW, Lee A, Unno T, Sadowsky MJ, Hur HG. Seasonal and

genotypic changes in Escherichia coli phylogenetic groups in the Yeongsan River

basin of South Korea. PLoS One 2014;9:5–12.

doi:10.1371/journal.pone.0100585.

47. Lewenza S, Abboud J, Poon K, Kobryn M, Humplik I, Bell JR, et al.

Pseudomonas aeruginosa displays a dormancy phenotype during long-term

survival in water 2018:1–19.

48. Hallatschek O, Hersen P, Ramanathan S, Nelson DR. Genetic drift at expanding

frontiers promotes gene segregation. Proc Natl Acad Sci 2007;104:19926–30.

doi:10.1073/pnas.0710150104.

49.   Herschend J, Damho ZB V, Marquard AM, Sven B, Sørensen SJ, Hä P, et al. A

628   meta-proteomics approach to study the interspecies interactions affecting

629   microbial biofilm development in a model community 2017:1–13.

630   doi:10.1038/s41598-017-16633-6.

631   50.   Berlanga M, Guerrero R. Living together in biofilms □: the microbial cell factory

632   and its biotechnological implications. Microb Cell Fact 2016:1–11.

633   doi:10.1186/s12934-016-0569-5.

634   51.   Stephens WZ, Wiles TJ, Martinez ES, Jemielita M, Burns AR, Parthasarathy R,

635   et al. Identification of population bottlenecks and colonization factors during

636   assembly of bacterial communities within the zebrafish intestine. MBio

637   2015;6:1–11. doi:10.1128/mBio.01163-15.

638   52.   Łukasik P, Nazario K, Van Leuven JT, Campbell MA, Meyer M, Michalik A, et

639   al. Multiple origins of interdependent endosymbiotic complexes in a genus of

640   cicadas. Proc Natl Acad Sci 2018;115:E226–35. doi:10.1073/pnas.1712321115.

641   53.   Chen X, He S, Liang Z, Li QX, Yan H, Hu J, et al. Biodegradation of

642   pyraclostrobin by two microbial communities from Hawaiian soils and metabolic

643   mechanism. J Hazard Mater 2018;354:225–30.

644   doi:10.1016/j.jhazmat.2018.04.067.

645   54.   Satapute P, Kaliwal B. Biodegradation of propiconazole by newly isolated

646   Burkholderia sp . strain BBK _ 9. 3 Biotech 2016;6:1–10. doi:10.1007/s13205-

647   016-0429-3.

648   55.   Rana S, Jindal V, Mandal K, Kaur G, Gupta VK. Thiamethoxam degradation by

649   Pseudomonas and Bacillus strains isolated from agricultural soils. Environ Monit

Assess 2015;187. doi:10.1007/s10661-015-4532-4.

56.   Ibrahim GGA, Amin MK, Hassan AA, El-Sheikh EA. IDENTIFICATION OF

PESTICIDES DEGRADING BACTERIA ISOLATED FROM EGYPTIAN

SOIL Chemicals and Reagents. Zagazig J Genet Biotechnol 2015;42:1129–43.

57.   Popp N, Schlömann M, Mau M. Bacterial diversity in the active stage of a

bioremediation system for mineral oil hydrocarbon-contaminated soils.

Microbiology 2006;152:3291–304. doi:10.1099/mic.0.29054-0.

58.   Roller M, Lucić V, Nagy I, Perica T, Vlahoviček K. Environmental shaping of

codon usage and functional adaptation across microbial communities. Nucleic

Acids Res 2013;41:8842–52. doi:10.1093/nar/gkt673.

59.   Brooks AN, Beer KD, Turkarslan S, Baliga NS, Li FY. Adaptation of cells to

new environments. Wiley Interdiscip Rev Syst Biol Med 2011;3:544–561.

doi:10.1002/wsbm.136.Adaptation.

60.   Komarova NL. Spatial interactions and cooperation can change the speed of

evolution of complex phenotypes. Proc Natl Acad Sci 2014;111:10789–95.

doi:10.1073/pnas.1400828111.

61.   Flemming HC, Wuertz S. Bacteria and archaea on Earth and their abundance in

biofilms. Nat Rev Microbiol 2019;17:247–60. doi:10.1038/s41579-019-0158-9.

668   Supporting information

669   **S1 Fig:** Number of sequences assigned by community and by taxonomic category and

670   their respective percentages within each community. M1: preheating planktonic

community; M2: postheating planktonic community; B1: preheating biofilm community; B2: postheating biofilm community; G1: preheating container biofilm community preheating; G2: postheating container biofilm community.

S1 Table: Agrochemicals whose containers were washed and the water used in the process to obtain the samples evaluated in this study.

S2 Table: Correlations of the functional characteristics identified in our analyses of the studied communities. M1: preheating planktonic community; M2: postheating planktonic community; B2: postheating biofilm community; r2: coefficient of determination; p: coefficient of significance.

Tank with water used to clean pesticide containers

Pesticide containers

Biofilm and planktonic amplicon sequencing

High-diversity planktonic communities

GAT AAAT CT GGTCTT ATT TCC

Biofilm collection from water previoulsy heated

β-Proteobacteria   α-Proteobacteria

M

γ-Proteobacteria   GD

TP

Acidobacteria_Gp4

FSD
TL RR TC
SM V AT
AA L N
SM C CV
X EZ OA
TP

Low-diversity biofilm communities

High potencial for degradation and biofilm pathways in biofilm communities

Genetic Drift