Controlling evolutionary dynamics

to optimize microbial bioremediation

Shota Shibasaki and Sara Mitri

Department of Fundamental Microbiology, University of Lausanne, Switzerland

Short running title: Optimizing microbial bioremediation

Author contribution: SS and SM conceived of the project, SS built and analyzed the model, SS and SM wrote the manuscript.

E-mail addresses: SS (shota.shibasaki@unil.ch), SM (sara.mitri@unil.ch)

Corresponding author's detail: sara.mitri@unil.ch

Acknowledgements: We thank Masato Yamamichi and Xiang-Yi Li for useful feedback on the earlier versions of the manuscript. SS is funded by the Nakajima foundation and the University of Lausanne and SM by European Research Council Starting Grant 715097.

Data availability: Python code and csv files used for simulations and analytic calculations are available at GitHub.

Conflict of interest: there is no conflict of interest.

Article type: Original Research

Controlling evolutionary dynamics
to optimize microbial bioremediation

Abstract

Some microbes have a fascinating ability to degrade compounds that are toxic for humans in a process called bioremediation. Although these traits help microbes survive the toxins, carrying them can be costly if the benefit of detoxification is shared by all surrounding microbes, whether they detoxify or not. Detoxification can thereby be seen as a public goods game, where non-degrading mutants can sweep through the population and collapse bioremediation. Here, we constructed an evolutionary game theoretical model to optimize bioremediation in a chemostat initially containing "cooperating" (detoxifying) microbes. We consider two types of mutants: "cheaters" that do not detoxify, and mutants that become resistant to the toxin through private mechanisms that do not benefit others. By manipulating the concentration and flow rate of a toxin into the chemostat, we identified conditions where cooperators can exclude cheaters that differ in their private resistance. However, eventually, cheaters are bound to invade. To overcome this inevitable outcome and maximize detoxification efficiency, cooperators can be periodically reinoculated into the population. Our study investigates the outcome of an evolutionary game combining both public and private goods and demonstrates how environmental parameters can be used to control evolutionary dynamics in practical applications.

Key words: public goods game, evolutionary game theory, eco-evolutionary feedback, chemostat, detoxification

1 Introduction

Microbes can be detrimental or vital to our health, our environment, and our economy. Much of applied microbiology strives to control these species, by promoting the growth of beneficial species, and suppressing that of harmful ones. We have achieved huge breakthroughs over centuries in eliminating pathogens and preventing dangerous diseases in humans, animals and plants. At the same time, microbes have played important roles in enhancing agriculture and food production (Wolfe and Dutton, 2015), and more recently in the production of biofuels or other chemicals (Antoni et al., 2007; Giri et al., 2020; Quin and Schmidt-Dannert, 2014; Ryan Georgianna and Mayfield, 2012), and in the degradation or "bioremediation" of toxic compounds, such as heavy metals or waste water (Atashgahi et al., 2018; Bertrand et al., 2015; Dixit et al., 2015; Kang et al., 2016; Zaccaria et al., 2020). Despite these exciting advances, these approaches remain only partially successful. In medical microbiology, the emergence of resistant pathogens has led to a life-threatening global health crisis (Tacconelli et al., 2018). On the engineering side, we lack sufficient understanding to maximize the benefits gained from microbes, in terms of production rates or the efficiency of degradation of toxic compounds (Giri et al., 2020).

In all the examples above, there are two obstacles for controlling microbes. First, microbes' response to

changes in their environment, and to the behavior of neighbouring cells may cause the extinction of species that contribute to community function or that protect against pathogens (ecological instability). Second, their large population size and short generation times mean that microbes quickly acquire mutations that can lead to evolutionary instability. Species can thereby gain resistance to antibiotics or lose their ability to perform a desired community function (Akita and Kamo, 2015; Bull and Barrick, 2017; Kumar et al., 1991; Rugbjerg et al., 2018b,a). In the bioremediation of heavy metals, for example, mutants that do not degrade these harmful compounds may invade the original population (Ellis et al., 2007; O'Brien et al., 2014) and exclude it. To optimize the functional efficiency of a microbial system, therefore, we need to consider both ecological and evolutionary dynamics of microbial populations (Schuster et al., 2010).

In this study, we focus in on a bioremediation problem, and develop a mathematical model to investigate the simple case of a single species degrading a toxic compound in a chemostat. We first explore its ecological and evolutionary stability, and second, use this knowledge to optimize the efficiency of toxin degradation over time. In our model, we assume that toxins are harmful to the microbes (e.g., heavy metals), who degrade them by secreting a costly product (e.g., extracellular enzymes). The benefit of degrading toxins is shared by the whole population as, for simplicity, our model does not include spatial structure. Toxin degradation can therefore be regarded as a public goods game (Samuelson, 1954; Broom et al., 2018), as defined in microbiology (Hummert et al., 2014; Smith and Schuster, 2019; West et al., 2007a). It is important to note that not all bioremediation systems correspond to public goods games (e.g., Smith et al. (1998); Röling et al. (2002)), but here we focus on a subset of these systems where the compound to be degraded is toxic to the microbes and its degradation is costly. In such a system, the evolutionary instability of bioremediation is expected (Ellis et al., 2007; O'Brien et al., 2014).

In our model, microbes can adopt one of four strategies: they can be product secretors that pay a cost to contribute to the public good (cooperators) or non-secretors that do not (cheaters) (see O'Brien et al. (2014) for an empirical example). In addition, microbes can be sensitive to the toxins, or can acquire resistance, for example, by activating efflux pumps to expel toxins from within the cell (Blair et al., 2015; Bottery et al., 2016; Rojo-Molinero et al., 2019), or thickening the cell wall. In essence, public good secretion can also be seen as a form of extracellular resistance to the toxin. In other words, here we consider toxin resistance through private or public means, whereby a cell benefits only itself or also the remaining population, respectively.

The population and evolutionary dynamics are then analyzed using evolutionary game theory, where a strategy is considered to be evolutionarily stable if it is not invaded by mutants with another strategy (Maynard Smith and Price, 1973). Evolutionary game theory typically considers the frequencies of strategies (i.e., frequencies sum to one), for example in the replicator dynamics (Cressman and Tao, 2014). Here, however, since toxin concentration decreases with the absolute number or density of degrader microbes (cooperators), our model describes the dynamics of the densities of strategies (as in

Hauert et al. (2006, 2008); Gokhale and Hauert (2016)). And since the microbes' death rate depends on toxin concentration in the environment and their resistance level, our model also includes environmental feedback (as in Gong et al. (2018); Tilman et al. (2020); Weitz et al. (2016)), where each strategy affects the environment differently, and the changing environment affects the fitness of each strategy differently.

We use this model to derive a protocol for optimal toxin degradation. We first show that cooperators that secrete toxin-degrading enzymes are excluded by cheaters that do not, if they have the same level of resistance to the toxin. This recapitulates a well-known result that can be explained by the tragedy of the commons (Hardin, 1968). However, we then show that cooperators can invade a population of cheaters if their level of toxin resistance is different. Since we assume that cheaters are unlikely to acquire double mutations leading to cooperators with a different resistance level, maintaining degradation is only possible if we periodically inoculate these cooperators back into the chemostat. The success of this approach relies on the ability of cooperators to invade cheaters of different resistance. We then calculate the values of the experimentally controllable parameters (inoculation probabilities of cooperators, chemostat dilution rates and inflowing toxin concentrations) that maximize the cumulative efficiency of detoxification.

In sum, our model combines population dynamics, evolutionary dynamics, and environmental feedback to optimize a population-level function. Integrating ecology and evolution into microbial public goods games is increasingly appreciated in microbial applications (Moreno-Fenoll et al., 2017; Sanchez and Gore, 2013). And while optimizing bioremediation is the case study we are considering here, our approach of controlling evolutionary dynamics by changing environmental parameters can be applied to many other microbial functions.

2 Model

In our scenario (Fig. 1A), cells can take on one of four strategies depending on whether they produce the enzymes that degrade the toxin (cooperate) or not (cheat), and whether the cells are resistant to the toxin (resistant) or not (sensitive): sensitive cooperator (sCo), sensitive cheater (sCh), resistant cooperator (rCo), and resistant cheater (rCh).

We begin by defining the bacterial population dynamics in our system. The dynamics of the density x of each strategy $i \in \{\text{sCo}, \text{sCh}, \text{rCo}, \text{rCh}\}$ in a chemostat is defined by growth, death and dilution out of the chemostat:

$$\frac{dx_i}{dt} = x_i \left[r_i \left(1 - \sum_j x_j \right) - \delta_i (T) - \alpha \right], \tag{1}$$

where r_i is the intrinsic growth rate of strategy i due to nutrients that are not explicitly defined in the model, $\delta_i (T)$ is the death rate of strategy i given toxin concentration T, and α is the dilution rate. The total densities of the four strategies $\sum_i x_i$ should be lower or equal to one in Eq (1); i.e., the carrying capacity is equal to one in the absence of death or dilution. In this formulation, a useful proxy for fitness

Figure 1: Schematic illustration of the model and examples of the dynamics

(A) In our scenario, a fluid with toxin concentration T_{in} flows into the chemostat while the same amount of fluid with toxin concentration T flows out. The dilution rate of the chemostat is α. Each cell can exhibit one of four strategies: sensitive cooperator (sCo), resistant cooperator (rCo), sensitive cheater (sCh), or resistant cheater (rCh). Cooperators produce enzymes that degrade the toxin, while cheaters do not. The toxin kills cells depending on its concentration, but resistant cells have a lower death rate compared to sensitive cells. Whether the cells are cooperators or cheaters is independent of their resistance level and vice versa. (B-E) Examples of the dynamics in the absence of mutation are shown (a.u. = "arbitrary units"). In each panel, the black solid line represents the toxin concentration T and the dashed black line the toxin concentration flowing into the chemostat T_{in}. Detoxification efficiency at each time-point is proportional to the vertical distance between dashed and solid black lines. Other colored lines represent the cell densities of one of the four strategies (solid dark green: sCo, dashed dark green: rCo, thick lime green: sCh). (B) sCo grows and degrades the toxins. (C) sCo is invaded and excluded by sCh. (D) rCo is invaded and excluded by sCo. (E) sCh is invaded and excluded by rCo. Note that the initial conditions in (C-E) are the stable equilibria of the mono-culture of the resident strategies, while in (B) we begin with a low density of sCo and $T(0) = T_{\text{in}}$. Parameter values are $\alpha = 0.1$, $T_{\text{in}} = 0.2$ (E) or 0.3 (otherwise), $f_{\text{max}} = 0.5$, $K_d = 0.2$, $r = 1$, $c_d = 0.15$, $c_r = 0.2$ (E) or 0.3 (otherwise), $d_{\text{max}} = 1$, $K_s = 0.2$ (E) or 0.3 (otherwise), $K_r = 0.6$, and $n = 1$ (E) or 3 (otherwise).

is the ratio between intrinsic growth and death at a given toxin concentration, whether death is by toxin or by dilution:

$$W_i(T) \equiv \frac{r_i}{\delta_i(T) + \alpha}. \tag{2}$$

At an equilibrium, $W_i(T) = 1/\left(1 - \sum_j x_j\right)$ should be satisfied for any strategy i that exists in the chemostat ($x_i > 0$). In addition, when $W_i(T) > W_j(T)$, strategy i increases faster or decreases slower than strategy j. For simplicity, this basic model assumes that strategies cannot mutate into each other. We extend it to include mutations in Appendix 5.

First, the intrinsic growth rates in this model differ depending on the costs each strategy pays. Cooperators pay a cost, c_d, for producing degrading enzymes, which are regarded as a public good since they reduce environmental toxicity and the death rate of all cells independently of their strategy. In addition, toxin resistance carries a cost, c_r. Such fitness costs, where resistant cells have lower fitness than sensitive ones in the absence of toxins, have been observed in many species (Andersson and Levin, 1999; Andersson and Hughes, 2010; San Millan and MacLean, 2017). In contrast to the production of degrading enzymes, however, where all cells benefit from decreased toxicity, the evolution of resistance can be regarded as an investment into a private good, where only the resistant cells themselves benefit. Assuming that the costs are additive, the intrinsic growth rate r_i of each strategy is defined as follows:

$$
\begin{align}
r_{\text{sCo}} &= r(1 - c_d) \tag{3a} \\
r_{\text{sCh}} &= r \tag{3b} \\
r_{\text{rCo}} &= r(1 - c_d - c_r) \tag{3c} \\
r_{\text{rCh}} &= r(1 - c_r), \tag{3d}
\end{align}
$$

where r is the maximum intrinsic growth rate.

Cellular death rate $\delta_i(T)$ increases with toxin concentration T, and is represented by a Hill equation as is common in models of death by drugs (Chou, 2006):

$$\delta_i(T; K) = d_{\max}\frac{T^n}{T^n + K_i^n}, \tag{4}$$

where $d_{\max}$ is the maximum death rate, K_i is the half maximal toxin concentration of strategy i, and n is the Hill coefficient, which determines the steepness of the function. Resistance can be modeled either by increasing K_i or decreasing n (Sampah et al., 2011). Here we assume that resistant cells have a larger K than sensitive cells: $K_r > K_s$, such that they reach $d_{\max}$ at a higher toxin concentration than the sensitive cells. Note that the toxin concentration T changes over time, as described below.

Due to the dilution in the chemostat, a proportion of cells of each strategy i flows out of the chemostat. The dilution rate into and out of the system is denoted by α.

To describe the population dynamics of each strategy, however, it is necessary to also formulate the dynamics of the toxin concentration because it affects the microbes' death rate, and because the toxin concentration changes over time as cooperators detoxify it. The dynamics of the toxin concentration T in the chemostat are defined by the concentration flowing into and out of the chemostat, and detoxification by cooperators:

$$\frac{dT}{dt} = \alpha T_{\text{in}} - f\left(x_{\text{Co}}\right) T - \alpha T, \tag{5}$$

where T_{in} is the toxin concentration flowing into the chemostat, and $f\left(x_{\text{Co}}\right)$ is the degradation rate which is assumed to follow a Michaelis-Menten function:

$$f\left(x_{\text{Co}}\right) = f_{\text{max}} \frac{x_{\text{Co}}}{x_{\text{Co}} + K_d}, \tag{6}$$

where $x_{\text{Co}} = x_{\text{sCo}} + x_{\text{rCo}}$; i.e., the sum of sensitive and resistant cooperators. Whether cooperators are resistant or sensitive has no impact on toxin degradation. K_d represents the density of cooperators x_{Co} that gives half the maximum of $f\left(x_{\text{Co}}\right)$. All parameters of the model are listed in Table 1.

As the goal of this study is to maximize the efficiency of detoxification, we define detoxification efficiency ϕ as the difference between the toxin concentration flowing into and out of the chemostat multiplied by the dilution rate:

$$\phi\left(\alpha, T_{\text{in}}, T\right) = \alpha\left(T_{\text{in}} - T\right). \tag{7}$$

With this definition, ϕ is proportional to the amount of detoxified liquid and is composed of the degree of detoxification and the amount of liquid flowing out of the chemostat. Although this equation gives the efficiency at any time t, we mainly focus on the efficiency at an equilibrium.

3 Results

3.1 All strategies can persist in mono-cultures with no mutation

We first analyse whether cooperators and cheaters can persist in mono-culture. Remember that only cooperators produce public goods that degrade the toxin and thereby increase the survival of all cells in the chemostat. When a few cooperators (either sensitive or resistant, $0 < x_i(0) \ll 1$, $i \in \{\text{sCo}, \text{rCo}\}$) are introduced into the system, they increase and converge to an equilibrium of positive density (Fig. 1B) if and only if

$$r_i > \delta_i\left(T_{\text{in}}\right) + \alpha \Leftrightarrow W_i\left(T_{\text{in}}\right) > 1, \tag{8}$$

where i is the focal strategy (see Appendix 1 for derivation). By solving $dT/dt = 0$ and $dx_i/dt = 0$, one can find a trivial equilibrium ($x_i = 0$) and one or more non-trivial equilibria ($T^*, x_i^* > 0$) that should

satisfy:

$$T^* = \frac{\alpha T_{\text{in}}}{\alpha + f(x_i^*)}, \tag{9a}$$

$$x_i^* = 1 - \frac{\delta_i(T^*) + \alpha}{r_i}, \tag{9b}$$

which we can calculate numerically using Newton's method.

In the absence of cooperators, we assume that the toxin concentration is equal to the incoming toxin because cooperators are the only degrader cells. In the case of a mono-culture of cheaters then, $T^* = T_{\text{in}}$ regardless of cell density, and their density at a stable equilibrium x_i^*, $i \in \{\text{sCh}, \text{rCh}\}$ is positive and given by Eq (9b) if inequality (8) holds (see Appendix 1 for details). From now on, we focus on conditions where cell density converges to a positive value in mono-culture (i.e. where inequality (8) holds).

3.2 Cheaters can invade a population of cooperators

Next, we ask what happens when a cheater mutant invades a population of cooperators at its non-trivial equilibrium or vice versa. Cheater mutants can invade and exclude a population of cooperators at any toxin concentration and independently of whether they are both sensitive or both resistant, as long as the resident and mutant have the same resistance levels (e.g., sCo and sCh) (Figs. 1C, and 2A). In contrast, cooperators are unable to invade a population of cheaters of the same resistance level at any toxin concentration. These findings recapitulate the classical result that cheaters will always dominate in a well-mixed environment (West et al., 2007b) because cooperators pay a cost for producing degrading enzymes, but are as sensitive to the toxin as cheaters. In other words, the tragedy of commons (Hardin, 1968) occurs in this case.

3.3 Toxin concentration determines invasion of sensitive and resistant cells

For mutants that differ in their private resistance level (e.g., sCo and rCo), the toxin concentration determines whether invasion succeeds or not (Figs. 2B-E). Intuitively, this is because the benefit of being resistant to the toxin is quite low when its concentration is very low. Similarly, when toxin concentration is very high, the death rate of the resistant strain is close to that of the sensitive strain and too high to compensate for the cost of private resistance. Under these conditions, sensitive cells can invade a population of resistant ones (Fig. 1D). Instead, resistant cells can invade a population of sensitive cells when the toxin concentration in the chemostat is intermediate. Two strategies that differ only in their resistance level never stably coexist (see Appendix 3 for derivation).

Figure 2: Short-term evolutionary dynamics of pairwise invasions

(A, F) Diagrams of pairwise invasion analysis when a single-mutation (A) or double-mutations (F) occur. A →
B represents that A is invaded by B, and the color of each arrow shows the condition for successful
invasion. Black arrows represent successful invasion regardless of the toxin concentration, while pink
and blue arrows represent toxin-concentration-dependent invasion (that invasion succeeds when the
toxin concentration is low or high, or when the toxin concentrations are intermediate, respectively).
(B-E, G-J) Examples of an invasion state space for each pair of strategies. A → B represents that A is a
resident strategy and B is an invader. Pairs of sCo and sCh, and rCo and rCh are omitted since
cheaters are fitter than cooperators in these pairs, regardless of the parameter values. In the yellow
areas, the resident strategies are not invaded by the invaders, while the invasion succeeds in the orange
areas. Each solid line is a boundary under which the resident strategy persists in mono-culture. The
dashed line in each panel represents where the fitness proxy W of resident and invader strategies are
equal at an equilibrium reached in a mono-culture of the resident strategy. Note that residents can
coexist with invaders under certain conditions (see Appendix 3). Parameter values in (B-E, G-J) are
$f_{\max} = 0.5$, $K_d = 0.2$, $r = 1$, $c_d = 0.15$, $c_r = 0.2$, $d_{\max} = 1$, $K_s = 0.3$, $K_r = 0.6$, and $n = 3$. Note that
panels (B-E, G-J) are examples of the state space given the parameter values; different parameter
values will show different invasion landscapes.

3.4 Cooperators can invade and coexist with cheaters of different resistance levels

Thus far, we have considered whether mutants can invade a resident population that differs in only one trait, their private or their public resistance (i.e. cooperative toxin degradation). While we assume that the time to reach an equilibrium following a single mutant invasion is shorter than the time for a second mutation to occur, we nevertheless explore invasions by such double mutants here (Fig. 2F). Depending on the concentration of toxins, rCo and sCh can invade each other's populations, as can sCo and rCh (Figs. 2G-J). This dependency on toxicity follows the same logic as for the invasion of a resistant mutant into a sensitive population described above: when the toxin concentration is intermediate, rCo have a much lower death rate than sCh, and the benefit of resistance exceeds the sum of the cost of cooperation c_d and resistance c_r. If, on the other hand, the toxin concentration is either too low or too high, resistance to the toxin does not provide enough of an advantage to overcome its cost, leading instead to the invasion of sCh into a population of rCo. The same logic, albeit with different thresholds can explain the invasion of sCo into rCh and vice versa (Figs. 2G-J). In sum, invasions of double mutants into resident populations that differ in both public and private resistance depends on toxin concentrations (see Appendix 2).

Once a mutant has invaded, whether it will coexist with the resident population is unclear because, as cooperators increase, toxin concentration decreases, which changes the fitness landscape. In other words, increasing cooperator density can decrease the fitness difference between cooperators and cheaters. In Appendix 3 we show that cooperators and cheaters of different resistance levels (e.g. rCo and sCh) can indeed stably coexist at certain parameter ranges. Nevertheless, these two coexisting strains can then be invaded by cheaters with different resistance (e.g. rCh), which excludes the other two strategies (see Appendix 4). In other words, cooperators are never evolutionarily stable because they can be invaded and excluded by cheaters of the same resistance level, as we show next.

3.5 In the long-term, cooperators are unlikely to be maintained

Having analyzed the outcomes of the invasion of all mutants in the short-term (Figs. 2A and F), we can now predict how the population in the chemostat will change in the long-term. Regardless of which type we start with, as cells mutate, the population will transition between different genotypic "states", which can be represented by the state transition diagram in Fig. 3. The probability of cooperators mutating into cheaters or vice versa is given by μ_1, while μ_2 is the probability to change the level of resistance. For simplicity, we assume that these mutation probabilities are very small. Once a mutation occurs, we use Eq (1) as outlined above to take us to the following equilibrium state. We assume that no further mutations will occur before the equilibrium is reached, but relaxing this assumption does not alter the overall dynamics (Appendix 7). For example, sCh will appear in the population of sCo with probability

Figure 3: Schematic illustration of the state transitions for long-term evolutionary dynamics

Arrows represent state transitions resulting from natural selection. Solid arrows show transitions that are independent of toxin concentration, and dashed arrows transitions that depend on toxin concentration, here depicted for a given T_{in} and α. The transition from sCo to rCo and vice versa do not occur if cooperators coexist with cheaters. Values along the arrows represent state transition probabilities.

$\mu_1 (1 - \mu_2)$ and exclude it. Then, rCh can appear in the population of sCh with probability $(1 - \mu_1) \mu_2$, but may invade or not, depending on the toxin concentration (Fig. 3).

Although in principle cooperators can invade a population of cheaters that differ in the level of resistance (e.g., rCo can invade sCh), (i) invasion success depends on α and T_{in}, (ii) double mutations are expected to be rare ($\mu_1 \mu_2$ is close to 0) and (iii) if a cheater mutant of the same resistance as the cooperator invades (e.g. rCh), it will dominate the population and replace the cooperators. Accordingly, it is very difficult to maintain cooperators in the chemostat due to natural selection. This brings us to one of the main findings of the study: even though cooperators and cheaters can coexist under some conditions, to maintain costly microbial detoxification, it is necessary to inoculate cooperators manually and to change α and T_{in} to favor their survival. Crucially, though, because cooperators are able to invade cheaters of opposite sensitivity, these inoculations can maintain cooperators – and thereby detoxification – in the short-term. In the following sections, we show how to control the values of α and T_{in} and inoculation probabilities to maximize the efficiency of detoxification.

3.6 Culture conditions can be controlled to optimize detoxification efficiency

Ultimately, our goal is to maximize the efficiency of detoxification ϕ, which depends on the absolute abundance of the two types of cooperators in the chemostat. In turn, these abundances can be controlled by changing the culture conditions through two parameters: the dilution rate α and the toxin concentration flowing into the chemostat T_{in}.

Figure 4: Optimal detoxification at each equilibrium

The efficiency of detoxification at an equilibrium state $\phi\left(\alpha, T_{\text{in}}, T\right)$ given α and T_{in} is represented by color in each panel: (A) only sCo, (B) coexistence of sCo with rCh, (C) only rCo, and (D) coexistence of rCo with sCh. The red stars represent the maximum efficiency of detoxification in each. In the areas above the dashed gray lines in each two left panels, the cooperator cannot persist in mono-culture (inequality 8). Parameter values are $f_{\max} = 0.5$, $K_d = 0.2$, $r = 1$, $c_d = 0.15$, $c_r = 0.3$, $d_{\max} = 1$, $K_s = 0.3$, $K_r = 0.5$, and $n = 1$ (A-B) or $n = 3$ (C-D). We used different values of n in the top and bottom rows to allow cooperators to coexist with cheaters with a different level of resistance.

To maximize the objective function in Eq (7), we consider three stable equilibrium states with different toxin concentrations flowing out of the chemostat: (i) $T = T_{\text{in}}$ when only cheaters are present, regardless of the values of α and T_{in}, (ii) $T = T_{ij}^{\dagger}$ when cooperators i coexist with cheaters j, which have different resistant levels, and (iii) $T = T_i^*$ when only one type of cooperators i is present. In the latter two cases, one can calculate the equilibria (analytically or numerically) and their corresponding detoxification efficiency ϕ for each culture condition (values of α and T_{in}). We can then find the optimal culture conditions that maximize this efficiency (Fig. 4), although the equilibrium can be ecologically unstable for some parameter values.

Intuitively, the maximum efficiency is larger in a mono-culture of cooperators than in a co-culture of cooperators and cheaters of different levels of resistance (see Appendix 6). If cheaters can be excluded from the population by changing α and T_{in}, the optimal strategy for cultivation is (i) to exclude the cheaters by adjusting the culture conditions, and then (ii) to change the culture conditions to maximize the productivity of a mono-culture of cooperators.

3.7 Inoculating cooperators to optimize detoxification efficiency

Above, we showed that even though they are unlikely to appear by double mutation, cooperators can invade a population of cheaters if their level of resistance is different (Fig. 3). Instead of waiting for these mutants to arise naturally, it would be more efficient to manually inoculate cooperators into the population, and to change α and T_{in} to allow them to invade successfully and to exclude the cheaters. Assuming that we cannot observe the prevalence of each strategy at will, the problem is how often to inoculate sensitive or resistant cooperators to maximize detoxification efficiency over time. If cooperator inoculation probabilities are too small, cheaters will dominate the population, leading to a detoxification efficiency of zero. If they are too large, we may inoculate cooperators unnecessarily (e.g., sCo into a mono-culture of sCo) or when they cannot invade (e.g., sCo into a mono-culture of sCh). Such unfavorable inoculations can be costly because they can require a higher in-flowing toxin concentration T_{in}, and result in reduced detoxification efficiency for some time.

To optimize cooperator inoculation probabilities m_1 and m_2 for the sensitive or resistant cooperators, respectively, we consider population state transitions as a Markov chain with discrete time-steps s, and find the values of m_1 and m_2 that maximize the total amount of toxin degradation (Figs. 5A, and A.9). Transitions in this Markov chain model can occur either due to mutations at probabilities $\boldsymbol{\mu} = (\mu_1, \mu_2)$ or inoculations at probabilities $\boldsymbol{m} = (m_1, m_2)$, resulting in the transition matrix $P(\boldsymbol{m}; \boldsymbol{\mu})$. The state distribution vector $\boldsymbol{\pi}(s) = (\pi_i(s))$ where $\pi_i(s)$ is the probability that the population is in state i at time step s ($\sum_i \pi_i(s) = 1$ for $s = 0, 1, \ldots, \infty$). Because the Markov chain is ergodic when cooperators can exclude cheaters that differ in their resistance level (Fig. 5A, but see Appendix 7 for a case where cooperators cannot exclude cheaters), the probability distribution of the population states converges to a unique stationary distribution $\boldsymbol{\pi}^*$ in the limit of $s \to \infty$, regardless of the initial distribution $\boldsymbol{\pi}(0)$:

$$\boldsymbol{\pi}^*(\boldsymbol{m}; \boldsymbol{\mu}) = \boldsymbol{\pi}^*(\boldsymbol{m}; \boldsymbol{\mu}) P(\boldsymbol{m}; \boldsymbol{\mu}). \tag{10}$$

Even though the transitions are probabilistic, we assume that the establishment of strategies following mutation or inoculation (i.e., short-term dynamics) is deterministic. Relaxing this assumption by introducing an establishment probability ϵ into the transition matrix $P(\boldsymbol{m}; \boldsymbol{\mu})$ does not change the ergodicity of the Markov chain, and we arrive at the stationary distribution in the same manner. In Appendix 7 we show how to calculate the expected cumulative efficiency of detoxification Φ from the beginning of the cultivation to time step s (defined in Eq (A.67)). This calculation is somewhat cumbersome, but for large s, Φ is approximately proportional (see Appendix 7) to the expected efficiency of detoxification at the stationary distribution $\boldsymbol{\pi}^*$:

$$\hat{\Phi}(\boldsymbol{m}; \boldsymbol{\mu}) \equiv \sum_i \phi_i \pi_i^*(\boldsymbol{m}; \boldsymbol{\mu}). \tag{11}$$

To maximize the expected cumulative detoxification efficiency Φ, therefore, we can calculate $\boldsymbol{m}$ that

 maximizes Eq (11). In Box 1 we show an example of how to use this approach in practice.

Box 1: An example of optimizing detoxification efficiency

Imagine that we have set up the experimental system described, and would like to compute the optimal inoculation probabilities. We first need to define the Markov chain to make predictions, and second, we need to experimentally measure the parameters of our bacterial strains, in particular their degradation efficiency ϕ_i at each of the different states i of the Markov chain.

To establish the Markov chain, we begin with a few simplifying assumptions: (i) that $\mu_2 = 0$, such that cooperators can only invade a population of cheaters that differ in the level of resistance by inoculation, and (ii) that mutations and the manual inoculation of a cooperator strategy can occur only in a mono-culture (i.e., at most two strategies can exist simultaneously in the population). We further assume that the parameters are in a range where at certain (α, T_{in}), (iii) sCo and rCo can mutually exclude each other, and (iv) sCo and rCo can exclude rCh and sCh, respectively. Under these assumptions (we relax (i) below and (ii)-(iv) in Appendix 7), the Markov chain consists of at least 14 states: four mono-culture situations, six transient situations where two strategies coexist, and four situations where the introduction of cooperators is unfavorable (see Fig. A.9 for the diagram). A simplified schematic of this model is shown in Fig. 5A.

By experimentally measuring mutation probabilities $\boldsymbol{\mu}$ and detoxification efficiencies $\boldsymbol{\phi} = (C\phi_i)$ for each state i, where C is a positive constant to change the time scale of ϕ_i into a discrete time step, we can calculate the probability distribution $\boldsymbol{\pi}(s, \boldsymbol{m}) = (\pi_i(s, \boldsymbol{m}))$ as a function of time step s and inoculation probabilities $\boldsymbol{m} = (m_1, m_2)$. Then, Dynamic Programming (DP) provides the optimal $\boldsymbol{m}$ that will maximize the cumulative expected detoxification efficiency Φ when the experiment finishes at time step s. Starting from a mono-culture of sCo, the two solid lines in Fig. 5B represent m_1 and m_2 provided by DP given some fictitious yet reasonable values of $\boldsymbol{\mu}$ and $\boldsymbol{\phi}$. At first, the optimal values of m_1 and m_2 are zero, because the state of the population is most likely to be a mono-culture of sCo, in which case inoculating cooperators would be pointless. However, as time passes, mutations will arise, and the population is likely to transition to a state of sCh mono-culture; then, the optimal values of m_1 and m_2 increase. At about 1,000 time steps, the optimal values of m_1 and m_2 converge to the values which maximize detoxification efficiency at the stationary distribution $\hat{\Phi}$ described by Eq (11). When $\mu_2 > 0$ (relaxing assumption (i)), the number of states increases and the state transition diagram becomes more complex. As long as the Markov chains is ergodic, however, it is possible to find the stationary distribution $\boldsymbol{\pi}^*$ and the optimal values of m_1 and m_2 that maximize Eq (11). We show how to find the optima for non-ergodic Markov chains in Appendix 7.

Table 1: List of parameters

Notation	Range	Description
α	$(0, 1]$	dilution rate of the chemostat
r	$(0, 1]$	maximum intrinsic growth rate of the microbe
c_d	$(0, 1]$	cost of cooperation (production of the degrading enzyme)
c_r	$(0, 1]$	cost of resistance to the toxin
$d_{\max}$	$(0, 1]$	maximum death rate by toxin
K_s	$[0, K_r)$	half maximal effective toxin concentration of the sensitive cells
K_r	$(K_s, 1]$	half maximal effective toxin concentration of the resistant cells
n	$(0, \infty]$	Hill coefficient
T_{in}	$(0, 1]$	toxin concentration flowing into the system
$f_{\max}$	$(0, 1]$	maximum degradation rate of the toxin
K_d	$(0, 1]$	half maximal effective cooperator density of the degradation rate
μ_1	$[0, 1]$	mutation probability in the function of detoxification
μ_2	$[0, 1]$	mutation probability in the resistance level
m_1	$[0, 1]$	inoculation probability of sCo
m_2	$[0, 1]$	inoculation probability of rCo

Figure 5: Optimization problem assuming that resistance mutations do not occur

(A) A simplified schematic illustration of the state transition diagram when $\mu_2 = 0$ and cooperators can exclude cheaters of different resistance levels. A → B represents a transition from state A to state B. Colored arrows represent transitions that occur through the introduction of sensitive and resistant cooperators in blue and pink, respectively. Dashed arrows indicate transient states where two strategies coexist. The full diagram containing 14 states is shown in Fig. A.9. (B) The optimal values of m_1 and m_2 (the blue and pink solid lines, respectively) which maximize the cumulative efficiency defined by Eq (A.67) calculated using Dynamic Programming. The two dashed lines represent the values of m_1 and m_2 which maximize Eq (11). The initial state of the population is a mono-culture of sCo. Around 1,000 time steps, the optimal values of m_1 and m_2 for Eq (A.67) converge to the values which maximize Eq (11). In practice, the detoxification efficiency at each of the 14 states of the model as well the mutation probabilities would be experimentally measured and plugged into the model to calculate the values of m_1 and m_2 that would maximize cumulative efficiency. The plot above was generated using the following fictitious values, as an illustration: $\mu_1 = 0.01$, $\mu_2 = 0$, $\boldsymbol{\phi} = \{0.4, 0.2, 0, 0.15, 0.3, 0.15, 0, 0.2, 0.35, 0.35, 0.3, 0, 0.2, 0\}$. See Appendix 7 for more detail.

4 Discussion

In this study, we have shown how to control the ecological and evolutionary dynamics of a microbial population growing in a chemostat in order to optimize the bioremediation of a toxic liquid. Public goods games where cooperators increase the growth rates of surrounding cells at a cost to themselves have been extensively studied, both empirically and theoretically (Allen et al., 2016; Hauert et al., 2006, 2008; Griffin et al., 2004; Gokhale and Hauert, 2016; Sanchez and Gore, 2013). Rather than increasing the growth rate of others, cooperators in our model degrade toxic compounds, which decreases the death rate of surrounding cells (O'Brien et al., 2014). This scenario enables us to introduce the evolution of resistance to the toxin, e.g. through efflux pumps, as a private good, which we base on studies of drug-dose effect and resistance to it (Chou, 2006; Sampah et al., 2011). Unsurprisingly, cheaters always exclude cooperators with the same private resistance level because detoxification is costly (West et al., 2007b). We show, however, that because the benefit of private resistance depends on the toxin concentration in the chemostat, cooperators can invade a population of cheaters that differ in their private resistance. The co-occurrence of two strains that differ both in their degradation ability as well as their resistance level is unlikely to suddenly arise by mutation, especially if we assume that mutations are rare. To maintain the degradation of toxins, therefore, it is necessary to periodically inoculate cooperators into the chemostat while changing the dilution rate and in-flowing toxin concentration to guarantee invasion success.

Optimal values for these parameters (cooperator inoculation probabilities, dilution rate, and in-flowing toxin concentration) that maximize the detoxification efficiency of the system can be calculated using our model. As input, the model requires experimental measurements of growth and death rates of the chosen microbe and its mutants (i.e., intrinsic growth rate of each strategy r_i, maximum death rate d_{max}, Hill coefficient n, median-effect toxin concentrations K_s, K_r, and degradation efficiencies ϕ of cooperators).

Our model and its results can also apply to problems other than bioremediation that involve survival in toxic environments. In essence, we are studying the evolutionary dynamics of public resistance (which is cooperative) and private resistance (which is not). Consider, analogously, two types of antibiotic resistance mechanisms: public mechanisms are costly and benefit the producing cell as well as its neighbors such as extracellular secretion of antibiotic-degrading enzymes (e.g. β-lactamases (Yurtsev et al., 2013)), and private resistance mechanisms only benefit the producing cells, such as efflux pumps. The evolutionary dynamics in a scenario whereby cells can switch between these different resistance mechanisms and being sensitive to the antibiotic correspond to Figs. 2A and F. In this case, however, an objective function would aim to minimize rather than maximize the densities of the most resistant strains. Another interesting aspect is that the benefits of resistance depend on toxin concentration in the chemostat, which is affected by the density of cooperators and the toxin concentration flowing into the chemostat. In other words, the public goods game affects the benefit of the private goods. This is why cooperators can invade a population of cheaters when they differ in their resistance level (Fig. 2F).

²⁹¹ Of course, our model relies on a number of assumptions and focuses only on a subset of possible biore-
²⁹² mediation systems. First, we only consider extracellular toxin degradation (e.g. by enzyme secretion),
²⁹³ while toxins can also be degraded inside cells (O'Brien and Buckling, 2015). For intra-cellular degrada-
²⁹⁴ tion, a different functional form of detoxification $f(x_{\mathrm{Co}})$ would be necessary, but we expect similar results
²⁹⁵ as long as this function increases monotonically with the density of cooperators. Indeed, the invasion
²⁹⁶ analysis is independent of the form of $f(x_{\mathrm{Co}})$. Similarly, we assume that toxins kill the microbes and
²⁹⁷ that their degradation does not contribute to growth. In reality, many compounds that are undesirable
²⁹⁸ for humans are instead used as substrates by microbes (Atashgahi et al., 2018). This latter case is simpler
²⁹⁹ than the one we consider here, since detoxification is no longer cooperative and there is no risk of cheaters
³⁰⁰ arising and collapsing the system. Finally, detoxification may carry a negligible cost, for example if it the
³⁰¹ toxic compound is neutralized by a change in pH, which occurs naturally due to a microbe's metabolism.

³⁰² Another issue is how to define detoxification efficiency ϕ. Rather than Eq (7) one could, for example,
³⁰³ define ϕ as the time needed for the toxins to decrease to a negligible concentration. This would change the
³⁰⁴ optimal culture conditions α and T_{in}, but not the procedure to find the optimal introduction probabities
³⁰⁵ of cooperators $\boldsymbol{m}$, which are independent of the formulation of ϕ. Our model also fixes some parameters,
³⁰⁶ such as the Hill coefficient n, which can evolve in reality (Sampah et al., 2011). Similarly, the cost
³⁰⁷ of resistance c_r can decrease over time due to compensatory evolution (Andersson and Hughes, 2010;
³⁰⁸ San Millan et al., 2014). Allowing these parameters to evolve would make it more difficult for sCo to
³⁰⁹ invade rCh because the relative fitness of rCh will increase.

³¹⁰ We also assume that our system is well-mixed and that there are no spatial gradients within the chemo-
³¹¹ stat. Spatial structure, for example, whereby detoxifying enzymes diffuse slowly through the chemostat
³¹² and have a patchy distribution can favor the coexistence of cooperators and cheaters (Allison, 2005).
³¹³ Indeed, previous empirical bioremediation studies have reported coexistence of cooperators with cheaters
³¹⁴ (Ellis et al., 2007; O'Brien et al., 2014). Theoretically, this may be due to a difference of resistance levels
³¹⁵ between cooperators and cheaters as we show here, but a simpler explanation would be the presence of
³¹⁶ spatial gradients. Relaxing the assumption of a perfectly well-mixed chemostat would make the persis-
³¹⁷ tence of cooperators easier. It may also increase the public benefit of toxin resistance, which we have
³¹⁸ considered to be private here (Rojo-Molinero et al., 2019).

³¹⁹ Finally, there may be other ways of periodically introducing cooperators. Experimentally, our constant
³²⁰ inoculation probabilities represent a situation where stock strains of cooperators would be manually added
³²¹ into the chemostat. If instead, multiple chemostats are running in parallel, another way of introducing
³²² cooperators would be to exchange certain amounts of fluids between chemostats. Ecologically, this would
³²³ correspond to migration among patches, and the optimal migration probabilities would depend on the
³²⁴ probability distribution of the different strategies in each chemostat. Comparing the optimal introduction
³²⁵ probabilities and the cumulative efficiency of detoxification between the model presented here and a multi-

chemostat system is left for future work.

In summary, we have combined an ecological model with evolutionary game theory to develop a protocol for the control and optimization of a bioremediation system by microbes, and guard it against collapse through the emergence of cheaters. More broadly speaking, our scenario motivates the integration of important elements from ecological models, such as population densities and environmental feedback, into evolutionary game theory. In essence, our model can be adapted to any practical applications involving costly microbial traits, where manipulating environmental conditions can be used to control evolutionary dynamics. Achieving this will allow us to better anticipate evolutionary change in microbial systems that we strive to control, whether this involves increasing toxin degradation as we have shown here, the production of public goods such as useful chemicals, or eliminating antibiotic resistant pathogens.

5 Data availability

Python code and csv files used for simulations and analytic calculations are available at GitHub, but we cannot show the url for double-blinding.

References

Akita, T. and Kamo, M. Theoretical lessons for increasing algal biofuel: Evolution of oil accumulation to avert carbon starvation in microalgae. *Journal of Theoretical Biology*, 380:183–191, 2015. ISSN 10958541. doi: 10.1016/j.jtbi.2015.05.027. URL http://dx.doi.org/10.1016/j.jtbi.2015.05.027.

Allen, R. C., McNally, L., Popat, R., and Brown, S. P. Quorum sensing protects bacterial co-operation from exploitation by cheats. *The ISME journal*, 10(JANUARY):1–11, 2016. ISSN 1751-7370. doi: 10.1038/ismej.2015.232. URL http://www.ncbi.nlm.nih.gov/pubmed/26744811.

Allison, S. D. Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. *Ecology Letters*, 8(6):626–635, 2005. ISSN 1461023X. doi: 10.1111/j.1461-0248.2005.00756.x.

Andersson, D. I. and Hughes, D. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nature Reviews Microbiology*, 8(4):260–271, 4 2010. ISSN 1740-1526. doi: 10.1038/nrmicro2319. URL http://dx.doi.org/10.1038/nrmicro2319http://www.nature.com/articles/nrmicro2319.

Andersson, D. I. and Levin, B. R. The biological cost of antibiotic resistance. *Current Opinion in Microbiology*, 2(5):489–493, 10 1999. ISSN 13695274. doi: 10.1016/S1369-5274(99)00005-3. URL http://linkinghub.elsevier.com/retrieve/pii/S1369527499000053.

Antoni, D., Zverlov, V. V., and Schwarz, W. H. Biofuels from microbes. *Applied Microbiology and Biotechnology*, 77(1):23–35, 10 2007. ISSN 0175-7598. doi: 10.1007/s00253-007-1163-x. URL http://link.springer.com/10.1007/s00253-007-1163-x.

Atashgahi, S., Sánchez-Andrea, I., Heipieper, H. J., van der Meer, J. R., Stams, A. J. M., and Smidt, H. Prospects for harnessing biocide resistance for bioremediation and detoxification. *Science*, 360(6390):743–746, 5 2018. ISSN 0036-8075. doi: 10.1126/science.aar3778. URL http://dx.doi.org/10.1016/B978-1-4377-2242-0/00034-1http://www.sciencemag.org/lookup/doi/10.1126/science.aar3778.

Bertrand, J.-C., Doumenq, P., Guyoneaud, R., Marrot, B., Martin-Laurent, F., Matheron, R., Moulin, P., and Soulas, G. Applied Microbial Ecology and Bioremediation. In Bertrand, J.-c., Caumette, P., Lebaron, P., Matheron, R., Normand, P., and Sime-Ngando, T., editors, *Environmental Microbiology: Fundamentals and Applications*, chapter 16, pages 659–753. Springer Netherlands, Dordrecht, 1st edition, 2015. ISBN 9789401791182. doi: 10.1007/978-94-017-9118-2.

Blair, J. M., Webber, M. A., Baylay, A. J., Ogbolu, D. O., and Piddock, L. J. Molecular mechanisms of antibiotic resistance. *Nature Reviews Microbiology*, 13(1):42–51, 2015. ISSN 17401534. doi: 10.1038/nrmicro3380.

Bottery, M. J., Wood, A. J., and Brockhurst, M. A. Selective Conditions for a Multidrug Resistance Plasmid Depend on the Sociality of Antibiotic Resistance. *Antimicrobial Agents and Chemotherapy*, 60(4):2524–2527, 4 2016. ISSN 0066-4804. doi: 10.1128/AAC.02441-15. URL http://aac.asm.org/lookup/doi/10.1128/AAC.02441-15.

Broom, M., Pattni, K., and Rychtář, J. Generalized Social Dilemmas: The Evolution of Cooperation in Populations with Variable Group Size. *Bulletin of Mathematical Biology*, pages 1–32, 12 2018. ISSN 0092-8240. doi: 10.1007/s11538-018-00545-1. URL http://link.springer.com/10.1007/s11538-018-00545-1.

Bull, J. J. and Barrick, J. E. Arresting Evolution. *Trends in Genetics*, 33(12):910–920, 2017. ISSN 13624555. doi: 10.1016/j.tig.2017.09.008. URL http://dx.doi.org/10.1016/j.tig.2017.09.008.

Chou, T.-c. Theoretical Basis, Experimental Design, and Computerized Simulation of Synergism and Antagonism in Drug Combination Studies. *Pharmacological Reviews*, 58(3):621–681, 9 2006. ISSN 0031-6997. doi: 10.1124/pr.58.3.10. URL http://pharmrev.aspetjournals.org/lookup/doi/10.1124/pr.58.3.10.

Cressman, R. and Tao, Y. The replicator equation and other game dynamics. *Proceedings of the National Academy of Sciences*, 111(Supplement_3):10810–10817, 2014. ISSN 0027-8424. doi: 10.1073/pnas.1400823111. URL http://www.pnas.org/cgi/doi/10.1073/pnas.1400823111.

Dixit, R., Wasiullah, Malaviya, D., Pandiyan, K., Singh, U., Sahu, A., Shukla, R., Singh, B., Rai, J., Sharma, P., Lade, H., and Paul, D. Bioremediation of Heavy Metals from Soil and Aquatic Environment: An Overview of Principles and Criteria of Fundamental Processes. *Sustainability*, 7(2):2189–2212, 2 2015. ISSN 2071-1050. doi: 10.3390/su7022189. URL http://www.mdpi.com/2071-1050/7/2/2189.

Ellis, R. J., Lilley, A. K., Lacey, S. J., Murrell, D., and Godfray, H. C. J. Frequency-dependent advantages of plasmid carriage by Pseudomonas in homogeneous and spatially structured environments. *ISME Journal*, 1(1):92–95, 2007. ISSN 17517362. doi: 10.1038/ismej.2007.11.

Giri, S., Shitut, S., and Kost, C. Harnessing ecological and evolutionary principles to guide the design of microbial production consortia. *Current Opinion in Biotechnology*, 62:228–238, 2020. ISSN 18790429. doi: 10.1016/j.copbio.2019.12.012. URL https://doi.org/10.1016/j.copbio.2019.12.012.

Gokhale, C. S. and Hauert, C. Eco-evolutionary dynamics of social dilemmas. *Theoretical Population Biology*, 111:28–42, 2016. ISSN 10960325. doi: 10.1016/j.tpb.2016.05.005. URL http://dx.doi.org/10.1016/j.tpb.2016.05.005.

Gong, L., Gao, J., and Cao, M. Evolutionary Game Dynamics for Two Interacting Populations in A Co-evolving Environment. In *2018 IEEE Conference on Decision and Control (CDC)*, pages 3535–3540. IEEE, 12 2018. ISBN 978-1-5386-1395-5. doi: 10.1109/CDC.2018.8619801. URL http://arxiv.org/abs/1806.03194https://ieeexplore.ieee.org/document/8619801/.

Griffin, A. S., West, S. a., and Buckling, A. Cooperation and competition in pathogenic bacteria. *Nature*, 430(7003):1024–7, 8 2004. ISSN 1476-4687. doi: 10.1038/nature02744. URL http://www.ncbi.nlm.nih.gov/pubmed/15329720.

Hardin, G. The Tradegy of the Commons. *Science*, 162(June):1243–1248, 1968. ISSN 0036-8075. doi: 10.1126/science.162.3859.1243.

Hauert, C., Holmes, M., and Doebeli, M. Evolutionary games and population dynamics: maintenance of cooperation in public goods games. *Proceedings of the Royal Society B: Biological Sciences*, 273(1600):2565–2571, 10 2006. ISSN 0962-8452. doi: 10.1098/rspb.2006.3600. URL http://rspb.royalsocietypublishing.org/cgi/doi/10.1098/rspb.2006.3717.

Hauert, C., Wakano, J. Y., and Doebeli, M. Ecological public goods games: Cooperation and bifurcation. *Theoretical Population Biology*, 73(2):257–263, 2008. ISSN 00405809. doi: 10.1016/j.tpb.2007.11.007.

Hummert, S., Bohl, K., Basanta, D., Deutsch, A., Werner, S., Theißen, G., Schroeter, A., and Schuster, S. Evolutionary game theory: cells as players. *Mol. BioSyst.*, 10(12):3044–3065, 2014. ISSN 1742-206X. doi: 10.1039/C3MB70602H. URL http://xlink.rsc.org/?DOI=C3MB70602H.

Kang, C.-h., Kwon, Y.-j., and So, J.-s. Bioremediation of heavy metals by using bacterial mixtures. *Ecological Engineering*, 89:64–69, 4 2016. ISSN 09258574. doi: 10.1016/j.ecoleng.2016.01.023. URL http://dx.doi.org/10.1016/j.ecoleng.2016.01.023https://linkinghub.elsevier.com/retrieve/pii/S0925857416300234.

Kumar, P. K. R., Maschke, H., Friehs, K., and Schügerl, K. Strategies for Improving Plasmid Stability in Bioreactors. *Trends in Biotechnology*, 9(8):279–284, 1991.

Maynard Smith, J. and Price, G. R. The Logic of Animal Conflict. *Nature*, 246(5427):15–18, 11 1973. ISSN 0028-836. doi: 10.1038/246015a0.

Moreno-Fenoll, C., Cavaliere, M., Martínez-García, E., and Poyatos, J. F. Eco-evolutionary feedbacks can rescue cooperation in microbial populations. *Scientific Reports*, 7(1):42561, 3 2017. ISSN 2045-2322. doi: 10.1038/srep42561. URL http://dx.doi.org/10.1038/srep42561http://www.nature.com/articles/srep42561.

O'Brien, S. and Buckling, A. The sociality of bioremediation: Hijacking the social lives of microbial populations to clean up heavy metal contamination. *EMBO reports*, 16(10):1241–1245, 2015. ISSN 1469-221X. doi: 10.15252/embr.201541064.

O'Brien, S., Hodgson, D. J., and Buckling, A. Social evolution of toxic metal bioremediation in Pseudomonas aeruginosa. *Proceedings of the Royal Society B: Biological Sciences*, 281(1787):20140858–20140858, 6 2014. ISSN 0962-8452. doi: 10.1098/rspb.2014.0858. URL http://rspb.royalsocietypublishing.org/content/281/1787/20140858http://rspb.royalsocietypublishing.org/cgi/doi/10.1098/rspb.2014.0858.

Quin, M. B. and Schmidt-Dannert, C. Designer microbes for biosynthesis. *Current Opinion in Biotechnology*, 29(1):55–61, 2014. ISSN 18790429. doi: 10.1016/j.copbio.2014.02.014. URL http://dx.doi.org/10.1016/j.copbio.2014.02.014.

Rojo-Molinero, E., Macià, M. D., and Oliver, A. Social Behavior of Antibiotic Resistant Mutants Within Pseudomonas aeruginosa Biofilm Communities. *Frontiers in Microbiology*, 10(MAR):1–11, 3 2019. ISSN 1664-302X. doi: 10.3389/fmicb.2019.00570. URL https://www.frontiersin.org/article/10.3389/fmicb.2019.00570/full.

Röling, W. F. M., Milner, M. G., Jones, D. M., Lee, K., Daniel, F., Swannell, R. J. P., and Head, I. M. Robust Hydrocarbon Degradation and Dynamics of Bacterial Communities during Nutrient-Enhanced Oil Spill Bioremediation. *Applied and Environmental Microbiology*, 68(11):5537–5548, 11 2002. ISSN 0099-2240. doi: 10.1128/AEM.68.11.5537-5548.2002. URL http://aem.asm.org/cgi/doi/10.1128/AEM.68.11.5537-5548.2002.

Rugbjerg, P., Myling-Petersen, N., Porse, A., Sarup-Lytzen, K., and Sommer, M. O. Diverse genetic error modes constrain large-scale bio-based production. *Nature Communications*, 9(1), 2018a. ISSN 20411723. doi: 10.1038/s41467-018-03232-w. URL http://dx.doi.org/10.1038/s41467-018-03232-w.

Rugbjerg, P., Sarup-Lytzen, K., Nagy, M., and Sommer, M. O. A. Synthetic addiction extends the productive life time of engineered Escherichia coli populations . *Proceedings of the National Academy of Sciences*, 115(10):201718622, 2018b. ISSN 0027-8424. doi: 10.1073/pnas.1718622115.

Ryan Georgianna, D. and Mayfield, S. P. Exploiting diversity and synthetic biology for the production of algal biofuels. *Nature*, 488(7411):329–335, 2012. ISSN 00280836. doi: 10.1038/nature11479.

Sampah, M. E. S., Shen, L., Jilek, B. L., and Siliciano, R. F. Dose-response curve slope is a missing dimension in the analysis of HIV-1 drug resistance. *Proceedings of the National Academy of Sciences*, 108(18):7613–7618, 2011. ISSN 0027-8424. doi: 10.1073/pnas.1018360108. URL http://www.pnas.org/cgi/doi/10.1073/pnas.1018360108.

Samuelson, P. A. The Pure Theory of Public Expenditure. *The Review of Economics and Statistics*, 36(4):387, 11 1954. ISSN 00346535. doi: 10.2307/1925895. URL https://www.jstor.org/stable/1925895?origin=crossref.

San Millan, A., Peña-Miller, R., Toll-Riera, M., Halbert, Z. V., McLean, A. R., Cooper, B. S., and Maclean, R. C. Positive selection and compensatory adaptation interact to stabilize non-transmissible plasmids. *Nature Communications*, 5:1–11, 2014. ISSN 20411723. doi: 10.1038/ncomms6208.

San Millan, A. and MacLean, R. C. Fitness Costs of Plasmids: a Limit to Plasmid Transmission. *Microbiology Spectrum*, 5(5), 9 2017. doi: 10.1128/microbiolspec.MTBP-0016-2017. URL http://www.asmscience.org/content/journal/microbiolspec/10.1128/microbiolspec.MTBP-0016-2017.

Sanchez, A. and Gore, J. Feedback between Population and Evolutionary Dynamics Determines the Fate of Social Microbial Populations. *PLoS Biology*, 11(4), 2013. ISSN 15449173. doi: 10.1371/journal.pbio.1001547.

Schuster, S., Kreft, J.-U., Brenner, N., Wessely, F., Theißen, G., Ruppin, E., and Schroeter, A. Cooperation and cheating in microbial exoenzyme production - Theoretical analysis for biotechnological applications. *Biotechnology Journal*, 5(7):751–758, 7 2010. ISSN 18606768. doi: 10.1002/biot.200900303. URL http://doi.wiley.com/10.1002/biot.200900303.

Smith, P. and Schuster, M. Public goods and cheating in microbes. *Current Biology*, 29(11):R442–R447, 6 2019. ISSN 09609822. doi: 10.1016/j.cub.2019.03.001. URL https://linkinghub.elsevier.com/retrieve/pii/S0960982219302660.

Smith, V. H., Graham, D. W., and Cleland, D. D. Application of resource-ratio theory to hydrocarbon biodegradation. *Environmental Science and Technology*, 32(21):3386–3395, 1998. ISSN 0013936X. doi: 10.1021/es9805019.

Tacconelli, E., Carrara, E., Savoldi, A., Harbarth, S., Mendelson, M., Monnet, D. L., Pulcini, C., Kahlmeter, G., Kluytmans, J., Carmeli, Y., Ouellette, M., Outterson, K., Patel, J., Cavaleri, M., Cox, E. M., Houchens, C. R., Grayson, M. L., Hansen, P., Singh, N., Theuretzbacher, U., Magrini, N., Aboderin, A. O., Al-Abri, S. S., Awang Jalil, N., Benzonana, N., Bhattacharya, S., Brink, A. J., Burkert, F. R., Cars, O., Cornaglia, G., Dyar, O. J., Friedrich, A. W., Gales, A. C., Gandra, S., Giske, C. G., Goff, D. A., Goossens, H., Gottlieb, T., Guzman Blanco, M., Hryniewicz, W., Kattula, D., Jinks, T., Kanj, S. S., Kerr, L., Kieny, M. P., Kim, Y. S., Kozlov, R. S., Labarca, J., Laxminarayan, R., Leder, K., Leibovici, L., Levy-Hara, G., Littman, J., Malhotra-Kumar, S., Manchanda, V., Moja, L., Ndoye, B., Pan, A., Paterson, D. L., Paul, M., Qiu, H., Ramon-Pardo, P., Rodríguez-Baño, J., Sanguinetti, M., Sengupta, S., Sharland, M., Si-Mehand, M., Silver, L. L., Song, W., Steinbakk, M., Thomsen, J., Thwaites, G. E., van der Meer, J. W., Van Kinh, N., Vega, S., Villegas, M. V., Wechsler-Fördös, A., Wertheim, H. F. L., Wesangula, E., Woodford, N., Yilmaz, F. O., and Zorzet, A. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *The Lancet Infectious Diseases*, 18(3):318–327, 2018. ISSN 14744457. doi: 10.1016/S1473-3099(17)30753-3.

Tilman, A. R., Plotkin, J. B., and Akçay, E. Evolutionary games with environmental feedbacks. *Nature Communications*, 11(1):915, 12 2020. ISSN 2041-1723. doi: 10.1038/s41467-020-14531-6. URL http://www.nature.com/articles/s41467-020-14531-6.

Weitz, J. S., Eksin, C., Paarporn, K., Brown, S. P., and Ratcliff, W. C. An oscillating tragedy of the commons in replicator dynamics with game-environment feedback. *Proceedings of the National Academy of Sciences*, 113(47):E7518–E7525, 2016. ISSN 0027-8424. doi: 10.1073/pnas.1604096113. URL http://www.pnas.org/lookup/doi/10.1073/pnas.1604096113.

West, S. A., Diggle, S. P., Buckling, A., Gardner, A., and Griffin, A. S. The Social Lives of Microbes. *Annual Review of Ecology, Evolution, and Systematics*, 38(1):53–77, 12 2007a. ISSN 1543-592X. doi: 10.1146/annurev.ecolsys.38.091206.095740. URL http://www.annualreviews.org/doi/10.1146/annurev.ecolsys.38.091206.095740.

West, S. A., Griffin, A. S., and Gardner, A. Evolutionary Explanations for Cooperation. *Current Biology*, 17(16):R661–R672, 8 2007b. ISSN 09609822. doi: 10.1016/j.cub.2007.06.004. URL http://linkinghub.elsevier.com/retrieve/pii/S0960982207014996.

Wolfe, B. E. and Dutton, R. J. Fermented foods as experimentally tractable microbial ecosystems. *Cell*,

161(1):49–55, 2015. ISSN 10974172. doi: 10.1016/j.cell.2015.02.034. URL http://dx.doi.org/10.1016/j.cell.2015.02.034.

Yurtsev, E. A., Chao, H. X., Datta, M. S., Artemova, T., and Gore, J. Bacterial cheating drives the population dynamics of cooperative antibiotic resistance plasmids. *Molecular Systems Biology*, 9(683): 1–7, 2013. ISSN 17444292. doi: 10.1038/msb.2013.39. URL http://dx.doi.org/10.1038/msb.2013.39.

Zaccaria, M., Dawson, W., Cristiglio, V., Reverberi, M., Ratcliff, L. E., Nakajima, T., Genovese, L., and Momeni, B. Designing a bioremediator: mechanistic models guide cellular and molecular specialization. *Current Opinion in Biotechnology*, 62:98–105, 2020. ISSN 18790429. doi: 10.1016/j.copbio.2019.09.006. URL https://doi.org/10.1016/j.copbio.2019.09.006.