Morphological and molecular evidence for range extension and first

occurrence of the Japanese seahorse, *Hippocampus mohnikei* (Bleeker 1853)

in a bay-estuarine system of Goa, central west coast of India

Sushant Sanaye[1], Rakhee Khandeparker[1], Rayadurga Anantha Sreepada[1*], Mamatha

Singanahalli Shivaramu[1], Harshada Kankonkar[1], Jayu Narvekar[2], Mukund

Gauthankar[1]

[1] Aquaculture Laboratory, Biological Oceanography Division, CSIR-National Institute of

Oceanography, Dona Paula, Goa–403 004, India

[2] Physical Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula, Goa–403

004, India

* Corresponding author

E-mail: sreepada@nio.org (RAS)

Abstract

Accurate information of taxonomy and geographic range of seahorse species (genus *Hippocampus*) is the first step in preparing threat assessments and designing effective conservation measures. Here, we report the range expansion and first occurrence of the Japanese seahorse, *Hippocampus mohnikei* (Bleeker, 1853) from the Mandovi estuarine ecosystem of Goa, central west coast of India (CWCI) based on morpho-molecular analyses. The morphometric and meristic traits particularly, short snout (29–35% head length), double cheek spine, low coronet and long tail (51.2–57.9% of standard length), 11 trunk rings, 37–39 tail rings, 15–16 dorsal and 12–14 fin rays observed in four collected specimens matched with the reported key diagnostic morphological criteria of vouchered specimens of *H. mohnikei*. The seahorse mitochondrial cytochrome oxidase subunit I (COI) and cytochrome *b* (Cyt *b*) genes were partially sequenced for conclusive genetic identification of the species under study. Molecular analysis showed that all four individuals of seahorse species clustered together suggesting a monophyletic lineage. Using the maximum similarity with GenBank database, maximum likelihood network and subsequent morphological analysis, the identity of the collected seahorse species was reconfirmed as *H. mohnikei*. With this new report, the geographic range of *H. mohnikei* extended significantly to the westward side from its previously known range. This new sighting of *H. mohnikei* could indicate a long-distance dispersal facilitated by due to prevailing oceanic circulation in the Indo-Pacific region or increased habitat suitability in bay-estuarine systems of Goa, CWCI. Comparison of the pair-wise genetic distances (Kimura 2-parameter) based on COI and Cyt *b* sequences revealed that the specimens examined in this study are genetically closer to *H. mohnikei* populations from Vietnam and Thailand than they are to those in Japan and China. To test the hypothesis

50   whether *H. mohnikei* are vagrants or previously unreported established population, long-term

51   inter-annual sampling and analyses are warranted.

Introduction

Seahorses (Syngnathiformes; Syngnathidae) belonging to the single genus, *Hippocampus*

(Rafinesque, 1810) are a fascinating and remarkable group of fishes with their unusual body

shape and their biology, with males incubating the fertilized eggs in a brood pouch [1]. They

are small, cryptic, and sedentary marine fishes that occur worldwide in shallow temperate to

tropical waters in a wide variety of habitats, including seagrass beds, estuaries, coral and

rocky reefs and mangroves [2−4]. Their remarkable ability to camouflage with structurally

complex habitats further reduces the risk of predation [5], while providing stealth for prey

capture [6]. However, these biological traits often also make them unnoticeable from coastal

habitat. Due to their cryptic nature, their ability to camouflage and the sparse distribution,

seahorses become almost unnoticeable and thus restricting the biologists for their

identification in dense coastal marine habitats.

The primary step in preparing threat assessments and designing effective conservation of

seahorse populations is highly dependent on the precise identification of individual species in

and beyond known geographic distributions [7]. Out of 44 recognized seahorse species

(Genus *Hippocampus*) throughout the world [8], *H. kelloggi*, *H. kuda*, *H. spinosissimus* and

H. trimaculatus are common in Indian coastal marine waters [9,10]. In addition, sightings

based on morphological identification of single specimens of *H. mohnikei* from the Palk Bay,

southeastern India [11], *H. borboniensis* [12] and *H. montebelloensis* [13] from Gulf of

Mannar, southeast coast of India and *H. camelopardalis* from Gulf of Kachchh, northwest

coast of India [14] have been reported. Another seahorse species, *H. histrix* is also suspected

to occur in Indian coastal waters [15]. The validity of *H. bornoinensis* and *H. montebelloensis*

as separate species however, is currently under revision due to their synonymity with *H. kuda*

and *H. zebra,* respectively [16].

The utility of morphological traits commonly used in diagnosis of *Hippocampus* spp. is

quite challenging as they lack certain key physical features (e.g. pelvic and caudal fins), and

have high variability of overlapping body proportions, color (camouflage) and skin filaments,

fin-ray and trunk-ring numbers may vary greatly within any given-species and between

species [3,17]. On the other hand, molecular methods have proved helpful in resolving

morphologically challenging seahorse taxonomy [16,18], phylogenetic relationships within

genus, *Hippocampus* [19,20], natural species boundaries [21−23] and genetic variability

[22,24] of many seahorse species. An integrated approach of combination of morphological

and genetic analyses [25] would aid in the management of demographically separate

populations as independent units and allow international legal mechanisms and international

agreements such as the convention on international trade in endangered species (CITES) to

work effectively [26].

Generally, seahorses are sedentary fish species with small home range and patchy

distribution in sheltered areas such as seagrasses and seaweeds [2,27,28]. However, instances

of limited migrations by seahorses in search of proper habitat, food or holdfasts have been

reported [28,29]. Evidences of long-distance dispersal of seahorse species via rafting

(floating debris and seaweeds) [2,12,30] and subsequent colonization [21] have also been

documented. The role of oceanic currents facilitating in long-distance dispersal and range

expansion of many seahorse species has also been highlighted [31−36].

The Japanese seahorse or Lemur-tail seahorse, *Hippocampus mohnikei* (Bleeker, 1853) is

a small coastal seahorse species (5−8 cm) inhabiting seagrasses, mangroves, oyster beds and

mussel farms [37]. Original records indicated that the distribution of *H. mohnikei* is restricted

or native to the coastal waters around Japan and Vietnam [17,38−40]. However, new records

of *H. mohnikei* from Cambodia, Malaysia, Thailand and Vietnam [37], along with recently

published studies from southeastern India [11] and Singapore [41], have greatly expanded the

known geographical range of *H. mohnikei* within Southeast Asia. As a consequence of

suspected reduction in population size to the tune of >30% over the past 10 years due to

fisheries exploitation [37,42−44], the status of *H. mohnikei* in the IUCN Red List of

Threatened Species has been listed as 'VULNERABLE' [45].

A probable occurrence of *H. mohnikei* along the central west coast of India (CWCI)

began in February 2017 when a fisherman posted a picture of seahorse incidentally caught in

gill net operated in the Chapora estuary, Goa (India) (15.6120° N, 73.7506° E) on social

media [S1 Fig] which was suspected to be an adult male of *H. mohnikei*. Sustained follow-up

surveys for its occurrence in the surrounding environs of the Chapora estuary did not yield in

any further reporting. Subsequently, four sub-adult specimens of *H. mohnikei*—hitherto

sighted only from southeast coast of India [11], caught as an incidental catch in bag net

fisheries during July, 2018 in the Mandovi estuarine system, Goa,CWCI have been examined

here.

This contribution describes the range expansion and first occurrence of *H. mohnikei* from

the western coastal India. To test further the hypothesis that the morphologically identified

seahorse species is *H. mohnikei*, two mitochondrial DNA gene regions (loci), cytochrome

oxidase subunit I (COI) and cytochrome *b* (Cyt *b*) were partially sequenced for

reconfirmation. Because of its rapid evolutionary rate and almost complete maternal

inheritance, mitochondrial DNA (mtDNA) has been widely used to identify both population

structure and genetic variability [46]. Within mtDNA, the COI and Cyt *b* genes are widely

used for constructing phylogeny of many taxa including inter-specific and intra-specific

variation in seahorse species owing to their slowest mutation and lower substitution rates

compared to other mtDNA genes [16,19,25]. Potential role of ocean currents on the range

expansion of *H. mohnikei* to north-western Asia have also been discussed. In the wake of the

vulnerability of seahorse populations to the threats such as habitat alteration/destruction and

fishing pressure globally, the present sighting of *H. mohnikei* in a bay-estuarine system of

CWCI is of considerable conservation and biogeographic significance.

Materials and Methods

Collection site and seahorses

The locality (Brittona, Goa, India, 15.3059°N, 73.5073°E, Fig 1) at which the examined

seahorse specimens were collected is a shallow water littoral environment in the proximity to

the mangrove-dominated Chorao Island in the Mandovi estuary, CWCI. The currents are

largely tide dominated, the tides being semi-diurnal with a mean amplitude of 1.3 m. Currents

are also influenced by the large seasonal freshwater influx [47] during the summer monsoon

season (June to September). Temporal variations in abiotic and biotic factors influenced by

monsoonal precipitation and seasonal upwelling render the collection location as one of the

most ecologically complex ecosystems [48,49]. Significant lowering of salinity occurs when

the freshwater discharge is at its maximum during monsoon. During the post- and pre-

monsoon periods, the salinity at the surface and bottom remains high (~25 ppt). The bottom

topography at the collection site is muddy and comprises mainly of silt and clay with rich

organic matter content [50]. The collection site is reported to harbor rich and diverse fishery

resources and its importance in supporting the life history stages of several marine teleosts

has also been well-documented [51].

Present report is based on four specimens of *H. mohnikei* (females, MK1 and MK2;

males, MK3 and MK4; Fig. 2. and Fig. 3) that were incidentally caught in bag nets attached

to the fixed stakes (total length, 25 m; mouth length, 5 m; cod end mesh size, 10 mm)

operated at the water depth of ~6 m in the Mandovi estuary during July 2018. Dead seahorses

landed as incidental catch in bag nets attached to the stakes operated by local fishermen in the

estuarine system formed the study material. No live seahorses were specifically caught for the

present study. Fin clips from these dead seahorses were used for molecular analysis. Since,

the present study did not involve any experimentation with live specimens, the approval of

the ethics committee for the usage of live animals for experiments is therefore

inconsequential.

Morphological analysis

Before subjecting them to a detailed analysis of morphological characteristics, seahorse

specimens were carefully photographed (Nikon D7200, Japan) and sexed via observation of

presence or absence of brood pouch. Morphometric measurements and meristic counts were

carried out following standardized protocols [15,17,52]. Morphometric characters such as

height (Ht, linear distance from the top of the head to the tip of the stretched tail), head length

(HL), trunk length (TrL), tail length (TaL) and standard length (SL = HL+TrL+TaL), were

measured. Measurements of snout length (SnL), ratio of head/snout lengths (HL/SnL), snout

depth (SnD), head depth (HD), pectoral fin base length (PL) and dorsal fin base length (DL)

were also conducted. All measurements were taken with the help of digital caliper to 0.1 mm

and repeated to ensure accuracy. Trunk depths on 4th (TD4) and 9th (TD9) trunk rings were

measured. The presence/absence of spines and dermal appendages was also noted. Meristic

characters such as the number of trunk rings (TrR), tail rings (TaR), dorsal fin rays (DF),

pectoral fin rays (PF) and anal fin rays (AF) were counted under stereo zoom microscope

(Olympus SZX7, Japan) and the values were confirmed by triplicate counting. Brood pouch

dimensions were measured in all male specimens.

Morphological identification of the collected seahorse specimens from Goa was based on

comparison of the salient morphometric and meristic characters with the reported diagnostic

features for different seahorse species [15,17,25]. For further confirmation, the morphological

data of Goa specimens was compared with all vouchered specimens of *H. mohnikei*

[11,15,25,53,54]. The voucher specimen collected from Goa is deposited in the national

marine biodiversity repository at CSIR-National Institute of Oceanography, Goa, India

(deposition ID: NIO1015/19).

Statistical analysis

Morphological data of *H. mohnikei* and two other commonly occurring seahorse species (*H.*

kuda and *H. trimaculatus*) in the coastal waters of India was evaluated by one-way analysis

of variance (ANOVA) followed by the Duncans multiple-range test [55]. The level of

significance was tested at 5.0%, represented as $P < 0.05$. A comparison of morphological

characters and meristic features of *H. mohnikei* (n = 4) with *H. kuda* (n = 10) and *H.*

trimaculatus (n = 10) was made through principal component analysis (PCA). PCA was

employed in which factor loadings based on components were used to determine the

morphometric factors using PRIMER software (version 6) after square-root transformation.

Molecular analysis: DNA extraction, PCR amplification and sequencing

For gene-based phylogenetic analysis, fragments of dorsal fin from freshly landed dead

seahorses were removed and preserved in 95% ethanol at −20°C until subjected to DNA

extraction. DNA extraction was carried out as described by Kumar et al. with a few

modifications [56]. In brief, lysis buffer and sodium dodecyl sulfate were added to the fin

tissue and vortexed. Further, RNase and Proteinase K were added to the mixture and

incubated for 15–20 minutes at room temperature. After the incubation period, the DNA was

extracted using phenol:chloroform:isoamyl alcohol (25:24:1) mixture. Precipitated DNA was

washed with 70% cold ethanol. The gel electrophoresis was performed to confirm the quality

and integrity of DNA.

Seahorse mitochondrial cytochrome oxidase subunit I (COI) gene was amplified using

primers,   forward   (5'TCAACTAATCACAAAGACATCGGCAC3′)   and   reverse

(5'ACTTCGGGGTGCCCAAAGAATC3′) [57]. Similarly, mitochondrial cytochrome *b* (Cyt

b) gene fragment was also amplified using PCR with seahorse-specific primers, forward

(5'AACYAGGACYAATGRCTTGA3') and reverse (5′GCASWAGGGAGGRKTTTAAC3′)

[43]. Reactions were carried out in a thermal cycler (Veriti® 96-well Thermal Cycler,

Applied Biosystems, CA, USA). The amplification of COI and Cyt *b* genes was carried out

by following cycling conditions as described by Lourie et al. with few modifications; an

initial denaturation step of 95°C for 2 minutes followed by 39 cycles of 94°C for 30s, 55°C

for 30s 72°C for 1 minute and a final extension at 72°C for 2 minutes [58]. Amplified PCR

products of both COI and Cyt *b* genes were verified on agarose gel (1.5%) and purified for

sequencing using PCR purification kit (Sigma Aldrich, USA). Gene sequence of the PCR

products was determined using a taq dye deoxy terminator cycle sequencing kit

(PerkinElmer, CA, USA) following supplier's protocols. Sequencing reaction products were

analyzed on 373 automated DNA Sequencer (Model 313 xl; Applied Biosystems, CA, USA).

Amplification of the mitochondrial COI and Cyt *b* genes resulted in fragments of 609 bp and

639 bp, respectively.

Phylogenetic analysis

Nucleotide sequences generated for four specimens collected from Goa (MK1, MK2, MK3

and MK4) were deposited in GenBank under accession numbers MN595216, MN595218,

MN595217 and MK330041.1 for COI and MN595213, MN595214, MN595215 and

MK112274.2 for Cyt *b* genes, respectively). Sequence integrity and genetic affinity with

known species was compared using the basic local alignment search tool (BLAST) of NCBI

[59] (http://www.ncbi.nih.gov/BLAST). To complete the analysis, sequences of Goa

specimens were aligned with 47 and 85 sequences of COI and and Cyt *b* genes, respectively

of all vouchered specimens of *H. mohnikei* retrieved from the GenBank database (accession

numbers shown in S1 Table). The mitochondrial COI and Cyt *b* gene sequences were aligned

using the ClustalW algorithm implemented in MEGA 7 [60]. To ascertain stable and reliable

relationships among the seahorse species, phylogenetic trees, separately for COI and Cyt *b*

genes were constructed through maximum likelihood (ML) method using K2P distances. The

analyses were performed in MEGA 7, and confidence levels were assessed using 1000

bootstrap replications. The alligator pipefish, *Syngnathoides biaculeatus* was used as an

outgroup species. Genetic distance values are used to estimate intraspecies and interspecies

kinships and also as a basis for phylogeny analysis. The degree of genetic divergence within

H. mohnikei from different geographical locations (S1 Table) was assessed using Kimura 2-

parameter (K2P) distance model [61] implemented in MEGA 7 [60]. Codon positions

included were $1^{st}+2^{nd}+3^{rd}$+noncoding. All ambiguous positions with alignment gaps and

missing data were removed from the sequences (pair-wise deletion option).

Prevailing ocean circulation

In order to understand the basin-wide circulation in the Indo-Pacific region (encompassing

South China Sea, East China Sea, and southeastern Bay of Bengal), monthly mean

climatology was prepared based on daily data on U (zonal) and V (meridional) components

of geostrophic current velocities derived from Aviso merged product for the years, 2007 –

2017. Altimeter products by Ssalto/Duacs and distributed by Aviso+, with support from Cnes

(https://www.aviso.altimetry.fr) were used. The climatological currents generated for the

winter season (December to February) from 2007 to 2017 for the Indo-Pacific region are

depicted in Fig. 4.

Results

Specimens examined

Colour of the head and body of freshly collected seahorse specimens (females, MK1 and

MK2; males, MK3 and MK4; NIO1015/19) was dark brown to dark black. The entire body

was covered with low spines and the coronet was observed to be short with five tiny

projections. Two prominent cheek spines and double rounded spines below the eye (each on

the either side of the head) were discernible in all four specimens. (Fig. 2 and Fig. 3.).

Morphological taxonomy

Measured height (Ht) and standard length (SL) of specimens varied between 39.2 and 54.2

mm (48.2 ± 6.5 mm) and 47.3 and 60.1 mm (54.95 ± 5.48 mm), respectively. TaL was

observed to be relatively long compared to the TrL of the body. The ratio of the head length

(HL) to snout length (SnL) of all four specimens varied between 3.2 and 3.5 (Table 1).

Number of trunk rings (TrR) in all specimens were 11 whereas, the number of tail rings

(TaR) varied narrowly between 37 and 39. The number of TrR supporting dorsal fin was two

and the number of TaR supporting dorsal fin was counted to be one in all specimens. Number

of dorsal fin pectoral fin and anal fin rays recorded were 15–16, 12–14 and 4, respectively.

Slight enlargements, respectively at 1^{st}, 4^{th}, 7^{th} and $10/11^{th}$TrR and 4^{th}, $7/8^{th}$, $9/10^{th}$,

$13/14^{th}$TaR were prominently discernible (Table 1).

Comparative morphology

We compared the morphological features of *H. mohnikei* with other two commonly occurring

seahorse species, *H. kuda* and *H. trimaculatus* occurring in coastal waters of India (Table 2).

Fresh specimens of *H. kuda* was orange to dark brown with reddish spots over the body with

single eye and cheek spine, rounded and backward curled coronet and deeper head and

rounded or less developed body spines [15,16]. *H. trimaculatus* has three prominent black

spots on the dorso-lateral surface of the first, fourth and seventh trunk rings [15,16] which are

clearly absent in *H. mohnikei*. Single eye and hooked like cheek spine also observed in *H.*

trimaculatus. Snout length (SnL) of *H. mohnikei* was shorter compared to the *H. kuda* and *H.*

trimaculatus. These morphological characters clearly distinguish *H. mohnikei* from *H. kuda*

and *H. trimaculatus* (Table 2).

PCA used to elucidate the factors contributing to morphological differences in *H.*

mohnikei, H. kuda and *H. trimaculatus* showed that many morphometric and meristic

characters play an important role in differentiating the seahorse species. Results of PCA

analysis revealed that 11 morphometric and meristic characters integrated into two principal

components (PC1 and PC2) which cumulatively contributed to 97% of total variance (S2

Table). The PC1 which explained 81% of total variation consisted of morphometric

characters such as SL, TaL, TD4, HL, TD9, TrL, HD, SnL and SnD while, the PC2 which

explained 16% of total variation consisted of the TaR and DF. The derived ordination

diagram after plugging standardized morphological data of all individuals of three seahorse

species into PCA is depicted in Fig. 5. PCA ordination diagram showed that three seahorse

species were clustered out distinctly in orientation space. Therefore, based on comparison of

morphological traits [15,16,25] and the results of PCA, the identity of seahorse specimens

collected from Goa was confirmed as *H. mohnikei*.

Morphological measurements revealed that all the four Goa specimens are consistent and

congruent with the salient morphological descriptions for type specimens of *H. mohnikei*

reported by Lourie et al. [15,17,25] for specimens collected from Vietnam and Japan.

Furthermore, meristic, morphometric, and key diagnostic morphological character

comparisons from vouchered specimens of *H. mohnikei* [11,15,25,53,54]*,* reconfirmed the

identity of seahorse specimens collected from Goa as *H. mohnikei* (Table 3).

Phylogenetic analysis and genetic distance

PCR for the mitochondrial COI and Cyt *b* genes of Goa seahorse specimens resulted in 609

bp and 639 bp amplicons, respectively. Maximum likelihood (ML) phylogenetic analysis

were performed using K2P distances with sequences of COI and Cyt *b* genes of seahorse

specimens from Goa (MK1, MK2, MK3 and MK4) with all downloaded sequences of

vouchered *H. mohnikei* from GenBank (S2 Table). ML trees of Goa seahorse specimens

(GenBank accession numbers, MK330041.1, MN595216.1, MN595217.1 and MN595218.1

for COI gene; GenBank accession numbers, MK112274.2, MN595213.1, MN595214.1 and

MN595215.1 for Cyt *b* gene) showed that all four individuals clustered together suggesting a

monophyletic lineage (Fig. 6 and Fig. 7). Furthermore, sequences of COI and Cyt *b* genes of

Goa specimens, respectively showed maximum phylogenetic similarities with *H. mohnikei*

(GenBank accession number: GQ502157.1) and with *H. mohnikei* (GenBank accession

numbers: EU179923.1 and EU179924.1) strongly supported by high bootstrap values of >95%

(Fig. 6 and Fig. 7).

Genetic distance describes the kinship distance between intraspecies and interspecies.

Calculated pair-wise genetic distance (K2P) values based on COI and Cyt *b* sequences of *H.*

mohnikei from Goa and other localities (Japan, Korea, China, Vietnam and Thailand)

revealed that *H. mohnikei* from Goa the show least genetic distances of 0.7% for both COI

and Cyt *b* genes with populations of *H. mohnikei* from Vietnam (COI) and Thailand (Cyt *b*),

respectively (Table 4 and Table 5). In contrast, Goa specimens showed highest genetic

distances of 1% for COI and 7% for Cyt *b* genes with populations of *H. mohnikei* from China

and Japan, respectively.

Ocean circulation

As seen from Fig. 4 that the circulation in the Indo-Pacific region shows a large-scale flow

moving from the East China Sea to the South China Sea. From the South China Sea, the

major part of the flow turns southeastward entering into the Java Sea and a small part turns

northwestward to eastern Bay of Bengal (BoB) and the Andaman Sea through Malacca Strait.

From eastern BoB, the surface waters flow westward as a part of the north equatorial current

(NEC). On encountering the coast of Sri Lanka, a part of the NEC move northward into the

northern BoB, while the rest moves westward and into the southern Arabian Sea, south of Sri

Lanka. The NEC feeds into the wintertime west India coastal current (WICC) and moves

waters from NEC towards north into the Arabian Sea along its eastern boundary.

Discussion

The geographic range of *Hippocampus mohnikei* is still unresolved as Lourie et al [15,16]

confirmed the nativity of this species to Japan and Vietnam and likely extending throughout

the South China Sea and the Gulf of Thailand [16,17,25]. New records on the occurrence of

H. mohnikei from many other non-native locations in north- and south-eastern coastal waters

of Asia [11,37,41,44,57,62] suggest its significant extended distribution in the Indo-Pacific

region (Fig. 8). The sighting of *H. mohnikei* in the Palk Bay, southeast coast of India [11] is

till date the known range limit towards northwestern Asian coastal waters. In southeastern

Asian coastal waters, *H. mohnikei* has been reported from Singapore [41] as well as from

Cambodia, Malaysia, Thailand and Vietnam [37]. In northeast Asia, *H. mohnikei* is reported

from China [44,57] and South Korea [62,63].

The present report on the occurrence of *H. mohnikei* in a bay-estuarine system of Goa,

CWCI as confirmed by both morphological and molecular analysis represents a significant

westward expansion compared to the previously known geographic range of this species [37].

Since the reporting of *H. mohnikei* solely based on morphological characters from the Palk

Bay, southeast coast of India [11], no further sightings of this species have been reported

from coastal marine waters of India. Therefore, a comparison of genetically distinct

populations of *H. mohnikei* inhabiting native (Japan and Vietnam) and extended ranges in the

Indo-Pacific region (Fig. 8) with Goa specimens gains significance. Lourie et al. reported that

a Vietnamese specimen of *H. mohnikei* was genetically distinct from *H. mohnikei* from Japan

[16,25].

Maximum likelihood (ML) tree constructed of Goa specimens with all vouchered

specimens of *H. mohnikei* based on COI gene using K2P distances showed maximum

phylogenetic proximity with specimen from Vietnam (GenBank accession number,

GQ502157.1) (Fig. 6). On the other hand, ML tree constructed with sequences of Cyt *b* gene

of Goa specimens showed maximum genetic similarities with *H. mohnikei* from Thailand

(GenBank accession numbers, EU179923.1 and EU179924.1) (Fig. 7). In both ML trees of

COI and Cyt *b* genes, phylogenetic proximity of all four Goa individuals with >95%

bootstrap support thus confirming the taxonomic identity of Goa specimens as *H. mohnikei*.

Furthermore, a comparison of pair-wise genetic distance values (K2P) based on sequence

COI (Table 5) and Cyt *b* (Table 6) genes showed that Goa specimens showed least genetic

distance with populations of *H. mohnikei* from Vietnam (K2P value, 0.7%) and Thailand

(K2P value, 0.7%), respectively. In contrast, Goa specimens showed highest genetic distances

of 1% for COI and 7% for Cyt *b* genes with populations of *H. mohnikei* from China and

Japan, respectively. Similar high pair-wise K2P values of sequences of Cyt *b* between *H.*

mohnikei populations from the Gulf of Thailand and those from China (13.54−14.46%) and

Japan (14.42−14.79%) have been recently reported [64].

A comparative assessment of K2P values of the present study with those from Vietnam

[65] and Gulf of Thailand [64], respectively for COI and Cyt *b* suggest that *H. mohnikei*

populations from Goa, Vietnam and the Gulf of Thailand might share a similar genetic

identity and genetically distinct of those from China and Japan. Also, a possible genetic

differentiation in populations of *H. mohnikei* inhabiting coastal marine waters of northeastern

Asia (Japan, Korea and China) and southeastern Asia (Vietnam, Thailand, Cambodia,

Malaysia, Singapore and India) may not be ruled out. Furthermore, least genetic divergence

recorded between different populations of *H. mohnikei* (Goa vs Vietnam, COI gene) and

(Goa vs. Thailand, Cyt *b* gene), the possibility of crypticity within *H. mohnikei* populations

from these localities is ruled out by applying 2% threshold genetic divergence for assessing

intra-specific differences [16]. In contrast, recent study by Panithanarak et al observed high

genetic variance between the populations of *H. mohnikei* from the Gulf of Thailand and

China (13.54–14.46%) and Japan (14.42–14.79%) suggested the possibility of cryptic species

within *H. mohnikei* population range [64]. Further studies, however are necessary to assess

the genetic variation within *H. mohnikei* populations in the entire Indo-Pacific region.

Recent studies have advanced our understanding of habitat associations of *H. mohnikei*

compared to what has been described in the IUCN Red List of Assessment [45]. Previously,

H. mohnikei was reported only in association with seagrass beds in Japan and in an estuary in

Vietnam [16,25]. Association of *H. mohnikei* populations with seagrass beds (*Zostera* spp.) in

Korea [62,63], Japan [66] and China [44] has been reported. Whereas the specimen reported

from the Palk Bay, southeastern India [11] was associated with two other additional genera of

seagrass species (*Halophila* and *Thalassia*). Recently, published data on the habitat

preferences of *H. mohnikei* from Southeast Asia [37] indicate its association with seagrass

beds of *Cymodocea* spp., *Enhalus* spp., *Halodule* spp., *Halophila* spp., *Syringodium* spp. and

Thalassia spp. Furthermore, these authors also report the association of *H. mohnikei* with

mangroves, oyster beds, mussel farms and sandy beaches, whereas, Lim reported its

occurrence in association with seaweeds in Singapore [41]. From these studies, it is evident

that *H. mohnikei* occupies a wide variety of habitats.

All previous reports on the habitat descriptions of *H. mohnikei* were located in the coastal

waters and bay-estuarine systems in the depth range of 0–10 m [11,17,25,37,44,62].

Interestingly, the occurrence of small juveniles of *H. mohnikei* in association with highly

dense live prey organisms in the offshore waters of Japan has also been reported [39]. The

present report of *H. mohnikei* is based on specimens caught as an incidental catch in bag nets

fixed to the stakes operated in the Mandovi estuary, Goa. Although precise information on

the habitat of *H. mohnikei*, could not be ascertained, it is worth describing the surrounding

environs at the site of collection. Predominantly, the estuarine stretches of the River Mandovi

have dense mangrove coverage and extensive oyster beds [67] and patches of seagrass,

Halophila ovalis and *H. beccarii* [68]. The channel network of Mandovi estuary is

characterized by islands (Chorao, Divar and Vanxim) and many sheltered areas interspersed

with dense mangrove ecosystem. The present occurrence of *H. mohnikei* in a bay-estuarine

system of Goa is not quite surprising as these are the typical habitats where this species has

been previously reported. For ascertaining habitat shift by geographically disconnected

populations of *H. mohnikei,* further intense studies on habitat specificities of different life

stages therefore, are warranted.

Food availability has been considered as one of the factors influencing the abundance and

distribution of seahorses [20]. Zooplankton such as *Oithona davisae*, Calanoida and *Penilia*

avirostris have been identified as the main prey organisms for juvenile and young *H.*

mohnikei in Tokyo Bay [39], whereas the adults have been reported to prey upon mainly

larger organisms such as gammarid and caprellid amphipods [63]. Nutrient-rich waters of the

Mandovi estuarine system sustain high primary productivity and abundance of zooplanktonic

prey organisms such as copepods [69−71] which are essential for the survival and

establishment of seahorse populations. Considering the prevailing environmental conditions,

mangrove shelter and abundance of prey organisms, the likelihood of establishment and

colonization of seahorse populations in these sheltered areas is quite high.

Factors such as habitat, diet, and even anthropogenic activities are the main reasons cited

for the migration of seahorses [72]. Furthermore, instances of seahorses for long distance

dispersal by means of rafting along with seaweeds or any other suitable holdfast under the

influence of ocean currents during juvenile stages have been documented [2,22]. A southward

migration of *H. histrix* ~1800 km under the influence of East Australian Current in the Great

Barrier Reef has been reported [33]. Furthermore, ~1100 km migration from the south to the

northeast coast of India by *H. fuscus* under the influence of East India Coastal Current has

been also reported [35]. A Direct evidence for passive long-distance migration of *H.*

trimaculatus at Malacca Strait has been documented recently [36].

Newborn juveniles of many seahorse species undergo a pelagic phase (planktonic)

ranging between two and six weeks depending on the species before they settle down into

sessile habitats (benthos) [2,73]. Consequently, planktonic phase juveniles are likely to

facilitate in widespread gene flow across geographically disconnected populations and

resulting in genetic homogeneity [74]. The occurrence of small juveniles of *H. mohnikei*

(15.1−45.5 mm) in association with highly dense live prey organisms in offshore waters of

Japan [39] suggests that *H. mohnikei* might undergo a prolonged planktonic phase.

Furthermore, a seasonal (warm and cold season) inshore-offshore migration by *H. mohnikei*

populations inhabiting Chinese coast has been confirmed [57]. Higher dispersal capabilities

of seahorse juveniles with floating debris, prey swirls and with seaweed rafts than adults has

been reported [2,30,36]. It has been postulated that marine species with greater longevity,

free swimming and feeding (planktotrophic) larval phases have relatively high dispersal

abilities and thus permit genetic exchange between populations [75]. A relatively prolonged

planktonic phase as in case of *H. mohnikei* may entail higher degree of dispersal under the

influence of oceanic circulation.

The confirmed occurrence of four sub-adult specimens (Mandovi estuary) and suspected

occurrence of an adult specimen of *H. mohnikei* in the proximately adjacent bay-estuarine

ecosystem (Chapora estuary) of Goa, CWCI could be a consequence of either long-distance

dispersal from Southeast Asian countries or stepping-stone mediated dispersal as revealed by

oceanic currents circulation in the Indo-Pacific region. It is clearly seen from Fig. 4 that water

mass from southeastern Asian coasts and east coast of India enters into the Arabian Sea as a

result of winter circulation [76]. In the absence of geographical barriers between *H. mohnikei*

populations of Goa and Southeast Asian countries, ocean currents flowing from regions

where sightings of *H. mohnikei* may contribute to wide-spread dispersal. Therefore, it is

plausible that oceanic currents may be aiding long-distance or stepping stone dispersal of *H.*

mohnikei populations from different geographical locations such as Cambodia, Malaysia,

Thailand and Vietnam [15,16,37,64], Singapore [41] and the Palk Bay, southeast coast of

India [11] to the west coast of India.

Conversation with local fishermen who operate stake nets at the collection site revealed

that sightings of seahorses are relatively more common during summer monsoon season than

in non-monsoonal months. During the summer monsoon period, a number of rivulets and

rivers connected to the Mandovi estuarine network discharge large amounts of freshwater

[77]. Consequently, it is possible that the rainwater discharge from sheltered areas—the

preferred seahorse habitats may force seahorses flow downstream along with currents due to

their poor swimming abilities.

It is well known that the west coast of India comes under the influence of substantial

coastal cold-water upwelling event during the summer monsoon season [48,77]. It has been

reported that the survival and dispersal of tropical marine species are known to be impacted

by a drastic decrease in water temperature associated with upwelling [78−80]. Consequently,

seahorse populations in western coastal India might undergo seasonal fluctuations [22]. A

significant decrement in abundance and diversity of demersal fishes during the summer

monsoon in a bay-estuarine system of the Mandovi river has been documented [51]. As a

consequence of the reduction in salinity and drop in temperature associated with monsoonal

precipitation and upwelling, respectively the seahorse populations might face limitations in

growth and recruitment along the west coast of India. Interestingly, unlike most subtropical

seahorse species, the greater tolerance by native populations of *H. mohnikei* to cold water in

the Western Pacific has been reported [31]. However, further temperature stress simulation

studies are required to ascertain whether geographically disconnected populations of *H.*

mohnikei also possess a similar trait or not.

Conclusions

Based on the new information presented in this paper, there is clear evidence that *H. mohniei*

has expanded its natural and reported geographic range. Recognizing the potential of

seahorses as a popular flagship species in estuarine conservation and sustainable management

[1,81,82], significant westward range expansion of *H. mohnikei* is of considerable

biogeographic and conservation importance. Intriguingly, the possibility of *H. mohnikei*

originating in India and subsequent dispersal to the east cannot be ruled out. Although there is

no published evidence to show that *H. mohnikei* originated in Japan/China, the original type

specimens studied were however, from Japan and Vietnam. At the same time, it may be plausible that *H. mohnikei* might be present in other localities, but has not been reported or overlooked. Further intense sampling and rigorous molecular analysis of *H. mohnikei* populations along the Indian coast would be therefore required to confirm whether these are vagrants or a previously unreported fairly established population.

Acknowledgements

The authors are grateful to Prof. Sunil Kumar Singh, Director, CSIR-National Institute of Oceanography, Goa (India) for encouragement and facilities. Thanks are due to local fishermen Mr. Harish Halarnkar for providing seahorse specimens. Thanks are also due to Dr. Mani Murali and Mr. Devanand Kavlekar for their help in preparing the maps. We would like to deeply thank Dr. Qiang Lin for sharing the morphometric data of *Hippocampus mohnikei* for comparison. The authors are thankful to Dr. Graham Short and Riley Pollom and two anonymous reviewers for their critical and constructive comments for improving the quality and presentation of the manuscript. This represents contribution No. xxxx of the CSIR-National Institute of Oceanography, Goa (India).

Author Contributions

Conceived and designed the experiments, writing-original draft: SVS, RAS, RDSK; Performed the experiments: SVS, MSS, HTK, MG; Data curation and formal analysis: SVS, RAS, MSS, RDSK, JN; Contributed reagents/materials/analysis tools: JN, HTK, MG; Visualization, Writing - review & editing: RAS, SVS, RDSK.

References

1. Sreepada RA, Desai UM, Naik S. The plight of Indian seahorses: Need for conservation and management. Curr Sci 2002; 82: 377–378.

2. Foster SJ, Vincent ACJ. Life history and ecology of seahorses: implications for conservation and management. J Fish Biol 2004; 65: 1–61.

3. Kuiter RH. Seahorses and Their Relatives. Aquatic Photographics, Seaford, Australia; 2009.

4. Froese R, Pauly D. FishBase.www.fishbase.org. Downloaded on 10/2018.

5. Harasti D, Martin-Smith K, Gladstone W. Ontogenetic and sex-based differences in habitat preferences and site fidelity of White's seahorse *Hippocampus whitei*. J Fish Biol 2014; 85: 1413–1428.

6. Hellyer C, Harasti D, Poore A. Manipulating artificial habitats to benefit seahorses in Sydney Harbour, Australia. Aquat Conserv Mar Fresh Ecosyst 2011; 21: 582–589.

7. Short G, Harasti D, Hamilton H. *Hippocampus whitei* Bleeker, 1855, a senior synonym of the southern Queensland seahorse *H. procerus* Kuiter, 2001: molecular and morphological evidence (Teleostei, Syngnathidae). ZooKeys 2019; 824: 109–133.

8. IUCN Seahorse, Pipefish and Stickleback Specialist Group (2019). https://www.iucn-seahorse.org (downloaded on 01/11/2019).

9. Salin KR, Mohankumaran NC. Resources and biodiversity of seahorses and the need for their conservation in India. Aquacult Asia 2006; 10: 3–8.

10. Murugan A, Dhanya S, Rajagopal S, Balasubramanian T. Seahorses and pipefishes of the Tamil Nadu coast. Curr Sci 2008; 95: 253–260.

11. Thangaraj M, Lipton AP. Occurrence of the Japanese seahorse *Hippocampus mohnikei* Bleeker 1854 from the Palk Bay coast of southeastern India. J Fish Biol 2007; 70: 310–312.

12. Krishnan TS, Kuiter RH, Patterson J. Occurrence of western Indian Ocean seahorse,

Hippocampus borboniensis Dumeril, 1870, in the Gulf of Mannar, Southeastern India.

Ind J Geo-Mar Sci 2011; 40: 642–644.

13. Yogeshkumar JS, Geetha S. A new species of *Hippocampus montebelloensis* (Family:

Syngnathidae) from the southeast coast of India. Int J Plant Anim Env Sci 2012; 2:

56–58.

14. Subburaman S, Murugan A, Goutham S, Kaul R, Prem Jyothi PVR, Balasubramanian

T. First distributional record of giraffe seahorse, *Hippocampus camelopardalis*

Bianconi 1854 (Family Syngnathidae) from Gulf of Kachchh waters, north west coast

of India. Indian J Geo-Mar Sci 2014; 43: 408–411.

15. Lourie SA, Foster SJ, Cooper EWT, Vincent ACJ. A Guide to the Identification of

Seahorses. Project Seahorse and TRAFFIC North America; 2004.

16. Lourie SA, Pollam RA, Foster SJ. A global revision of the seahorses *Hippocampus*

Rafinesque 1810 (Actinopterygii: Syngnathiformes): taxonomy and biogeography

with recommendations for further research. Zootaxa 2016; 4146: 1–66.

17. Lourie SA, Vincent ACJ, Hall HJ. Seahorses: an identification guide to the world's

species and their conservation. Project seahorse, London, UK; 1999.

18. Silveira RB, Siccha-Ramirez R, Silva JRS, Oliveira C. Morphological and molecular

evidence for the occurrence of three *Hippocampus* species (Teleostei: Syngnathidae)

in Brazil. Zootaxa 2014; 3861: 317–332.

19. Casey SP, Hall HJ, Stanley HF, Vincent ACJ. The origin and evolution of seahorses

(genus *Hippocampus*): a phylogenetic study using the cytochrome b gene of

mitochondrial DNA. Mol Phylogenet Evol 2004; 30: 261–272.

20. Wang X, Han X, Zhang Y, Liu S, Lin Q. Phylogenetic analysis and genetic structure
of the seahorse, *Hippocampus fuscus* from the Arabian and Red Sea based on
mitochondrial DNA sequences. Mitochondrial DNA A 2019; 30: 165–171.

21. Lourie SA, Green DM, Vincent ACJ. Dispersal, habitat differences and comparative
phylogeography of South East Asian seahorses (Syngnathidae: *Hippocampus*). Mol
Ecol 2005; 14: 1073–1094.

22. Teske PR, Hamilton H, Palsboll PJ, Choo CK, Gabr H, Lourie SA. et al. Molecular
evidence for long-distance colonization in an Indo-Pacific seahorse lineage. Mar Ecol
Prog Ser 2005; 286: 249–260.

23. Boehm JT, Waldman J, Robinson JD, Hickerson MJ. Population genomics reveals
seahorses (*Hippocampus erectus*) of the western mid-Atlantic coast to be residents
rather than vagrants. PloS One 2015; 10: e0116219.
http://dx.doi.org/10.1371/journal.pone.0116219.

24. Saarman NP, Louie KD, Hamilton H. Genetic differentiation across eastern Pacific
oceanographic barriers in the threatened seahorse *Hippocampus ingens*. Conserv
Genet 2010; 11: 1989–2000.

25. Lourie SA, Pritchard JC, Casey SP, Ky TS, Hall HJ. et al. The taxonomy of
Vietnam's exploited seahorses (family Syngnathidae). Biol J Linn Soc 1999; 66: 231–
256.

26. Woodall LC, Otero-Ferrer F, Correia M, Curtis JMR, Garrick Maidment N. et al. A
synthesis of European seahorse taxonomy, population structure, and habitat use as a
basis for assessment, monitoring and conservation. Mar Biol 2018; 165: 1–19.

27. Curtis JMR, Vincent ACJ. Life history of an unusual marine fish: survival, growth
and movement patterns of *Hippocampus guttulatus* Cuvier 1829. J Fish Biol 2006; 68:
707–733.

28. Caldwell I, Vincent ACJ. A sedentary fish on the move: effects of displacement on

long-snouted seahorse (*Hippocampus guttulatus* Cuvier) movement and habitat use.

Environ Biol Fish 2013; 96: 67–75.

29. Correia M, Caldwell IR, Koldewey HJ, Andrade JP, Palma J. Seahorse

(Hippocampinae) population fluctuations in the Ria Formosa Lagoon, south Portugal.

J Fish Biol 2015; 87: 679– 690.

30. Luzzatto DC, Estalles ML, Diaz De Astarloa JM. Rafting seahorses: the presence of

juvenile *Hippocampus patagonicus* in floating debris. J Fish Biol 2013; 83: 677–681.

31. Han SY, Kim JK, Tashiro F, Kai Y, Yoo JT. Relative importance of ocean currents

and fronts in population structures of marine fish: a lesson from the cryptic lineages of

the *Hippocampus mohnikei* complex. Mar Biodiv 2019; 49: 263–275.

32. Luzzatto DC, Pujol MG, Figueroa D, Diaz De Astarloa JM. The presence of the

seahorse *Hippocampus patagonicus* in the deep waters: additional evidence of the

dispersive capacity of the species. Mar Biodivers Rec 2014; 7: e71.

33. Harasti D. Range extension and first occurrence of the thorny seahorse *Hippocampus*

histrix in New South Wales, Australia. Mar. Biodivers Rec 2015; 8:

https://doi.org/10.1017/S1755267215000263.

34. Harasti D. Southward range extension of the great seahorse (*Hippocampus kelloggi*

Jordan and Snyder, 1901) in Australia. J Appl Ichthyol 2017; 1–3

https://doi.org/10.1111/jai.13414.

35. Mahapatro D, Mishra RK, Panda S. Range extension of a vulnerable Sea horse

Hippocampus fuscus (Actinopterygii: Syngnathidae) on the north-eastern Bay of

Bengal coast. Mar Biodivers Rec 2017; 10:6. DOI 10.1186/s41200-017-0108-z.

36. Wong NLWS, The JC. Direct observation of drifting juvenile seahorse *Hippocampus*

trimaculatus (Leach, 1814) beneath the offshore surface at Malacca Strait. Check List

2017; 13: 993–996.

37. Aylesworth L, Lawson LM, Laksanawimol P, Ferber P, Loh TL. New records of the

Japanese seahorse *Hippocampus mohnikei* in Southeast Asia lead to updates in range,

habitat and threats. J Fish Biol 2016; 88: 1620–1630.

38. Masuda H, Amaoka K, Araga C, Uyeno T, Yoshino T. The Fishes of the Japanese

Archipelago. Tokyo: Tokai University Press; 1984.

39. Kanou K, Kohno H. Early life history of a seahorse, *Hippocampus mohnikei,* in

Tokyo Bay Japan. Ichthyol Res 2001; 48: 361–368.

40. Suda Y, Inoue T, Nakamura M, Masuda N, Doi H, Murai T. Nearshore ichthyofauna

in the intermediate sandy beach, Doigahama Beach, Yamaguchi Prefecture, Japan. J

Nat Fish Univ 2004; 52: 11–29.

41. Lim K. New Singapore record of the seahorse, *Hippocampus mohnikei*. Singapore

Biodiver Rec 2013; 2013: 68–69.

42. Laksanawimol P, Petpiroon S, Damrongphol P. Trade of seahorses, *Hippocampus*

spp. (Actinopertygii: Syngnathiformes: Syngnathidae), on the East Coast of the Gulf

of Thailand. Acta Ichthyol Piscata 2013; 3: 229–235.

43. Chang CH, Jang-Liaw NH, Lin YS, Fang YC, Shao KT. Authenticating the use of

dried seahorses in the traditional Chinese medicine market in Taiwan using molecular

forensics. J Food Drug Analy 2013; 21: 310–316.

44. Zhang Y, Pham NK, Zhang H, Lin J, Lin Q. Genetic variations in two seahorse

species (*Hippocampus mohnikei* and *Hippocampus trimaculatus)*: evidence for

Middle Pleistocene population expansion. PloS One 2014; 9: e105494.

45. Pollom R. *Hippocampus mohnikei*. The IUCN Red List of Threatened Species 2017:

http://dx.doi.org/10.2305/IUCN.UK.20173.RLTS.T41005A54907304.en43.

Downloaded on 10 November 2018.

46. Song Z, Song J, Yue B. Population genetic diversity of Prenant's schizothoracin,

Schizothorax prenanti, inferred from the mitochondrial DNA control region. Environ

Biol Fish 2008, 81: 247–252.

47. Shetye SR, *Gouveia AD, Singbal* SY, Naik CG, Sundar D. et al. Propagation of tides

in the Mandovi–Zuari estuarine network. Proc Indian Acad Sci (Earth Planet Sci)

1995; 104: 667–682.

48. Qasim SZ, Sen Gupta R. Environmental characteristics of the Mandovi-Zuari

estuarine system. Estuar Coast Shelf Sci *1*981; 13: 557–578.

49. Madhupratap M, Nair KNV, Gopalkrishnan TC, Haridas P, Nair KKC. et al. Arabian

sea oceanography and fisheries of the west coast of India. *Curr Sci 2001; 81: 3*55–

361.

50. Ansari ZA, Sreepada RA, Dalal SG, Ingole BS, Chatterji A. Environmental influences

on the trawl catches in a bay-estuarine system of Goa, west coast of India. Estuar

Coast Shelf Sci 2003; 56: 503–515.

51. Ansari ZA, Chatterji A, Ingole BS, Sreepada RA, Rivonker CU. et al. Structure of

inshore demersal fish community at Goa. Estuar Coast Shelf Sci 1995; 41: 593–610.

52. Lourie S. Measuring seahorses. Project Seahorse Technical Report No.4, Version 1.0.

Project Seahorse, Fisheries Centre, University of British Columbia; 2003.

53. Zhang Y, Qin G, Wang X, Qiang L. A new species of seahorse (Teleostei:

Syngnathidae) from the South China Sea. Zootaxa 2016; 4170: 384–392.

54. Wibowo K, Tashiro S, Motomura H. Seahorses of the genus *Hippocampus*

(Syngnathidae) from Kagoshima Prefecture, southern Japan, with a new record of *H.*

mohnikei Bleeker, 1853. Nature of Kagoshima 2017; 43:71–76.

55. Underwood AJ. Experiments in Ecology: their logical and interpretation using

Analysis of Variance. Cambridge University Press. Cambridge, UK; 1997.

56. Kumar R Singh PJ, Nagpure NS, Kushwaha B, Srivastava SK. et al. A non-invasive

technique for rapid extraction of DNA from fish scales. Ind J Exp Biol 2007; 45: 992–

997.

57. Qin G, Zhang Y, Ho ALFC, Zhang Y, Lin Q. Seasonal distribution and reproductive

strategy of seahorse. ICES J Mar Sci 2017; 74: 2170–2179.

58. Lourie SA, Vincent ACJ. A marine fish follows Wallace's line: the phylogeography

of the three-spot seahorse (*Hippocampus trimaculatus*, Syngnathidae, Teleostei) in

Southeast Asia. J Biogeogr 2004; 31: 1975–1985.

59. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search

tool. J Mol Biol 1990; 215: 403–410.

60. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis,

version 7.0 for bigger datasets. Mol Biol Evol 2016; 33: 1870–1874.

61. Kimura M. A simple method for estimating evolutionary rates of base substitutions

through comparative studies of nucleotide sequences. J Mol Evol 1980; 16: 111–120.

62. Choi Y, Rho S, Park H, Kang D. Population characteristics of two seahorses,

Hippocampus coronatus and *Hippocampus mohnikei*, around seagrass beds in the

southern coastal waters of Korea. Ichthyol Res 2012; 59: 235–241.

63. Kwak S, Huh S, Kim H. Feeding habits of *Hippocampus mohnikei* in an eelgrass

(*Zostera marina*) bed. Korean J Ichthyol 2008; 20: 112–116.

64. Panithanarak T, Karuwanjaroen R. A preliminary study of genetic differentiation in
the Japanese seahorse (*Hippocampus mohnikei*) populations: a possibility of cryptic
species. Burapha Sci J 2019; 24: 124–137.

65. Hou F., Wen L, Peng C, Guo J. Identification of marine traditional Chinese medicine
dried seahorses in the traditional Chinese medicine market using DNA barcoding.
Mitochondrial DNA A 2018; 29(1):107–112.

66. Otsuka Y, Suzuki H, Akagawa I. Occurrence, gonad morphology and maturity of
Japanese seahorse *Hippocampus mohnikei* in Matsushima Bay, Japan. J School Mar
Sci Tech Tokai Univ 2009; 7: 11–22.

67. Wafar S. Ecology of mangroves along the estuaries of Goa; Ph. D. Thesis, Karnataka
University, Dharward, India; 1987.

68. Jagtap TG. Distribution of seagrasses along Indian coast. Aquac Bot 1991; 40: 379–
386.

69. Padmavati G, Goswami SC. Zooplankton ecology in the Mandovi-Zuari estuarine
system of Goa, west coast of India. Ind J Mar Sci 1996; 25: 268–273.

70. Dalal SG, Goswami SC. Temporal and ephemeral variations in copepod community
in the estuaries of Mandovi and Zuari– west coast of India. J Plankton Res 2001; 23:
19–26.

71. Das PB. Zooplankton assemblages in spatial stretches of a tropical estuary. Int J Res
Appl Sci Engg Technol 2017; 5: 1789–1799.

72. Qin G, Johnson C, Zhang Y, Zhang H, Yin J, Miller G. et al. Temperature-induced
physiological stress and reproductive characteristics of the migratory seahorse
Hippocampus erectus during a thermal stress simulation. Biol Open 2018; 25:
7(6).bio032888.

73. Scales H. Advances in the ecology, biogeography and conservation of seahorses
(genus *Hippocampus*). Prog Phys Geog 2010; 34: 443–458.

74. Bohonak AJ. Dispersal, gene flow, and population structure. Q Rev Biol 1999; 74:
21–45.

75. Palumbi SR. Genetic divergence, reproductive isolation, and marine speciation. Annu.
Rev Ecol Evol S 1994; 25: 547–572.

76. Wyrtki K. Physical oceanography of the southeast Asian waters. Scientific results of
maritime investigations of the South China Sea and the Gulf of Thailand 1959–1961.
Naga Report, 2. Scripps Institution of Oceanography, La Jolla, CA; 1961.

77. Shetye SR, Gouveia AD, Shenoi SSC, Michael GS, Sundar D, Almedia AM. et al.
The coastal current off western India during the northeast monsoon. Deep-Sea Res
1991; 38: 1517–1529.

78. Fleminger A. The Pleistocene equatorial barrier between the Indian and Pacific
oceans and a likely cause for Wallace's Line. In Pierrot-Bults AC, van der Spoel S,
Zahuranec BJ, Johnson RK editors, Pelagic marine science. UNESCO Tech Pap Mar
Sci 1986; 49: 84–97.

79. Maree RC, Whitfield AK, Booth AJ. Effect of water temperature on the biogeography
of South African estuarine fishes associated with the subtropical/warm temperate
subtraction. S Afr J Sci 2000; 96: 184–188.

80. Bowen BW, Bass AL, Rocha LA, Grant WS, Robertson DR. Phylogeography of the
trumpetfishes *(Aulostomus)*: ring species complex on a global scale. Evolution 2001;
55: 1029–1039.

81. Martin-Smith KM, Vincent ACJ. Seahorse declines in the Derwent estuary, Tasmania,
in the absence of fishing pressure. Biol Conserv 2005; 123: 533–545.

82. Shokri MR, Gladstone W, Jelbart J. The effectiveness of seahorses and pipefish

(Pisces: Syngnathidae) as a flagship group to evaluate the conservation value of

estuarine seagrass beds. Aquat Conserv 2009; 19: 588–595.

Supporting Information

S1 Fig. Picture of *H. mohnikei* posted on social media. Male specimen of *Hippocampus*

mohnikei caught by gill net in the Chapora estuary, Goa, India

S1 Table. The COI and Cyt *b* gene sequences. GenBank accession numbers and sources of

the mitochondrial gene sequences (COI and Cyt *b*) of seahorse species used for

construction of phylogenetic trees.

S2 Table. The COI and Cyt *b* gene sequences. GenBank accession numbers and sources of

the mitochondrial gene sequences (COI and Cyt *b*) of seahorse species used for

construction of phylogenetic trees.

Table 1. Detailed morphology of *H. mohnikei*. Morphological characteristics of four

specimens of *Hippocampus mohnikei* collected from the Mandovi estuary, Goa, central west

coast of India (MK1 and MK2 = Female; MK3 and MK4 = Male).

	MK1	MK2	MK3	MK4
Morphometric characters (mm)				
Height (Ht)	39.2	54.2	48.1	51.3
Standard length (SL)	47.3	60.1	55.2	57.2
Trunk length (TrL)	16.1	19.4	15.2	15.1
Tail length (TaL)	24.2	32.1	31.2	33.1
Head length (HL)	8.5	9.1	9.2	8.7
Snout length (SnL)	2.5	3.1	3.3	2.5
HL/SnL ratio	3.4	3.2	3.3	3.5
Snout depth (SnD)	1.5	2.1	2.1	2.1
Head depth (HD)	5.5	7.1	6.9	7.2
Eye orbit	1.5	2.1	2.2	2.1
Trunk depth between 4^{th} and 5^{th} trunk ring (TD4)	3.1	4.2	3.9	4.1
Trunk depth between 9^{th} and 10^{th} trunk ring (TD9)	4.7	6.1	5.8	5.9
Pectoral fin base length	2.1	2.1	2.1	2.2
Dorsal fin base length	3.5	4.1	4.1	3.9
Anal fin base length	0.5	0.5	0.5	0.5
Pouch length	-	-	6.2	6.6
Pouch depth	-	-	3.1	2.5
Pouch width	-	-	3.2	3.3
Meristic features (numbers)				
Trunk Rings (TrR)	11	11	11	11
Tail Rings (TaR)	37	38	37	39
TrR supporting dorsal fin	2	2	2	2
TaR supporting dorsal fin	1	1	1	1
Cheek spines	2	2	2	2
Eye spines	2	2	2	2
Enlargement of trunk Rings	1, 4, 7, 11	1, 7, 10	1, 4, 7, 11	1, 4, 7, 10
Enlargement of tail Rings	4, 7, 10, 13	4, 8, 10, 13	4, 7, 9, 13	4, 7, 10, 14
Dorsal fin rays	15	16	16	16
Pectoral fin rays	12	14	14	14
Anal fin rays	4	4	4	4

Table 2. Morphological comparison of seahorses. Comparison of morphological characteristics of *H. mohnikei* with other two seahorse species, (*H. trimaculatus* and *H. kuda)*. Sample size for each species shown in parentheses. Mean±SD denoted with different superscripts significantly differ from each other (P < 0.05).

Morphometrics (mm)	*H. mohnikei* (*n* = 4)	*H. kuda* (*n* = 10)	*H. trimaculatus* (*n* = 10)
Height (Ht)	48.2 ± 6.50^a	157.73 ± 19.24^b	93.06 ± 14.71^c
Standard Length (SL)	54.95 ± 5.48^a	182.65 ± 11.64^b	109.87 ± 6.45^c
Trunk Length (TrL)	16.45 ± 2.02^a	47.83 ± 2.54^b	26.40 ± 2.12^c
Tail Length (TaL)	30.15 ± 4.04^a	104.43 ± 9.07^b	61.70 ± 3.40^c
Head Length (HL)	8.83 ± 0.41^a	30.39 ± 1.88^b	21.70 ± 1.77^c
Snout Length (SnL)	2.67 ± 0.21^a	12.06 ± 1.04^b	8.90 ± 0.99^c
HL/SnL ratio	3.33 ± 0.14^a	2.54 ± 0.16^b	2.45 ± 0.19^{bc}
Snout Depth	1.95 ± 0.30^a	4.25 ± 0.41^b	3.45 ± 0.16^c
Head Depth	6.68 ± 0.79^a	14.80 ± 1.84^b	9.50 ± 0.53^c
TD4	3.83 ± 0.50^a	12.16 ± 1.19^b	6.75 ± 0.86^c
TD9	5.63 ± 0.63^a	18.68 ± 1.51^b	9.55 ± 0.90^c
Pectoral fin base length	2.03 ± 0.1^a	7.07 ± 1.2^b	4.75 ± 0.6^c
Dorsal fin base length	3.88 ± 0.3^a	15.73 ± 1.8^b	9.80 ± 0.8^c
Meristic counts			
Trunk Rings (TrR)	11 ± 0.00	11 ± 0.00	11 ± 0.00
Tail Rings (TaR)	38 ± 1.00^a	36 ± 1.00^b	41 ± 1.00^c
TrR supporting Dorsal Fin	2 ± 0.00	2 ± 0.00	2 ± 0.00
TaR supporting Dorsal Fin	1 ± 0.00	1 ± 0.00	1 ± 0.00
Cheek Spines	2 ± 0.00^a	1 ± 0.00^b	1 ± 0.00^c
Eye spines	2 ± 0.00^a	1 ± 0.00^b	1 ± 0.00^c
Dorsal fin rays	16 ± 1.00^a	18 ± 1.00^b	20 ± 1.00^c
Pectoral fin rays	14 ± 1.00^a	16 ± 1.00^b	17 ± 1.00^c

Table 3. Comparison of morphological characters. Comparison of diagnostic morphological characters (range and mean) of vouchered specimens of *Hippocampus mohnikei*. Values in parentheses are mean.

Data source	Present Study	Lourie et al. [15,25]	Thangaraj and Lipton [11]	Zhang et al. [53]	Wibwo et al. [54]
Voucher/field Number	NIO1015/19	RMNH7259	CMFRI No. 146	SCSMBC007418–19	KAUM-1. 17724
Locality	Goa, west coast of India	Vietnam	Southeast coast of India	Yellow Sea, China	Kagoshima, Southern Japan
No. of samples (N)	04	04	01	16	01
Morphometrics (mm)					
Standard Length (SL)	47.3–60.1 (54.95)	54.0–57.0	70.0	52.5–83.0 (66.4)	-
Trunk Length (TrL)	15.1–19.4 (16.45)		22.0	15.0–23.0 (19.2)	-
Tail Length (TaL)	24.2–33.1 (30.15)		38.0	28.5–46.0 (36.7)	-
Head Length (HL)	8.5–9.2 (8.83)		10.0	9.0–14.0 (10.5)	-
Snout Length (SnL)	2.5–3.3 (2.67)		3.0	1.8–3.9 (2.4)	
HL/SnL ratio	3.2–3.5 (3.35)	2.8–3.9 (3.0)	3.33	3.59–5.75 (4.5)	3.1
Snout Depth	1.5–2.1 (1.95)	-	2.5	-	-
Head Depth	5.5–7.2 (6.68)	-	7.0	-	-
TD4	3.1–4.2 (3.83)	-	5.5	-	-
TD9	4.7–6.1 (5.63)	-	8.0	-	-
Meristic counts					
Trunk Rings (TrR)	11	11	11	11	11
Tail Rings (TaR)	3737–39 (38)	37–40 (38)	38	37–40 (38)	38
TrR supporting dorsal fin	2	2	-	-	-
TaR supporting dorsal fin	1	1	-	-	-
Cheek Spines	2	2	2		2
Rounded spines below eye	Double	Double	Single	-	-
Dorsal fin rays (DF)	15–16	15–16	14	16	15
Pectoral fin rays	12–14	12–14 (13)	12	12–14 (13)	12
Anal fin rays (AF)	4	-	-	4	-

Table 4. Pair-wise genetic distance based on COI gene sequences. Pair-wise genetic distances (Kimura 2-parameter model) between *Hippocampus mohnikei* populations of Goa and the Indo-Pacific region based on COI gene sequences. Values in parentheses indicate percentage.

Localities of *H. mohnikei* with GenBank accession numbers	Goa	Japan	Vietnam	China
Goa Specimens (MN595216–MN595218; MK330041.1)	0.00			
Japan (GQ502158.1–GQ502159)	0.008 (0.8)	0.00		
Vietnam (GQ502157.1)	0.007 (0.7)	0.009 (0.9)	0.00	
China (KP140138–KP140181)	0.010 (1.0)	0.003 (0.3)	0.010 (1.0)	0.00

772 **Table 5. Pair-wise genetic distance based on Cyt *b* gene sequences.** Pair-wise genetic distances (Kimura 2-parameter model) between

773 *Hippocampus mohnikei* populations of Goa and the Indo-Pacific region based on Cyt *b* gene sequences. Genetic divergence of > 2% within

774 same locality separated out. Values in parentheses indicate percentage.

Localities of *H. mohnikei* with GenBank Accession numbers	Goa	Japan1	Japan2	Korea1	Korea2	China	Thailand
Goa Specimens (MN595213.1, MN595214.1, MN595215.1 and MK112274.2)	0.00						
Japan1 (AF192688.1‒AF192689.1; KT731898.1‒KT731909.1)	0.067 (6.7)	0.00					
Japan2 (KX017612.1‒KX017625.1)	0.070 (7.0)	0.071 (7.1)	0.00				
Korea1 (KT731897.1–KT731920.1; KT731924.1–KT731926.1; KT731929.1; KT731932.1–KT731935.1; KT731940.1–KT731941.1; KT731944.1–KT731952.1 and KT731955.1)	0.066 (6.6)	0.040 (4.0)	0.044 (4.4)	0.00			
Korea2 (KT731921.1–KT731923.1; KT731928.1; KT731930.1–KT731931.1; KT731936.1–KT731937.1, KT731943.1 and KT731955.1)	0.067 (6.7)	0.004 (0.4)	0.012 (1.2)	0.066 (6.6)	0.00		
China (KT946929.1–KT946937.1; KC527556.1 and KC527584.1)	0.066 (6.6)	0.037 (3.7)	0.042 (4.2)	0.057 (5.7)	0.004 (0.4)	0.00	
Thailand (EU179923.1–EU179924.1)	0.007 (0.7)	0.001 (0.1)	0.001 (0.1)	0.001 (0.1)	0.001 (0.1)	0.001 (0.1)	0.00

Fig 1. Sampling localities of *H. mohnikei*. A map showing major rivers and estuarine

systems of Goa (central west coast of India). Location of a suspected adult male *H.*

mohnikei in Chapora estuary (●); collection site of confirmed report of *H. mohnikei* in the

Mandovi estuary (★).

Fig 2. Collected seahorses from Goa. Sub-adult specimens of *Hippocampus mohnikei*

(Bleeker 1853) (females, MK1 and MK2; males, MK3 and MK4; NIO1015/19) from the

Mandovi estuary, Goa, central west coast of India.

Fig 3. Close-up view of *H. mohnikei*. *Hippocampus mohnikei (*Bleeker 1853) (A-male;

B-female) from the Mandovi estuary, Goa, central west coast of India (NIO1015/19).

Fig. 4. Oceanic circulation in Indo-Pacific region. Climatological surface currents

during winter season (December–February) for the Indo-Pacific region. Red dots indicate

the geographic locations of distribution of *Hippocampus mohnikei.* (WICC = West Indian

Coastal Current, EICC = East India Coastal Current, NEC= North Equatorial Current and

SCSWC = South China Sea Warm Current).

Fig. 5. PCA analysis scores. Scores of factor loadings plot of principal component

analysis of three species, *H. mohnikei*, *H. kuda* and *H. trimaculatus* based on

morphological data.

Fig. 6. Phylogenetic relationships of Goa specimens with other seahorse species based
on COI gene sequences. ML phylogenetic tree of Goa specimens and all vouchered
Hippocampus mohnikei based on mitochondrial cytochrome oxidase subunit I gene (COI)
sequences using K2P distances with 1000 bootstrap iterations. Numbers at nodes indicate
bootstrap support values and nodes with >50% bootstrap support values are shown. The
alligator pipefish, *Syngnathoides biaculeatus* (KY066152.1) served as an out-group
species. Scale bar = genetic distance of 0.05. ML = Maximum likelihood

Fig. 7. Phylogenetic relationships of Goa specimens with other seahorse species based on Cyt *b* gene sequences. ML phylogenetic tree of Goa specimens and all vouchered *Hippocampus mohnikei* based on mitochondrial cytochrome *b* gene (Cyt *b*) sequences using K2P distances with 1000 bootstrap iterations. Numbers at nodes indicate bootstrap support values and nodes with >50% bootstrap support values are shown. The alligator pipefish, *Syngnathoides biaculeatus* (EF189167.1) served as an out-group species. Scale bar = genetic distance of 0.05. ML = Maximum likelihood

829 **Fig. 8. Reported sightings of *H. mohnikei* in Indo-Pacific region.** Map showing native
830 range, recent sightings and all reported sightings for *Hippocampus mohnikei* in the Indo-
831 Pacific region. Sightings are classified as new (this study, ★), recently published (2004 –
832 2016, (▲) and original records [15] (*). Map was modified from Aylesworth et al. [37]
833 and re-created in the World Geodetic System (WGS) 84 datum.