Morphological and molecular evidence for range extension and first

occurrence of the Japanese seahorse, *Hippocampus mohnikei* (Bleeker 1853)

in a bay-estuarine system of Goa, central west coast of India

Sushant Sanaye[1], Rakhee Khandeparker[1], Rayadurga Anantha Sreepada[1*], Mamatha

Singanhalli Shivaramu[1], Harshada Kankonkar[1], Jayu Narvekar[2]

[1]Aquaculture Laboratory, Biological Oceanography Division, CSIR-National Institute of

Oceanography, Dona Paula, Goa–403 004, India

[2]Physical Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula, Goa–403

004, India

*sreepada@nio.org; sreepada63@gmail.com

Abstract

Accurate information of taxonomy and distribution range of seahorse species (genus

Hippocampus) is the first step in preparing threat assessments and designing effective

conservation measures. Here, we report the range expansion and first occurrence of the

Japanese seahorse, *Hippocampus mohnikei* (Bleeker, 1853) from the Mandovi estuarine

ecosystem of Goa, central west coast of India based on morpho-molecular analyses. The

morphometric and meristic traits particularly, short snout length, double cheek spine, low

coronet and long tail observed in all four collected specimens matched with the reported

diagnostic morphological criteria of *H. mohnikei*. The seahorse mitochondrial cytochrome

oxidase subunit I (COI, 609 bp) and cytochrome *b* (Cyt *b*, 639 bp) genes were partially

sequenced for conclusive genetic identification of the species under study. Using the

maximum similarity with GenBank database, maximum likelihood network and subsequent

morphological analysis, the identity of the collected seahorse species was reconfirmed as *H.*

mohnikei. With this new report, the distributional range of *H. mohnikei* extended significantly

to the westward side from its hitherto known range of Japan and Vietnam.The possible role of

prevailing oceanic circulation in the Indo-Pacific region in facilitating dispersal and range

extension of *H. mohnikei* has been highlighted. Comparison of the pair-wise genetic distances

(Kimura 2-parameter) based on COI and Cyt *b* sequences revealed that the conspecificity of

Goa specimens with populations of *H. mohnikei* from Vietnam and Thailand rather than

Japan. To test the hypothesis whether *H. mohnikei* populations are vagrants or residents of a

bay-estuarine system, long-term inter-annual sampling and analyses are warranted.

Introduction

Seahorses (Syngnathiformes; Syngnathidae) belonging to the single genus, *Hippocampus*

(Rafinesque, 1810) are a fascinating and remarkable group of fishes with their unusual body

shape and their biology, with males incubating the fertilized eggs in a brood pouch [1]. They

are small, cryptic, and sedentary marine fishes that occur worldwide in shallow temperate to

tropical waters in a wide variety of habitats, including seagrass beds, estuaries, coral and

rocky reefs and mangroves [2−4]. Their remarkable ability to camouflage with structurally

complex habitats further reduces the risk of predation [5], while providing stealth for prey

capture [6]. However, such biological traits often also make them unnoticeable from coastal

habitat, particularly in those regions where seahorse trade is absent. Often overlooked by

most fishermen, cryptic nature, ability to camouflage and the sparse distribution restrict

biologists to identify their presence in thick coastal marine habitats.

Out of 41 valid seahorse species (Genus *Hippocampus*) throughout the world [7],

Hippocampus kelloggi, *H. kuda*, *H. spinosissimus* and *H. trimaculatus* are most common in

Indian coastal marine waters [8,9]. In addition, sightings based on morphological

identification of single specimens, *H. mohnikei* from the Palk Bay, southeastern India [10], *H.*

borboniensis [11] and *H. montebelloensis* [12] from Gulf of Mannar, southeast coast of India

and *H. camelopardalis* from Gulf of Kachchh, northwest coast of India [13] have been

reported. Another seahorse species, *H. histrix* is also suspected to occur in Indian coastal

waters [14].

The utility of morphological traits commonly used in diagnosis of *Hippocampus* spp. is

quite challenging as they lack certain key physical features (e.g. pelvic and caudal fins), high

variability and overlapping of body proportions, colour (camouflage) and skin filaments

within and between species [3,15]. On the other hand, molecular methods have proved

helpful in resolving morphologically challenging seahorse taxonomy [7,16], phylogenetic

relationship within genus, *Hippocampus* [17,18], natural species boundaries [19−21] and

genetic variability [20,22] of many seahorse species. An integrated approach of combination

of morphological and genetic analyses [23] would aid in the management of demographically

separate populations as independent units and allow international legal mechanisms and

international agreements such as the convention on international trade in endangered species

(CITES) to work effectively [24].

Generally, seahorses are sedentary fish species with small home range and patchy

distribution in sheltered areas such as seagrasses and seaweeds [2,25,26]. However, instances

of small migration of seahorses in search of proper habitat, food or holdfasts have been

reported [26,27]. Evidence of long-distance dispersal of seahorse species via rafting (floating

debris and seaweeds) [2,28,9] and subsequent colonisation [20] have been documented. The

role of oceanic currents facilitating as stepping-stones in long-distance dispersal and range

expansion of many seahorse species has also been highlighted [29−34].

The Japanese seahorse or Lemur-tail seahorse, *Hippocampus mohnikei* (Bleeker, 1853) is

a small coastal seahorse species (5–8 cm) inhabiting seagrasses, mangroves, oyster beds and

mussel farms along the coasts of continental Asia from southeastern India to Korea and Japan

[35]. Previous reports [15,36−38] indicated that the distribution of *H. mohnikei* is restricted or

native to the coastal waters around Japan and Vietnam. As a consequence of recent

exploitations [35,39−41], the status of *H. mohnikei* in the IUCN Red List of Threatened

Species has been listed as 'vulnerable' [42].

A probable occurrence of *H. mohnikei* along the central west coast of India (CWCI) began

in February 2017 when a fisherman posted a picture of seahorse incidentally caught in gill net

operated in the Chapora estuary, Goa (India) (15.6120° N, 73.7506° E) on social media [S1

Fig.] which was suspected to be an adult male of *H. mohnikei*. Sustained follow-up surveys

for its occurrence in the surrounding environs of the Chapora estuary did not yield in any

further reporting. This contribution describes the range expansion and first occurrence of *H.*

mohnikei from the Mandovi estuarine ecosystem, Goa, CWCI. To test further the hypothesis

that the morphologically identified seahorse species is *H. mohnikei*, gene markers (COI and

Cyt *b* genes) were employed for reconfirmation. Potential role of ocean currents on the range

expansion of *H. mohnikei* to north-western Asia have also been discussed. In the wake of the

vulnerability of seahorse populations to the threats such as habitat alteration/destruction and

fishing pressure globally, the present sighting of *H. mohnikei* in a bay-estuarine system is of

considerable conservation and biogeographic significance.

Materials and Methods

Collection site and seahorses

The collection site of seahorses (Brittona, Goa, India, 15.3059°N, 73.5073° E, Fig 1) is a

shallow water littoral environment which is in the proximity to the mangrove-dominated

Chorao Island in the Mandovi estuary, CWCI. The currents are largely tide dominated, the

tides being semi-diurnal with a mean amplitude of 1.3 m. Currents are also influenced by the

large seasonal freshwater influx [43] during the summer monsoon season (June to

September). Temporal variations in abiotic and biotic factors influenced by monsoonal

precipitation and seasonal upwelling render the collection location as one of the most

ecologically complex ecosystems [44,45]. Significant lowering of salinity occurs when the

freshwater discharge is at its maximum during monsoon. During the post- and pre-monsoon

periods, the salinity at the surface and bottom remains high (~25 ppt). The bottom topography

at the collection site is characterized by muddy and comprises mainly of silt and clay with

rich organic matter content [46]. The collection site is reported to harbour rich and diverse

fishery resources and its importance in supporting the life history stages of several marine

teleosts has also been well-documented [47]. Present report is based on four specimens of *H.*

mohnikei (Fig. 2. and Fig. 3) that were incidentally caught in bag nets attached to the fixed

stakes (total length, 25 m; mouth length, 5 m; cod end mesh size, 10 mm) operated at the

water depth of ~6 m in the Mandovi estuary during July 2018. Freshly landed dead seahorses

as incidental catch in bag nets attached to the stakes operated by local fishermen in the

estuarine system formed the study material. No live seahorses were specifically caught for the

present study. Fin clips from these dead seahorses were used for molecular analysis. The

present study did not involve any experimentation with live animals and hence ethics

committee's approval for the usage of live animals for experiments is inconsequential.

Fig 1. Sampling localities of *H. mohnikei*. A map showing major rivers and estuaries of Goa

(central west coast of India). Location of a suspected adult male *H. mohnikei* in Chapora

estuary (●); collection site of confirmed report of *H. mohnikei* in the Mandovi estuary (★).

Fig 2. Collected seahorses from Goa. Specimens of *Hippocampus mohnikei* (MK1 and

MK2 = females; MK3 and MK4 = males) collected from the Mandovi estuary, Goa, central

west coast of India.

Fig 3. Close-up of *H. mohnikei*. A closer view of *Hippocampus mohnikei* (A-male; B-

Female) collected from the Mandovi estuary, Goa, central west coast of India.

Morphological analysis

Before subjecting them to a detailed analysis of morphological characteristics, seahorses were

carefully photographed (Nikon D7200, Japan) and sexed via observation of presence or

absence of brood pouch. Morphometric measurements and meristic counts were carried out

as per standardized protocols [14,15,48]. Morphometric characters such as height (Ht, linear

distance from the top of the head to the tip of the stretched tail), standard length (SL), head

length (HL), trunk length (TrL) and tail length (TaL) were measured. Measurements of snout

length (SnL), ratio of head/snout lengths (HL/SnL), snout depth (SnD), head depth (HD),

pectoral fin base length (PL) and dorsal fin base length (DL) were also conducted. All

measurements were taken with the help of digital caliper to 0.1 mm and repeated to ensure

accuracy. Coronet height (CH), trunk depths on 4th trunk ring (TD4) and 9th trunk ring (TD9)

were measured. The presence/absence of spines and dermal appendages was also noted.

Meristic characters such as the number of trunk rings (TrR), tail rings (TaR), dorsal fin rays

(DF), pectoral fin rays (PF) and anal fin rays (AF) were counted under stereo zoom

microscope (Olympus SZX7, Japan) and the values were confirmed by three time counting.

Brood pouch dimensions were also taken in case of male specimens.

The salient morphometric and meristic characters of the collected seahorse specimens

were compared with the reported diagnostic features for different seahorse species [14,15,23].

The voucher material is deposited in the national marine biodiversity repository at CSIR-

National Institute of Oceanography, Goa, India (deposition ID: NIO1015/19).

Statistical analysis

Morphological data of *H. mohnikei* and two other commonly occurring seahorse species (*H.*

kuda and *H. trimaculatus*) was evaluated by one-way analysis of variance (ANOVA)

followed by the Duncan's multiple-range tests [49]. The level of significance was tested at

5.0%, represented as $P < 0.05$. A comparison of morphological characters and meristic

features of *H. mohnikei* (n = 4) with *H. kuda* (n = 10) and *H. trimaculatus* (n = 10) was made

through principal component analysis (PCA). PCA was employed in which factor loadings

based on components were used to determine the morphometric factors using PRIMER

software (version 6) after square-root transformation.

Molecular analysis: DNA extraction, PCR amplification and sequencing

For gene-based phylogenetic analysis, fragments of dorsal fin from fresh seahorse were

removed and preserved in 95% ethanol at −20°C until subjected to DNA extraction. DNA

extraction was carried out as described [50] with a few modifications. In brief, lysis buffer

and sodium dodecyl sulfate were added to the fin tissue and vortexed. Further, RNase and

Proteinase K were added to the mixture and incubated for 15–20 minutes at room

temperature. After the incubation period, the DNA was extracted using

phenol:chloroform:isoamyl alcohol (25:24:1) mixture. Precipitated DNA was washed with

70% cold ethanol. The gel electrophoresis was performed to confirm the quality and integrity

of DNA.

Seahorse mitochondrial cytochrome oxidase subunit I (COI) gene was amplified using

primers, forward (5'TCAACTAATCACAAAGACATCGGCAC3′) and reverse

(5'ACTTCGGGGTGCCCAAAGAATC3′) [51]. Similarly, mitochondrial cytochrome *b* (Cyt

b) gene fragment was also amplified using PCR with seahorse-specific primers, forward

(5'AACYAGGACYAATGRCTTGA3') and reverse (5′GCASWAGGGAGGRKTTTAAC3′)

[40]. Reactions were carried out in a thermal cycler (Veriti® 96-well Thermal Cycler,

Applied Biosystems, CA, USA). The amplification of COI and Cyt *b* genes was carried out

by following cycling conditions as described [52] with few modifications; an initial

denaturation step of 95°C for 2 minutes followed by 39 cycles of 94°C for 30s, 55°C for 30s

72°C for 1 minute and a final extension at 72°C for 2 minutes. Amplified PCR products of

both COI and Cyt *b* genes were verified on agarose gel (1.5%) and purified for sequencing

using PCR purification kit (Sigma Aldrich, USA). Gene sequence of the PCR products was

determined using a taq dye deoxy terminator cycle sequencing kit (PerkinElmer, CA, USA)

following supplier's protocols. Sequencing reaction products were analyzed on 373

automated DNA Sequencer (Model 313 xl; Applied Biosystems, CA, USA). Amplification of

the mitochondrial COI and Cyt *b* genes resulted in fragments of 609 bp and 639 bp,

respectively.

Phylogenetic analysis

Sequences of Goa specimens (GenBank accession numbers MK330041.1 and MK112274.2

for COI and Cyt *b* genes, respectively) were compared using the basic local alignment search

tool (BLAST) of NCBI [53] (http://www.ncbi.nih.gov/BLAST). Sequences of Goa specimens

were aligned with those of morphologically similar seahorse species (*H. coronatus* and *H.*

mohnikei) and the most commonly occurring species in the Indo-Pacific region (*H.*

camelopardalis, *H. comes, H. histrix, H. kelloggi, H. kuda, H. sindonis, H. spinosissimus* and

H. trimaculatus) retrieved from the GenBank database. The mitochondrial COI and Cyt *b*

gene sequences were aligned using the ClustalW algorithm implemented in MEGA 7 [54]. To

ascertain stable and reliable relationships among the seahorse species, phylogenetic trees

were constructed through maximum likelihood (ML) method. The analyses were performed

in MEGA 7, and confidence levels were assessed using 1000 bootstrap replications. The

alligator pipefish, *Syngnathoides biaculeatus* was used as an outgroup species for

phylogenetic analysis to differentiate between different seahorse species. The genetic

distances of *H. mohnikei* from different geographical locations were calculated using Kimura

2-parameter (K2P) distance model [55] implemented in MEGA 7 [54]. Codon positions

included were 1st+2nd+3rd+noncoding. All ambiguous positions with alignment gaps and

missing data were removed from the sequences (pair-wise deletion option).

Prevailing ocean circulation

In order to understand the basin-wide circulation in the Indo-Pacific region (encompassing

South China Sea, East China Sea, and southeastern Bay of Bengal), monthly mean

climatology was prepared based on daily data on U (zonal) and V (meridional) components

of geostrophic current velocities derived from Aviso merged product for the years,

2007–2017. Altimeter products by Ssalto/Duacs and distributed by Aviso+, with support

from Cnes (https://www.aviso.altimetry.fr) were used.

Results

Material examined

Colour of the head and body of freshly collected seahorse specimens was almost dark brown

to dark black. The entire body was covered with low spines and the coronet was observed to

be short with five tiny projections. Two prominent cheek spines and two small rounded

spines below the eye were discernible in all four specimens. (Fig 2. and Fig 3.).

Morphological taxonomy

Data on detailed morphometric measurements and meristic counts recorded on four seahorse

specimens from Goa are presented in Table 1. Measured height (Ht) and standard length (SL)

of specimens varied between 39.2 and 54.2 mm (48.2 ± 6.5 mm) and 47.3 and 60.1 mm

(54.95 ± 5.48 mm), respectively. Trunk lengths (TrL) and tail lengths (TaL) varied between

15.1 and 19.4 mm (16.45 ± 2.02 mm) and 24.2 and 33.1 mm (30.15 ± 4.04 mm), respectively.

TaL was observed to be relatively long compared to the TrL of the body. The ratio of the

head length (HL) to snout length (SnL) of all specimens varied between 3.2 and 3.5.

Number of trunk rings (TrR) in all specimens were 11 whereas, the number of tail rings

(TaR) varied narrowly between 37 and 39. The number of TrR supporting dorsal fin was two

and the number of TaR supporting dorsal fin was counted to be one in all specimens. Number

of dorsal fin rays, pectoral fin rays and anal fin rays recorded were 15–16, 12–14 and 4,

respectively. Slight enlargements, respectively of 1st, 4th, 7th, and 10/11thTrR and 4th, 7/8th,

9/10th, 13/14thTaR were prominently discernible. The measured morphometric and meristic

characters of Goa specimens are consistent with the morphological descriptions for *H.*

mohnikei reported by Lourie et al. [14,15,23] for specimens collected from Vietnam and

Japan.

Table 1. Detailed morphology of *H. mohnikei*. Morphological characteristics of four

specimens of *Hippocampus mohnikei* collected from the Mandovi estuary, Goa, central west

coast of India (MK1 and MK2 = Female; MK3 and MK4 = Male).

	MK1	MK2	MK3	MK4
Morphometric characters (mm)				
Height (Ht)	39.2	54.2	48.1	51.3
Standard length (SL)	47.3	60.1	55.2	57.2
Trunk length (TrL)	16.1	19.4	15.2	15.1
Tail length (TaL)	24.2	32.1	31.2	33.1
Head length (HL)	8.5	9.1	9.2	8.7
Snout length (SnL)	2.5	3.1	3.3	2.5
HL/SnL ratio	3.4	3.2	3.3	3.5
Snout depth (SnD)	1.5	2.1	2.1	2.1
Head depth (HD)	5.5	7.1	6.9	7.2
Eye orbit	1.5	2.1	2.2	2.1
Trunk depth between 4th and 5th trunk ring (TD4)	3.1	4.2	3.9	4.1
Trunk depth between 9th and 10th trunk ring (TD9)	4.7	6.1	5.8	5.9
Pectoral fin base length	2.1	2.1	2.1	2.2
Dorsal fin base length	3.5	4.1	4.1	3.9

Anal fin base length	0.5	0.5	0.5	0.5
Pouch length	-	-	6.2	6.6
Pouch depth	-	-	3.1	2.5
Pouch width	-	-	3.2	3.3

Meristic features (numbers)

Trunk Rings (TrR)	11	11	11	11
Tail Rings (TaR)	37	38	37	39
TrR supporting dorsal fin	2	2	2	2
TaR supporting dorsal fin	1	1	1	1
Cheek spines	2	2	2	2
Eye spines	2	2	2	2
Enlargement of trunk Rings	1, 4, 7, 11	1, 7, 10	1, 4, 7, 11	1, 4, 7, 10
Enlargement of tail Rings	4, 7, 10, 13	4, 8, 10, 13	4, 7, 9, 13	4, 7, 10, 14
Dorsal fin rays	15	16	16	16
Pectoral fin rays	12	14	14	14
Anal fin rays	4	4	4	4

Comparative morphology

A comparative morphological features of *H. mohnikei* and other two commonly occurring

seahorse species (*H. kuda* and *H. trimaculatus*) is presented in Table 2. Fresh specimens of

H. kuda was orange to dark brown with reddish spots over the body with single eye and

cheek spine, rounded and backward curled coronet and deeper head and rounded or less

developed body spines [14,15]. *H. trimaculatus* has three prominent black spots on the dorso-

lateral surface of the first, fourth and seventh trunk rings [14,15] which are clearly absent in

H. mohnikei. Single eye and hooked like cheek spine also observed in *H. trimaculatus*. Snout

length (SnL) of *H. mohnikei* was shorter compared to the *H. kuda* and *H. trimaculatus*. These

morphological characters clearly distinguishes *H. mohnikei* from *H. kuda* and *H.*

trimaculatus.

PCA used to elucidate the factors contributing to morphological differences in *H.*

mohnikei, H. kuda and *H. trimaculatus* showed that many morphometric and meristic

characters play an important role in differentiating the seahorse species. Results of PCA

analysis revealed that 11 morphometric and meristic characters integrated into two principal

components (PC1 and PC2) which cumulatively contributed to 97% of total variance (Table

3). The PC1 which explained 81% of total variation consisted of morphometric characters

such as SL, TaL, TD4, HL, TD9, TrL, HD, SnL and SnD while, the PC2 which explained

16% of total variation consisted of the TaR and DF. The derived ordination diagram after

plugging standardized morphological data of all individuals of three seahorse species into

PCA is depicted in Fig. 4. PCA ordination diagram showed that three seahorse species were

clustered out distinctly in orientation space. Therefore, based on comparison of

morphological traits [14,15,23] and the results of PCA, the identity of seahorse specimens

collected from Goa was confirmed as *H. mohnikei*.

Table 2. Morphological comparison of seahorses. Comparison of morphological characteristics of *H.*
mohnikei with other two seahorse species, (*H. kuda* and *H. trimaculatus)*. Sample size for each species are
shown in parentheses, Mean±SD denoted with different superscripts in same row significantly differ from
each other (P < 0.05).

Morphometrics (mm)	*H. mohnikei* (*n* = 4)	*H. kuda* (*n* = 10)	*H. trimaculatus* (*n* = 10)
Height (Ht)	48.2 ± 6.50^a	157.73 ± 19.24^b	93.06 ± 14.71^c
Standard Length (SL)	54.95 ± 5.48^a	182.65 ± 11.64^b	109.87 ± 6.45^c
Trunk Length (TrL)	16.45 ± 2.02^a	47.83 ± 2.54^b	26.40 ± 2.12^c
Tail Length (TaL)	30.15 ± 4.04^a	104.43 ± 9.07^b	61.70 ± 3.40^c
Head Length (HL)	8.83 ± 0.41^a	30.39 ± 1.88^b	21.70 ± 1.77^c
Snout Length (SnL)	2.67 ± 0.21^a	12.06 ± 1.04^b	8.90 ± 0.99^c
HL/SnL ratio	3.33 ± 0.14^a	2.54 ± 0.16^b	2.45 ± 0.19^{bc}
Snout Depth	1.95 ± 0.30^a	4.25 ± 0.41^b	3.45 ± 0.16^c
Head Depth	6.68 ± 0.79^a	14.80 ± 1.84^b	9.50 ± 0.53^c
TD4	3.83 ± 0.50^a	12.16 ± 1.19^b	6.75 ± 0.86^c
TD9	5.63 ± 0.63^a	18.68 ± 1.51^b	9.55 ± 0.90^c
Pectoral fin base length	2.03 ± 0.1^a	7.07 ± 1.2^b	4.75 ± 0.6^c
Dorsal fin base length	3.88 ± 0.3^a	15.73 ± 1.8^b	9.80 ± 0.8^c
Meristic features (numbers)			
Trunk Rings (TrR)	11 ± 0.00	11 ± 0.00	11 ± 0.00
Tail Rings (TaR)	38 ± 1.00^a	36 ± 1.00^b	41 ± 1.00^c
TrR supporting Dorsal Fin	2 ± 0.00	2 ± 0.00	2 ± 0.00
TaR supporting Dorsal Fin	1 ± 0.00	1 ± 0.00	1 ± 0.00
Cheek Spines	2 ± 0.00^a	1 ± 0.00^b	1 ± 0.00^c
Eye spines	2 ± 0.00^a	1 ± 0.00^b	1 ± 0.00^c
Dorsal fin rays	16 ± 1.00^a	18 ± 1.00^b	20 ± 1.00^c
Pectoral fin rays	14 ± 1.00^a	16 ± 1.00^b	17 ± 1.00^c

287 **Table 3. PCA analysis of seahorse species.** Loadings of components matrix and

288 contributions of each axis to total variance ranked as per their coefficients (eigen values $> \pm$

289 0.30 are included; values in parentheses indicate variance explained by each principal

290 component).

Morphological characters	Coefficients
Principal component 1 (81%)	
SL	0.334
TaL	0.333
TD4	0.329
TD9	0.327
HL	0.327
TrL	0.325
HD	0.324
SnL	0.318
SnD	0.314
Principal component 2 (16%)	
TaR	0.663
DF	0.655

308 **Fig 4. PCA analysis scores.** Scores of factor loadings plot of principal component analysis of

309 three species, *H. mohnikei*, *H. kuda* and *H. trimaculatus* based on morphological data.

Phylogenetic analysis and genetic distance

PCR for the mitochondrial COI and Cyt *b* genes resulted in 609 bp and 639 bp amplicons,

respectively. Phylogenetic trees constructed with sequences of COI and Cyt *b* genes of Goa

specimens with those of morphologically similar and most abundant seahorse species in the

Indo-Pacific region are depicted in Fig. 5 and Fig. 6, respectively. Sequences of COI

(GenBank accession number MK330041.1) and Cyt *b* (GenBank accession number

MK112274.2) genes of Goa specimens showed similarities of 100% with *H. mohnikei*

(GenBank accession number: GQ502157.1) and 93% with *H. mohnikei* (GenBank accession

number: EU179924.1), respectively. Based on maximum similarities, the genetic identity of

Goa specimens was confirmed as *H. mohnikei*. Genetic distance describes the kinship

distance between intraspecies and interspecies. Calculated pair-wise genetic distance (K2P)

values based on COI and Cyt *b* sequences of *H. mohnikei* from Goa and other localities

(Japan, Korea, China, Vietnam and Thailand) are presented in Table 4 and Table 5,

respectively. Relatively, *H. mohnikei* from Goa showed the least genetic distances of 6.5%

(COI) and 5.4% (Cyt *b*) with populations of *H. mohnikei* from Vietnam (COI) and Thailand

(Cyt *b*), respectively.

Fig 5. Phylogenetic relationships of Goa specimen with other seahorses species based on

COI gene sequences. Phylogenetic relationships of Goa specimen (MK330041.1) with

closely related species based on cytochrome oxidase sub-unit I gene (COI) constructed

through maximum likelihood method with bootstrap values (1000 replicates) as percentage at

the nodes. The alligator pipefish, *Syngnathoides biaculeatus* (KY066152.1) served as an

outgroup species. Scale bar = genetic distance of 0.05.

Fig. 6. Phylogenetic relationships of Goa specimen with other seahorses species based on

Cyt *b* gene sequences. Phylogenetic relationships of Goa specimen (MK112274.2) and

closely related seahorse species based on cytochrome *b* gene (Cyt *b*) constructed through

maximum likelihood method with bootstrap values (1000 replicates) as percentage at the

nodes. The alligator pipefish, *Syngnathoides biaculeatus* (EF189167.1) served as an out-

group species. Scale bar = genetic distance of 0.05.

Table 4. Pair-wise genetic distance based on COI gene sequences. Pair-wise genetic

distances (Kimura 2-parameter model) based on COI gene sequences of *Hippocampus*

mohnikei populations between Goa and the Indo-Pacific region. Values in parentheses

indicate percentage.

Localities of *H. mohnikei* with GenBank Accession number	1	2	3	4
1 Goa, India (MK330041.1)	0.000			
2 Japan (GQ502158.1)	0.085 (8.5)	0.000		
3 China (KX689046.1)	0.100 (10.0)	0.032 (3.2)	0.000	
4 Vietnam (GQ502157.1)	0.065 (6.5)	0.092 (9.2)	0.104 (10.4)	0.000

354 **Table 5. Pair-wise genetic distance based on Cyt *b* gene sequences.** Pair-wise genetic

355 distances (Kimura 2-parameter model) based on Cyt *b* gene sequences of *Hippocampus*

356 *mohnikei* populations between Goa and the Indo-Pacific region. Values in parentheses

357 indicate percentage.

Localities of *H. mohnikei* with GenBank Accession number	1	2	3	4	5
1 Goa, India (MK112274.2)	0.000				
2 South Korea (KT731897.1)	0.126 (12.6)	0.000			
3 China (KC527556.1)	0.128 (12.8)	0.006 (0.6)	0.000		
4 Japan (AF192689.1)	0.134 (13.4)	0.050 (5.0)	0.047 (4.70)	0.000	
5 Thailand (EU179924.1)	0.054 (5.4)	0.132 (13.2)	0.128 (12.8)	0.140 (14.0)	0.000

Ocean circulation

The climatological currents generated for the winter season (December to February) from

2007 to 2017 for the Indo-Pacific region are depicted in Fig. 7. As seen from the figure that

the circulation shows a large-scale flow moving from the East China Sea to the South China

Sea. From the South China Sea, the major part of the flow turns southeastward entering into

the Java Sea and a small part turns northwestward to eastern Bay of Bengal (BoB) and the

Andaman Sea through Malacca Strait. From eastern BoB, the surface waters flow westward

as a part of the north equatorial current (NEC). On encountering the coast of Sri Lanka, a part

of the NEC move northward into the northern BoB, while the rest moves westward and into

the southern Arabian Sea, south of Sri Lanka. The NEC feeds into the wintertime west India

coastal current (WICC) and moves waters from NEC towards north into the Arabian Sea

along its eastern boundary.

Fig. 7. Oceanic circulation in Indo-Pacific region. Climatological surface currents during

winter season (December–February) for the Indo-Pacific region. Red dots indicates the

geographic locations of distribution of *Hippocampus mohnikei.* (WICC= West Indian Coastal

Current, EICC= East India Coastal Current, NEC= North Equatorial Current and SCSWC=

South China Sea Warm Current).

Discussion

The distributional range of *Hippocampus mohnikei* is still unresolved as [14,15] confirmed

the nativity of this species to Japan and Vietnam and likely extending throughout the South

China Sea and the Gulf of Thailand [14,15,23]. Recent reports on the occurrence of *H.*

mohnikei from many other non-native locations in north- and south-eastern coastal waters of

Asia [10,35,41,51,56,57] suggest its significantly extended distribution in the Indo-Pacific

region (Fig. 8). The sighting of *H. mohnikei* in the Palk Bay, southeast coast of India [10] is

till date the known range limit towards northwestern Asian coastal waters. In southeastern

Asian coastal waters, *H. mohnikei* has been reported from Singapore [57] as well as from

Cambodia, Malaysia, Thailand and Vietnam [35]. Towards northeast Asian coastal waters

[56,58] reported *H. mohnikei* in southern Korea, whereas, [41,51] reported *H. mohnikei* from

China.

Fig. 8. Reported sightings of *H. mohnikei* in Indo-Pacific region. Map showing native

range, recent sightings and all reported sightings for *Hippocampus mohnikei* in the Indo-

Pacific region. Sightings are classified as new (this study, ★), recently published (2004–2016,

(▲) and original records [14] (*). Map was modified from Aylesworth et al. [35] and re-

created in the World Geodetic System (WGS) 84 datum.

The present report on the occurrence of *H. mohnikei* in a bay-estuarine system of Goa,

CWCI as confirmed by both morpho-molecular analysis is the most recent represents a

significant westward expansion compared to the previously known distributional range of this

species [35]. Since the reporting of *H. mohnikei* solely based on morphological characters

from the Palk Bay, southeast coast of India [10], no further sightings of this species have

been reported from coastal marine waters of India. Therefore, a comparison of genetically

distinct populations of *H. mohnikei* inhabiting native (Japan and Vietnam) and extended

ranges in the Indo-Pacific region (Fig. 8) with Goa specimens gains significance. Lourie et al.

[7,23] reported that a Vietnamese specimen of *H. mohnikei* was genetically distinct from *H.*

mohnikei from Japan. Results of phylogenetic constructed of sequences of COI (GenBank

accession number, MK330041.1) and Cyt *b* (GenBank accession number MK112274.2)

genes of Goa specimens showed maximum similarities of 100% with (GenBank accession

number GQ502157.1) and 93% (GenBank accession number EU179924.1) with *H. mohnikei*

from Vietnam and Thailand, respectively. Furthermore, a comparison of pair-wise genetic

distance values (K2P) based on sequences COI (Table 4) and Cyt *b* (Table 5) genes revealed

the conspecificity of Goa specimens with populations of *H. mohnikei* from Vietnam (K2P

value, 6.5%) and Thailand (K2P value, 5.4%) rather than Japan (K2P values, 8.5% and

13.4% for COI and Cyt *b* sequences, respectively). In contrast, high pair-wise K2P values of

sequences of Cyt *b* between *H. mohnikei* populations from the Gulf of Thailand and those

from China (13.54−14.46%) and Japan (14.42−14.79%) have been recently reported [59]. A

comparative assessment of K2P values of the present study with those from the Gulf of

Thailand [59] suggest that *H. mohnikei* populations from Goa and the Gulf of Thailand might

be genetically distinct from China and Japan. Also, a possible genetic differentiation in

populations of *H. mohnikei* inhabiting coastal marine waters of northeastern Asia (Japan,

Korea and China) and southeastern Asia (Vietnam, Thailand, Cambodia, Malaysia, Singapore

and India) may not be ruled out. Furthermore, least K2P values between disjunct populations

of *H. mohnikei* (Goa vs Vietnam, COI gene) and (Goa vs. Thailand, Cyt *b* gene) may reflect

on the possibility of cryptic species within *H. mohnikei* populations [7,23,59] which needs

further confirmation.

Recent studies have advanced our understanding on habitat associations of *H. mohnikei*

compared to what has been described in the IUCN Red List Assessment [42]. Previously, *H.*

mohnikei was reported only in association with seagrass beds in Japan and in an estuary in

Vietnam [15,23]. Association of *H. mohnikei* populations with seagrass beds (*Zostera* spp.) in

Japan [58,60], Korea [56] and China [41] has been reported. Whereas the specimen reported

from the Palk Bay [10] was associated with two other additional genera of seagrass

(*Halophila* and *Thalassia*). Recently, published report on the habitat preferences of *H.*

mohnikei from Southeast Asia [35] indicate its association with seagrass beds of *Cymodocea*

spp., *Enhalus* spp., *Halodule* spp., *Halophila* spp., *Syringodium* spp. and *Thalassia* spp.

Furthermore, these authors also report the association of *H. mohnikei* with mangroves, oyster

beds, mussel farms and sandy beaches, whereas, [57] has reported the its occurrence in

association with seaweeds in Singapore. From these studies, it is evident that *H. mohnikei*

occupies a wide variety of habitats.

All previous reports on the habitat descriptions of *H. mohnikei* were located in the coastal

waters and bay-estuarine systems in the depth range of 0–10 m [10,15,23,35,41,56].

Interestingly, the occurrence of small juveniles of *H. mohnikei* in association with highly

dense live prey organisms in the offshore waters of Japan has also been reported [37]. The

present report of *H. mohnikei* is based on specimens caught as an incidental catch in bag nets

fixed to the stakes operated in the Mandovi estuary, Goa. Although precise information on

the habitat of *H. mohnikei,* could not be ascertained, it is worth describing the surrounding

environs at the site of collection. Predominantly, the estuarine stretches of the River Mandovi

have dense mangrove coverage and extensive oyster beds [61] and patches of seagrass,

Halophila ovalis and *H. beccarii* [62]. The channel network of Mandovi estuary is

characterised by islands (Chorao, Divar and Vanxim) and many sheltered areas interspersed

with dense mangrove ecosystem. The present occurrence of *H. mohnikei* in a bay-estuarine

system of Goa is not quite surprising as these are the typical habitats where this species has

been previously reported. For ascertaining habitat shift by disjunct populations of *H.*

mohnikei. further intense studies on habitat specificities of different life stages therefore, are

warranted.

Food availability has been considered as one of the factors influencing the abundance and

distribution of seahorses [19]. Zooplankton such as *Oithona davisae*, Calanoida and *Penilia*

avirostris have been identified as the main prey organisms for juvenile and young *H.*

mohnikei in Tokyo Bay [37], whereas the adults have been reported to prey upon mainly

larger organisms such as gammarid and caprellid amphipods [58]. Nutrient-rich waters of the

Mandovi estuarine system sustain high primary productivity and abundance of zooplanktonic

prey organisms such as copepods [63−65] which are essential for the survival and

establishment of seahorse populations. Considering the prevailing environmental conditions,

mangrove shelter and abundance of prey organisms, the likelihood of establishment and

colonization of seahorse populations in these sheltered areas is quite high.

Factors such as habitat, diet, and even anthropogenic activities are the main reasons cited

for the migration of seahorses [66]. Furthermore, instances of seahorses for long distance

dispersal by means of rafting along with seaweeds or any other suitable holdfast under the

influence of ocean currents during juvenile stages have been documented [2,20]. Southward

migration of *H. histrix* ~1800 km under the influence of East Australian Current in the Great

Barrier Reef has been reported [31]. Furthermore, ~1100 km migration from the south to the

northeast coast of India by *H. fuscus* under the influence of East Indian Coastal Current has

been also reported [33]. Direct evidence for passive long-distance migration of *H. trimaculatus* at Malacca Strait has been documented recently [34].

Newborn juveniles of many seahorse species undergo a pelagic phase (planktonic) ranging between two and six weeks depending on the species before they attain a settlement phase (benthic) [2,67]. Consequently, planktonic phase juveniles may facilitate in widespread gene flow across geographically disjunct populations and resulting in genetic homogeneity [68]. The occurrence of small juveniles of *H. mohnikei* (15.1−45.5 mm) in association with highly dense live prey organisms in offshore waters of Japan [37] suggests that *H. mohnikei* might undergo a prolonged planktonic phase. Furthermore, a seasonal (warm and cold season) inshore-offshore migration by *H. mohnikei* populations inhabiting Chinese coast has been confirmed [51]. Higher dispersal capabilities of seahorse juveniles with floating debris, prey swirls and with seaweed rafts than adults has been reported [2,28,34]. It has been postulated that marine species with greater longevity, free swimming and feeding (planktotrophic) larval phases have relatively a high dispersal abilities and thus promoting genetic exchange between populations [69]. A relatively prolonged planktonic phase as in case of *H. mohnikei* may facilitate the high degree of dispersal under the influence of oceanic circulation.

The confirmed occurrence of four sub-adult specimens (Mandovi estuary) and suspected occurrence of an adult specimen of *H. mohnikei* in the proximately adjacent bay-estuarine ecosystem (Chapora estuary) of Goa, CWCI could be as a consequence of either long-distance dispersal from Southeast Asian countries or stepping-stone mediated dispersal as revealed by oceanic currents circulation in the Indo-Pacific region. It is clearly seen from Fig. 7 that water mass from southeastern Asian coasts and east coast of India enters into the Arabian Sea as a result of winter circulation [70]. In the absence of geographical barriers between *H. mohnikei* populations of Goa and Southeast Asian countries, ocean currents

flowing from regions where sightings of *H. mohnikei* may contribute to wide dispersal.

Therefore, it is plausible that oceanic currents may be aiding long-distance or stepping stone

dispersal of *H. mohnikei* populations from different geographical locations such as

Cambodia, Malaysia, Thailand and Vietnam [14,15,35,59], Singapore [57] and the Palk Bay,

India [10] to the west coast of India.

Conversation with local fishermen who operate stake nets at the collection site revealed

that sightings of seahorses are relatively more common during summer monsoon season than

in non-monsoonal months. During the summer monsoon period, a number of rivulets and

rivers connected to the Mandovi estuarine network discharge large amounts of freshwater

[71]. Consequently, it may be possible that the rainwater discharge from sheltered areas (the

preferred seahorse habitats) may entail seahorses to flow downstream along with currents due

to their poor swimming abilities. It is well known that the west coast of India comes under the

influence of substantial coastal cold-water upwelling event during the summer monsoon

season [45,71]. It has been reported that the survival and dispersal of tropical marine species

are known to be impacted by a drastic decrease in water temperature associated with

upwelling [72−74]. Consequently, seahorse populations in western coastal India might

undergo seasonal fluctuations [20]. A significant decrement in abundance and diversity of

demersal fishes during the summer monsoon in a bay-estuarine system of the Mandovi river

has been documented [47]. As a consequence of the reduction in salinity and drop in

temperature associated with monsoonal precipitation and upwelling, respectively the seahorse

populations might face limitations in growth and recruitment along the west coast of India.

Interestingly, unlike most subtropical seahorse species, the greater tolerance by native

populations of *H. mohnikei* to cold water inhabiting in the Western Pacific has been reported

[29]. However, further temperature stress simulation studies are required to ascertain whether

translocated populations of *H. mohnikei* also possess a similar trait or not.

Conclusions

Recognising the potential of seahorses as a popular flagship species in estuarine conservation and sustainable management [1,75,76], significant westward range expansion of *H. mohnikei* is of considerable biogeographic and conservation importance. Further intense sampling and analysis of *H. mohnikei* populations along the Indian coast are therefore required to confirm whether the present occurrence of *H. mohnikei* in a bay-estuarine system, CWCI represents (i) an isolated translocation or (ii) naturally occurring facilitated by oceanic circulation or (iii) the first sighting of an unreported of fairly established populations of this vulnerable seahorse species.

Acknowledgements

The authors are grateful to the Director, CSIR-National Institute of Oceanography, Goa (India) for encouragement and facilities. Thanks are due to local fishermen Mr. Harish Halarnkar for providing seahorse specimens. Thanks are also due to Dr. Mani Murali and Mr. Devanand Kavlekar for their help in preparing the maps. This represents contribution No. xxxx of the CSIR-National Institute of Oceanography, Goa (India).

Author Contributions

Conceived and designed the experiments: SVS RAS RDSK: Performed the experiments: SVS RAS RDSK MSS HTK. Analyzed the data: SVS RAS MSS RDSK JN. Contributed reagents/materials/analysis tools: JN HTK. Wrote the paper: RAS SVS RDSK.

References

1. Sreepada RA, Desai UM, Naik S. The plight of Indian seahorses: Need for conservation

and management. Curr Sci 2002; 82: 377–378.

2. Foster SJ, Vincent ACJ. Life history and ecology of seahorses: implications for

conservation and management. J Fish Biol 2004; 65: 1–61.

3. Kuiter RH. Seahorses and Their Relatives. Aquatic Photographics, Seaford, Australia;

2009.

4. Froese R, Pauly D. FishBase.www.fishbase.org. Downloaded on10/2018.

5. Harasti D, Martin-Smith K, Gladstone W. Ontogenetic and sex-based differences in habitat

preferences and site fidelity of White's seahorse *Hippocampus whitei*. J Fish Biol 2014;

85: 1413–1428.

6. Hellyer C, Harasti D, Poore A. Manipulating artificial habitats to benefit seahorses in

Sydney Harbour, Australia. Aquat Conserv Mar Fresh Ecosyst 2011; 21: 582–589.

7. Lourie SA, Pollam RA, Foster SJ. A global revision of the seahorses *Hippocampus*

Rafinesque 1810 (Actinopterygii: Syngnathiformes): taxonomy and biogeography with

recommendations for further research. Zootaxa 2016; 4146: 1–66.

8. Salin KR, Mohankumaran NC. Resources and biodiversity of seahorses and the need for

their conservation in India. Aquacult Asia 2006; 10: 3–8.

9. Murugan A, Dhanya S, Rajagopal S, Balasubramanian T. Seahorses and pipefishes of the

Tamil Nadu coast. Curr Sci 2008; 95: 253–260.

10. Thangaraj M, Lipton AP. Occurrence of the Japanese seahorse *Hippocampus mohnikei*

Bleeker 1854 from the Palk Bay coast of southeastern India. J Fish Biol 2007; 70: 310–

312.

11. Krishnan TS, Kuiter RH, Patterson J. Occurrence of western Indian Ocean seahorse,

Hippocampus borboniensis Dumeril, 1870, in the Gulf of Mannar, Southeastern India. Ind

J Geo-Mar Sci 2011; 40: 642–644.

12. Yogeshkumar JS, Geetha S. A new species of *Hippocampus montebelloensis* (Family:

Syngnathidae) from the southeast coast of India. Int J Plant Anim Env Sci 2012; 2: 56–

58.

13. Subburaman S, Murugan A, Goutham S, Kaul R, Prem Jyothi PVR, Balasubramanian T.

First distributional record of giraffe seahorse, *Hippocampus camelopardalis* Bianconi

1854 (Family Syngnathidae) from Gulf of Kachchh waters, north west coast of India.

Indian J Geo-Mar Sci 2014; 43: 408–411.

14. Lourie SA, Foster SJ, Cooper EWT, Vincent ACJ. A Guide to the Identification of

Seahorses. Project Seahorse and TRAFFIC North America; 2004.

15. Lourie SA, Vincent ACJ, Hall HJ. Seahorses: an identification guide to the world's

species and their conservation. Project seahorse, London, UK; 1999.

16. Silveira RB, Siccha-Ramirez R, Silva JRS, Oliveira C. Morphological and molecular

evidence for the occurrence of three *Hippocampus* species (Teleostei: Syngnathidae) in

Brazil. Zootaxa 2014; 3861: 317–332.

17. Casey SP, Hall HJ, Stanley HF, Vincent ACJ. The origin and evolution of seahorses

(genus *Hippocampus*): a phylogenetic study using the cytochrome b gene of

mitochondrial DNA. Mol Phylogenet Evol 2004; 30: 261–272.

18. Wang X, Han X, Zhang Y, Liu S, Lin Q. Phylogenetic analysis and genetic structure of

the seahorse, *Hippocampus fuscus* from the Arabian and Red Sea based on mitochondrial

DNA sequences. Mitochondrial DNA A 2019; 30: 165–171.

19. Lourie SA, Green DM, Vincent ACJ. Dispersal, habitat differences and comparative

phylogeography of South East Asian seahorses (Syngnathidae: *Hippocampus*). Mol Ecol

2005; 14: 1073–1094.

20. Teske PR, Hamilton H, Palsboll PJ, Choo CK, Gabr H, Lourie SA. et al. Molecular

evidence for long-distance colonization in an Indo-Pacific seahorse lineage. Mar Ecol

Prog Ser 2005; 286: 249–260.

21. Boehm JT, Waldman J, Robinson JD, Hickerson MJ. Population genomics reveals

seahorses (*Hippocampus erectus*) of the western mid-Atlantic coast to be residents rather

than   vagrants.   PloS   One   2015;   10:   e0116219.

http://dx.doi.org/10.1371/journal.pone.0116219.

22. Saarman NP, Louie KD, Hamilton H. Genetic differentiation across eastern Pacific

oceanographic barriers in the threatened seahorse *Hippocampus ingens*. Conserv Genet

2010; 11: 1989–2000.

23. Lourie SA, Pritchard JC, Casey SP, Ky TS, Hall HJ. et al. The taxonomy of Vietnam's

exploited seahorses (family Syngnathidae). Biol J Linn Soc 1999; 66: 231–256.

24. Woodall LC, Otero-Ferrer F, Correia M, Curtis JMR, Garrick Maidment N. et al. A

synthesis of European seahorse taxonomy, population structure, and habitat use as a basis

for assessment, monitoring and conservation. Mar Biol 2018; 165: 1–19.

25. Curtis JMR, Vincent ACJ. Life history of an unusual marine fish: survival, growth and

movement patterns of *Hippocampus guttulatus* Cuvier 1829. J Fish Biol 2006; 68: 707–

733.

26. Caldwell I, Vincent ACJ. A sedentary fish on the move: effects of displacement on long-

snouted seahorse (*Hippocampus guttulatus* Cuvier) movement and habitat use. Environ

Biol Fish 2013; 96: 67–75.

27. Correia M, Caldwell IR, Koldewey HJ, Andrade JP, Palma J. Seahorse (Hippocampinae)

population fluctuations in the Ria Formosa Lagoon, south Portugal. J Fish Biol 2015; 87:

679– 690.

28. Luzzatto DC, Estalles ML, Diaz De Astarloa JM. Rafting seahorses: the presence of

juvenile *Hippocampus patagonicus* in floating debris. J Fish Biol 2013; 83: 677–681.

29. Han SY, Kim JK, Tashiro F, Kai Y, Yoo JT. Relative importance of ocean currents and

fronts in population structures of marine fish: a lesson from the cryptic lineages of the

Hippocampus mohnikei complex. Mar Biodiv 2019; 49: 263–275.

30. Luzzatto DC, Pujol MG, Figueroa D, Diaz De Astarloa JM. The presence of the seahorse

Hippocampus patagonicus in the deep waters: additional evidence of the dispersive

capacity of the species. Mar Biodivers Rec 2014; 7: e71.

31. Harasti D. Range extension and first occurrence of the thorny seahorse *Hippocampus*

histrix in New South Wales, Australia. Mar. Biodivers Rec 2015; 8:

https://doi.org/10.1017/S1755267215000263.

32. Harasti D. Southward range extension of the great seahorse (*Hippocampus kelloggi*

Jordan and Snyder, 1901) in Australia. J Appl Ichthyol 2017; 1–3

https://doi.org/10.1111/jai.13414.

33. Mahapatro D, Mishra RK, Panda S. Range extension of a vulnerable Sea horse

Hippocampus fuscus (Actinopterygii: Syngnathidae) on the north-eastern Bay of Bengal

coast. Mar Biodivers Rec 2017; 10:6. DOI 10.1186/s41200-017-0108-z.

34. Wong NLWS, The JC. Direct observation of drifting juvenile seahorse *Hippocampus*

trimaculatus (Leach, 1814) beneath the offshore surface at Malacca Strait. Check List

2017; 13: 993–996.

35. Aylesworth L, Lawson LM, Laksanawimol P, Ferber P, Loh TL. New records of the

Japanese seahorse *Hippocampus mohnikei* in Southeast Asia lead to updates in range,

habitat and threats. J Fish Biol 2016; 88: 1620–1630.

36. Masuda H, Amaoka K, Araga C, Uyeno T, Yoshino T. The Fishes of the Japanese

Archipelago. Tokyo: Tokai University Press; 1984.

37. Kanou K, Kohno H. Early life history of a seahorse, *Hippocampus mohnikei,* in Tokyo

Bay Japan. Ichthyol Res 2001; 48: 361–368.

38. Suda Y, Inoue T, Nakamura M, Masuda N, Doi H, Murai T. Nearshore ichthyofauna in

the intermediate sandy beach, Doigahama Beach, Yamaguchi Prefecture, Japan. J Nat

Fish Univ 2004; 52: 11–29.

39. Laksanawimol P, Petpiroon S, Damrongphol P. Trade of seahorses, *Hippocampus* spp.

(Actinopertygii: Syngnathiformes: Syngnathidae), on the East Coast of the Gulf of

Thailand. Acta Ichthyol Piscata 2013; 3: 229–235.

40. Chang CH, Jang-Liaw NH, Lin YS, Fang YC, Shao KT. Authenticating the use of dried

seahorses in the traditional Chinese medicine market in Taiwan using molecular

forensics. J Food Drug Analy 2013; 21: 310–316.

41. Zhang Y, Pham NK, Zhang H, Lin J, Lin Q. Genetic variations in two seahorse species

(*Hippocampus mohnikei* and *Hippocampus trimaculatus)*: evidence for Middle

Pleistocene population expansion. PloS One 2014; 9: e105494.

42. Pollom R. *Hippocampus mohnikei*. The IUCN Red List of Threatened Species 2017:

http://dx.doi.org/10.2305/IUCN.UK.20173.RLTS.T41005A54907304.en43.   Downloaded

on 10 November 2018.

43. Shetye SR, Gouveia AD, Singbal SY, Naik CG, Sundar D. et al. Propagation of tides in

the Mandovi–Zuari estuarine network. Proc Indian Acad Sci (Earth Planet Sci) 1995; 104:

667–682

44. Qasim SZ, Sen Gupta R. Environmental characteristics of the Mandovi-Zuari estuarine

system. Estuar Coast Shelf Sci 1981; 13: 557–578.

45. Madhupratap M, Nair KNV, Gopalkrishnan TC, Haridas P, Nair KKC. et al. Arabian sea

oceanography and fisheries of the west coast of India. Curr Sci 2001; 81: 355–361.

46. Ansari ZA, Sreepada RA, Dalal SG, Ingole BS, Chatterji A. Environmental influences on

the trawl catches in a bay-estuarine system of Goa, west coast of India. Estuar Coast Shelf

Sci 2003; 56: 503–515.

47. Ansari ZA, Chatterji A, Ingole BS, Sreepada RA, Rivonker CU. et al. Structure of inshore

demersal fish community at Goa. Estuar Coast Shelf Sci 1995; 41: 593–610.

48. Lourie S. Measuring seahorses. Project Seahorse Technical Report No.4, Version 1.0.

Project Seahorse, Fisheries Centre, University of British Columbia; 2003.

49. Underwood AJ. Experiments in Ecology: their logical and interpretation using Analysis

of Variance. Cambridge University Press. Cambridge, UK; 1997.

50. Kumar R Singh PJ, Nagpure NS, Kushwaha B, Srivastava SK. et al. A non-invasive

technique for rapid extraction of DNA from fish scales. Ind J Exp Biol 2007; 45: 992–

997.

51. Qin G, Zhang Y, Ho ALFC, Zhang Y, Lin Q. Seasonal distribution and reproductive

strategy of seahorse. ICES J Mar Sci 2017; 74: 2170–2179.

52. Lourie SA, Vincent ACJ. A marine fish follows Wallace's line: the phylogeography of

the three-spot seahorse (*Hippocampus trimaculatus*, Syngnathidae, Teleostei) in

Southeast Asia. J Biogeogr 2004; 31: 1975–1985.

53. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search

tool. J Mol Biol 1990; 215: 403–410.

54. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis,

version 7.0 for bigger datasets. Mol Biol Evol 2016; 33: 1870–1874.

55. Kimura M. A simple method for estimating evolutionary rates of base substitutions

through comparative studies of nucleotide sequences. J Mol Evol 1980; 16: 111–120.

56. Choi Y, Rho S, Park H, Kang D. Population characteristics of two seahorses, *Hippocampus coronatus* and *Hippocampus mohnikei*, around seagrass beds in the southern coastal waters of Korea. Ichthyol Res 2012; 59: 235–241.

57. Lim K. New Singapore record of the seahorse, *Hippocampus mohnikei*. Singapore Biodiver Rec 2013; 2013: 68–69.

58. Kwak S, Huh S, Kim H. Feeding habits of *Hippocampus mohnikei* in an eelgrass (*Zostera marina*) bed. Korean J Ichthyol 2008; 20: 112–116.

59. Panithanarak T, Karuwanjaroen R. A preliminary study of genetic differentiation in the Japanese seahorse (*Hippocampus mohnikei*) populations: a possibility of cryptic species. Burapha Sci J 2019; 24: 124–137.

60. Otsuka Y, Suzuki H, Akagawa I. Occurrence, gonad morphology and maturity of Japanese seahorse *Hippocampus mohnikei* in Matsushima Bay, Japan. J School Mar Sci Tech Tokai Univ 2009; 7: 11–22.

61. Wafar S. Ecology of mangroves along the estuaries of Goa; Ph. D. Thesis, Karnataka University, Dharward, India; 1987.

62. Jagtap TG. Distribution of seagrasses along Indian coast. Aquac Bot 1991; 40: 379–386.

63. Padmavati G, Goswami SC. Zooplankton ecology in the Mandovi-Zuari estuarine system of Goa, west coast of India. Ind J Mar Sci 1996; 25: 268–273.

64. Dalal SG, Goswami SC. Temporal and ephemeral variations in copepod community in the estuaries of Mandovi and Zuari– west coast of India. J Plankton Res 2001; 23: 19–26.

65. Das PB. Zooplankton assemblages in spatial stretches of a tropical estuary. Int J Res Appl Sci Engg Technol 2017; 5: 1789–1799.

66. Qin G, Johnson C, Zhang Y, Zhang H, Yin J, Miller G. et al. Temperature-induced physiological stress and reproductive characteristics of the migratory seahorse

Hippocampus erectus during a thermal stress simulation. Biol Open 2018; 25:

7(6).bio032888.

67. Scales H. Advances in the ecology, biogeography and conservation of seahorses (genus

Hippocampus). Prog Phys Geog 2010; 34: 443–458.

68. Bohonak AJ. Dispersal, gene flow, and population structure. Q Rev Biol 1999; 74: 21–45.

69. Palumbi SR. Genetic divergence, reproductive isolation, and marine speciation. Annu.

Rev Ecol Evol S 1994; 25: 547–572.

70. Wyrtki K. Physical oceanography of the southeast Asian waters. Scientific results of

maritime investigations of the South China Sea and the Gulf of Thailand 1959–1961. Naga

Report, 2. Scripps Institution of Oceanography, La Jolla, CA; 1961.

71. Shetye SR, Gouveia AD, Shenoi SSC, Michael GS, Sundar D, Almedia AM. et al. The

coastal current off western India during the northeast monsoon. Deep-Sea Res 1991; 38:

1517–1529.

72. Fleminger A. The Pleistocene equatorial barrier between the Indian and Pacific oceans

and a likely cause for Wallace's Line. In Pierrot-Bults AC, van der Spoel S, Zahuranec

BJ, Johnson RK editors, Pelagic marine science. UNESCO Tech Pap Mar Sci 1986; 49:

84–97.

73. Maree RC, Whitfield AK, Booth AJ. Effect of water temperature on the biogeography of

South African estuarine fishes associated with the subtropical/warm temperate subtraction.

S Afr J Sci 2000; 96: 184–188.

74. Bowen BW, Bass AL, Rocha LA, Grant WS, Robertson DR. Phylogeography of the

trumpetfishes *(Aulostomus)*: ring species complex on a global scale. Evolution 2001; 55:

1029–1039.

75. Martin-Smith KM, Vincent ACJ. Seahorse declines in the Derwent estuary, Tasmania, in

the absence of fishing pressure. Biol Conserv 2005; 123: 533–545.

76. Shokri MR, Gladstone W, Jelbart J. The effectiveness of seahorses and pipefish (Pisces:

Syngnathidae) as a flagship group to evaluate the conservation value of estuarine seagrass

beds. Aquat Conserv 2009; 19: 588–595.

Supporting Information

S1 Fig. Picture of *H. mohnikei* posted on social media. Male specimen of *Hippocampus*

mohnikei caught by gill net in the Chapora estuary, Goa, India