CRISPR-Cas9 modified bacteriophage for treatment of *Staphylococcus aureus* induced osteomyelitis and soft tissue infection

Leah K. Horstemeyer, [1] JooYoun Park,[2] Elizabeth Swanson,[3] Mary C. Beard, [1] Emily M. McCabe, [1] Anna Rourke,[1] Keun Seok Seo,[2] Lauren B. Priddy[1]*

[1]Department of Agricultural and Biological Engineering, Mississippi State University, Mississippi State, MS, USA

[2]Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University, Mississippi State, MS, USA

[3]Department of Clinical Sciences, College of Veterinary Medicine, Mississippi State University, Mississippi State, MS, USA

*corresponding author

Email: lbpriddy@abe.msstate.edu, (662) 325-5988, Department of Agricultural and Biological Engineering, Mississippi State University, 130 Creelman Street, Mississippi State, MS, USA 39762

Abstract

Osteomyelitis, or bone infection, is often induced by antibiotic resistant *Staphylococcus aureus* strains of bacteria. Although debridement and long-term administration of antibiotics are the gold standard for osteomyelitis treatment, the increase in prevalence of antibiotic resistant bacterial strains limits the ability of clinicians to effectively treat infection. Bacteriophages (phages), viruses that effectively lyse bacteria, have gained recent attention for their high specificity, non-toxicity, and the low likelihood of resistance development by pathogens. Previously, we have shown that CRISPR-Cas9 genomic editing techniques could be utilized to expand bacteriophage host range and enhance bactericidal activity through modification of the tail fiber protein, as well as improve safety with removal of major virulence genes. In a dermal infection study, these CRISPR-Cas9 phages reduced bacterial load relative to unmodified phage. Thus, we hypothesized this bacteriophage would be effective to mitigate infection from a biofilm forming *S. aureus* strain *in vitro* and *in vivo*. *In vitro*, qualitative fluorescent imaging demonstrated superiority of phage to conventional vancomycin and fosfomycin antibiotics against *S. aureus* biofilm. Quantitative antibiofilm effects increased over time for fosfomycin, phage, and fosfomycin-phage (dual) therapeutics delivered via alginate hydrogel. We developed an *in vivo* rat model of osteomyelitis and soft tissue infection that was reproducible and challenging and enabled longitudinal monitoring of infection progression. Using this model, phage (with and without fosfomycin) delivered via alginate hydrogel were successful in reducing soft tissue infection but not bone infection, based on bacteriological, histological, and scanning electron microscopy analyses. Notably, the efficacy of phage at mitigating soft tissue infection was equal to that of high dose fosfomycin. Future research may

utilize this model as a platform for evaluation of therapeutic type and dose, and alternate delivery

vehicles for osteomyelitis mitigation.

Introduction

For nearly a century, antibiotics have been a vital resource utilized by clinicians to

eliminate infection, with nearly 270 million prescriptions dispensed in 2015 alone.[1] Antibiotics

are utilized for a variety of infections, from common otitis externa ("swimmers ear") to severe

endocarditis, pneumonia, meningitis or osteomyelitis. Although antibiotics are typically able to

clear infection, antibiotic resistant strains of bacteria continue to emerge. It is not as lucrative,

nor as feasible, for pharmaceutical companies to develop novel antibiotics at the rates that these

multi-drug resistant (MDR) bacterial strains are isolated. Nationally, approximately $2.2 billion

is spent annually to treat MDR bacterial infections.[2] By 2050, it is estimated that nearly 10

million people could die each year due to resistant strains of bacteria.[3]

Staphylococcus aureus (*S. aureus*), a gram-positive bacterial strain, is one of the most

commonly isolated and arguably one of the most detrimental pathogens with antibiotic

resistance. One of the most common antibiotic resistant strains of *S. aureus* is methicillin-

resistant *S. aureus* (MRSA). MRSA alone was responsible for over 80,000 reported infections in

2011 alone, of which 11,285 resulted in death.[4] *S. aureus* is able to achieve antibiotic

resistance with genomic changes such as altered synthesis of peptidoglycan, a major component

of the bacterial cell wall. Additionally, some strains of *S. aureus* can produce biofilms, an

extracellular polymeric matrix including dead bacterial cells, which surrounds and protects the

living, underlying layer of *S. aureus*.[5] These biofilms can be difficult to penetrate, and

oftentimes require surgical intervention to remove.

Difficulties in treating osteomyelitis, or the infection of bone, have been exacerbated by

the rise of antibiotic resistant bacterial strains, particularly *S. aureus* strains, which are the most

common cause of bone infection.[6] Of diabetic foot ulcers, which occur in 25% of diabetic

patients, approximately 20% will spread to nearby bony hosts and result in osteomyelitis.[7] As

diabetic diagnoses continue to increase in the United States with an expected 55 million to be

afflicted by 2030, osteomyelitis infections will be an ongoing challenge for the healthcare

community.[8] It is essential that new therapeutics be engineered and tested, for rapid translation

into clinical use.

Bacteriophages (phages), or viruses that kill their bacterial hosts, are one class of

therapeutics that have gained attention in recent years due to their high specificity, non-toxicity,

and abundancy in nature.[9,10] Phages have been used for decades in Eastern Europe but have

not yet been adopted in the United States or other countries. This may be due to public concern

regarding elective viral use, issues concerning commercial phage production, and/or the ability to

fund and validate clinical trials.[11] Nonetheless, the potential benefits of this treatment have

been indicated by results of clinical trials of phages for treating diabetic foot ulcers, chronic

otitis, and urinary tract infections[11–13]. In April 2019, data from clinical trials were published

from Sydney, Australia, where intravenous (IV) administration of phage was utilized for

Staphylococcus infection treatment. Marked reduction of *staphylococci* with no adverse events

were reported.[14] In the United States as of January 2019, IV administration of phage for

ventricular assist device infection treatment received approval for phase I/II clinical trials.[15]

Collectively, these clinical trials demonstrate the efficacy of bacteriophage therapeutics and

suggest their potential utility against MDR bacterial strains.

Hydrogels are a commonly used, easily tailored delivery vehicle for therapeutics for a

wide variety of ailments, including osteomyelitis[6,16,17]. Alginate hydrogels are injectable,

well characterized, and biocompatible.[18,19] Furthermore, bacteriophages have been

successfully delivered to sites of infection with various hydrogel-based delivery systems in

previous studies.[16,17,20]

Although the high specificity of phages can be beneficial for treating a known, single

species, specificity of these viruses can make polymicrobial infection mitigation challenging. In

the clinical scenario, it is ideal for health care providers to administer one broad-spectrum drug

immediately upon patient presentation, rather than spend time identifying the causative agents of

infection. Previously, we have utilized CRISPR-Cas9 to modify temperate bacteriophage, which

effectively removed major virulence genes and expanded host range via modifications to the tail

fiber protein (which codes host specificity). *In vitro* testing revealed the improvements of

bacteriophage bactericidal activity due to this CRISPR-Cas9 system.[21]Within 6h of treatment,

the CRISPR-Cas9 phage effectively killed 1×10^5 CFU *S. aureus* culture. With native,

unmodified phage treatment, the culture was found to increase to approximately 1×10^9 CFU.

Similar effects were noted in an *in vivo* dermal infection study, where CRISPR-Cas9 phage

treatment resulted in nearly complete mitigation of dermal infections (~1 log CFU/g tissue),

while treatment with unmodified phage resulted in a significantly higher bacterial load (~3.5 log

CFU/g tissue).[21]

The objectives of the present work were: (i) to develop a green fluorescent protein (GFP)

integrated *S. aureus* strain (ATCC 6538-GFP), modify bacteriophage using CRISPR-Cas9, and

evaluate the bactericidal efficacy of our CRISPR-Cas9 modified bacteriophage *in vitro*,

compared to conventional antibiotics, and (ii) to develop an *in vivo* model of osteomyelitis and

soft tissue infection using this biofilm forming *S. aureus* strain, and use it to assess the

antimicrobial effects of bacteriophage, antibiotic, and dual bacteriophage-antibiotic therapies via

histological, radiographic, and bacteriological analyses. Our hypothesis was that CRISPR-Cas9

modified bacteriophage would be effective against *S. aureus* infection *in vitro* and in the femur

and contiguous soft tissue *in vivo*.

Materials and Methods

Bacterial Strain(s) and Culture

For a stable quantification of biofilm, *S. aureus* strain ATCC 6538 was genetically

modified to contain chromosomally integrated green fluorescent protein (GFP), as previously

described.[22] Briefly, *S. aureus* strain ATCC 6538 was transduced with a temperature sensitive

plasmid pTH100 harboring the GFP gene by electroporation and cultured in a brain heart

infusion (BHI) agar plate supplemented with chloramphenicol (BHI-CM) at 30°C, a plasmid

replication permissive temperature. To promote the first homologous recombination and cure

pTH100, a single colony grown in a BHI-CM plate was transferred to a fresh BHI-CM plate and

cultured at 42°C, a plasmid replication non-permissive temperature. To promote the second

homologous recombination, which removed the plasmid and resulted in a loss of

chloramphenicol resistance but maintained the GFP phenotype, a single colony was inoculated

into BHI broth and cultured at 37°C overnight. A serial dilution of culture was inoculated onto a

BHI plate and incubated at 37°C overnight. A GFP positive single colony checked by ultraviolet

lamp was randomly selected and streaked onto BHI and BHI-CM. A colony that was both GFP

positive and sensitive to chloramphenicol, indicating the integration of the GFP gene into the

chromosome and removal of plasmid, was selected for experiments (ATCC 6538-GFP).

Preparation of Alginate Hydrogels

All alginate gels were initially prepared at a 3% (w/v) concentration, for ultimate dilution to 2% after loading them with therapeutic. A 3% alginate mixture (w/v) was made with alginic acid powder (Sigma-Aldrich) and alpha Minimum Essential Medium (αMEM, Gibco) then left overnight at room temperature. This solution was sterile filtered (0.2 μm, Pall) and transferred into 1mL syringes. Therapeutics were then added directly to the alginate. The crosslinker, calcium sulfate (0.21g $CaSO_4$ / mL distilled H_2O) was loaded into a separate 1mL syringe and was mixed vigorously with the alginate solution for approximately one minute. Hydrogels were kept at 4°C or on ice until use.

Synthesis of CRISPR-Cas9 Modified Bacteriophages

S. aureus strain RF122 harboring CRISPR-Cas9 modified bacteriophage was cultured in BHI broth to the mid-exponential phase (OD600 at 0.3).[21] To induce CRISPR-Cas9 modified bacteriophage, mitomycin C (1 μg/mL, Sigma-Aldrich) was added to the culture and further incubated at 30°C with shaking at 80 RPM. A complete lysis of culture typically occurred within 2-3 hours. The clear lysate was sterilized with syringe filers (0.22 μm, Nalgene). The concentration of phage was calculated by determining the plaque-forming units using a soft agar (0.5%, w/v) overlaying method.[21]

Kirby-Bauer Analyses

To analyze the bactericidal activity of therapeutics, a Kirby-Bauer assay was performed as previously described, with slight modifications.[23] Stock solutions of fosfomycin (50 mg/mL) and phage (~10MOI/mL) were prepared in phosphate buffered saline (PBS). Using these stock solutions, a total of 10μL of: (i) fosfomycin, (ii) phage, (iii) dual: fosfomycin (5μL) and phage (5 μL), or (iv) PBS alone were directly applied to bacterial lawns, without the use of

171   disks as traditionally described. The applied solutions were allowed to set undisturbed for

172   approximately 5-10 minutes at room temperature, and were then incubated at 37°C for 24h. The

173   zones of inhibition were then measured and recorded.

Qualitative and Quantitative Bactericidal Activity on Biofilms

For qualitative *in vitro* evaluation of antibiofilm efficacy, a 6-well tissue culture plate was

pre-coated with 2% human serum for 24 hours, after which *Staphylococcus aureus* ATCC 6538-

GFP was cultured in tryptic soy broth (TSB) supplemented with 2% glucose for 72 hours. After

gentle washing with PBS, TSB supplemented with vancomycin (256, 512, or 1024 μg/mL),

fosfomycin (16, 64, 128 μg/mL) or bacteriophage (5, 10, or 25 multiplicity of infection (MOI))

was added to the biofilm and incubated for 24 hours. After gentle washing with PBS three times,

remaining biofilm indicated by GFP signal was measured using Cytation 5 plate reader (BioTek).

To quantify the antibiofilm activity of selected therapeutics delivered by alginate

hydrogels, fosfomycin, phage, or dual therapeutic was loaded in 2% alginate hydrogel and

overlaid on top of the biofilms. As a control, empty 2% alginate hydrogel was used. BHI broth

was added and cultures were incubated at 37°C for 24 h. After removing BHI broth, the entire

2% alginate hydrogel and biofilm were harvested and vigorously washed with PBS by

centrifugation to remove residual therapeutic. A serial dilution in PBS was plated onto BHI

plates to determine viable bacterial counts.

Rat Osteomyelitis Model

All procedures (Fig. 1) were performed in accordance with the Institutional Animal Care

and Use Committee (IACUC) of Mississippi State University. Charles River Sprague Dawley

female rats, 13 weeks old, were housed with 12h light/dark cycles and were provided food and

water *ad libitum.* Rats were administered slow release buprenorphine (1.0-1.2 mg/kg BW,

ZooPharm) pre-operatively for pain relief. Rats were anesthetized with isoflurane at an initial

concentration of 2-3%, and maintained at 1-2%. After sterile preparation of the left hindlimb by

fur removal and alcohol and chlorhexidine scrubs, the skin was incised with an anterior

approach, from the level of mid-diaphysis to the patella, along the lower half of the femur. The

muscle tissue was separated using blunt dissection along the muscle bundle divisions on the

anterolateral side of the femur. In the mid-diaphysis, a 1.2 mm (diameter) bicortical defect was

created with a pneumatic drill (Conmed Hall), and a #65 drill bit (McMaster-Carr). To mimic

contamination of orthopedic screws with *S. aureus* occurring in development of osteomyelitis *in*

vivo, sterile orthopedic screws (Antrin Miniature Specialties, #00-90) were placed into 200 μL of

a bacterial suspension (~$1x10^8$ CFU) of ATCC 6538-GFP for approximately 5-10 minutes

(average 6.5 min). The screw was then placed into a 96-well plate to dry for up to 6 minutes

(average 4 min). The bacterial load of these contaminated screws was approximately $5x10^4$ CFU,

determined by placing screws into 1mL of PBS, vigorously vortexing to elute bacteria from the

screw, then serially diluting the eluents for bacterial counting on BHI agar plates. *In vitro*

characterization of bacterial load based on (i) the time screws remained in culture (soak time)

and (ii) dry time was performed using the same procedure as for bacterial counts from *ex vivo*

screws. To assess the effect of soak time, dry time was kept at a constant 5 minutes, and

similarly for the effect of dry time, soak time was kept constant at 5 minutes. To complete the *in*

vivo procedure, the superficial fascia lata and skin were closed with sutures. Longitudinal

monitoring of infection at days 1, 3, and 6 post-infection was performed via radiographs with

fluorescent overlays using the IVIS Lumina XRMS II system (PerkinElmer).

After a 7 day infection period, the area was accessed along the original incision line. The

infected screw was removed and placed into 1mL PBS or fixative for bacterial counting or SEM,

respectively. Then, 100 µL of fosfomycin (3 mg), phage (MOI 3), dual (3 mg fosfomycin and

MOI 3 phage), or PBS loaded into 2% alginate hydrogel was injected into the lateral end of the

bicortical defect, with excess hydrogel pooling in the medial, underlying soft tissue.

On day 8, approximately 24 hours post-treatment, animals were sacrificed via CO_2

inhalation. The hindlimb was initially cleaned with chlorhexidine, and sterile instruments were

used to disarticulate the femur and adjacent soft tissues for further evaluation. For bacterial

counting, bone samples were initially minced using sterile bone rongeurs and further processed

using a homogenizer (Cole-Parmer, LabGEN7, 30s at setting 2-3, 30s at setting 9-10). Soft tissue

samples were minced using sterile surgical scissors, then homogenized (30s at setting 2-3, 30s at

setting 7-8). Following initial processing, homogenates were vortexed (2000 RPM, 1 minute),

diluted as necessary, spread onto BHI agar plates, and incubated for 24 hours at 37°C for

enumeration, with a detection limit set at 25-250 colonies.

Fig 1. Overview of *in vivo* experimental procedure. (A-B) On day 0 (infection surgery day), a

bicortical defect (drill hole) was generated in the mid-diaphysis of the left femur. A

contaminated orthopedic screw was then fastened into this space, and left for 7 days to generate

robust osteomyelitis and soft tissue infection. At day 7, the orthopedic screw was removed and

100µL of therapeutic(s) were injected into the defect space. At day 8, 24h after treatment, soft

tissues and bone samples were collected for histology, scanning electron microscopy, and

bacterial counts. (C) At days 1, 3, and 6, IVIS imaging was performed to track infection

progression.

Electron Microscopy and Histological Analyses

Screw samples for electron microscopy analysis were collected during revision surgeries

on day 7 immediately prior to application of treatment, and placed directly into a fixative

consisting of 5% glutaraldehyde, 2% paraformaldehyde (w/w) in a sodium cacodylate buffer (a

modified "Karnovsky's" solution).[24] For electron microscopy analysis of infected bone, a

representative control femur (empty alginate group) was collected at day 8, broken along the

screw line with sterile bone rongeurs, and immediately placed in fixative for 24h. For both the

screw and bone samples, no dehydration series was performed in order to preserve the biofilm.

Prior to imaging, both samples were placed onto stubs with carbon tape and sputter coated

(Quorom Tech Model # SC7640) with platinum at 30 mA and 3.5 kV for 3-5 minutes. All

samples were then imaged using FESEM (Carl Zeiss AG-SUPRA 40).

For histological analyses, the infected femur and adjacent soft tissues were placed into

10% formalin for 48h, at ~20°C. Bone samples were decalcified for 5 days in Kristensen's

solution, then rinsed and placed into 10% formalin.[25] Tissues were routinely processed,

embedded in paraffin, sectioned at 5µm, and Gram or hematoxylin and eosin (H&E) stained.

Statistical Analyses

All statistical analyses were performed using either GraphPad Prism 8 or SAS software

systems. For the Kirby-Bauer assay, a one-way analysis of variance (ANOVA) with Tukey's

multiple comparisons test was performed. For the *in vitro* antibiofilm assay, a two-way ANOVA

with Tukey's multiple comparisons were performed. For bacterial counts from directly prepared

and *ex vivo* orthopedic screws, a one-way ANOVA was performed, with Sidak's multiple

comparisons. All aforementioned statistical analyses were performed using GraphPad Prism 8

(GraphPad Software, Inc.). For bone and soft tissue *ex vivo* bacterial counts, general linear

models using PROC MIXED in SAS (SAS Institute, Inc.) were performed, with pairwise

comparisons using Tukey's (comparing treatment groups to one another) or Dunnett's

(comparing each treatment to control) tests. An alpha level of 0.05 was used to determine

statistical significance for all methods. Data are presented as mean ± standard deviation (SD).

Results

Integration of GFP into *S. aureus*

Longitudinal analyses of infection progression and regression is an ideal tool for new

model generation. Plasmids harboring reporter genes such as luminescence and fluorescence

have been most commonly used; however, it is necessary to constantly provide antibiotic

selective pressure to prevent a loss of plasmid, which is not achievable with *in vivo* models. In

this study, we integrated the GFP reporter gene into the genome of *S. aureus* ATCC 6538 strain

for the stable and accurate assessment of bacterial growth. All *S. aureus* ATCC 6538-GFP

colonies grown in BHI plates without antibiotic selective pressure were highly fluorogenic over a

span of 7 days (Fig. 2A). Furthermore, ATCC 6538-GFP recovered from an *ex vivo* orthopedic

screw at day 7 post-infection was still highly fluorogenic (Fig. 2B). These results demonstrated

that *S. aureus* ATCC 6538 chromosomally integrated with GFP can be used for real-time

monitoring of bacterial proliferation *in vitro* and *in vivo*.

Fig 2. Integration of GFP into ATCC 6538. (A) Phenotypic expression confirming integration

of GFP into ATCC 6538 was stable over a 7 day period. (B) A *Staphylococcus aureus* CFU

isolated from a contaminated screw *ex vivo* at day 7 revealed GFP expression continued after

growth over a week *in vivo*.

Kirby-Bauer Analyses

For initial investigation of selected therapeutics, a Kirby-Bauer assay was performed

(Fig. 3A). All therapeutics—fosfomycin, phage, and dual—had a larger zone of inhibition than

the PBS control ("a", $p<0.0001$), which generated no zone of inhibition ("N.D.") Dual and

fosfomycin also resulted in a zone of inhibition greater than phage treatment ("b," $p<0.0001$).

Fig 3. *in vitro* Analyses of Therapeutic Bactericidal Activity. (A) From the Kirby-Bauer assay,

all therapeutics delivered via PBS had a greater antibacterial effect than the PBS control ("a",

$p<0.0001$), which generated no zone of inhibition ("N.D."). Dual and fosfomycin therapeutics

also generated a zone of inhibition greater than phage treatment ("b," $p<0.0001$). (B)

Vancomycin (256-1024 µg/mL), fosfomycin, (16-128 µg/mL) and bacteriophage (MOI 5-25)

delivered via PBS revealed varied bactericidal activity, where green indicated bacterial vitality

and black indicated a lack of bacterial presence. Interestingly, vancomycin appeared to have little

to no efficacy on biofilms at all concentrations. Fosfomycin, in contrast, showed efficacy at 64

and 128 µg/mL, a dose range approximately one-tenth the vancomycin doses utilized.

Bacteriophage was effective at an MOI of 10, indicated by the black panel revealing no viable *S.*

aureus. (C) Compared to the empty alginate group, alginate-loaded fosfomycin, phage, and dual

therapeutic-treated biofilms had lower bacterial loads at 6, 12, and 24 hours, except the

fosfomycin group at 6 hours ("c," $p<0.05$). Interestingly, all groups (fosfomycin, phage, dual,

and empty alginate gel) had lower growth at all time points compared to the PBS control, i.e. the

empty alginate gel exerted a killing effect ("s," $p<0.05$). Within individual treatment groups over

time, increased antimicrobial effects were observed. The fosfomycin treated biofilms were

different at 6 and 12 hours, and at 12 and 24 hours (**$p<0.0001$). Phage treated groups were

different at 6 and 12 hours, and at 6 and 24 hours (*$p<0.05$). Dual treated biofilms were different

at 6 and 12 hours, at 6 and 24 hours, and at 12 and 24h (**$p<0.0001$). The empty alginate and

PBS controls resulted in no changes over time.

Qualitative and Quantitative Bactericidal Activity on Biofilms

Biofilms are generally considered the greatest agent of osteomyelitis treatment failure,

and thus are important to consider when developing therapeutics. For this reason, it is important

to evaluate the efficacy of novel therapeutics on robust biofilms, for translation into relevant

clinical scenarios. Antibiofilm efficacy of therapeutics *in vitro* was characterized utilizing two

different analyses: (i) qualitative fluorescent (phenotypic) assessment and (ii) quantitative

bacterial counting. Surprisingly, vancomycin, one of the most commonly utilized antibiotics for

difficult osteomyelitis cases, appeared to have little or no effect on biofilms, as indicated by the

presence of green fluorescing *S. aureus* (Fig. 3B). Fosfomycin is a small molecular weight (138

g/mol) broad-spectrum antibiotic and promising therapeutic option against biofilm.[26] Here,

fosfomycin appeared to remove biofilm at 64 and 128 µg/mL, doses much lower than

vancomycin. The CRISPR-Cas9 modified bacteriophage has dual killing mechanisms: (i) a direct

lysis of target bacteria by holin or murein hydrolase, and (ii) CRISPR-Cas9 nuclease activity.

From qualitative fluorescent analyses, it was determined than a phage MOI of ~10 was effective

in clearing biofilm infection (Fig. 3B).

Alginate is a versatile biopolymer used for prolonged, localized availability of

therapeutic.[27] Antibiofilm assays with bacterial counts were utilized to qualitatively assess the

effects over time of selected therapeutics delivered via alginate (Fig. 3C). Compared to the

empty alginate group, fosfomycin, phage, and dual therapeutic-treated biofilms had significantly

lower bacterial loads at 6, 12, and 24 hours, except the fosfomycin group at 6 hours ("c,"

$p<0.05$). Interestingly, all groups (fosfomycin, phage, dual, and empty alginate gel) were

significantly lower at all time points compared to the PBS treatment, i.e. the empty alginate gel

exerted a killing effect ("s", $p<0.05$). In all treatment groups where alginate was loaded with

therapeutic(s), antibiofilm effects increased over time. The fosfomycin treated biofilms were

different at 6 and 12 hours, and at 12 and 24 hours (**p<0.0001). Phage treated groups were

different at 6 and 12 hours, and at 6 and 24 hours (*p<0.05). Dual treated biofilms were different

at 6 and 12 hours, at 6 and 24 hours, and at 12 and 24h (**p<0.0001). As expected, the empty

alginate and PBS controls did not change over time.

Bacterial Load on Orthopedic Screws

To generate consistent infection in the osteomyelitis model, contaminated orthopedic

screw preparation had to first be characterized. Two parameters of screw preparation were

evaluated: soak time and dry time. Soak time (5-20 min) of screws appeared to increase

somewhat proportionally with respect to bacterial load (Fig. 4A). Dry times from 0-10 min

appeared to have little effect on bacterial load; at 20 min, decreased viability of *S. aureus* was

observed (Fig. 4B).

In the *in vivo* model, orthopedic screws removed at day 7 to allow injection of therapeutic

into the infected defect space were analyzed for bacterial load to confirm all treatment groups

began with a similar extent of infection. Bacterial counts from *ex vivo* screws indicated similarly

severe infection among all samples (treated immediately following), with an average 8.19×10^4

CFU/mL bacterial load (Fig. 4C). Per what would become individual treatment groups,

calculated averages were: 1.05×10^5, 7.50×10^4, 8.20×10^4, and 6.57×10^4 for fosfomycin, phage,

dual, and control groups, respectively. No significant differences between any groups were

observed.

Fig 4. Effect of soak time and dry time on bacterial load of orthopedic screws. (A) Soak time

of sterile orthopedic screws in *Staphylococcus aureus* appeared to linearly relate to ultimate

bacterial load. (B) Dry time did not appear to decrease ultimate bacterial load, until 20 minutes

of dry time. (A-B) The ranges of soak time and dry time used for preparation of screws for the *in*

vivo study are indicated by the red portions of the x-axes. The average time for soak and dry

times for orthopedic screws used *in vivo* is marked by an "x". (C) Bacterial counts collected

from orthopedic screws at day 7 *ex vivo* indicated a similar ultimate bacterial load among what

would become different treatment groups.

Scanning Electron Microscopy

Representative SEM images of screws collected at day 7 post-infection revealed an

abundance of gram positive cocci (*S. aureus*), as expected, along the distal portion of the screw

and within the ridges of the screw throughout its length (Fig. 5A-B). In bone fragments collected

along the screw line (defect site) of an untreated (empty alginate) control sample at day 8 (24h

post-treatment), gram positive cocci, presumably *S. aureus,* were visible. (Fig. 5C-D).

Fig 5. Bacteria on *ex vivo* orthopedic screws and bone. (A-B) Gram positive cocci, and what

appears to be biofilm, was evident on the distal portion (A) and between threads (B) of an *ex vivo*

orthopedic screw excised at day 7. (C-D). Femur sample adjacent to the defect/screw site

collected at day 8 from untreated (empty alginate) control revealed dispersed gram positive

cocci.

Bacterial Load Within Bone and Soft Tissue

The average bone bacterial counts per treatment group were as follows: (i) control: 4.197

$\pm$ 0.289, (ii) fosfomycin: 3.401 $\pm$ 0.924, (iii) phage: 4.076 $\pm$ 0.268, and (iv) dual: 3.607 $\pm$ 0.316

(Log_{10}(CFU)), Fig. 6A). Fosfomycin bacterial counts were lower than empty alginate control

(p=0.0083) and phage (p=0.0486).

The average soft tissue bacterial counts per treatment group were as follows: (i) control:

4.713 $\pm$ 0.331, (ii) fosfomycin: 4.146 $\pm$ 0.377, (iii) phage: 4.160 $\pm$ 0.516, and (iv) dual: 4.201 $\pm$

0.556 (Log$_{10}$(CFU)), Fig. 6B). Bacterial counts were lower in fosfomycin (p=0.0225), phage

(0.0265), and dual (p=0.0430) treated groups compared to empty alginate control.

Fig 6. Bacterial counts from bone and soft tissue. (A) Bacterial counts from bone tissue

harvested from fosfomycin treated animals were lower than those for empty alginate

(**p=0.0083) and phage (*p=0.0486) groups. (B) In soft tissue, bacterial loads were reduced in

all three treatment groups—fosfomycin (*p=0.0225), phage (*p=0.0265), and dual

(*p=0.0430)—compared to empty alginate.

Histology

Bone samples stained with H&E or Gram revealed strong evidence for development of a

severe osteomyelitis infection. The areas at the site of the screw were characterized by extensive

remodeling within the medullary cavity, with replacement of marrow cells by a central area of

neutrophils surrounded by fibrovascular proliferation and reactive bone. Within the cortex, at the

site of screw placement, was mild bone necrosis characterized by empty lacunae and bone loss.

Along the periosteal surface there was locally extensive proliferation of woven bone (periosteal

proliferation). Additionally, abundant gram-positive cocci were localized within the bone. No

differences in the extent of infection among any groups were apparent. The outcome of the

fosfomycin-treated bone histology varied from the other samples, due to sample damage during

processing. Ultimately, this prevented the collection of images along the screw line (where the

majority of bacteria and inflammation were localized), as seen with the other groups. However,

the proliferation of woven bone, as a reactive process on the cortical surface, is visible (Fig. 7B).

Within the phage-treated bone sample, a linear track of gram-positive bacteria in the bone at the

original site of the screw line was visible (Fig. 7C). In the dual-treated sample, neutrophilic

inflammation surrounded by reactive bone and fibrosis was observed (Fig.7D left, *).

Fig 7. Histology of infected bones one day post-treatment. (A, left) At the site of the screw of the empty alginate control was marked neutrophilic inflammation (arrows) with bone loss, surrounded by reactive bone and fibrosis (*, bar=50μm). (A, right) A higher magnification of the area of bone loss with large numbers of neutrophils (bar=5μm). Inset demonstrating gram positive bacteria within and outside macrophages (arrow, bar=5μm). (B) Within the fosfomycin treated bone, periosteal proliferation of woven bone was noted (arrows, bar=50μm). (C, left) A linear track in the bone at the site of the screw (arrows) with abundant neutrophils and reactive bone and fibrosis was observed (arrows, H&E, bar=5μm). (C, right) Gram staining demonstrating aggregates of basophilic bacteria along the screw site (bar=50μm). Inset is a higher magnification of the bacteria (bar=5μm). (D, left) At the site of the screw was marked neutrophilic inflammation with bone loss (arrows), surrounded by reactive bone and fibrosis (*, bar=50μm). (D, right) A higher magnification of the area of bone loss with large numbers of neutrophils (bar=50μm). Inset demonstrating gram positive bacteria within and outside macrophages (arrow, bar=5μm).

Discussion

In this manuscript, a previously developed CRISPR-Cas9 modified bacteriophage therapeutic, which was successful in treating external dermal infection[21], was evaluated as a therapeutic for internal osteomyelitis and contiguous soft tissue infection in a rat model using a biofilm forming strain of *S. aureus*. For real time monitoring of *S. aureus,* GFP was chromosomally integrated into *S. aureus* ATCC 6538 strain by homologous recombination. We demonstrated that *S. aureus* ATCC 6538 strain carrying the GFP gene stably maintained the GFP phenotype without antibiotic selective pressure *in vitro* and *in vivo*.

The therapeutic effects of vancomycin, fosfomycin, CRISPR-Cas9 modified

bacteriophage, and fosfomycin-bacteriophage dual treatments were evaluated against biofilm *in*

vitro. Vancomycin is considered a last resort treatment for *S. aureus* infection. Recent guidelines

recommend vancomycin trough concentrations between 15 and 20 µg/mL for effectively

controlling *S. aureus* infection.[28] As biofilms are typically more difficult to treat than

planktonic bacteria, a much higher concentration of vancomycin (256-1024 µg/mL) was used

here. Although the planktonic *S. aureus* ATCC 6538 strain was sensitive to vancomycin at 2

µg/mL within BHI broth (data not shown), in biofilm form, it was highly resistant to vancomycin

even at 1024 µg/mL. Biofilms consist of a group of bacteria and their byproducts such as

extracellular polymeric substances (EPS) including proteins, DNA, RNA, polysaccharides, and

peptidoglycans. These EPS materials provide physical barriers to penetration of antibiotics to the

inner viable population of bacteria in the biofilm. Vancomycin, a large glycopeptide antibiotic

with a molecular weight of 1,449 g/mol, binds to the D-ala-D-ala terminal amino acid at the stem

of pentapeptide crosslinking peptidoglycan for efficacy[29]. Thus, resistance to vancomycin by

biofilm may be explained by poor penetration of vancomycin due to its bulky size, which could

have led to entrapment at the peptidoglycan layer of biofilm. In contrast, fosfomycin showed

better efficacy against biofilm at much lower (64 and 128 µg/mL) doses. Fosfomycin is a small

bactericidal antibiotic with a molecular weight of 138 g/mol. It interferes with the first step of

peptidoglycan synthesis by inhibiting the phosphoenolpyruvate synthetase[30]. Thus, the

enhanced fosfomycin efficacy could be explained by better penetration of fosfomycin due to its

small size and its inhibition of the first step of peptidoglycan synthesis. From qualitative

fluorescent analyses, it was determined than a phage MOI of ~10 was effective in clearing

biofilm infection. This is similar and in some cases an improvement upon *in vitro* evaluation of

phage treatments discussed in literature, with biofilm eradication reported with MOI 10-

100.[21,31,32] Alginate hydrogel served as an effective delivery vehicle, enabling increasing

effects against biofilm over a 24h period, for fosfomycin, phage, and dual treatments *in vitro*.

Previously, we have observed similar sustained effects of bone morphogenetic protein-2 released

from and retained within alginate hydrogels *in vitro* and *in vivo*.[18,27]

For further evaluation of the CRISPR-Cas9 phage, motivated by augmented *in vitro*

efficacy relative to antibiotic controls, we developed a clinically relevant model of implant-

related osteomyelitis. In human cases of osteomyelitis, chronic infection is diagnosed after a 6-

week period of infection, while our model had only a 1-week infection period. Based on SEM

images, 1 week appeared to be sufficient to induce severe infection, including biofilm, in this rat

model. By culturing *S. aureus* on orthopedic screws, infection was localized to the femur and

surrounding soft tissue, as indicated by fluorescent imaging and histology. Fluorescent imaging

served as a qualitative tool for longitudinal infection progression/regression, although no direct

correlation between radiance output and bacterial load were observed (data not shown). This

could be attributed at least in part to a residual GFP signal that likely exists after bacterial cell

death, due to the persistence of the GFP. Clinically, debridement accompanied by long-term

antibiotic administration is the gold standard for osteomyelitis treatment.[33] In this study, we

have avoided debridement altogether so as to limit potential clearing of infection from any

source other than the therapeutics delivered. For future studies, debridement may be included to

more readily mimic the clinical scenario and enable evaluation of larger antibacterial materials

such as scaffolds or putties.

From bacterial counts performed on excised soft tissues, it was determined that severe

soft tissue infection accompanied the expected high bacterial load in the bone samples. In

clinical cases of osteomyelitis, soft tissue infection is a common pathological finding of

osteomyelitis infection progression[34–36]. In this model, soft tissue infection likely developed

due to the distal end of the orthopedic screw resting freely within the soft tissue medial to the

defect site. On excised orthopedic screws collected on day 7, scanning electron microscopy

indicated a purulent, thick biofilm layer of *Staphylococci* on the end of the orthopedic screw.

Based on *in vitro* results, the process of orthopedic screw preparation can be used to tailor the

extent of infection, as soaking the screws for a shorter period of time would be expected to

introduce less *S. aureus* into the bone and as a result induce a less severe infection. Histology

results support the development of severe osteomyelitis infection, with disease hallmarks such as

neutrophilic inflammation, reactive bone, fibrosis, and gram-positive bacteria. Within the 24-

hour time frame of this study, no differences among treatment groups were apparent. If later time

points were evaluated, the differences noted in bacterial counting would likely be more readily

reflected histologically.

Although only the fosfomycin group resulted in reduced bacterial load in the femur, in

soft tissue, all three treatments, including phage alone and phage with fosfomycin (dual) led to

lower bacterial counts compared to empty alginate. It should be noted than an extremely high

dose of fosfomycin (3g) was administered to the rat femur in this study. In humans, a 3g oral

dose is recommended for treatment of urinary tract infections.[26] Conversely for bacteriophage

dose, although a minimum effective MOI of ~10 was observed *in vitro*, *in vivo* only MOI of ~3

was able to be delivered due to: (i) the volume of alginate hydrogel delivered to the small defect

site (100 µL total, but only ~10 µL fit into the defect itself), and (ii) the thicker consistency of

phage solution, limiting the concentration that could be prepared in alginate hydrogel.

Collectively, these discrepancies in dosing likely limited the efficacy of phage treatment alone in

osteomyelitis mitigation relative to antibiotic with and without phage. Furthermore, the

canaliculi of cortical bone may provide an ideal environment for bacteria to evade treatment.[37]

It should also be noted that in this study, a one-time 100 µL treatment was applied locally.

Clinically, osteomyelitis is treated via debridement and systemic antibiotics for several

weeks.[33] Similarly, success with bacteriophage therapy has been associated with continuous,

prolonged delivery of the virus. A bacteriophage cocktail was used to successfully clear femoral

infection with four intraperitoneal doses of phage (100 µL of ~$2x10^{12}$ plaque forming units

(PFU)/mL) over the span of 48 hours.[38] Another study adopted a treatment regimen for tibial

osteomyelitis consisting of a once daily $3x10^8$ PFU/mL intramuscular bacteriophage injection for

14 days, which resolved the infection.[39] Recently, a case report was published describing the

success of a weekly injection of bacteriophage over a seven-week period for human digital

osteomyelitis.[36] Collectively, these data suggest that sustained, localized concentrations of

phage may be necessary for efficacy in treatment of bone infection. In future studies, a greater

initial dose of phage therapeutic should be considered, or a longer duration of treatment achieved

with a delivery vehicle capable of tailored release of therapeutic. Given a higher phage dose

and/or prolonged availability, it is possible that the efficacy of phage observed here *in vitro* could

be matched *in vivo*. Furthermore, it may be more advantageous to use alternating doses of the

antibiotic and phage therapeutic over time, rather than a combined simultaneous

application.[40,41] In this study, no additive effect of fosfomycin or phage was observed in the

dual treatment group. Only a 24h treatment period was evaluated, which may have limited the

effect of our selected therapeutics, as later time points may have allowed therapeutics, especially

those containing phages (which must replicate for optimal bactericidal activity), to have a

cumulative effect. Nonetheless, our challenging composite tissue infection model enabled

efficient, rapid testing of antimicrobial therapeutics using a biofilm forming strain of *S. aureus*.

Despite the prevalence and severity of osteomyelitis, no bacteriophage-based treatment

for the disease has reached clinical trials in the United States. As populations of MDR-bacteria

continue to spread and new strains are isolated, engineering novel therapeutics will be essential

to augment the dwindling list of effective, available antibiotics. Phage therapy has great potential

to fill this niche, as phages can be made readily and at a low cost. Using CRISPR-Cas9

technology as in this study, phages can be modified to have a wide host range.[21] By

contributing to the pipeline of bacteriophage therapeutic evaluation compared to traditional

antibiotics, the goal of this work was to demonstrate efficacy of phage against bone and soft

tissue infection. Enhanced bactericidal activity of CRISPR-Cas9 phage was demonstrated *in*

vitro against biofilm, when compared to conventionally used vancomycin and fosfomycin

antibiotics. Subsequently, an implant-related model of osteomyelitis and soft tissue infection was

established, confirmed with histological, radiographic, and scanning electron microscopy

analyses. Although phage did not mitigate bone infection 24h post-treatment, soft tissue infection

was reduced 24h following treatment with bacteriophage, and notably to the same extent as

treatment with high dose antibiotic. To improve bacteriological outcomes in the future, further

investigations of delivery vehicles and optimal dosing are warranted.

531 Acknowledgements

This research was supported by the xxxxxxx and the xxxxxx. The authors would like to thank

Dr. Jean Feugang and the USDA-ARS Biophotonics Initiative (58-6402-3-018) for supporting

our use of the IVIS Lumina XR system, Dr. Alicia Olivier for histological analyses, and Drs.

Lucy Senter and Bridget Willeford for veterinary care. Thanks also to Jamie Walker, Delisa

Pennell, Dr. Hayley Gallaher, Weitong Chen, Kali Sebastian, Kristen Lacy, Christine Grant,

Drew Moran, Ryan Yingling, Alex Feaster, Hannah Bostick, Christina Moffett, Sonja Jensen,

Anna Marie Dulaney, and Luke Tucker for assistance with surgeries.

Citations

1. Center for Disease Control and Prevention. Antibiotic Use in the United States, 2017:

Progress and Opportunities. US Dep Heal Hum Serv . 2017;

doi:10.1073/pnas.1417106112

2. Thorpe KE, Joski P, Johnston KJ. Antibiotic-Resistant Infection Treatment Costs Have

Doubled Since 2002, Now Exceeding $2 Billion Annually. Health Aff (Millwood). United

States; 2018;37: 662–669. doi:10.1377/hlthaff.2017.1153

3. O'Neill J. Antimicrobial Resistance : Tackling a crisis for the health and wealth of nations.

Rev Antimicrob Resist. 2014;

4. Tom Frieden. Antibiotic Resistance Threats. Cdc. 2013; doi:CS239559-B

5. Archer NK, Mazaitis MJ, William Costerton J, Leid JG, Powers ME, Shirtliff ME.

Staphylococcus aureus biofilms: Properties, regulation and roles in human disease.

Virulence. 2011. doi:10.4161/viru.2.5.17724

6. Johnson CT, Wroe JA, Agarwal R, Martin KE, Guldberg RE, Donlan RM, et al. Hydrogel

delivery of lysostaphin eliminates orthopedic implant infection by *Staphylococcus aureus*

and supports fracture healing. Proc Natl Acad Sci. 2018; doi:10.1073/pnas.1801013115

7. Geraghty T, LaPorta G. Current health and economic burden of chronic diabetic

osteomyelitis. Expert Review of Pharmacoeconomics and Outcomes Research. 2019.

doi:10.1080/14737167.2019.1567337

8. Rowley WR, Bezold C, Arikan Y, Byrne E, Krohe S. Diabetes 2030: Insights from

Yesterday, Today, and Future Trends. Popul Health Manag. 2017;

doi:10.1089/pop.2015.0181

9. Abatángelo V, Peressutti Bacci N, Boncompain CA, Amadio AA, Carrasco S, Suárez CA,

et al. Broad-range lytic bacteriophages that kill Staphylococcus aureus local field strains.

PLoS One. 2017; doi:10.1371/journal.pone.0181671

10. Tang Z, Huang X, Sabour PM, Chambers JR, Wang Q. Preparation and characterization of

dry powder bacteriophage K for intestinal delivery through oral administration. LWT -

Food Sci Technol. 2015; doi:10.1016/j.lwt.2014.08.012

11. Furfaro LL, Payne MS, Chang BJ. Bacteriophage Therapy: Clinical Trials and Regulatory

Hurdles. Front Cell Infect Microbiol. 2018; doi:10.3389/fcimb.2018.00376

12. Wright A, Hawkins CH, Änggård EE, Harper DR. A controlled clinical trial of a

therapeutic bacteriophage preparation in chronic otitis due to antibiotic-resistant

Pseudomonas aeruginosa; a preliminary report of efficacy. Clin Otolaryngol.

Wiley/Blackwell (10.1111); 2009;34: 349–357. doi:10.1111/j.1749-4486.2009.01973.x

13. Leitner L, Sybesma W, Chanishvili N, Goderdzishvili M, Chkhotua A, Ujmajuridze A, et

al. Bacteriophages for treating urinary tract infections in patients undergoing transurethral

resection of the prostate: A randomized, placebo-controlled, double-blind clinical trial.

BMC Urol. 2017; doi:10.1186/s12894-017-0283-6

14. Fabijan AP, Lin RCY, Ho J, Maddocks S, Iredell JR. Safety and Tolerability of

Bacteriophage Therapy in Severe Staphylococcus aureus

Infection. bioRxiv. 2019; 619999. doi:10.1101/619999

15. Voelker R. FDA Approves Bacteriophage TrialFDA Approves Bacteriophage TrialNews

From the Food and Drug Administration. JAMA. 2019;321: 638.

doi:10.1001/jama.2019.0510

16.   Johnson C, Dinjaski N, Prieto M, García A. Bacteriophage Encapsulation in

584   Poly(Ethylene Glycol) Hydrogels Significantly Reduces Bacteria Numbers in an Implant-

585   Associated Infection Model of Bone Repair. Igarss 2014. 2014; doi:10.1007/s13398-014-

586   0173-7.2

587   17.   Bean JE, Alves DR, Laabei M, Esteban PP, Thet NT, Enright MC, et al. Triggered

588   Release of Bacteriophage K from Agarose/Hyaluronan Hydrogel Matrixes by

589   Staphylococcus aureus Virulence Factors [Internet]. CHEMISTRY OF MATERIALS. pp.

590   7201–7208. doi:10.1021/cm503974g

591   18.   Priddy LB, Chaudhuri O, Stevens HY, Krishnan L, Uhrig BA, Willett NJ, et al. Oxidized

592   alginate hydrogels for bone morphogenetic protein-2 delivery in long bone defects. Acta

593   Biomater. 2014/06/17. 2014;10: 4390–4399. doi:10.1016/j.actbio.2014.06.015

594   19.   Krishnan L, Priddy LB, Esancy C, Li MTA, Stevens HY, Jiang X, et al. Hydrogel-based

595   Delivery of rhBMP-2 Improves Healing of Large Bone Defects Compared With

596   Autograft. Clin Orthop Relat Res. 2015; doi:10.1007/s11999-015-4312-z

597   20.   Souza GR, Yonel-Gumruk E, Fan D, Easley J, Rangel R, Guzman-Rojas L, et al. Bottom-

598   up assembly of hydrogels from bacteriophage and Au nanoparticles: The effect of Cis-

599   and trans-acting factors. PLoS One. 2008; doi:10.1371/journal.pone.0002242

600   21.   Park JY, Moon BY, Park JW, Thornton JA, Park YH, Seo KS. Genetic engineering of a

601   temperate phage-based delivery system for CRISPR/Cas9 antimicrobials against

602   Staphylococcus aureus. Sci Rep. The Author(s); 2017;7: 44929. Available:

603   http://dx.doi.org/10.1038/srep44929

22.   De Jong NWM, Van Der Horst T, Van Strijp JAG, Nijland R. Fluorescent reporters for

markerless genomic integration in Staphylococcus aureus. Sci Rep. 2017;

doi:10.1038/srep43889

23. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a

standardized single disk method. Am J Clin Pathol. England; 1966;45: 493–496.

24. Karnovsky MJ. A Formaldehyde-Glutaraldehyde Fixative of High Osmolality for Use in

Electron Microscopy. The American Society for Cell Biology Source: The Journal of Cell

Biology. 1965.

25. Kristensen HK. An Improved Method of Decalcification. Stain Technol. Taylor &

Francis; 1948;23: 151–154. doi:10.3109/10520294809106242

26. Dijkmans AC, Zacarías NVO, Burggraaf J, Mouton JW, Wilms E, van Nieuwkoop C, et

al. Fosfomycin: Pharmacological, Clinical and Future Perspectives. Antibiotics. 2017;

doi:10.3390/antibiotics6040024

27. Krishnan L, Priddy LB, Esancy C, Klosterhoff BS, Stevens HY, Tran L, et al. Delivery

vehicle effects on bone regeneration and heterotopic ossification induced by high dose

BMP-2. Acta Biomater. 2017; doi:10.1016/j.actbio.2016.12.012

28. Lodise TP, Lomaestro B, Graves J, Drusano GL. Larger vancomycin doses (at least four

grams per day) are associated with an increased incidence of nephrotoxicity. Antimicrob

Agents Chemother. 2008; doi:10.1128/AAC.01602-07

29. Wang F, Zhou H, Olademehin OP, Kim SJ, Tao P. Insights into key interactions between

vancomycin and bacterial cell wall structures. ACS Omega. 2018;

doi:10.1021/acsomega.7b01483

30. Takahata S, Ida T, Hiraishi T, Sakakibara S, Maebashi K, Terada S, et al. Molecular

mechanisms of fosfomycin resistance in clinical isolates of Escherichia coli. Int J

Antimicrob Agents. 2010; doi:10.1016/j.ijantimicag.2009.11.011

31.   Alves DR, Gaudion A, Bean JE, Perez Esteban P, Arnot TC, Harper DR, et al. Combined

Use of Bacteriophage K and a Novel Bacteriophage To Reduce Staphylococcus aureus

Biofilm Formation. Appl Environ Microbiol. 2014; doi:10.1128/aem.01789-14

32.   Pincus NB, Reckhow JD, Saleem D, Jammeh ML, Datta SK, Myles IA. Strain specific

phage treatment for Staphylococcus aureus infection is influenced by host immunity and

site of infection. PLoS One. 2015; doi:10.1371/journal.pone.0124280

33.   Nelson CL, McLaren SG, Skinner RA, Smeltzer MS, Thomas JR, Olsen KM. The

treatment of experimental osteomyelitis by surgical debridement and the implantation of

calcium sulfate tobramycin pellets. J Orthop Res. 2002; doi:10.1016/S0736-

0266(01)00133-4

34.   Taki H, Krkovic M, Moore E, Abood A, Norrish A. Chronic long bone osteomyelitis:

diagnosis, management and current trends. Br J Hosp Med. 2016;

doi:10.12968/hmed.2016.77.10.c161

35.   Berbari EF, Kanj SS, Kowalski TJ, Darouiche RO, Widmer AF, Schmitt SK, et al. 2015

Infectious Diseases Society of America (IDSA) Clinical Practice Guidelines for the

Diagnosis and Treatment of Native Vertebral Osteomyelitis in Adults[a]. Clinical Infectious

Diseases. 2015. doi:10.1093/cid/civ633

36.   Fish R, Kutter E, Bryan D, Wheat G, Kuhl S. Resolving Digital Staphylococcal

Osteomyelitis Using Bacteriophage—A Case Report. Antibiotics. 2018;

doi:10.3390/antibiotics7040087

37.   de Mesy Bentley KL, Trombetta R, Nishitani K, Bello-Irizarry SN, Ninomiya M, Zhang

L, et al. Evidence of Staphylococcus Aureus Deformation, Proliferation, and Migration in

651   Canaliculi of Live Cortical Bone in Murine Models of Osteomyelitis. J Bone Miner Res.

652   2017; doi:10.1002/jbmr.3055

653 38. Kishor C, Mishra RR, Saraf SK, Kumar M, Srivastav AK, Nath G. Phage therapy of

654   staphylococcal chronic osteomyelitis in experimental animal model. Indian J Med Res.

655   2016; doi:10.4103/0971-5916.178615

656 39. Ibrahim OMS, Sarhan SR, Salih SI. Activity of Isolated Staphylococcal Bacteriophage in

657   Treatment of Experimentally Induced Chronic Osteomyelitis in Rabbits. Adv Anim Vet

658   Sci. 2016; doi:10.14737/journal.aavs/2016/4.11.593.603

659 40. Lopes A, Pereira C, Almeida A. Sequential Combined Effect of Phages and Antibiotics on

660   the Inactivation of Escherichia coli. Microorganisms. MDPI; 2018;6: 125.

661   doi:10.3390/microorganisms6040125

662 41. Tagliaferri TL, Jansen M, Horz H-P. Fighting Pathogenic Bacteria on Two Fronts: Phages

663   and Antibiotics as Combined Strategy  [Internet]. Frontiers in Cellular and Infection

664   Microbiology . 2019. p. 22. Available:

665   https://www.frontiersin.org/article/10.3389/fcimb.2019.00022

A

Generation of a bicortical defect → Placement of a contaminated screw → Removal of screw at 7 days → Injection of therapeutic into the bicortical defect

B

Infection Surgery:
Placement of a Contaminated Screw into a Bicortical Defect

Treatment Surgery:
Removal of the Screw and Application of Treatment(s)

IVIS Imaging

IVIS Imaging

IVIS Imaging

IVIS Imaging, Histology, and Bacterial Counts

Activity

Day

0   1   3   6   7   8

C

Representative Pre-treatment Longitudinal IVIS Images

Day 1

Day 3

Day 6

$\times 10^8$

4.0

3.5

3.0

2.5

Radiance
$(p/sec/cm^2/sr)$

A *in vitro* **stability of ATCC 6538-GFP**   **B** *ex vivo stability* **of ATCC 6538-GFP**

A Kirby-Bauer Assay

Zone of Inhibititon (mm)

B Qualitative Antibiofilm Assay

Vancomycin Fosfomycin CRISPR-Cas9 Phage

TSB

C Quantitative Antibiofilm Assay

$\log_{10}$ (CFU)

Fosfomycin
Phage
Dual
Empty Alginate
PBS

A Effect of Soak Time on Screw Bacterial Load

B Effect of Dry Time on Screw Bacterial Load

C Bacteria Recovered from *ex vivo* Orthopedic Screws at Day 7