Genetic architecture influences when and how hybridization contributes to colonization

Bryan Reatini[1,2,] Todd J. Vision[1,3,4]

[1]Department of Biology, University of North Carolina at Chapel Hill, NC 27599-3280

[2]ORCID:0000-0001-8643-4296

[3]ORCID:0000-0002-6133-2581

[4]*To whom correspondence should be addressed*: tjv@unc.edu

Running title: Genetic architecture, hybridization and colonization

Keywords: genetic architecture, hybridization, colonization

Abstract

The role of genetic architecture in adaptation to novel environments has received considerable attention when the source of adaptation variation is *de novo* mutation. Relatively less is known when the source of adaptive variation is inter- or intraspecific hybridization. We model hybridization between divergent source populations and subsequent colonization of an unoccupied novel environment using individual-based simulations in order to understand the influence of genetic architecture on the timing of colonization and the mode of adaptation. We find that two distinct categories of genetic architecture facilitate rapid colonization but that they do so in qualitatively different ways. For few and/or tightly linked loci, the mode of adaptation is via the recovery of adaptive parental genotypes. With many unlinked loci, the mode of adaptation is via the generation of novel hybrid genotypes. The first category results in the shortest colonization lag phases across the widest range of parameter space, but further adaptation is mutation limited. The second category takes longer and is more sensitive to genetic variance and dispersal rate, but can facilitate adaptation to environmental conditions which exceed the tolerance of parental populations. These findings have implications for understanding the origins of biological invasions and the success of hybrid lineages.

Introduction

A persistent question regarding the source of adaptive variation leading to colonization is why intra and interspecific admixture – hereafter collectively referred to as hybridization – sometimes leads to increased colonization success and sometimes does not (Bock et al., 2015; Dlugosch et al., 2015). Hybridization can contribute to colonization by producing an overall increase in standing genetic variation, by transferring adaptive alleles between genetic backgrounds (adaptive introgression), or by producing novel genotypes in hybrids (transgressive segregation) (Pfennig et al., 2016). Many species complexes are known in which hybridization may have contributed to colonization in these ways (Ellstrand and Schierenbeck, 2000; Schierenbeck and Ellstrand, 2008). However, hybridization doesn't always contribute to increased colonization success, and identifying the genetic factors which may influence when and how hybridization contributes to colonization has become a central objective of the field of invasion genetics (Bock et al., 2015).

A second persistent question in invasion genetics is what the genetic architecture of so-called invasiveness traits tends to be (Bock et al., 2015; Dlugosch et al., 2015). Genetic architecture here is defined as the number of loci governing a quantitative trait, their positions within the genome, and the interactions among different alleles and loci. Much is already known about the genetic architecture of adaptation to heterogeneous environments when adaptation stems from *de novo* mutation. For example, models of adaptation across linear gradients have demonstrated the importance of sufficient genetic variance (Kirkpatrick and Barton, 1997), two patch models have demonstrated the importance of mutations of large effect (Gomulkiewicz et al., 1999; Holt et al., 2003), and a recent evaluation of adaptation across an approximately continuous landscape demonstrated the potential importance of both high genetic variance and

mutations of large effect (Gilbert and Whitlock, 2017). However, range expansion may commonly be facilitated by gene flow rather than *de novo* mutation, and studies on the genetic architecture of colonization and range expansion stemming from hybridization are currently lacking.

Here, we wish to examine how genetic architecture itself influences when and how hybridization contributes to colonization success. Hybrid invasions present a unique combination of population genetic forces which may interact with genetic architecture in important ways. For example, selection on beneficial variation within a population receiving migrants may be rendered ineffective due to genetic swamping – the homogenizing effect of persistent gene flow (García-Ramos and Kirkpatrick, 1997; Yeaman and Otto, 2011; Yeaman and Whitlock, 2011). The strength of genetic swamping can be modulated by genetic architecture, with architectures dominated by alleles with large selection coefficients and/or tightly linked loci being more resistant than architectures dominated by alleles with small selection coefficients and/or unlinked loci (Tigano and Friesen, 2016; Yeaman and Whitlock, 2011). Yeaman (2015) also demonstrated that architectures characterized by many alleles of small effect may be robust to genetic swamping when genetic variance is sufficiently high. For this reason, the genetic architectures of adaptive traits in hybrid populations may be filtered by genetic swamping.

When hybridization occurs between divergent lineages in a new environment, as would be the case between introduced lineages or between an introduced and native lineage, colonization success may depend on the ability to recover and maintain adaptive variation in the face of persistent migration from these divergent source populations. In fact, the time required for natural selection to drive adaptive genetic variation to high frequency is a proposed explanation for one of the most ubiquitous phenomena in invasion biology; the initial lag phase in which

introduced populations persist in low numbers before rapid growth (Crooks, 2005; Crooks and Soulé, 1999). Indeed, using a source-sink model incorporating adaptation from mutation, Holt et al. (2003) demonstrated that transient occupancy of a sink can persist for a long time before punctuated growth following adaptation. Given these findings, it is reasonable to hypothesize that factors such as genetic architecture, which influence the recovery of adaptive variation in newly formed hybrid populations, may influence the duration of the lag phase. If the lag phase is sufficiently long, we may not witness its resolution within our limited window of observation, and conclude that hybridization has failed to contribute to colonization. We hereafter refer to the idea that the genetic architecture of invasiveness traits influences the contribution of hybridization to colonization – as the *architecture hypothesis of hybrid invasion*.

Here we test the architecture hypothesis of hybrid invasion using individual based simulations. Specifically, we test two underlying predictions: (1) that colonization lag phase duration is sensitive to the genetic architecture of invasiveness traits, (2) that architectures resistant to genetic swamping will produce relatively short lag phases. Given our results, we then test an additional prediction: the filtering effect of colonization on genetic architecture influences the ability of resulting colonist lineages to occupy and adapt to more extreme environments – an important characteristic for hybrid speciation and adaptive radiation.

Methods

We simulated colonization in a source-sink scenario involving dispersal from two divergent source populations using quantiNemo2 (Neuenschwander et al., 2008). Generations were discrete and consisted of four life cycle events in the following order: reproduction, offspring dispersal, aging, and regulation of adults. We initiated polymorphic populations in two separate patches with differing environmental optima, limiting dispersal for a 10,000 generation *burn-in period* to allow the two populations to reach mutation-selection-drift equilibrium in their respective patches. We then allowed one-way dispersal from these two source populations into a novel environment (the sink) for 500 generations – a timeframe relevant for most contemporary invasions – which we refer to as the *colonization phase*. We assume that occupancy of the novel environment by either source population alone is dependent on recurrent immigration. Thus it will serve as a sink for maladapted migrants from the source populations, at least initially.

We simulated two quantitative traits and subjected them to stabilizing selection. For the first trait, one source population generated an adaptive genotype during the burn-in period relative to the optimum of the sink patch while the other source population generated a maladaptive genotype (Figure 1). The opposite was true for the second trait, resulting in each source population possessing an adaptive genotype for one trait and a maladaptive genotype for the other with respect to the trait optimum in the sink. In this way, all of the standing genetic

Figure 1. The landscape of adaptive optima for the two quantitative traits. The sink shares optima with one source populations for each trait. The secondary sink has optima more extreme than those of the original source patches.

variation necessary to adapt to the sink is present in the two source populations at the start of the colonization phase, but it must be recovered via hybridization between immigrants in the sink.

We evaluated a range of differences in environmental optima (ΔZ_{opt} of 6, 8, and 10) between source patches, where Z_{opt} represents the optimal phenotype for each quantitative trait. For reference, one unit of Z_{opt} is equivalent to two-fold the variance of the mutation effect size distribution for 1-locus architectures. This range is intermediate between smaller values of ΔZ_{opt} that yield successful colonization by a single source population in the absence of hybridization, and larger values that fail to colonize under a variety of genetic architectures. The range of values of ΔZ_{opt} tested here therefore encompasses the environmental optima relevant for our analysis of hybridization's contribution to colonization given the other assumptions we make about the strength of selection (discussed below).

Reproduction and dispersal

Reproduction followed a Wright-Fisher Model with a variable population size, N, of diploids in which random mating occurred with any individual in the same patch, including self-mating, with probability $1/N$. Fecundity for each individual was drawn from a Poisson distribution with a mean of four. Growth within each patch was governed by the fecundity of its individuals, and each patch had a carrying capacity of 1000 individuals. We allowed one-way dispersal from source populations into the sink at dispersal rates of $d=10^{-2}$ and $d=10^{-3}$ during the colonization phase, where dispersal rate is the probability of an offspring migrating to the sink. Our justification for using these rates is as follows; based on our previous assumptions, the expected number of migrants N_{m} entering the sink per generation can be written as $N_{\mathrm{m}} = 2Kfd$. This simplifies to $N_{\mathrm{m}} = 8000d$, given that carrying capacity $K = 1000$ for both source populations

and fecundity f has a mean of four. At $d=10^{-1}$, $N_m=800$, which could result in the sink reaching

carrying capacity solely due to the influx and chance survival of migrants, which would not be of

interest. At the other extreme of $d=10^{-4}$, $N_m=0.8$, which may not provide sufficient opportunity

for hybridization to occur within 500 generation. We therefore limit our analysis to intermediate

values of $d=10^{-2}$ and $d=10^{-3}$, for which $N_m=80$ and $N_m=8$, respectively. This provides opportunity

for hybridization between migrants while still allowing us to evaluate the effect of an order of

magnitude difference in propagule pressure. After dispersal, aging occurred, which removed the

adults from each patch and converted the offspring into adults for the next non-overlapping

generation. Population regulation then followed, randomly culling the number of new adult

individuals in each patch that exceeded the carrying capacity.

Selection and genetic architecture

Selection acted on offspring survivorship, with an individual's fitness modeled as its survival

probability before aging and reproduction. Stabilizing selection acted on the phenotype of both

quantitative traits within each patch. Trait fitness W_t was defined by the standard Gaussian

function $W_t = e^{-\frac{\left(P-Z_{opt}\right)^2}{2\omega^2}}$, where P is the value of the quantitative trait, Z_{opt} is the optimal trait

value of the patch, and ω^2 is the width of the fitness function. We assume ω^2 to be 7.5 and

environmental variance (V_E) to be one, following Gilbert et al. (2017). These values were used

throughout all simulations. We then calculate the fitness of an individual W as the product of the

trait fitness for the two simulated traits: $W = \prod_{i=1}^{2} W_i$.

We simulated quantitative traits under four baseline schemes of genetic architecture

characterized by the number of loci (1, 10, 100, and 1000 loci), assuming free recombination and

additive alleles. We then modified these baseline architectures in subsequent analyses to include

variations in linkage and epistasis (Table 1). Mutations acted in a stepwise fashion where the

effect of a new mutation was added to the previous allelic value. The variance of the normal

distribution from which mutational effects were drawn was scaled by the number of loci

underlying the trait (Table 1), such that the effect size of mutations generated during the burn-in

period depended upon the model being tested (*i.e.,* few mutations of large effect for architectures

with fewer loci and many mutations of small effect for architectures with many loci). We

compared mutation rates of $\mu=10^{-4}$ and $\mu=10^{-5}$ in order to evaluate the effect of equilibrium

values of V_g within source populations on colonization success, following Yeaman (2015), who

found that these two rates produced notable differences in the capacity for adaptation when

genetic swamping was strong. In order to prevent adaptation from mutation during the

colonization phase, we only allowed mutation during the burn-in period.

Analysis

We measured lag phase duration as the number of generations between the initiation of

dispersal into the sink (transient occupancy) and the time at which carrying capacity in the sink

was reached (permanent residency). We used mean genotypic value, G, as a metric to track

adaptation within the sink population during the colonization phase. It was calculated as $G =$

$\sum_{i=1}^{L} a_i + a_{i'}$, where L is the number of loci and a_i and $a_{i'}$ are the two alleles at the i^{th} locus of a

diploid individual. This formula is modified in section 3.2.2. to incorporate epistasis.

Loci	mutation rate	allelic variance	linkage clusters	epistatic variance
1	10^{-4}	0.5	U	0
10	10^{-4}	0.05	U	0
100	10^{-4}	0.005	U	0
1000	10^{-4}	0.0005	U	0
1	10^{-5}	0.5	U	0
10	10^{-5}	0.05	U	0
100	10^{-5}	0.005	U	0
1000	10^{-5}	0.0005	U	0
10	10^{-5}	0.05	1	0
100	10^{-5}	0.005	1	0
100	10^{-5}	0.005	10	0
1000	10^{-5}	0.0005	1	0
1000	10^{-5}	0.0005	10	0
10	10^{-5}	0.05	U	0.05
10	10^{-5}	0.05	U	0.5
10	10^{-5}	0.05	U	2.25

Table 1. Genetic architectures tested in simulations of hybrid colonization. Allelic variance is the variance of the distribution of mutation effect sizes. *U* denotes unlinked loci.

Given that mutation was not allowed during the colonization phase, sampling the entirety of both source populations at the end of the burn-in period allowed us to determine the origin of every allele in the metapopulation – with the exception of alleles found in both source populations at the beginning of the colonization phase. Using these data, we characterized the genetic architecture of adaptation to the sink by measuring the frequency, effect size, and origin of each adaptive allele within the sink immediately after the lag phase. To be included in the analysis, we required an adaptive allele to have a frequency greater than a dispersal dependent threshold, calculated as the maximum frequency of a new migrant allele entering the sink from one of the two source populations:

$$\max (\text{frequency}) = \frac{\left((2N_{\text{source}}f)d\right)}{2N_{\text{sink}}}$$

where f is fecundity and d is dispersal rate. Since N_{source} and N_{sink} are both K after the lag phase

and f is assumed to have a mean of four throughout all simulations, this simplifies to

$\text{max(frequency)} = 4d$.

We then quantified the relative impact of each of these adaptive alleles in the sink by taking

the product of its frequency and effect size, a quantity we refer to as the *weight* of an allele. By

summing the weights of adaptive alleles originating from each source population, we were able

to estimate the relative contribution of each source population to adaptation in the sink. When

adaptation in the sink proceeds primarily by the recovery of parental genotypes, this is reflected

by the total weight of adaptive alleles being primarily from one source. By contrast, when

adaptation is achieved through the generation of hybrid genotypes, the contribution from the two

source populations is more even. All analyses were averaged across 100 simulation replicates for

each scheme of genetic architecture. Analyses of variance (ANOVAs) were then performed to

determine the effect of genetic architecture on lag phase duration.

Results

The 'when' and the 'how' of hybrid colonization

In order to test the architecture hypothesis of hybrid invasion, we first focused on its

primary underlying prediction: colonization lag phase duration is sensitive to the genetic

architecture of invasiveness traits. Lag phase duration for the four baseline schemes of genetic

architecture (1, 10, 100, or 1000 loci, assuming free recombination and additivity within and

between loci) varied with the magnitude of the differences between patch optima (ΔZ_{opt}), the

mutation rate during the burn-in period (which in turn affected the equilibrium values of V_g in the

source populations), and the dispersal rate (Figure 2). In mild environments, which we define as

having a relatively small ΔZ_{opt} of 6, mean colonization lag phase was on the order of ten generations for all genetic architectures and all parameter values, and thus did not differ between architectures. The lag phase when colonizing moderate environments ($\Delta Z_{opt}=8$) varied depending on the genetic architecture and other assumptions of the model (Figure 3A). In more extreme environments ($\Delta Z_{opt}=10$), 1-locus architectures were capable of colonizing the sink but all other architectures failed (i.e., the lag phase exceeded 500 generations) for the parameter values we

tested. For subsequent analyses, we use $\Delta Z_{opt}=8$, since for that parameter value, all genetic architectures showed some sensitivity to the other parameters.

Overall, the lag phase was shorter for the 1-locus simulations compared to all other architectures under all parameter values tested. For architectures with many loci, a low burn-in mutation rate produced longer lag phases and, when combined with high dispersal rate, resulted in a failure to colonize the sink during colonization phase. A significant three-way interaction was found between the number of loci, the dispersal rate, and the mutation rate. In order to analyze the underlying factors contributing to this interaction, simple second-order interactions were analyzed by performing two-way ANOVAs between each pair

Figure 2. Mean colonization lag phase and 95% confidence intervals for each baseline architecture assuming different values of ΔZ_{opt} (white=6, grey=8, black=10), dispersal rate d (diamonds=10^{-2}, squares=10^{-3}) and mutation rate ($\mu=10^{-4}$ in A, $\mu=10^{-5}$ in B). For $\Delta Z_{opt}=8$, significant interactions were found between locus number, mutation rate, and dispersal rate.

of factors (loci, μ and d) at each level of the third factor (Table 3). Significant simple second-order interactions were found between mutation and dispersal for 100 and 1000 loci, between loci and dispersal at $\mu=10^{-5}$ (Figure 2B), and between loci and mutation at both levels of dispersal. For all architectures, lag phase duration closely corresponded with the timing of adaptation to the sink, as shown for the case of high mutation and dispersal rates in Figure 3B.

	Type II sum of squares	Degrees of freedom	F	p
dispersal	39,649	1	80.495	< 2.2e-16
loci	17,581,779	3	1189.814	< 2.2e-16
mutation	14,335,878	1	2910.461	< 2.2e-16
dispersal-by-loci	614,310	3	41.572	< 2.2e-16
dispersal-by-mutation	615,323	1	124.922	< 2.2e-16
loci-by-mutation	12,812,203	3	867.042	< 2.2e-16
dispersal-by-loci-by-mutation	653,133	3	44.2	< 2.2e-16
residual error	780,2212	1,584		

Table 2. Multifactorial ANOVA for the response variable of colonization lag time and the fixed factors of dispersal rate, mutation rate, and the number of loci.

	Type II sum of squares	Degrees of freedom	F	p
dispersal:mutation for 1 locus				
dispersal	951.7	1	366.129	< 2.2e-16
mutation	1.8	1	0.701	0.403
dispersal:mutation	5.1	1	1.948	0.164
residuals	1029.4	396		
dispersal:mutation for 10 loci				
dispersal	4823	1	1.418	0.23448
mutation	44711	1	13.143	0.00033
dispersal:mutation	4090	1	1.202	0.27356
residuals	1347178	396		

	dispersal:mutation for 100 loci			
dispersal	229393	1	26.52	4.12e-07
mutation	13671876	1	1580.65	< 2.2e-16
dispersal:mutation	241228	1	27.89	2.12e-07
residuals	3425205	396		
	dispersal:mutation for 1000 loci			
dispersal	775632	1	101.4	< 2.2e-16
mutation	13431492	1	1756.1	< 2.2e-16
dispersal:mutation	1023132	1	133.8	< 2.2e-16
residuals	3028799	396		
	loci:dispersal at $\mu=10^{-4}$			
loci	539545	3	91.015	< 2.2e-16
dispersal	11974	1	6.06	0.014
loci:dispersal	6101	3	1.029	0.379
residuals	1565009	792		
	loci:dispersal at $\mu=10^{-5}$			
loci	29854436	3	1263.64	< 2.2e-16
dispersal	999839	1	126.96	< 2.2e-16
loci:dispersal	1261342	3	53.39	< 2.2e-16
residuals	6237202	792		
	loci:mutation at $d=10^{-2}$			
loci	11845016	3	1966	< 2.2e-16
mutation	10445649	1	5202	< 2.2e-16
loci:mutation	9299236	3	1544	< 2.2e-16
residuals	1590229	792		
	loci:mutation at $d=10^{-3}$			
loci	6351073	3	269.9	< 2.2e-16
mutation	4505552	1	574.4	< 2.2e-16
loci:mutation	4166099	3	177.1	< 2.2e-16
residuals	6211983	792		

Table 3. Two-way ANOVAs for the response variable of colonization lag time and each pair of fixed factors of dispersal rate, mutation rate, and the number of loci at each level of the third factor.

In addition to quantitatively affecting the duration of the lag phase, the different genetic

architectures led to qualitatively different modes of adaptation to the sink (Figure 3C). We

Figure 3. A) Duration of the colonization lag phase (in generations) for each simulation replicate (black dots) under different dispersal rates *d* and mutation rates μ assuming ΔZ_{opt}=8. B) Recovery of mean genotypic value in the sink for each simulation replicate (black lines), the end of the lag phase for each replicate (grey bars), and mean lag phase duration (white diamonds) – taken from high mutation rate and high dispersal rate simulations from panel A. C) Total contribution to adaptation (in terms of total weight of adaptive alleles) in the sink for each source population averaged across simulation replicates from panel B.

analyzed the genetic architecture of adaptation under high mutation and dispersal rates, since the

lag phase duration was minimized with these parameter values for all genetic architectures. For

the 1-locus and 10-loci simulations, adaptation to the sink was based on the same alleles that contributed to adaptation in the source populations. Specifically, 100% of alleles contributing to adaptation in the sink originated from their respective adapted source population for 1-locus architectures. For 10-loci simulations, the mean contribution across replicates was 98.2% for both traits. In contrast, adaptation to the sink for 100-loci and 1000-loci architectures proceeded predominantly by the generation of novel hybrid genotypes. For 100-loci, the mean contribution of adaptive alleles originating from source population one was 33.1% for trait one and 67.1% for trait two. For 1000-loci, the mean contribution from source population one was 45.3% for trait one and 54.5% for trait two. Thus, two modes of adaptation in the sink are observed; the recovery of parental genotypes as seen for 1 and 10-loci architectures, and the generation of hybrid genotypes for 100 and 1000-loci architectures.

Linkage

Above, we have assumed free recombination between all loci and additive contributions to phenotype for the sake of simplicity, but both linkage and epistasis may influence adaptation in newly formed hybrid populations (Abbott et al., 2013; Harrison and Larson, 2014; Yeaman and Whitlock, 2011). Linkage between loci reduces the recombination rate, and therefore may reduce the strength of genetic swamping when there are many loci (Tigano and Friesen, 2016). Since genetic swamping from source populations appears to be the force that impeded colonization for 100-loci and 1000-loci architectures when the burn-in mutation rate was low, we hypothesized that linkage could rescue these architectures under such conditions. To test this hypothesis, we incorporated varying degrees of linkage into 10, 100, and 1000 locus architectures. All loci for each trait were clustered within either 1 or 10 chromosomes. Chromosomes were assumed to be

100 cM in length. Loci were spaced at 1 cM intervals in all cases except that of 1000 loci on a

single chromosome, in which spacing was 0.1 cM. Effectively, all the loci affecting each trait

were grouped into either one or ten clusters of linked loci (Table 1). Importantly, there was free

recombination between the loci of the two different traits.

Incorporating linkage rescued the 100-loci and 1000-loci architectures under conditions of

low burn-in mutation rate and high dispersal, with one cluster of linked loci resulting in shorter

lag phases than ten clusters (Figure 4A). Linkage also resulted in a shift in the mode of

adaptation from the generation of novel hybrid genotypes towards the recovery of parental

genotypes (Figure 4B).

Figure 4. A) Duration of colonization lag phase for each simulation replicate incorporating varying degrees of linkage and assuming low burn-in mutation rate ($\mu=10^{-5}$) and high dispersal ($d=10^{-2}$). B) Genetic architecture of adaptation to the sink assuming 1 or 10 linkage clusters from panel A in comparison to baseline simulations with no linkage (U=unlinked) assuming high mutation rate ($\mu=10^{-4}$), and high dispersal rate ($d=10^{-2}$).

Epistasis

Epistasis may influence adaptation in a hybrid zone by creating novel interactions between

divergent migrant genotypes. If epistatic interactions are favored by selection, epistasis may

result in shorter lag phases and a shift in the mode of adaptation to the generation of novel

genotypes. While if such interactions are disfavored, lag phase duration may be lengthened while

parental genotypes are recovered. In order to test these predictions, we incorporated epistasis into the model such that the genotype of an individual was determined by the combination of the additive effects of all alleles and an epistatic effect unique to each multilocus genotype. The genotypic value of an individual G was computed as $G = \epsilon + \sum_{i=1}^{nbLoci} G_i$ where ϵ represents the epistatic effect of the individuals genotype, and G_i represents the additive effect of locus i. The epistatic effect for each unique multilocus genotype was drawn from a normal distribution, and we tested a range of variances of that distribution (V_ϵ=0.05, 0.5, 2.25). The lowest value in this range reflects the variance of the distribution from which mutation effects were drawn for baseline 10-loci simulations, the middle value represented that base variance multiplied by the ten loci, and the highest value represented that base variance multiplied by the number of pairwise combinations between ten loci. We constrained our test of epistasis to ten locus simulations because computing the epistatic effects for 100- and 1000-loci simulations was not computationally feasible. For the same reason, we also limited the maximum number of alleles at each locus to 55, the minimum value possible within quantiNemo2. We assumed the lower burn-in mutation rate and the higher dispersal rate, since these were the conditions for which there was strong genetic swamping in previous simulations.

Overall, the inclusion of epistasis resulted in a greater percentage of replicates that failed to colonize the sink: 11%, 33%, and 19% of replicates for V_ϵ=0.05, 0.5, 2.25, respectively, in comparison to 1% for the baseline architecture without epistasis. Incorporating epistasis resulted in significantly greater variance in colonization lag phase (σ^2=2.3e4, 4.3e4, and 3.4e4) for V_ϵ=0.05, 0.5, and 2.25 respectively) compared to the baseline architecture (σ^2=5.4e3) (Table 3, Figure 5A). Given the heterogeneity of variance, Games-Howell *post hoc* tests were performed to compare the means. Lag phase duration did not differ significantly among the nonzero values

of V_ϵ, but mean lag phase duration was significantly longer for the two highest values of V_ϵ=0.5 and V_ϵ=2.25 in comparison to the baseline architecture without epistasis (Table 3). We did not observe a shift in the mode of adaptation for those replicates that did succeed in colonizing the sink (Figure 5B).

Figure 5. A) Duration of colonization lag phase for each simulation replicate for the three values of epistatic variance (V_ϵ) compared with the baseline simulations with no epistasis and assuming a low burn-in mutation rate (μ=10^{-5}) and high dispersal (d=10^{-2}). B) Proportion of adaptive alleles originating from each source population for the three values of V_ϵ in comparison to baseline

	Degrees of freedom	F	p
epistasis	3	8.671	1.39e-05
residual error	396		

Post hoc Games-Howell tests

pair	difference	lower	upper	t	degrees of freedom	p
0.05-0	31.66	-12.45	75.77	1.87	142.13	0.247
0.5-0	96.87	39.64	154.10	4.41	123.40	1.3e-4
2.25-0	53.49	1.79	105.19	2.69	129.44	3.96e-2
0.5-0.05	65.21	-1.57	131.99	2.53	182.26	5.84e-2
2.25-0.05	21.83	-40.32	83.98	0.91	191.41	0.799
2.25-0.5	-43.38	-115.28	28.52	1.56	195.43	0.402

Table 3. Effects of epistasis on colonization lag time. Top: Levene's test for homogeneity of variance for varying levels of epistasis ($V_\epsilon = 0$, 0.05, 0.5, and 2.25). Bottom: pairwise comparisons of means using *post hoc* Games-Howell tests.

Introgression vs. transgressive segregation

So far, we have interrogated how genetic architecture may influence the timing of colonization and mode of adaptation by hybrid lineages, but we have not considered post-colonization range expansion and evolution. In section 3.1, we observed two primary, and mutually exclusive, modes of adaptation to the sink: either the recovery of parental genotypes within a trait (as seen for 1- and 10-loci architectures) or the generation of hybrid genotypes (as seen for 100- and 1000-loci architectures). Since we assumed additive contributions to the genotypic value for those baseline simulations, we can infer that complementary gene action explains the presence of adaptive alleles from both sources for 100- and 1000-loci architectures. Complementary gene action can enable the expression of transgressive phenotypes in hybrid populations if each parental population contributes alleles differing in sign (Rieseberg et al., 1999). This leads us to predict that, for 100 and 1000-loci architectures, but not for 1 and 10-loci architectures, hybrid lineages will be capable of further adaptation to optima exceeding that of either source patch.

In order to test this prediction, we conducted simulations in which an additional patch (a secondary sink) was added adjacent to the primary sink with an environmental optimum twice that of the primary sink for each trait (Figure 1, $\Delta Z_{opt}=16$ for the secondary sink). We allowed one-way dispersal from the primary to the secondary sink. We then monitored colonization and adaptation to the secondary sink by tracking occupancy and mean genotypic value over time. Since mutation was prohibited during the colonization phase, a hybrid population capable of

adapting to the extreme optima of the secondary sink would necessarily do so via transgressive segregation of standing variation. We tested baseline architectures in the absence of linkage assuming a high burn-in mutation rate ($\mu=10^{-4}$) and high dispersal rate ($d=10^{-2}$), since these parameters yielded the shortest lag phases across all architectures in section 3.1.

In agreement with our prediction, we found that the 100-loci and 1000-loci architectures were capable of colonizing the secondary sink while the 1-locus and 10-loci architectures were unable to do so, and that colonization of the secondary sink corresponded with adaptation to its extreme optima (Figure 6).

Figure 6. Mean genotypic value in the secondary sink plotted over time for each simulation replicate (black lines). Genotypic values of 4/-4 represent the optima of source patches, whereas genotypic values of 8/-8 represent the optima of the secondary sink. Grey bars denote when carrying capacity was reached in the secondary sink for each simulation replicate, and white diamonds mark the mean.

Discussion

The steady accumulation of examples of colonization and invasion by hybrid populations in recent decades (Ellstrand and Schierenbeck, 2000; Schierenbeck and Ellstrand, 2008) has made it a priority in the field of invasion genetics to understand the genetic factors underpinning the contribution of hybridization to colonization (Bock et al., 2015). It is now understood that

hybridization can contribute to colonization by producing an overall increase in standing genetic variation, the transfer of adaptive alleles between genetic backgrounds (adaptive introgression), and/or the production of novel genotypes in hybrids (transgressive segregation) (Pfennig et al., 2016).

Our aim here was to test, using individual based simulations, how the architecture of invasiveness traits can influence the contribution of hybridization to colonization. We focused on two underlying predictions: (1) that colonization lag phase duration would be sensitive to genetic architecture and (2) that architectures resistant to genetic swamping would produce shorter lag phases. Given our results, we then tested an additional prediction that the filtering effect of colonization on genetic architecture would influence the ability of resulting colonist lineages to occupy and adapt to more extreme environments.

We found two distinct categories of genetic architecture that facilitate rapid colonization by hybrid populations, doing so in fundamentally different ways. The first category included architectures with few loci and/or tightly linked loci. The mode of adaptation observed for such architectures was the recovery of adaptive parental genotypes. Because this pattern is akin to adaptive introgression, we refer to this as an 'adaptive introgression-type architecture'. The second category included architectures with many freely recombining loci and high V_g, in which the mode of adaptation was the generation of novel hybrid genotypes. Because this second pattern is more akin to transgressive segregation, we refer to this as a 'transgressive segregation-type architecture'.

We found that the extremes within these categories produced the shortest lag phases, with a single locus or a single linkage group producing shorter lag phases than 10 loci or 10 linkage groups for 'adaptive introgression'-type architectures, and 1000 loci producing shorter lag phases

than 100 loci for 'transgressive segregation'-type architectures. These results support our first

prediction that lag phase duration is sensitive to the genetic architecture of invasiveness traits.

The second prediction, that genetic architectures resistant to genetic swamping produce

shorter lag phases, is supported by two lines of evidence. First, when V_g in the source

populations was high, architectures with many loci succeeded in colonization at both low and

high dispersal rates, but when V_g in source populations was low, increased dispersal resulted in

colonization failure across all replicates. The increase in lag duration with increased dispersal

suggests that the establishment of adaptive genotypes was hindered by a greater influx of

maladaptive alleles. This is consistent with previous studies of adaptation across clines that have

shown that such architectures are vulnerable to genetic swamping (Yeaman and Otto, 2011;

Yeaman and Whitlock, 2011). Our observation that high V_g counteracted increased dispersal, and

in fact reduced lag phase significantly for 1000-loci architectures, is consistent with Yeaman

(2015), who found that the effects of genetic swamping may be mitigated by sufficiently high V_g

for architectures dominated by alleles of small effect. Our model demonstrates that this

mitigating effect of V_g on the strength of genetic swamping can significantly reduce lag phase

duration for hybrid invasions with such architectures.

The second line of evidence supporting the prediction that genetic architectures resistant to

genetic swamping produce shorter lag phases is that incorporating linkage rescued some

simulations in which colonization would otherwise fail. In particular, for the many-locus

architectures, when V_g was low and dispersal was high, colonization failed in the case of free

recombination but succeeded in the presence of linkage. Incorporating linkage also resulted in a

corresponding shift in the mode of adaptation towards the recovery of parental genotypes. This

suggests that by reducing recombination between loci, linkage facilitates the recovery of

adaptive parental haplotypes despite a steady influx of maladaptive migrant alleles, which is consistent with previous research demonstrating that linkage mitigates the effects of genetic swamping in hybridizing populations (Yeaman and Whitlock, 2011).

A consequence of the longer lag phases observed for architectures prone to swamping is that colonization of the sink often failed to occur within our limited window of observation. For instance, many-loci architectures failed to colonize the sink when V_g was low and dispersal was high. This suggests that these architectures may be under-represented among those contributing to successful hybrid invasions, particularly for recent introductions and organisms with long generation times, because the time since introduction would often be too short. Our results also suggest that, for those hybrid invasions that *are* successful, invasion-related traits would be enriched for 'adaptive introgression'-type architectures, since these facilitate rapid colonization across the widest range of parameter space.

Strong epistatic interactions may, on average, hinder colonization despite accelerating it in some instances. For 10-loci architectures, we found that the inclusion of epistasis produced greater variance in colonization lag phase and a greater percentage of replicates that failed to colonize the sink. Lag phase was significantly longer when epistatic variance was high in comparison to baseline architectures without epistasis. Interestingly, epistasis had no observable effect on the mode of adaptation to the sink. Further study will be needed to determine if these findings can be generalized to architectures with more loci.

Our results point to an interesting a tradeoff between the modes of adaptation associated with the two extremes of swamping-resistant architectures. 'Adaptive introgression'-type architectures produce the shortest invasion lag phases across the widest range of parameter space, with 1-locus and single linkage group architectures producing lags with mean duration on

the order of ten generations (Figures 2 and 4). Such a capacity to rapidly colonize environments within the bounds of parental trait values may result in initial invasions that spread rapidly. However, such architectures necessarily limit adaptive traits to those that can be obtained by reconstituting parental genotypes, and further adaptation to novel environments beyond the bounds of parental phenotypes may be mutation-limited (Figure 6). By contrast, 'transgressive segregation'-type architectures produce relatively longer lag phases and fail to colonize the sink under some conditions of V_g and dispersal. Once established, however, these hybrid lineages demonstrate a capacity to adapt to environmental conditions beyond the range of parental populations. Transgressive segregation-type architectures thus may facilitate invasions that can spread into a wider variety of novel habitats (Figure 6).

As noted recently by Marques *et al.* (2019), admixture-derived large-effect haplotypes and the generation of novel genotypes via transgressive segregation are may be frequent sources of adaptive variation for speciation and adaptive radiation. While our simulations do not interrogate the contribution of admixture variation to reproductive isolation *per se*, the colonization of novel or extreme environments by hybrids can promote ecological differentiation between hybrid and parental populations, which in turn fosters conditions favorable for speciation and adaptive radiation (Abbott et al., 2013; Seehausen, 2004). Given our finding that only successful hybrid colonists with 'transgressive segregation'-type architectures were capable of further colonization and adaptation to more extreme environments, we would predict that such colonists are more likely to generate new hybrid species and adaptive radiations than those with 'adaptive introgression'-type architectures.

We recognize that the model we present is a crude representation of the complexity of a hybrid invasion, and we have made many simplifying assumptions in order to focus on the role

of genetic architecture. For one, we assume the absence of genetic incompatibilities (ie. Dobzhansky-Muller Incompatibilities or DMIs) between the source populations. DMIs are known to influence the genomic landscape of introgression (Harrison and Larson, 2014), and undoubtedly play an important role in determining when and how hybridization contributes to colonization success. By assuming a lack of DMIs, our model therefore most closely represents hybridization between recently diverged populations.

A second assumption is that we can consider the two traits of interest in isolation from each other and in isolation from other traits not directly related to colonization. We deliberately prevented selective interference between the two adaptive traits in our model by assuming free recombination between them. Empirical evidence does suggest that range expansions may commonly involve relatively few adaptive traits (Bock et al., 2015), but it may be important to consider both interference between those traits and, especially in hybrids, other traits under selection in the hybrids. With many segregating polymorphisms under selection, there is considerable opportunity for selective interference (Comeron and Kreitman, 2002; McVean and Charlesworth, 2000), which could delay or prevent adaptation to a novel environment. In a hybrid, segregating loci with formerly fixed differences, affecting traits unrelated to colonization *per se*, may also be newly visible to selection. Thus, not only could selective interference between loci from the two different traits potentially hinder colonization, but this would be exacerbated in a more realistic scenario with many more traits segregating for genetic variation in the hybrids.

A third assumption is that new mutations do not accumulate during the colonization phase and thus do not contribute to colonization of the sink. Range expansion stemming from *de novo* mutation has been thoroughly studied theoretically (Gilbert and Whitlock, 2017; Gomulkiewicz

et al., 1999; Holt et al., 2003; Kirkpatrick and Barton, 1997), and could feasibly contribute to adaptation within novel environments on timescales relevant for contemporary invasions (Bock et al., 2015; Dlugosch et al., 2015). Moreover, the time required for beneficial mutations to accumulate is another potential genetic cause of invasion lag phases (Crooks and Soulé, 1999; Holt et al., 2003). In reality, hybrid invasions may involve a combination of mutation and gene flow, and our model only focuses on the latter.

A final assumption is that the lag phase is limited only by local adaptation. Although this was a necessary assumption in order to interrogate the potential role of genetic architecture on lag phase duration, there are other factors that could contribute to the invasion lag phase. These include overcoming Allee effects (Aikio et al., 2010), the time to reach sexual maturity (Wangen and Webster, 2006), changes to the receiving environment (Crooks, 2005; Crooks and Soulé, 1999), and, as discussed above, the accumulation of new mutations.

Despite these simplifying assumptions, our results allow us to make a number of robust conclusions. Primarily, we demonstrate that genetic architecture can influence when and how hybridization contributes to colonization. We find that two categories of swamping-resistant genetic architectures produce short colonization lag phases, and that they do so by qualitatively different modes of adaptation. We also find that there is a tradeoff between the two categories of colonist genetic architectures which may influence future adaptation to more extreme environments.

Acknowledgements

We thank Morgan Holder and Wesley Price for assistance in conducting initial explorations of the simulation model and software.

Authorship contributions

BR and TJV conceived the work and designed the methodology. BR carried out the simulations and performed data analysis. BR initially drafted, and BR and TJV jointly revised the manuscript.

Literature Cited

Abbott, R., Albach, D., Ansell, S., Arntzen, J.W., Baird, S.J.E., Bierne, N., Boughman, J., Brelsford, A., Buerkle, C.A., Buggs, R., et al. (2013). Hybridization and speciation. J. Evol. Biol. *26*, 229–246.

Aikio, S., Duncan, R.P., and Hulme, P.E. (2010). Lag-phases in alien plant invasions: separating the facts from the artefacts. Oikos *119*, 370–378.

Bock, D.G., Caseys, C., Cousens, R.D., Hahn, M.A., Heredia, S.M., Hübner, S., Turner, K.G., Whitney, K.D., and Rieseberg, L.H. (2015). What we still don't know about invasion genetics. Mol. Ecol. *24*, 2277–2297.

Comeron, J.M., and Kreitman, M. (2002). Population, Evolutionary and Genomic Consequences of Interference Selection. Genetics *161*, 389–410.

Crooks, J.A. (2005). Lag times and exotic species: The ecology and management of biological invasions in slow-motion. Ecoscience *12*, 316–329.

Crooks, J.A., and Soulé, M.E. (1999). Lag times in population explosions of invasive species: causes and implications. Invasive Species Biodivers. Manag. Based Pap. Present. NorwayUnited Nations UN Conf. Alien Species 2nd Trondheim Conf. Biodivers. Trondheim Nor. 1-5 July 1996 103–125.

Dlugosch, K.M., Anderson, S.R., Braasch, J., Cang, F.A., and Gillette, H.D. (2015). The devil is in the details: genetic variation in introduced populations and its contributions to invasion. Mol. Ecol. *24*, 2095–2111.

Ellstrand, N.C., and Schierenbeck, K.A. (2000). Hybridization as a stimulus for the evolution of invasiveness in plants? Proc. Natl. Acad. Sci. *97*, 7043–7050.

García-Ramos, G., and Kirkpatrick, M. (1997). Genetic Models of Adaptation and Gene Flow in Peripheral Populations. Evolution *51*, 21–28.

Gilbert, K.J., and Whitlock, M.C. (2017). The genetics of adaptation to discrete heterogeneous environments: frequent mutation or large-effect alleles can allow range expansion. J. Evol. Biol. n/a-n/a.

Gilbert, K.J., Sharp, N.P., Angert, A.L., Conte, G.L., Draghi, J.A., Guillaume, F., Hargreaves, A.L., Matthey-Doret, R., and Whitlock, M.C. (2017). Local Adaptation Interacts with Expansion Load during Range Expansion: Maladaptation Reduces Expansion Load. Am. Nat. *189*, 368–380.

Gomulkiewicz, R., Holt, R.D., and Barfield, M. (1999). The Effects of Density Dependence and Immigration on Local Adaptation and Niche Evolution in a Black-Hole Sink Environment. Theor. Popul. Biol. *55*, 283–296.

Harrison, R.G., and Larson, E.L. (2014). Hybridization, Introgression, and the Nature of Species Boundaries. J. Hered. *105*, 795–809.

Holt, R.D., Gomulkiewicz, R., and Barfield, M. (2003). The phenomenology of niche evolution via quantitative traits in a "black-hole" sink. Proc. R. Soc. B Biol. Sci. *270*, 215–224.

Kirkpatrick, M., and Barton, N.H. (1997). Evolution of a Species' Range. Am. Nat. *150*, 1–23.

Marques, D.A., Meier, J.I., and Seehausen, O. (2019). A Combinatorial View on Speciation and Adaptive Radiation. Trends Ecol. Evol.

McVean, G.A., and Charlesworth, B. (2000). The effects of Hill-Robertson interference between weakly selected mutations on patterns of molecular evolution and variation. Genetics *155*, 929–944.

Neuenschwander, S., Hospital, F., Guillaume, F., and Goudet, J. (2008). quantiNemo: an individual-based program to simulate quantitative traits with explicit genetic architecture in a dynamic metapopulation. Bioinformatics *24*, 1552–1553.

Pfennig, K.S., Kelly, A.L., and Pierce, A.A. (2016). Hybridization as a facilitator of species range expansion. Proc. R. Soc. B Biol. Sci. *283*, 20161329.

Rieseberg, L.H., Archer, M.A., and Wayne, R.K. (1999). Transgressive segregation, adaptation and speciation. Heredity *83*, 363–372.

Schierenbeck, K.A., and Ellstrand, N.C. (2008). Hybridization and the evolution of invasiveness in plants and other organisms. Biol. Invasions *11*, 1093.

Seehausen, O. (2004). Hybridization and adaptive radiation. Trends Ecol. Evol. *19*, 198–207.

Tigano, A., and Friesen, V.L. (2016). Genomics of local adaptation with gene flow. Mol. Ecol. *25*, 2144–2164.

Wangen, S.R., and Webster, C.R. (2006). Potential for multiple lag phases during biotic invasions: reconstructing an invasion of the exotic tree Acer platanoides. J. Appl. Ecol. *43*, 258–268.

Yeaman, S. (2015). Local Adaptation by Alleles of Small Effect. Am. Nat. *186*, S74–S89.

Yeaman, S., and Otto, S.P. (2011). Establishment and Maintenance of Adaptive Genetic Divergence Under Migration, Selection, and Drift. Evolution *65*, 2123–2129.

Yeaman, S., and Whitlock, M.C. (2011). The Genetic Architecture of Adaptation Under Migration–Selection Balance. Evolution *65*, 1897–1911.