Long-term reciprocal gene flow in wild and domestic geese reveals complex domestication history

Marja E. Heikkinen[1,2], Minna Ruokonen[1§], Thomas A. White[1,3], Michelle M. Alexander[4], İslam Gündüz[5], Keith

M. Dobney[6,7,8], Jouni Aspi[1], Jeremy B. Searle[2]*, Tanja Pyhäjärvi[1,9]*

§ deceased

* contributed equally to this paper

[1] Department of Ecology and Genetics, PO Box 3000, Fi-90014 University of Oulu, Finland

[2] Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York 14853, USA

[3] CMPG, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland

[4] University of York, BioArCh, Environment Building, Wentworth Way, Heslington, York, YO10 5NG

[5] Department of Biology, Faculty of Arts and Sciences, University of Ondokuz Mayis, Samsun, Turkey

[6] Department of Archaeology, Classics and Egyptology, University of Liverpool, 12–14 Abercromby Square,

Liverpool L69 7WZ, UK

[7] Department of Archaeology, University of Aberdeen, St Mary's, Elphinstone Road, Aberdeen, AB24 3UF,

UK

[8] Department of Archaeology, Simon Fraser University, Burnaby, B.C. V5A 1S6, 778-782-419, Canada

[9] Biocenter Oulu, PO Box 5000, Fi-90014 University of Oulu, Finland

Author for correspondence:

Marja E. Heikkinen

email: marja.e.heikkinen@oulu.fi

ORCID:

MEH, 0000-0002-7051-5654; MMA, 0000-0001-8000-3639; İG, 0000-0002-6436-5397; KMD, 0000-0001-

9036-4681; JA, 0000-0002-2451-3201; JBS, 0000-0001-7710-5204; TP, 0000-0001-6958-5172

Abstract

Hybridisation can substantially impact genetic diversity of domesticates, but this situation has not been

widely studied. We analysed the impact of hybridisation on the genome-wide genetic variation in current

populations of the European domestic goose and its wild progenitor: the greylag goose (*Anser anser*). Our

dataset consisted of 58 wild greylag and 75 domestic geese genotyped for 33,527 single nucleotide

polymorphisms (SNPs) identified by genotyping-by-sequencing (GBS). Demographic reconstruction and

clustering analysis suggested that divergence between wild and domestic geese around 5,300 generations

ago was followed by long-term genetic exchange, and that greylag populations have 3.2–58.0% admixture

proportions with domestic geese, that also varies geographically. Many modern European breeds share

considerable (> 10%) ancestry with Chinese domestic geese (derived from the swan goose *A. cygnoid*). This

study shows a domestication process progressing despite continued and pervasive gene flow from the wild

form.

Keywords: *Anser anser*, domestication, domestic goose, genotyping-by-sequencing, hybridisation,

population genomics

1. Introduction

Reproductive isolation is a defining feature of speciation and yet hybridisation between species is an

important general phenomenon in trait evolution [1,2]. Among birds, the Anseriformes (ducks, geese and

swans) show particularly pervasive hybridisation, 41.6% to > 60% of species hybridising with each other

[3,4]. Domestication generates differentiated gene pools and reproductive isolation between domestics

and their wild progenitor, but hybridisation between domestic and wild forms has been well demonstrated

in both plants [1,5] and animals [6,7]. The impacts include genetic and trait enrichment of domestics, for

instance, in chicken the acquisition of a yellow skin phenotype is a result of past mating between red

junglefowl and grey junglefowl [8]. In geese, the high tendency for hybridisation among wild species is also

shown among wild and domestic forms [9,10], creating an exciting opportunity to study the complex

dynamics of hybridisation and domestication.

The domestic geese of the world (European and Chinese forms) are derived from two different wild

species: the greylag (*Anser anser*) and the swan goose (*A. cygnoid*) [11,12]. *A. anser* and *A. cygnoid* shared a

common ancestor about 3.4 Mya [13], but are still able to hybridise [4], and some domestic breeds are

reportedly hybrid [14]. The greylag has been divided into the western, nominate subspecies *A. a. anser*

(Linnaeus, 1758) with a European breeding range and the eastern subspecies *A. a. rubrirostris* (Swinhoe,

1871) breeding further east, although the subspecific boundary is not well defined, and mitochondrial DNA

has not been found to distinguish them [10]. Of these subspecies, *rubrirostris* is larger and lighter coloured

than *anser* [15] and has a pink bill and cold pink legs in contrast to the orange bill and flesh-coloured legs of

anser, the bill colour used as primary evidence in favour of the original domestication of *rubrirostris* [16]. As

with all domesticates, domestic geese varieties are morphologically more diverse than their wild

counterparts, particularly in plumage and body size [14].

The current knowledge about goose domestication relies largely on ancient texts and archaeological

evidence. Questions about where and when domestication took place, the genetic changes associated with

it and the later history of domestic geese, however, remain largely unresolved [10]. There are depictions

from the New Kingdom of Egypt that suggest geese were already fully domesticated by the 18[th] Dynasty

(1450-1341 BCE). The earliest reliable reference to domestic geese in western Eurasia is Homer's Odyssey

(first half of 8[th] century BCE) and geese were certainly well-established poultry by Roman times [17].

Genetic diversity in the mitochondrial DNA (mtDNA) of greylag and European domestic geese

showed reduced diversity in the domestics [10] which may result from an early domestication bottleneck

or, alternatively, later breed formation. There is a particular mitochondrial haplogroup common in the

domestics [10], and archaeological domestic goose bones from the High Medieval (11[th]-13[th] century CE) of

Russia belonged to that haplogroup [18].

MtDNA relationships between extant Chinese and European domestic goose breeds confirm that the

former, excluding one breed, have swan goose ancestry, whereas European domestic goose and the

Chinese Yili breed have greylag ancestry [12,19,20]. However, Chinese mtDNA haplotypes may occasionally

occur in European domestics, and vice versa [10,19].

Genomic data can be much more powerful than mtDNA in hybridisation inference, and has been

applied, for instance, to New World cattle, which along with their taurine ancestry were shown to have a

greater proportion of indicine ancestry than previously assumed [21]. Interpretation is still challenging and

it is best to infer jointly the genetic impact of initial domestication and subsequent hybridisation of wild and

domestic populations, as the latter can obscure domestic-wild genetic relationships and may also give a

false impression of the number of times a species has been domesticated [22–24].

Here we investigate goose domestication history using genome-wide single nucleotide polymorphism

(SNP) data from thousands of loci, obtained by genotyping-by-sequencing (GBS). We used 56 and 50

samples of greylag and domestic geese from a previous mtDNA study [10], together with 2 new Turkish

greylag and 25 new domestic specimens. We studied the interplay between domestication and

hybridisation by addressing the following questions: i) what is the extent of genetic differentiation amongst

wild and domestic geese? ii) what is the approximate time of domestication? and iii) what is the role of

intra- and interspecific hybridisation in goose domestication history and the current genetic makeup?

2. Materials and methods

a) Sampling

The greylag wild-collected samples derive widely from Eurasia (Figure 1, Table S1). The European domestic

goose samples represented 14 different breeds (Table S1) and individuals that did not belong to any

recognised breed or were presumed to be crosses between European and Chinese domestic geese. Some

specimens were reported to be Chinese domestic geese. The domestic samples were obtained from local

breeders in Denmark, Sweden and the UK and those from Turkey were collected directly by the authors.

b) DNA extraction and GBS library construction

GBS [25] libraries were constructed at the Cornell Biotechnology Resource Center (BRC) following DNA

extraction with the DNeasy Blood and Tissue Kit (QIAGEN) with RNase treatment. Each individual DNA

sample and an adaptor with a unique barcode were combined in a 96-well plate along with a common

adaptor. Samples were treated with the EcoT-22I (ATGCAT) restriction enzyme to create fragmented DNA.

Barcoded adapters and common adapters with matching sticky ends were ligated to each sample with T4

DNA ligase. The samples were pooled and purified with a QIAquick PCR Purification Kit (QIAGEN). PCR

amplification of the library used primers complementary to barcoded and common adapters with products

purified as above, and the samples were 100 bp SE-sequenced with Illumina HiSeq 2000/2500 at the BRC.

c) GBS pipeline and SNP calling

Raw sequence reads were run through the Command Line Interface of the Tassel 5 GBS v2 Discovery and

Production pipelines [26]. Details about the pipelines and SNP calling are in the Electronic Supplementary

Material, ESM. Good quality reads were recorded as tags and aligned to the *A. cygnoid domesticus*

GenBank assembly (AnsCyg_PRJNA183603_v1.0 GCF_000971095.1) [27] using the Burrows-Wheeler Aligner

with default settings [28]. After running the raw data through the pipelines, 69,865 SNPs were obtained.

The SNPs were subjected to additional filtering using VCFtools [29]. We removed indels, loci with

more than two alleles and invariant loci. However, invariant loci were retained for phylogenetics, informing

about greylag-swan goose divergence. After preliminary analyses loci with observed heterozygosity over

0.75 were removed as potential paralogs. Individuals with more than 20% missing data across loci were

removed. The final dataset consisted of 33,527 biallelic SNPs and 133 individuals (58 wild and 75 domestic).

d) Population structure analyses

Population clustering and structure was analysed with STRUCTURE 2.3.4 [30] and Principal Component

Analysis (PCA) [31]. For the whole dataset, STRUCTURE was run with 1000 burn-in steps followed by 10 000

iterations of MCMC for data collection for K = 1-10 allowing admixture with five replicates of each run to

reach convergence. For the STRUCTURE analyses done separately on greylags and European domestic

geese, see ESM. An admixture model with correlated allele frequencies among populations [32] was used in

all STRUCTURE analyses and the iterations were automated with StrAuto 1.0 [33]. We applied both

likelihood of K and ΔK [34] of successive K values to determine the optimal number of clusters, using

STRUCTURE HARVESTER [35]. CLUMPP 1.1.2 [36] was used to align the assignments from different

replicates of K and DISTRUCT 1.1 [37] for visualisation. A PCA was performed with the prcomp function in R

[38] and the significance of eigenvalues determined based on the Tracy-Widom distribution [22,31].

A neighbor-joining tree was constructed for phylogenetic analysis, with pairwise distance between

individuals obtained with the R package ape [39] based in 40,191 loci. The *A. cygnoid* reference genome

and the invariant sites that differed from it were included in the tree construction.

e) Tests for admixture and simulations of demographic history

The history of admixture was tested with a 3-Population test f_3(C; A, B) implemented in AdmixTools 4.1 [40,

ESM]. This method offers a formal test to explain observed patterns of admixture in a target population

without an outgroup. For identification of admixture between Chinese and European domestics, Grey and

White Chinese were combined to represent the Chinese, and the Landes breed that had minimum

indication of admixture in STRUCTURE was chosen to represent the European domestic source population.

Different models of demographic history were tested with fastsimcoal2 ver 2.6 [41]. We excluded all

SNPs that had missing data within the whole data set and executed the analyses with a site frequency

spectrum (SFS) based on 9,212 SNPs. As there are no estimates of the genetic diversity per base pair for

greylags, we estimated the proportions of variable and monomorphic sites in the data. From the BAM file

with –depth option in SAMtools 1.7 [42], we estimated 9,801,382 bp covered with GBS tags. We then

mimicked the filtering steps done for the biallelic SNPs to reduce the total number of sites in equivalent

proportions. We removed the same number of sites that corresponded to the number of SNPs that were

removed because they were indels, had more than 2 alleles or had heterozygosity over 0.75. Since some of

the SNPs were removed from this analysis due to missing data in some individuals, we removed an equal

proportion of sites from the total number of sites as well. The final SFS had 1,681,316 sites.

To infer the demographic history, we chose a subset of individuals from both wild-collected greylags

and domestic geese to represent the genetic variation in both groups. Therefore, 11 greylags with > 90.8%

of greylag ancestry and 15 domestic geese with > 91.4% of European domestic goose ancestry were

selected for the analysis. The mutation rate for the simulations was 1.38×10^{-7} [43]. The parameter

estimation for each model tested involved 100,000 simulations and 40 conditional maximization (ECM)

cycles. The parameters for each model were estimated with 100 independent runs to obtain the global

maximum. The models tested were i) simple divergence of two populations with no migration, ii)

divergence of two populations with migration and iii) divergence of two populations with two different

migration matrices (Figure 2). The best model was selected based on Akaike's weight of evidence as in [41].

For parametric bootstrapping 100 SFS were simulated with the parameter estimates obtained from the real

SFS, followed by maximum likelihood estimation with 50 independent runs for each bootstrap SFS. The 95%

confidence intervals were obtained from the bootstrap data for each estimated parameter.

f) The estimation of genetic diversity

Genetic diversity and pairwise F_{ST} values were investigated with the *hierfstat* R package [44]. Expected

heterozygosity (H_E) was calculated for each locus and population and averaged across loci. Difference in

average H_E between greylags and European domestics was tested with a two-sample t-test with the Welch

correction for non-homogeneity of variance [45]. For comparing the genetic diversity among wild and

domestics, only pure greylag populations (< 10% admixture with domestic geese) and pure European

domestic geese (< 10% admixture with Chinese domestic geese) were used to avoid hybridisation effects on

the estimates. The admixture proportions were obtained from STRUCTURE.

The variance components across loci for hierarchical F-statistics for pure greylags and pure European

domestics were estimated using locus-by-locus analysis of molecular variance (AMOVA) implemented in

Arlequin 3.5.2.1 [46]. The significance was tested with 16 000 permutations.

3. **Results**

a) Population structure

There was clear divergence between greylags and both European and Chinese domestic geese according to

STRUCTURE and PCA (Figure 3a-b). At $K = 2$, populations/breeds are clustered based on their status (wild or

domestic) and, at $K = 3$, domestic geese are further separated into European and Chinese. At $K = 4$, the

fourth cluster is admixed within greylag populations but none of the individuals are unanimously assigned

to that cluster. The likelihood was highest for $K = 3$. These results were supported by PCA as the first two

PCs out of 14 significant PCs ($p < 0.05$) were enough to separate the three groups (wild, European domestic,

Chinese domestic) from each other (Figure 3a). Overall the greylag populations showed 3.2% - 23.5%

admixture proportions with European domestic geese when $K = 3$. In contrast, not all European domestic

geese showed admixture with greylags and the admixture percentages ranged from 0.0 to 8.4%. At $K = 3$

many European domestic goose breeds showed mixed ancestry with Chinese domestic geese (0.0-27.1%).

The neighbor-joining tree repeated the major patterns observed with STRUCTURE and PCA revealing

a star shaped phylogeny and confirming that the domestic and greylag geese largely form different clades

(ESM, Figure S1). Surprisingly, the Chinese domestic geese were closer to European domestic geese and

greylags, than to the swan goose reference genome. In addition, one greylag from Turkey was more closely

related to the Chinese domestic geese than other greylags, also indicated by admixture proportions from

STRUCTURE. Further, two Crested Faroese individuals and four domestics from the USA (2 unknown and 2

Toulouse crosses) were closer to Chinese than European domestic geese. These six individuals also showed

high proportions of admixture with Chinese domestics in the STRUCTURE analysis.

Some further population structure was observed within both greylags and domestic geese, when

analysed separately with STRUCTURE and PCA. Geographically, greylags differentiated by subspecies (ESM,

Figures S2-S3). STRUCTURE indicated little differentiation among European domestic geese, but the PCA

revealed separation between the European breeds and the Turkish domestic geese (ESM, Figures S4-S5).

b) Admixture and the time of domestication

Although, STRUCTURE implied considerable mixed ancestry from multiple genetic clusters for Dutch and

Turkish greylags, the f_3 analysis did not confirm admixture for these populations. The two Turkish greylag

samples came from the same area as our NW-Turkish domestic population, which among Turkish domestic

geese showed highest admixture with greylags (2.2%), but admixture was not confirmed with the f_3 test.

The f_3 analysis confirmed admixture of domestic geese in line with STRUCTURE results. Most notably,

the African breed is a hybrid between European and Chinese domestic geese (Z-score -6.399), unexpected

as this breed has been assumed to have originated solely from swan goose. The European-Chinese hybrid

status of the Kholmogor and Steinbacher breeds was also confirmed (Z-scores of -8.933 and -5.349,

respectively). The Kholmogor breed also fell halfway between European and domestic geese both in

STRUCTURE and PCA, whereas the Steinbacher was genetically closer to European domestic geese in the

PCA. However, the Diepholzer breed, which reportedly is also a hybrid, was not confirmed as such in our

analysis. Other domestic breeds/groups with admixture status in STRUCTURE were also confirmed to have

a European-Chinese admixture when a Z-score threshold of -3 (roughly corresponding to $p < 0.01$) was

used: Sebastopol, Toulouse cross, Domestic NY, Embden, Tufted Roman (Table S2, Figure 3c). The Crested

Faroese breed gave indication of admixture based on STRUCTURE analysis and the f_3 test supported this (Z-

score of -2.228, p < 0.05). Surprisingly, the Northern Turkish domestic population was not admixed with

Chinese domestic geese in STRUCTURE but f_3 analysis gave a contrasting signal (Z-score -2.459, p < 0.05).

The demographic model that best fit our data suggested divergence of greylag and domestic geese

with a recent migration rate change (Table 1, Table S3). The model suggested divergence around 5319

generations ago (95% confidence intervals (CI): 1538-8225) with asymmetric but close to equal migration

rates from greylags to domestic geese following divergence. About 159 (79-462) generations ago, the

migration matrix changed, suggesting higher gene flow from domestic geese to greylags towards modern

times. Given an estimated generation time for these geese of about 3 years, this suggests divergence about

14 000 BCE and gene flow shift about 480 years ago.

c) Genetic diversity

An AMOVA was used to partition genetic diversity among greylag vs. domestic (group level), and among

populations (greylag) and among breeds (domestics), and within population levels (Table 2). The fixation

index between greylag and domestic geese was 0.158 and there was also significant differentiation among

greylag populations/domestic breeds. The average pairwise F_{ST} between greylag populations and domestic

breeds was 0.197, among greylag populations 0.088 and among domestic breeds 0.174 (Table S4).

The genetic diversity measured as average H_E was higher in pure greylags (0.146) than pure European

domestic geese (0.096) (Welch's t-test, degrees of freedom (df) = 10.594, $p = 3.91 \times 10^{-5}$). The difference in

average H_E remained when non-pure greylag and non-pure European domestics were included in the

comparison (0.156 vs. 0.107; Welch's t-test, df = 19.28, p = 0.000418) (Table S1). The Steinbacher and

African breeds (Table S1) were excluded from both estimates because the Steinbacher is known to be a

European-Chinese cross and the African has swan goose ancestry. The average H_E ranged from 0.140

(Denmark) to 0.150 (Kazakhstan) in pure greylags and from 0.047 (Landes) to 0.123 (Domestic N-Turkey) in

pure European domestics. The average H_E was higher in greylag populations that showed high admixture

proportions with domestic geese, Netherlands (0.157, 13.8% European domestic, 2.2% Chinese domestic)

and Turkey (0.236, 23.5% European domestic, 34.5% Chinese domestic), but not significantly so (Welch's t-

test, df = 1.002, p = 0.421, average H_E: pure greylags 0.146 vs non-pure greylags 0.196). The average H_E was

also higher in European domestics that showed high admixture proportions with Chinese domestics, i.e.

Crested Faroese (0.117, 17% Chinese domestic), Sebastopol (0.133, 11.1% Chinese domestic), Domestic NY

(0.143, 78.6% European domestic, 21.4% Chinese domestic) and Toulouse cross (0.151, 72.9% European

domestic, 27.1% Chinese domestic), and the difference was also statistically significant (Welch's t-test, df =

9.0991, p = 0.0039, average H_E: pure European domestics 0.096 vs non-pure European domestics 0.136).

4. Discussion

We studied the dynamics of domestication and hybridisation in grey (*Anser*) geese using genome-wide SNP

data. The results demonstrated genetic divergence between Eurasian wild greylag and European domestic

geese with long-term genetic exchange between them. We also inferred temporal changes in the direction

of gene flow. The degree of hybridisation between greylag and domestic geese also varied geographically.

Surprisingly, several domestic goose breeds also showed a substantial genetic contribution of Chinese

domestic geese. We also provide insights about the origin and the timing of goose domestication.

a) Genetic diversity and differentiation of greylag and European domestic geese

Domestic species often show reduced genetic diversity compared to their wild ancestor, attributable to

genetic drift during population bottlenecks of initial domestication, combined with subsequent artificial

selection associated with breed formation [47]. Domestic geese appear to follow the same trend. We found

European domestic geese to have lower H_E than wild greylags.

European domestic geese are genetically distinct from their wild progenitor but no more so than for

other domestic birds. The average pairwise F_{ST} values between greylag populations and domestic goose

breeds were lower than between red junglefowl and domestic chicken populations [48], and domestic

geese are less distinctive than domestic pigeons [49]. Among domestic geese, the Turkish are particularly

interesting. From mtDNA, the Turkish domestic geese stand out as the most genetically variable group [10],

and although this is less evident from GBS, among the pure European domestic geese the Northern Turkish

showed the highest average H_E. The f_3 analysis indicates a history of admixture with Chinese domestics for

this population, which may explain its high genetic diversity.

We found a genetic separation between European and Near Eastern populations of greylags that

aligned with the western and eastern subspecies (*A. a. anser* and *A. a. rubrirostris*) [50], a distinction which

could not be made based on mtDNA [10]. Hybridisation between the western and eastern subspecies is

suggested from admixture in Dutch and Danish greylags in STRUCTURE. There is historical evidence for the

introduction of *rubrirostris* to Belgium in 1954 and to Netherlands in 1960s [9,51]; thus, *rubrirostris* genes

may have originated from the recently introduced gene pool spreading to Denmark.

b) When and where were geese domesticated?

Traditional views on goose domestication claim it first occurred in the eastern Mediterranean (possibly

Egypt) around the 3rd Millennium BCE [17,52]. Domestication of chicken and perhaps pigeon took place

earlier, but domestication of duck later, at least in Europe [23]. Demographic modelling suggests that the

wild greylag and related domestic lineages split approximately 5,300 generations ago placing domestication

origins at 14 000 BCE assuming a 3-year generation time [15]. This estimated genetic divergence time is,

admittedly, considerably earlier than any evidence for animal domestication except dog. It is important to

note that the estimated divergence times have large confidence intervals and merely indicates the split

between the ancestors of contemporary wild and domestic lineages. It is most likely that our demographic

modelling reflects the early divergence of different lineages of greylags, only one of which contributed to

later domestication. The subsequent reduction or even disappearance of that wild lineage means that,

despite wide geographical sampling, the possible modern wild population(s) of the greylag progenitor to

domestic geese was not sampled in this study and must, therefore, be sought by ancient DNA approaches.

It is, however, tempting to make our own determination of the location of goose domestication from

the modern data presented here. Given that genetic diversity would be expected to be highest in the

'domestication centre' and reduce with increasing distance from there, the high mtDNA diversity of Turkish

domestic geese means the eastern Mediterranean remains a strong candidate for the origin of goose

domestication, and our nuclear data accord with that. However, as we have shown, hybridisation between

wild and domestic geese can also generate high genetic diversity both within and outside the original

domestication location.

c) The role of intra- and interspecific hybridisation in goose domestication history

i. Evidence of current hybridisation

Domestic animals and their wild relatives are often observed to interbreed, and this is also true for geese.

Both field observations and mtDNA results [9,10] suggested some current hybridisation between domestic

and greylag geese. Genome-wide analysis covering multiple greylag populations and domestic breeds

revealed a considerable impact of hybridisation on genetic diversity of both wild and domestic geese.

Hybridisation is particularly prevalent in certain geographical regions. Dutch and especially Turkish

wild greylag samples had more shared genetic affiliation with domestics than Scandinavian and Finnish

greylag populations (Figure 3b). Some regions may offer more hybridisation opportunities, e.g. climate may

allow greylags to be sedentary year-round and be favourable for keeping domestic geese. The Netherlands,

for instance, lies on the Atlantic flyway offering breeding, staging and wintering areas for greylags [53,54].

Since pair-bonding of geese generally occurs on wintering grounds [55], hubs for migrating geese such as

the Netherlands may permit population mingling. Nevertheless, the f_3 test did not support a simple history

of admixture for the Netherlands. Patterson et al. [40] have stated that population-specific drift may mask

the signal of admixture in such analyses, leading to a non-negative f_3. The f_3 model is relatively simple, with

only two sources, and may not catch the signal of admixture in the Dutch greylag population because of the

previous contribution of *rubrirostris*, which was not included in the model.

Based on ringing data most greylag populations in Scandinavia follow the Atlantic flyway - some of

the geese wintering in the Netherlands and others in southwest Spain. However, Finnish greylags favour

the Central European flyway and winter in North Africa, with a minority of Finnish greylags using the

Atlantic Flyway [53,54]. The Finnish populations of greylag showed the lowest admixture proportions with

domestic geese (S-Finland 3.2% domestic goose, N-Finland 3.3% domestic geese) among greylag

populations. Rearing geese is not a popular practice in Finland and they constituted less than 5% of poultry

kept in Finland in 2014 [56]. The Norwegian populations showed only slightly higher admixture proportions

with domestic geese, although domestic mtDNA haplotype ANS19 was detected from a wild greylag

collected in Finnmark, Norway [57]. This haplotype is a partial sequence of the D5 haplotype identified by

Heikkinen et al. [10], and identical to that found in White Roman domestic geese [58].

Inferring the hybridisation patterns in the Turkish greylags is more complicated, as Turkish greylags

indicate hybridisation with both Chinese and European domestics. Both greylags sampled in Turkey showed

considerable admixture with domestic geese. One of them appeared genetically as a hybrid of European

and Chinese domestic goose with only a small proportion of greylag ancestry, whereas the other one was a

more equal mix of European domestic goose and greylag supplemented by a considerable Chinese

domestic goose ancestry. There is some indication of hybridisation between greylags and domestic geese

within that area as the domestic geese sampled from the same area showed some admixture with greylags,

but this was not confirmed with f_3 analysis. These results may reflect a local practice of keeping captive

greylags within a flock of domestic geese. Another possibility is that the Turkish greylags have hybridised

with some unsampled distinct greylag population and simply appear genetically like domestic geese due to

lack of representation of the unsampled wild population. The greylag population breeding and wintering in

the Black Sea region is not well monitored [59].

ii. Long-term hybridisation

Domestication can be seen as an analogy of speciation where an animal population transforms to an

ecotype that is adapted to the human niche [23] and at later stages of domestication is perpetuated with

reproductive isolation in the form of selection managed by humans [60]. However, this reproductive

isolation may not be complete e.g., in pig [7]. While the genetic divergence of the greylag and its domestic

descendant is evident, our results suggest extensive long-term genetic exchange between them. In

addition, the demographic modelling suggests that the gene flow patterns have changed over time.

Initially, gene flow was greater from greylag to domestic geese. Possibly early goose herders

supplemented the domestic gene pool with wild-caught greylags. Alternatively, wild ganders may have

interbred with domestic geese and contributed to the domestic gene pool. It is unlikely that the early

stages of goose domestication were rigorously managed, allowing matings outside the domestic gene pool.

Both would explain the observed higher gene flow from greylag to domestic geese in the early stages of

domestication. It is in the farmers' interest to keep the domestic geese and wild geese reproductively

isolated to keep control over the traits that are being selected, but artificial selection of traits would have

become possible only after the domestic gene pool had been established. After that, it may occasionally be

beneficial to restock the flock to maintain enough genetic diversity. The natural tendency for imprinting in

geese facilitates this practice. Later, the balance of gene flow shifted towards greylags. This likely occurred

after goose-keeping became well-established in the Medieval period [17]. The rise in number of domestic

geese may have allowed an increase in domestic goose escapees resulting in increased gene flow from

domestic geese back to greylags.

Furthermore, not only have domestic geese admixed with wild greylags but also European and

Chinese domestic geese have hybridised. Hybridisation with ancestral species or closely related species is a

rule rather than an exception in domestic species, e.g., the genetic makeup of chicken derives from

multiple different species of *Gallus* [8]. Similarly, the genetic make-up of domestic geese seems to derive

from two closely related species. This hybridisation with Chinese domestic geese may have introduced

some traits not present in greylags to European domestic geese and vice versa.

5. Conclusion

The genome-wide data reported here reveals that wild and domesticated European and Chinese greylag

and swan goose have intermingled relationships where selection, domestication and hybridisation all play a

role. The two independent domestication events of closely related species provide an excellent system to

further investigate the relative roles of mutation, selection and gene flow. This includes investigation of

potentially adaptive gene flow between the domesticates and identification of shared genome regions

subject to anthropogenic selection and extent of convergent evolution in a domestication setting. A more

detailed analysis of the relationships of wild greylag and swan geese and their domestic descendants would

require wide geographic sampling of both wild species and the domestic breeds combined with an

application of ancient biomolecules and advanced biometrics applied to the remains of geese recovered

from archaeological sites.

Authors' contributions.

MEH conceived the study, contributed to data collection, analysed the data and drafted the manuscript.

MR acquired the funding, conceived the study and contributed to data collection. TAW contributed to data

collection and participated to data analysis. MMA and İG contributed to data collection. KMD, JA, JBS and

TP conceived the study and contributed to writing and interpretation of data, with TP also participating in

data analysis. All authors, excluding MR, reviewed, improved and approved the manuscript.

Competing interests.

The authors declare no conflict of interest.

Funding.

This work was supported by the Academy of Finland (grant no. 131673 to MR, grant no 283609 to JA and

grant no. 287431 to TP); and the Cornell Center for Comparative and Population Genomics (3CPG) to JBS.

The Emil Aaltonen Foundation and Oulun läänin talousseuran maataloussäätiö are acknowledged for the

personal grants awarded to MEH.

Acknowledgements.

We thank the Ministeriet for Fødevarer, Landbrug og Fiskeri in Denmark, the Svenska Lanthönsklubben in

Sweden and the Goose Club in the UK for their help in obtaining the domestic goose samples. Silvia

Markova is acknowledged for her help with the laboratory work. We are grateful for Tiina Mattila for her

help with the fastsimcoal2 analysis. Lastly, we thank the CSC – IT Center for Science in Finland for providing

computing resources.

References

1. Arnold ML. 2004 Natural hybridization and the evolution of domesticated, pest and disease

organisms. *Mol. Ecol.* **13**, 997–1007. (doi:10.1111/j.1365-294X.2004.02145.x)

2. Abbott R, Albach D, Ansell S, Arntzen JW, Baird SJE, Bierne N, et al. 2013 Hybridization and

speciation. *J. Evol. Biol.* **26**, 229–246. (doi:10.1111/j.1420-9101.2012.02599.x)

3. Grant PR, Grant BR. 1992 Hybridization of bird species. *Science* **256**, 193–197.

(doi:10.1126/science.256.5054.193)

4. Ottenburghs J, van Hooft P, van Wieren SE, Ydenberg RC, Prins HHT. 2016 Hybridization in geese: a

review. *Front. Zool.* **13**, 20. (doi:10.1186/s12983-016-0153-1)

5. Janzen GM, Wang L, Hufford MB. 2019 The extent of adaptive wild introgression in crops. *New

Phytol.* **221**, 1173-1664. (doi:10.1111/nph.15457)

6. Godinho R, Llaneza L, Blanco JC, Lopes S, Álvares F, García EJ, et al. 2011 Genetic evidence for

multiple events of hybridization between wolves and domestic dogs in the Iberian Peninsula. *Mol.*

Ecol. **20**, 5154–5166. (doi:10.1111/j.1365-294X.2011.05345.x)

7.   Frantz LAF, Schraiber JG, Madsen O, Megens H-J, Cagan A, Bosse M, et al. 2015 Evidence of long-
term gene flow and selection during domestication from analyses of Eurasian wild and domestic pig
genomes. *Nat. Genet.* **47**, 1141–1148. (doi:10.1038/ng.3394)

8.   Eriksson J, Larson G, Gunnarsson U, Bed'hom B, Tixier-Boichard M, Strömstedt L, et al. 2008
Identification of the yellow skin gene reveals a hybrid origin of the domestic chicken. *PLoS Genet.* **4**,
e1000010. (doi:10.1371/journal.pgen.1000010)

9.   Kuijken E, Devos K. 1996 The status of the greylag goose *Anser anser* in Flanders, Belgium. *Wetlands*
International Goose Specialist Group Bulletin **26**, 8.

10.   Heikkinen ME, Ruokonen M, Alexander M, Aspi J, Pyhäjärvi T, Searle JB. 2015 Relationship between
wild greylag and European domestic geese based on mitochondrial DNA. *Anim. Genet.* **46**, 485–497.
(doi:10.1111/age.12319)

11.   Delacour J. 1954 *The waterfowl of the world : Vol. 1, The magpie goose, whistling ducks, swans and*
geese, sheldgeese and shelducks. 2nd ed. London: Country Life.

12.   Shi X-W, Wang J-W, Zeng F-T, Qiu X-P. 2006 Mitochondrial DNA cleavage patterns distinguish
independent origin of Chinese domestic geese and western domestic geese. *Biochem. Genet.* **44**,
237–245. (doi:10.1007/s10528-006-9028-z)

13.   Ottenburghs J, Megens H-J, Kraus RHS, Madsen O, van Hooft P, van Wieren SE, et al. 2016 A tree of
geese: a phylogenomic perspective on the evolutionary history of True Geese. *Mol. Phylogenet. Evol.*
101, 303–313. (doi:10.1016/j.ympev.2016.05.021)

14.   Buckland R, Guy G, editors. 2002 Goose production. In *FAO Animal Production and Health Paper -*
154, Rome: Food and Agriculture Organization of the United Nations.

15.   Cramp S, Simmons KEL, editors. 1977 *Handbook of the birds of Europe, the Middle East, and North*
Africa: The birds of the Western Palearctic. Vol. 1: Ostrich-Ducks. New York: Oxford University Press.

16.   Kear J. 1990 *Man and wildfowl*. London: T & AD Poyser Ltd.

17.   Albarella U. 2005 Alternate fortunes? The role of domestic ducks and geese from Roman to
Medieval times in Britain. In Feathers, grit and symbolism (ed. G Grupe, J Peters), *Documenta*
Archaeobiologiae **3**, 249–258.

18.   Honka J, Heino M, Kvist L, Askeyev I, Shaymuratova D, Askeyev O, et al. 2018 Over a thousand years
of evolutionary history of domestic geese from Russian archaeological sites, analysed using ancient
DNA. *Genes* **9**, 367. (doi:10.3390/genes9070367)

19.   Sun J, Zhang S, He D-Q, Chen S-Y, Duan Z-Y, Yao Y-G, et al. 2013 Matrilineal genetic structure of
domestic geese. *J. Poult. Sci.* **51**, 130–137. (doi:10.2141/jpsa.0120152)

20.   Ren T, Liang S, Zhao A, He K. 2016 Analysis of the complete mitochondrial genome of the Zhedong
White goose and characterization of NUMTs: reveal domestication history of goose in China and
Euro. *Gene* **577**, 75–81. (doi:10.1016/j.gene.2015.11.018)

21.   McTavish EJ, Decker JE, Schnabel RD, Taylor JF, Hillis DM. 2013 New World cattle show ancestry
from multiple independent domestication events. *Proc. Natl. Acad. Sci. USA* **110**, E1398–E1406.
(doi:10.1073/pnas.1303367110)

22.   van Heerwaarden J, Doebley J, Briggs WH, Glaubitz JC, Goodman MM, de Jesus Sanchez Gonzalez J,
et al. 2011 Genetic signals of origin, spread, and introgression in a large sample of maize landraces.
Proc. Natl. Acad. Sci. USA **108**, 1088–1092. (doi:10.1073/pnas.1013011108)

23.   Larson G, Fuller DQ. 2014 The evolution of animal domestication. *Annu. Rev. Ecol. Evol. Syst.* **45**,
115–136. (doi:10.1146/annurev-ecolsys-110512-135813)

24.   Marshall FB, Dobney K, Denham T, Capriles JM. 2014 Evaluating the roles of directed breeding and
gene flow in animal domestication. *Proc. Natl. Acad. Sci. USA* **111**, 6153–6158.
(doi:10.1073/pnas.1312984110)

25.   Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. 2011 A robust, simple
genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS One* **6**, e19379.
(doi:10.1371/journal.pone.0019379)

26.   Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q, et al. 2014 TASSEL-GBS: a high
capacity genotyping by sequencing analysis pipeline. *PLoS One* **9**, e90346.
(doi:10.1371/journal.pone.0090346)

27.   Lu L, Chen Y, Wang Z, Li X, Chen W, Tao Z, et al. 2015 The goose genome sequence leads to insights
into the evolution of waterfowl and susceptibility to fatty liver. *Genome Biol*. **16**, 89.
(doi:10.1186/s13059-015-0652-y)

28.   Li H, Durbin R. 2009 Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**, 1754–1760. (doi:10.1093/bioinformatics/btp324)

29.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. 2011 The variant call format
and VCFtools. *Bioinformatics* **27**, 2156–2158. (doi:10.1093/bioinformatics/btr330)

30.   Pritchard JK, Stephens M, Donnelly P. 2000 Inference of population structure using multilocus
genotype data. *Genetics* **155**, 945–959.

31.   Patterson N, Price AL, Reich D. 2006 Population structure and eigenanalysis. *PLoS Genet.* **2**, e190.

(doi:10.1371/journal.pgen.0020190)

32. Falush D, Stephens M, Pritchard JK. 2003 Inference of population structure using multilocus
genotype data: linked loci and correlated allele frequencies. *Genetics* **164**, 1567–1587.

33. Chhatre VE, Emerson KJ. 2017 StrAuto: automation and parallelization of STRUCTURE analysis. *BMC*
Bioinformatics **18**, 192. (doi:10.1186/s12859-017-1593-0)

34. Evanno G, Regnaut S, Goudet J. 2005 Detecting the number of clusters of individuals using the
software structure: a simulation study. *Mol. Ecol.* **14**, 2611–2620. (doi:10.1111/j.1365-
294X.2005.02553.x)

35. Earl DA, VonHoldt BM. 2012 STRUCTURE HARVESTER: a website and program for visualizing
STRUCTURE output and implementing the Evanno method. *Conserv. Genet. Resour.* **4**, 359–361.
(doi:10.1007/s12686-011-9548-7)

36. Jakobsson M, Rosenberg NA. 2007 CLUMPP: a cluster matching and permutation program for
dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**,
1801–1806. (doi:10.1093/bioinformatics/btm233)

37. Rosenberg NA. 2003 distruct: a program for the graphical display of population structure. *Mol. Ecol.*
Notes **4**, 137–138. (doi:10.1046/j.1471-8286.2003.00566.x)

38. R Core Team. 2017 *R: a language and environment for statistical computing*. Vienna, Austria: R
Foundation for Statistical Computing.

39. Paradis E, Claude J, Strimmer K. 2004 APE: analyses of phylogenetics and evolution in R language.
Bioinformatics **20**, 289–290. (doi:10.1093/bioinformatics/btg412)

40. Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, et al. 2012 Ancient admixture in
human history. *Genetics* **192**, 1065–1093.

41. Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. 2013 Robust demographic inference
from genomic and SNP data. *PLoS Genet.* **9**, e1003905. (doi:10.1371/journal.pgen.1003905)

42. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. 2009 The sequence alignment/map
format and SAMtools. *Bioinformatics* **25**, 2078–2079. (doi:10.1093/bioinformatics/btp352)

43. Pujolar JM, Dalén L, Olsen RA, Hansen MM, Madsen J. 2018 First de novo whole genome sequencing
and assembly of the pink-footed goose. *Genomics* **110**, 75–79. (doi:10.1016/j.ygeno.2017.08.008)

44. Goudet J. 2005 hierfstat, a package for R to compute and test hierarchical F-statistics. *Mol. Ecol.*
Notes **5**, 184–186. (doi:10.1111/j.1471-8286.2004.00828.x)

45. Welch BL. 1938 The significance of the difference between two means when the population

variances are unequal. *Biometrika* **29**, 350. (doi:10.2307/2332010)

46.   Excoffier L, Lischer HEL. 2010 Arlequin suite ver 3.5: a new series of programs to perform population
genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* **10**, 564–567. (doi:10.1111/j.1755-
0998.2010.02847.x)

47.   Moyers BT, Morrell PL, McKay JK. 2018 Genetic costs of domestication and improvement. *J. Hered.*
109, 103–116. (doi:10.1093/jhered/esx069)

48.   Kanginakudru S, Metta M, Jakati R, Nagaraju J. 2008 Genetic evidence from Indian red jungle fowl
corroborates multiple domestication of modern day chicken. *BMC Evol. Biol.* **8**, 174.
(doi:10.1186/1471-2148-8-174)

49.   Stringham SA, Mulroy EE, Xing J, Record D, Guernsey MW, Aldenhoven JT, et al. 2012 Divergence,
convergence, and the ancestry of feral populations in the domestic rock pigeon. *Curr. Biol.* **22**, 302–
308. (doi:10.1016/j.cub.2011.12.045)

50.   Scott DA, Rose PM. 1996 *Atlas of Anatidae populations in Africa and Western Eurasia*. Wetlands
International Publication No. 41. Wetlands International, Wageningen, The Netherlands.

51.   Rooth J. 1971 The occurrence of the greylag goose *Anser anser* in the western part of its distribution
area. *Ardea* **59**, 17–27.

52.   Zeuner FE. 1963 *A history of domesticated animals*. New York and Evanston: Harper & Row.

53.   Madsen J, Cracknell G, Fox T, editors. 1999 *Goose populations of the Western Palearctic. A review of*
status and distribution. Wetlands International Publication No. 48. Wetlands International,
Wageningen, The Netherlands; National Environmental Research Institute, Rønde, Denmark.

54.   Andersson Å, Follestad A, Nilsson L, Persson H. 2001 Migration patterns of Nordic greylag geese
Anser anser. *Ornis Svecica* **11**, 19–58.

55.   Rohwer FC, Anderson MG. 1988 Female-biased philopatry, monogamy, and the timing of pair
formation in migratory waterfowl. In *Current Ornithology* (ed. RF Johnston), pp. 187-221. Boston,
MA: Springer US. (doi:10.1007/978-1-4615-6787-5_4)

56.   Official Statistics of Finland (OSF): Number of livestock [e-publication]. Helsinki: Natural Resources
Institute Finland [referred: 17.12.2016]. Access method: http://www.stat.fi/til/klm/index_en.html
2016.

57.   Pellegrino I, Cucco M, Follestad A, Boos M. 2015 Lack of genetic structure in greylag goose (*Anser*
anser) populations along the European Atlantic flyway. *PeerJ* **3**, e1161. (doi:10.7717/peerj.1161)

58.   Wang CM, Way TD, Chang YC, Yen NT, Hu CL, Nien PC, et al. 2010 The origin of the White Roman

goose. *Biochem. Genet.* **48**, 938–943. (doi:10.1007/s10528-010-9374-8)

59.   Fox AD, Ebbinge BS, Mitchell C, Heinicke T, Aarvak T, Colhoun K, et al. 2010 Current estimates of
goose population sizes in western Europe, a gap analysis and an assessment of trends. *Ornis Svecica*
20, 115–127.

60.   Zeder MA. 2012 The domestication of animals. *J. Anthropol. Res.* **68**, 161–190.
(doi:10.3998/jar.0521004.0068.201)

61.   BirdLife International and Handbook of the Birds of the World (2016) 2016. *Anser anser*. The IUCN
Red List of Threatened Species. Version 2018-1.

549   Table 1 The parameter values that maximise likelihood estimates (MLE) of the preferred demographic

550   model for goose domestication history (see text) with their 95% confidence intervals (CI) and the search

551   range for the parameter values.

Model	Number of parameters	Parameter	MLE	95% CI	Search range
Divergence with 2 migration matrices	9	ANCSIZE	1112	291.95 - 9030.75	100 - 100000
		T_1	5319	1538.80 - 8225.45	100 - 20000
		$M1_{WD}$	5.35×10^{-4}	2.96×10^{-4} - 6.75×10^{-4}	1×10^{-5} - 20
		$M1_{DW}$	4.25×10^{-4}	1.73×10^{-4} - 6.03×10^{-4}	1×10^{-5} - 20
		T_2	159	79.65 - 462.95	50 - 10000
		$M2_{WD}$	6.69×10^{-4}	4.57×10^{-4} - 9.69×10^{-4}	1×10^{-5} - 20
		$M2_{DW}$	1.72×10^{-3}	1.33×10^{-3} - 2.15×10^{-3}	1×10^{-5} - 20
		N_{DOM}	959	879.95 - 1073.25	50 - 100000
		N_{WILD}	2504	2341.60 - 2695.55	50 - 100000

ANCSIZE, effective population size of ancestral population; T_1, time of divergence in generations; N_{DOM}, effective population size for domestic geese; N_{WILD}, effective population size for greylags; T_2, estimate of time in generations when the migration matrix switched; $M1_{WD}$ migration rate from wild to domestic following T_1; $M1_{DW}$ migration rate from domestic to wild following T_1; $M2_{WD}$ migration rate from wild to domestic following T_2; $M2_{DW}$ migration rate from domestic to wild following T_2.

553 Table 2 Hierarchical analysis of molecular variance (AMOVA) of greylags and their domestic descendants,
554 considering pure populations of greylags (first group) and pure breeds of European domestic geese (second
555 group).

Source of variation	Sum of squares	Variance components	Percentage variation	Fixation indices
Among groups	47565.119	431.4291	15.8	F_{CT} = 0.158[**]
Among populations and breeds within groups	82960.489	302.51404	11.1	F_{SC} = 0.131[**]
Within populations and breeds	345889.821	2003.45893	73.2	F_{ST} = 0.268[**]
Total	476415.429	2737.40207		

[**] $p < 0.001$

Figure 1 Map showing the sampling sites for wild greylags used in this study. The breeding area of the

species is shown on darker grey. The sampling sites in Kazakhstan were combined for analyses (one sample

per location) and the sampling sites in Southern Finland included combined samples from the

geographically close sites of Västanfjärd, Nauvo (shown) and Kemiö (shown). The Iranian samples were

collected during the wintering season. Map modified from IUCN [61].

565   Simple divergence model   Divergence with migration   Divergence with 2 migration matrices

566   Figure 2 Demographic histories of goose domestication as tested with fastsimcoal2.

Figure 3 The genetic divergence and hybridisation patterns in greylag and domestic geese. Population

status and names labelled as in Table S1. The colours in a) and b) are associated to different groups as

follows: greylags (blue), European domestics (green) and Chinese domestics (red). a) The first three

principal components summarising the genetic variation in geese (percentage explained by each PC is

shown). b) STRUCTURE assignment plots for K=2, K=3 and K=4. Each vertical line represents one individual

with K number of colours indicating proportion of ancestry from the inferred clusters. c) Plot relating to the

f_3 (Table S2) values obtained for each population. The more negative the f_3, the more significant is Z-score

in favour of admixture. The f_3 values were not calculated for Landes and the Chinese geese, as they were

used as source populations, thus they were given an f_3 value of 0.