Reaching around obstacles accounts for uncertainty in coordinate transformations

Parisa Abedi Khoozani[1,2], Dimitris Voudouris[4,5], Gunnar Blohm[1,2,3], Katja Fiehler[4,5]

[1]Centre for Neuroscience Studies, Queen's University, Kingston, Ontario, Canada
[2]Canadian Action and Perception Network (CAPnet), Toronto, Ontario, Canada
[3]Association for Canadian Neuroinformatics and Computational Neuroscience (CNCN), Kingston, Ontario, Canada
[4]Center for Mind, Brain, and Behaviour, Marburg University, Marburg, Germany
[5]Psychology and Sport Sciences, Justus Liebig University, Giessen, Germany

Abstract

When reaching to a visual target, humans need to transform the spatial target representation into the coordinate system of their moving arm. It has been shown that increased demands in such coordinate transformations, for instance when the head is rolled toward one shoulder, lead to higher movement variability and influence movement decisions. However, it is unknown whether the brain incorporates such added variability to adjust movements when it is necessary. We designed an obstacle avoidance task in which participants had to reach to a visual target without colliding with an obstacle. We introduced different coordinate transformation demands by varying head roll (straight, 30° clockwise and 30° counterclockwise). In agreement with previous studies, we observed that the reaching variability increased when the head was tilted. In addition, participants systematically changed their obstacle avoidance behavior with head roll. In particular, they changed the preferred direction of passing the obstacle and increased the error margins indicated by stronger movement curvature. Indeed, participants' reaching movements did not differ in the number of collisions between the head roll and the head straight conditions. These results suggest that the brain takes the added movement variability during head roll into account and compensates for it by adjusting the reaching trajectories.

Introduction

Transforming retinal information to the coordinate system of the moving arm is crucial for performing visually guided movements, e.g. reaching (Buneo & Andersen, 2006; Buneo et al., 2002; Cohen & Andersen, 2002; Engel, Flanders, & Soechting, 2002; Knudsen, 2002; Knudsen, du Lac, & Esterly, 1987; Lacquaniti & Caminiti, 2003; Soechting & Flanders, 1992). It has been suggested that coordinate transformations should be considered as stochastic processes that add uncertainty to the transformed signals (Alikhanian, Carvalho, & Blohm, 2015; McGuire & Sabes, 2009; Sober & Sabes, 2003, 2005). Furthermore, it has been shown that stochasticity in coordinate transformations propagates to the movement resulting in increased movement variability (Abedi Khoozani & Blohm, 2018; Burns, Nashed, & Blohm, 2011; Burns & Blohm, 2010; Schlicht & Schrater, 2007); however, it is unknown if the brain accounts for potential consequences of such added movement variability while planning and executing reaching movements.

Accurate coordinate transformations rely on the estimation of three-dimensional (3D) body pose (Blohm & Crawford 2007). This requires an internal model of different body parts with regard to each other, e.g.

eye relative to head translation, and an estimation of joint angles, e.g. head rotation. While internal
models are learned and -most likely- do not change, the estimation of the joint angle can arise from two
sources: 1) afferent sensory signals and 2) efferent copies of motor commands. Both signals are
corrupted with uncertainty in sensory processing and variability of neuronal spiking (Faisal, Selen, &
Wolpert, 2009). Several studies have suggested that varying body pose, e.g. rolling the head, increases
behavioral variability (Abedi Khoozani & Blohm, 2018; Burns, Nashed, & Blohm, 2011; Burns & Blohm,
2010; Schlicht & Schrater, 2007). For instance, Burns and Blohm (2010) showed that rolling the head to
either shoulder results in higher goal-directed reaching variability compared to straight head reaching.
The authors argued that this increased variability stems from the signal-dependent noise in coordinate
transformations. However, another interpretation can be that since humans perform reaching mostly in
an upright posture, the difference in variability can arise from the lack of experience, or less familiarity,
in the rolled condition (Sober & Körding, 2012). To differentiate between these two speculations, we
have previously asked human participants to perform visually guided reaching movements in both rolled
head condition and with a neck load (Abedi Khoozani & Blohm, 2018). Our rationale was that if
familiarity determined variability, then neck load should have no effect; conversely, active inference of
head roll angles should result in larger (resp. smaller) variability due to increased (resp. decreased)
signal-dependent noise due to muscle activity (Cordo et al., 2002; Faisal et al., 2008; Lechner-
Steinleitner, 1978; Sadeghi et al., 2007; Scott & Loeb, 1994) in case of resistive (resp. assistive) load
conditions. Since larger joint angle estimates and muscle activations are accompanied with higher
uncertainty (Blohm & Crawford, 2007; Van Beuzekom & Van Gisbergen, 2000; Wade & Curthoys, 1997),
both head roll and neck load manipulations should result in noisier coordinate transformations. Our
result supported the hypothesis that signal-dependent noise in coordinate transformations increases
movement variability (Abedi Khoozani & Blohm, 2018).However, it is unknown if the brain incorporates
this added movement variability in reaching movements when it compromises task performance.

One approach to investigate whether the brain is accounting for the added movement variability caused
by stochastic coordinate transformations is to perform reaching movements in constrained
environments, i.e. in the presence of obstacles. A failure in accounting for the added variability should
result in behavioral consequences, i.e. obstacle collisions. In general, humans are successful in avoiding
obstacles and they do so by accounting for several factors such as sensory uncertainty (Cohen, Biddle, &
Rosenbaum, 2010), e.g. visual uncertainty, motor noise (Cohen et al., 2010; Hamilton & Wolpert, 2002),
and biomechanical costs (Cohen et al., 2010;   Sabes, 1997; Sabes, Jordan, & Wolpert, 1998; Voudouris,
Smeets, & Brenner, 2012). For instance, Cohen et al. (2010) showed that both higher visual uncertainty
and increased motor noise resulted in increased distance from the obstacle (increased safety margins).
In addition, removing visual feedback results in slower reaction times (Khan, Elliott, Coull, Chua, & Lyons,
2002) and decreased endpoint accuracy (Chua & Elliott, 1993; Heath, 2005; Heath, Westwood, &
Binsted, 2004; Khan et al., 2002) in reaching. These results show that obstacle avoidance is sensitive to
many factors such as sensory accuracy, systemic noise (i.e. motor noise), associated biomechanical
costs, and task requirements, providing an optimal test bed to evaluate if increased noise induced by
stochastic coordinate transformations is considered during reaching.

To investigate whether humans can compensate for the higher movement variability caused by
stochastic coordinate transformations, we designed a reaching task to a visual target while avoiding an
obstacle. To modulate the uncertainty in coordinate transformation, participants performed the
reaching movements with different head orientations (30° toward right shoulder (clockwise; CW), 0°,
and 30° toward the left shoulder (counterclockwise; CCW)). As shown in previous studies, varying the
head roll results in higher movement variability (Abedi Khoozani & Blohm, 2018; Alikhanian et al., 2015;
Burns & Blohm, 2010). Therefore, we expected either higher collision rate for the rolled head compared
to straight head condition, if the added movement variability was ignored, or compensatory behavior
(e.g., increased error margins), if the added movement variability was considered. Furthermore,
previous studies showed that providing visual feedback of the moving hand will alleviate the effect of
noisy coordinate transformations (Blohm & Crawford, 2007). Therefore, we asked participants to
perform reaching movements with or without visual feedback. We expected to observe lower
compensation, if there is any, for the visual feedback condition compared to the no-visual feedback
condition. In agreement with previous studies, we observed that movement variability increased when
the head was tilted. In both feedback conditions, this was accompanied by a change in the preferred
direction of passing the obstacle and increased the error margin, while the collision rate remained
unaffected. We conclude that the brain accounts for the added uncertainty due to coordinate
transformations and compensates for it whenever it compromises task performance.

Materials and methods

Participants

We collected data from 18 healthy humans (10 female) aged between 19 to 38 years (M = 25 years) with
normal or corrected to normal vision. All participants were right handed by self-report, and free of any
known neurological issues. The experiment was approved by the Justus Liebig University Giessen,
general board of ethics and all participants gave their written consents. They received monetary
compensation (8 € / hour) or course credits for their participation.

Apparatus and stimuli

Participants were seated in front of a workspace that comprised a robotic setup with a graspable handle
(vBot; Howard, Ingram, & Wolpert, 2009), a monitor, and a mirror. A helmet with a protruding long stick
and a measuring framework were used to control for the head roll in each condition (Figure 1A). Visual
stimuli were presented on the monitor and reflected in the mirror, which was placed above the robot
(Figure 1B).

A mirror was placed horizontally in front of the participants. This mirror prevented vision of the arm so
that participants were unable to see the movements that they performed below the mirror. The visual
stimuli were presented on a monitor and reflected on the mirror that was placed below it (Figure 1B).
Four disks and the visual instructions were presented in each trial. Two discs (0.5 cm diameter) were
blue, each serving as starting and target position. Both were located in the middle of the screen laterally
(X-position; which was the same as the middle of the body). The starting and the target positions were 9
and 11 cm closer and further from the centre of the screen, respectively, resulting in a distance of 20 cm

between the two. A third disc (1.8 cm diameter), colored red, served as the obstacle. It was presented in
the middle of the screen and in five possible lateral positions: at the center (in line with the starting and
the target position) or shifted by 1.8 cm or 3.6 cm either to the right or left of that central position
(Figure 1C). To simulate a physical obstacle, a repulsive force-field (between 0 to 40N, faster movements
toward the center resulted in higher force) from the center of the obstacle was applied (i.e. vBotDisc).
Consequently, the handle could not move into the obstacle. Finally, a fourth, white disk (0.5 cm
diameter) represented the position of the robotic handle, whenever visual feedback was provided. A
visual instruction that was prompting participants to move to the starting position at the beginning of
the trial was presented at the centre of the screen. Visual stimuli were implemented using Psychtoolbox
(MATLAB 2015) while C++ programming was used for implementing the force-field and programming
the robot.

Figure 1. Overview of the experimental setup. A) Participants performed the reaching task in three head
orientations (30° CCW, 0, and 30° CW). Here only the framework of the robotic setup and the head orientation
setup is displayed, B) The vBot setup consisted of a mirror that displayed the task instructions from the monitor.
Below the mirror, a robot handle was placed that could be freely moved by the participants. C) Participants
brought their hand to the start point and reached to the target while avoiding hitting the obstacle in the center.
Obstacles were randomly shifted to the left or right, along the X-position, across trials. Possible shift positions are
depicted by "×" with the letters indicating the shifts; ML: most leftward, L: leftward, C: central, R: rightward, MR:
most rightward, D) Example trial: first, participants were instructed to bring their hand to the starting position (1),
as soon as they arrived at the starting position (2), the target position and the obstacle (here central) appeared and
participants were instructed to move to the target position in less than 1000 ms (3).

Procedure

In order to reach to the viewed target, participants grasped the robot's handle with their right hand
while resting their forehead on the workspace's framework in front of the screen. They performed the
task in three possible head orientations: 30° CCW, 0°, 30° CW. Each head orientation condition was
performed with and without visual feedback of their reaching hand. The visual feedback was provided
by a moving white disc (0.5 cm diameter) on the screen representing the robot handle position. This
resulted in six combinations, each of which was presented in separate blocks of trials. At the beginning

of each block, the experimenter positioned the participant's head in the respective orientation. Before the start of each trial, participants grasped the robot handle and brought it to a fixed starting position using visual feedback of the robot's handle. After positioning the hand on the starting position, the target and an obstacle were simultaneously displayed on the mirror. Participants were instructed to immediately start reaching towards the visual target while avoiding the obstacle. If participants were able to reach the target without hitting the obstacle in less than 1000 ms from the moment of target presentation, the visual target would turn green indicating that the trial was successful. Otherwise the target would turn red indicating that the trial was aborted and would be repeated later. At the end of each trial all visual stimuli disappeared and the next trial started with the appearance of the starting position.

Before starting the experiment, participants performed a short practice block with 20 trials. Within each of the six blocks, each obstacle position was presented 48 times, resulting in a total of 240 trials per block, and thus in a total of 1440 trials for the complete experiment, which lasted roughly 60 minutes. The combination of the head angle and visual feedback for each block was chosen based on Latin squares method to counterbalance among all conditions (Jacobson & Matthews, 1998).

Data analysis

All offline analyses were performed using MATLAB 2018. For each trial, trajectories were normalized using functional data analysis (Ramsay & Silverman, 2005): B-splines were fit to each dimension of the raw data (x and y) and over-sampled to 2000 equally-spaced time points. A central differential algorithm was used to calculate hand velocity and acceleration. Before each derivation, a low-pass filter (autoregressive forward-backward filter, cutoff frequency = 50 Hz) was used to smooth the data. We calculated the movement onset by finding the moment of 25% and 75% of the trial's peak velocity and then extrapolating a line between these two moments until this line crossed that trial's baseline velocity as this was measured by averaging the velocity during the first 200 ms of the trial (for futher details see Brenner & Smeets, 2018). The reaction time (RT) was calculated by subtracting the time of movement onset from the time that the target appeared. The trial was ended as soon as the participants' hand distance from the center of the target was less than 0.5 cm when hand visual feedback was provided. When visual feedback of the hand was absent, the end of the trial was defined as the moment when the participants' hand reached the position in depth of the target and was less than 4 cm away from its lateral position. The movement duration (MD) was calculated as the time difference between the end of the trial and movement onset. Trials with RT < 100 ms (predictive movements) and RT+MD > 1000 ms (too slow) were removed from the analysis (1.7%) and considered as invalid trials.

To assess the effect of varying head orientation on movement behavior, we calculated movement variability across the whole reaching trajectory. To do so, the normalized trajectories of each participant were averaged separately for each obstacle position and movement direction (rightward or leftward of that obstacle). Since movements were predominantly along the Y-position direction, we only calculated the standard deviation of the handle's lateral position across trials for each of the 2000 normalized steps. Then, we calculated the boundaries of the averaged trajectories by adding/subtracting the standard deviation to/from the mean of the trajectory along the X-position (Figure 2A). Finally, we

calculated the movement variability as the area between the trajectory boundaries. We performed
these steps separately for each participant, head orientation, visual condition, and movement direction
for each obstacle position. We considered the calculated movement variability for different directions of
each obstacle valid only if we had sufficient data. That is if the number of movements in a certain
direction for an obstacle was less than 10% of the overall movements for that obstacle, we considered
the movement variability invalid.

In the next step, we assessed if the added movement variability due to rolling the head had a tangible
effect on the movement strategies. To do so, we considered two parameters: the direction of passing
the obstacle (i.e. around the right side or the left side of the obstacle) and the distance while passing the
obstacle. For movement direction, we calculated the percentage of rightward movements and expected
to see higher rightward movements for the obstacle shifted to the left and the reverse for the obstacle
shifted to the right and similar percentage of rightward and leftward movements for the central
obstacle. We hypothesized that rolling the head should modulate the preferred movement direction;
most noticeably for the central obstacle. With regards to distance from the obstacle, we hypothesized
that reaching trajectories should deviate further away from the obstacle in the head roll conditions in
order to compensate for the added movement variability. This should be reflected in a larger curvature
which is more noticeable for the central obstacle (β: expansion biases; Figure 2C). Based on our earlier
findings and due to under- or over-compensation for head roll (Abedi Khoozani & Blohm, 2018), we
further expected that movement trajectories for straight head conditions should fall symmetrically
between the trajectories for the two head roll conditions (α: rotational biases; Figure 2B). In our data,
both expansion and rotational biases are combined, however, each will affect the trajectory differently.
That is, rotational biases are expected to be symmetrical around the straight head condition, while
expansion biases should cause the deviation curvature in the same direction, regardless of head roll
direction. To separate rotational and expansion biases from each other we employed the following
method. First, we assumed that both expansion and rotational biases are multiplicative; therefore, the
shift caused by each of them is a multiplication of the maximum curvature of the normalized trajectory
for the head straight condition with the same direction and visual condition as the rest of the considered
trajectories in the calculation (mc_{H0}). Therefore, using the following assumptions $mc_{HR/L} = \beta *$
$|mc_{H0}|$ and $mc_{HR/L} \propto \alpha * \widetilde{HR}$ (where $\widetilde{HR}$ is the estimated head angle for the head roll conditions),
we first calculated the shifts in trajectories (the difference of the averaged trajectories between straight
head and 30° CW/CCW head orientations) for each head orientation caused by expansion (Δx_{β}) and
rotation (Δx_{α}) as following:

$$\Delta x_{\beta,R} = \frac{\Delta x_{R,HL} + \Delta x_{R,HR}}{2} \ and \ \Delta x_{\alpha,R} \propto \frac{\Delta x_{R,HL} - \Delta x_{R,HR}}{2}$$

, where the first subscripts indicate the movement direction (R: rightward and L: leftward) and the
second ones indicate the head roll direction (HL: 30°CCW and HR: 30°CW). The visualization of the
variables for the rightward movement direction is provided in Figure 2D.

For expansion biases, we calculated the percentage of expansion as following:

$\quad \beta_R = 100 * (\frac{\Delta x_{\beta,R}}{mc_{R,H0}})$

, therefore, positive values indicate expansion while negative values indicate shrinkage.

For rotational biases, we calculated the angular value as following:

$$\alpha_R = \alpha_{R,H0} - \alpha_{R,HR} = \tan^{-1}(\frac{Y_{mc_{R,H0}}}{mc_{H0}}) - \tan^{-1}(\frac{Y_{mc_{R,HR}}}{mc_{H0} + \Delta x_{\alpha,R}})$$

Here, for simplification, we assumed that the rotational effect on horizontal values is small and
therefore $Y_{mc_{R,HR}} \approx Y_{mc_{R,H0}}$. Similar calculations can be applied for the leftward movements.

Figure 2. Variability, rotational, and expansion biases calculations. A) The overall movement variability was
calculated as the area between the movement boundaries. The boundaries were calculated for each participant
separately by adding/subtracting the standard deviation of the lateral movements from the average for all 2000
samples along the trajectory, B) rotational biases (α): we hypothesized that rolling the head creates rotational
biases resulting in symmetrical shifts of the trajectories for the 30° CW/CCW head orientations (colored solid lines)
around the straight head (black solid line), C) expansion biases (β): varying head orientation increases movement
variability resulting in hand trajectories moving away from the obstacle, D) combination of rotational biases and
expansion biases result in an asymmetry in the shifted trajectories for the 30° CW/CCW head orientations (colored
solid lines) compared to the straight head (black solid line).

Finally, to assess if participants were able to compensate for the effect of varying head orientation, we
calculated the movement speed and the collision rate via dividing the number of collisions with the
obstacle by the total number of valid trials for each head roll and visual feedback condition.

The data and analysis code of this manuscript is available at OSF (https://osf.io/tf8p5/) and GitHub
(https://github.com/Parisaabedi/Obstacle-Avoidance).

Statistical analysis

We used JASP (https://jasp-stats.org/) to perform the statistical analyses. To examine the effect of head
orientation (0 and 30° C/CW), obstacle position (most leftward, leftward, central, rightward, and most
rightward), and visual feedback (with and without) on the above-mentioned dependent variables

(movement variability, movement direction, rotational and expansion biases, movement speed, and
collision rate), we deployed repeated measures ANOVA. Significant differences between the conditions
were further investigated with two-sample paired t-tests and the reported p-values were Bonferroni-
Holm corrected. All the saved file of the performed statistical analysis is available at GitHub
(https://github.com/Parisaabedi/Obstacle-Avoidance).

Results

The objective of this study was to investigate whether humans compensate for movement variability
caused by increased demands in coordinate transformations. To this aim, we asked participants to reach
to a visual target without colliding with an obstacle while having different head orientations (30°
CW/CCW and 0°). We hypothesized that participants should increase their safety margins, i.e. stronger
trajectory curvature, in the 30° CW/CCW head orientations compared to the straight head to
compensate for the increased uncertainty. In addition, participants could also decrease their movement
velocity to compensate for higher uncertainty.

In the first step, we demonstrate that participants indeed adapted their movement strategies to
compensate for the higher variability. Figure 3 illustrates the trajectories of two example participants
(#2 and #16). Both participants were able to successfully avoid the obstacle in all head orientations (the
collision rate increase was less than 2%), however, each of the two participants showed a different
movement behavior in the rolled head orientations (green and blue) compared to the straight head
(black). Specifically, participant #2 moved further away from the obstacle to successfully reach to the
target (increased error margin). In contrary, participant #16 kept the same distance from the obstacle,
but instead decreased the peak velocity. Based on these results, one can speculate that humans might
compensate for the added variability due to stochastic coordinate transformations using different
strategies.

Figure 3. Obstacle avoidance strategies of two participants. The data is shown for the central obstacle position (red circle) without visual feedback. The left panel illustrates the trajectory data as well as the velocity data for participant #2. This participant moved further away from the obstacle, specifically for the rightward movements, in the head roll conditions (green and blue solid lines) compared to straight head condition (solid black line). The two panels on the sides illustrate a zoomed version of the trajectories. The peak velocity did not change for the rightward movements while increased for the leftward movements. The right panel illustrates the behavior of participant #16. In contrary to participant #2, this participant decreased the peak velocity and also decreased the distance from the obstacle, especially for leftward movements.

Rolling the head increased movement variability

Previous studies demonstrate that rolling the head increases movement variability (Burns & Blohm, 2010; Abedi Khoozani & Blohm, 2018). Therefore, in the first step we investigated the effect of varying head orientation on movement variability depending on the visual feedback of the hand. To remind the reader, we calculated the movement variability as the surrounding area between the lateral deviations from the averaged trajectory. Since we didn't expect to observe any effect of obstacle position on the movement variability, we performed a 3 (head orientation) × 2 (visual feedback) repeated measures ANOVA. We observed a main effect of head roll (F(2,34)=4.39, p = 0.020, η^2=0.205), a main effect of visual feedback (F(1,17)=31.36, p < 0.001, η^2=0.648) and no interaction between head roll and visual

feedback (F(2,34)=14.08, p = 0.403). Post-hoc t-test for the head orientation effect revealed a significant
increase of movement variability for the CW head orientation compared to straight head (t(17) = 2.729,
p = 0.043, Cohen's d = - 0.643), but the difference between CCW vs straight head was slightly more
variable (t(17) = 2.171, p = 0.089). Unsurprisingly, there was no difference between CCW vs CW (t(17)=
1.040, p = 0.313). Thus, we confirmed previous work that rolling the head increased the movement
variability. In the next step we assessed if the participants adapted their movement behavior to
compensate for the increased movement variability.

Figure 4. Effect of varying head orientation on movement variability. A) Visual feedback condition:
varying the head orientation did not affect the movement variability. B) Without visual feedback
condition: participants showed different effects of varying head orientation on their movement
variability. Some participants showed increased movement variability while others showed decreased
movement variability. Error bars are standard deviations.

Participants adapted their obstacle avoidance behavior for head roll conditions

To further explore the effect of head roll on movement strategies, we determined the following
parameters: Directional changes (preferred direction to pass the obstacle), rotational and expansion
biases, and collision rate. We found that all participants adapted their movement direction and
expansion biases in the 30° CW/CCW head orientations to successfully reach to the target with similar
collision rates as in the straight head orientation.

Directional changes

Figure 5 depicts the percentage of rightward movements for different head orientations, obstacle
positions, and visual feedback conditions. Varying the head roll changed the preference in passing the
obstacle from a certain side (left vs right). Rolling the head CCW led to a tendency to pass the obstacle
from the right side, while rolling the head CW changed the tendency to pass the obstacle from the left
side. Unsurprisingly, shifting the obstacle to the right or left of the central position changed the
preferred direction of the movement. For example, when the centrally placed obstacle was shifted to
the right, participants preferred to pass it from its left side and vice versa. Lastly, having visual feedback
of the movement did not seem to influence the passing side. The 3 (head orientation) × 5 (obstacle
position) × 2 (visual feedback) repeated measures ANOVA revealed a main effect of head orientation
($F_{(2,34)} = 12.564$, $p < 0.001$, $\eta^2 = 0.43$), a main effect of obstacle position ($F_{(4,68)} = 290.279$, $p < 0.001$,
$\eta^2 = 0.95$), and an interaction between head orientation and obstacle position ($F_{(8,136)} = 405.711$, $p <$
0.001, $\eta^2 = 0.29$). As can been seen in Figure 5 and revealed from the statistical analysis, there is no
difference between the two obstacles configurations on the left (most leftward and leftward) or on the
right of the central obstacle (most rightward and rightward). As there was no effect of visual feedback (p
$= 0.963$) and no interaction between visual feedback and any other conditions (all p's > 0.2), we
collapsed the percentage of rightward movements across the visual feedback conditions as well as
across the two leftward and the two rightward obstacle configurations (Figure 3C). The repeated
measure ANOVA for the collapsed data for the central obstacle revealed a main effect of head
orientation ($F_{(2,68)} = 15.91$, $p < 0.001$, $\eta^2 = 0.48$). Post-hoc t-test showed a significant difference
between the straight head and the CW head orientation ($t(17) = 3.076$, $p = 0.021$, Cohen's $d = 0.73$),
between the straight head and the CCW head orientation ($t(17) = 2.589$, $p = 0.019$, Cohen's $d = 0.61$),
and between the CW and the CWW head orientation ($t(17) = 5.703$, $p < 0.001$, Cohen's $d = 1.344$). These
results demonstrate that participants opted for more rightward and leftward crosses compared to the
straight head when rolling the head CCW and CW, respectively.

Figure 5. Percentage of rightward movements. Head roll caused changes in the preferred direction of
movement A) with and B) without visual feedback. In both conditions rolling the head CCW increased
the tendency in passing the obstacle from the right side. Obstacle positions: ML: most leftward, L:
leftward, C: central, R: rightward, MR: most rightward. Open circles representing single participants. C)
Since there was no difference between the leftward obstacle shifts (ML and L) and between rightward
obstacle shifts (R and ML), we collapsed the leftward and rightward shifts. The star indicates statistical
significance with $p < 0.05$.

Rotational and Expansion biases

We showed that increasing the coordinate transformation demands affects the obstacle avoidance
strategy as reflected in the preferred direction to cross an obstacle. In the next step, our goal was to
examine the effect of increased movement variability on the error margins.

To do so, we first needed to select the trajectories for which we had enough data. As shown in Figure 5,
varying obstacle position modulated the percentage of rightward movements independent of head
orientations. For instance, for most leftward obstacles more than 95% of the movements passed the
obstacle from the right side. Therefore, only very few leftward trajectories occurred in such cases which
is not sufficient for further analysis. As a result, from now on, we only considered the reasonable
preferred direction, i.e. direction with more than 10% of the overall movements, for non-central
obstacle positions, i.e. rightward movements for obstacles shifted to the left and vice versa.

In addition to changing the preferred direction, increasing the safety margin by increasing the trajectory
curvature could also compensate for the increased variability caused by head roll. However, we expect
to observe the increased curvature only when it is necessary, i.e. when the distance between the
moving hand and the obstacle is not high enough to avoid hitting the obstacle. For instance, as
mentioned earlier, in the absence of visual feedback, participant #2 only increased the curvature in the
head rolled conditions when passing the obstacle for rightward but not for leftward movements (Figure
3, left panel). One possible explanation could be that removing visual feedback caused a higher safety
margin for leftward movements in the straight head condition (difference between solid and dotted line
in figure 6F-H) which resulted in no need for further requirement to increase the curvature in the head
roll conditions.

To investigate the effect of head roll on movement trajectories for all participants, we plotted the
pooled trajectories for each obstacle for both with and without visual feedback conditions (Figure 6A-E
and Figure 6F-J respectively). Since the effects were more prominent when visual feedback was
withdrawn, we will focus on this condition. As illustrated in Figure 6F-J, rolling the head created both
rotational (blue and green trajectories are shifted in opposite side of the black) and expansion (the
trajectory shifts are not symmetrical; i.e. green trajectory being close to black one for Figure 6I-J) biases
when visual feedback of the hand was removed. The effect of head roll was more noticeable for
rightward (Figure 6I-J) compared to leftward movements (Figure 6F-G). However, one needs to consider
that the shift caused by removing the visual feedback for leftward movements while the head was
straight was already stronger than for the rightward movements; the difference between the black solid
line (without visual feedback) and the black dotted line (with visual feedback). This observation is in line
with previous studies (Chapman & Goodale, 2008; De Haan et al., 2014; Menger, Dijkerman, et al., 2013;
Menger et al., 2012; Ross et al., 2018; Ross et al., 2015), and as Menger et al. (2013) illustrated mainly
due to the degree of obstructiveness of the obstacle. This indicates that people adapt their
compensatory behavior if it is necessary. In other words, if the safety margins for the leftward
movements were sufficiently large, there is no further need to increase the margins in the presence of
higher uncertainty.

Figure 6. Participants showed both rotational and expansion biases caused by varying head orientation. A-E: Movement trajectories for different head orientations and obstacle positions when visual feedback is provided. Trajectories are averaged across participants. The shaded areas indicate the standard errors of the mean. The obstacle is depicted as the red circle. A: most rightward obstacle, B: rightward obstacle, C: central obstacle, D: leftward obstacle, E: most leftward obstacle. Providing visual feedback of the hand abolished the rotational biases but not the expansion biases. F-J: Movement trajectories for different head orientations and obstacle positions when visual feedback is removed. Obstacle positions are equal to the ones in A-E. The shift due to removing the visual feedback

(difference between the black dotted line and the solid line) is most prominent for leftward movements
([F] and [G]) compared to rightward movements ([I] and [J]). Rightward movements showed both
rotational and expansion biases while leftward movements are mainly overlapping.

To quantify the effect of head orientation on safety margins, we first separated the rotational biases
(due to misestimation of the head angle) from the expansion biases (due to uncertainty in head angle
estimation). For details of this calculation please see "Materials and Methods" section. We performed
the calculations for each individual participant and each obstacle, separately for each visual condition. In
the next step, we combined the three leftward movements (for MR, R, and C obstacle positions) and
similarly the three rightward movements (for ML, L, and C obstacle positions). Figure 7 demonstrates
the changes in rotational (Figure 7A) and expansion biases (Figure 7B) for different movement
directions.

Regarding the rotational biases, similar to previous studies (Burns & Blohm, 2010; Abedi Khoozani &
Blohm, 2018), we considered a positive value of the rotational bias as an indicator of overestimated
head angle and vice versa for negative rotational biases. In the presence of visual feedback, the
rotational biases were not significantly higher than zero for both rightward movements (t(17) = 0.926, p
= 0.367, Cohen's d = 0.22) and leftward movements (t(17) = 2.115, p = 0.050, Cohen's d = 0.50). On the
other hand, when visual feedback was removed, we observed an overestimation of head angle for the
rightward movements (t(17) = 2.521, p = 0.022, Cohen's d = 0.59) and no statistically significant
rotational biases for leftward movements (t(17) = 0.646, p = 0.527, Cohen's d = 0.15).

Figure 7. Quantification of rotational and expansion biases caused by rolling the head. A) rotational
biases caused by misestimation of the head angle, B) expansion biases caused by increased uncertainty
due to head roll. . Error bars are standard errors of the mean. Stars represent statistical significance, p <
0.05.

Regarding the expansion bias, a positive value indicates that participants increased the movement
curvature to create a higher safety margin. Consequently, a zero expansion bias means no change in
movement curvature for the CW/CCW head orientations compared to the straight head orientation and
a negative expansion bias indicates a smaller distance between movement trajectory and the obstacle
for the CW/CCW head orientations than the straight head orientation. Additionally, for better

visualization we used the expansion/shrinkage percentage; i.e. +20% is equivalent to $\beta = 1.2$ and -20%

is equivalent to $\beta = 0.8$. As can be seen in Figure7B, participants significantly increased their movement

curvature while rolling their head compared to the straight head for rightward movements both with

visual feedback (t(17) = 3.082, p = 0.007, Cohen's d = 0.73) and without visual feedback (t(17) = 4.142, p

< 0.001, Cohen's d = 0.98). For leftward movements, the movement curvature was increased when

visual feedback was provided (t(17) = 3.388, p = 0.003, Cohen's d = 0.80). However, when visual

feedback was withdrawn, curvature of leftward movements did not differ significantly for different head

orientations (t(17) = 1.537, p = 0.143, Cohen's d = 0.36), possibly because such movements had already

a large safety margin from the obstacle (see Figure 6F-H). In contrary, for rightward movements, having

no visual feedback did not cause higher movement curvature; therefore, it was necessary to increase

movement curvature for the head roll conditions to increase the safety margin to successfully avoid the

obstacle.

Movement speed

As mentioned before, increasing movement speed is another compensation strategy to counteract

increased movement variability due to rolling the head. However, we did not find any changes in

movement speed for any of the experimental conditions (all p > 0.1).

Adapted behavior resulted in the same collision rate for different head orientations

As demonstrated in the previous sections, participants adapted their behavior (movement directions

and safety margins) for different head orientations. It is logical to assume that the purpose of this

adapted behavior was to compensate for the increased movement variability due to head roll. We did

not expect to observe any difference in the collision rate for the different obstacle positions, therefore,

we pooled the data across the obstacle positions and assessed if head orientation or visual feedback

affected the collision rate. The 3 (head orientation) x 2 (visual feedback) repeated measures ANOVA

revealed no main effect of head orientation (F(2,34) = 0.100, p = 0.905, η^2=0.006), a main effect of visual

feedback (F(1,17) = 12.831, p = 0.002, η^2=0.430), and no interaction between the two (F(2,34) = 1.044, p

= 0.363, η^2=0.058). As illustrated in Figure 8, removing visual feedback caused an increase in the collision

rate. However, in both visual conditions, the collision rate remained the same for different head

orientations indicating that participants were able to successfully compensate for the added variability

due to varying head orientations.

Figure 8. Effect of head orientation on collision rate. A) Visual feedback condition: participants were
able to successfully perform the reaching task showing a low collision rate for different head
orientations. B) No visual feedback condition: removing the visual feedback increased the overall
collision irrespective of the head orientation. . Error bars are standard deviations.

Discussion

The goal of the current study was to assess whether and how humans account for the added movement uncertainty induced by stochastic coordinate transformations in goal-directed movements. To this aim, we asked human participants to reach to visual targets while avoiding obstacles with different head orientations (straight and 30° CW/CCW) and visual feedback configurations (with/without visual feedback). We hypothesized that if humans are compensating for the increased uncertainty caused by stochastic coordinate transformations, varying head orientation should not affect their performance (i.e. same collision rate for all head orientations). If that was true, we hypothesized to observe compensatory effects in the trajectories, such as increased error margins (increased curvature), for the rolled compared to the straight head conditions. As expected, rolling the head increased movement variability. To accommodate this increased variability, participants adapted their movement behavior by varying their preferred movement direction and increasing their safety margins from the obstacle (based on collision likelihood). Consequently, the collision rate remained the same for all head orientations. Thus, humans considered the increased movement variability resulting from stochastic coordinate transformations in their goal-directed movements.

The main assumption of the current study is that the stochasticity of coordinate transformations propagates to the final motor output. This assumption is based on numerous studies demonstrating that uncertainty in coordinate transformations causes higher variability in movement execution (Abedi Khoozani & Blohm, 2018; Biguer, Prablanc, & Jeannerod, 1984; Bock, 1986, 1993; Burns & Blohm, 2010; Henriques el al., 1998; Henriques & Crawford, 2000; Lewald & Ehrenstein, 2002; McGuire & Sabes, 2009; Schlicht & Schrater, 2007; Schütz, Henriques, & Fiehler, 2013; Sober & Sabes, 2003, 2005; Vaziri, Diedrichsen, & Shadmehr, 2006). For instance, when reaching to visual targets of different eccentricities with respect to gaze fixation, reaching movements overshoot the target in the absence of visual feedback (Bock, 1986, 1993; Henriques et al., 1998; Henriques & Crawford, 2000; Lewald & Ehrenstein, 2002; Vaziri et al., 2006). It has been suggested that these overshoots likely arise from noise in transforming the visual estimate of the target to the proprioceptive estimate of the hand (Dessing et al., 2012). Furthermore, McGuire and Sabes (2009) showed that the gaze-dependent errors vary based on the target's effector (visual or proprioceptive or both) as well as the available information of the initial hand position (with or without visual feedback). They showed that the gaze-dependent reaching errors are only observable for visual targets and are abolished for proprioceptive targets, suggesting that the transformation of a visual target to the coordinate frame of the arm systematically affects reaching movements. Based on the evidence that accurate coordinate transformations rely on the estimation of body geometry (Blohm & Crawford, 2007) and to elaborate more on the effect of stochastic coordinate transformations on reaching movements, previous studies varied the reliability of the head angle estimations via rolling the head and/or loading the neck (Abedi Khoozani & Blohm, 2018; Burns & Blohm, 2010). Both factors biased reaching movements and increased the movement variability compared to a control condition (e.g. straight head and no neck load). All together, we, therefore, argue that there is a clear propagation of uncertainty caused by stochastic coordinate transformations to the performed reaching movements.

If stochastic coordinate transformations cause higher movement variability, does the brain account for
such noise when planning and executing goal-directed movements? In the following, we argue that it is
rather unlikely that the brain dismisses such nuisances.

We observed two main strategies to compensate for the increased movement variability caused by
stochastic coordinate transformations: (1) changes in the preferred direction of passing the obstacle,
and (2) increased safety margins. With regards to strategy (1), we believe that it is caused by signal-
dependent noise. Since for the rightward/leftward obstacles one direction is distinctly dominant (e.g.
rightward direction for the obstacle shifted to the most leftward positions), we only focus on the central
obstacle in which the likelihood of passing the obstacle from the right- or left side was (almost) at
chance level. To elaborate more on why changing the preferred direction will facilitate the effect of
coordinate transformations, we exemplarily consider the 30° CW head orientation. In this configuration,
participants preferred to pass the obstacle from the left side (Figure 5). It has been shown that humans
move their gaze to specific task related landmarks (e.g. possible contact point with obstacle) during
reaching movements to gain spatial information for movement control (Johansson et al., 2001).
Consequently, for 30° CW head orientation, a rightward eye rotation is required in order to have a more
accurate view of the right side of the screen (with regard to the body and screen midpoint) while such
rotation is not required for left side of the screen. Meanwhile the accurate coordinate transformation
relies on the estimation of body geometry (Blohm & Crawford, 2007), here both head and eye angles.
Additionally, it has been established that the noise associated with this estimation is signal-dependent
(Abedi Khoozani & Blohm, 2018; Burns & Blohm, 2010; Schlicht & Schrater, 2007); that is the higher the
amplitude of the signal, the noisier the estimation. Therefore, the extra rotations-translation of the eye,
required for the right side of the screen, may result in noisier eye-in-head orientation estimations and,
consequently, noisier coordinate transformations. Accordingly, to decrease the uncertainty associated
with the coordinate transformation it is logical to pass the obstacle on the left side, which is also in
accordance with our data (see Figure 5C). Hence, we argue that a likely explanation is that participants
systematically adapted their preferred movement direction to decrease the uncertainty accompanied by
required coordinate transformations and ultimately the likelihood of hitting the obstacle.

With regards to strategy (2), we observed increased safety margins (i.e. expansion biases) when the
head was tilted except for movements performed on the left side of the obstacles in the absence of
visual feedback. The asymmetry in increasing safety margins between rightward and leftward
movements was unexpected. To explain this pattern of results it is noteworthy to emphasize that the
increase in trajectory curvature due to removing the visual feedback is much less for rightward
compared to leftward passes (the comparison between dotted line and solid line in Figure 6B), which is
also in accordance with previous findings (Chapman & Goodale, 2008; De Haan et al., 2014; Menger et
al., 2013; Menger et al., 2012; Ross et al., 2018; Ross et al., 2015). Menger et al. (2013) demonstrated
that the observed asymmetry cannot be explained by motor lateralization and argued that the observed
behavior is most likely due to the degree of intrusiveness of an obstacle for the whole arm (Menger et
al., 2013; Menger et al., 2012). In other words, when humans are passing an obstacle, they accurately
calculate the likelihood of hitting the obstacle with their whole moving arm (Voudouris et al., 2012).
Furthermore, the expansion biases persist even when the visual feedback of the hand was available

during the movement. While it has been argued that providing visual feedback of the hand will remove
the biases caused by gaze-shifts and more generally by coordinate transformations (Brown, Marlin, &
Morrow, 2015; Dessing et al., 2012; Saunders, 2004; Saunders & Knill, 2003), we argue that the signal-
dependent noise still persists in the system. In other words, while the extra source of information (i.e.
visual information of the hand position) decreases the amount of uncertainty, it is not fully abolished.
Similarly, Ross et al. (2015) observed that varying the fixation while the visual feedback was available
caused participants to veer away from the fixated obstacle as opposed to free viewing or central
fixation. The authors speculated that the observed pattern can be explained by the misestimation of the
target position on the retina, however, we argue that the observed veering away from the fixated
obstacle might be better explained by stochastic coordinate transformations: Given that varying the
gaze position will result in higher uncertainty in eye-in-head orientation estimation and consequently in
noisier movements, it is logical to increase the safety margin to decrease the likelihood of obstacle
collision.

As mentioned above, in our study participants increased their safety margin for tilted head conditions if
the initial safety margin (straight head condition curvature) could not account for the added uncertainty
due to stochastic coordinate transformations. More specifically, since the curvature for the straight
head condition is already high enough for leftward movements without visual feedback, we did not
observe any expansion biases. This can be explained by including the movement signal-dependent noise
in the calculation of collision likelihood and motor planning (Hamilton & Wolpert, 2002; Harris &
Wolpert, 1998). Harris and Wolpert (1998) proposed a theoretical framework, called task optimization in
the presence of signal-dependent noise (TOPS), and showed that including signal-dependent noise
provides a general framework which can explain both saccadic and point-to-point arm movements. In a
later study, Hamilton and Wolpert (2002) extended this framework and showed that TOPS can also
predict the trajectories generated in an obstacle avoidance task. Based on these observations, they
proposed that controlling the statistics of the movement (such as minimizing the endpoint error) while
accounting for signal-dependent noise might offer a unifying principle for goal-directed movements.
Therefore, it is also crucial to have a better understanding of the nature of such signal-dependent noise
corrupting the movements. While previous studies (Hamilton & Wolpert, 2002; Harris & Wolpert, 1998;
Van Beers, Baraduc, & Wolpert, 2002) mainly associated the signal-dependent noise with the amplitude
of the motor command, we argue that the processing, e.g. coordinate transformations, required for
generating the motor command can also cause movement variability and therefore, it is important to
account for such noise in the motor circuitry.

While the role of coordinate transformations in the motor planning stage is demonstrated through
many studies (Abedi Khoozani & Blohm, 2018; Burns & Blohm, 2010; McGuire & Sabes, 2009; Sober &
Sabes, 2003, 2005), surprisingly, its impact is fully ignored in the optimal motor control field. This study
provided the first evidence that the brain accommodates for the added movement variability due to
uncertainty in coordinate transformations. However, it is not clear at what stage during motor planning
and execution such accommodations might occur. Based on the optimal control theory, the motor
system selects the appropriate control law to calculate the motor command based on the desired task
goal (e.g. grab a pen) and the current system states (i.e. limb position). Since both motor commands and

sensory signals of motor performance are corrupted with noise, the optimal state estimator uses both
sensory signals (feedback circuitry) and an efference copy of the motor commands (feedforward
circuitry) to estimate the current state of the limb (Scott & Norman, 2003; Scott, 2004; Todorov, 2004;
Todorov & Jordan, 2002). Within this framework, however, it is unknown in which coordinates each of
these processes (feedback and feedforward) can be carried out. It has been shown that it is beneficial to
plan movements in multiple coordinate frames (McGuire & Sabes, 2009). In addition, the feedback of
the movement can be presented in different coordinate frames, e.g. the visual feedback of the hand in
retinal coordinates and the proprioceptive feedback of the hand in body coordinates. Thus, it is not
trivial which coordinate system should be used for implementing optimal feedback control. For instance,
all signals could be transformed and then combined in one coordinate frame (e.g. visual, proprioceptive,
or an intermediate frame). Alternatively, all signals could be transformed into the other signal's
coordinate frame (i.e. visual to proprioceptive and vice versa) and the error signal is generated in all
coordinate frames in parallel (similar to generating movement plans in multiple coordinate systems).
Further modeling and experimental studies are required to investigate the role of coordinate
transformations in the optimal motor control framework. Such studies have implications not only in the
motor control field, but also in perception, decision making as well as applicable fields such as brain
machine interfaces and robotics.

All in all, we believe that uncertainty in coordinate transformations results from signal-dependent noise
and propagates to motor behavior. Additionally, the brain accounts for such noise during motor
planning, and possibly execution, and adapts the behavior whenever such noise can compromise
performance.

Acknowledgement

We would like to thank Marie Mosebach for her help in data collection. This project was funded by
NSERC (Canada), DFG IRTG 1901 "The brain in action" (Germany), and DAAD (Germany).

References

Abedi Khoozani, P., & Blohm, G. (2018). Neck muscle spindle noise biases reaches in a multisensory
integration task. *Journal of Neurophysiology*, *120*(3), 893–909.
https://doi.org/10.1152/jn.00643.2017

Alikhanian, H., Carvalho, S. R., & Blohm, G. (2015). Quantifying effects of stochasticity in reference frame
transformations on posterior distributions. *Frontiers in Computational Neuroscience*, *9*, 82.
https://doi.org/10.3389/fncom.2015.00082

Battaglia, P. W., & Schrater, P. R. (2007). Humans trade off viewing time and movement duration to
improve visuomotor accuracy in a fast reaching task. *Journal of Neuroscience*, *27*(26), 6984–6994.
https://doi.org/10.1523/JNEUROSCI.1309-07.2007

Biguer, B., Prablanc, C., & Jeannerod, M. (1984). The contribution of coordinated eye and head
movements in hand pointing accuracy. *Experimental Brain Research*, *55*, 462–469.

Blohm, G., & Crawford, J. D. (2007). Computations for geometrically accurate visually guided reaching in
3-D space. *Journal of Vision*, *7*(5), 4. https://doi.org/10.1167/7.5.4

Bock, O. (1986). Contribution of retinal versus extraretinal signals towards visual localization in goal-
directed movements. *Experimental Brain Research*, *64*, 476–482.

Bock, O. (1993). Localization of objects in the peripheral visual field. *Behavioural Brain Research*, *56*(1),
77–84. https://doi.org/10.1016/0166-4328(93)90023-J

Brenner, E., & Smeets, J. B. J. (2018). How can you best measure reaction times? *Journal of Motor
Behavior*, 25, 1–10. https://doi.org/10.1080/00222895.2018.1518311

Brown, L. E., Marlin, M. C., & Morrow, S. (2015). On the contributions of vision and proprioception to
the representation of hand-near targets. *Journal of Neurophysiology*, *113*(2), 409–419.
https://doi.org/10.1152/jn.00005.2014

Buneo, C. A., & Andersen, R. A. (2006). The posterior parietal cortex: sensorimotor interface for the
planning and online control of visually guided movements. *Neuropsychologia*, *44*(13), 2594–2606.
https://doi.org/10.1016/j.neuropsychologia.2005.10.011

Buneo, C. A., Jarvis, M. R., Batista, A. P., & Andersen, R. A. (2002). Direct visuomotor transfomrations for
reaching. *Nature*, *416*(3), 632–636.

Burns, J. K., Nashed, J. Y., & Blohm, G. (2011). Head roll influences perceived hand position. *Journal of
Vision*, *11*(9), 3. https://doi.org/10.1167/11.9.3

Burns, J. K., & Blohm, G. (2010). Multi-sensory weights depend on contextual noise in reference frame
transformations. *Frontiers in Human Neuroscience*, *4*, 221.
https://doi.org/10.3389/fnhum.2010.00221

Chapman, C. S., & Goodale, M. A. (2008). Missing in action: The effect of obstacle position and size on
avoidance while reaching. *Experimental Brain Research*, *191*(1), 83–97.
https://doi.org/10.1007/s00221-008-1499-1

Chua, R., & Elliott, D. (1993). Visual regulation of manual aiming. *Human Movement Science*, *12*(4), 365–

401. https://doi.org/10.1016/0167-9457(93)90026-L

Cohen, R. G., Biddle, J. C., & Rosenbaum, D. A. (2010). Manual obstacle avoidance takes into account
visual uncertainty, motor noise, and biomechanical costs. *Experimental Brain Research*, *201*(3),
587–592. https://doi.org/10.1007/s00221-009-2042-8

Cohen, Y. E., & Andersen, R. A. (2002). A common reference frame for movement plans in the posterior
parietal cortex. *Nature Reviews Neuroscience*, *3*(7), 553–562. https://doi.org/10.1038/nrn873

De Haan, A. M., Van der Stigchel, S., Nijnens, C. M., & Dijkerman, H. C. (2014). The influence of object
identity on obstacle avoidance reaching behaviour. *Acta Psychologica*, *150*, 94–99.
https://doi.org/10.1016/j.actpsy.2014.04.007

Dessing, J. C., Byrne, P. A., Abadeh, A., & Crawford, J. D. (2012). Hand-related rather than goal-related
source of gaze-dependent errors in memory-guided reaching. *Journal of Vision*, *12*(11), 17.
https://doi.org/10.1167/12.11.17

Engel, K. C., Flanders, M., & Soechting, J. F. (2002). Oculocentric frames of reference for limb movement.
Archives Italiennes de Biologie, 140(3), 211-219.

Faisal, A. A., Selen, L. P., & Wolpert, D. M. (2008). Noise in the nervous system. *Nature reviews.*
Neuroscience, *9*(4), 292–303. doi:10.1038/nrn2258

Hamilton, A. F. D. C., & Wolpert, D. M. (2002). Controlling the statistics of action: obstacle avoidance.
Journal of Neurophysiology, *87*(5), 2434–2440. https://doi.org/10.1152/jn.00875.2001.

Harris, C. M., & Wolpert, D. M. (1998). Signal-dependent noise determines motor planning. *Nature*,
394(6695), 780-784. https://doi.org/10.1038/nature03031.1.

Heath, M. (2005). Role of limb and target vision in the online control of memory-guided reaches. *Motor*
Control, *9*, 281–309. https://doi.org/10.1123/mcj.9.3.281

Heath, M., Westwood, D. A., & Binsted, G. (2004). The control of memory-guided reaching movements
in peripersonal space. *Motor Control*, *8*, 76–106. https://doi.org/10.1123/mcj.8.1.76

Henriques, D. Y., Klier, E. M., Smith, M. A., Lowy, D., & Crawford, J. D. (1998). Gaze-centered remapping
of remembered visual space in an open-loop pointing task. *The Journal of Neuroscience, 18*(4),
1583–1594.

Henriques, D. Y. P., & Crawford, J. D. (2000). Direction-dependent distortions of retinocentric space in
the visuomotor transformation for pointing. *Experimental Brain Research*, *132*(2), 179–194.
https://doi.org/10.1007/s002210000340

Howard, I. S., Ingram, J. N., & Wolpert, D. M. (2009). A modular planar robotic manipulandum with end-
point torque control. *Journal of Neuroscience Methods*, *181*(2), 199–211.
https://doi.org/10.1016/j.jneumeth.2009.05.005

Jacobson, M., & Matthews, P. (1998). Generating uniformly distributed random Latin squares. *Journal of*
Combinatorial Designs, *4*(6), 405–437. https://doi.org/10.1002/(SICI)1520-
6610(1996)4:6<405::AID-JCD3>3.0.CO;2-J

Johansson, R. S., Westling, G., Bäckström, A., & Flanagan, J. R. (2001). Eye-hand coordination in object
manipulation. *The Journal of Neuroscience, 21(17), 6917–6932.*
https://doi.org/10.1523/JNEUROSCI.21-17-06917.2001
668

Khan, M. A., Elliott, D., Coull, J., Chua, R., & Lyons, J. (2002). Optimal control strategies under different
feedback schedules: Kinematic evidence. *Journal of Motor Behavior*, *34*(1), 45–57.
https://doi.org/10.1080/00222890209601930

Knudsen, E. (2002). Computational maps in the brain. *Annual Review of Neuroscience*, *10*(1), 41–65.
https://doi.org/10.1146/annurev.neuro.10.1.41

Knudsen, E. I., du Lac, S., & Esterly, S. D. (1987). Computational maps in the brain. *Annual Review of*
Neuroscience, *10*, 41–65. https://doi.org/10.1146/annurev.ne.10.030187.000353

Lacquaniti, & Caminiti. (2003). Visuo-motor transformations for arm reaching. *European Journal of*
Neuroscience, *10*(1), 195–203. https://doi.org/10.1046/j.1460-9568.1998.00040.x

Lewald, J., & Ehrenstein, W. H. (2002). Auditory-visual spatial integration: A new psychophysical
approach using laser pointing to acoustic targets. *The Journal of the Acoustical Society of America*,
104(3), 1586–1597. https://doi.org/10.1121/1.424371

McGuire, L. M. M., & Sabes, P. N. (2009). Sensory transformations and the use of multiple reference
frames for reach planning. *Nature Neuroscience*, *12*(8), 1056–1061.
https://doi.org/10.1038/nn.2357

Menger, R., Dijkerman, H. C., & Van der Stigchel, S. (2013). The effect of similarity: non-spatial features
modulate obstacle avoidance. *PLoS ONE* 8(4): e59294.
https://doi.org/10.1371/journal.pone.0059294

Menger, R., Van der Stigchel, S., & Dijkerman, H. C. (2012). How obstructing is an obstacle? The
influence of starting posture on obstacle avoidance. *Acta Psychologica*, 141, 1-8.
https://doi.org/10.1016/j.actpsy.2012.06.006

Menger, R., Van Der Stigchel, S., & Dijkerman, H. C. (2013). Outsider interference: No role for motor
lateralization in determining the strength of avoidance responses during reaching. *Experimental*
Brain Research, *229*(4), 533–543. https://doi.org/10.1007/s00221-013-3615-0

Ramsay, J., & Silverman, B. W. (2005). *Functional data analysis*. New York, NY: Springer.
https://doi.org/10.1007/b98888

Ross, A. I., Schenk, T., Billino, J., Macleod, M. J., & Hesse, C. (2018). Avoiding unseen obstacles:
Subcortical vision is not sufficient to maintain normal obstacle avoidance behaviour during
reaching. *Cortex*, *98*, 177–193. https://doi.org/10.1016/j.cortex.2016.09.010

Ross, A. I., Schenk, T., & Hesse, C. (2015). The effect of gaze position on reaching movements in an
obstacle avoidance task. *PLoS ONE*, *10*(12), e0144193.
https://doi.org/10.1371/journal.pone.0144193

Sabes, P. N. (1997). Obstacle avoidance and a perturbation sensitivity model for motor planning. *The*
Journal of Neuroscience, *17*(18), 7119–7128.

Sabes, P. N., Jordan, M. I., & Wolpert, D. M. (1998). The role of inertial sensitivity in motor planning. *The*
Journal of Neuroscience, *18*(15), 5948–5957. https://doi.org/10.1523/jneurosci.18-15-05948.1998

Saunders, J. A. (2004). Visual feedback control of hand movements. *The Journal of Neuroscience*, *24*(13),
3223–3234. https://doi.org/10.1523/jneurosci.4319-03.2004

Saunders, J. A., & Knill, D. C. (2003). Humans use continuous visual feedback from the hand to control
fast reaching movements. *Experimental Brain Research*, *152*(3), 341–352.
https://doi.org/10.1007/s00221-003-1525-2

Schlicht, E. J., & Schrater, P. R. (2007). Impact of coordinate transformation uncertainty on human
sensorimotor control. *Journal of Neurophysiology*, *97*(6), 4203–4214.
https://doi.org/10.1152/jn.00160.2007

Schütz I., Henriques D.Y.P., & Fiehler K. (2013) Gaze-centered spatial updating in delayed reaching even
in the presence of landmarks. *Vision Research*, 87, 46–52

Scott, S. H., & Norman, K. E. (2003). Computational approaches to motor control and their potential role
for interpreting motor dysfunction. *Current Opinion in Neurology*, 16:693–698.
10.1097/01.wco.0000102631.16692.71

Scott, S. H. (2004). Optimal feedback control and the neural basis of volitional motor control. *Nature*
Reviews Neuroscience, *5*(7), 532–545. https://doi.org/10.1038/nrn1427

Sober, S. J., & Körding, K. P. (2012). What silly postures tell us about the brain. *Frontiers in Neuroscience*,
6, 5–6. https://doi.org/10.3389/fnins.2012.00154

Sober, S. J., & Sabes, P. N. (2003). Multisensory integration during motor planning. *The Journal of*
Neuroscience, *23*(18), 6982–6992.

Sober, S. J., & Sabes, P. N. (2005). Flexible strategies for sensory integration during motor planning.
Nature Neuroscience, *8*(4), 490–497. https://doi.org/10.1038/nn1427

Soechting, J. F., Flanders, M. (1992). Moving in three dimensional space: frames of reference, vectors,
and coordinate systems. *Annual Review of Neuroscience*, 167–191.

Todorov, E. (2004). Optimality principles in sensorimotor control. *Nature Neuroscience*, *7*(9), 907–915.
https://doi.org/10.1038/nn1309

Todorov, E., & Jordan, M. I. (2002). Optimal feedback control as a theory of motor coordination. *Nature*
Neuroscience, *5*(11), 1226–1235. https://doi.org/10.1038/nn963

Van Beers, R. J., Baraduc, P., & Wolpert, D. M. (2002). Role of uncertainty in sensorimotor control.
Philosophical Transactions of the Royal Society B: Biological Sciences, *357*(1424), 1137–1145.
https://doi.org/10.1098/rstb.2002.1101

Vaziri, S., Diedrichsen, J., & Shadmehr, R. (2006). Why does the brain predict sensory consequences of
oculomotor commands? Optimal integration of the predicted and the actual sensory feedback. *The*
Journal of Neuroscience, *26*(16), 4188–4197. https://doi.org/10.1523/jneurosci.4747-05.2006

Voudouris, D., Smeets, J. B. J., & Brenner, E. (2012). Do obstacles affect the selection of grasping points?

739   *Human Movement Science, 31*(5), 1090–1102. https://doi.org/10.1016/j.humov.2012.01.005