How to Deal with Multiplicity in Neuroimaging?
A Case for Global Calibration

Gang Chen[*a], Paul A. Taylor[a], Robert W. Cox[a], and Luiz Pessoa[b,c,d]

[a]Scientific and Statistical Computing Core, National Institute of Mental Health, USA
[b]Department of Psychology, University of Maryland, College Park, USA
[c]Department of Electrical and Computer Engineering, University of Maryland, College Park, USA
[d]Maryland Neuroimaging Center, University of Maryland, College Park, USA

Abstract

Neuroimaging faces the daunting challenge of multiple testing – an instance of *multiplicity* – that is associated with two other issues to some extent: low inference efficiency and poor reproducibility. Typically, the same statistical model is applied to each spatial unit independently in the approach of massively univariate modeling, and corrections for multiple testing are applied to account for the existence of tens of thousand of simultaneous statistical inferences. In dealing with multiplicity, the general strategy employed in the field is the same regardless of the specifics: trust the local effect estimates (e.g., voxel-wise effect magnitude) while adjusting the extent of statistical evidence at the global level to account for the overall false positive rate. However, in this approach, modeling efficiency is compromised because each spatial unit (e.g., voxel, region, matrix element) is treated as an isolated and independent unit during massively univariate modeling. In addition, the required step of multiple testing "correction" by taking into consideration spatial relatedness, or *neighborhood leverage* (such as the extent of statistical evidence), can only partly recoup statistical efficiency, resulting in potentially excessive penalization, as well as arbitrariness due to thresholding procedures. The present paper reviews how Stein's paradox (1956) motivates a Bayesian multilevel (BML) approach that, rather than fighting multiplicity, embraces it to our advantage through a *global calibration* process among spatial units. Global calibration is accomplished via a Gaussian distribution for the cross-region effects whose properties are not a priori specified, but a posteriori determined by the data at hand through the BML model. Our framework therefore incorporates multiplicity as integral to the modeling structure, not a separate correction step. The central objective is to turn multiplicity into a strength by improving model efficiency and the overall predictive accuracy across all spatial units simultaneously. In addition, our proposal reverberates with recent proposals to eliminate the dichotomization of statistical evidence ("significant" vs. "non-significant"). We thus hope that it can help improve the interpretability of study findings, as well as promote reporting the full gamut of results (not only "significant" ones), thereby enhancing research transparency and reproducibility.

Introduction

Neuroimaging techniques often measure multiple signals simultaneously, such as from electro- or magneto-encephalogram sensors, or at spatial locations across the brain in functional magnetic resonance imaging (FMRI). In the former case, one may have tens to hundreds of nodes, while in the latter, one may have signals at over 100,000 spatial units (voxels and/or grayordinates) in the brain. A researcher will then be interested in evaluating how signals vary according to experimental conditions. This is not your typical statistical inference scenario with, for example, behavioral data. So, how does one proceed?

---

[*]Corresponding author. E-mail address: gangchen@mail.nih.gov

In FMRI, the conventional analysis at the group level adopts a so-called *massively univariate analysis* approach with a two-step procedure[1]. First, one performs a statistical analysis at every location separately but usually with the same model (or design matrix). The analysis itself can be a simple *t*-test (do signals during conditions A and B differ convincingly?) or a more elaborate linear model (ANOVA, GLM, or mixed-effects model) with several predictors and interaction terms (is there robust evidence for a three-way interaction?). But testing at multiple locations simultaneously forces the investigator to rethink her approach of controlling for the overall false positive rate (FPR) under the null hypothesis significance testing (NHST) framework. One is not simply worried about errors at a single location but at all locations of the dataset *at the same time* – hence the need for a second step. This multiple testing problem has led to beautiful and creative solutions by the neuroimaging community in the past quarter of century, including random field theory (Worsley et al., 1992), Monte Carlo simulations (Forman et al., 1995), and permutation testing (Nichols and Holmes, 2001; Smith and Nichols, 2009). However, we believe that the current approaches have important shortcomings.

Consider the two-step procedure. In the first step, by performing analysis at every location separately, one is assuming that all spatial units are unrelated to each other; in other words, they do not share information. Accordingly, there will be as many models specified as the number of the spatial units, leading to potential overfitting and model inefficiency. A model specifies how the data are related to the effect in question, such as the difference of two means in a *t*-test or GLM. Whatever type of model is adopted in the first step, the investigator is forced by the presence of simultaneous inferences to follow up with a second step to adequately control for the overall chance of misidentification in the reported results. As mentioned, a series of correction methods have been developed in the past.

In the present paper, we argue that such a massively univariate approach leads to information waste and inefficient modeling. There may be a misperception outside the neuroimaging community that statistical analysis with brain data is untrustworthy, as if multiple testing were usually not corrected due to the misreading of the oft-noted dead salmon study that won the Ig Nobel prize (Bennet et al., 2010). The reality might be the opposite because of the modeling inefficiency. The multiplicity challenges are far from trivial and remain active research topics, as evinced in a recent paper that stirred substantial controversy (Eklund et al., 2016). Against this backdrop, we propose an *integrative approach* that incorporates all the spatial units into a single, unified Bayesian multilevel (BML) model, leading to potentially improved inference efficiency through *global calibration*. The BML framework also facilitates transparency, reporting a study's full results instead of artificial dichotomization ("significant vs. non-significant"), and model validation. Accordingly, the modeling framework resonates with critical assessments of the NHST framework which have intensified in recent years (Nuzzo, 2014; Wasserstein and Lazar, 2016; McShane et al., 2018; Amrhein et al., 2019).

Controlling overall FPR under the conventional framework

We start with a brief review of how multiplicity is handled under NHST in neuroimaging. The most basic correction for multiple testing is obtained via Bonferroni correction, namely by raising the bar for statistical evidence at each spatial unit by dividing the desired significance level by the number of spatial units (e.g., setting $\alpha = 0.0000005$ if tests will be performed at $100,000$ locations). However, this approach assumes that effects at every spatial location are fully independent, an assumption that is violated in the case of FMRI data, making this classical approach highly conservative in general (i.e., there are fewer independent tests than the number of voxels). Investigators recognized early on that the probability that a clique of spatial units (such as a cluster of voxels) would be "active" together was much smaller than that of individual units. Accordingly, such spatial relatedness (i.e., voxel adjacency) has motivated most methodologies aimed at addressing multiplicity. Typical correction efforts belong to two main categories: 1) controlling for family-wise error, so that the overall FPR at the whole-brain or cluster level is approximately at the nominal value, and 2) controlling for false discovery rate (FDR), which targets the expected proportion of identified items ("discoveries") that are incorrectly labeled (Benjamini and Hochberg, 1995). FDR can be used to handle a needle-in-haystack problem, where a small number of effects exists among a sea of zero effects in, for example, bioinformatics. However, FDR is usually conservative for typical neuroimaging data and thus is less widely adopted.

Multiple testing correction methods tend to be "add on" procedures that are separate from the individual statistical models of interest. Therefore, they contain an element of *arbitrariness*, as evidenced by the existence of varied "correction" techniques across the field. Conceptually, the situation is similar to that of defining an island using "pixels" in a satellite image as a piece of sub-continental land above a fluctuating sea level (Fig. 1): the definition of the island will depend on

---

[1] The two steps discussed here should not be confused with the "two steps" of first estimating regression coefficients at the individual-participant level and then employing those at a second step of statistical testing across participants.

Figure 1: Schematic illustration of handling multiplicity via neighborhood leveraging procedures. The process is illustrated by considering three hypothetical spatial clusters that can be imagined as three sub-continental landmasses. The "pass" or "fail" of a cluster depends on its size (cross-section extent) at a particular voxel-wise threshold (or sea level; showcased by the blue segment length). In other words, only the neighborhood defined by that particular threshold matters in the sense that the spatiotemporal information contained in the data is summarized by a snapshot (a cluster defined by a cutoff). At the level of t_1, clusters A and B survive (green "+" sign) based on the spatial threshold T_1 while C does not (red "×" sign) because of its small size. In contrast, at the stringent level t_3, both clusters A and B fail while C survives. Note that the surviving clusters strongly depend on the "sea level" adopted, and no single case is ideal. In particular, if an anatomical region is intrinsically small, clustering will often fail to reveal it unless the statistical evidence is unusually strong.

the water-level threshold adopted, which itself may vary with tide, season, geological time, and a minimal area setting (e.g., 10,000 square meters). In the conventional statistics framework, the thresholding bar ideally plays the role of winnowing the wheat (true effect[2]) from the chaff (random noise), and a p-value of 0.05 is commonly adopted as a benchmark for comfort in most fields. However, a problem facing correction methods for multiple testing is the arbitrariness surrounding thresholding and that of the cut-off value itself (why not use 0.04 or 0.06, for instance?). Thresholding is a binary operator that splits the outcome into categories in a way that in practice entails the belief that the effect is artificially divided into "real vs. non-existing"; that is, data are separated into those that the investigator "cares about" and the "rest".

Available methods

At present, there are four general approaches of neighborhood leverage to handling multiplicity in neuroimaging: two of them can be called "cluster-based" and two can be called "permutation-based". The two cluster-based methods are Random Field Theory (as used in SPM) and Monte Carlo simulations (as adopted in AFNI): starting with a voxel-wise probability threshold (e.g., 0.01, 0.001) at the voxel level, a spatial-extent threshold is subsequently determined that specifies the minimal cluster size that should be believed, therefore controlling the overall FPR at the cluster level – the idea is that one should believe in the islands, not in the image pixels. The voxel-level p-value and the cluster size trade-off against each other, with a higher statistical voxel-level threshold leading to a smaller cluster size cutoff. Thus, a small region of activation can only gain ground with a low p-value at the voxel level, while large regions with a relatively large p-value at the voxel level may fail to survive the criterion (Fig. 1 at the threshold t_3). Similarly, a lower statistical threshold (higher p-value) requires a larger cluster volume, so that smaller regions have little chance of reaching the survival level (Fig. 1 at the level t_1 or t_2). (For the investigator attracted to small regions of the brain, such as the amygdala or thalamic subnuclei, this poses a considerable headache.) The arbitrariness of the statistical threshold at the voxel level poses another challenge to the investigator: one typically will lose spatial specificity with a low threshold since small regions that are contiguous will become part of large clusters of activation (at times spanning hundreds of cubic milliliters in the brain); but note that the ability to detect large regions of activation is compromised when a high statistical threshold is chosen. A recent critique of the approach of cluster formation (Eklund et al., 2016) has resulted in a trend to require a higher statistical bar in neuroimaging (a voxel-wise threshold below 0.001). However, shifting the threshold does not affect the arbitrariness associated with the thresholding procedure itself: it is a corollary of dichotomization.

---

[2]Note that the concept of *true* effect only makes sense under the modeling framework at hand.

Alternative methods exist that employ permutation testing procedures. The maximum statistic approach (Nichols and Holmes, 2001), an early version, starts with the construction of a permutation-based null distribution of a maximum statistic (either maximum testing statistic or maximum cluster size) at a predetermined primary threshold. The original data are assessed against the null distribution, and the top winners at a designated rate (e.g., 5%) are declared as the surviving ones. While the approach is effective at maintaining the nominal FPR level, as in the parametric case above, the results will depend on the threshold employed, often strongly so. The maximum statistic approach has been extended to the analysis of matrix data, implemented in packages such as network-based statistics (NBS) (Zalesky et al., 2010) and the CONN toolbox (Whitfield-Gabrieli and Nieto-Castanon, 2012). Again, thresholding decisions may lead to disparate results when a different primary threshold is adopted – even as the same data are analyzed.

More recent permutation-based approaches take into consideration both effect magnitude and spatial extent (Smith and Nichols, 2009), and have been implemented in programs such as Randomise and PALM in FSL using threshold-free cluster enhancement (Smith and Nichols, 2009), and in 3dttest++ in AFNI using equitable thresholding and clusterization (Cox, 2019). A similar approach has also been borrowed in the analysis of matrix data (Bagio et al., 2018). These techniques circumvent the problem of having to choose a primary voxel-level threshold, thus ameliorating the issue of requiring a predetermined threshold in cluster-based methods. Nevertheless, the approach may come with some loss in statistical efficiency. For example, it is possible for a spatial clique to survive a cluster-based method that utilizes a primary threshold but not survive the permutation approach that combines both signal strength and statistical evidence (e.g., permutation versus cluster-based approach adopting a primary threshold at t_3 in Fig. 1), especially when the spatial clique sizes are relatively homogeneous. On the other hand, the permutation approach might achieve higher sensitivity when the spatial size varies substantially (e.g., permutation versus cluster-based approach adopting a primary threshold at t_2 in Fig. 1).

Regardless of the specifics of each adjustment method, the effect of interest is assessed through a model applied separately at each spatial unit under the current modeling framework. Yet, each individual unit is only meaningful when it belongs to a surviving cluster. Accordingly, the investigator cannot attach a specific uncertainty level (e.g., confidence interval under NHST) at particular locations because of the indivisible nature of the spatial cluster (Woo and Wager, 2015). As what survives is the entire cluster, the statistical significance at the voxel level is undefined.

Other issues

Another challenge facing investigators involves potential "mid-analysis" changes of correction method, or more broadly of "data domain". When a whole-brain voxel-wise analysis fails to allow a region to pass the peer-accepted threshold, is it now legitimate to go through additional analysis steps? For example, the existing literature may legitimately point towards a set of target brain regions, so should the investigator now focus on those regions, or possibly even a single "star" brain area (e.g., amygdala, accumbens, fusiform gyrus, etc.)? Changing the *model space*, such as the space of multiple testing, may seem to "improve" the strength of statistical evidence (Kruschke, 2010), despite the fact that the original data remains exactly the same! Whereas trying multiple analysis strategies until "statistical significance" is reached is clearly unacceptable, where should the line be drawn between legitimate "trying to understand one's data" and "data snooping" or p-hacking? To claim that a study would come to a halt based on a "first pass" of data analysis is admittedly naive, and does not acknowledge the actual way that science proceeds in the laboratory.

A final issue with the massively univariate modeling strategy is that it implicitly makes the assumption that the investigator is fully ignorant about the distribution of effect sizes across space. Specifically, conventional statistical inferences under NHST assume that all potential effects have the same probability of being observed; it is assumed that they follow a uniform distribution from $-\infty$ to $+\infty$ (Gelman et al., 2014). However, in practice, many types of data tend to have a density of roughly Gaussian characteristics with a bell-shaped distribution exhibiting a single peak and near symmetry (which is not surprising given that the central limit theorem applies in many situations). The information waste resulting from the assumption of uniform instead of Gaussian distribution leads to inefficient modeling due to overfitting. The assumption of uniform distribution of effect sizes should be viewed in contrast to the adoption, indeed in the same conventional methods of statistical inference, of a Gaussian distribution when experimental entities are concerned (e.g., participants, animals). In fact, per the maximum entropy principle, the most conservative distribution is the Gaussian if the data have a finite variance (McElreath, 2016). Accordingly, the question that needs to be considered is as follows: Do we truly believe that effects across spatial units are uniformly dispersed across all possible values? If the answer is "no" (e.g., extremely high or low values are nearly unattainable), why cannot we assume that the effects across spatial units follow a Gaussian, as routinely practiced for the measuring units of subjects that are considered as random samples from a hypothetical

pool, instead of adopting the stance of "full ignorance"? To anticipate, the Bayesian framework described here allows distributional assumptions to be incorporated in a principled manner, and to be validated against the observed data.

Stein's paradox and hierarchical modeling

Charles Max Stein (1956) discovered a counterintuitive phenomenon in statistics and in decision theory: When three or more effects are estimated individually, the overall accuracy of each is worse than an integrative approach that shares information across the effects.

Predicting basketball player performance

Suppose that the shooting rate of LeBron James' field goals during the 2018-2019 National Basketball Association (NBA) season is 51.0%. If one has to guess his field goal rate during the next season without any other information, his current efficiency would be the best (unbiased) estimate, even though his future performance may be better or worse, of course. The same would be true for any other player. Now consider the prediction of the future success rate of James as part of the top 50 NBA players. The surprising fact is that more accurate predictions can be produced compared to those simply using their current percentages as future predictions, as rigorously proven by James and Stein (1961). In other words, by considering each player as a member of a pool of the top 50 performers, we can make overall more accurate predictions than directly using each player's current performance as a prediction (Fig. 2). Thus, although we can reasonably assume that all players' performances are independent of one another, it is better to pool information across all players to make a prediction for each individual player. The pooling is considered "partial" when some information is borrowed from other players when predicting each individual player, in contrast to "complete" pooling[3] in which all players are assigned the same prediction (e.g., overall average) . Although Stein's paradox was originally proven for the case of Gaussian distributions, it has since been generalized to other distributions and even multivariate cases.

The moral we can learn from the NBA player example is that the fundamental difference between the two modeling approaches lies in our willingness to apply available knowledge. The at-first intuitive approach of using each player's current performance to predict future percentage (no pooling, Fig. 2A; note that current and predicted values are the same) assumes that we do not have any knowledge about the distribution of performances, and thus treat each player as an independent entity. In fact, such stance of "total ignorance" is equivalent to making the assumption that shooting rate performance is uniformly distributed. That is to say, all possible values are credible, including very low (e.g., 10%) and very high (e.g., 90%) shooting rates, which are actually not observed in practice.

It is instructive to consider the problem of simultaneously predicting the performance of the set of 50 players as a type of multiplicity problem. In this case, it is not only the individual accuracy (cf. voxel-wise p-value under NHST) that matters the most, but the overall accuracy (cf. overall FPR) that is of concern. As stated, one approach is to treat each player as independent from the rest, similar to the massively univariate approach in neuroimaging (Fig. 2A). As players do not simply repeat their performance, the observed future performance (generated hypothetically) will deviate from the preceding season to some extent (as indicated by the purple lines connecting squares to diamonds in Fig. 2A). We go through this exercise again but now pretend that we could time-travel to the future and take a sneak peek at the independent shooting rates y_i of the top 50 NBA players. If we predicted players' future performance based on current performance via linear regression, we would obtain the fit displayed in Fig. 2C (green line). The same data as used in Fig. 2A were employed, which was generated based on the assumption that future shooting rates follow a Gaussian distribution. If we plot the linear regression predictions in a manner that follows that of Fig. 2A, we can see that the predicted values differ from the current one, as can be seen by the slanted green lines in Fig. 2B. The role of the Gaussian distribution assumption of future performance can be visualized as a sort of elastic band, whereby predictions are mutually informative. Notably, predictions for outlying players (high and low scorers) are shrunk toward the group mean via the implicit partial pooling of the procedure.

In the present case, our willingness to apply prior knowledge (that shooting percentages typically follow a Gaussian distribution) transformed the problem into a simple GLM system, thereby achieving a higher overall accuracy in Fig. 2B (see the comparison between the two sums of squared residuals) than that with the assumption of no prior knowledge (represented by a uniform distribution) in Fig. 2A. Notably, the prior information (i.e., the distribution assumption) only sets the general shape while the specifics of the shape (i.e., mean and variance) can be determined from actual data through

---

[3]Complete pooling can viewed as specifying a prior with a zero variance.

Figure 2: Geometric illustration of Stein's paradox with simulated data of the top basketball players. Here, we pretend that we could time-travel to the future and take a sneak peek at the independent shooting rates y_i of the players, which are assumed to follow a Gaussian distribution: $y_1, y_2, .., y_{50} \overset{iid}{\sim} N(m_y, \sigma_y^2)$ (here, $m_y = 0.4, \sigma_y = 0.1$). LeBron James's data point is labeled as "LBJ". (A) Simple scheme in which future performance of the top 50 players is predicted (values represented by squares) to be the same as in the current year (circles). The actual "future" values are also shown (diamonds). The purple dash lines link the predicted and actual future values. (B) The performance of the top 50 players is predicted through simple regression. The slanted green lines, corresponding to the green fitted line in (C), illustrate partial pooling, which interestingly also corresponds to the concept of regression to the mean established by Francis Galton. For the hypothetical dataset at hand, note that the prediction of the next season for James as one of the top 50 players is 49.0%, a value that is slightly downgraded from his current 51.0%. (C) Two scenarios are contrasted for the top 50 players. First, the same as in panel (A): using each player's current performance to predict future values gives the diagonal line fit $x = y$ (red line). Second, we could imagine having future data and predicting future y_i values with the current data x_i as an explanatory variable. In this case, the ordinary least squares solution (green line) would produce a fitted line shallower than the diagonal (dotted red line), which illustrates the partial pooling effect (see, Stigler 1990). (D) What would happen if we predicted the performance of the top 50 players as part of a bigger pool of the top 100 players? The effect of partial pooling remains evident by the shallower line of regression fit (sold blue line). Although the prediction of the top 50 players as part of the top 100 players is different from that limited to the top 50 players (dashed green line, same as the solid green line in panel C), the prediction difference is relatively small due to the adaptive nature of the Gaussian distribution assumed for the data.

(A) Counties with highest kidney cancer death rates   (B) Counties with lowest kidney cancer death rates

Figure 3: The U. S. counties with the highest (A) and the lowest (B) 10% death rates for kidney cancer during 1980-1989 are shown with blots. Republished with permission of Taylor & Francis Group LLC, from Gelman et al. (2013); permission conveyed through Copyright Clearance Center, Inc.

the model (and estimated via ordinary least squares, or maximum likelihood). Thus, partial pooling is adaptive in the sense that information about centrality and spread of the players' performances is not required as part of the prior information, but is determined from the data. The adaptivity of the Gaussian prior can be further illustrated as follows. The prediction for the top 50 players through regression is different when they are considered as a subset of the top 100 players. However, the difference due to the addition of the 51th-100th players is relatively small (Fig. 2D).

Epidemiology of kidney cancer

An example conceptually closer to the situation in neuroimaging involves epidemiological survey data regarding the highest and lowest cancer rates across the United States (Fig. 3). Examination of the map indicates that the highest kidney cancer rates were observed in relatively sparsely populated regions of the country (Fig. 3A). One then might be tempted to infer that geographical factors explain the high rates, such as restricted access to healthcare, low utility infrastructure, or higher doses of radiation. But it is difficult to reconcile these explanations with the observation that counties with the lowest kidney cancer death rates also tend to be less populated (Fig. 3B). A more parsimonious explanation of the data, which is consistent with both maps, is based on the small sample size of rural counties, combined with the relatively low mortality rate due to kidney cancer (which is a rare disease). To see this, suppose that the national average death rate is around 1 per 2,000 people (per 10 years). In a county with a population of about 1,000 people, one can imagine observing zero or one deaths by chance alone, which would place the county into the map with the lowest or highest rate, respectively.

So, based on the survey data, how should we make reasonable inferences about *all* the individual counties? Two questions can be posed: 1) Should we really believe that a resident in a specific county has a higher (or lower) probability of dying of kidney cancer than the national average? 2) Can we infer that a resident in a county in Fig. 3A (or Fig. 3B) has a much higher (or lower) probability of dying of kidney cancer than a person in a county in Fig. 3B (or Fig. 3A)? That is, would we be able to confidently recommend that someone move from a county in Fig. 3A to one in Fig. 3B, or to discourage someone from moving from a county in Fig. 3B to one Fig. 3A? The answers would appear to be "no" in both cases.

The epidemiological distribution of kidney cancer deaths can be seen as involving a multiplicity problem over the several thousands of districts. A natural solution might be to borrow the idea of neighborhood leverage in neuroimaging (that is, spatial clustering), and to utilize the relatedness among adjacent counties to modulate the statistical evidence for clusters of counties. Note, however, that this leveraging approach only adjusts the extent of statistical evidence, and it does not modify the death rate estimates themselves (at county or cluster level). In other words, the conventional neuroimaging approach to handling multiplicity does not address two types of estimation errors that are important to consider (Gelman and Tuerlinckx, 2000). One is the so-called *type M* error, which refers to a potential over- or under-estimation of the effect magnitude. The second is the so-called *type S* error, which refers to getting the sign of a comparison wrong (does county A have a higher or lower death rate than county B?). A potential solution is to employ a methodology that prevents the estimation process from being swayed by larger fluctuations.

Consideration of Stein's paradox in the context of the kidney cancer case suggests potential strategies to address the multiplicity issue. One possibility is to not place trust only in isolated bits of survey data (at the county level), but instead to incorporate all counties into a Bayesian model with a hierarchical structure that leverages death rate information across counties (Gelman et al., 2014). The BML model reflects the central concept behind Stein's paradox:

Figure 4: Illustration of the concept and interpretation for type I (FPR), type II, type S and type M errors. Suppose that there is a hypothetical noncentral t_{20} distribution (solid black curve) for a true effect (blue vertical line) of 0.7 (A) or 0.3 (B) and a standard error of 1.0. Under the null hypothesis (red vertical line and dot-dashed green curve), two-tailed testing with a type I error of 0.05 leads to having thresholds at ± 2.09; FPR $= 0.05$ corresponds to the null distribution's total area of the two tails (marked with red diagonal lines). The power (shaded in blue) is the total area of the t_{20} distribution for the true effect (black curve) beyond these thresholds, which is 0.10 (A) or 0.06 (B). The *type S* error is the ratio of the blue area in the true effect distribution's left tail beyond the threshold of -2.09 to the total area of both tails, which is 4% (A) or 21% (B) (i.e., the ratio of the "statistically significant" area in the wrong-signed tail to that of the total "statistically significant" area). If a random draw from the t_{20} distribution under the true effect happens to be 2.2 (small gray square), it would be identified as statistically significant at the 0.05 level, and the resulting *type M* error would quantify the magnification of the estimated effect size as $2.2/0.7 \approx 3.1$ (A) or $2.2/0.3 \approx 7.3$ (B), which is substantially larger than unity. These two plots aptly demonstrate the importance of controlling the errors of incorrect sign and incorrect magnitude when a large amount of variability exists in the data. High variance is bound to occur when we deal with the multiplicity issue embedded with massively univariate modeling, which is further exacerbated by scenarios such as small sample size, noisy data, unaccounted for cross-subjects variability and suboptimal alignment to a standard template. More systematic exploration and comparison between the conventional (type I and type II) and the new (type S and type M) sets of errors can be found in Gelman and Tuerlinckx (2000) and Chen et al. (2019b).

We should be somewhat sceptical about individual inferences of effect estimates and their uncertainties, especially extreme ones. Furthermore, we should adopt some information sharing – what is called partial pooling or shrinkage because of the tendency to reduce extreme values – which is possible through a hierarchical model that shares information across all, not just neighboring, spatial units. One may feel uneasy about partial pooling because of its introduction of *bias* to individual effect estimates[4]. The seemingly puzzling phenomenon of Stein's paradox aptly captures the biased predictions about James' future performance among his NBA peers (remember that the unbiased estimates should follow the $y = x$ red dotted lines in Fig. 2). In the example, we shifted our focus from the accuracy of one particular effect (James' future shooting rate; or, in the cancer case, the death rate at one specific county) to the *overall* predictive accuracy among all effects (overall accuracy among all the 50 players or among all US counties).

To recapitulate, a fundamental difference between the conventional, isolated modeling approach and the integrative BML method is that the former seeks statistical unbiasedness at each spatial unit, while accepting a daunting FPR problem that requires corrective procedures. Having an unbiased estimator (the quintessential example is probably the sample mean as an estimate of the population mean) may sound appealing, but it is not always the most desirable property, as recognized by statisticians and applied scientists for some time. This is especially the case when sample size is not large or when noise potentially overwhelms the effect, conditions all too often encountered by experimentalists in general and neuroimagers in particular. Briefly, in many circumstances, an unbiased estimator may have high variance. In such cases, it can fluctuate non-trivially from sample to sample. Adopting a biased estimator, instead, may be beneficial if it has lower variance; thus, although the estimator does not tend, in the limit (in practice with rather large sample sizes) to the "true" value of the effect of interest, it can prove advantageous in many practical settings.

As schematically demonstrated in Fig. 2, the introduction of a Gaussian distribution introduces a mild constraint when estimating all players' performances simultaneously (their shooting rates are not free to vary everywhere, but are informed by the entire set of performances in a Gaussian fashion). In fact, many conventional methodologies employ constraints to their estimation procedures. More technically, they benefit from regularization (Tikhonov et al., 1998), which involves introducing additional information in order to solve an ill-posed problem or to prevent overfitting. For example, ridge regression, LASSO (Tibshirani, 2009), elastic nets, and linear mixed-effects framework introduce additional constraints to the estimated regression coefficients in addition to minimizing residuals (e.g., the LASSO attempts to keep the overall "budget" of weights under a certain value: $\sum_j |\beta_j| \leq k$). It is also interesting to note that global calibration is employed in meta analysis (Glass, 1976). When summarizing multiple studies of an effect, there is no correction for multiplicity of statistical evidence, but instead effects are weighted based on their respective relative reliability.

Some properties of the BML to be described further below include the following. First, the multiplicity issue is

---

[4]Recall that, in statistics, the bias of an estimator is the difference between this estimator's expected value and the "true" value of the parameter being estimated. An estimator with zero bias is thus called unbiased, otherwise it is called biased.

automatically dissolved given that a *single* model is employed – there is no "multiple" to correct! In addition to being conservative about extreme effects due to the shrinkage phenomenon, there is only one overall posterior that is formulated as a joint distribution in a high-dimensional parameter space (the dimensionality of which equals the number of parameters in the model). No multiplicity is incurred because 1) all the spatial units are included in one model and regularized through a reasonable prior, and 2) only one overall posterior distribution is utilized to predict various effects of interest. In fact, the analyst can make inference for as many effects of interest (including comparisons) as desired because each effect of interest is simply a marginal distribution (or a perspective projection) of the overall high-dimensional joint posterior distribution (like various angles of projecting the earth onto a 2D map). Second, an important focus becomes addressing type M and type S errors (see Fig. 4) (Gelman and Tuerlinckx, 2000), both of which are interpretable in practice. Third, model efficiency and performance can be directly assessed and compared straightforwardly through predictive accuracy and cross validations, similar to techniques such as least squares, maximum likelihood and information criterion. One predictive accuracy indicator is the Watanabe-Akaike information criterion, for instance, via leave-one-out cross-validation (Vehtari et al., 2017), and another one involves "posterior predictive checks", which simulate replicated data under the fitted model so as to graphically compare actual data to model predictions.

Focusing on the research hypothesis instead of fighting a "strawman"

Before applying the BML approach to FMRI data, it is instructive to briefly compare the frameworks of NHST inference (as commonly employed by experimentalists) and the proposed Bayesian approach. Consider a scenario in which a single one-sample t-test with 20 degrees of freedom (e.g., 21 subjects) is employed in the NHST setting. The null hypothesis H_0 is that the population mean is zero. Suppose that the data indicate that $t_{20} = 2.85$. If H_0 were true, the probability of observing a t-value so extreme is rather low ($p = 0.01$ in Fig. 5, left). The t-value value thus provides a measure of "surprise": How surprising would it be to observe such an extreme value in a world in which H_0 were really true? In this setting, one estimates the extent of surprise $P(\text{data} \mid H_0)$. By custom if $P < 0.05$, one declares that the effect is "present", with the all-important "significance" label.

The Bayesian framework aims to answer a different, though, related question: What is the probability of a research hypothesis H_R based on the observed data, $P(H_R \mid \text{data})$? Note the difference in what is being "measured" and what is "given" in this proposition, as opposed to the preceding NHST case. Such a probability can be computed by using Bayes' rule (Kruschke, 2010). In a typical setting the research hypothesis H_R refers to an effect or parameter θ (e.g., "population mean") being positive or negative (e.g., $H_R : \theta > 0$). An attractive property of this framework is that it is not typically formulated to generate a binary decision ("real effect" vs. "noise", or "significant" vs. "not significant") but instead to obtain the entire probability density distribution associated with $P(\theta \mid \text{data})$ (Fig. 5, right). This posterior distribution is interpreted in a natural way even though it may take getting used to for those who are unfamiliar with Bayesian inferences. For example, $P(0.1 < \theta < 0.2 \mid \text{data})$ is the area under the curve within the effect interval $(0.1, 0.2)$. For convenience, one can also indicate "tail probabilities" associated with values commonly used in the literature. For example, the two-sided area indicated in green is 0.05 and the two-sided area indicated in yellow-plus-green is 0.1; these would be analogous to 95% and 90% confidence intervals, respectively, in the NHST framework. Here, we see that the value $\theta = 0$ lies inside the left green tail; thus, the probability that the parameter exceeds zero is $P(\theta > 0 \mid \text{data}) = 0.99$. One can then use the probability in question to *emphasize* or summarize the extent (e.g., "strong", "moderate", or "weak") of statistical evidence. But we stress that the goal is to quantify and qualify the evidence, not to make a binary decision in terms of "passes threshold" versus "fails to pass threshold". In this manner, the value of $P(H_R \mid \text{data})$ is not used for declaring that a result is "real" based on a threshold, but as the amount of evidence on a continuum. There is no need for thresholds and indeed one is encouraged not to use them – in the end they are arbitrary.

Applications of Bayesian multilevel modeling in neuroimaging

We now illustrate the application of Bayesian multilevel modeling to neuroimaging data. We perform group analysis as a prediction process for all spatial units simultaneously in a single model, which assesses the statistical evidence for the research hypothesis H_R based on the available data, that is, $P(H_R \mid \text{data})$.

Figure 5: Different probability definition and focus between conventional and Bayesian frameworks. Statistical inferences under NHST usually pivot around the "weirdness indicator" of the p-value (left curve): green (or yellow) tails symbolize a two-sided significance level of 0.05 (or 0.1). If the data renders a t_{20} value of 2.85 (small gray square), reaching the comfort zone (green or yellow area) with a two-tailed p-value of 0.01 (the probability of obtaining such data with a t_{20} value at least as extreme if the effect were truly zero), one may declare to have strong evidence for the effect with an FPR threshold of 0.05 under the NHST framework. In contrast, inferences under the Bayesian framework directly address the research interest (right curve): what is the probability of the effect magnitude being greater than 0 with the data at hand (greater than 0.99 in this case)? There are different ways to provide a point estimate of centrality from the posterior distribution such as mean, median and mode (or maximum a posteriori probability (MAP)) under the Bayesian framework, while such point estimates are usually the same due to symmetry under the theoretical distribution of NHST. However, Bayesian inferences tend to emphasize more the uncertainty of an effect, not its point estimate. The green and yellow tails of the posterior density mark the extent of statistical evidence associated with the two specific (two-sided 95% and 90%) uncertainty intervals, and the dotted dark green line shows the median or 50% quantile of the posterior density. Notice 1) the x-axis is different between the two densities (standardized value for NHST and effect in physical dimension for Bayesian paradigm), 2) $P(\text{data} \mid \text{zero effect})$ under NHST is conceptually and numerically most of the time not the same as $1 - P(\text{effect} > \text{ or } < 0 \mid \text{data})$ under the Bayesian framework, 3) the null distribution under NHST has a smooth and regular shape due to the assumption of a standard curve in the model as a prior while the irregular posterior distribution is formulated through random samples through Markov chain Monte Carlo simulations under Bayesian framework, and 4) compared to the conventional confidence interval that is flat and inconvenient to interpret, the posterior density provides much richer information such as spread, shape and skewness.

Region-based analysis

In a recent study (Chen et al., 2019b), we developed a BML framework for region-based group analysis. As BOLD responses approximately share the same scale, the approach allows information to be pooled and calibrated across regions to jointly make inferences at individual regions. The approach was applied to an FMRI dataset of 124 subjects, each of which had effect estimates at 21 regions. For each individual, a seed-based correlation analysis was performed with the right temporo-parietal junction as the seed (i.e., the correlation between each region and the seed). The question of interest was as follows: What is the relationship between such correlations and individual differences in a questionnaire-measure of theory of mind?

For comparison, the same data were analyzed via conventional whole-brain voxel-wise analysis, which illustrated the difficulty of detecting surviving clusters. With a primary voxel-wise p-value threshold of 0.05, 0.01 or 0.005, four clusters were observed after correction through Monte Carlo simulations (Cox et al., 2017); with a primary threshold of 0.001, only two clusters survived (Xiao et al., 2019). In contrast, with BML analysis 8 out of the 21 regions exhibited considerable evidence (Fig. 6). In a Bayesian framework, effect inference is summarized via its entire distribution, the so-called posterior density distribution. (Note that as there is no closed-form solution for the distribution, it must be determined via computational algorithms, such as Markov chain Monte Carlo (MCMC) sampling methods.) Similar to traditional approaches, one can highlight distributional tail areas to denote quantiles of interest (such as 95%), which one may use to label effects as "moderate", "strong", and so on. Consider the case of the R TPJp region (top, middle panel of Fig. 6). In this case, the probability of the effect being less than or equal to zero (blue vertical line) is rather small, as it is clearly within the green (0.05) tail. In other words, the probability that the effect is greater than 0 is the area to the right of the blue line (in this case, 0.99). In contrast, in the case of the R Insula (top, right panel of Fig. 6), the probability that the effect lies in the central white area of the plot (which includes zero) is 90%. Now consider the case of the L SFG region (second row, middle panel of Fig. 6). One possibility is to disregard this region given that the zero-effect line does not reside in the colored tails, which is the approach typically adopted under the common practice of dichotomizing. But close inspection of the posterior reveals that the probability that its effect is greater than zero is about 0.93, and the experimentalist may thus wish to deem the results as providing a "moderate" amount of statistical evidence. Regardless of the linguistic terms utilized, one does not have to dichotomize the results, and in fact the investigator is encouraged to report the full posterior[5]. It is also instructive to explicitly compare BML results with those obtained with conventional

---

[5]It is probably advisable to avoid the "codification" of linguistic labels ("moderate", "strong", etc.) that are tied to particular probability values (say, 95%). Naturally, considering values along a continuous numerical scale is an advantage of mathematics over discrete language usage.

Figure 6: Posterior density distributions of the slope effect derived from BML with 124 subjects at 21 regions. The vertical blue line indicates zero effect; orange and green tails mark the regions beyond the 90% and 95% uncertainty (compatibility or quantile) intervals, respectively. If results highlighting is desirable, one can claim the regions with strong evidence of slope effect as the blue line being within the color tails, as indicated with orange and green dot-dashed boxes. Compared to the conventional confidence interval that is flat and inconvenient to interpret, the posterior density provides much richer information about each effect such as spread, shape and skewness. Relative to the conventional whole-brain voxel-wise analysis that rendered with only two surviving clusters (Xiao et al., 2019) based on the primary voxel-wise p-value threshold of 0.001, the BML showed a much higher inference efficiency with 8 regions that could be highlighted with strong evidence. To illustrate the conventional dichotomization pitfall through a common practice of thresholding at 0.05, the region of L SFG also elicited some extent of slope effect with a moderate amount of statistical evidence: the probability that its effect is greater than zero is about 0.93 conditional on the data and the BML model.

Figure 7: Comparisons of results between the conventional region-based GLM and BML in box plots. For each region, the left bar shows the GLM result that was inferred with the region as an isolated entity while the right bar corresponds to the BML inference. With one GLM for each of the 21 regions, none of the regions would survive FPR correction under NHST. The slope effect is shown with the horizontal black bar in the middle of each box as the mean for GLM or median for BML; each box (or whisker pair) represents the 90% (or 95%) confidence interval for GLM (dashed, left) or uncertainty interval for BML (solid, right). The shrinkage or pooling impact of BML calibrates and "drags" most, but not all regions (see R TPJp).

(A) GLM posterior predictive density   (B) BML posterior predictive density

Figure 8: Model performance comparisons through posterior predictive checks and cross validations between conventional univariate GLM (A) and BML (B). The subfigures A and B show the posterior predictive density overlaid with the raw data from the 124 subjects at the 21 ROIs for GLM and BML, respectively: solid black curve is the raw data at the 21 ROIs with linear interpolation while the fat curve in light blue is composed of 500 sub-curves each of which corresponds to one draw from the posterior distribution based on the respective model. The differences between the solid black and light blue curves indicate how well the respective model fits the raw data. BML fitted the data clearly better than GLM at the peak and both tails as well as the skewness because pooling the data from both ends toward the center through shrinkage clearly validates our adoption of BML.

GLM estimates (Fig. 7). We can see that the BML estimates are pulled to the center (with the exception of region R TPJp), an overall behavior analogous to the one described for the basketball shooting performances.

An appealing aspect of Bayesian modeling is that the performance and efficiency of a model can be readily compared to alternative models with the use of cross validation tools such as information criteria and posterior predictive checks. The intuition of visual examination on predictive accuracy is that, if a model is reasonable, the data generated based on the model should look similar to the raw data at hand. For example, the massively univariate approach with GLM yielded a poor fit (Fig. 8A), whereas a much better fit was obtained with BML (Fig. 8B).

Matrix-based analysis

We now illustrate the BML framework in the context of another type of data that are of great interest to neuroimaging investigators, namely those summarized in *matrix* form. The most common example is that involving time series correlations between regions (e.g., as in resting state studies), inter-subject correlations among subjects (e.g., as employed for naturalistic scanning), or similarity measures (e.g., representational similarity, latent semantic analysis). In a typical case, an investigator may have tens or even hundreds of ROIs and wish to examine the correlation structure in the data. Consider the case in which the brain is partitioned into 100 ROIs, leading to a 100×100 correlation matrix. Given sym-

---

Nevertheless, linguistic descriptors are probably benign as long as the posterior distribution is fully presented.

Figure 9: Characterizing the inter-region correlation structure of group brain data. A) Correlations between two pairs of brain regions, r_{ij}, are not independent when they share a common region. Thus, when simultaneously estimating multiple correlations, such relatedness needs to be modeled and accounted for. B) Making inferences about correlations leads to a *multiplicity* problem, in particular how to account for the simultaneous inferences of all effects under NHST. In a Bayesian framework, multiplicity relates to the problem of modeling all correlations simultaneously by invoking information sharing or partial pooling. C) Within a Bayesian multilevel framework, it is possible to frame the problem in terms of capturing the population-level effect of (1) brain region, R_q, (2) region pair, r_{ij}, and (3) subject k. The characterization of the effect of brain regions is a unique contribution of our framework, which allows investigators to reveal a region's "importance" within a principled statistical framework.

Figure 10: Posterior density plots of region effects (in Fisher's z-value). Each posterior distribution indicates the probability of observing region effects. The orange and green tails mark areas outside the two-sided 90% and 95% quantile intervals, respectively; the blue vertical lines indicate the zero region effect. Consider a region such as the BNST_L (top row, third column): the zero region effect lies in the left green area, indicating that the probability that the effect is positive is greater or equal to 0.975 (conversely, the probability that the effect is negative is ≤ 0.025). The same is true for the Thal_R (second row, second column). In these two cases, there is strong statistical evidence of a region effect, as indicated with green dot-dashed boxes. Two other regions (BNST_R and Thal_L; orange dot-dashed boxes) exhibited moderate statistical evidence of a region effect (the blue vertical lines were each within the orange band). Four more regions forming contralateral pairs of regions (BF_L and BF_R, aIns_L and aIns_R) plus SMA_R also exhibited some statistical evidence as they were close to the typical "convenience" thresholds. Note that the posterior density provides rich information about each effect distribution, including shape, spread and skewness. Unlike the conventional confidence interval that is flat and inconvenient to interpret, it is valid to state that, conditional on the data and model, with probability, say, 95%, the region effect lies in its 95% posterior interval. Inferences about region pairs are shown in Fig. 11. Region abbreviations: BF: basal forebrain; BNST: bed nucleus of the stria terminalis; IFG: inferior frontal gyrus; IPG: inferior parietal gyrus; Ins: insula; MPFC: medial prefrontal cortex; SMA: Supplementary motor area; Thal: thalamus. Other abbreviations: a: anterior; p: posterior; m: medial; L: left; R: right.

Figure 11: Comparisons of region pair effects between univariate GLM (left) and BML (right) for a matrix dataset. Density plots like Fig. 10 are possible at the region-pair level, but only a summarized version is presented here due to space limitation. The empty entries along the diagonal correspond to the correlation value of 1. The effect magnitude (Fisher's z-value) is symbolized with both circle area and color scheme (colorbar, far right). The impact of partial pooling (or shrinkage) under BML is evident as the effects for most region pairs are "pulled" toward the middle relative to their GLM counterparts. Not to dichotomize but to highlight the extent of the statistical evidence, region pair are colored with a green or orange background based on the strength of statistical evidence associated with 95% or 90% two-sided quantiles. The high efficiency of BML becomes obvious when compared with the conventional GLM. With BLM, 33 region pairs exhibited moderate to strong statistical evidence and they formed a subset of those 62 region pairs declared under GLM. With GLM, 62 region pair were identified as statistically significant (one-sided, 0.05; green and orange boxes) without correction for multiple testing; when cluster-level correction was applied to GLM (FPR of one-sided 0.05 with permutations), no region pairs survived. A distinct feature of BML from GLM, as a natural consequence of the effect decomposition in BML, is the capability of inferring at the region level (Chen et al., 2019c) as presented in Fig. 10.

metry, one is interested in $100 \times 99/2$ entries (or more generally $\frac{1}{2}n(n-1)$ with n ROIs). Although the number of ROIs here is not particularly high, the number of pairwise inferences is 4950, which entails a non-trivial correction for multiple testing. Along the lines described in the preceding section, we developed an approach to model all pairwise correlations simultaneously. Given that a single model is involved, correcting for multiplicity is not required.

The central issue of matrix-based data analysis is to make inferences for each *region pair* which poses considerable challenges. First, correlation values are, by definition, computed over pairs of random variables (e.g., average BOLD time series for FMRI data), but the existence of "shared regions" implies that some pairwise correlations are not independent, namely they are correlated themselves (Fig. 9A). Because the correlation between regions R_i and R_j and that between regions R_j and R_k share a common region R_j, proper modeling requires accounting for such covariance between region pairs. A second challenge concerns of course the problem of multiple testing (Fig. 9B), which has been tackled via permutations, where a null distribution is generated to declare which region pairs survive thresholding (Zalesky et al., 2010).

We applied our BML approach to data from a previous cognitive-emotional task (Choi et al., 2012). Briefly, participants performed a response-conflict task (similar to the Stroop task) under safe and threat conditions. A unique feature of the approach developed is that it allows the assessment of effects at the *individual-region level* (Fig. 10). Because the effect of each region pair is decomposed in terms of multiple components including that for each individual region involved, a region-level effect that characterizes the "contribution" of a region can be inferred by BML model. Such an effect, presented with its uncertainty (and the entire posterior distribution), allows the assessment of region importance in a manner that is statistically more informative than those metrics commonly used in graph-theoretic analysis such as degrees and hubs.

Figure 11 compares the results based on the conventional GLM approach and those based on our BML proposal. The GLM results were obtained through a massively univariate analysis with $(16 \times 15)/2$ independent models, one for each region pair. The BML results were derived by fitting a single model to half of the off-diagonal elements in the correlation matrix. The impact of partial pooling (i.e., information sharing) under BML can be observed by noticing the smaller effect sizes among many region pair. In other words, the effects for most region pairs are "pulled" toward the "middle" relative to their GLM counterparts. The inference efficiency of BML can be appreciated by noticing that 33 region pairs exhibited "moderate" to "strong" statistical evidence; they formed a subset of the 62 region pairs that *individually* were below the

0.05 threshold under GLM. At the same time, when cluster-level correction for multiple testing was applied at FPR of one-sided 0.05 with permutations (Zalesky et al., 2010), no region pairs survived in the GLM case.

Discussion

The bulk of statistical training focuses on the conventional NHST paradigm, and so researchers are accustomed to accepting the framework unquestioningly. The "surprise" measure encapsulated by $P(\text{data} \mid H_0)$ has become a target goal of data analysis that takes its aim at a "straw man" scenario of "zero effect" under NHST (Fig. 5, left). Unlike situations with truly dichotomous categories (e.g., guilt or innocence in a courtroom trial), most effects or mechanisms under investigation in neuroimaging – and arguably biology more generally – exist along a continuum (e.g., Gonzalez-Castillo et al., 2012), and therefore will be considerably distorted if binarized (as in statistically "significant" vs. "non-significant").

As discussed by others in the past, a more natural objective of the experimentalist would be to focus on the research hypothesis H_R of interest, which can be assessed through $P(H_R \mid \text{data})$ (Fig. 5, right). With rather benign model assumptions, Bayesian methodology allows just that. Furthermore, advances in Bayesian methods in the past two decades are encouraging a reexamination of some entrenched practices. For example, unbiasedness is generally considered to be a "nice," if not requisite, property for an estimator. However, it might not always be the most important property. For one, when it is difficult to reach a satisfactory sample size (a nearly universal reality of experimentalists), some extent of statistical bias at the individual-entity level (e.g., participant) might be a cost worth incurring to achieve an overall higher predictive accuracy at the collective level.

Limitations of NHST applications in neuroimaging

Much has been written about NHST over the past few decades. Here, we briefly enumerate a few aspects that are relevant in the present context.

(1) **Vulnerability to misconception**. The disconnect between the p-value and the probability of research interest often leads to conceptual confusion (Nuzzo, 2014). For example, an investigator is likely more interested in the probability of a positive (or negative) effect given the data, $P(H_R \mid \text{data})$, (Fig. 5, right), which is conceptually different from, but often mistakenly construed as, the p-value.

(2) **Arbitrariness due to dichotomization**. As the underlying physiological or neurological effect is in all likelihood intrinsically continuous, the introduction of arbitrary demarcation through hard thresholding results in both information loss and distortion. Due to the common practice of NHST and the adoption of significance level as a publication-filtering criterion, a statistically non-significant result is often misinterpreted as a non-existent effect (the absence of evidence is equated with the evidence of absence). Indeed, although even fairly introductory students of statistics will explicitly be aware of this problem, in experimental literature, the practice of describing non-significant results as non-existent effects is puzzlingly widespread.

Hard thresholding leads to conceptual issues such as the following one: Is the difference between a statistically significant result and a non-significant one itself statistically significant (Fig. 12)? Translated into the neuroimaging context, when blobs of spatial units are relatively near but below the adopted threshold (given the cluster size or the integration of cluster size and statistical evidence), available software implementations hide them from the investigator (they are left uncolored, much like a location inside a ventricle). Should those "activation blobs" really be considered as totally devoid of evidence?

(3) **Arbitrariness due to model space**. When the whole-brain voxel-wise analysis fails to reveal the expected regions at currently acceptable levels of statistical evidence, how should the experimenter proceed? Indeed, investigators may feel that it is justifiable to adopt "small volume correction" by limiting the data domain, somehow; for example, by focusing on regions of interest outlined in prior literature, or possibly particular regions of outstanding interest (such as the nucleus accumbens in the domain of reward processing). But how to proceed in those circumstances is unclear as, in reality, few researchers are prepared to publicly describe their work as "purely exploratory" at that point.

(4) **Overstated estimates and distorted inference**. Reporting only statistically significant results tends to produce overestimated or inflated effect sizes (Cremers et al., 2017), as illustrated in the kidney cancer example (Fig. 3); in fact, the current publication practice of only allowing for results with stringent statistical evidence leads to many problems (Amrhein et al., 2019). Suppose that a brain region cannot survive the currently accepted correction criterion (cluster- or permutation-based) but still presents moderate amount of statistical evidence. Such result would have difficulty in getting reported under the current reviewing process, and its absence from the literature may waste the important information

Figure 12: Arbitrariness resulting from dichotomization. Two hypothetical effects that independently follow Gaussian distributions: $N(m_1, \sigma_1^2)$ (blue) and $N(m_2, \sigma_2^2)$ (red), respectively, where $m_1 = 0.2, \sigma_1 = 0.1, m_2 = 0.4, \sigma_2 = 0.3$. Under NHST the first effect would be considered statistically significant with a one-sided (or two-sided) p-value of 0.023 (or 0.045); in contrast, the second effect would not be viewed statistically significant given a one-sided (or two-sided) p-value of 0.091 (or 0.18). On the other hand, the difference between the two effects is not statistically significant with a one-sided p-value of 0.26; in fact, the second effect is more likely larger than the first one with a probability of 0.74.

that could be utilized as evidence for future studies. If the region fails to reach the designated statistical criterion among four out of ten such studies, the impact of dichotomization on meta analysis could also be substantial. We believe that the reporting of results and publication acceptance should not be based on a dichotomous decision rule in terms of a predetermined statistical significance level (e.g., 0.05).

(5) **Disregard for effect size and unavailability of uncertainty measures**. Because neuroimaging studies perform inferences across tens of thousands of spatial locations, the practice in the field is to present maps that employ colors symbolizing statistical values (Chen et al., 2017). The unavailability of effect sizes leads to crude meta analyses, and renders power analysis challenging in neuroimaging. What is more, the statistic (e.g., t-value) and the associated uncertainty (standard error or confidence interval) are frequently not directly interpretable at the spatial unit level because of the extra step required to adjust for multiple testing. In fact, as discussed, each spatial clique either passes or fails the criterion for surviving; accordingly, no specific uncertainty can be assigned to the effect at the spatial unit level.

(6) **Lack of spatial specificity**. Whereas the conventional massively univariate approach attempts to determine statistical evidence at each spatial unit, the procedures to handle multiple testing render statistical inference viable only at the level of clusters of activation. Surviving clusters have different spatial extents, and in practice one may observe some that are quite large. As the unit of statistical inference is the entire cluster, the investigator loses the ability to refer to individual voxels, as shown in the common practice of locating a region via its "peak" voxel.

(7) **Penalizing intrinsically small regions**. Compared to the excessive Bonferroni correction, neighborhood-leveraging procedures offer a relatively effective approach at handling multiplicity. However, the conventional massively univariate approach inefficiently models each individual spatial unit separately, and the compensation for the incurred multiplicity cannot fully recover the lost efficiency, resulting in loss of statistical power as the process attempts to achieve nominal FPR levels. The combination of statistical evidence and spatial extent adopted in recent permutation-based methods (Smith and Nichols, 2009; Cox, 2019) provides a principled approach to address the arbitrariness of primary thresholding, but the approach still discriminates against spatially small regions. For example, between two brain regions with comparable statistical strength, the anatomically larger one would be more likely to survive; and between a scenario involving an isolated region and another with two or more contiguous regions, the former may fail to survive the current filtering methods even when locally exhibiting stronger statistical strength.

Modeling frameworks and conceptual differences

Should we consider an effect of a particular entity under investigation (player, county, brain location) to be unknown but fixed if it can be discovered, or to be a varying quantity to be assessed relative to a background of information? In the former case, we are "objective" in the sense of fully trusting the empirical data and proceed by assuming that all potential findings are equiprobable (e.g., activation strength is as likely to be 0.1% as it is to be 100% signal increase). In the latter case, we are "subjective" and assume that particular results are more or less likely with, for example, the assumption of physical symmetry or exchangeability among the regions (e.g., activation strength would follow a Gaussian distribution with its mean and standard deviation inferred later by the model as a data generator). More broadly, it is pertinent to comment on statistical modeling that addresses two categories of uncertainty: epistemic and aleatoric. If an effect is considered intrinsically fixed and simply awaits improved measurement methods to be determined, uncertainty due to current lack of

knowledge (or data) about its effect size is said to be *epistemic*. Once enough information is available, we should be able to discover, estimate, or approximate the effect with shrinking error bars. Paradigmatic examples include measurement errors of constants in physics, such as Newton's g, the speed of light, or Planck's constant. In contrast, uncertainty associated with an effect is said to be *aleatoric* when it is rooted in the intrinsic randomness of the phenomenon itself (in simplistic terms, if an effect sometimes has one value and sometimes has other values, uncertainty is considered to be aleatoric).

In the context of conventional statistical models (t-test, GLM, ANOVA and LME), epstemic quantities are modeled as *fixed effects*, such as when estimating intercept and slopes. The underlying reasoning is that these effects are believed to capture "real" properties of the population (e.g., activation strength, or association between brain activation and behavior), such that absence of information about each fixed effect is temporary, and its uncertainty will be reduced once the effect can be characterized better and estimated through an improved model (and sufficient data). In contrast, aleatoric quantities are modeled as *random effects*. The assumption is that the uncertainty of these quantities reflects the elusive nature of fluctuations among measuring entities (e.g., devices, subject properties). Under the conventional modeling framework, random effects are generally of no interest to the investigator (e.g., residuals in GLM and subject-specific effects under LME) but are included in the model to account for data variability and to allow generalizations from the current sample to a hypothetical population.

Although at first pass the distinction between epistemic and aleatoric variability appears clear-cut, consider the following example concerning an object's length. Initially, we might choose to conceptualize "length" as an intrinsic, fixed characteristic, and our need for multiple measurements to estimate its value comes from epistemic uncertainties such as inaccuracy of the measurement device and human error. However, in practice length also depends on environmental conditions such as temperature, pressure, and humidity; because these affect the underlying value of the quantity itself, such considerations push us into the realm of aleatoric uncertainties. While the conceptual choice will make little difference on outcome in this simple scenario, one should note how consideration of pragmatic considerations render many effects appropriate for aleatoric classification[6].

To recapitulate, the focus of the Bayesian paradigm is not on the inbuilt metaphysical nature of the effect under study; rather, we treat *all* model parameters (including those that are typically coded as fixed effects under the conventional framework) as random variables by expressing our knowledge and uncertainty through probability distributions. Consider the following example relevant in the context of neuroimaging. For the population of all humans, do we expect there to be a fundamentally fixed, underlying BOLD response in the amygdala (say, 0.538% signal increase) when contrasting images of fearful vs. neutral faces? Should we assume that all the factors contributing to uncertainty of measuring this effect in an experiment (e.g., genetic variability, experimental context, mood, alertness, anxiety, etc.) are epistemic? We suggest that it is reasonable to treat the uncertainty associated with any effect, regardless of the investigator's interest, as aleatoric within the scope of the BML framework and determine its magnitude uncertainty based on a (large enough) sample of participants. Indeed, the range of aleatoric uncertainty can be reported as a quantile interval from the posterior distribution (or other forms of statistical summary as in conventional approaches). In other words, instead of forcing a view of brain response as exact, intrinsic values across *all* human beings, we make statistical inferences based on an integrative model that renders a representative distribution of the effect. The Bayesian framework thus helps dissolve the distinctions between the uncertainty types. Model parameters such as intercept and slopes are no longer treated as unknown fixed effects at the population level, but are pragmatically considered to be aleatoric, much like random effects such as subjects under the conventional framework. More broadly, a guiding principle of Bayesian statistics is that the state of knowledge about anything unknown should be expressed by a probability distribution.

A little more technically, the move away from assuming epistemic uncertainties ("fixed but unknown") allows us first to incorporate spatial units as levels of a factor into LME modeling, and then to transition from LME to BML (for technical details, see Chen et al., 2019b). Without operating within the BML framework, the individual spatial units would be typically treated as independent entities (isolated locations) and modeled in parallel under the massively univariate modeling. Consider the example of inferring an effect at each region pair with matrix data. In a traditional LME model each region pair is treated as a random effect, and no inference can be made about individual regions and region pairs other than the intercept (common effect shared by all regions and all subjects), which is typically not very informative in practice. In contrast, within the BML framework, the effect of each region pair is modeled as the contribution of each involved region plus their interaction. In doing so, we can derive both the individual effects *and* their combined contribution as a pair by

---

[6]Indeed, the penalty for considering uncertainties as aleatoric is rather modest: in the limit that non-epistemic contributions vanish, estimate uncertainty simply decreases. In other words, if one can model most contributions to a given variable of interest, all that remains is measurement variability.

sampling from the posterior joint distribution (Chen et al., 2019c). Thus, the distinction between fixed- and random-effects characterized in the conventional LME framework is mapped to, in the BML context, the distinction between information pooling across regions and, separately, across-subject variability.

Historically, much has been discussed about the Bayesian approach in the context of "subjectivity" versus "objectivity." Bayesian methods are frequently linked with the former, and in fact in a negative fashion. As all statistical models are subjective in the sense of idealizing or approximating reality, we favor adopting a pragmatic stance. Certainly, analytic decisions require scientific evaluation, including specific aspects of data processing specifics (amount of spatial smoothing, choice of data included or excluded, model validation, etc.) and uncertainty assignment through a probability distribution. By considering Stein's paradox and related examples, we hope to have motivated the idea that the "objectivity" of effect inferences (i.e., statistical unbiasedness) should not be the sole criterion in adopting a statistical approach. In fact, the Gaussian prior incorporated in the scenarios of the basketball players, counties, and ROIs, is a model assumption much like those in conventional statistical methods (e.g., Gaussian distribution for residuals and subjects). Note that the Gaussian prior only stipulates the distribution shape, and its specific parameters are actually determined a posteriori through the model conditioning on the data. As experimentalists, we generally have prior knowledge about an effect of interest (e.g., a BOLD percent change of 100% is unrealistic for most contrasts). Adopting a weakly informative prior adds a small amount of real-world information to the model – it provides a "nudge". Naturally, the analyst should explicitly report all model assumptions, including the priors adopted. Importantly, the performance of a Bayesian model can be evaluated via graphical tools such as posterior predictive checks (Fig. 8) as well as via cross-validation. Taken together, we believe that the global calibration approach can contribute to improving neuroimaging research in multiple ways, including by reducing thresholding-related arbitrariness, increasing model efficiency, and enhancing reproducibility.

Potential and limitations of BML applicability in neuroimaging

Table 1 compares the conventional massively-univariate approach and the multilevel framework described here.

The investigator's intention concerning the data space has a considerable impact on the extent of statistical evidence in conventional NHST (Kruschke, 2010). For example, if the investigator decides to perform a whole-brain voxel-wise analysis, or to target a few dozen regions of interest, the associated corrections for multiple testing might result in quite different results (from not being able to report a single voxel to reporting noteworthy results across multiple regions). How does BML fare in this respect?

The BML framework tends to be less vulnerable to choices of data space as long as the prior distribution is weakly informative (which is the recommendation under typical scenarios). In most cases, the largest impact of information sharing (partial pooling) will be observed only in the most extreme estimates. For example, in the case of the top basketball players, choosing a considerably broader Gaussian prior would reduce shrinkage, that is, the extent to which the smallest and largest estimates are pulled toward the mean. Now, suppose we changed our "intentions" and decided to analyze the top 100 players instead of top 50 (perhaps a reviewer of the study found the number 50 somewhat arbitrary and suggested 100 as less arbitrary since it is a perfect square). The impact on the original estimates would be fairly small as long as the added players' performance did not substantially deviate from the original distribution assumption (e.g., Gaussian). Provided that the investigator's objective is the overall predictive accuracy, partial pooling is expected to produce similar results (Fig. 2D illustrates this behavior for the case of pooling among the top 50 NBA players versus the top 100 players). Accordingly, in a neuroimaging study, if the set of spatial units changes because of evolving research focus (e.g., adding or subtracting regions), the impact would typically be small; again, as long as the Gaussian assumption still reasonably holds.

We also wish to emphasize the importance of reporting the continuous spectrum of statistical evidence. In the applications described in previous sections, effect uncertainty was quantified via quantile intervals of the posterior distribution, which provide a straightforward and intuitive representation of variability. This approach works well when the number of estimate involved is relatively small (say, 10-20) but is more challenging for larger datasets. Given that neuroimaging data analysis typically involves larger data structures, one may choose to highlight a subset of the effects, particularly those with "strong" or "moderate" evidence. However, these convenient linguistic labels should be operationally defined and not viewed as carrying categorical information ("active" vs. "not active").

The BML models described here were defined at the level of brain regions. Could the approach be extended and be applied to a whole-brain voxel-wise level? At present the answer is "no" given the computational cost of fitting very large multilevel models. However, this type of analysis may be feasible in the not distant future with the use of Graphical Processing Units and within-chain parallelization (Stan Development Team, 2019). Nevertheless, there are a few disadvan-

Table 1: Comparisons of assumptions and properties of massively univariate analysis and Bayesian multilevel modeling

	Massively Univariate Analysis	Bayesian Multilevel Modeling
Number of models	number of units plus correction for multiple comparisons	one
Sharing of information	each unit is independent	units are exchangeable and loosely regularized
Focus of error control	overall type I (i.e., FPR)	type S (sign) and type M (magnitude)
Strategy for multiplicity	FPR correction (control for inflated statistical evidence)	partial pooling (control for inflated effect sizes)
Effects uncertainty	epistemic (effect is intrinsic and fixed, uncertainty from measurement error, etc.)	aleatoric (effect has inherent variability)
Effect inferences	locally unbiased, no calibration involved because of implicit assumption of uniform distribution	locally biased, globally calibrated through a regularizing (Gaussian) distribution
Framing of hypotheses	$P(\text{data} \mid H_0)$: test "surprise" of having the observed data under the null hypothesis H_0 scenario	$P(H_R \mid \text{data})$: test evidence for research hypothesis H_R given the observed data
Inference method	perform NHST with a binary decision based on an FPR-adjusted threshold	assess statistical evidence $P(H_R \mid \text{data})$ through posterior distribution: highlight but no hide
Model efficiency	local (e.g., unbiasedness of each unit, statistical power)	global (cross-validation, posterior predictive checks)

tages associated with whole-brain voxel-wise analysis that could be complemented through a region-based approach. For instance, analysis at the voxel level is susceptible to the quality of inter-subject alignment, an increasing challenge as the field keeps pushing for smaller and smaller spatial units (e.g., less than 1 mm isotropic voxels). In addition, inferences based on region-based analysis are region-specific; in contrast, statistical evidence at the voxel level is not necessarily well-aligned with the anatomical or functional definitions, leading to ambiguous or difficult inferences.

A potential concern about the BML approach is the underlying assumption about the relationships among the spatial units. Although independence is not required, exchangeability[7] is needed to guarantee that no differential information among spatial units is available (it also allows approximating the prior distribution across regions in terms of a mixture of identical and independent distributions per de Finetti's theorem). The assumption of exchangeability is also advantageous because a prior distribution with fewer adjustable parameters will be associated with "sharper" posterior densities (i.e., one with a more discernible peak) than more complex prior distributions (Jefferys and Berger, 1992). Nevertheless, it is to be expected that some regions in the brain will share more information between them (relative to other regions), such as anatomically adjacent or interhemispheric (i.e., homotopic) regions. Therefore, a fruitful direction of research will be to relax the assumption of exchangeability so as to capture potential forms of information sharing between groups of regions. In particular, the multilevel correlation structure among spatial units may be informed by the functional organization of brain regions, such as a gradient between sensorimotor and transmodal areas (Huntenburg et al., 2018).

Conclusions

Neuroimaging investigators confront the issue of multiplicity front and center. The analysis challenges posed by this problem are inherently linked to the conventional strategy of modeling the data in a univariate manner: Each spatial unit (e.g., voxel, region) is modelled independently. In the present paper, we described how this approach leads to modeling inefficiency and information loss given the required procedures for handling multiplicity. As an alternative framework, we presented a Bayesian multilevel modeling framework that makes inferences across spatial units by pooling information across them and achieving better overall predictive accuracy. Instead of a multiplicity of models (one for each spatial unit), a single integrative model is employed to embrace multiplicity to modeling advantage. It is our hope that the convergence of theoretical progress combined with algorithm development will stimulate the growth of Bayesian modeling in neuroimaging. Bayesian models also have the potential to encourage reporting experimental findings more fully, instead of dichotomizing them into "significant" and "non-significant" bins, thus reducing information loss in the literature while enhancing both transparency and reproducibility.

Acknowledgments

The research and writing of the paper were supported (GC, PAT, and RWC) by the NIMH and NINDS Intramural Research Programs (ZICMH002888) of the NIH/HHS, USA. LP's research is supported in part by the National Institute

---

[7]Exchangeability assumes that no differential information is available across the measuring entities (e.g., regions and subjects in neuroimaging) in the model. A set of entities is exchangeable if its joint probability distribution is a symmetric function of the entities; thus, if the order or sequence of the entities is permuted, the joint distribution would not change. Exchangeability captures the distributional symmetry among the entities in a sense that does not require independence: independent and identically distributed set of entities is exchangeable, but not vice versa.

References

Amrhein, V., Greenland, S., McShane, B., 2019. Scientists rise up against statistical significance. Nature 567, 305-307.

Baggio, H.C., Abos, A., Segura, B., Campabadal, A., Garcia-Diaz, A., Uribe, C., Compta, Y., Marti, M.J., Valldeoriola, F., Junque, C., 2018. Statistical inference in brain graphs using threshold-free network-based statistics. Human Brain Mapping 39(6).

Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B 57, 289-300.

Bennett, C.M., Baird, A.A., Miller, M.B., Wolford, G.L., 2010. Neural correlates of interspecies perspective taking in the post-mortem Atlantic Salmon: An argument for multiple comparisons correction. Journal of Serendipitous and Unexpected Results.

Calhoun, V., Adali, T., Pearlson, G., Pekar, J., 2001. A Method for Making Group Inferences From Functional MRI Data Using Independent Component Analysis. Hum.Brain Map. 14:140-151.

Chen, G., Cox, R.W., Glen, D.R., Rajendra, J.K., Reynolds, R.C., Taylor, P.A., 2019a. A tail of two sides: Artificially doubled false positive rates in neuroimaging due to the sidedness choice with t-tests. Human Brain Mapping 40(3):1037-1043.

Chen, G., Taylor, P.A., Cox, R.W., 2017. Is the statistic value all we should care about in neuroimaging? Neuroimage 147:952-959.

Chen, G., Cox, R.W., Glen, D.R., Rajendra, J.K., Reynolds, R.C., Taylor, P.A., 2019a. A tail of two sides: Artificially doubled false positive rates in neuroimaging due to the sidedness choice with t-tests. Human Brain Mapping 40(3):1037-1043.

Chen G, Xiao Y, Taylor PA, Riggins T, Geng F, Redcay E, 2019b. Handling Multiplicity in Neuroimaging through Bayesian Lenses with Multilevel Modeling. Neuroinformatics (in press). https://rdcu.be/bhhJp

Chen, G., Bürkner, P.-C., Taylor, P.A., Li, Z., Yin, L., Glen, D.R., Kinnison, J., Cox, R.W., Pessoa, L., 2019c. An Integrative Approach to Matrix-Based Analyses in Neuroimaging. Human Brain Mapping (in press). https://doi.org/10.1101/459545

Chen, G., Taylor, P.A., Qu, X., Molfese, P.J., Bandettini, P.A., V, R.W., Finn, E.S., 2019d. Untangling the Relatedness among Correlations, Part III: Extending Model Capabilities of Inter-Subject Correlation Analysis for Naturalistic Scanning. Under review.

Choi, J. M., Padmala, S., Pessoa, L., 2012. Impact of state anxiety on the interaction between threat monitoring and cognition. Neuroimage 59(2) 1912-1923.

Cox, R.W., Chen, G., Glen, D.R., Reynolds, R.C., Taylor, P.A., 2017. FMRI Clustering in AFNI: False-Positive Rates Redux. Brain Connect. 7(3):152-171.

Cox, R.W., 2019. Equitable Thresholding and Clustering. Brain Connectivity. In press.

Cremers, H.R., Wager, T.D., Yarkoni, T., 2017. The Relation Between Statistical Power and Inference in fMRI. PLoS ONE 12(11):e0184923

Forman, S.D., Cohen, J.D., Fitzgerald, M., Eddy, W.F., Mintun, M.A., Noll, D.C., 1995. Improved assessment of significant activation in functional magnetic resonance imaging (fMRI): use of a cluster-size threshold. Magn Reson Med. 33:636-647.

Gelman, A., Carlin, J.B., Stern, H.S., Dunson, D.B., Vehtari, A., Rubin, D.B., 2014. Bayesian data analysis. Third edition. Chapman & Hall/CRC Press.

Gelman, A., Tuerlinckx, F., 2000. Type S error rates for classical and Bayesian single and multiple comparison procedures. Computational Statistics 15: 373-390.

Glass, G.V., 1976. Primary, secondary, and meta-analysis of research. Educational Researcher. 5(10): 3-8.

Gonzalez-Castillo, J., Saad, Z.S., Handwerker, D.A., Inati, S.J., Brenowitz, N., Bandettini, P.A., 2012. Whole-brain, time-locked activation with simple tasks revealed using massive averaging and model-free analysis. PNAS 109(14):5487-5492.

Huntenburg, J.M., Bazin, P.L., Margulies, D.S., 2018. Large-scale gradients in human cortical organization. Trends in cognitive sciences, 22(1):21-31.

James, W., Stein, C., 1961. Estimation with quadratic loss. Proc. Fourth Berkeley Symposium 1, 361-380.

Jeffreys, W.H., Berger, J.O., 1992. Sharpening Ockham's Razor on a Bayesian strop. American Scientist 80.

Jenkinson, M., Beckmann, C.F., Behrens, T.E., Woolrich, M.W., Smith, S.M., 2012. FSL. NeuroImage, 62:782-90.

Kriegeskorte, N., Simmons, W.K., Bellgowan, P.S.F., Baker, C.I., 2009. Circular analysis in systems neuroscience: the dangers of double dipping. Nat. Neurosci. 12:535–540.

Kinnison, J., Padmala, S., Choi, J.M., Pessoa, L., 2012. Network analysis reveals increased integration during emotional and motivational processing. J Neurosci. 32(24):8361-8372.

Kruschke, J.K., 2010. Bayesian data analysis. Wiley Interdisciplinary Reviews: Cognitive Science 1(5):658-676.

McElreath R., 2016, Statistical Rethinking: A Bayesian Course with Examples in R and Stan. Chapman & Hall/CRC Press. https://www.overleaf.com/project/5cb0b178e51a152e6d1cefc1

McShane, B.B., Gal, D., Gelman, A., Robert, C., Tackett, J.L., 2019. Abandon Statistical Significance.

Nichols, T.E., Holmes, A.P., 2001. Nonparametric permutation tests for functional neuroimaging: a primer with examples. Hum Brain Mapp. 15(1):1-25.

Pinheiro, J.C., Bates D.M., 2000. Mixed-Effect Models in S and S-plus. Springer-Verlag, New York.

R Core Team, 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Santosa, F., Symes, W.W., 1986. Linear inversion of band-limited reflection seismograms. SIAM Journal on Scientific and Statistical Computing. SIAM. 7(4):1307–1330.

Smith, S.M., Nichols, T.E., 2009. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. Neuroimage. 44(1):83-98.

Stan Development Team, 2019. Stan Modeling Language Users Guide and Reference Manual, Version 2.19.0. http://mc-stan.org

Stein, C., 1956. Inadmissibility of the usual estimator for the mean of a multivariate distribution. Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability 1: 197-206.

Stigler, S.M., 1990, A Galtonian perspective on shrinkage estimators. Statistical Science 5(1):147-155.

Tikhonov, A.N., Leonov, A.S., Yagola, A.G., 1998. Nonlinear ill-posed problems. London: Chapman & Hall.

Vehtari, A., Gelman, A., and Gabry, J., 2017. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. Statistics and Computing. 27(5):1413-1432.

Whitfield-Gabrieli, S., and Nieto-Castanon, A., 2012. Conn: A functional connectivity toolbox for correlated and anticorrelated brain networks. Brain Connectivity. doi:10.1089/brain.2012.0073

Worsley, K.J., Marrett, S., Neelin, P., Evans, A.C., 1992. A three-dimensional statistical analysis for CBF activation studies in human brain. Journal of Cerebral Blood Flow and Metabolism, 12:900-918.

Xiao, Y., Geng, F., Riggins, T., Chen, G., Redcay, E., 2019. Neural correlates of developing theory of mind competence in early childhood. NeuroImage 184:707-716.

Zalesky, A., Fornito, A., Bullmore, E.T., 2010. Network-based statistic: identifying differences in brain networks. NeuroImage 53:1197-1207.