Interaction between induced and natural variation at *oil yellow1* delays flowering in maize

Rajdeep S. Khangura[†§‡], Bala P. Venkata[†2], Sandeep R. Marla[†1], Michael V. Mickelbart[†‡], Singha

Dhungana[3], David M. Braun[3], Brian P. Dilkes[§‡], Gurmukh S. Johal[†‡]

[†]Department of Botany and Plant Pathology, Purdue University, West Lafayette, Indiana 47907

[‡]Center for Plant Biology, Purdue University, West Lafayette, Indiana 47907

[§]Department of Biochemistry, Purdue University, West Lafayette, Indiana 47907

[3]Division of Biological Sciences, Interdisciplinary Plant Group, Missouri Maize Center,

University of Missouri, Columbia, Missouri 65211

Current address

[1]Department of Agronomy, Kansas State University, Manhattan, Kansas 66506

[2]Donald Danforth Plant Science Center, St. Louis, Missouri 63132

Corresponding Authors

Brian P. Dilkes, 170 S. University Street, Purdue University, West Lafayette, Indiana 47907,

Phone: (765) 494 2584, Email: bdilkes@purdue.edu

Gurmukh S. Johal, 915 W. State Street, Purdue University, West Lafayette, Indiana 47907, Phone:

(765) 494-4448, Email: gjohal@purdue.edu

Abstract

We previously demonstrated that maize (*Zea mays*) locus *very oil yellow1 (vey1)* encodes a

putative cis-regulatory expression polymorphism at the magnesium chelatase subunit I gene (aka

oil yellow1) that strongly modifies the chlorophyll content of the dominant *Oy1-N1989* mutants.

The *vey1* allele of Mo17 inbred line reduces chlorophyll content in the mutants leading to reduced

photosynthetic output. *Oy1-N1989* mutants in B73 flowered four days later than wild-type.

Enhancement of *Oy1-N1989* by the Mo17 allele at the *vey1* QTL delayed flowering further,

resulting in detection of a flowering time QTL in two bi-parental mapping populations crossed to

Oy1-N1989. The near isogenic lines of B73 harboring the *vey1* allele from Mo17 delayed flowering

of *Oy1-N1989* mutants by twelve days. Just as previously observed for chlorophyll content, *vey1*

had no effect on flowering time in the absence of the *Oy1-N1989* allele. Loss of chlorophyll

biosynthesis in *Oy1-N1989* mutants and enhancement by *vey1* reduced CO_2 assimilation. We

attempted to separate the effects of photosynthesis on the induction of flowering from a possible

impact of chlorophyll metabolites and retrograde signaling by manually reducing leaf area.

Removal of leaves, independent of the *Oy1-N1989* mutant, delayed flowering but surprisingly

reduced chlorophyll contents of emerging leaves. Thus, defoliation did not completely separate

the identity of the signal(s) that regulates flowering time from changes in chlorophyll content in

the foliage. These findings illustrate the necessity to explore the linkage between metabolism and

the mechanisms that connect it to flowering time regulation.

KEYWORDS flowering time; photosynthesis; epistasis, setaria; sorghum

Introduction

As sexual reproduction is necessary for the production of seeds and fruits, the onset of flowering

in angiosperms has been a key focus for plant biologists working on ornamental, horticultural, and

other crop species (Lang 1952; Zeevaart 1962; Searle 1965). The onset of reproductive

development in angiosperms is affected by a change in meristem identity. The vegetative to floral

transition of meristems commits plant development to production of floral organs and sexual

reproduction. The integration of signals to correctly time this transition is key to plant fitness.

Unsurprisingly, endogenous and environmental cues regulate flowering time (Amasino and

Michaels 2010; Cho *et al.* 2017; Minow *et al.* 2018). A critical environmental cue is the duration

of the light period, or photoperiod. The photoperiodic responses of plants influence the vegetative

to floral transition and the mechanisms of this response have been a focus of intensive research for

over a century (Klebs, 1918). Multiple non-photoperiodic cues as well as endogenous signals,

sometimes called the autonomous pathway, are also critical to floral transition. Endogenous signals

including hormones and the carbohydrate status of the plant can also play a critical role in the

regulation of flowering time (Corbesier *et al.* 1998; Moghaddam and Ende 2013). But

environmental influences, such as light quality, alter hormone biosynthesis (Lang 1957; Evans and

Poethig 1995; Mutasa-Göttgens and Hedden 2009), and light intensity dramatically influences

photosynthesis and thereby carbohydrate status (Chen *et al.* 2004). These stimuli converge through

the same floral integrators (named FLOWERING LOCUS T (FT) and FLOWERING LOCUS D

(FD) in *Arabidopsis thaliana)* for which orthologs have been identified in many flowering plants

(Abe *et al.* 2005; Wigge *et al.* 2005; Corbesier *et al.* 2007; Meng *et al.* 2011; Zhang *et al.* 2016).

Accumulation of FT and FD gene products trigger the vegetative shoot apical meristems to acquire

the competency to become inflorescence meristems and produce flowers in part via the activation

of MADS-box transcription factors that control meristem identity through APETALA1 (Abe *et al.*

2005; Wigge *et al.* 2005). In Arabidopsis, FT is regulated by CONSTANS (CO) in response to

both circadian regulation and photoperiodic responses. CONSTANS itself is regulated by the

MADS-box transcription factor SUPPRESSOR OF CONSTANS1 (SOC1) (Samach *et al.* 2000).

Maize was domesticated from teosinte (*Zea mays* ssp. *parviglumis* or spp. *mexicana*) in

Central America (Doebley *et al.* 1997; Wang *et al.* 1999). Strong selection on time to flower

contributed to the adaptation of maize to different latitudes (Salvi *et al.* 2007; Huang *et al.* 2017;

Swarts *et al.* 2017). Flowering of teosinte is promoted by short-day conditions. In contrast,

temperate maize germplasm is relatively day-neutral and flowering is primarily under the control

of the autonomous pathway (Coles *et al.* 2010). Several mutant and QTL studies have identified

loci affecting flowering time in maize (Buckler *et al.* 2009; Coles *et al.* 2010; Steinhoff *et al.* 2012;

Bouchet *et al.* 2017). Some of these loci include: *indeterminate1* (*id1*; Colasanti et al. 1998); *early*

phase change (*epc*; Vega et al. 2002); the maize FLORICAULA/LEAFY orthologs *zfl1* and *zfl2*

(Bomblies et al. 2003); *delayed flowering1* (*dlf1*; Muszynski et al. 2006); the cis-element

polymorphism *vegetative transitioning1* (*vgt1*; Salvi et al. 2007) that regulates a downstream

APETALA2-like transcription factor *zmrap2.7*; *zea mays mads4* (*zmm4*; Danilevskaya et al. 2008);

zmcct10 (Hung *et al.* 2012); *zea mays centroradiales8* (*zcn8*; Meng et al. 2011); *zea mays mads1*

(*zmmads1*; Alter et al. 2016); and *zea mays mads69* (Liang *et al.* 2019). Many of these encode the

maize orthologs of genes identified as regulators of flowering in Arabidopsis. For example, *dlf1*

and *zcn8* encode homologs of the Arabidopsis flowering time determinants FD and FT,

respectively. *id1* encodes a zinc-finger transcription factor acting upstream of both DLF1 and

ZCN8 (Kozaki *et al.* 2004; Muszynski *et al.* 2006; Meng *et al.* 2011). *zmm4* is an activator of

flowering that is part of a conserved syntenic pair of MADS box genes in the grasses, with *zmm24*

as the neighboring gene, and encodes one of two maize paralogs of the wheat flowering time and

vernalization response locus VRN1 (Danilevskaya *et al.* 2008). *zmm4* acts downstream of *dlf1* and

id1 in the control of flowering time in maize. *zmmads1* is a functional homolog of the Arabidopsis

flowering time and circadian rhythm regulator *soc1* (Alter *et al.* 2016).

One important endogenous signal that contributes to flowering time is the carbohydrate

allocation status (Ohto *et al.* 2001; Seo *et al.* 2011; Eveland and Jackson 2012; Wahl *et al.* 2013;

Cho *et al.* 2018). In maize, mutants that are compromised in either sugar export from source tissues

or loading sucrose into the phloem flower later than their congenic wild-type siblings (Braun *et al.*

2006; Baker and Braun 2008; Ma *et al.* 2008; Slewinski *et al.* 2009; Slewinski and Braun 2010).

This is not limited to maize, as starch-deficient Arabidopsis mutants exhibit delayed flowering

(Corbesier *et al.* 1998). Trehalose-6-phosphate (T6P) has been implicated as a reporter of the

energy status and Arabidopsis mutants effected in this metabolite also exhibit altered flowering

time (Paul 2008; Wahl et al. 2013; Seo et al. 2011). T6P and sucrose are positively correlated, and

low levels of T6P results in delayed flowering in Arabidopsis (Wahl *et al.* 2013). Remarkably, the

carbohydrate status and transcriptional regulatory genes controlling flowering time may be directly

linked in maize. The *id1* flowering time mutants alter carbohydrate partitioning in leaves and

accumulate more sucrose and starch (Coneva et al. 2012). As a result, maize flowering time gene

ID1 has been proposed to act as a carbohydrate status sensor that influences flowering time

(Coneva, Zhu, and Colasanti 2007; Minow et al. 2018). Remarkably, the promoter of the T6P

biosynthetic gene TPS1 is a predicted target of the ID1 DNA-binding protein and low levels of

T6P were observed in *id1* mutants (Minow *et al.* 2018).

If sugars are critical for floral transitioning in plants, then manipulation of photosynthesis

should alter flowering. Magnesium chelatase (MgChl) is a hetero-oligomeric enzyme complex that

catalyzes the first committed step of chlorophyll biosynthesis by conversion of protoporphyrin IX

(PPIX) into magnesium-PPIX (Walker and Weinstein 1991; Gibson et al. 1995). We previously

identified a cis-acting expression polymorphism at the *oil yellow1* (*oy1*) locus (Sawers *et al.* 2006)

associated with a QTL called *very oil yellow1* (*vey1*) that modifies the chlorophyll content of semi-

dominant *Oy1-N1989* mutants (Khangura *et al.* 2019). The effect of *vey1* is only visible in the

presence of *Oy1-N1989,* indicating that this natural variant has a cryptic effect on the function of

the MgChl complex. We used controlled crosses to segregate *Oy1-N1989* and the modifier alleles

at *vey1* to generate populations of maize with a range of chlorophyll contents. We used this

variation in chlorophyll to explore the effects of chlorophyll content on net CO_2 assimilation.

These changes in chlorophyll accumulation resulted in changes in net CO_2 assimilation and

photosynthetically-fixed carbon accumulation. Remarkably, we noticed that flowering time across

material with differing photosynthetic rates and chlorophyll contents was dramatically different.

We observed that *Oy1-N1989* mutants flowered later and that the reduced chlorophyll

accumulation was associated with a delay in flowering time. Similar to the cryptic effects of *vey1*

on chlorophyll content, the modifier allele had no effect on flowering time in the absence of the

Oy1-N1989 mutant allele. Chlorophyll content was consistently associated with earlier flowering

and partial rescue of chlorophyll accumulation in the *Oy1-N1989* mutant by the *vey1* QTL

accelerated flowering in the mutants but had no effect on wild-type siblings. In addition to

measurements of net CO_2 assimilation, the premature senescence of maize leaves induced by sugar

accumulation, following sink removal, was also reduced by *Oy1-N1989* and further reduced by

vey1 alleles that decrease chlorophyll content and net CO2 assimilation. The effect of reduced

photosynthate accumulation on flowering time was not specific to *Oy1-N1989* as mechanical

removal of leaves, to reduce plant leaf area, also delayed flowering time. Thus, all of our results

are consistent with an integrative measure of carbon assimilation linking energy status and

flowering time in maize.

Materials and Methods

Plant materials

The origin of *Oy1-N1989* mutant allele has been previously described (Khangura *et al.* 2019). The

mutant allele is maintained in B73 background and is propagated by crossing heterozygous

mutants (*Oy1-N1989/oy1*) to the wild-type siblings. The introgressed stock of *Oy1-N1989* into

B73 was used for crossing to various mapping populations. A total of 216 intermated B73 × Mo17

population (IBM-RILs), and 251 synthetic 10 doubled haploid lines (Syn10-DH) were crossed as

ear-parents with *Oy1-N1989/oy1*:B73 pollen. The pollen of *Oy1-N1989/oy1*:B73 plants were also

crossed on to the ears of 35 B73-Mo17 near-isogenic lines (BM-NILs) for QTL validation. Tables

S1-S3 contain the full list of IBM-RILs, Syn10-DH, and BM-NILs used to develop F_1 hybrid

populations.

Field trials

The field experiments were conducted at the Purdue Agronomy Center for Research and Education

(ACRE) in West Lafayette, Indiana. Each cross was evaluated as a single plot of 12-16 plants.

Each plot derived from crosses with *Oy1-N1989/oy1*:B73 tester segregated for both mutant and

wild-type siblings in approximately 1:1 ratio. Seeds were sown with tractor-driven seed planter

with plot length of 3.84 meters (m) with alley length of 0.79 m, and inter-row spacing of 0.79 m.

Rainfall was sufficient throughout the growing season and eliminated any need of irrigation.

Standard crop management practices at Purdue in terms of fertilizer, pest, and weed control for

growing field maize were adopted.

Each experiment was divided into blocks. Progenies of crosses of B73 and Mo17 ears to

Oy1-N1989/oy1:B73 pollen were used as parental checks in each block. Parental checks were

randomized with each block. The IBM-RILs F_1 populations were evaluated as a single replication

in 2013. Syn10-DH F_1 populations were evaluated in 2016 with two randomized replications

planted in randomized complete block design (RCBD). Hybrid progenies of *Oy1-N1989/oy1*:B73

with BM-NILs were screened in 2016 with five replications planted in RCBD.

Field phenotyping and data collection

The mutant plants in each plot were identified visually as pale plants. The chlorophyll content in

mutant and wild-type plants was approximated using a CCM-200 plus (Opti-Sciences Inc., Hudson,

NH) as described in Khangura *et al.* 2019. We previously demonstrated correlation of 0.94

between CCM-200 plus values and chlorophyll *a*, chlorophyll *b*, and total chlorophyll contents

(Khangura *et al.* 2019), and therefore used CCM-200 plus values (CCM) as a proxy for chlorophyll

content in the materials described here. CCMI refers to measurements taken with CCM-200 plus

instrument from 25-30 days after sowing and CCMII refers to measurements taken 45-50 days

after sowing. Mutants were tagged around V5-V7 stages of development. Tagging is necessary as

suppression of the *Oy1-N1989* mutant phenotype by *vey1* interferes with visual classification of

mutant and wild-type siblings at maturity. Flowering time in each F_1 population was recorded

separately on the wild-type and mutant plants. The date at which roughly half of the wild-type or

mutant plants in a given plot were shedding pollen was recorded as the date of anthesis for a given

genotype in that plot. Similarly, the date at which half of the wild-type or mutant plants had

emerging silks was recorded as the date of silking for the respective genotype in that plot. The

dates of anthesis and silking for both wild-type and mutant genotypes were then subtracted from

the date of planting to obtain respective wild-type or mutant days to anthesis (DTA), and days to

silking (DTS). Further, the difference between DTA and DTS was used to derive the anthesis-

silking interval (ASI); ASI = DTA-DTS. Wild-type and mutant traits are denoted with a prefix WT

and MT, respectively. Ratio and differences of these flowering times values were also calculated

as MT/WT and WT-MT, respectively.

Induced leaf senescence assay: A total of 15 F_1 populations derived from B73-like NILs ×

Oy1-N1989/oy1:B73 cross were used for this experiment. Seven of these B73-like NILs carried an

introgression of *vey1* from Mo17 (*vey1^{Mo17}*), whereas the other eight NILs had the B73 genotype

at *vey1* (*vey1^{B73}*). These NIL populations were planted in the field with at least two replications of

each genotype planted in RCBD. The procedure for this experiment was adapted from Sekhon *et*

al. 2012. Briefly, for the suppressing NIL, F_1 crosses of primary and secondary ears of both wild-

type and mutant plants were covered with shoot bags before silk emergence. After 3-4 days of

tassel shedding, roughly half of the shoot bags were removed, and these ears were allowed to open

pollinate. Staggered rows of B73 were planted on both sides of the experimental row in the field

to ensure pollination of exposed ears. The day of shoot exposure was marked as 0 days after

anthesis (DAA). For the enhancing B73 NILs, a late set of suppressed NIL F_1 was used as a control.

A similar strategy of shoot bagging of ears before silk emergence and ear exposure after 3-4 days

of tassel shedding was done. The plants that showed tassel maturity on the same day were tagged

and subsequently used for observation.

Genotypic and gene expression data

The genotypic data and other public datasets on various mapping populations used in this

study have been described previously (Khangura *et al.* 2019). Briefly, the public marker dataset

for IBM-RILs was obtained from MaizeGDB with 2,178 markers (Sen *et al.* 2010). The markers

were reduced to 2,156 after removing duplicate variants, with ~13.3 percent of missing data in the

final dataset. Expression data of *oy1* locus in IBM-RILs were obtained from a public repository of

the National Science Foundation grant (GEPR: Genomic Analyses of shoot meristem function in

maize;   NSF   DBI-0820610;

https://ftp.maizegdb.org/MaizeGDB/FTP/shoot_apical_meristem_data_scanlon_lab/). This data

consists of the expression of maize genes in the tissue derived from the shoot apex of 14 days old

IBM-RILs seedlings. The expression data from each gene is normalized to reads per kilobase of

transcript per million mapped reads (RPKM). Genotypic data consisting of 6611 SNPs for Syn10-

DH lines was obtained from Liu et al. 2015. This dataset had no missing genotypes and was used

as such for QTL analyses. Genotypes of the B73-Mo17 Near Isogenic Lines (BM-NILs) used for

QTL validation were obtained from Eichten *et al.* 2011.

Molecular analyses

OY1 sequencing: The genomic DNA and cDNA preparation derived from second and third fully-

expanded leaf of maize inbred lines were used to amplify the genic sequence of the *oy1* locus. The

detail description of methods and primers used for PCR amplification and sequence analyses of

the PCR products has been described previously (Khangura *et al.* 2019). Briefly, the PCR

amplicons were sequenced using Illumina and Sanger sequencing platforms at Purdue Genomic

Core Facility.

Allele-specific expression (ASE) assay: Three replications of B73-Mo17 near-isogenic lines

(BM-NILs) × *Oy1-N1989/oy1*:B73 F_1 progenies were grown in the field at Purdue Agronomy

Farm along with parental F_1 checks. Mutant siblings derived from four B73-like NIL crossed with

Oy1-N1989/oy1:B73 were selected for the ASE experiment. These NILs consisted of two B73-

like NILs with $vey1^{Mo17}$, and other two B73-NILs with $vey1^{B73}$ genotype. Leaf tissue was harvested

from the top fully-expanded leaf at the V3 developmental stage from the mutant siblings of the

four B73-like NIL F_1 plots. For each biological replicate, tissue was pooled from 4-5 seedlings to

make one sample. The samples were stored at -80 ˚C until needed. The procedure of total RNA

isolation, cDNA synthesis, and the ASE assay has been described previously in detail (Khangura

et al. 2019). Briefly, total RNA was isolated using a modified Phenol/Lithium chloride protocol

(Eggermont et. al. 1996). One µg of DNase treated RNA was used to synthesize cDNA using

iScriptTM Select cDNA synthesis kit (catalog#170-8896, Bio-Rad, Hercules, CA) according to

the manufacturer's recommendations. The sensitivity of the ASE assay was tested using the

genomic DNA control that was prepared in the ratio of 1:1 (derived from B73 × Mo17 F_1 hybrid),

1:2, and 2:1 from B73 and Mo17 DNA. PCR was conducted using the forward oligonucleotide 5'-

TCACCGTCTGCAATGTCGCCGCTC   -3'   and   reverse   oligonucleotide   5'-

AGTATGCCCCTGTTGGCCTTGGCG -3' under following reaction conditions with 30 cycles of

polymerization (94˚C for 30s, 56˚C for 30s, 72 ˚C for 30s and final extension for 2 minutes) to

amplify the targeted region of OY1. The primer pair used in this assay flanked the SNP that causes

the L176F amino acid change that resulted in the *Oy1-N1989* mutant allele. PCR products were

sequenced on a MiSeq instrument (Illumina, San Diego, CA) at the Purdue Genomics Core Facility.

Allele count of the reads mapped to small reference sequence derived from B73 was done using

the GATK packages (DePristo *et al.* 2011). Read counts derived from GATK was used to calculate

allele-specific expression. As expected, control samples derived from genomic DNA displayed

bias in the read counts in a DNA dosage-dependent manner. Genomic DNA derived from B73 ×

Mo17 F_1 hybrids resulted in roughly 1:1 read counts demonstrating no bias in the assay.

Statistical analyses

The exploratory data analysis was done using JMP 13.0 (SAS Institute Inc. 2016). Averages of

various traits from line-cross populations, IBM-RILs and Syn10-DH, were subjected to QTL

analyses. QTL detection was done using a single interval mapping function "scanone" in R/qtl, a

software package implemented in R (Broman *et al.* 2003; R core Team 2013). Remarkably similar

results were obtained with composite interval mapping function "cim" in R/qtl (data not shown).

All the QTLs reported in this study are derived from "scanone" function.

Defoliation assay

The defoliation experiments were performed using maize inbred B73, sorghum (*Sorghum bicolor*)

inbred BTx623, and green foxtail (*Setaria viridis*) inbred A10.1. These experiments were

conducted under the greenhouse conditions using mogul base high-pressure sodium lamps (1000

Watts) as the supplemental light source for L:D cycle of 16:8 hours, with the temperature around

28ºC (day-time) and 20ºC (night-time). The maize inbred line B73 was defoliated at V3 leaf stage.

All the leaves were removed from the base, slightly above the ligule. Sorghum inbred BTx623

plants with three to four fully opened leaves (~20 days after sowing in the greenhouse) were

defoliated in a similar way. All fully expanded leaves at ~15 days after planting were removed in

Setaria plants. In addition, all tillers were also defoliated similar to the main stem in *Setaria*. The

flowering time was recorded on maize as described above. For sorghum and *Setaria*, the date of

head emergence on every plant was recorded, and it was deducted from the date of planting to

obtain days to heading.

Non-structural carbohydrate (NSC) quantification

The soluble sugars and starch were quantified from the mutant siblings of four B73-NILs × *Oy1-*

N1989/oy1:B73 F$_1$ population. Two of these NILs carried *vey1^{Mo17},* while the other two had *vey1^{B73}*

genotype. Plants were grown in the field with three replications using RCBD. Top fully-expanded

leaf at the V3 stage was harvested at 1:00 PM and transferred to liquid nitrogen. Leaf tissues were

stored at -80°C until needed. Leaf samples were ground into a fine powder and ~100 mg of the

powder was used to extract sucrose, glucose, fructose, and starch using a previously described

method (Leach and Braun 2016). The quantification of these NSC was done using a method as

described (Leach *et al.* 2017). Briefly, high-performance anion exchange (HPAE) chromatography

(ICS-5000, Thermo-Fisher Scientific) was used to analyze the neutral fraction of the extract. The

HPAE chromatography equipped with a SA10 column was maintained at 45°C. An isocratic

potassium hydroxide solution (10 mmol/L) at a constant flow rate of 1.5mL/min^{-1} was used as an

eluent. Quadrupole wave form as per the manufacturer's recommendation was used for detection

with a disposable gold electrode maintained at 25°C. Sugar standards were used to construct a

standard curve, and samples were diluted to ensure that the detected values fell within the scope

of this curve.

Gas-exchange measurements

Gas-exchange measurements were taken on the field-grown plants during the summer of 2017 at

the Purdue ACRE farm. The gas-exchange measurements were taken on the third leaf on plants at

the V3 stage between 11 AM and 1 PM using LICOR LI-6400XT open photosynthesis system (LI-

COR Inc., Lincoln, NE, USA). Four B73-NILs × *Oy1-N1989/oy1*:B73, two with *vey1^{Mo17}*, were

used for these measurements. Plants were grown in a randomized complete block design of three

replicated blocks. For each genotype nine plants were measured as three replicates in each of the

three blocks. The following instrument conditions were maintained throughout the measurement

period: an artificial light source with an intensity of 1700 μmol photosynthetically active radiation

(PAR) m^{-2} s^{-1}, air temperature of ~31 ℃, CO_2 concentration of 400 mL L^{-1}, air flow of 400 μmol

s^{-1}, and relative humidity of 50-60%. Leaf temperatures varied from 32-34℃ during the

measurements.

Chlorophyll fluorescence measurements were taken on the same leaves used for gas-

exchange using a Handy PEA (Hansatech Instruments Ltd., Norfolk, UK). Leaves were dark-

adapted for 20-30 min using leaf clips before taking measurements. The saturation pulse rate of

3000 μmol m^{-2} s^{-1} was used to measure the emission of chlorophyll fluorescence. The initial

chlorophyll fluorescence yield (F_0), the variable chlorophyll fluorescence yield (F_v), and the

maximum chlorophyll fluorescence yield (F_m) were recorded. The maximum photochemical

efficiency of PSII in dark-adapted leaves was obtained by calculating the ratio of F_v/F_m.

Data availability

All phenotypic data from the QTL and NIL populations are attached to this manuscript as

supplemental tables S1-S3. All marker data was previously used in Khangura *et al.* 2019 and made

available to the public via Figshare (doi 10.25387/G3.7370948).

Results

Negative affect of *Oy1-N1989* on flowering time is exacerbated by Mo17

While preparing the material for our previous study (Khangura *et al.* 2019), we noticed that *Oy1-*

N1989 exhibited a consistent delay in flowering. If this was due to variation in the accumulation

of chlorophyll, or downstream consequences of changes in chlorophyll content, we expect that

contribution of the *vey1* enhancing QTL from Mo17 should make this more severe. Heterozygous

Oy1-N1989 mutant plants in the Mo17 × B73 hybrid genetic background flower up to two-weeks

later than wild-type siblings. The *Oy1-N1989* mutants also flower later in an isogenic inbred B73

background; however, the delay is less than half with mutants flowering four days after wild-type

siblings (Figure 1; Table S4). The wild-type B73 and Mo17 F_1 hybrid plants flower earlier than

the wild-type B73 inbred plants (Table S4). Maize is protandrous in nature; tassels mature earlier

than the ear-inflorescence. The effect of *Oy1-N1989* and *vey1* on flowering time was similar for

both anthesis and silk emergence. The window of difference in maturity of the tassel and ear

inflorescence, measured as ASI, is wider in the wild-type siblings compared to the mutants in

Mo17 × B73 hybrid background. However, the wild-type and mutant siblings are not different in

B73 inbred background. This is consistent with the *vey1* allele from Mo17 impacting all

phenotypes recorded for *Oy1-N1989* allele to date, including delayed flowering.

Delayed flowering time of *Oy1-N1989* mutants in B73 × Mo17 mapping populations maps to

vey1

To test whether the effect in the hybrids was the result of the *vey1* locus, or some other factor in

the Mo17 background, we recording flowering time in wild-type and mutant plants of the F_1

progenies of *Oy1-N1989/oy1*:B73 pollen-parent with IBM-RILs and Syn10-DH. The crossing

scheme used for line-cross experiments have been described before (Khangura *et al.* 2019).

Hereafter, the hybrid populations developed from these crosses will be referred to as IBM-RILs

F_1 and Syn10-DH F_1 populations (Figure 2).

Pairwise correlations were calculated between previously reported chlorophyll index

measures (Khangura *et al.* 2019) and flowering time traits collected from the same plants (Tables

S5 and S6). The chlorophyll index was measured at two time points CCMI (25-30 days after

sowing) and CCMII (45-50 days after sowing). In both the IBM-RILs and Syn10-DH F_1

populations, significant positive correlations were observed between wild-type and mutant

flowering traits. In both populations, chlorophyll measurements in the wild-type siblings

(WT_CCMI and WT_CCMII) did not show any correlation with the mutant flowering time traits

(Figure 3, Table S5, and S6). In the IBM-RILs crosses, wild-type CCMII displayed a weak

significant negative correlation with wild-type DTA and DTS. Similarly, in the Syn10-DH crosses

wild-type CCMI displayed a significant weak negative correlation with wild-type DTA and DTS.

This indicates that the phenomena observed in our mutants, reduced chlorophyll content associated

with a delay in flowering time, could be seen in the wild-type siblings although the genetic basis

for the variation in wild-type chlorophyll was not resolved to the *oy1* locus (Khangura *et al.* 2019).

The most dramatic and obvious correlation were the negative correlation between CCM trait values

(CCMI and CCMII) and flowering time in the mutant siblings in both IBM-RILs and Syn10-DH

F_1 populations. The mutants in these populations also showed a clear increase in mean values for

days to anthesis and silking compared to wild-type siblings (Figure 4).

The frequency distribution plot of days to anthesis in mutant siblings of IBM-RILs and

Syn10-DH F_1 populations displayed a bimodal distribution, suggesting a single locus segregating

in these populations as the basis of flowering time suppression of *Oy1-N1989* mutants by B73

genotype (Figure 5). QTL mapping detected a single QTL on chromosome 10 at similar linkage

positions for all mutant derived flowering traits (Figure 5; Tables S7 and S8). This corresponds to

the *vey1* locus that we previously described as a major QTL that controls chlorophyll pigment

biosynthesis in the presence of the *Oy1-N1989* allele (Khangura *et al.* 2019). QTL mapping for

number of days to flower (anthesis and silking) in the mutant heterozygotes (MT_DTA and

MT_DTS), ratio of days to flower derived from mutant and wild-type siblings (Ratio_DTA and

Ratio_DTS), and the difference in days to flower between the wild-type and mutant siblings

(Diff_DTA and Diff_DTS) all detected *vey1* as the sole QTL in the population exceeding a

permutation-estimated significance threshold (alpha<0.05) in these traits and explaining ~40-48%

of the phenotypic variation in the IBM-RILs crosses, and >65% variation in the flowering traits in

Syn10-DH F_1 crosses. As expected, the *Oy1-N1989* enhancing *vey1^{Mo17}* allele was associated with

a delay in flowering time in the mutant hybrid siblings (Tables S7 and S8). In addition to *vey1*, an

additional QTL controlling mutant anthesis and silking was detected in the Syn10-DH F_1

population on chromosome 2 which we call *other oy1 flowering time locus1* (*oof1*). This QTL

explained ~7-8% variation with the Mo17 allele at this locus resulting in ~2-3 days of delay in

reproductive maturity of mutant siblings.

Because of the very large effect of *vey1* on flowering time, and the mild segregation

distortion at this locus in both bi-parental populations (Khangura *et al.* 2019), weak QTL might be

detected by spurious linkage between chr2 and chr10 markers. To test this, we carried out multiple

regressions using top markers at *oof1* and *vey1* as independent variables and DTA in the Syn10-

DH as a dependent variable (data not shown). In this analysis, both *vey1* and *oof1* remained a

significant factor in the multiple regression and explained 62% and 2% of the variance in flowering

time, respectively. Inclusion of a *vey1 × oof1* interaction term did not improve the fit of the model,

did not eliminate the significance of the *oof1* term, and the interaction term was not a significant

variable. Therefore, we propose that *oof1* is a novel QTL contingent upon the *Oy1-N1989* mutation

and genetically independent of *vey1*.

The mechanism(s) of the flowering time delays in *Oy1-N1989* mutants does not seem similar

to the effect of drought stress which differentially alters the time to silking and anthesis, commonly

measured by the anthesis-silking interval (ASI). The ASI of mutant and wild-type siblings were

not discernable and no QTL were detected for ASI in the mutants. In our conditions, we did not

detect any QTL for flowering time in wild-type siblings (Tables S7 and S8). Thus, detection of

vey1 and *oof1* in these QTL analyses was conditioned on the presence of the *Oy1-N1989* allele

suggesting a role for either photosynthesis or chlorophyll metabolites in the regulation of flowering

time.

We further tested the effect of the *vey1* critical region using the BM-NIL F_1 population that

has been previously described. We observed that introgression of *vey1* locus from Mo17 resulted

in delaying flowering time in both B73 and Mo17 isogenic backgrounds, whereas *vey1*

introgression from B73 into Mo17 can restore the negative effect of Mo17 genotype at this position

on flowering time (Figure 6, Figure S3, and Table S3). No effect of *vey1* introgression from either

parent was visible on the flowering time of the wild-type siblings. Representative plants showing

the delayed reproductive maturity of B73-like NILs carrying *vey1^{Mo17}*, and reciprocal introgression

of *vey1^{B73}* into Mo17 background displayed the opposite effect on flowering time (Figure 6). Thus,

these results clearly show the single locus effect of *vey1* on flowering time in maize in *Oy1-N1989*-

contingent manner in both isogenic inbred (B73) and hybrid (Mo17 × B73) background.

Expression polymorphism in B73-like NILs is consistent with cis-acting regulatory

polymorphism at *vey1*

Our previous study looking at the suppression of *Oy1-N1989* mutant phenotype using chlorophyll

accumulation identified a cis-eQTL at *oy1* in the IBM-RILs population (Khangura *et al.* 2019).

Normalized expression (RPKM) of OY1 derived from 14 days old shoot apices of IBM-RILs (Li

et al., 2013; 2018) were used for this analysis. The top marker, isu085b, used in the detection of

this cis-eQTL was also one of the top significant markers for mutant flowering time traits (Figure

S1 and Table S9). Regression of OY1 RPKM and flowering traits collected in IBM-RILs F_1

population identified a significant linear relationship between gene expression in wild-type inbred

lines and flowering time measurements from mutant F_1 siblings (Figure S2 and Table S9). Roughly

21% of the variation in mutant DTA could be explained by OY1 expression in the IBM-RILs shoot

apices. OY1 RPKM did not predict variation in wild-type DTA.

The allele specific-expression (ASE) assay in our previous study identified bias in expression

with the wild-type *oy1* allele from Mo17 displaying relatively lower expression (Khangura *et al.*

2019). Our previous work compared mutant plants in two different genetic backgrounds which can

complicate the interpretations due to trans effect in the hybrid background. To overcome this

limitation, an ASE assay was designed to test bias at *oy1* using near-isogenic lines in the B73

background. These B73-like NILs consisted of two independent NILs with *vey1^{Mo17}* introgression,

and two independent NILs with B73 genotype at *vey1*. This material was grown in the field and

expression of *oy1* was quantified in the top fully-expanded leaf at the V3 developmental stage.

Consistent with the previous ASE results, a significantly greater proportion of expression was

derived from the *Oy1-N1989* mutant allele when the wild-type *oy1* allele was contributed by

vey1^{Mo17} introgression as compared to the isogenic mutant siblings carrying the wild-type *oy1*

allele from *vey1^{B73}* locus (Table S10).

Net CO_2 assimilation and sugar metabolism is reduced in *Oy1-N1989* mutants in *vey1*-

dependent manner

We measured net CO_2 assimilation, sub-stomatal CO_2, photosystem II fluorescence, and

photosynthate accumulation in enhanced and suppressed *Oy1-N1989/oy1* mutants. A previous

study in maize has shown that reduction in chlorophyll levels in the leaves leads to a reduction in

photosynthetic rate (Huang *et al.* 2009). A similar reduction in photosynthetic rate should be

displayed by *Oy1-N1989/oy1* mutants and the Mo17 allele should show a greater reduction in

photosynthesis compared to the B73 allele. We tested this using F_1 progenies derived from four

B73-like NILs that were crossed to *Oy1-N1989/oy1*:B73 pollen. Two of these NILs had *vey1*^{Mo17}

introgression, whereas the other two had *vey1*^{B73} genotype. The negative effect of *vey1*^{Mo17}

introgression on chlorophyll accumulation was visible in the *Oy1-N1989* mutant allele background

(Figure 6). As expected, photosynthetic rate (A) was reduced in mutants as compared to wild-type

siblings, and mutants were modified further by the *vey1* genotype (Table S11). Photosystem II

efficiency (Fv/Fm) measurements indicated higher photo-oxidative damage to the photosystem in

enhanced mutant plants compared to the suppressed mutants. Wild-type siblings of all four B73-

like NIL F_1 progenies showed no statistically significant difference for chlorophyll and gas-

exchange measurements. This indicates that, just as for chlorophyll content, the *vey1* QTL affects

net CO_2 assimilation in the presence of the *Oy1-N1989* mutant allele. No differences in stomatal

conductance (g_s) or transpiration (E) was observed in the mutant plants, however, severe mutant

plants showed significantly higher accumulation of intracellular CO_2 (Ci) compared to the

suppressed mutants suggesting the failure of the enhanced mutant plants to uptake carbon (Table

S11).

These differences in photosynthetic rate should result in a decrease of non-structural

carbohydrates. The levels of non-structural carbohydrates were determined from leaves of the

same mutant plants that were used for gas-exchange measurements. Levels of sucrose, glucose,

fructose, and starch were measured in these samples. All of these showed a significant reduction

in the enhanced mutants compared to the suppressed mutants in the B73 isogenic background

(Figure 7 and Table S12). Lower levels of sugars and starch in the enhanced mutants are consistent

with the observation of low photosynthetic rate in these genotypes compared to the suppressed

mutant NILs.

Defoliation in *Zea mays*, *Sorghum bicolor*, and *Setaria viridis* delays flowering time

The existing literature suggests sugars and carbohydrate metabolism play an important role in

regulating flowering time in plants (Ohto *et al.* 2001; Seo *et al.* 2011; Wahl *et al.* 2013). We

hypothesized that removal of source tissue should mimic sugar starvation observed in *Oy1-N1989*

mutants. As a result, defoliation should result in delayed flowering in defoliated (treated) plants

compared to non-defoliated (control) plants. We tested this hypothesis using the wild-type inbred

strains in maize and two other monocot species: *Sorghum bicolor* (sorghum) and *Setaria viridis*

(green foxtail). Each species was defoliated at an early vegetative stage when only a few leaves

had fully expanded. The intent was to deprive plants of leaf area and photosynthate early in

development to separate the loss of chlorophyll and the loss of photosynthate, which are coupled

in the *Oy1-N1989* mutant study. Defoliation delayed flowering in all the plant species tested

(Figure 8). Maize, sorghum, and green foxtail displayed a delay in flowering by about 13, 7, and

4 days, respectively, compared to the control plants. Therefore, based on the defoliation study in

multiple monocot species we propose that early season defoliation results in the removal of source

tissue that might be critical for vegetative to floral transition in plants. Surprisingly, one-week

post-defoliation, newly emerged leaves displayed a pale leaf color. Chlorophyll estimation using

CCM found a reduction in the leaf greenness in the defoliated treatments compared to the control

samples (Figure 8). The newly emerging leaves of maize, sorghum, and green foxtail showed

~35%, ~48%, and ~58% reduction in chlorophyll, respectively, in the treated plants compared to

the control plants. Subsequent leaves emerging from the defoliated treatments displayed normal

leaf color suggesting recovery of the plants from defoliation stress. Thus, the delay in flowering

time is likely due to a loss of resources (carbon reserves) but cannot be formally distinguished

from a change in chlorophyll accumulation.

Leaf senescence is suppressed by *Oy1-N1989* mutants in a *vey1*-dependent manner

Leaf senescence can be induced in maize by sucrose accumulation in the leaves. This can be

accomplished genetically by disrupting sucrose transport (Braun *et al.* 2006; Baker and Braun

2008; Slewinski *et al.* 2009) or by preventing the maize ears from acting as a sink (Allison and

Weinmann 1970; Sekhon *et al.* 2012, 2019). In maize, previous studies have shown that leaf

senescence caused by pollination prevention or ear removal before pollination is genotype-

dependent (Ceppi et al. 1987; Allison and Weinmann 1970). We found that levels of free sugars

and starch were significantly lower in the young leaves of enhanced *Oy1-N1989* mutant B73 NILs

compared to the suppressed mutant NILs (Figure 8). If the accumulation of sugars in the leaves is

a pre-requisite for leaf senescence, we expected delayed senescence in the *Oy1-N1989* mutants

and for this to vary in a *vey1* dependent manner. We tested the effect of this differential sugar

accumulation in mutant B73-like NILs on induced leaf senescence by pollination avoidance. We

observed that 30 days-after-anthesis (DAA), the top leaves of the unpollinated wild-type B73

plants showed complete senescence of the leaves with only a few green patches. Unpollinated

Oy1-N1989 mutant plants in the suppressed B73 NIL background did not show such an extreme

phenotype but rather, showed only a few patches of anthocyanin accumulation and cell death by

30 DAA (Figure 9). On the other hand, unpollinated mutant plants in B73 NIL background with

vey1^{Mo17} introgression did not show any sign of leaf senescence at 30 DAA. At 42 DAA,

unpollinated wild-type and mutants with and without *vey1^{Mo17}* genotype showed leaf senescence,

however, the enhancement of *Oy1-N1989* by the *vey1^{Mo17}* resulted in substantially less cell death

and anthocyanin accumulation.

Discussion

Our previous work identified a modifier, *vey1*, that affects the chlorophyll accumulation in *Oy1-*

N1989/oy1 heterozygotes (Khangura *et al.* 2019). The *vey1* polymorphism(s) are common natural

variant(s) linked to the *oy1* locus of maize. We proposed that *vey1* results from cis-acting

regulatory polymorphisms that affect the expression of OY1. The suppression of the mutant, and

accumulation of chlorophyll, follows the proportion of wild-type and mutant transcript levels

(Table S10 and Khangura *et al.* 2019). In this work, we describe a delay in flowering in the *Oy1-*

N1989 mutants and demonstrate that *vey1* encodes a strong modifying QTL altering the flowering

time of mutant siblings in all mapping populations (Figures 5 and 6; Tables S7 and S8). Just as the

detection of *vey1* for CCM was contingent on *Oy1-N1989* mutant allele in the background, there

was no effect of *vey1* on flowering time in wild-type siblings. We observed the same marker,

isu085b, had the strongest effect on chlorophyll content, OY1 transcript abundance in shoot apices

(Khangura *et al.* 2019) and variation in flowering time of IBM-RILs x *Oy1-N1989* mutant F_1

mutant siblings (Table S9 and Figure S2). Taking these observations together, we propose that the

cis-acting eQTL at *oy1* is affecting chlorophyll level in *Oy1-N1989* mutants, and that the alteration

in photosynthesis through perturbed chlorophyll metabolism affects flowering time variation in

mutant siblings in these populations.

Our previous work on *vey1* has focused on the cryptic nature of the variation, and the

interaction between the *vey1* QTL and the *Oy1-N1989* mutant allele. However, the experiments

presented here clearly transcend this limitation and identify chlorophyll content as a heretofore

unappreciated correlate, if not determinant, of flowering time affecting even in wild-type siblings

(Figure 3; Tables S5 and S6). We observed that CCM values, especially CCMII in IBM-RILs and

CCMI in Syn10-DH F_1 populations, were negatively correlated with wild-type flowering time

measurements. This suggests a role for the normal variation in chlorophyll contents in the control

of flowering time in maize either via changes to photosynthetic output or signaling. As no QTL

were detected for CCM in the wild-type siblings in our previous work (Khangura *et al.* 2019), it

also suggests that mechanism responsible for CCM and flowering time covariation is likely

independent of *vey1*. This is further strengthened by the absence of an effect of *vey1* on flowering

time in the wild-type siblings in our NILs (Table S3 and Figure S3). It is possible that flowering

time QTL segregating in the B73 × Mo17 cross are fortuitously linked to other loci affecting CCM

in wild-type plants, which did not pass the threshold of detection in our QTL experiments.

Otherwise there should be no formal relationship between CCM and flowering time in other

populations with greater recombination or different alleles, such as in an association panel. Further

experiments are required to validate or reject the causal nature of the relationships between CCM

and flowering time in wild-type siblings of this material.

Even though Syn10-DH and IBM-RILs are derived from the same parents, they differ in the

method of development and rates of recombination (Liu et al. 2015; Hussain et al. 2007; Lee et al.

2002). Thus, these two populations yield different levels of resolution for QTL detection. Our

previous analysis to fine map *vey1* QTL using CCM values showed Syn10-DH to have a higher

mapping resolution than the IBM-RILs population (Khangura *et al.* 2019). We also observed

different QTL for flowering time in these populations. QTL analysis in Syn10-DH F_1 population

detected the *oof1* QTL on chromosome 2 affecting MT_DTA and MT_DTS as well as the major-

effect locus *vey1* (Table S8). The detection of *oof1* was dependent on the presence of the *Oy1-*

N1989 mutant, but was neither contingent on nor displayed any epistatic interactions with *vey1* as

assessed by multiple regression. Thus, *oof1* appears to be a novel locus of independent mechanism

affecting flowering time in the *Oy1-N1989* mutants. This QTL on chromosome 2 was not detected

for any other trait in the Syn10-DH and IBM-RILs testcross populations with *Oy1-N1989/oy1*:B73

in the current study nor was it identified as a modifier of chlorophyll content in our previous study

(Khangura *et al.* 2019). The amount of phenotypic variation explained by *vey1* in flowering time

was higher in Syn10-DH experimental material than the IBM-RILs testcross population by ~20%

(Tables S7 and S8). Greater variation in flowering time in the IBM-RILs testcross progenies that

could not be modeled by marker genotypes may result from higher residual heterozygosity or

greater rates of pollen contamination during the single seed descent (SSD) procedure used to

generate the IBM-RILs (Lee et al. 2002). Heterozygosity is expected to be negligible in Syn10-

DH population because of the DH procedure employed to fix allele segregation during population

development (Hussain *et al.* 2007). In addition, we had a moderate increase in sample size in the

Syn10-DH (251 lines) as compared to the IBM-RILs (216 lines) that is expected to result in a

marginally greater power to detect QTL in the Syn10-DH F_1 populations.

Chlorophyll levels are correlated with the rate of photosynthesis in plants (Huang *et al.* 2009).

Some controversy has been reported in mutants of soybean affected in an ortholog of *oy1* with

some reports showing little impact on photosynthesis (Sakowska *et al.* 2018) and others clearly

demonstrating a linear relationship between chlorophyll variation and photosynthesis (Walker *et*

al. 2017). The reasons for conflicting conclusions results from soybean mutants are not fully clear

but it warrants some caution in making simplistic interpretations about the impact of chlorophyll

deficient mutants on photosynthesis. In our study, net CO_2 assimilation was associated with the

severity of the *Oy1-N1989* mutant phenotype. The effect of chlorophyll loss on CO_2 assimilation

measurements using a LICOR instrument was somewhat non-linear. A nearly 5-fold reduction in

CCM in *Oy1-N1989* NILs carrying a B73 allele at *vey1* resulted in in a 22% reduction in net CO_2

assimilation while the 10-fold reduction in CCM in *Oy1-N1989* NILs carrying a Mo17 allele at

vey1 resulted in a 64% reduction in net photosynthetic rate, compared to their isogenic wild-type

siblings (Table S11). Ultimately, both reductions resulted in less accumulation of free sugars and

starch, with a substantially greater reduction in the enhanced mutant NILs (Figure 7; Tables S11

and S12).

The allelic interactions of *Oy1-N1989* and wild-type *oy1* alleles are consistent with the

inductive role of carbohydrate status on floral transition (Ohto et al. 2001; Seo et al. 2011; Wahl

et al. 2013; Minow et al. 2018). Additional work exploring proposed carbohydrate signaling

metabolites, such as T6P and organic acids, and analyses of the downstream floral integrators such

as the FT orthologs of maize are still needed to link our results to the existing models of floral

transition regulation in maize (Minow *et al.* 2018). One complexity that we observed in our data

is that the alternate experiments such as defoliation were not able to distinguish photosynthate

levels from changes in chlorophyll (Figure 8). The mechanical removal of leaves resulted in

changes to CCM. Two other subunits of magnesium chelatase are encoded by genes that were

identified in Arabidopsis as *genomes uncoupled* mutants, *gun4* and *gun5*, with altered retrograde,

plastid-to-nuclear, signaling with defects in chlorophyll metabolism (Susek *et al.* 1993; Mochizuki

et al. 2001; Larkin *et al.* 2003). Multiple studies have demonstrated retrograde signaling mutants

with defects in chlorophyll metabolism (Hernández-Verdeja and Strand 2018) and circadian

rhythm (Jones 2019). Retrograde signaling events are also associated with cell death, chloroplast

development, and etiolation but it acts via a number of modifiers of flowering, including

phytochromes, phytochrome signaling components, blue light perception via *crytochrome1,* and

circadian rhythmicity of key genes that affect flowering time. While retrograde signaling is an

attractive alternative model to sugar signaling for the phenomena reported here, mutants in

orthologs of *oy1* in both monocots and dicots (Mochizuki *et al.* 2001; Gadjieva *et al.* 2005) have

been tested for a *genomes uncoupled* phenotype and they did not perturb retrograde signaling. This

makes it unlikely that *Oy1-N1989* and *vey1* are altering flowering time via aberrant retrograde

signals. Additional experiments are necessary to clearly separate the effects of chlorophyll

metabolite levels from photosynthate levels to independently test their effects on flowering time

in maize.

We have known that defoliation is practiced in some maize nurseries to stagger flowering

time and permit intercrossing of lines with divergent reproductive maturities. We had presumed

that this was based on published research. Remarkably, we were not able to find a reference for

this practice. Previous work on defoliation in maize has looked at the effect of early and late season

defoliation on growth, and yield components but did not report flowering times (Crookston and

Hicks 1988; Pearson and Fletcher 2009). As a result, Figure 8 provides information that was

informally shared within the maize genetics research community but not described formally. In

addition, we extended this observation from maize to both sorghum and green foxtail, indicating

that defoliation can achieve staggered flowering times in these species as well. This may be a

general feature of grasses, but appears to not be universal in angiosperms as early-season

defoliation of photoperiod-sensitive strawberries did not affect flowering time (Guttridge 1959).

Surprisingly, the defoliation of maize, sorghum, and green foxtail resulted in reduced chlorophyll

accumulation in the new leaves. As a result, a mechanical treatment that was expected to only

affect photosynthetic leaf area, and not chlorophyll content, served to highlight the interconnected

nature of plant metabolism: large changes to any feature of central metabolism or plant physiology

result in large changes to all of central metabolism and plant physiology. Carefully conceived

narrow manipulation of the metabolites and additional studies of mechanisms of flowering time

modulation are required to determine how chlorophyll metabolism influences flowering time.

Sugar export and phloem loading mutants in maize that carry lesions in *tie-dyed1*, *tie-dyed2*,

sucrose export defective1, *psychedelic*, and *sucrose trasporter1* have all been shown to delay

flowering time (Braun *et al.* 2006; Baker and Braun 2008; Ma *et al.* 2008; Slewinski *et al.* 2009;

Slewinski and Braun 2010). We also observed declines in sugar levels in *Oy1-N1989* mutants

(Figure 7 and Table S12) and a delay in flowering time (Figure 1). It is tempting to consider these

mutants as a demonstration that sugar signaling can work independent of chlorophyll, but these

mutants also display low chlorophyll contents (Braun *et al.* 2006; Baker and Braun 2008; Ma *et*

al. 2008; Slewinski *et al.* 2009; Slewinski and Braun 2010). Curiously, *sucrose export defective1*

encodes tocopherol cyclase (Porfirova *et al.* 2002; Sattler *et al.* 2003). Tocopherol and chlorophyll

share the phytol side chain, and the salvage pathway for phytol side chains from chlorophyll can

contribute to tocopherol accumulation in both maize and Arabidopsis (Ischebeck *et al.* 2006;

Schelbert *et al.* 2009; Diepenbrock *et al.* 2017). While it is attractive to try and unify these findings,

chlorophyll breakdown products were only rate limiting for tocopherol synthesis in senescent

tissues of Arabidopsis (Zhang *et al.* 2014). Furthermore, in *Oy1-N1989* mutants the phytol

precursors should be abundant as this pool is not being consumed by chlorophyll biosynthesis.

Future experiments that more carefully explore these metabolites, for instance, via other mutants

that do not simultaneously affect multiple pathways are required. The finding that defoliation

impacts flowering time and the same biochemical and molecular responses as observed in the

photosynthetically compromised *Oy1-N1989* mutants may also provide mechanistic insights for

this phenomenon.

Accumulation of sugars in the leaves due to sink disruption has been proposed to induce leaf

senescence in a variety of angiosperms, including maize and Arabidopsis (Allison and Weinmann

1970; Ceppi *et al.* 1987; Pourtau *et al.* 2006; Sekhon *et al.* 2012, 2019). In maize, leaf senescence

is triggered when pollination is prevented and sucrose accumulates in leaves due to the lack of the

sink activity of a pollinated ear (Ceppi et al. 1987; Allison and Weinmann 1970). This sucrose-

dependent leaf senescence is genotype-dependent, and B73 is particularly susceptible to this

phenomenon. Genetic inheritance of the induced senescence phenomenon in maize inbred line

B73 is proposed to be under the control of a single dominant locus (Ceppi *et al.* 1987). The leaf

senescent phenotypes of the sucrose export mutants demonstrate that excessive photosynthate

accumulation can cause tissue to senesce regardless of ear presence (Braun *et al.* 2006; Baker and

Braun 2008; Ma *et al.* 2008; Slewinski *et al.* 2009). Induced leaf senescence by sink removal or

sugar application shows some overlap of biochemical and molecular mechanisms with natural

senescence in plants (summarized in Sekhon et al. 2012). A study of gene expression in the B73

inbred of maize identified senescence associated genes that exhibit gene expression changes during

pollination-prevented leaf senescence (Sekhon et al. 2012). We found that leaf senescence could

be prevented or delayed by the suppression of photosynthesis in the *Oy1-N1989* mutant and further

modulated by *vey1* variants. We expect that future gene expression studies in *Oy1-N1989* mutants

and sugar export mutants will identify genes consistently impacted by lower and higher

chlorophyll levels, presumably including senescence associated genes and genes regulating

carbohydrate metabolism.

Conclusions

Flowering time in plants is regulated by environmental and endogenous cues. In this study, we

demonstrate that the interaction between natural and induced variation at the *oy1* locus of maize,

encoding the magnesium chelatase subunit I responsible for the first committed step of chlorophyll

biosynthesis, regulates flowering time. The effect of *oy1* variation on flowering time and other

physiological traits is consistent with the previously proposed and widely-accepted role of

photosynthetically derived sugars on flowering time. The previously identified *vey1* QTL, a

putative cis-acting polymorphism(s) in the regulatory region proximal to *oy1*, modified the

flowering time delay in the *Oy1-N1989* mutant heterozygotes. Further, *vey1* and *Oy1-N1989*

affected gas-exchange parameters, chlorophyll fluorescence, sugar accumulation, and starch

accumulation in the immature leaves of *Oy1-N1989* mutant plants. Thus, all data are consistent

with *Oy1-N1989* altering flowering time by perturbing carbon fixation via photosynthesis.

Acknowledgements

We would like to thank the National Science Foundation for NSF grant 1444503 awarded to G.S.J.

and B.P.D which helped fund this research. We thank the Purdue ACRE and ICSC staff members

Mr. Jim Beaty and Mr. Jason Adams for help with planting, and management of the field

experiments described in this work. We also thank past Johal lab members for help with planting

and data collection. We would like to thank Mike Gosney for helping with gas-exchange

measurements. We gratefully acknowledge the bobcat operator for dumping loads of gravel on our

office walls while we attempted to write this paper. Solidarity with all.

REFERENCES

Abe, M., Y. Kobayashi, S. Yamamoto, Y. Daimon, A. Yamaguchi *et al.*, 2005 FD , a bZIP Protein Mediating Signals from the Floral Pathway Integrator FT at the Shoot Apex. Science 309: 1052–1056.

Allison, J. C. S., and H. Weinmann, 1970a Effect of Absence of Developing Grain on and Senescence of Maize Leaves. Plant Physiol. 46: 435–436.

Alter, P., S. Bircheneder, L.-Z. Zhou, U. Schlüter, M. Gahrtz *et al.*, 2016 Flowering Time-Regulated Genes in Maize Include the Transcription Factor ZmMADS1. Plant Physiol. 172: 389–404.

Amasino, R. M., and S. D. Michaels, 2010 The Timing of Flowering. Plant Physiol. 154: 516–520.

Baker, R. F., and D. M. Braun, 2008 Tie-dyed2 functions with tie-dyed1 to promote carbohydrate export from maize leaves. Plant Physiol. 146: 1085–1097.

Bomblies, K., R.-L. Wang, B. A. Ambrose, R. J. Schmidt, R. B. Meeley *et al.*, 2003 Duplicate FLORICAULA/LEAFY homologs zfl1 and zfl2 control inflorescence architecture and flower patterning in maize. Development 130: 2385–2395.

Bouchet, S., P. Bertin, T. Presterl, P. Jamin, D. Coubriche *et al.*, 2017 Association mapping for phenology and plant architecture in maize shows higher power for developmental traits compared with growth influenced traits. Heredity (Edinb). 118: 249–259.

Braun, D. M., Y. Ma, N. Inada, M. G. Muszynski, and R. F. Baker, 2006 tie-dyed1 Regulates carbohydrate accumulation in maize leaves. Plant Physiol. 142: 1511–22.

Broman, K. W., H. Wu, S. Sen, and G. a. Churchill, 2003 R/qtl: QTL mapping in experimental crosses. Bioinformatics 19: 889–890.

Buckler, E. S., J. B. Holland, P. J. Bradbury, C. B. Acharya, P. J. Brown *et al.*, 2009 The genetic architecture of maize flowering time. Science 325: 714–718.

Ceppi, D., M. Sala, E. Gentinetta, A. Verderio, and M. Motto, 1987 Genotype-Dependent Leaf Senescence in Maize INHERITANCE AND EFFECTS OF POLLINATION-PREVENTION. Plant Physiol. 85: 720–725.

Chen, M., J. Chory, and C. Fankhauser, 2004 Light Signal Transduction in Higher Plants. Annu. Rev. Genet. 38: 87–117.

Cho, L. H., R. Pasriga, J. Yoon, J. S. Jeon, and G. An, 2018 Roles of Sugars in Controlling Flowering Time. J. Plant Biol. 61: 121–130.

Cho, L. H., J. Yoon, and G. An, 2017 The control of flowering time by environmental factors. Plant J. 90: 708–719.

Colasanti, J., Z. Yuan, and V. Sundaresan, 1998 The indeterminate gene encodes a zinc finger protein and regulates a leaf-generated signal required for thetransition to flowering in maize. Cell 93: 593–603.

Coles, N. D., M. D. McMullen, P. J. Balint-Kurti, R. C. Pratt, and J. B. Holland, 2010 Genetic

control of photoperiod sensitivity in maize revealed by joint multiple population analysis.
Genetics 184: 799–812.

Coneva, V., D. Guevara, S. J. Rothstein, and J. Colasanti, 2012 Transcript and metabolite signature
of maize source leaves suggests a link between transitory starch to sucrose balance and the
autonomous floral transition. J. Exp. Bot. 63: 5079–5092.

Coneva, V., T. Zhu, and J. Colasanti, 2007 Expression differences between normal and
indeterminate1 maize suggest downstream targets of ID1, a floral transition regulator in maize.
J. Exp. Bot. 58: 3679–3693.

Corbesier, L., P. Lejeune, and G. Bernier, 1998 The role of carbohydrates in the induction of
flowering in Arabidopsis thaliana: Comparison between the wild type and a starchless mutant.
Planta 206: 131–137.

Corbesier, L., C. Vincent, S. Jang, F. Fornara, Q. Fan et al., 2007 FT protein movement contributes
to long-distance signaling in floral induction of Arabidopsis. Science 316: 1030–1033.

Crookston, R. K., and D. R. Hicks, 1988 Effect of Early Defoliation on Maize Growth and Yield:
an Eleven-year Perspective. Crop Sci. 28: 371–373.

Danilevskaya, O. N., X. Meng, D. A. Selinger, S. Deschamps, P. Hermon et al., 2008 Involvement
of the MADS-Box Gene ZMM4 in Floral Induction and Inflorescence Development in Maize.
Plant Physiol. 147: 2054–2069.

DePristo, M. A., E. Banks, R. Poplin, K. V Garimella, J. R. Maguire et al., 2011 A framework for
variation discovery and genotyping using next-generation DNA sequencing data. Nat. Genet.
43: 491–498.

Diepenbrock, C. H., C. B. Kandianis, A. E. Lipka, M. Magallanes-Lundback, B. Vaillancourt et
al., 2017 Novel Loci Underlie Natural Variation in Vitamin E Levels in Maize Grain. Plant
Cell 29: 2374–2392.

Doebley, J., A. Stec, and L. Hubbard, 1997 The evolution of apical dominance in maize. Nature
386: 485–488.

Eichten, S. R., J. M. Foerster, N. de Leon, Y. Kai, C.-T. Yeh et al., 2011 B73-Mo17 near-isogenic
lines demonstrate dispersed structural variation in maize. Plant Physiol. 156: 1679–90.

Evans, M. M., and R. S. Poethig, 1995 Gibberellins promote vegetative phase change and
reproductive maturity in maize. Plant Physiol. 108: 475–87.

Eveland, A. L., and D. P. Jackson, 2012 Sugars, signalling, and plant development. J. Exp. Bot.
63: 3367–3377.

Gadjieva, R., E. Axelsson, U. Olsson, and M. Hansson, 2005 Analysis of gun phenotype in barley
magnesium chelatase and Mg-protoporphyrin IX monomethyl ester cyclase mutants. Plant
Physiol. Biochem. 43: 901–908.

Gibson, L. C. D., R. D. Willows, C. G. Kannangara, D. von Wettstein, and C. N. Hunter, 1995
Magnesium-protoporphyrin chelatase of Rhodobacter sphaeroides: reconstitution of activity
by combining the products of the bchH, -I, and -D genes expressed in Escherichia coli. Proc.
Natl. Acad. Sci. U. S. A. 92: 1941–1944.

Guttridge, C. G., 1959 Evidence for a Flower Inhibitor and Vegetative Growth Promoter in the

Strawberry. Ann. Bot. 23: 351–360.

Hernández-Verdeja, T., and Å. Strand, 2018 Retrograde Signals Navigate the Path to Chloroplast
Development. Plant Physiol. 176: 967–976.

Huang, M., T. L. Slewinski, R. F. Baker, D. Janick-Buckner, B. Buckner *et al.*, 2009 Camouflage
patterning in maize leaves results from a defect in porphobilinogen deaminase. Mol. Plant 2:
773–89.

Huang, C., H. Sun, D. Xu, Q. Chen, Y. Liang *et al.*, 2017 ZmCCT9 enhances maize adaptation to
higher latitudes. Proc. Natl. Acad. Sci. 115: E334–E341.

Hung, H.-Y., L. M. Shannon, F. Tian, P. J. Bradbury, C. Chen *et al.*, 2012 ZmCCT and the genetic
basis of day-length adaptation underlying the postdomestication spread of maize. Proc. Natl.
Acad. Sci. 109: E1913–E1921.

Hussain, T., P. Tausend, G. Graham, and J. Ho, 2007 Registration of IBM2 SYN10 Doubled
Haploid Mapping Population of Maize. J. Plant Regist. 1: 81.

Ischebeck, T., A. M. Zbierzak, M. Kanwischer, and P. Dörmann, 2006 A salvage pathway for
phytol metabolism in Arabidopsis. J. Biol. Chem. 281: 2470–2477.

Jones, M. A., 2019 Retrograde signalling as an informant of circadian timing. New Phytol. 221:
1749–1753.

Karl, J., 2016 Maize Leafy1 45ft. Maize Genet. Coop. Newsl. 90: 1–15.

Khangura, R. S., S. Marla, B. P. Venkata, N. J. Heller, G. S. Johal *et al.*, 2019 A Very Oil Yellow1
Modifier of the Oil Yellow1-N1989 Allele Uncovers a Cryptic Phenotypic Impact of Cis-
regulatory Variation in Maize. G3 9: 375–390.

Kozaki, A., S. Hake, and J. Colasanti, 2004 The maize ID1 flowering time regulator is a zinc finger
protein with novel DNA binding properties. Nucleic Acids Res. 32: 1710–1720.

Lang, A., 1952 Physiology of Flowering. Ann. Rev. Plant. Physiol. 3: 265–306.

Lang, A., 1957 The effect of gibberellin upon flower formation. Proc. Natl. Acad. Sci. 43: 709–
717.

Larkin, R. M., J. M. Alonso, J. R. Ecker, and J. Chory, 2003 GUN4, a regulator of chlorophyll
synthesis and intracellular signaling. Science 299: 902–906.

Leach, K. A., and D. M. Braun, 2016 Soluble Sugar and Starch Extraction and Quantification from
Maize (*Zea mays*) Leaves. Curr. Protoc. Plant Biol. 1: 139–161.

Leach, K. A., T. M. Tran, T. L. Slewinski, R. B. Meeley, and D. M. Braun, 2017 Sucrose
transporter2 contributes to maize growth, development, and crop yield. J. Integr. Plant Biol.
59: 390–408.

Lee, M., N. Sharopova, W. D. Beavis, D. Grant, M. Katt *et al.*, 2002 Expanding the genetic map
of maize with the intermated B73 x Mo17 (IBM) population. Plant Mol. Biol. 48: 453–61.

Liang, Y., Q. Liu, X. Wang, C. Huang, G. Xu *et al.*, 2019 ZmMADS69 functions as a flowering
activator through the ZmRap2.7-ZCN8 regulatory module and contributes to maize flowering
time adaptation. New Phytol. 221: 2335–2347.

Liu, H., Y. Niu, P. J. Gonzalez-Portilla, H. Zhou, L. Wang *et al.*, 2015 An ultra-high-density map

as a community resource for discerning the genetic basis of quantitative traits in maize. BMC
Genomics 16: 1078.

Ma, Y., R. F. Baker, M. Magallanes-Lundback, D. Dellapenna, and D. M. Braun, 2008 Tie-dyed1
and Sucrose export defective1 act independently to promote carbohydrate export from maize
leaves. Planta 227: 527–538.

Meng, X., M. G. Muszynski, and O. N. Danilevskaya, 2011 The FT-Like ZCN8 Gene Functions
as a Floral Activator and Is Involved in Photoperiod Sensitivity in Maize. Plant Cell 23: 942–
960.

Minow, M. A. A., L. M. Ávila, K. Turner, E. Ponzoni, I. Mascheretti *et al.*, 2018 Distinct gene
networks modulate floral induction of autonomous maize and photoperiod-dependent teosinte.
J. Exp. Bot. 69: 2937–2952.

Mochizuki, N., J. a Brusslan, R. Larkin, a Nagatani, and J. Chory, 2001 Arabidopsis genomes
uncoupled 5 (GUN5) mutant reveals the involvement of Mg-chelatase H subunit in plastid-
to-nucleus signal transduction. Proc. Natl. Acad. Sci. U. S. A. 98: 2053–2058.

Moghaddam, M. R. B., and W. Van den Ende, 2013 Sugars, the clock and transition to flowering.
Front. Plant Sci. 4: 1–6.

Muszynski, M. G., T. Dam, B. Li, D. M. Shirbroun, Z. Hou *et al.*, 2006 delayed flowering1
Encodes a basic leucine zipper protein that mediates floral inductive signals at the shoot apex
in maize. Plant Physiol. 142: 1523–1536.

Mutasa-Göttgens, E., and P. Hedden, 2009 Gibberellin as a factor in floral regulatory networks. J.
Exp. Bot. 60: 1979–1989.

Ohto, M.-A., K. Onai, Y. Furukawa, E. Aoki, T. Araki *et al.*, 2001 Effects of Sugar on Vegetative
Development and Floral Transition in Arabidopsis. Plant Physiol. 127: 252–261.

Paul, M. J., 2008 Trehalose 6-phosphate: a signal of sucrose status. Biochem. J. 412: 1–2.

Pearson, A., and A. L. Fletcher, 2009 Effect of total defoliation on maize growth and yield. Agron.
New Zeal. 39: 1–6.

Porfirova, S., E. Bergmuller, S. Tropf, R. Lemke, and P. Dormann, 2002 Isolation of an
Arabidopsis mutant lacking vitamin E and identification of a cyclase essential for all
tocopherol biosynthesis. Proc. Natl. Acad. Sci. 99: 12495–12500.

Pourtau, N., R. Jennings, E. Pelzer, J. Pallas, and A. Wingler, 2006 Effect of sugar-induced
senescence on gene expression and implications for the regulation of senescence in
Arabidopsis. Planta 224: 556–568.

R core Team, 2013 R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, http://www.R-project.org/.

Sakowska, K., G. Alberti, L. Genesio, A. Peressotti, G. Delle Vedove *et al.*, 2018 Leaf and canopy
photosynthesis of a chlorophyll deficient soybean mutant. Plant Cell Environ. 41: 1427–1437.

Salvi, S., G. Sponza, M. Morgante, D. Tomes, X. Niu *et al.*, 2007 Conserved noncoding genomic
sequences associated with a flowering-time quantitative trait locus in maize. Proc. Natl. Acad.
Sci. U. S. A. 104: 11376–11381.

Samach, A., H. Onouchi, S. E. Gold, G. S. Ditta, Z. Schwarz- *et al.*, 2000 Distinct Roles of

CONSTANS Target Genes in Reproductive Development of Arabidopsis. Science 288:
1613–1616.

Sattler, S. E., E. B. Cahoon, S. J. Coughlan, and D. Dellapenna, 2003 Characterization of
Tocopherol Cyclases from Higher Plants and Cyanobacteria . Evolutionary Implications for
Tocopherol Synthesis and Function. Plant Physiol. 132: 2184–2195.

Sawers, R. J. H., J. Viney, P. R. Farmer, R. R. Bussey, G. Olsefski et al., 2006 The maize Oil
yellow1 (Oy1) gene encodes the I subunit of magnesium chelatase. Plant Mol. Biol. 60: 95–
106.

Schelbert, S., S. Aubry, B. Burla, B. Agne, F. Kessler et al., 2009 Pheophytin Pheophorbide
Hydrolase (Pheophytinase) Is Involved in Chlorophyll Breakdown during Leaf Senescence
in Arabidopsis. Plant Cell 21: 767–785.

Searle, N. E., 1965 Physiology of Flowering. Ann. Rev. Plant. Physiol. 16: 97–118.

Sekhon, R. S., K. L. Childs, N. Santoro, C. E. Foster, C. R. Buell et al., 2012 Transcriptional and
Metabolic Analysis of Senescence Induced by Preventing Pollination in Maize. Plant Physiol.
159: 1730–1744.

Sekhon, R. S., C. Saski, R. Kumar, B. Flinn, F. Luo et al., 2019 Integrated Genome-Scale Analysis
Identifies Novel Genes and Networks Underlying Senescence in Maize. Plant Cell
tpc.00930.2018.

Sen, T. Z., L. C. Harper, M. L. Schaeffer, C. M. Andorf, T. E. Seigfried et al., 2010 Choosing a
genome browser for a Model Organism Database: surveying the Maize community. Database
2010: 1–9.

Seo, P. J., J. Ryu, S. K. Kang, and C. M. Park, 2011 Modulation of sugar metabolism by an
INDETERMINATE DOMAIN transcription factor contributes to photoperiodic flowering in
Arabidopsis. Plant J. 65: 418–429.

Slewinski, T. L., and D. M. Braun, 2010 The Psychedelic genes of maize redundantly promote
carbohydrate export from leaves. Genetics 185: 221–232.

Slewinski, T. L., A. Garg, G. S. Johal, and D. M. Braun, 2010 Maize SUT1 functions in phloem
loading. Plant Signal. Behav. 5: 1–4.

Slewinski, T. L., R. Meeley, and D. M. Braun, 2009 Sucrose transporter1 functions in phloem
loading in maize leaves. J. Exp. Bot. 60: 881–892.

Steinhoff, J., W. Liu, J. C. Reif, G. Della Porta, N. Ranc et al., 2012 Detection of QTL for
flowering time in multiple families of elite maize. Theor. Appl. Genet. 125:.

Susek, R. E., F. M. Ausubel, and J. Chory, 1993 Signal Transduction Mutants of Arabidopsis
Uncouple Nuclear CAB and RBCS Gene Expression from Chloroplast Development. Cell 74:
787–799.

Swarts, K., R. M. Gutaker, B. Benz, M. Blake, R. Bukowski et al., 2017 Genomic estimation of
complex traits reveals ancient maize adaptation to temperate North America. Science 357:
512–515.

Vega, S. H., M. Sauer, J. A. J. Orkwiszewski, and S. Poethig, 2002 The early phase change gene
in Maize. Plant Cell 14: 133–147.

Wahl, V., J. Ponnu, A. Schlereth, S. Arrivault, T. Langenecker *et al.*, 2013 Regulation of flowering
by Trehalose-6-phosphate signalling in Arabidopsis thaliana. Science 339: 704–708.

Walker, B. J., D. T. Drewry, R. A. Slattery, A. VanLoocke, Y. B. Cho *et al.*, 2017 Chlorophyll can
be reduced in crop canopies with little penalty to photosynthesis. Plant Physiol. 176: 1215–
1232.

Walker, C. J., and J. D. Weinstein, 1991 In vitro assay of the chlorophyll biosynthetic enzyme Mg-
chelatase: resolution of the activity into soluble and membrane-bound fractions. Proc. Natl.
Acad. Sci. U. S. A. 88: 5789–5793.

Wang, R. L., A. Stec, J. Hey, L. Lukens, and J. Doebley, 1999 The limits of selection during maize
domestication. Nature 398: 236–239.

Wigge, P. A., M. C. Kim, K. E. Jaeger, W. Busch, M. Schmid *et al.*, 2005 Integration of Spatial
and Temporal Information During Floral Induction in Arabidopsis. Science 309: 1056–1059.

Zeevaart, J. A. D., 1962 Physiology of Flowering. Science 137: 723–731.

Zhang, W., T. Liu, G. Ren, S. Hörtensteiner, Y. Zhou *et al.*, 2014 Chlorophyll Degradation: The
Tocopherol Biosynthesis-Related Phytol Hydrolase in Arabidopsis Seeds Is Still Missing.
Plant Physiol. 166: 70–79.

Zhang, L., H. Yu, S. Lin, and Y. Gao, 2016 Molecular Characterization of FT and FD Homologs
from Eriobotrya deflexa Nakai forma koshunensis. Front. Plant Sci. 7: 1–10.

Figure 1. Reproductive maturity is delayed by *Oy1-N1989* allele in maize (A) Wild-type (left) and mutant (right) sibling from Mo17 x *Oy1-N1989/oy1*:B73 cross (reproduced from Figure 1b from Khangura *et. al.*, 2019), black scale bar = 50 cm. (B) Wild-type (left) and mutant (right) sibling from Mo17 x *Oy1-N1989/oy1*:Mo17 cross. Close-up view of the emerging tassel of mutant siblings (C) from panel A, and (D) from panel B. Distribution of (E) days to anthesis, (F) days to silking, and (G) Anthesis-silking interval (ASI) in the wild-type and mutant siblings in B73 and Mo17 x B73 (Mo17/B73) hybrid genetic backgrounds.

Figure 2. Crossing scheme to map *Oy1-N1989* enhancer/suppressor loci using IBM-RILs (n=216) and Syn10-DH (n=251) populations. Black and blue colors indicate B73 and Mo17 genotypes. Chromosome 10 of the heterozygous pollen parent *Oy1-N1989/+*:B73 is shown with a golden spot indicating *Oy1-N1989* mutant allele. The resulting F_1 progenies from these crosses are depicted here with state of chromosome 10 for each F_1 testcross showing segregation of wildtype and mutant (with the golden spot) siblings.

Figure 3. Pairwise correlation matrix between wildtype CCMII (WT_CCMII), mutant CCMII (MT_CCMII), wildtype days to anthesis (WT_DTA), and mutant days to anthesis (MT_DTA) in (A) IBM-RILs x *Oy1-N1989/oy1*:B73, and (B) Syn10-DH x *Oy1-N1989/oy1*:B73 F_1 test cross populations.

Figure 4. Distribution of days to flower (anthesis and silking) in the wildtype and mutant siblings in (A) IBM-RILs x *Oy1-N1989/oy1*:B73, and (B) Syn10-DH x *Oy1-N1989/oy1*:B73 F$_1$ test cross populations. Abbreviations: wildtype days to anthesis (WT_DTA), wildtype days to silking (WT_DTS), mutant days to anthesis (MT_DTA), and mutant days to silking (DTS).

Figure 5. Distribution of days to 50% anthesis of the mutant siblings (MT_DTA) in
(A) Syn10-DH x *Oy1-N1989/oy1*:B73, and (B) IBM-RILs x *Oy1-N1989/oy1*:B73 F_1 population. Genome wide QTL plot of MT_DTA in (C) Syn10-DH, and (D) IBM-RILs F_1 population. X-axis indicates the chromosome number and Y-axis indicates the logarithm of odds (LOD) of tested markers. Black horizontal bar indicates permutation testing based threshold for QTL detection.

Figure 6. Effect of *vey1* locus from B73 and Mo17 on flowering time in reciprocal isogenic backgrounds
(A) Delayed maturity of the heterozygous mutants in isogenic Mo17 (BC$_7$ generation) inbred background, compared to the wildtype siblings. Mutant and wildtype siblings in b030 (B73-like NIL with *vey1^{B73}*) shows early flowering while the isogenic B73 introgression of the *vey1^{Mo17}* locus in b094 exhibits delayed flowering.
(B) Early maturity in the heterozygous mutant in m097 (Mo17-like NIL with *vey1^{B73}*) compared to the mutant in B73 x Mo17 F$_1$ hybrid background. Measuring stick on the left in both panels is 243 cm.

Figure 7. Effect of *vey1* on leaf soluble sugars and starch content in *Oy1-N1989/oy1* heterozygotes in the isogenic B73 NILs. NILs b135, and b185 have *vey1^B73^* introgression, whereas b189, and b094 have *vey1^Mo17^* introgression.

Figure 8. Effect of defoliation on flowering time and leaf color in (A-C) maize, (D-F) sorghum, and (G-I) green foxtail. Asterisk indicates significant difference between treatment means using student's t-test at p<0.05.

Figure 9. Effect of *vey1* on senescence induced by pollination prevention. Pictures of the representative primary ear leaf derived from plants at 30 and 40 days-after-anthesis (DAA) either with open pollinated (top panels: A, C, and E) or unpollinated (bottom panels: B, D, and F) ears. Representative primary ear leaf from (A-B) B73 wildtype (left) and mutant sibling (right) at 30DAA, (C-D) wildtype (left) and mutant (right) sibling from F_1 between B73-like NIL-b107 (with *vey1^Mo17* introgression) X *Oy1-N1989/oy1*:B73 at 30 DAA, (E-F) wildtype (B73-like NIL-b107), mutant B73-like NIL-b135 with *vey1^B73* (middle), and mutant B73-like NIL-b107 with *vey1^Mo17* (right) at 42 DAA. Scale bar in each figure is 6.35 cm.