RESEARCH ARTICLE

Lignin-based resistance to *Cuscuta campestris* parasitism in

Heinz resistant tomato cultivars

Min-Yao Jhu[1*], Moran Farhi[1,2*], Li Wang[1,3], Richard N. Philbrook[1], Michael S.

Belcher[4,5], Hokuto Nakayama[1], Kristina S. Zumstein[1], Steven D. Rowland[1], Mily

Ron[1], Patrick M. Shih[1,4,6,7], Neelima R. Sinha[1@].

[1] Department of Plant Biology, University of California, Davis, CA, 95616, United

States.

[2] Research, Archer Daniels Midland Company, 1554 Drew Avenue, Davis, California,

95618 USA.

[3] College of Forestry, Beijing Forestry University, Beijing, 100083, PR China.

[4] Feedstocks Division, Joint BioEnergy Institute, Emeryville, CA, United States.

[5] Department of Plant and Microbial Biology, University of California, Berkeley,

Berkeley, CA, United States.

[6] Genome Center, University of California, Davis, Davis, CA, United States.

[7] Environmental Genomics and Systems Biology Division, Lawrence Berkeley National.

[@] Corresponding author: Neelima R. Sinha

Email: nrsinha@ucdavis.edu.

[*] These authors contributed equally to the work.

Short title: Lignin-based resistance blocks dodder entry

One-sentence summary: Four key regulators confer lignin accumulation in the tomato

stem cortex blocking *C. campestris* host penetration upon infection.

The author responsible for distribution of materials integral to the findings presented in

this article in accordance with the policy described in the Instructions for Authors

(www.plantcell.org) is: Neelima R. Sinha (nrsinha@ucdavis.edu).

Abstract

Cuscuta species (dodders) are common agriculturally destructive parasitic angiosperms. These parasitic plants use haustoria as physiological bridges to extract nutrients and water from hosts. *Cuscuta campestris* has a broad host range and wide geographical distribution. While some wild tomato relatives are resistant, cultivated tomatoes are generally susceptible to *C. campestris* infestations. However, some Heinz tomato hybrid cultivars exhibit resistance to this parasite. The stem cortex in these resistant lines responds with local lignification upon *C. campestris* attachment, preventing parasite entry into the host. We compared gene expression patterns under *C. campestris* infestation in resistant and susceptible cultivars and identified *LIF1* (*Lignin Induction Factor 1*, an *AP2*-like transcription factor), *SlMYB55*, and *CuRLR1* (*Cuscuta R-gene for Lignin-based Resistance 1*, a *CC-NBS-LRR*) as key factors conferring host resistance. Transient overexpression results suggest that *SlMYB55* and *LIF1* directly regulate cortical lignification. We also identified *SlWRKY16*, a transcription factor that is upregulated in all tested tomato cultivars upon *C. campestris* attachment. *SlWRKY16* acts as a negative regulator of *LIF1* function. Moreover, *CuRLR1* responds to a large protein molecule from *C. campestris* extracts and may play a role in signaling or function as a receptor for receiving *C. campestris* signals to regulate lignification-based resistance.

Main Text

Introduction

Parasitic plants directly attach to hosts using specialized haustorial organs known as haustoria. These connections function as physiological bridges to extract nutrients and water from the hosts, making most traditional herbicides and control methods, including management of soil fertility, hand weeding, and sanitation, either too cost-intensive, labor-intensive or ineffective in regulating parasitic plant infestations. Therefore, parasitic angiosperms are among the most devastating pests, reducing the yields of agricultural crops each year by billions of dollars worldwide (Agrios, 2005; Yoder and Scholes, 2010). Members of the *Cuscuta* genus (family Convolvulaceae), also known as dodders, occur worldwide and *Cuscuta* infestations in tomato alone lead to 50–72% yield-reductions (Yaakov et al., 2001). Despite serious agricultural problems caused by *Cuscuta*, our understanding of the interactions between *Cuscuta* and its hosts is relatively limited compared to our knowledge of pathogenic fungi, bacteria, and viruses. Only recently, the first receptor (*CuRe1*, Solyc08g016270) from *Cuscuta* was identified in tomatoes (Hegenauer et al., 2016). *CuRe1* initiates PAMP (Pathogen-associated molecular pattern)-triggered immunity (PTI) to *Cuscuta reflexa*. However, plants that lack *CuRe1* are still fully resistant to *C. reflexa*. This result indicates that other layers of defense, besides *CuRe1,* must also be involved in the responses to these parasites. Therefore, further investigating the potential multilayered resistance mechanisms will aid in developing parasitic plant-resistant crops.

Potential immune responses and defense responses to parasitic plants have been observed in several crop plants, including rice, tomato, cowpea, and sunflower (Mutuku et al., 2015; Hegenauer et al., 2016; Duriez et al., 2019; Su et al., 2020). Notably, most previous reports indicated that hypersensitive response is the major mechanism that contributes to plant immunity to parasitic plants (LANE et al., 1993; Hegenauer et al., 2016; Su et al., 2020). A few studies indicated that secondary cell-wall modification and formation also contribute to defense in root parasitic plants (Yoshida and Shirasu, 2009; Mutuku et al., 2019). Plants often modify their cell walls in response to pathogen

infection and herbivore feeding (Moura et al., 2010). Among different modifications,

lignification is considered a major mechanism for resistance in plants (Vance et al., 1980;

Moura et al., 2010; Bellincampi et al., 2014; Malinovsky et al., 2014). Lignified cell

walls have higher mechanical strength, are impermeable to water, and less accessible to

cell wall-degrading enzymes (Bhuiyan et al., 2009; Barros et al., 2015). However, how

secondary cell-wall modification and lignin are involved in the defense responses toward

stem parasitic plants still remain largely unknown. Thus, we specifically investigated host

cell wall composition and the lignin biosynthesis pathways aiming to discover the

potential additional layers of resistance to *Cuscuta* spp.

Cuscuta campestris (*C. campestris*) attacks a wide range of crop species

worldwide (Lanini and Kogan, 2005). Although tomatoes are usually susceptible

(Ashton, 1976), some cultivars are resistant to *Cuscuta* (Hembree et al., 1999; Yaakov et

al., 2001). These dodder-resistant tomatoes were used to identify genes involved in

defense responses. We discovered that the resistance response in these cultivars is based

on lignification in the stem cortex upon *C. campestris* infection. Recent work described

the involvement of lignin in the resistance responses to root parasitic plants (Labrousse et

al., 2001; Kusumoto et al., 2007; Cui et al., 2018), but the regulatory mechanisms remain

unknown. Based on comparative transcriptomics and gain-of-function studies in

susceptible cultivars, we identified two transcription factors, *SlMYB55* (Solyc10g044680)

and *LIF1* (Solyc02g079020, *Lignin induction Factor 1*, an *AP2*-like protein), that

regulate the biosynthesis of lignin in the cortex. Moreover, *CuRLR1* (Solyc04g009110, a

CC-NBS-LRR) may be engaged in signaling or function as a receptor for *C. campestris*,

leading to lignification-based resistance. Overexpression of *CuRLR1* in susceptible

tomato only induced strong lignification upon *C. campestris* attachment or *C. campestris*

extract injection. We also identified *SlWRKY16* (Solyc07g056280) as a negative regulator

of the lignin-based resistance. CRISPR-mediated mutants of *SlWRKY16* showed

lignification in the cortex and were more resistant to *C. campestris*. Furthermore, we

noticed that *CuRLR1* responds to a large protein molecule from *C. campestris* extracts,

which might represent potential novel *C. campestris* signals or effectors. In summary, we

discovered that these four key regulators control the lignin-based resistance response in

the stem cortex upon *C. campestris* infection which blocks *C. campestris* strands from

parasitizing selected Heinz tomato cultivars.

Results

Response to *C. campestris* in the resistant cultivars

While most tomato cultivars can be parasitized by *C. campestris*, the Heinz hybrids 9492 and 9553 (H9492 and H9553) exhibit resistance to dodders (Yaakov et al., 2001). *C. campestris* strands grew well on the susceptible H1706 (genome sequenced) and H9775 (Heinz hybrid 9775 – closely related to the resistant cultivars) (Figure 1A). On the other hand, *C. campestris* strands could not form good attachments with H9492 and H9553, and haustoria detached from the host stem, preventing parasite growth (Figure 1B). Based on the biomass ratio of *C. campestris* and host (relative growth rate), H9492 and H9553 cannot support long-term (>45 days) growth of *C. campestris*, in contrast to H9775 and H1706 (Figure 1C).

To identify the basis for resistance, we analyzed dodder attachment on susceptible and resistant lines using histology and cell wall-specific staining with Toluidine Blue O and Phloroglucinol-HCl (Liljegren, 2010). Upon challenging these different cultivars with *C. campestris* strands, lignin accumulation in the stem cortex was observed in the resistant H9492 and H9553, but not in the susceptible H9775 and H1706 (Figure 1D – 1L). The resistance mechanism involved local lignification in the stem cortex, creating a barrier to haustorium penetration, and dodder attachment on the resistant cultivars (Figure 1D – 1G). Little to no lignin accumulates in the cortex of both resistant and susceptible cultivars without *Cuscuta* attachment (Figure 1L). In addition, *Cuscuta* attachment sites usually cause some wounding responses and programmed cell death in both resistant and susceptible cultivars (Figure 1M).

Identifying the key time point in host-parasite interactions

Changes in the levels of Salicylic Acid and Jasmonic Acid have been reported at 36 to 48 hours after attachment (Runyon et al., 2010). To capture the earliest responses to dodder parasitism, we performed a time-course RNA-Seq analysis on 0, 1, 2, 3, and 4 DPA (days post attachment) of *C. campestris* on tomatoes H1706 (susceptible). At these stages, the dodder strands were not embedded in the host and could be removed to collect the attached stem area. Maximal transcriptional changes peaked at 4 DPA (Supplemental

Figure 1, Supplemental Data Set 1), suggesting that the Differentially Expressed (DE)

genes include core genes involved in the early response to *C. campestris* infection.

Accordingly, we chose 4 DPA for further gene expression analysis of resistant and

susceptible cultivars.

Gene expression in the resistant and susceptible host response to *C. campestris*

We challenged the resistant H9492 and H9553, and susceptible H9775 and H1706 with

C. campestris strands. We collected stem tissues at 4 DPA for RNA-seq and differential

gene analysis in dodder infested versus uninfested plants. In principal component analysis

(PCA) on the transcriptomes of resistant and susceptible cultivars (Supplemental Figure

2), PC1 accounted for 44% of the variation and significantly clustered the data into two

separate sets: infested and non-infested samples. However, PCA did not separate

different cultivars into distinct genotypic groups. Thus the transcriptional differences in

response to *C. campestris* likely involve a small number of genes.

Next, we conducted differential gene expression (DGE) analyses by comparing *C.*

campestris infested and uninfested host plants using an interaction design model (design

model = infested or uninfested condition + genotype + condition: genotype). Based on

our communication with the Kraft Heinz Company, both H9492 and H9553 were

developed in the same breeding program. However, H9553 is more resistant to *C.*

campestris than H9492 based on the relative growth rate results (Figure 1C). Therefore,

we suspected that enhanced resistance to *C. campestris* is due to the alterations in key

regulatory genes. We selected 113 genes that were differentially expressed (Supplemental

Data Set 2) between infested H9775 (susceptible) and infested H9553 (resistant) with

adjusted p value cutoff < 0.1 and log2 fold change > 1. Consistent with our observations

of lignin accumulation in resistant tomato cultivars upon *C. campestris* infestation

(Figure 1), many of these genes are known to be involved in the lignin biosynthetic

pathway, including three Laccase genes (LACs, Solyc05g052340, Solyc09g010990,

Solyc10g085090) and Caffeoyl CoA 3-O-methyltransferase (CCOMT, Solyc01g107910)

(Supplemental Figure 3). To narrow down the potential candidates regulating resistance,

we focused on transcription factors (TF - based on gene annotations) as possible key

regulators of lignin biosynthesis pathways, and membrane located or cytosolic receptors

which may receive signals from *C. campestris*. Using these two criteria, we selected three
candidate genes for further study, including a TF related to *AP2*, a *SlMYB55* TF*,* and a
gene encoding an N-terminal coiled-coil nucleotide-binding site leucine-rich repeat
protein (*CC-NBS-LRR*) (Supplemental Figure 3).

Transient overexpression of candidate genes using Virus-based Gene Expression
(VGE)
The Heinz cultivar is highly recalcitrant to transformation. The resistant lines are F1
hybrids in the Heinz background and show increased expression of these three candidate
genes upon *C. campestris* infestation. The susceptible cultivars show reduced expression
of these key regulators upon *C. campestris* infestation. To evaluate if the candidate genes
can confer lignification-based resistance in susceptible tomato cultivars, we cloned *GUS,*
AP2-like, *SlMYB55* and *CC-NBS-LRR* genes into Virus-based Gene Expression (VGE)
vectors (map in Supplemental Figure 1; sequence in Supplemental Data Set 9) for
transient overexpression in the susceptible H1706. We saw significant GUS expression in
the stem around the injection site (Supplemental Figure 4), and lack of lignification due
to the process of injection itself (Figure 2A and D). Therefore, we used *GUS*-injected
plants as our mock controls for VGE experiments. We sectioned and stained injected
stems with lignin-specific Phloroglucinol-HCl for lignin detection. VGE with *SlMYB55*
and *AP2*-like successfully overexpressed *SlMYB55* and *AP2*-like in the first internode
near the injection site and induced stem lignification in the susceptible H1706 (Figure
2B–2C and 2G). We named this *AP2*-like protein *LIF1* (*Lignin Induction Factor* 1) based
on the ability of *LIF1* (and *SlMYB55*) to induce lignin biosynthesis in the cortex.

In contrast, the H1706 plants with VGE of *CC-NBS-LRR* were very similar to
those with GUS VGE (Figure 2E). Previous studies indicated that many genes in the
NBS-LRR family encode intracellular receptors that detect pathogens and trigger defense
signaling (Padmanabhan et al., 2013). We suspected that this *CC-NBS-LRR* might also
play a role in signaling or function as a receptor for signals from *Cuscuta* that are needed
to initiate subsequent defense responses, such as lignin accumulation in the resistant
cultivars. Hence, we compared the response differences between *Cuscusta* infested and
uninfested susceptible H1706 with *CC-NBS-LRR* VGE (Figure 2E – 2F and 2H). Our

results showed the overexpression of *CC-NBS-LRR* only induced lignification upon *C.*
campestris attachment (Figure 2H), and suggest that direct or indirect perception of *C.*
campestris signals by this *CC-NBS-LRR* leads to lignification-based resistance. Thus, we
named this gene *CuRLR1* (*Cuscuta R-gene for Lignin-based Resistance 1*). Furthermore,
VGE of *SlMYB55*, *LIF1*, and *CuRLR1* with *C. campestris* all induced lignin accumulation
in the cortex and made the susceptible cultivar H1706 more resistant to *C. campestris*
(Figure 2I – 2L, Supplemental Figure 5, and Supplemental Data Set 3).

Lignin is a complex polymer and phloroglucinol-HCl staining is a fast and
efficient lignin detection method but it only detects the cinnamaldehyde end groups of
lignin, preferentially staining the G and S-type aldehyde form monolignols (Pomar et al.,
2002; Cass et al., 2015). Therefore, we also conducted an acetyl bromide assay to
determine total lignin content. Based on our results, the overexpression of *SlMYB55* and
LIF1 both increased total lignin content. Surprisingly, the overexpression of *CuRLR1*
slightly increased the total lignin content even without *Cuscuta* signals. With *Cuscuta*
signals, the total lignin content was much higher in *CuRLR1* overexpressing plants
(Figure 2M). We used HPLC to show that p-coumarate and trans-ferulate are both
increased in *CuRLR1* overexpressed plants, but the samples with *Cuscuta* signals have
much higher levels of these two precursors than the samples without *Cuscuta* signals
(Supplemental Figure 6). PYRO-GC analysis showed that samples from *CuRLR1*
overexpressing plants without *Cuscuta* signals have the largest percentage of H-lignin
and the largest concentration of coumarate derivatives (Figure 2N). H-lignin has been
correlated with both stress response as well as defense from pathogen intrusion because
this is a form of "defense" lignin that can be generated and deposited more rapidly than G
or S lignin (Zhang et al., 2007; Moura et al., 2010; Liu et al., 2018). Our results show that
CuRLR1 overexpression alone leads to an increase in the upstream steps of the lignin
biosynthesis pathway producing H-type monolignols, while *Cuscuta* signals may actually
up-regulate the final biosynthesis pathways leading to G-type and S-type monolignols
(Figure 2M and 2N). Since H-lignin and coumarate are not incorporated into lignin as
aldehydes they are not detected by phloroglucinol staining.

Regulatory mechanisms and networks leading to resistance responses

Since both H9492 and H9553 hybrid cultivars arose in the same breeding program, enhanced resistance to dodders observed in these two cultivars is likely due to the presence of some unique sequence polymorphisms in these cultivars. The resistance-specific nucleotide polymorphisms (SNPs) could contribute to the regulation or function of our candidate genes, so we specifically identified SNPs that are common in H9553 and H9492 but different from H9775 (Supplemental Data Set 4). There were no resistance-specific SNPs in coding regions of our candidate genes except one SNP located in a *LIF1* exon. This resistance-specific SNP changes 251 Lysine (K, in H1706) to 251 Glutamine (Q, in H9553). However, based on our protein domain prediction using InterProScan (Jones et al., 2014; Mitchell et al., 2018) and protein structure analysis using Phyre2 (Kelley et al., 2015), this amino acid replacement is not located in any known protein domains or structures. We also conducted PROVEAN (Protein Variation Effect Analyzer) analysis, and this K251Q variant only has a 0.619 PROVEAN score, indicating that this is a neutral variant. We therefore specifically focused on resistance-specific SNPs in the promoter regions of our candidate genes (Supplemental Data Set 5). Our SNP analysis detected several resistance-specific SNPs in the *LIF1* promoter region, but no resistance-specific SNPs were detected in other candidate gene promoter regions (< 5 kb upstream). One resistance-specific SNP was detected in the *CuRe1* promoter region (> 5 kb upstream) located at a putative YABBY binding site. However, this SNP is also located 1184bp upstream of *ULP1* (Solyc08g016275) and may regulate expression of this neighboring gene instead of *CuRe1*. Therefore, we focused on the *LIF1* promoter region and conducted transcription factor (TF) binding site predictions. Based on our phylogenetic network analysis (Solís-Lemus et al., 2017) using 500 kb around the LIF1 resistance-specific SNP enriched region, these SNPs might be introgressed from wild tomato species (likely coming from *S. galapagense* and/or *S. pennellii*, Supplemental Figure 7). One of these resistance-specific SNPs is located right at a WRKY binding W-box cis-element (TTGACY-core motif (Ciolkowski et al., 2008; Chen et al., 2019)) (Supplemental Figure 8 and Supplemental Data Set 6). This SNP is predicted to interrupt WRKY binding, likely leading to *LIF1* expression differences between resistant and

253   susceptible cultivars upon *C. campestris* attachment. Hence, we also focused on potential

254   WRKY TFs in our selected gene lists.

255   Based on our hypothesis, the genes that are differentially expressed upon *C.*

campestris attachment in both resistant cultivars may be regulated by the three candidate

genes that we selected. Therefore, to understand the relationships between the candidate

genes and their targets, we conducted DGE analysis with ANOVA and selected 10939

differentially expressed genes (DEG) with FDR less than 0.1 (Supplemental Data Set 7).

Next, we used Barnes-Hut t-distributed stochastic neighbor embedding (BH-SNE) to

generate gene clusters using RSMod (a pipeline developed by us, script included in code

availability) (Ranjan et al., 2016). In this analysis 5941 DEGs are clustered into 48

groups based on their gene expression patterns and 4998 DEGs are in the noise group.

Among the 48 gene clusters generated (Supplemental Data Set 8), five clusters were

selected based on their GO (Gene Ontology) enrichment terms and the candidate genes

they included (Figure 3A-B and Supplemental Data Set 8). The GO term of cluster 39 is

"DNA binding", which includes potentially key TFs, like *MYB55*. The GO term of

cluster 11 is "lignin biosynthetic and catabolic process", which encapsulates the observed

resistance phenotypes, and includes Caffeoyl CoA 3-O-methyltransferase (*CCOMT*) and

three Laccase (*Lac*) genes identified in our model based approach (Supplemental Data

Set 2). Cluster 23 includes a *Cinnamoyl-CoA reductase* gene (Solyc03g097170, *CCR*)

and is enriched in the "xyloglucan:xyloglucosyl transferase activity" GO term, which

may indicate potential cell wall modifications. "Response to biotic stimulus" is the GO

term enriched in cluster 17, which also includes a previous identified Cuscuta receptor,

CuRe1. Additionally, we noticed *SlWRKY16* is always clustered with *CuRe1*. We also

noticed that *SlWRKY16* was highly upregulated at 4 DPA in all four Heinz cultivars

(Supplemental Figure 9A). Host tissues surrounding haustoria from the tomato M82

cultivar also show upregulated expression of *SlWRKY16* at 4 DPA in our time-course

data with FDR < 0.1 and real-time qPCR data (Supplemental Figure 9B). Thus,

SlWRKY16 is a commonly upregulated host response gene across different cultivars and

may play an important role in the transduction of *C. campestris* signals upon host

attachment. Furthermore, one of the resistance-specific SNPs in the *LIF1* promoter region

mentioned above, is located at a WRKY transcription factor W-box (TTGACY-core
motif) binding site, which is also the predicted *SlWRKY16* binding site based on
homologous genes in the phylogenetic tree of the WRKY domain at the Plant
Transcription Factor Database (Jin et al., 2016). Taking all these criteria together, we
included *SlWRKY16* in our candidate genes for further analysis.

We focused on these 1676 genes in clusters 11, 17, 23, 39, 46 and included
CuRLR1 (Figure 3A-B and Supplemental Data Set 8) to construct gene co-expression
networks (GCNs) for different treatments and cultivars to identify central hub genes
(script included in code availability). Surprisingly, *CuRLR1*, *SlWRKY16* and *CuRe1* had
few connections or almost no connection with other genes in the GCN in susceptible
cultivars with *Cuscuta* attachments (Figure 3C-D). However, *CuRe1* and *SlWRKY16*
became central hub genes in resistant cultivars only upon *C. campestris* attachments and
connected with *CuRLR1* (Figure 3E-F, Supplemental Figure 10). Based on our DNA-Seq
analysis, we cannot detect any resistance-specific SNPs in the promoter regions or coding
regions of *CuRe1* and *CuRLR*1, and *SlWRKY16* (Supplemental Data Set 5). This result
indicates that the differential expression of *CuRe1* and *CuRLR*1, and *SlWRKY16* may be
controlled by trans-regulatory factors or protein interactions. Based on our GCN analysis
and DNA-Seq analysis results, we propose that all tomato cultivars have *Cuscuta*
receptors CuRe1 (Hegenauer et al., 2016) and CuRLR1, and SlWRKY16 is a key factor
in the transduction of *C. campestris* signals upon attachment of the parasite to the host.
However, the differential expression patterns and connections of these three genes
determine the different resistance responses in these Heinz susceptible and resistant
cultivars.

Functional characterization of *SlWRKY16* by CRISPR/Cas9 knockouts and VGE

To validate the function of *SlWRKY16* and its role in lignification-based
resistance, we produced stable *SlWRKY16* edited M82 lines using the CRISPR/Cas9
targeted gene knockout system (Pan et al., 2016). Our homozygous null mutants were
generally smaller than M82 wild type (Figure 4A and 4B) even though *wrky16* and M82
wild type are at the same developmental stage (Figure 4G). However, *wrky16* plants are
more resistant to *C. campestris* than M82 wild type (Figure 4C – 4F). Homozygous

wrky16 lines continuously produce cortical lignin and have a stronger resistance to *C.*
campestris attachment (Figure 4C – 4F and 4H).

Based on our SNP analysis and transcription factor binding site prediction results,
we propose the hypothesis that the resistance-specific SNP which is located at a WRKY
binding site in the *LIF1* promoter region could interrupt SlWRKY16 protein binding,
leading to *LIF1* expression differences between resistant and susceptible cultivars upon
C. campestris attachment. Therefore, we conducted qPCR analysis to determine the
expression levels of *LIF1* in both susceptible M82 wild type and *wrky16* tomatoes (M82
background). We observed an increase in *LIF1* expression in resistant *wrky16* tomatoes
compared to susceptible M82 tomatoes (Supplemental Figure 11A). Considering LIF1 is
an AP2/B3-like transcription factor, any elevation in *LIF1* expression could potentially
lead to large differences in the downstream gene expression pathways.

To evaluate the interaction between *SlWRKY16* and the other three candidate
genes, we transiently overexpressed *LIF1*, *SlMYB55*, *CuRLR1,* and *GUS* controls in the
susceptible H1706, M82 wild type and resistant *wrky16* tomatoes. In the *GUS* transient
overexpression control group, we observed that *wrky16* plants accumulate much more
lignin than H1706 and M82 wild type (Supplemental Figure 11B). Overexpression of
LIF1 induced more lignification in H1706, M82 and *wrky16* plants. This result shows
additive effects of loss of *SlWRKY16* function and overexpression of *LIF1* in lignification
responses (Supplemental Figure 11B), suggesting that *SlWRKY16* may not only regulate
LIF1 expression at the transcriptional level, but also may regulate LIF1 protein function
by other mechanisms. Also, overexpression of *MYB55* induced more lignification in
wrky16 plants but not in M82 (Supplemental Figure 11B), indicating that the loss of
SlWRKY16 function in M82 allow more lignin accumulation upon *MYB55*
overexpression. This result also suggests subtle differences between cultivars and that
SlWRKY16 may act upstream of *MYB55*.

However, overexpression of *CuRLR1* with *C. campestris* infection was able to
induce more lignification in M82, but not in *wrky16* tomatoes (Supplemental Figure 12).
This result suggests that either *CuRLR1* and *SlWRKY16* are in the same pathway with

WRKY16 downstream of CuRLR1, or that *CuRLR1* and *SlWRKY16* are in two
independent pathway that may influence each other. This hypothesis matches with the
gene coexpression networks we built, which show that *CuRLR1* and *SlWRKY16* are
peripherally positioned in resistant cultivars in the *Cuscuta* treated condition, with
multiple layers of genes connecting them. In order to elucidate other layers of regulation
between these genes, we conducted protein-protein interaction investigations.

Subcellular localization and interactions between the candidate proteins

One described mechanism for triggering innate immunity following TMV
infection in tobacco involved interaction and subsequent nuclear localization of the SPL6
TF with the TIR-NBS-LRR receptor (Padmanabhan et al., 2013; Padmanabhan and
Dinesh-Kumar, 2014). Therefore, we investigated the potential interactions between our
candidates and their protein subcellular localization to uncover potential regulatory
mechanisms. Based on our results using translational GFP fusions, LIF1 and SlWRKY16
are located mainly in the nucleus (Figure 5A), while *CuRLR1* is located in both the
nucleus and the cytosol. Bimolecular fluorescence complementation (BiFC) experiments
with split YFP using transient infiltration in *N. benthamiana* leaves show that the LIF1
and SlWRKY16 proteins interact and get localized to the cytoplasm (Figure 5B).
Interactions between other combinations, CuRLR1-LIF1, CuRLR1-SlWRKY16, or
CuRLR1-CuRe1, were not detected in our experiments. To verify the interaction between
LIF1 and SlWRKY16 proteins and their subcellular localizations when they interact with
each other, we also co-expressed the fusion proteins LIF1-GFP and SlWRKY16-RFP to
observe their subcellular localizations. We found that GFP and RFP signals are located
mainly in the nucleus when we only overexpress LIF1-GFP or SlWRKY16-RFP in
separate *N. benthamiana* leaves. However, when we co-express LIF1-GFP and
SlWRKY16-RFP in the same leaves, GFP and RFP signals mostly overlap in the
cytoplasm (Figure 5C). These results not only further confirm that LIF1 and SlWRKY16
proteins may interact with each other and stay in the cytosol, but also validate our
hypothesis that SlWRKY16 can regulate LIF1 expression at both the transcriptional and
protein interaction levels.

Analysis of the *Cuscuta* signal using *Cuscuta* extract injections

To further discern the nature of the major signals that trigger lignification-based resistance, we injected the first internode of the resistant H9553 with *Cuscuta* extracts subjected to different treatments (Figure 6). Untreated or filtered *Cuscuta* extract injections induced the accumulation of lignin in the cortex region (Figure 6B-C). On the other hand, alteration of *Cuscuta* extract pH from 5.8 to 9 abolished lignin accumulation (Figure 6D-E), suggesting either instability or sequestration of the *Cuscuta* signaling molecules in alkaline conditions. In addition, heat-treated extract and proteases-treated extract could not trigger the lignification response (Figure 6F-J). Moreover, *Cuscuta* extract injections also induced lignin accumulation in H1706 with VGE overexpressing *CuRLR1*, but not in H1706 (Supplemental Figure 13). This result indicates that *CuRLR1* may be able to either sense some unknown factors in Cuscuta extract or some part of the response to these factors and lead to lignin-based resistant responses. Furthermore, filtration of extracts through devices with different molecular weight cutoffs indicates that fractions smaller than 30kD cannot trigger strong lignification response (Supplemental Figure 14). Thus, the active *Cuscuta* signal for induction of lignin-based resistance is larger than 30kD but smaller than 100kD, and distinct from the previously identified *Cuscuta* signal that binds *CuRe1* (Hegenauer et al., 2016).

Discussion

Cuscuta spp. cause massive loss in infested tomato fields in the United States. Our study reveals the underlying molecular genetic mechanisms for lignin-based resistance responses in resistant tomato cultivars. Lignin is a complex phenolic polymer, which is generated from three major monolignols, paracoumaryl alcohol, coniferyl alcohol, and sinapyl alcohol, using covalent crosslinks formed via free radical polymerization (Ferrer et al., 2008). Accumulation of lignin in plant stems or roots has been shown to reinforce plant resistance to invading herbivores, parasites and pathogens (Reimers and Leach, 1991; Gayoso et al., 2010; Taheri and Tarighi, 2012; War et al., 2012; Dhakshinamoorthy et al., 2014; Kumari et al., 2016; Zhang et al., 2019). Lignification at the host-parasite interface in roots has been reported in plants that are resistant to root parasitic plants (Goldwasser et al., 1999; PÉRez-De-Luque et al., 2005; Cameron et al., 2006; Lozano-Baena et al., 2007). However, for stem parasitic plants, most research has focused on hypersensitive response or necrosis as the major mechanisms for host plant defense (LANE et al., 1993; Hegenauer et al., 2016; Su et al., 2020). A few reports of incompatible reactions between tomato plants and *Cuscuta reflexa* characterized by a visible brownish plaque at infection sites, suggested this might be due to suberized or lignified cell walls (Sahm et al., 1995). However, previous studies also indicate that different *Cuscuta* species can have diverse host-parasite interactions with the same host species (Ranjan et al., 2014; Kaiser et al., 2015; Hegenauer et al., 2016). For example, although cultivated tomatoes (*S. lycopersicum*) are generally resistant to *Cuscuta reflexa* (Sahm et al., 1995; Hegenauer et al., 2016), most domesticated tomato cultivars are in considered susceptible to *C. campestris*. Therefore, using tomato cultivars that have been bred for resistance to *C. campestris* could help us reveal the multilayered resistance mechanisms to *Cuscuta* spp. and will aid in developing parasitic plant-resistant crops. Here, we identified a strong lignin-based resistance response toward *C. campestris* attack in specific Heinz tomato cultivars, adding another layer on the previous reported hypersensitive-type response mechanism.

Three key genes, *LIF1*, *SlMYB55*, and *CuRLR1* regulate lignin accumulation in the cortex. Of these, *CuRLR1* responded to *Cuscuta* signals and further reinforced lignin

deposition in the resistant cultivars. The *Cuscuta* signals that trigger the lignin-based defense responses appear to be large heat-sensitive proteins (30kD-100kD, Figure 6), and distinct from the previously identified small modified peptide *Cuscuta* signal (~2kD) that is recognized by *CuRe1*, an LRR receptor-like serine/threonine-protein kinase (RLP) (Hegenauer et al., 2016). In conclusion, we propose a new multilayered model for *Cuscuta* resistance response in tomato (Figure 7). *CuRLR1* is a cytosolic factor, which either receives large signaling molecules from *C. campestris* as a receptor, or may be a novel factor which plays a role in downstream signal transduction upon sensing *Cuscuta* signals and triggering a lignin-based resistance response (Figure 7, red-labeled pathway). Based on previous studies, NBS-LRRs are usually located in the cytoplasm and nucleus and likely to recognize pathogen effectors to induce effector-triggered immunity (ETI) (Dodds and Rathjen, 2010). This matches want our observed CuRLR1 subcellular localization (Figure 5) and might also explain why the *Cuscuta* signals that trigger the lignin-based defense responses are in different size range compared with the previously identified *Cuscuta* signal. Our research results shed light on a potential ETI pathway in parasitic plant resistance and provides the foundation for further future studies into how these various layers of resistance connect.

Transient overexpression of *SlMYB55* and *LIF1* induced lignin accumulation in the cortex, placing *SlMYB55* and *LIF1* as positive regulators in the lignin biosynthesis pathway (Figure 7, pink and yellow-labeled pathway). Other yet undiscovered *Cuscuta* receptors or factors may induce *SlMYB55* and *LIF1* expression upon *Cuscuta* attachment. On the other hand, *wrky16* plants showed lignin accumulation and stronger resistance to *Cuscuta*, suggesting that *SlWRKY16* is a negative regulator of this lignin-based resistance pathway (Figure 7, green-labeled pathway). Based on our DNA-Seq, BiFC, and subcellular localization data (Figure 5, Supplemental Figure 8), we propose that *SlWRKY16* regulates the function of *LIF1* by a combination of inhibition of *LIF1* transcription and physical capture of LIF1 proteins to block their entry into the nucleus (Figure 7, yellow and green-labeled pathway). *CuRe1* is reported to mediate PAMP/MAMP-triggered immunity (PTI/MTI) (Hegenauer et al., 2016) (Figure 7, blue-labeled pathway). GCN analysis indicates a coexpression connection between *CuRe1* and

SlWRKY16 (Figure 3A-D). *CuRe1* and *SlWRKY16* both became central hub genes in
resistant cultivars upon *Cuscuta* attachments (Figure 3D), suggesting the hypothesis that
SlWRKY16 may act downstream of *CuRe1* (Figure 7, blue-labeled pathway). Thus, we
envision crosstalk between different resistance pathways that may be triggered together
to enhance host defense responses.

We systemically investigated the lignin-based resistance response in crop plants
toward stem parasitic plants and our finding has potential implications for enhancing crop
resistance to parasitic plants. Notably, overexpression of the CuRLR1 protein induced
upregulation of lignin precursors, but extensive lignin accumulation was only triggered
by *Cuscuta* extracts. This indicates that introducing high level of CuRLR1 protein would
provide resistance to *C. campestris*, but would not trigger the crop to continuously spend
a lot of resources in producing large amount of cortical lignin and stunted growth. The
identification of *CuRLR1* might provide a path forward to introduce resistance into other
important crops that are also attacked by *C. campestris*.

Materials and Methods

<u>Plant materials used in the study</u>

We obtained four different cultivars from Dr. Rich Ozminkowski at HeinzSeed,
including the Heinz hybrid cultivars 9492 and 9553 (H9492 and H9553), and the related
susceptible Heinz hybrid cultivar 9775 (H9775) and the sequenced susceptible Heinz
cultivar 1706 (H1706). Our *Cuscuta* was originally collected from tomato field in
California, and we obtained seeds from W. Thomas Lanini. This *Cuscuta* was previously
identified as *Cuscuta pentagona* (Yaakov et al., 2001), which is a closely related species
to *Cuscuta campestris*(Costea et al., 2015b). To clear up the confusion, we extract DNA
to verify species by molecular phylogenetics. Based on phylogenetic analysis of plastid
trnL-F, rbcL sequences, and nrITS, nrLSU sequences (Stefanović et al., 2007; García et
al., 2014; Costea et al., 2015b), we confirmed that our experimental species is *Cuscuta*
campestris (Supplementary Figure 9-13). According to our results, our *Cuscuta*
campestris isolate is most similar to *Cuscuta campestris* 201 voucher Rose 46281 WTU
from USA, CA (Supplemental Figure 15-19) that is published by Costea *et al.* in 2015
(Costea et al., 2015a).

<u>Histology and Cell Wall-Specific Staining</u>

For preparing the sections at the *C. campestris* attachment area and Agroinjection
sites on tomato stems, we hand-sectioned plants at 200 to 500 μm thickness using razor
blades and kept these sections in 4°C water before staining. For preparing the sections of
haustoria attached to hairy roots, we fixed samples in 7% Plant Tissue Culture Agar and
used Lancer Vibratome Series 1000 to prepare 100 μm sections and kept these sections in
4°C water.

For Phloroglucinol-HCl Staining, we followed the published protocols (Liljegren,
2010; Pradhan Mitra and Loqué, 2014) with some modifications. To prevent plasmolysis
during staining, we added an ethanol dehydration process before staining, which is
described as follows: we removed the water and then immersed sections in cold 30%
ethanol and then 60% ethanol for 5 minutes each. We prepared phloroglucinol-HCl stain

(Ph-HCl) or Wiesner stain by preparing a 2:1 mixture of 100% EtOH and concentrated HCl and dissolving powdered phloroglucinol into this solution at a final concentration of 3% w/v. After removing the 60% ethanol, we added phloroglucinol-HCl solution dropwise to the Petri dishes, and let the sections sit in the stain for 5 minutes. The lignified areas of the sections stain bright red within 30 seconds of immersion in the stain. After removing the phloroglucinol-HCl and adding 60% ethanol back, we imaged the sections in the petri dish on a white background using a Zeiss SteREO Discovery, V12 microscope and Nikon Eclipse E600 microscope.

For Toluidine Blue O Staining, we used a published protocol with some modifications (O'Brien et al., 1964). We immersed the sections in the stain for 30 seconds, and then washed with water three times for 30 seconds each. After removing the agar from around the sections using forceps, we mounted the sections with water on a slide and imaged using a Nikon Eclipse E600 microscope.

Image analysis of stem and haustorium sections

To quantify the lignin content of each section, we analyzed images using the image processing software ImageJ (Schneider et al., 2012). We added a Gaussian blur with a sigma radius of 2.00 to reduce image noise. We set the color space of the image to L*a*b* to generate histograms that measure lightness, green-red contrast, and blue-yellow contrast of the image. We adjusted the lightness filter to allow histogram coordinates ranging from zero to the peak of the image histogram, and the green-red filter to allow from the histogram peak to 255, and the blue-yellow filter to allow all histogram coordinates. These coordinates filter for red areas on the image, corresponding to lignified areas in the stem sections. We measured the total area of lignification, then selected areas corresponding to the lignified xylem of the stem and measured this area. We subtracted the xylem area from the total lignin area to calculate the cortex lignin area.

DNA-seq library construction for resistant and susceptible Heinz cultivars

DNA was extracted from the leaves of three weeks old seedlings using GeneJET Plant Genomic DNA Purification Mini Kit (Thermo Scientific, Waltham, MA, USA)

DNA-Seq libraries were prepared using an in-house protocol modified from Breath

Adapter Directional sequencing (BrAD-seq) (Townsley et al., 2015). First, 5 µg of

genomic DNA was fragmented using a Covaris E220 (Covaris, Inc. Woburn, MA, USA)

with the following settings: Peak Incident Power (W) 140; Duty Factor 10%; Cycles per

Burst 200 and Treatment Time (s) 90 to obtain an average fragment size of 400 base

pairs. Next, the fragmented DNA was end-repaired and A-tailed in a single reaction using

DNA End Repair Mix and Taq DNA polymerase (New England Biolabs). Y-type

adapters were ligated and an enrichment PCR was performed with as in BrAD-seq

(Townsley et al., 2015) using 7 cycles. Individual libraries were quantified by PCR and

pooled to equaled amounts. After a final library cleanup with AMPure beads (Beckman

Coulter, Brea, CA, USA), DNA-seq libraries were sequenced at the California Institute

for Quantitative Biosciences (QB3) at the University of California, Berkeley using the

HiSeq 4000 platform at 150 Paired Read (PR). (Illumina Inc. San Diego, CA, USA).

<u>Resistant and susceptible Heinz cultivar DNA-seq SNP analysis, promoter binding site</u>

<u>analysis, and protein domain and structure prediction</u>

For single nucleotide polymorphism (SNP) analysis, we mapped DNA-seq read

data to sequenced H1706 tomato genome itag 3.0 to identify SNPs using CLC Genomics

Workbench 11 (QIAGEN, https://www.qiagenbioinformatics.com/). Next, we compared

the SNPs across the resistant and susceptible tomato cultivars and focus on finding the

SNPs that are common in H9553 and H9492 but different from H9775. In other words,

we focus on the SNPs that exist in resistant cultivars and named these SNPs as "resistant

specific SNPs" (Supplemental Data Set 4). Among our 4 candidate genes, *LIF1* was the

only gene that has resistant specific SNPs in the promoter region (Supplemental Data Set

5). In order to identify potential introgression regions, we conducted phylogenetic

network analysis using the PhyloNetworks package (Solís-Lemus et al., 2017) in the Julia

environment on XSEDE (Dahan et al., 2014) with 500 kb of sequence around the LIF1

resistance-specific SNP enriched region (Supplemental Figure 7). To identify potential

transcription factor binding sites in the *LIF1* promoter region, we used PlantPAN 3.0

(http://PlantPAN.itps.ncku.edu.tw) (Chow et al., 2015; Chow et al., 2018) "TF/TFBS

Search" and "Promoter Analysis". We also predicted the *SlWRKY16* binding site based

on the homologous genes in the phylogenetic tree of WRKY domains on the Plant
Transcription Factor Database (PlantTFDB v5.0, http://planttfdb.gao-lab.org;
Phylogenetic Tree for Solanum lycopersicum WRKY Family: http://planttfdb.gao-
lab.org/phylo_tree.php?sp=Sly&fam=WRKY) (Jin et al., 2016). To determine the
consequences of the K251Q amino acid replacement in the LIF1 protein, we predicted
potential protein domains of LIF1 using InterProScan
(https://www.ebi.ac.uk/interpro/search/sequence/) (Jones et al., 2014; Mitchell et al.,
2018). We conducted protein 3D structure prediction and analysis using Phyre2 (Protein
Homology/analogY Recognition Engine v 2.0,
http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index) (Kelley et al., 2015). We
also predicted whether K251Q amino acid substitution is deleterious or neutral using
PROVEAN (Choi, 2012; Choi et al., 2012) (Protein Variation Effect Analyzer,
http://provean.jcvi.org/seq_submit.php).

Timecourse RNA-Seq library construction and analysis

We challenged H1706 tomato cultivars with strands of *C. campestris* and
collected stem tissues at 1, 2, 3, 4 days post attachment (DPA) and 0 DPA as negative
controls. Following this, we constructed strand-specific poly-A based libraries for RNA-
seq (Townsley et al., 2015). We conducted sequencing of these libraries on two lanes on
Illumina HiSeq 2000 at 50bp Single Read (SR).

We used using CLC Genomics Workbench 11 (QIAGEN) for following RNA-seq
analysis. First, we mapped resistant and susceptible cultivar RNA-seq read data to
sequenced H1706 tomato genome itag 3.0. To see the general pattern across libraries, we
conducted principal component analysis (PCA) of gene expression across different DPA.
Next, we used ANOVA comparison with DPA factors and cutoff FDR < 0.1 to select
differentially expressed genes (DEG) list. Then, we drew Venn diagrams of DEGs at
different DPA libraries. 0 DPA libraries are without *Cuscuta* treatments and serve as the
negative control for comparisons. The cutoff of these DEGs are FDR < 0.1 and fold
change > 1.5. Following, we constructed a heat map of DEGs across different DPA

libraries. Euclidean distance and complete linkage are used for this clustering analysis
(Supplemental Figure 1).

<u>Resistant and susceptible cultivar RNA-Seq library construction and interaction model-</u>
<u>based analysis</u>

We challenged the resistant and susceptible tomato cultivars with strands of *C.*
campestris and collected stem tissues at 4 days post attachment (DPA). Following this,
we constructed strand-specific poly-A based libraries for RNA-seq from the four tomato
cultivars, including the resistant cultivars H9492 and H9553, and the susceptible cultivars
H9775 and H1706. We conducted sequencing of these libraries on two lanes on Illumina
HiSeq 4000 at 100bp Single Read (SR). We first mapped reads to sequenced H1706
tomato genome itag 2.4. To investigate gene expression changes across the resistant and
susceptible cultivars in dodder infested versus uninfested plants, we conducted a PCA
with K-means clustering using the normalized read counts of sequences mapped to the
tomato transcriptome (Figure 3). Next, we defined differentially expressed genes with the
Bioconductor package DESeq2 employing an interaction design (design = ~ Condition +
Genotype + Condition: Genotype). Following this, we focused on these 113 genes that
display expression changes upon dodder infestation that are different in H9492 and
H9553 compared to H1706 and H9775. Within these 113 genes, we picked our three
candidate genes based on gene annotation and functions.

<u>Barnes-Hut clustering analysis and gene coexpression network analysis for resistant and</u>
<u>susceptible cultivar RNA-Seq</u>

In order to get a more comprehensive differentially expressed gene (DEG) list, we
mapped resistant and susceptible cultivar RNA-seq read data to sequenced H1706 tomato
genome itag 3.0 by using CLC Genomics Workbench 11 (QIAGEN). Next, we used
ANOVA comparison with both factors, all cultivars and with/without *Cuscuta* treatments,
and cutoff FDR < 0.1 to select DEG list. In these 10939 genes, we applied Barnes-Hut t-
distributed stochastic neighbor embedding (BH-SNE) using RSMod package (script
included in code availability) generated 85 gene clusters based on their gene expression
patterns. Based on their GO (Gene Ontology) enrichment terms and their included

candidate genes, five clusters were selected for further analysis (Supplemental Data Set 8, yellow-labeled genes). We use these selected genes to build gene coexpression networks by using the R script (script included in code availability) that was modified from our previously published method (Ichihashi et al., 2014). We constructed gene coexpression networks for different *C. campestris* treatments in susceptible and resistant cultivars with normal quantile cutoff = 0.997.

Virus-based Gene Expression (VGE) in tomatoes

For preparing the binary vector for plant transient expression that carries the ToMoV DNA, we used a modified pSP72-TAV (Gilbertson et al., 1993; Hou and Gilbertson, 1996) that is lacking the coat protein ORF (CP) and have a restriction enzyme multisite in which a Gateway® cassette (Thermo Fischer Scientific) was cloned by In-Fusion (Takara) in the *NcoI* site. The whole replicon fragment of TAV-GW was amplified from this vector and cloned into a binary vector to generate a vector for Agrobacterium-mediated transient expression in plants. Since the ToMoV DNA-B (Carrying the viral movement protein (MP) and the CP are missing, this clone is not an infectious clone and is only serves as a viral replicon by replicating via rolling circle mechanism (Stenger et al., 1991).

For transiently expressing our candidate genes in the susceptible tomato cultivar H1706, we used this Virus-based Gene Expression (VGE) vector pTAV (Supplemental Figure 20). We cloned *GUS*, *LIF1*, *SlMYB55* and *CuRLR1* genes into pTAV and transformed these into thermo-competent *Agrobacterium tumefaciens* by heat-shock-transformation. For culturing Agrobacterium and preparing agroinjection, we followed the previously published protocol (Vel et al., 2009) with some modifications. For each experiment, we started from growing transformed Agrobacterium on Lysogeny broth (LB) agar plates with appropriate antibiotic selections at 30° C for 2 days. Following this, we inoculated 10 mL liquid LB with transformed Agrobacteria (AGL1) and incubated at 30° C for 16 hours with 200 r.p.m. shaking. We diluted the primary cultures 1:5 into Induction Media (Vel et al., 2009) supplemented with appropriate antibiotic selections and 200 µM acetosyringone, and then incubated them at 30° C for 24 hours with 200

r.p.m. shaking. When the O.D. 600 of the culture was around 1, we harvested the

transformed Agrobacteria by centrifuging at 3000 x g for 10 minutes and then

resuspended Agrobacteria in Inoculation Buffer (10 mM 2-[N-Morpholino] ethane

sulfonic acid (MES), 10 mM MgCl$_2$, 200 µM acetosyringone and 0.5 mM dithiothreitol)

to an O.D. 600 of 1 culture. Next, we injected this transformed Agrobacterium culture

into the first internode of tomato stems using a syringe equipped with a 0.8 mm x 38.1

mm MonoJect needle.

Preparation of *C. campestris* extracts and injection protocols

For one mL *C. campestris* extracts, we collected 100 mg of the stem tissue in

microcentrifuge tubes from *C. campestris* growing on H1706 tomato plants. We used the

BioSpec Mini-Beadbeater to grind the liquid nitrogen-frozen tissue with five 2.3 mm

diameter BioSpec zirconia beads and 1.0 mm diameter BioSpec zirconia beads in the

tubes for 1 minute, and then mixed with one mL deionized water. To remove the plant

tissue debris, we centrifuged extracts for 30 seconds at 5000 r.c.f., and used only the

supernatant for untreated extract injections. For heat-treated extracts, we heated at 95 °C

for 5 minutes. For pH treated extracts, we adjusted the pH to 9 by adding 0.1M NaOH.

For filtered extracts, we filtered untreated extracts through a VWR 0.2 µm sterile syringe

filter. We injected different treated extracts into the first internode of tomato stems using

a syringe equipped with a 0.8 mm x 38.1 mm MonoJect needle. Furthermore, we used

3K, 10K, 30K, and 100K Amicon® Ultra Centrifugal Filter Devices to filter *Cuscuta*

extracts. Then, we use flow through extracts to do injection on H9553 stems to test the

size of *Cuscuta* signals.

Protein interaction, subcellular localization and co-localization of fusion candidate

proteins

For protein interaction assays, we performed *in vivo* using bimolecular

fluorescence complementation (BiFC) system (Kerppola, 2008). The plasmids were

constructed by using the Gateway-compatible BiFC vectors SPDK1794 (*p35S::cCitrine*)

and SPDK1823 (*p35S::nCitrine*). The leaves of four-week-old *Nicotiana (N.)*

benthamiana were injected with different combinations of *Agrobacterium (A.)*

tumefaciens GV3101 containing the transient expression vectors (Fang and Spector,

2010).

For subcellular localization and co-localization of fusion candidate proteins, we

performed with transient expression fluorescent fusion proteins *in vivo*. The plasmids

were constructed by using the Gateway-compatible vectors pGWB5 (*p35S::GFP*) and

pGWB660 (*p35S::TagRFP*). The leaves of four-week-old *N. benthamiana* were injected

with the *A. tumefaciens* GV3101 strain containing one of the plasmids (Sparkes et al.,

2006). To verify the interaction between LIF1 and SlWRKY16 proteins and determine

their subcellular localizations when they interact with each other, we also co-expressed

the fusion proteins LIF1-GFP (*p35S::LIF1-GFP*) and SlWRKY16-RFP

(*p35S::SlWRKY16-TagRFP*). To reduce gene☐silencing and enhance transient expression

of our candidate proteins, we co-expressed the fusion proteins with p19 obtained from

Professor Bo Liu's Lab at University of California, Davis. Three individual plants and

three adult leaves of each plant were used for each treatment. Fluorescence was observed

2 – 5 days after transfection by a Confocal Laser Scanning Platform Zeiss LSM710

(Zeiss, Germany).

<u>Lignin content and composition analysis</u>

Tomato stem material from specified treatments was flash-frozen in liquid

nitrogen and lyophilized at -50°C and ≤ 0.1 mbar for 48 hours in a 6L FreeZone 6

Benchtop Freeze Dry System (Labconco Corp., Kansas City, MO, USA). Lyophilized

material was pulverized in a 2ml screw top microcentrifuge tube with a glass bead for 10

minutes at a frequency of 20/s in a TissueLyser (Retsch ballmill, Qiagen, Venlo,

Netherlands). The material was AIR prepped and destarched using protocols from Barnes

and Anderson (Barnes and Anderson, 2017). Acetyl bromide analysis was carried out

using the same protocol with one exception: samples were incubated at 50°C with gentle

swirling every 10 minutes for 2 hours to limit the degradation of xylan, which can lead to

the over quantification of lignin present in the sample. Lignin quantification reactions

were performed in a 10 mm light path quartz cuvette (VWR, Radnor, PA, USA, catalog

number 414004-062) and absorbance measurements were taken on a SPECTRAmax plus

384 (Molecular Devices, San Jose, CA, USA). %ABSL was calculated using the extinction coefficient of 17.2 (Chen and Dixon, 2007). Presently, no extinction coefficient is available for tomato stem lignin so the extinction coefficient for tobacco stem lignin was used. Cell wall-bound aromatics and HPLC analysis of liberated aromatics was performed as described by Eudes et al. (Eudes et al., 2015). The remaining cell wall material post-aromatic extraction, now enriched for lignin, was then washed with water 3 times and dried at 30°C overnight. The chemical composition of the lignin enriched cell wall material was analyzed by pyrolysis-gas chromatography (GC)/mass spectrometry(MS) using a previously described method with some modifications (Eudes et al., 2015). Pyrolysis of biomass was performed with a Pyroprobe 6200 (CDS Analytical Inc., Oxford, PA, USA) connected with GC/MS (GCMS-QP2010 Ultra Gas Chromatograph Mass Spectrometer, Shimadzu corp., Kyoto, Japan) equipped with an AgilentHP-5MS column (30 m 9 0.25 mm i.d., 0.25 lm film thickness). The pyrolysis was carried out at 550 °C. The chromatograph was programmed from 50 °C (1 min) to 300 °C at a rate of 30 °C/min; the final temperature was held for 10 min. Helium was used as the carrier gas at a constant flow rate of 1 mL/min. The mass spectrometer was operated in scan mode and the ion source was maintained at 300 °C. The compounds were identified by comparing their mass spectra with those of the NIST library and those previously reported (Ralph and Hatfield, 1991; Gutiérrez et al., 2006). Peak molar areas were calculated for the lignin degradation products, and the summed areas were normalized per sample.

Data availability

All data is available in the main text or the supplementary materials. All DNA-Seq and RNA-Seq raw data are deposited on NCBI SRA PRJNA550259.

Code availability

All R scripts and package for analysis are deposited on GitHub. R script for RNA-Seq interaction model-based analysis deposited on GitHub (Link: https://github.com/MinYaoJhu/Moran-s-RNA-Seq-analysis-script). R script and RSMod package for Barnes-Hut clustering analysis deposited on GitHub (Link: https://github.com/sdrowland/RSMod). R script for RNA-Seq gene coexpression network

723   analysis deposited on GitHub (Link: https://github.com/Hokuto-GH/gene-coexpression-

724   network-script).

Author Contributions

M.F. initiated the project with supervision from N. S. and established the connection with

Heinz. M.-Y.J. conducted phylogenetic analysis and verified *Cuscuta* species. M.-Y.J.,

M.F., and R.N.P. conducted histochemical and biomass analysis. M.R. modified the VGE

vector and M.F. cloned candidate genes into VGE vectors. M.-Y.J., R.N.P., and L.W.

conducted VGE overexpression experiments and analysis for all candidate genes. S.D.R.

and M.-Y.J. developed lignin section area quantification methods. M.-Y.J. produced

VGE overexpressed samples and M.S.B. performed acetyl bromide, HPLC and Pryo-GC-

MS analysis with P.M.S.'s supervision. M.F. made DNA-Seq and RNA-Seq libraries.

M.F. conducted preliminary SNP analyses and model-based DEG analyses and picked

primary candidate genes. M.-Y.J. performed SNP analyses, protein domain and structure

prediction, PROVEAN analysis for DNA-Seq data, and ANOVA DEG analyses, BH-

SNE clustering to select candidate genes and built GCNs for RNA-Seq data. S.D.R.

developed the RSMod R package. H.N., S.D.R., and M.-Y.J. modified the GCN analysis

script and M.-Y.J. construct GCNs. M.-Y.J. and S.D.R. performed PhyloNetwork

analysis. M.-Y.J. and L.W. conducted *wrky16* plant phenotyping and VGE

overexpression experiments on *wrky16*. M.-Y.J. and L.W. performed subcellular

localization and BIFC experiments. M.-Y.J. performed LIF1-GFP and SlWRKY16-RFP

co-localization experiments. M.-Y.J., R.N.P., and L.W. conducted *Cuscuta* extract

injection experiments. M.-Y.J. wrote the initial manuscript draft for the main text and

made figures and tables with primary editing from N. S.. M.-Y.J., R.N.P., L.W., M.S.B.,

M. R., and H.N. wrote the initial draft for materials and methods. M.-Y.J., M.F., R.N.P.,

L.W., M.S.B., H.N., K.Z., P.M.S., and N.R.S. contributed to the editing of this

manuscript. N.R.S. supervised the whole project.

Acknowledgments

751   We are grateful to S. Dinesh-Kumar, A. B. Britt, S. Brady, and D. Runcie for their input

752   on this research, and S. Dinesh-Kumar, A. B. Britt, S. Brady, and E. D. Marable for

editing suggestions. We thank C. Wong, C. Ito, Z. Jaramillo, and J. Lu for helping with some parts of experiments or image analysis, and Axtell Lab and N. Johnson for providing information to help with verifying *Cuscuta* species, and Liu Lab and Y. Lee for providing p19, pGWB5 and pGWB660 plasmid. We also thank R. Ozminkowski, Manager of Agricultural Research from HeinzSeed for providing seeds of three Heinz cultivars, and the UC Davis Plant Transformation Facility for generating transgenic plants, and CyVerse for online data storage. This work used the Extreme Science and Engineering Discovery Environment (XSEDE) on the Pittsburgh Supercomputing Center (PSC) at the Regular Memory (Bridges) servers through allocation IBN200015. We thank A. Eudes for his recommendations and guiding conversations in lignin analysis, and J. Ortega for her technical support with the pyro-GC/MS instrument. **Funding:** This work was funded by USDA-NIFA (2013-02345). M. J. was supported by Yen Chuang Taiwan Fellowship, Taiwan Government Scholarship (GSSA), Elsie Taylor Stocking Memorial Fellowship, and Katherine Esau Summer Graduate Fellowship program; M.F. was supported by United States-Israel Binational Agricultural Research and Development Fund (postdoctoral fellowship no. FI–463–12), M.S.B. and P.M.S. were supported by the DOE Joint BioEnergy Institute (http://www.jbei.org) supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy.

Competing interests: The authors declare that they have no competing interests.

References

Agrios, G.N. (2005). chapter thirteen - PLANT DISEASES CAUSED BY PARASITIC HIGHER PLANTS, INVASIVE CLIMBING PLANTS, AND PARASITIC GREEN ALGAE. In Plant Pathology (Fifth Edition) (San Diego: Academic Press), pp. 705-722.

Ashton, F.M. (1976). Cuscuta Spp. (Dodder): A Literature Review of Its Biology and Control. (Division of Agricultural Sciences, University of California).

Barnes, W.J., and Anderson, C.T. (2017). Acetyl Bromide Soluble Lignin (ABSL) Assay for Total Lignin Quantification from Plant Biomass. Bio-protocol **7,** e2149.

Barros, J., Serk, H., Granlund, I., and Pesquet, E. (2015). The cell biology of lignification in higher plants. Annals of botany **115,** 1053-1074.

Bellincampi, D., Cervone, F., and Lionetti, V. (2014). Plant cell wall dynamics and
wall-related susceptibility in plant–pathogen interactions. Frontiers in Plant
Science **5**.
Bhuiyan, N.H., Selvaraj, G., Wei, Y., and King, J. (2009). Role of lignification in plant
defense. Plant signaling & behavior **4,** 158-159.
Cameron, D.D., Coats, A.M., and Seel, W.E. (2006). Differential Resistance among
Host and Non-host Species Underlies the Variable Success of the Hemi-parasitic
Plant Rhinanthus minor. Annals of Botany **98,** 1289-1299.
Cass, C.L., Peraldi, A., Dowd, P.F., Mottiar, Y., Santoro, N., Karlen, S.D.,
Bukhman, Y.V., Foster, C.E., Thrower, N., Bruno, L.C., Moskvin, O.V.,
Johnson, E.T., Willhoit, M.E., Phutane, M., Ralph, J., Mansfield, S.D.,
Nicholson, P., and Sedbrook, J.C. (2015). Effects of PHENYLALANINE
AMMONIA LYASE (PAL) knockdown on cell wall composition, biomass
digestibility, and biotic and abiotic stress responses in Brachypodium. Journal of
Experimental Botany **66,** 4317-4335.
Chen, F., and Dixon, R.A. (2007). Lignin modification improves fermentable sugar
yields for biofuel production. Nature Biotechnology **25,** 759.
Chen, X., Li, C., Wang, H., and Guo, Z. (2019). WRKY transcription factors:
evolution, binding, and action. Phytopathology Research **1,** 13.
Choi, Y. (2012). A fast computation of pairwise sequence alignment scores between a
protein and a set of single-locus variants of another protein. In Proceedings of the
ACM Conference on Bioinformatics, Computational Biology and Biomedicine
(Orlando, Florida: Association for Computing Machinery), pp. 414–417.
Choi, Y., Sims, G.E., Murphy, S., Miller, J.R., and Chan, A.P. (2012). Predicting the
Functional Effect of Amino Acid Substitutions and Indels. PLOS ONE **7,** e46688.
Chow, C.-N., Lee, T.-Y., Hung, Y.-C., Li, G.-Z., Tseng, K.-C., Liu, Y.-H., Kuo, P.-L.,
Zheng, H.-Q., and Chang, W.-C. (2018). PlantPAN3.0: a new and updated
resource for reconstructing transcriptional regulatory networks from ChIP-seq
experiments in plants. Nucleic Acids Research **47,** D1155-D1163.
Chow, C.-N., Zheng, H.-Q., Wu, N.-Y., Chien, C.-H., Huang, H.-D., Lee, T.-Y.,
Chiang-Hsieh, Y.-F., Hou, P.-F., Yang, T.-Y., and Chang, W.-C. (2015).
PlantPAN 2.0: an update of plant promoter analysis navigator for reconstructing
transcriptional regulatory networks in plants. Nucleic Acids Research **44,** D1154-
D1160.
Ciolkowski, I., Wanke, D., Birkenbihl, R.P., and Somssich, I.E. (2008). Studies on
DNA-binding selectivity of WRKY transcription factors lend structural clues into
WRKY-domain function. Plant Mol Biol **68,** 81-92.
Costea, M., García, M.A., Baute, K., and Stefanović, S. (2015a). Entangled
evolutionary history of Cuscuta pentagona clade: A story involving hybridization
and Darwin in the Galapagos. TAXON **64,** 1225-1242.
Costea, M., García, M.A., Baute, K., Stefanovi, and Sa. (2015b). Entangled
evolutionary history of Cuscuta pentagona clade: A story involving hybridization
and Darwin in the Galapagos. Taxon **64,** 1225-1242.
Cui, S., Wada, S., Tobimatsu, Y., Takeda, Y., Saucet, S.B., Takano, T., Umezawa,
T., Shirasu, K., and Yoshida, S. (2018). Host lignin composition affects

830   haustorium induction in the parasitic plants Phtheirospermum japonicum and
831   Striga hermonthica. New Phytologist **218,** 710-723.

832 **Dahan, M., Towns, J., Cockerill, T., Dahan, M., Foster, I., Gaither, K., Grimshaw,**
833   **A., Hazlewood, V., Lathrop, S., Lifka, D., Peterson, G., Roskies, R., Scott, J.,**
834   **and Wilkins-Diehr, N.** (2014). XSEDE: Accelerating scientific discovery.
835   Computing in Science and Engineering **16,** 62-74.

836 **Dhakshinamoorthy, S., Mariama, K., Elsen, A., and De Waele, D.** (2014). Phenols
837   and lignin are involved in the defence response of banana (Musa) plants to
838   infection. Nematology **16,** 565-576.

839 **Dodds, P.N., and Rathjen, J.P.** (2010). Plant immunity: towards an integrated view of
840   plant–pathogen interactions. Nature Reviews Genetics **11,** 539-548.

841 **Duriez, P., Vautrin, S., Auriac, M.-C., Bazerque, J., Boniface, M.-C., Callot, C.,**
842   **Carrère, S., Cauet, S., Chabaud, M., Gentou, F., Lopez-Sendon, M., Paris,**
843   **C., Pegot-Espagnet, P., Rousseaux, J.-C., Pérez-Vich, B., Velasco, L., Bergès,**
844   **H., Piquemal, J., and Muños, S.** (2019). A receptor-like kinase enhances
845   sunflower resistance to Orobanche cumana. Nature Plants **5,** 1211-1215.

846 **Eudes, A., Sathitsuksanoh, N., Baidoo, E.E.K., George, A., Liang, Y., Yang, F.,**
847   **Singh, S., Keasling, J.D., Simmons, B.A., and Loqué, D.** (2015). Expression of
848   a bacterial 3-dehydroshikimate dehydratase reduces lignin content and improves
849   biomass saccharification efficiency. Plant Biotechnology Journal **13,** 1241-1250.

850 **Fang, Y., and Spector, D.L.** (2010). BiFC Imaging Assay for Plant Protein-Protein
851   Interactions. Cold Spring Harbor Protocols **2010,** pdb.prot5380.

852 **Ferrer, J.L., Austin, M.B., Stewart, C., and Noel, J.P.** (2008). Structure and function
853   of enzymes involved in the biosynthesis of phenylpropanoids. Plant Physiology
854   and Biochemistry **46,** 356-370.

855 **García, M.A., Costea, M., Kuzmina, M., and Stefanović, S.** (2014). Phylogeny,
856   character evolution, and biogeography of Cuscuta (dodders; Convolvulaceae)
857   inferred from coding plastid and nuclear sequences. American Journal of Botany
858   **101,** 670-690.

859 **Gayoso, C., Pomar, F., Novo-Uzal, E., Merino, F., and Martínez de Ilárduya, Ó.**
860   (2010). The Ve-mediated resistance response of the tomato to Verticillium dahliae
861   involves H2O2, peroxidase and lignins and drives PALgene expression. BMC
862   Plant Biology **10,** 232.

863 **Gilbertson, R.L., Hidayat, S.H., Paplomatas, E.J., Rojas, M.R., Hou, Y.-M., and**
864   **Maxwell, D.P.** (1993). Pseudorecombination between infectious cloned DNA
865   components of tomato mottle and bean dwarf mosaic geminiviruses. Journal of
866   General Virology **74,** 23-31.

867 **Goldwasser, Y., Hershenhorn, J., Plakhine, D., Kleifeld, Y., and Rubin, B.** (1999).
868   Biochemical factors involved in vetch resistance toOrobanche aegyptiaca.
869   Physiological and Molecular Plant Pathology **54,** 87-96.

870 **Gutiérrez, A., Rodríguez, I.M., and del Río, J.C.** (2006). Chemical Characterization of
871   Lignin and Lipid Fractions in Industrial Hemp Bast Fibers Used for
872   Manufacturing High-Quality Paper Pulps. Journal of Agricultural and Food
873   Chemistry **54,** 2138-2144.

Hegenauer, V., Fürst, U., Kaiser, B., Smoker, M., Zipfel, C., Felix, G., Stahl, M., and Albert, M. (2016). Detection of the plant parasite *Cuscuta reflexa* by a tomato cell surface receptor. Science **353,** 478-481.

Hembree, K.J., Lanini, W., and Va, N. (1999). Tomato varieties show promise of dodder control. In Proc. Calif. Weed Sci. Soc, pp. 205-206.

Hou, Y.M., and Gilbertson, R.L. (1996). Increased pathogenicity in a pseudorecombinant bipartite geminivirus correlates with intermolecular recombination. J Virol **70,** 5430-5436.

Ichihashi, Y., Aguilar-Martínez, J.A., Farhi, M., Chitwood, D.H., Kumar, R., Millon, L.V., Peng, J., Maloof, J.N., and Sinha, N.R. (2014). Evolutionary developmental transcriptomics reveals a gene network module regulating interspecific diversity in plant leaf shape. Proceedings of the National Academy of Sciences **111,** E2616-E2621.

Jin, J., Tian, F., Yang, D.-C., Meng, Y.-Q., Kong, L., Luo, J., and Gao, G. (2016). PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. Nucleic Acids Research **45,** D1040-D1045.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., and Hunter, S. (2014). InterProScan 5: genome-scale protein function classification. Bioinformatics **30,** 1236-1240.

Kaiser, B., Vogg, G., Fürst, U.B., and Albert, M. (2015). Parasitic plants of the genus Cuscuta and their interaction with susceptible and resistant host plants. Frontiers in Plant Science **6,** 45.

Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N., and Sternberg, M.J.E. (2015). The Phyre2 web portal for protein modeling, prediction and analysis. Nature Protocols **10,** 845-858.

Kerppola, T.K. (2008). Bimolecular Fluorescence Complementation (BiFC) Analysis as a Probe of Protein Interactions in Living Cells. Annual Review of Biophysics **37,** 465-487.

Kumari, C., Dutta, T.K., Banakar, P., and Rao, U. (2016). Comparing the defence-related gene expression changes upon root-knot nematode attack in susceptible versus resistant cultivars of rice. Scientific Reports **6,** 22846.

Kusumoto, D., Goldwasser, Y., Xie, X., Yoneyama, K., Takeuchi, Y., and Yoneyama, K. (2007). Resistance of red clover (Trifolium pratense) to the root parasitic plant Orobanche minor is activated by salicylate but not by jasmonate. Ann Bot **100,** 537-544.

Labrousse, P., Arnaud, M.C., Serieys, H., Bervillé, A., and Thalouarn, P. (2001). Several Mechanisms are Involved in Resistance of Helianthus to Orobanche cumana Wallr. Annals of Botany **88,** 859-868.

LANE, J.A., BAILEY, J.A., BUTLER, R.C., and TERRY, P.J. (1993). Resistance of cowpea [Vigna unguiculata (L.) Walp.] to Striga gesnerioides (Willd.) Vatke, a parasitic angiosperm. New Phytologist **125,** 405-412.

Lanini, W.T., and Kogan, M. (2005). Biology and Management of Cuscuta in Crops.

Liljegren, S. (2010). Phloroglucinol Stain for Lignin. Cold Spring Harbor Protocols **2010,** pdb.prot4954.

Liu, Q., Luo, L., and Zheng, L. (2018). Lignins: Biosynthesis and Biological Functions in Plants. International journal of molecular sciences **19,** 335.

Lozano-Baena, M.D., Prats, E., Moreno, M.T., Rubiales, D., and Pérez-de-Luque, A. (2007). Medicago truncatula as a Model for Nonhost Resistance in Legume-Parasitic Plant Interactions. Plant Physiology **145,** 437.

Malinovsky, F.G., Fangel, J.U., and Willats, W.G.T. (2014). The role of the cell wall in plant immunity. Frontiers in Plant Science **5**.

Mitchell, A.L., Attwood, T.K., Babbitt, P.C., Blum, M., Bork, P., Bridge, A., Brown, S.D., Chang, H.-Y., El-Gebali, S., Fraser, M.I., Gough, J., Haft, D.R., Huang, H., Letunic, I., Lopez, R., Luciani, A., Madeira, F., Marchler-Bauer, A., Mi, H., Natale, D.A., Necci, M., Nuka, G., Orengo, C., Pandurangan, A.P., Paysan-Lafosse, T., Pesseat, S., Potter, S.C., Qureshi, M.A., Rawlings, N.D., Redaschi, N., Richardson, L.J., Rivoire, C., Salazar, G.A., Sangrador-Vegas, A., Sigrist, C.J A., Sillitoe, I., Sutton, G.G., Thanki, N., Thomas, P.D., Tosatto, S.C E., Yong, S.-Y., and Finn, R.D. (2018). InterPro in 2019: improving coverage, classification and access to protein sequence annotations. Nucleic Acids Research **47,** D351-D360.

Moura, J.C.M.S., Bonine, C.A.V., De Oliveira Fernandes Viana, J., Dornelas, M.C., and Mazzafera, P. (2010). Abiotic and Biotic Stresses and Changes in the Lignin Content and Composition in Plants. Journal of Integrative Plant Biology **52,** 360-376.

Mutuku, J.M., Yoshida, S., Shimizu, T., Ichihashi, Y., Wakatake, T., Takahashi, A., Seo, M., and Shirasu, K. (2015). The WRKY45-Dependent Signaling Pathway Is Required For Resistance against Striga hermonthica Parasitism. Plant Physiology **168,** 1152-1163.

Mutuku, J.M., Cui, S., Hori, C., Takeda, Y., Tobimatsu, Y., Nakabayashi, R., Mori, T., Saito, K., Demura, T., Umezawa, T., Yoshida, S., and Shirasu, K. (2019). The Structural Integrity of Lignin Is Crucial for Resistance against Striga hermonthica Parasitism in Rice. Plant Physiology **179,** 1796-1809.

O'Brien, T.P., Feder, N., and McCully, M.E. (1964). Polychromatic staining of plant cell walls by toluidine blue O. Protoplasma **59,** 368-373.

Padmanabhan, M.S., and Dinesh-Kumar, S.P. (2014). The conformational and subcellular compartmental dance of plant NLRs during viral recognition and defense signaling. Current Opinion in Microbiology **20,** 55-61.

Padmanabhan, M.S., Ma, S., Burch-Smith, T.M., Czymmek, K., Huijser, P., and Dinesh-Kumar, S.P. (2013). Novel Positive Regulatory Role for the SPL6 Transcription Factor in the N TIR-NB-LRR Receptor-Mediated Plant Innate Immunity. PLOS Pathogens **9,** e1003235.

Pan, C., Ye, L., Qin, L., Liu, X., He, Y., Wang, J., Chen, L., and Lu, G. (2016). CRISPR/Cas9-mediated efficient and heritable targeted mutagenesis in tomato plants in the first and later generations **6,** 24765.

PÉRez-De-Luque, A., Rubiales, D., Cubero, J.I., Press, M.C., Scholes, J., Yoneyama, K., Takeuchi, Y., Plakhine, D., and Joel, D.M. (2005). Interaction between Orobanche crenata and its Host Legumes: Unsuccessful Haustorial Penetration and Necrosis of the Developing Parasite. Annals of Botany **95,** 935-942.

Pomar, F., Merino, F., and Barceló, A.R. (2002). O-4-Linked coniferyl and sinapyl
aldehydes in lignifying cell walls are the main targets of the Wiesner
(phloroglucinol-HCl) reaction. Protoplasma **220,** 0017-0028.
Pradhan Mitra, P., and Loqué, D. (2014). Histochemical Staining of Arabidopsis
thaliana Secondary Cell Wall Elements. Journal of Visualized Experiments :
JoVE**,** 51381.
Ralph, J., and Hatfield, R.D. (1991). Pyrolysis-GC-MS characterization of forage
materials. Journal of Agricultural and Food Chemistry **39,** 1426-1437.
Ranjan, A., Ichihashi, Y., Farhi, M., Zumstein, K., Townsley, B., David-Schwartz,
R., and Sinha, N.R. (2014). De Novo Assembly and Characterization of the
Transcriptome of the Parasitic Weed Dodder Identifies Genes Associated with
Plant Parasitism. Plant Physiology **166,** 1186-1199.
Ranjan, A., Budke, J.M., Rowland, S.D., Chitwood, D.H., Kumar, R., Carriedo, L.,
Ichihashi, Y., Zumstein, K., Maloof, J.N., and Sinha, N.R. (2016). eQTL
Regulating Transcript Levels Associated with Diverse Biological Processes in
Tomato. Plant Physiology **172,** 328-340.
Reimers, P.J., and Leach, J.E. (1991). Race-specific resistance to Xanthomonas oryzae
pv. oryzae conferred by bacterial blight resistance gene Xa-10 in rice (Oryza
sativa) involves accumulation of a lignin-like substance in host tissues.
Physiological and Molecular Plant Pathology **38,** 39-55.
Runyon, J.B., Mescher, M.C., Felton, G.W., and De Moraes, C.M. (2010). Parasitism
by Cuscuta pentagona sequentially induces JA and SA defence pathways in
tomato. Plant, Cell & Environment **33,** 290-303.
Sahm, A., Pfanz, H., Grünsfelder, M., Czygan, F.-C., and Proksch, P. (1995).
Anatomy and Phenylpropanoid Metabolism in the Incompatible Interaction of
Lycopersicon esculentum and Cuscuta reflexa. Botanica Acta **108,** 358-364.
Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25
years of image analysis. Nature Methods **9,** 671.
Solís-Lemus, C., Bastide, P., and Ané, C. (2017). PhyloNetworks: A Package for
Phylogenetic Networks. Molecular Biology and Evolution **34,** 3292-3298.
Sparkes, I.A., Runions, J., Kearns, A., and Hawes, C. (2006). Rapid, transient
expression of fluorescent fusion proteins in tobacco plants and generation of
stably transformed plants. Nature Protocols **1,** 2019-2025.
Stefanović, S., Kuzmina, M., and Costea, M. (2007). Delimitation of major lineages
within Cuscuta subgenus Grammica (Convolvulaceae) using plastid and nuclear
DNA sequences. American Journal of Botany **94,** 568-589.
Stenger, D.C., Revington, G.N., Stevenson, M.C., and Bisaro, D.M. (1991).
Replicational release of geminivirus genomes from tandemly repeated copies:
evidence for rolling-circle replication of a plant viral DNA. Proceedings of the
National Academy of Sciences of the United States of America **88,** 8029-8033.
Su, C., Liu, H., Wafula, E.K., Honaas, L., de Pamphilis, C.W., and Timko, M.P.
(2020). SHR4z, a novel decoy effector from the haustorium of the parasitic weed
Striga gesnerioides, suppresses host plant immunity. New Phytologist **226,** 891-
908.
Taheri, P., and Tarighi, S. (2012). The Role of Pathogenesis-Related Proteins in the
Tomato-Rhizoctonia solani Interaction. Journal of Botany **2012,** 6.

Townsley, B.T., Covington, M.F., Ichihashi, Y., Zumstein, K., and Sinha, N.R.
(2015). BrAD-seq: Breath Adapter Directional sequencing: a streamlined, ultra-
simple and fast library preparation protocol for strand specific mRNA library
construction. Frontiers in Plant Science **6**.
Vance, C.P., Kirk, T.K., and Sherwood, R.T. (1980). Lignification as a Mechanism of
Disease Resistance. Annual Review of Phytopathology **18,** 259-288.
Vel, squez, A., C, Chakravarthy, S., and Martin, G.B. (2009). Virus-induced Gene
Silencing (VIGS) in Nicotiana benthamiana and Tomato**,** e1292.
War, A.R., Paulraj, M.G., Ahmad, T., Buhroo, A.A., Hussain, B., Ignacimuthu, S.,
and Sharma, H.C. (2012). Mechanisms of plant defense against insect
herbivores. Plant Signaling & Behavior **7,** 1306-1320.
Yaakov, G., Lanini, W.T., and Wrobel, R.L. (2001). Tolerance of Tomato Varieties to
Lespedeza Dodder. Weed Science **49,** 520-523.
Yoder, J.I., and Scholes, J.D. (2010). Host plant resistance to parasitic weeds; recent
progress and bottlenecks. Current Opinion in Plant Biology **13,** 478-484.
Yoshida, S., and Shirasu, K. (2009). Multiple layers of incompatibility to the parasitic
witchweed, Striga hermonthica. New Phytologist **183,** 180-189.
Zhang, S.-H., Yang, Q., and Ma, R.-C. (2007). Erwinia carotovora ssp. carotovora
Infection Induced "Defense Lignin" Accumulation and Lignin Biosynthetic Gene
Expression in Chinese Cabbage (Brassica rapa L. ssp. pekinensis). Journal of
Integrative Plant Biology **49,** 993-1002.
Zhang, Y., Wu, L., Wang, X., Chen, B., Zhao, J., Cui, J., Li, Z., Yang, J., Wu, L.,
Wu, J., Zhang, G., and Ma, Z. (2019). The cotton laccase gene GhLAC15
enhances Verticillium wilt resistance via an increase in defence-induced
lignification and lignin components in the cell walls of plants. Molecular Plant
Pathology **20,** 309-322.
1037

Figures

Figure 1 | The comparison of resistance responses to *C. campestris* in tomato cultivars. (A) *C. campestris* grows on the susceptible H9775, (B) and cannot attach on the resistant H9553. (C) The biomass ratio of host and *C. campestris* (*Cuscuta* weight/ tomato weight) on different cultivars. Data were assessed using pair-wise comparisons with Tukey test. P-values between "a" and "b" are < 0.05; H1706, n = 9; H9775, n = 10, H9492, n = 10, H9553 n = 7. Data were collected at 45 days post attachment (DPA). (D-G) 100 μm vibratome longitudinal sections of *C. campestris* haustoria attaching to H1706 (D-E) and H9553 (F-G), and stained with Toluidine Blue O. Lignin is stained as blue. (D and F) Scale bar, 40 μm. (E and G) Scale bar, 10 μm. (H-O) are ~300 μm sections of the haustoria attachment sites stained with Phloroglucinol-HCl. Scale bar, 1 mm. Lignin is stained as red. Stem cross-sections of H1706 (H and L), H9775 (I and M), H9492 (J and N), and H9553 (K and O) without *C. campestris* treatment (labeled with -Cc) and with *C. campestris* attached (labeled with +Cc). (P) Cortex lignin area percentage in different cultivars. Data presented are assessed using multiple comparisons with Dunnett's test. "*": p-values < 0.05, "**": p-values < 0.01. (Q) Cell death area percentage in different cultivars. (P-Q) H1706-Cc, n = 20; H1706+Cc, n = 38; H9775-Cc, n = 19; H9775+Cc, n = 40; H9492-Cc, n = 16; H9492+Cc, n = 38; H9553-Cc, n = 17; H9553+Cc, n = 30. Data were collected at 14 DPA.

Figure 2 | Virus-based Gene Expression (VGE) in tomato H1706. (A-F) ~300 μm

stem sections near injection sites, Phloroglucinol-HCl stains lignin red. VGE of *GUS* (A),

LIF1 (B), *SlMYB55* (C), and *CuRLR1* (E) in stem without *C. campestris*. VGE of *GUS*

(D) and *CuRLR1* (F) with *C. campestris*. (G) Cortex lignin area percentage in VGE of

LIF1 and *SlMYB55* (n = 12 each) and (H) *CuRLR1* with and without *C. campestris* (*GUS*-

Cc, n = 18; *GUS*+Cc, n =28; *CuRLR1*-Cc, n = 47; *CuRLR1*+Cc, n = 53). (G-H) Data were

collected at 7 days post injection (DPI) and 14 days post attachment (DPA). Data are

assessed using Dunnett's test with *GUS*-Cc as negative control. "*": p-values < 0.01. (I-

L) VGE of *CuRLR1*, *LIF1* and *SlMYB55* induces cortical lignin making H1706 resistant

to *C. campestris*. Scale bar, 30 μm. Samples were collected at 7 DPI and 6 DPA. (M)

Acetyl bromide assay for lignin in VGE stems. Samples were collected at 7 DPI and 6

DPA. Data are assessed using Dunnett's test. "*": p-values < 0.05, "**": p-values < 0.01.

Biological replicates for *GUS*, n = 18; *SlMYB55*, n = 10; *LIF1*, n = 18, *CuRLR1*-Cc, n =

18; *CuRLR1*+Cc, n = 18. Technical replicates for acetyl bromide assay; *GUS*, n = 11;

SlMYB55, n = 3; *LIF1*, n = 13; *CuRLR1*-Cc, n = 11; *CuRLR1*+Cc, n = 11. (N) PYRO-GC

assay for monolignols in *CuRLR1* VGE samples with and without *C. campestris*. Samples

were collected at 7 DPI and 6 DPA. Biological replicates collected from first internodes;

GUS, n = 8; *LIF1*, n = 8, *CuRLR1*-Cc, n = 18; *CuRLR1*+Cc, n = 18. PYRO-GC assay

technical replicates; *GUS*, n = 3; *LIF1*, n = 3; *CuRLR1*-Cc, n = 5; *CuRLR1*+Cc, n = 5.

1078

Figure 3 | Barnes-Hut t-distributed stochastic neighbor embedding (BH-SNE)

generated gene clusters and gene co-expression network (GCN) analysis. (A) BH-

SNE generated gene clusters based on their gene expression patterns. Total cluster

(module) number was 48. (B) The candidate genes that are included in the clusters are

labeled with their corresponding colors. The selected gene clusters for GCN are labeled

in yellow (CuRel and WRKY16 cluster, cluster 17), red (CCOMT and LAC cluster,

cluster 11), blue (MYB55 cluster, cluster 39), pink (LIF1 cluster, cluster 46), green (CCR

cluster, cluster 23) colors. *CuRLR1* is in the noise cluster, and is labeled in grey color.

Parameters used in this analysis: perplexity (perp) = 20, lying = 250, cutoff = 20, seed =

2. (C-F) Gene co-expression networks (GCNs) of four different Heinz susceptible and

resistant cultivars upon *C. campestris* treatments. Based on BH-SNE analysis, 1676 genes

in cluster 11, 17, 23, 39, 46 and CuRLR1 were selected for building GCNs. +Cc indicates

with *C. campestris* infection treatments. The genes that are listed at the left of the GCN

and not labeled in the network are the genes that have no coexpression connections with

all the other genes in the list.

1094

Figure 4 | The role of *SlWRKY16* in *Cuscuta* resistance using CRISPR/Cas9 gene
knockouts. (A-H) Samples and data were collected at 7 DPA. (A-B) overall phenotype
comparison between M82 and homozygous *SlWRKY16* CRISPR lines (*wrky16*). Scale
bar, 2 cm. (C-D) *C. campestris* growing on M82 and *wrky16*. (E-F) ~300 μm hand
sections of M82 and *wrky16* stems near *Cuscuta* attachment site stained with
Phloroglucinol-HCl. Lignin is stained red. (G) Leaf number of *wrky16* and M82.
Biological replicates, n = 18 for each. (H) Cortex lignin area percentage in M82 and
wrky16 stems. Data presented are assessed using student's t-test. "*" indicates p-value is
less than 0.01. Replicates: M82, n = 33; *wrky16*, n = 34.

1104

Figure 5 | Subcellular localization of candidate genes and protein-protein
interactions. (A) Subcellular localizations of LIF1, SlWRKY16, and CuRLR1 proteins.
(B) Verification of protein-protein interactions and locations of SlWRKY16 and LIF1 by
Bimolecular Fluorescence Complementation (BiFC). The gene with *cCitrine* fusion is
listed before the "+" sign and the gene with *nCitrine* fusion is listed after the "+" sign.
(C) Co-expression of fusion protein LIF1-GFP and SlWRKY16-RFP to observe
subcellular localizations. Yellow color in the merged panel indicates that GFP and RFP
signals are overlapped.

Figure 6 | *C. campestris* extract injections to detect *Cuscuta* signals. (A-I) ~300 μm hand sections of resistant H9553 stems near injection sites stained with Phloroglucinol-HCl. Lignin is stained red. Data were collected at 7 days post injection (DPI). The H9553 plants are injected with (A) water, (B) untreated *C. campestris* extract, pH 5.8, (C) *C. campestris* extract filtered with 0.2 μm filter, (D) pH 9 water, and (E) pH 9 *C. campestris*

extract, (F) heat-treated *C. campestris* extract (95°C for 30 minutes), (G) Protease E-
treated *C. campestris* extract, (H) Protease K-treated *C. campestris* extract, and (I)
Trypsin-treated *C. campestris* extract. (J) Percentage of lignified cortex area in total stem
area. The samples injected with water serve as negative controls. Different treated or
untreated *C. campestris* extracts are compared to negative controls. Data presented are
assessed using pair-wise comparisons with Tukey test. P-value of the contrasts between
"a" and "b" are less than 0.01. Replicates: water, n = 22; untreated extract, n = 13; filtered
extract, n = 24; heat-treated extract, n = 12; pH 9 water, n = 11; pH 9 extract, n = 12;
Protease E-treated, n = 12; Protease K-treated, n = 12; Trypsin-treated, n = 12.

1128

Figure 7 | Model of *C. campestris* resistance response in tomato cultivars. Red-labeled pathway: newly identified cytosolic *CuRLR1*, which may receive large signaling molecules from *C. campestris* or play a role in signal transduction upon *Cuscuta* perception. This triggers downstream signal transduction and induces a lignin-based resistance response. This resistant response may be an effector-triggered immunity (ETI). Pink and yellow-labeled pathway: *SlMYB55* and *LIF1* function as positive regulators in the lignin biosynthesis pathway. Yellow and green-labeled pathway: *SlWRKY16* and *LIF1* mediated lignin-based resistant responses and with a potential connection to *CuRe1*. Blue-labeled pathway: previously identified *CuRe1* mediated PAMP/MAMP-triggered immunity (PTI/MTI) pathway.