Ovarian hormone status dictates the neuroinflammatory and behavioural consequences of sub-chronic stress exposure in middle-aged female mice

Rand S. Eid[1,3], Stephanie E. Lieblich[2,3], Sarah J. Wong[2,3], and Liisa A.M. Galea[1,2,3]

[1] Graduate program in Neuroscience, University of British Columbia, Vancouver, BC
[2] Department of Psychology, University of British Columbia, Vancouver, BC
[3] Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver, BC

Corresponding author:

Dr. Liisa Galea
Djavad Mowafaghian Centre for Brain Health
University of British Columbia
2215 Wesbrook Mall
Vancouver, BC
Canada, V6T 1Z3
Tel: +1 (604) 822 6536
Email: liisa.galea@ubc.ca

Abstract

Ovarian hormones influence the outcomes of stress exposure and are implicated in stress-related disorders including depression, yet their roles are often complex and seemingly contradictory. Importantly, depression and stress exposure are associated with immune dysregulation, and ovarian hormones have immunomodulatory properties. However, how ovarian hormones can influence the inflammatory outcomes of stress exposure is poorly understood. Here, we examined the effects of long-term ovariectomy on the behavioural and neuroinflammatory outcomes of sub-chronic stress exposure in middle-aged mice. Briefly, sham-operated and ovariectomized mice were assigned to non-stress groups or exposed to 6 days of variable stress. Mice were assessed on a battery of behavioural tests, and cytokine concentrations were quantified in the frontal cortex and hippocampus. Postsynaptic density protein-95 and phosphorylated mitogen-activated protein kinases (MAPKs) were also measured in the frontal cortex. Long-term ovariectomy modified the central cytokine profile by robustly reducing cytokine concentrations in the frontal cortex and modestly increasing concentrations in the hippocampus. Under non-stress conditions, long-term ovariectomy also reduced extracellular signal-regulated kinase (ERK) phosphoprotein expression in the frontal cortex and increased some measures of depressive-like behaviour. The effects of sub-chronic stress exposure were however more pronounced in sham-operated mice. Notably, in sham-operated mice only, sub-chronic stress exposure increased IL-1β and IL-6:IL-10 ratio in the frontal cortex and hippocampus and reduced pERK1/2 expression in the frontal cortex. Further, although sub-chronic stress exposure increased anhedonia-like behaviour regardless of ovarian status, it increased passive-coping behaviour in sham-operated mice only. These data indicate that long-term ovariectomy has potent effects on the central cytokine milieu and mitigates the neuroinflammatory and behavioural effects of sub-chronic stress exposure in middle-aged mice. These findings therefore suggest that the immunomodulatory properties of ovarian hormones are of relevance in the context of stress and possibly depression.

1   **Introduction**

Major Depressive Disorder (MDD), a leading cause of disability and ill-health worldwide (Friedrich,

2017), is approximately twice more prevalent in women than men (Kessler and Bromet, 2013; Salk et

al., 2017). In addition to this robust difference in prevalence rates, disease presentation and

pathophysiology differ significantly between men and women (reviewed in Eid et al., 2019). Although

there is an encouraging increase in studies investigating sex differences in depression and in animal

models of stress exposure, studies that target female-specific factors that may contribute to risk or

resilience to depression remain scarce. Ovarian hormones play a role in mood regulation and influence

depression in women (Brummelte and Galea, 2010; Hantsoo and Epperson, 2015; Soares, 2013) and

therefore represent one such target. For example, there is an increased susceptibility to develop

depression during the reproductive years in women compared to men (Gutiérrez-Lobos et al., 2002).

These data provide the basis for the hypothesis that ovarian hormones may increase risk for depression

in women. Other lines of evidence, however, support the contrasting hypothesis that ovarian hormones

may afford resilience against depression. Specifically, events that are characterized by large reductions

in ovarian hormones, including the postpartum and perimenopause, are associated with an increased

risk for depression in women (Cohen et al., 2006; Freeman et al., 2004; Hendrick et al., 1998; Soares,

2014). In rodent studies, ovariectomy alone increases depressive-like behaviour in the short term and

increases susceptibility to the development of depressive-like behaviour in response to chronic stress

exposure in the long term (Lagunas et al., 2010; Li et al., 2014; Mahmoud et al., 2016). Taken together,

this literature highlights the complex and seemingly paradoxical roles of ovarian hormones in mood

regulation and depression. Importantly, ovarian hormones regulate a wide range of physiological

processes and systems that are compromised in depression, including stress response and immune

systems (Goel et al., 2014; Klein and Flanagan, 2016). Therefore, examining the interplay between

endocrine and immune systems in the context of stress exposure may clarify the role of ovarian

hormones in depression.

There is mounting evidence of immune dysregulation in depression (Hodes et al., 2015; Miller

and Raison, 2016). At least a subpopulation of individuals with MDD present with increased markers

of inflammation, supported by several meta-analyses which highlight that IL-6 and TNF-α are

increased in the blood and cerebral spinal fluid (Dowlati et al., 2010; Haapakoski et al., 2015; Liu et al.,

2012; Wang and Miller, 2018). Further, poor antidepressant efficacy has been linked to an inability to

normalize dysregulated inflammatory processes (Syed et al., 2018). Stress exposure, a major risk factor

for depression (Kendler et al., 1999), can also activate inflammatory responses in the brain and

periphery (Felger et al., 2015; Gouin et al., 2012). Importantly, sex differences exists across many

immune processes, mediated in part by sex hormones (Klein and Flanagan, 2016). Several reports

document the anti-inflammatory actions of estrogens in the brain; in response to inflammatory

challenges *in vitro* and *in vivo*, estradiol or selective estrogen receptor modulators (SERMs) ameliorate

microglia inflammatory activity and the secretion of proinflammatory cytokines including IL-1β, TNF-

α, and IL-6 (Smith et al., 2011; Suuronen et al., 2005; Tapia-Gonzalez et al., 2008; Vegeto et al., 2001).

Despite this, the role of ovarian hormones at the intersection of depression and immune dysregulation

is not well known. This is particularly important during the transition to menopause, a time of

heightened risk for MDD (Cohen et al., 2006; Freeman et al., 2004) that is also marked by substantial

changes in ovarian and immune function (Burger et al., 2002; Giefing-Kröll et al., 2015). The current

study targets this gap in the literature by examining how the ovarian hormone milieu can influence the

depressogenic and inflammatory consequences of stress exposure in middle-aged mice.

A better understanding of endocrine-immune interactions in the pathogenesis of depression may

be achieved by investigating downstream intracellular signaling pathways that are elicited by stress

exposure. Mitogen-activated protein kinase (MAPK) pathways represent a logical target, as they

regulate a wide range of cellular processes in response to diverse stimuli, including hormones and

cytokines (Cargnello and Roux, 2011). Further, studies in humans and animal models of depression

indicate that MAPK signaling pathways are compromised in the disorder (Hollos et al., 2018; Miller

and Raison, 2006; Wang and Mao, 2019). The extracellular signal-regulated kinase (ERK) subfamily of

MAPKs has received most attention in the context of depression (Wang and Mao, 2019), however the

stress-activated protein kinases, c-Jun NH_2-terminal kinase (JNK) and p38 MAPK, may also be

implicated (Hollos et al., 2018; Miller and Raison, 2006). Here, we quantified phosphorylated MAPKs

to explore cellular mechanisms that may underly the role of ovarian hormones in risk or resilience in

the face of stress exposure.

Disruptions in synaptic function and number are evidenced following stress and thought to be

key factors in the pathogenesis of stress-related disorders such as depression (Duman et al., 2016). The

prefrontal cortex (PFC) may be particularly vulnerable; in addition to reports of reduced PFC volume

in depression (Drevets, 2000; Drevets et al., 2008), post-mortem studies find reductions in the number

of synapses and in the expression of synapse-related genes in the PFC (Feyissa et al., 2009; Kang et al.,

2012). In rodent models, chronic stress exposure results in dendritic remodeling in the frontal cortex

and hippocampus (reviewed in McEwen et al., 2016), in addition to alterations is spines and synaptic

markers (Orlowski et al., 2012; Workman et al., 2013). Importantly, neuronal remodeling in the frontal

cortex and hippocampus in response to stress is influenced by sex and sex hormones (Galea et al., 1997; Shansky et al., 2010, 2009). Further, inflammatory processes could be implicated in synaptic alterations in stress and depression, as synaptic pruning is mediated by microglia during development and possibly in disease (Hong et al., 2016; Paolicelli et al., 2011). Therefore, examining synaptic markers may also clarify endocrine-immune interactions in stress and depression.

In this study, we used female mice to examine the effects of long-term ovariectomy on the behavioural and inflammatory consequences of sub-chronic stress exposure in middle age. We chose middle age because few studies have examined this important time point in females, and because perimenopause represents a time of increased vulnerability to depression in women. We measured cytokine concentration in the periphery, the hippocampus, and frontal cortex, areas known to be susceptible to stress exposure and compromised in depression (Howard et al., 2019; McKinnon et al., 2009). We also quantified phosphorylated MAPKs and postsynaptic density protein 95 (PSD-95) in the frontal cortex. We hypothesized that ovarian hormone deprivation will dictate behavioural outcomes and modulate cytokines, cell signaling proteins, and PSD-95 expression in response to stress exposure.

Method

Animals and surgery

Young-adult female C57BL/6N mice were purchased from Charles River Laboratories (Quebec, Canada). Mice received bilateral ovariectomy (OVX; n = 14) or sham surgery (Sham; n = 14) at Charles River Laboratories at 8 weeks of age and arrived at our facility at 9 weeks of age. Mice were group housed (2-3/cage) in a temperature- and humidity- controlled colony room (21 ± 1°C; 50 ± 10% humidity), maintained on a 12-hour light/dark cycle (lights on at 07:00 h), and provided *ad libitum* access to food and water. Mice were left undisturbed, apart from weekly cage changing, until the beginning of further experimental manipulations at 11 months of age. The reason mice were tested at middle age was two fold: 1) this is a time of heightened vulnerability to depression in women (Cohen et al., 2006; Freeman et al., 2004) that few animal studies target, and 2) previous rodent studies suggest that differences in susceptibility to stress exposure may emerge only after longer periods of ovarian hormone deprivation (Lagunas et al., 2010). All procedures were approved by the Animal Care Committee at the University of British Columbia and were performed in accordance with the ethical guidelines set by the Canadian Council on Animal Care.

Sub-chronic variable stress exposure

In each ovarian status condition (OVX, Sham), half the mice were exposed to 6 days of sub-chronic variable stress, adapted from previous methods (Hodes et al., 2015; Labonté et al., 2017). We chose a 6-day paradigm to examine group differences in early indicators of immune dysregulation in response to stress. Mice in the stress condition were transferred to a separate colony room and allowed to acclimatize for two weeks prior to stress exposure. Stress-exposed mice were housed separately to avoid the transmission of stress-induced olfactory cues to non-stressed mice (Brechbuhl et al., 2013). The protocol consisted of three stressors, applied once per day on days 1-3, and repeated on days 4-6 (see timeline in **Figure 1**). Mice were exposed to foot shock stress on days 1 and 4 (100 shocks at 0.45mA, randomly distributed across 1 hour), tail suspension stress on days 2 and 5 (1 hour, suspended using laboratory tape, mid-way across the tail), and restraint stress on days 3 and 6 (1 hour, in a well-ventilated 50ml falcon tube, in the home cage). Mice in non-stress groups were handled daily for 5 minutes across days 1-6, but otherwise left undisturbed. Body weight was monitored daily across the stress exposure period and in non-stressed controls.

Behavioural testing

A battery of behavioural tests was conducted between experimental days 7 and 9 (**Figure 1**). All mice underwent behavioural testing, and except for the sucrose preference test, all tests were conducted under red light conditions and in designated testing rooms. Animals were acclimatized to the rooms for 1 hour prior to testing.

Splash test (ST). Self-grooming behaviour in the ST was used as an index of self care-like and motivational behaviour, adapted from previous methods (Hodes et al., 2015; Isingrini et al., 2010). Mice were sprayed three times on the back with a 10% sucrose solution (in tap water), then individually placed in an empty clean cage for 5 mins. The test was filmed, and time spent grooming was scored by an experimenter blinded to condition. Immediately after the self-grooming test, mice were individually housed in clean cages, deprived of food in preparation for the novelty suppressed feeding test, and returned to their colony room. Mice remained individually housed for the remainder of the experiment as per Hodes et al., 2015.

Novelty Suppressed Feeding (NSF). The NSF test was used to assess anxiety-like behaviour, and testing was performed according to published protocols (Hodes et al., 2015; Santarelli et al., 2003). Mice were food-deprived overnight, then individually placed in an arena (50x50x20cm) covered with clean bedding and a single pellet of regular chow in the centre. Latency to start feeding was measured

as an index of anxiety-like behaviour. Mice that did not feed within 10 minutes were removed from the

arena and assigned a latency of 600 seconds in the analysis. Bedding was replaced between mice. To

account for potential appetite differences between groups, home cage latency to feed and amount of

food consumed within 5 minutes were measured.

Sucrose Consumption Test (SCT). The SCT was used to assess anhedonia-like behaviour and

performed according to previous protocols (Hodes et al., 2015). Mice were first acclimatized to two

bottles in the home cage, both filled with tap water. After 24 hours and following the NSF test, the

bottles were replaced with two new bottles, one containing tap water and the other 1% sucrose (in tap

water). The right-left position was counterbalanced between animals in each condition. The bottles

were removed 24 hours later, weighed, then returned to the cage for an additional 24 hours with the

right-left position switched for each mouse. Importantly, to avoid hunger-related changes in sucrose

consumption, sucrose was presented after food was returned to the home cage following NSF.

Forced Swim Test (FST). The FST was used to assess stress-coping behaviour and was conducted

according to standard protocols (Can et al., 2012). Mice were individually placed in a 4L glass beaker

filled with clean water at 25□±□0.5°C to a depth of 15cm for a duration of 6 minutes. The test was

filmed, and an observer blinded to experimental condition scored two aspects of passive-coping

behaviour: 1) latency to first bout of immobility and 2) amount of time spent immobile across the

duration of the test. Passive-coping behaviour was defined as immobility with only movements

necessary to remain floating or to keep the head above water.

Figure 1. Experimental timeline. Mice received bilateral ovariectomy or sham surgery at 8 weeks of age. At 11 months of age, mice in the stress condition were subjected to foot shock stress on days 1 and 3, tail suspension stress on days 2 and 4, and restraint stress on days 3 and 6. All mice were tested on the splash test on day 7, the novelty suppressed feeding test on day 8, the forced swim test on day 9, and the sucrose consumption test on days 8-10 (starting after NSF on day 8). Tissue was collected on day 10.

Tissue collection and processing

Mice were euthanized via rapid decapitation, and trunk blood was collected into EDTA-coated tubes

then centrifuged at 4°C and 10g for 15 minutes and plasma stored at -80°C. Immediately following

decapitation, the frontal cortex and hippocampus were micro-dissected on a cold surface, flash-frozen

on dry ice, then stored at -80°C. Frontal cortex was collected anterior to the genu of the corpus

collosum, and the entire rostral-caudal extent of the hippocampus was collected. Adrenal glands were

also collected and weighed immediately.

Electrochemiluminescence immunoassay kits from Meso Scale Discovery (MSD; Rockville, MD) were

used according to manufacturer instructions for cytokine, PSD-95, and cell signaling protein

measurements. A Sector Imager 2400 (MSD) was used to read the plates, and data was analyzed using

the Discovery Workbench 4.0 software from MSD. Frontal cortices and hippocampi from all mice were

homogenized individually using an Omni Bead Ruptor (Omni international, Kennesaw, GA) with

200µl and 150µL of cold lysis buffer, respectively. Homogenates were centrifuged at 4°C and 10g for

15 minutes, and stored at -80°C. For all assays (cytokine, PSD-95, cell signaling phosphoproteins),

frontal cortex and hippocampus values were normalized to total protein concentrations, which were

quantified using the Pierce Micro BCA Protein Assay Kit (ThermoFisher Scientific) used according to

manufacturer instructions, with samples run in triplicates. Cell signaling proteins and PSD-95 were

only quantified in the frontal cortex, as hippocampal tissue remaining after cytokine quantification was

insufficient for these assays.

Cytokine quantification in brain and plasma

V-PLEX proinflammatory Panel 1 kits (Mouse) from MSD (Rockville, MD) were used to measure

cytokine concentrations in plasma, frontal cortex, and hippocampus. Plates arrived pre-coated with

primary antibodies for the simultaneous quantification of Interferon- γ (IFN-γ), Interleukin-1β (IL-1β),

Interleukin-2 (IL-2), Interleukin-4 (IL-4), Interleukin-5 (IL-5), Interleukin-6 (IL-6), chemokine (C-X-C

motif) ligand 1 (CXCL1), Interleukin-10 (IL-10), Interleukin-12p70 (IL-12p70), and tumor necrosis

factor-α (TNF-α). This panel was chosen because it allows for the quantification of several pro- and

anti-inflammatory cytokines as well as the chemokine CXCL1, thus provides a comprehensive view of

the cytokine milieu. All samples were run in duplicates, and sample and secondary antibody

incubations were performed according to manufacturer instructions. Lower limits of detection (LLOD),

which differed between analytes and plates (2 plates total), were as follows: IFN-γ: 0.033-0.076 pg/ml;

IL-1β: 0.048-0.069 pg/ml; IL-2: 0.32-0.79pg/ml; IL-4: 0.077-0.091 pg/ml; IL-5: 0.097-0.10 pg/ml; IL-

6: 0.83-0.85 pg/ml; CXCL1: 0.15-0.17 pg/ml; IL-10: 0.18-0.62 pg/ml; IL -12p70: 11.8-13.1 pg/ml;

TNF-α: 0.082-0.096 pg/ml. In plasma, IL -12p70 was not detectable in 75% of samples, and IL-4 was

not detectable in 79% of samples, and thus were not included in analyses.

Cell signaling phosphoprotein and PSD-95 quantification in the frontal cortex

179 Cell signaling phosphoproteins (phospho-ERK1/2 (pERK1/2), pMEK1/2, pJNK, pp33, and pSTAT3)

180 and PSD-95 were measured in frontal cortex samples using kits from MSD (Rockville, MD). Primary

181 antibody pre-coated plates were used for sample and secondary antibody incubations according to

182 manufacturer's instructions, with samples run in duplicates, and results reported after normalization to

183 total protein concentrations in each sample.

Radioimmunoassays for hormone quantification in plasma

Plasma corticosterone concentrations from samples collected at euthanasia were quantified using a

commercially available radioimmunoassay kit used in accordance with manufacturers' instructions

(corticosterone double-antibody radioimmunoassay kit, MP biomedicals, Solon, OH). To avoid

handling-induced elevations in circulating glucocorticoids, trunk blood samples were collected within 2

minutes of the experimenter entering the colony room. The inter- and intra-assay coefficients of

variation were below 10%.

An Ultra-Sensitive Estradiol Radioimmunoassay kit (Beckman Coulter, Prague, Czech Republic) was

used according to manufacturers' instructions to measure 17β-estradiol concentrations in plasma

samples collected at euthanasia. Assay sensitivity is 2.2pg/ml, and average inter- and intra-assay

coefficients of variation were below 10%.

Vaginal cytology for estrous cycle assessment

Estrous cycle stage can affect behaviour, measures of neural plasticity, and immune function (Beagley

and Gockel, 2003; Meziane et al., 2007; Woolley et al., 1990), thus vaginal lavage samples were

obtained on behavioural testing days, and estrous stage was determined according to previous methods

(Cora et al., 2015). Plasma was collected on the last day for estradiol quantification. We expected

irregular estrous cycling in at least a proportion of sham-operated mice as they were middle-aged at

testing. Mice in persistent diestrus, persistent estrus, or that displayed abnormalities in the length or

order of estrous cycle stages were all classified as irregularly cycling, as we have done previously

(Galea et al., 2018). Ovariectomized mice were also lavaged to control for potential effects of the

procedure.

Statistical analyses

All statistical analyses were performed using Statistica software (Tulsa, OK). Behavioural measures,

cytokine concentrations, IL-6:IL-10 ratio, corticosterone concentrations, PSD-95, and cell signaling

protein levels were each analyzed using factorial analysis of variance (ANOVA), with ovarian status

(OVX, Sham) and stress condition (stress, non-stress) as the between-subject factors. Body mass was

used as a covariate when analyzing sucrose consumption, immobility in the forced swim test, self-

grooming behaviour in the splash test, and adrenal mass. Estrous cycle stage (proestrus, non-proestrus)

was used as a covariate in behavioural analyses. 17β-estradiol concentrations were used as a covariate

in corticosterone, cytokine, cell signaling, and PSD-965 analyses. Covariate effects are only mentioned

when significant. Post-hoc analyses utilized Newman-Keul's comparisons and any *a priori*

comparisons were subjected to a Bonferroni correction. A Chi-square test was used to compare the

frequency of mice that had regular estrous cycles between sham-operated groups. Pearson's

correlations were performed between variables of interest. Outliers that fell more than 2.5 standard

deviations away from the mean were removed from analyses. Principal Component Analyses (PCA)

were used to reduce the cytokine data into a smaller number of uncorrelated variables and to obtain

information about the amount of variance accounted for by potential cytokine networks within the data.

PCAs were followed by ANOVAs on individual principal component scores as we have done

previously (Eid et al., 2019a), to assess the effects of ovarian status and stress exposure on cytokine

networks.

Results

Body mass was significantly reduced across days of stress exposure regardless of ovarian status

Long-term ovariectomy significantly increased body mass ($F(1, 24)=22.72$, $p<0.0001$), therefore to

investigate the effects of sub-chronic stress exposure we calculated body mass as a percentage of mass

on experimental day 1. Body mass percentage decreased significantly across the 6 days of stress

exposure ($F(4, 96)=32.967$, $p<0.0001$; stress condition by day interaction; **Fig. 2**). Specifically, in

stress-exposed groups, there was a significant decline on day 3 relative to day 2, and on every day

thereafter (p's <0.006). Body mass percentage was also significantly different between stress and non-

stress groups on days 3-6 (p's <0.007). There were also significant main effects of stress condition and

day (p's<0.0001), a trend toward a significant day by ovarian status interaction ($p=0.064$), but no other

significant main effects or interactions (p's >0.095).

Figure 2. Body mass across days, shown as a percentage of body mass on experimental day 1. Regardless of ovarian status, body mass decreased significantly across days of sub-chronic stress exposure; * indicates p<0.007, significantly different from day-matched non-stress groups, and stress groups on prior day/s of stress exposure. Data in means ± standard error of the mean. OVX, ovariectomized.

Long-term ovariectomy reduced circulating 17β-estradiol concentrations but stress exposure did

not affect 17β-estradiol concentrations or estrous cycling

Plasma 17β-estradiol concentrations were significantly reduced by long-term ovariectomy (F(1,

18)=9.20, p=0.007; main effect of ovarian status). Stress condition did not affect 17β-estradiol

concentrations, nor did the interaction of stress condition and ovarian status (all p's >0.2; **Table 1**).

Within sham-operated groups, there was no significant difference in the frequency of mice that had

regular estrous cycles ($\chi^2(1) = 0.31$, p = 0.58; **Table 1**).

Table 1. Plasma 17β-estradiol concentrations from trunk blood at decapitation and percentage of sham-operated mice with regular estrous cycles.

Group	17β-estradiol (pg/ml)	% of mice regularly cycling
Sham, non-stress	10.34. ± 1.85	28.57
Sham, stress	10.32 ± 2.20	42.86
OVX, non-stress	4.30 ± 0.96	N/A
OVX, stress	7.52 ± 0.71	N/A

Data in means ± standard error of the mean. OVX, ovariectomized.

Basal plasma corticosterone concentrations and adrenal mass did not differ significantly between

groups

Basal plasma corticosterone concentrations did not differ significantly between groups (all p's >0.3;

Table 2), but there was a significant covariate effect of plasma 17β-estradiol concentrations (p =

0.027). Similarly, adrenal mass was not significantly affected by ovarian status, stress exposure, nor
their interaction (p's>0.3; **Table 2**).

Table 2. Basal plasma corticosterone concentrations and adrenal mass.

Group	CORT (pg/ml)	Adrenal mass (mg)
Sham, non-stress	42.48 ± 8.99	6.99 ± 0.66
Sham, stress	33.79 ± 6.70	8.06 ± 0.53
OVX, non-stress	31.92 ± 6.88	8.50 ± 0.88
OVX, stress	39.58 ± 8.62	8.16 ± 0.69

Data in means ± standard error of the mean. OVX, ovariectomized; CORT, corticosterone.

Ovariectomy reduced latency to immobility under non-stress conditions, and sub-chronic stress
exposure reduced latency to immobility in sham-operated mice only

Under non-stress conditions, ovariectomy significantly decreased latency to immobility in the FST
(p=0.01; a priori comparisons; ovarian status by stress condition interaction: $F(1, 18)=1.8372$, p=0.19).
Further, exposure to sub-chronic stress significantly decreased the latency to immobility in sham-
operated (p=0.012) but not in ovariectomized mice (p=0.54). There was also a significant main effect
of stress condition (p=0.025) but not of ovarian status (p=0.22). The percentage of time spent immobile
in FST did not significantly differ between groups (all p's >0.09; **Table 3)**

Sub-chronic stress exposure decreased sucrose consumption regardless of ovarian status

Stress exposure significantly reduced sucrose consumption ($F(1, 23)=4.87$, p=0.038; main effect of
stress condition; **Fig. 3B**). There was also a trend for ovariectomy to reduce sucrose consumption ($F(1,$
$23)=3.98$, p=0.067), but no other significant effects (p's>0.4). In comparison, water consumption was
not significantly affected by body mass, ovarian status, stress condition, nor their interaction (all p's
>0.4; **Table 3**).

Figure 3. Latency to first bout of immobility in the Forced Swim Test and sucrose consumption in the Sucrose Consumption Test. **(A)** Ovariectomy significantly reduced latency to immobility under non-stress conditions, and sub-chronic stress exposure reduced latency to immobility in sham-operated mice only; # indicates p=0.01 and *indicates p=0.012. **(B)** Stress exposure decreased sucrose consumption, regardless of ovarian status, and there was a trend for ovariectomy to reduce sucrose consumption; * indicates p=0.038, main effect of stress condition. FST = forced swim test; OVX = ovariectomy. Data in means + standard error of the mean

262 **Sub-chronic stress exposure and ovarian status did not significantly affect behaviour in the**

263 **splash and novelty suppressed feeding tests**

264 Anxiety-like behavior as measured in the novelty suppressed feeding test and self-grooming behaviour

265 in the splash test were not significantly altered by ovariectomy, sub-chronic stress exposure, nor their

266 interaction (all p's >0.1; **Table 3**).

Table 3. Behavioural data.

Group	% time spent immobile in FST	Latency to feed in NSF (s)	% time spent grooming in ST	H_2O consumption (g)
Sham, non-stress	64.96 ± 5.37	542.83 ± 36.35	14.35 ± 3.04	8.29 ± 0.81
Sham, stress	58.79 ± 2.94	521.29 ± 40.01	19.98 ± 1.53	7.00 ± 0.44
OVX, non-stress	73.62 ± 2.05	507.86 ± 59.72	14.71 ± 4.31	8.00 ± 0.79
OVX, stress	65.88 ± 3.55	600.00 ± 0.00	18.32 ± 2.78	9.14 ± 1.26

Data in means ± standard error of the mean. OVX, ovariectomized; FST, forced swim test; NSF, novelty suppressed feeding; ST, splash test.

267 **Principal component (PC) analyses in cytokine data:**

268 **Frontal cortex: PC1 scores were reduced by long-term ovariectomy, and PC2 scores were**

269 **increased by stress exposure in sham-operated mice only**

270 The model generated 3 principal components, accounting for 81.8% of the variance within the frontal

271 cortex cytokine dataset. Variance explained by principal component 1 (PC1) = 52.8%, PC2 = 17.1 %,

and PC3 = 11.9%. Factor loadings are shown in **Table 4**. ANOVA results reveal that ovariectomy significantly reduced PC1 scores ($F_{(1, 24)}$=5.23, p=0.031; main effect of ovarian status; **Fig. 4A**), but there was no significant main effect of stress condition, nor an ovarian status by stress condition interaction (p's >0.6). As such we will refer to PC1 as "Ovarian hormone sensitive" cytokines. Stress significantly increased PC2 scores in sham (p = 0.008), but not ovariectomized mice (p = 0.51; a priori comparisons, ovarian status by stress condition interaction: $F_{(1, 24)}$=2.526, p<0.13; **Fig. 4B**). There was also a significant main effect of stress condition (p = 0.019), but not of ovarian status (p = 0.55). For PC3 scores, there were no significant main effects of ovarian status or stress condition, nor a significant interaction (p's >0.19). Thus, in the frontal cortex we identify 8 cytokines that were proportionally more sensitive to ovarian hormone status based on component loadings(IL-2, IL-4 IL5, IL-6, IL-10, TNF-α, IL-12p70 and the chemokine CXCL1) and 3 cytokines sensitive to both stress and ovarian hormone status (IL-1β, TNF-α and IL-4).

Hippocampus: Stress exposure increased PC2 scores regardless of ovarian status, and there were trends for long-term ovariectomy to increase PC1 and decrease PC2 scores

The model generated 3 principal components, accounting for 81.4% of the variance within the hippocampus cytokine dataset. Variance explained by principal component 1 (PC1) = 58.0%, PC2 = 14.8%, and PC3 = 8.6%. Factor loadings are shown in **Table 4**. ANOVAs reveal a weak trend for ovariectomy to increase PC1 scores ($F_{(1, 23)}$=3.02, p=0.096; main effect of ovarian status; **Fig. 4C**), and no significant main effect of stress condition, nor an ovarian status by stress condition interaction (p's >0.3). Stress significantly increased PC2 scores regardless of ovarian status ($F_{(1, 23)}$=4.40, p=0.047; **Fig. 4D**), and there was a trend toward significance for ovariectomy to decrease PC2 scores ($F_{(1, 23)}$=3.67, p=0.068), but no significant stress condition by ovarian status interaction (p > 0.6). For PC3 scores, there were no significant main effects of ovarian status or stress condition, nor a significant interaction (p's >0.3). Thus, these analyses in the hippocampus identify 8 cytokines that were sensitive to ovarian hormone status based on component loadings (IL-2, IL-4 IL5, IL-6, IL-10, TNF-α, IL-12p70, IFN-γ) and 3 cytokines sensitive to stress (IL-1β, TNF-α and the chemokine CXCL1).

Plasma: PC scores did not differ significantly between groups

The model generated 2 principal components, accounting for 48.2% of the variance within the plasma cytokine dataset. Variance explained by principal component 1 (PC1) = 29.0%, and PC2 = 19.2%. Factor loadings are shown in **Table 4**. ANOVAs reveal no significant main effects of stress condition or ovarian status, nor a significant interaction for PC1 or PC2 scores (p's > 0.06; **Fig. 4E-F**).

Table 4. Principal component analyses loading table

Cytokine	Frontal cortex			Hippocampus			Plasma	
	PC1	**PC2**	**PC3**	**PC1**	**PC2**	**PC3**	**PC1**	**PC2**
IFN-γ	0.22	-0.30	**0.91**	**0.85**	-0.03	-0.24	**0.59**	0.17
IL-10	**0.93**	-0.21	0.25	**0.90**	-0.19	0.09	-0.25	0.25
IL-1β	0.22	**0.85**	0.36	0.45	**0.58**	-0.35	**0.59**	**-0.67**
IL-2	**0.90**	0.00	-0.08	**0.89**	-0.19	-0.17	**0.73**	-0.25
IL-5	**0.87**	-0.23	0.08	**0.65**	-0.30	0.71	0.24	**0.65**
IL-6	**0.77**	0.10	-0.09	**0.88**	-0.007	0.19	**0.67**	0.24
CXCL1	**0.73**	0.35	-0.30	0.17	**1.13**	**0.44**	**0.65**	0.03
TNF- α	**0.75**	**0.47**	0.18	**0.64**	**0.54**	-0.31	0.26	**0.67**
IL-12p70	**0.90**	-0.11	-0.17	**0.92**	-0.16	0.064	N/A	N/A
IL-4	**0.49**	**-0.66**	0.04	**0.89**	-0.11	0.09	N/A	N/A

High loadings indicated in bold.

Figure 4. Principal component analyses scores in cytokine data. **(A)** Long-term ovariectomy significantly decreased Principal Component (PC) 1 scores in frontal cortex; * indicates p=0.031, main effect of ovarian status. **(B)** Exposure to sub-chronic stress increased PC2 scores in the frontal cortex in sham-operated mice only; * indicates p=0.008, significantly higher than non-stress sham-operated group. **(C)** There was a weak trend (p=0.096) for ovariectomy to increase PC1 scores in the hippocampus. **(D)** Sub-chronic stress exposure increased PC2 scores in the hippocampus, regardless of ovarian status, and there was a trend for ovariectomy (p=0.064) to reduce PC2 scores; * indicates p=0.047, main effects of stress condition. There were no significant group differences in PC1 or PC2 scores in plasma **(E-F)**. PC = principal component; OVX = ovariectomy. Data in means + standard error of the mean.

Individual cytokine ANOVAs:

Long-term ovariectomy reduced the concentrations of several cytokines in the frontal cortex

Long-term ovariectomy significantly reduced the concentrations of the following cytokines/chemokine in the frontal cortex: TNF-α ($F_{(1, 24)}$=5.41, p=0.029; **Fig. 5A**); CXCL1 ($F_{(1, 24)}$=4.44, p=0.046; **Fig. 5B**); IL-10 ($F_{(1, 22)}$=9.80, p=0.0049; **Fig. 5C**); IL-12p70 ($F_{(1, 24)}$=6.40, p=0.018; **Fig. 5D**), IFN-γ ($F_{(1, 23)}$=5.89, p=0.023; **Fig. 5E**), and IL-6 ($F_{(1, 24)}$=4.16, p=0.05; **Fig. 5F**). For each of these cytokines, there was no significant main effect of stress condition, and no stress condition by ovarian status interaction (p's > 0.2).

Sub-chronic stress exposure significantly increased IL-1β in the frontal cortex of sham-operated mice only

Exposure to stress increased IL-1β concentrations in the frontal cortex of sham-operated (p=0.018, a priori comparison) but not ovariectomized mice (p=0.98; $F_{(1, 24)}$=3.1706, p=0.088; stress condition by ovarian status interaction; **Fig. 5G**). There was also a trend toward a significant main effect of stress exposure to increase IL-1β (p= 0.082), but no significant main effect of ovarian status (p = 0.32).

Sub-chronic stress exposure significantly decreased IL-4 in the frontal cortex

Stress exposure significantly reduced IL-4 concentrations in the frontal cortex ($F_{(1, 24)}$=5.28, p=0.031; main effect of condition; **Fig. 5H**). This effect was driven but a larger decrease in sham-operated mice, as a priori comparisons reveal a trend toward a significant difference between sham-operated groups (p = 0.041) but not between ovariectomized groups (p = 0.28). There was also a trend toward significance for ovariectomy to reduce IL-4 (p = 0.085), but no significant stress condition by ovarian status interaction (p = 0.46). There were no significant effects of stress condition or ovarian status, nor an interaction to affect IL-2 (P's > 0.25; **Fig. 5I**) or IL-5 (p's>0.25; **Fig. 5J**) concentrations in the frontal cortex.

Figure 5. Cytokine concentrations in the frontal cortex, normalized by total protein concentrations. Long-term ovariectomy reduced the concentrations of TNF-α **(A)** CXCL1 **(B)**, IL-10 **(C)**, IL-12p70 **(D)**, IFN-γ **(E)**, and IL-6 **(F)** regardless of stress exposure; * indicates significant main effects of ovarian status (p's <0.05). **(G)** Stress exposure increased IL-1β in the frontal cortex of sham-operated mice only; * indicates p = 0.018, significantly higher than non-stress sham group. **(H)** Stress exposure significantly reduced IL-4; * indicates p = 0.031, main effect of stress condition. Concentrations of IL-2 **(I)** and IL-5 **(J)** did not differ significantly between groups. OVX, ovariectomized; IL, interleukin; CXCL1, chemokine (C-X-C motif) ligand 1; TNF-α, tumor necrosis factor-α; IFN-γ, Interferon- γ. Data in means + standard error of the mean.

Long-term ovariectomy increased IL-2, IL-5, and IL-6 concentrations in the hippocampus

Long-term ovariectomy significantly increased IL-2 ($F_{(1, 22)}=8.92$, $p<0.007$; **Fig. 6A**), IL-5 ($F_{(1, 22)}=8.37$, $p=0.008$; **Fig. 6B**), and IL-6 ($F_{(1, 23)}=5.80$, $p=0.024$; **Fig. 6C**) concentrations in the hippocampus. For each of these cytokines, there was no significant main effect of stress condition, nor a stress condition by ovarian status interaction (all p's >0.18).

Sub-chronic stress exposure increased IL-1β in the hippocampus of sham-operated mice only, and decreased IL-12p70 concentrations regardless of ovarian status

Sub-chronic stress exposure increased IL-1β concentrations in the hippocampus of sham-operated ($p=0.04$) but not ovariectomized mice ($p=0.78$; $F_{(1, 23)}=1.82$, $p=0.19$; a priori comparison; stress condition by ovarian status interaction; **Fig. 5D**). There was a weak trend toward significance for a main effect of stress condition ($p=0.09$), and no significant main effect of ovarian status ($p>0.58$). Regardless of ovarian status, exposure to sub-chronic stress significantly decreased hippocampal IL-12p70 concentrations ($F_{(1, 23)}=4.38$, $p=0.048$; **Fig. 6E**). There was no significant main effect of ovarian status nor an ovarian status by stress condition interaction (p's >0.1). There was also a weak trend for stress exposure to reduce IL-10 ($p=0.082$; **Fig. 6F**), but no significant group differences for hippocampal IL-4 (p's > 0.2; **Fig. 6G**), IFN- γ (p's > 0.29; **Fig. 6H**), TNF-α (p's > 0.27; **Fig. 6I**), or CXCL1 (p's > 0.3; **Fig. 6J**).

Figure 6. Cytokine concentrations in the hippocampus, normalized by total protein concentrations. Long-term ovariectomy significantly increased concentrations of IL-2 **(A)**, IL-5 **(B)**, and IL-6 **(C)** in the hippocampus; * indicates p<0.03, main effect of ovarian status**. (D)** Stress exposure increased IL-1β in the hippocampus of sham-operated mice only; * indicates p = 0.04, significantly higher than non-stress sham group. **(E)** Sub-chronic stress exposure significantly decreased IL-12p70 concentrations, regardless of ovarian status; * indicates p<0.05, main effect of stress condition. Hippocampal concentrations of IL-10 **(F)**, IL-4 **(G)**, IFN- γ **(H)**, TNF-α **(I)**, and CXCL1 **(J)** did not significantly differ between groups. OVX, ovariectomized; IL, interleukin; CXCL1, chemokine (C-X-C motif) ligand 1; TNF-α, tumor necrosis factor-α; IFN-γ, Interferon- γ. Data in means + standard error of the mean.

Ovariectomy significantly increased plasma CXCL1 concentrations

Ovariectomy significantly increased CXCL1 concentrations in plasma ($F_{(1,20)}$=6.62, p=0.018; main effect of ovarian status; **Fig 7A**), but there was no significant main effect of stress condition (p=0.23) nor an ovarian status by stress condition interaction (p = 0.53). There were trends toward significance for an ovarian status by stress condition interaction to affect plasma IL-10 ($F_{(1, 20)}$=3.85, p=0.064; **Fig. 7B**) and IL-6 ($F_{(1, 19)}$=3.2520, p=0.087; **Fig. 7C**) concentrations, but no significant main effects of ovarian status nor stress condition in both cases (p's > 0.19). There were no significant main effects of stress condition or ovarian status nor a significant interaction for plasma IL-5 (all p's >0.29; **Fig. 7D**), TNF-α (p's > 0.14; **Fig. 7E**), IFN-γ (p's > 0.4; **Fig. 7F**), IL-2 (p's > 0.6; **Fig. 7G**), or IL-1β (p's > 0.2; **Fig. 7H**).

Figure 7. Cytokine concentration in plasma. **(A)** Ovariectomy significantly increased CXCL1 concentrations in plasma; * indicates p= 0.018, main effect of ovarian status. There were no significant group differences in plasma concentrations of IL-10 **(B)**, IL-6 **(C)**, IL-5 **(D)**, TNF-α **(E)**, IFN-γ **(F)**, IL-2 **(G)**, or IL-1β **(H)**. OVX, ovariectomized; IL, interleukin; CXCL1, chemokine (C-X-C motif) ligand 1; TNF-α, tumor necrosis factor-α; IFN-γ, Interferon- γ. Data in means + standard error of the mean.

Sub-chronic stress exposure increased IL-6:IL-10 ratio in the frontal cortex and hippocampus in

sham-operated mice only

IL-6:IL-10 ratios were analyzed as an indicator of pro- to anti-inflammatory cytokine balance as

previously described (de Brito et al., 2016; Wood et al., 2015). In both brain regions, sub-chronic stress

exposure significantly increased IL-6:IL-10 ratio in sham-operated mice only (frontal cortex: p =

0.033; ovarian status by stress condition interaction: $F_{(1, 22)}$=4.72, p=0.041, **Fig 8A**; hippocampus: p =

0.015; a priori comparisons; stress condition by ovarian status interaction: $F_{(1, 23)}=1.9553$, p=0.17,

Fig. 8B). In the hippocampus, there was also a significant main effect of condition, with stress

increasing the ratio (p=0.025), but not of ovarian status (p=0.87), and there were no significant main

effects in the frontal cortex (p's >0.12). Finally, under non-stress conditions, ovariectomy significantly

increased IL-6:IL-10 ratio in the frontal cortex only (p = 0.014; **Fig. 8A**). For plasma IL-6:IL-10 ratios,

there were no significant main effects, nor a significant interaction (all p's >0.3).

Figure 8. Interleukin-6 to interleukin-10 ratio in the frontal cortex **(A)**, hippocampus **(B)**, and plasma **(B)**. **(A)** Ovariectomy significantly increased IL-6:IL-10 ratio in the frontal cortex, and sub-chronic stress exposure increased IL-6:IL-10 ratio in sham-operated mice only. * indicates p= 0.014 and # indicates p=0.033, significantly higher than non-stress sham group. **(B)** sub-chronic stress exposure increased IL-6:IL-10 ratio in the hippocampus in sham-operated mice only. # indicates p=0.015, significantly higher than non-stress sham group. **(C)** There were no significant group differences in plasma IL-6:IL-10 ratio. OVX, ovariectomized; IL, interleukin. Data in means + standard error of the mean.

pERK1/2 and pMEK1/2 were significantly reduced by long-term ovariectomy and the effects of

stress were dependent on ovarian status

Ovariectomy significantly decreased pERK1/2 and pMEK1/2 expression under non-stress conditions

relative to sham-operated mice (both p's = 0.032). Sub-chronic stress exposure significantly decreased

pERK1/2 expression in sham-operated mice (p= 0.023), but not in ovariectomized mice (p=0.35;

ovarian status by stress condition interaction: $F_{(1, 24)}=5.68$, p=0.025 , **Fig. 9A**). Conversely, sub-

chronic stress exposure significantly increased pMEK1/2 expression in ovariectomized mice (p=

0.042), but not in sham-operated mice (p = 0.31; significant ovarian status by stress condition

interaction $F_{(1, 24)}=5.065$, p=0.034, **Fig 9B**). A similar, albeit non-significant trend was observed for

pp38 expression (ovarian status by stress condition interaction ($F_{(1, 24)}=3.844$, p=0.062; **Fig. 9C**), and

there were no significant main or interaction effects for pJNK or pSTAT3 expression (all p's >0.14; **Fig**

9D-E).

Figure 9. Cell signaling phosphoprotein expression in the frontal cortex, normalized by total protein levels. (**A**) Under non-stress conditions, ovariectomy decreased pERK1/2 expression, and stress exposure decreased pERK1/2 expression in sham-operated mice only; * indicates $p = 0.032$, significant difference between non-stress groups; # indicates $p = 0.023$, relative to non-stress sham-operated mice. (**B**) Under non-stress conditions, ovariectomy decreased pMEK1/2 expression, and stress exposure increased pMEK1/2 expression in ovariectomized mice only; * indicates $p = 0.032$, significant difference between non-stress groups; # indicates $p = 0.042$, relative to non-stress ovariectomized mice. There were no significant group differences in pp38 (**C**), pJNK (**D**), and pSTAT3 (**F**) expression. Data in means + standard error of the mean. OVX, ovariectomized; ERK, extracellular signal-regulated kinase; JNK, c-Jun NH$_2$-terminal kinase. Data in means + standard error of the mean.

PSD-95 expression was not significantly affected by ovariectomy or sub-chronic stress exposure

Expression of the postsynaptic scaffolding protein PSD-95 was utilized as a synaptic marker and a proxy measure of excitatory synapse density (Wainwright et al., 2016). PSD-95 expression was previously found to be reduced by approximately 40% in the PFC of individuals with depression (Feyissa et al., 2009), Here, PSD-95 expression in the frontal cortex was not significantly affected by stress condition, ovarian status, nor their interaction (all p's >0.4; **Table 5**).

Table 5. Postsynaptic density protein-95 expression in the prefrontal cortex**.**

Group	PSD-95 (signal/ug protein)
Sham, non-stress	376.11 ± 53.53
Sham, stress	412.56 ± 54.68
OVX, non-stress	415.88 ± 31.09
OVX, stress	445.36 ± 18.38

Data in means ± standard error of the mean. OVX, ovariectomized; PSD-95, postsynaptic density protein-95.

Reduced latency to immobility in the forced swim test was associated with increased IL-6:IL-10 in the frontal cortex and hippocampus

Pearson's correlations were performed between behavioural measures and markers of neuroinflammation that were significantly affected by stress exposure in an ovarian status-dependent manner. We found that reduced latency to immobility in the forced swim test was significantly associated with increased IL-6:IL-10 in the frontal cortex (r = - 0.43, p = 0.048; **Fig. 10A**) and hippocampus (r = - 0.43, p = 0.044; **Fig. 10B**). However, there were no significant correlations between latency to immobility and IL-1β concentrations in the frontal cortex (r = - 0.11, p = 0.6) or hippocampus (r = 0.04, p = 0.85).

Figure 10. Association between passive-coping behaviour and pro- to anti-inflammatory cytokine balance in the frontal cortex **(A)** and hippocampus **(B)**. Reduced latency to immobility was significantly associated with increased IL-6:IL-10 in the frontal cortex (**A**; r = - 0.43, p = 0.048) and hippocampus (**B**; r = - 0.43, p = 0.044). FST, forced swim test; IL, interleukin.

Discussion

Here, we report profound effects of long-term ovariectomy on the central cytokine milieu in middle-aged female mice, evidenced by reduced cytokine concentrations in the frontal cortex but increased cytokine concentrations in the hippocampus. Long-term ovariectomy in middle-age also reduced phosphorylated ERK1/2 and MEK1/2 expression in the frontal cortex. Along with these neural changes in the frontal cortex and hippocampus, long-term ovariectomy in middle-age resulted in a modest increase in depressive-like behaviour under non-stress conditions. We further show ovarian status-dependent effects of sub-chronic stress exposure on cytokine concentrations in the frontal cortex and hippocampus, as seen by increased IL-1β and a shift toward a pro-inflammatory cytokine bias (IL-6:IL-10) in sham-operated mice only. Sub-chronic stress exposure also decreased expression of

phosphorylated ERK1/2 in the frontal cortex in sham-operated mice only. Importantly, this was coupled

with a greater behavioural susceptibility to sub-chronic stress in sham-operated mice. These data

suggest that ovarian hormones in middle-age exert a powerful influence on the neuroimmune

environment in a region- dependent manner, and may dictate certain behavioural and

neuroinflammatory consequences of sub-chronic stress exposure.

Long-term ovariectomy modifies the cytokine signature under non-stress conditions and in

response to sub-chronic stress exposure

We observed brain region-specific effects of long-term ovariectomy (9 months) on cytokine

concentrations, with reductions in the frontal cortex and modest elevations in the hippocampus. Indeed,

more than 50% of the variance within the frontal cortex cytokine data was explained by Principal

Component 1 (PC1), which appeared to be largely accounted for by ovarian status. Ovarian status also

affected the hippocampal cytokine milieu, although to a lesser extent than in the frontal cortex. Overall,

the principal component loadings indicated 7 overlapping cytokines were important contributors to the

first principle component (PC1) that was modified by long-term ovariectomy albeit in opposing

directions between the frontal cortex and hippocampus (IL-2, IL-4, IL-5, IL6, IL-10, IL-12p70, TNFα).

In terms of individual ANOVA analyses more, distinct sets of cytokines were affected by long-term

ovariectomy in the two brain regions, with the exception of IL-6 which was increased in the

hippocampus but decreased in the frontal cortex. The effects of sub-chronic stress exposure on cytokine

concentrations in the frontal cortex and hippocampus were less pronounced. This is not surprising

given the sub-chronic nature of the stress paradigm (6 days), thus we would expect longer durations of

stress exposure to result in larger alterations in the cytokine milieu. However, component loadings

indicate IL-1β, TNF-α and the chemokine CXCL1 as important contributors to PC2 in the

hippocampus, which showed significant effects of sub-chronic stress exposure. It is important to note

that all mice underwent behavioural testing, including the forced swim test which is an acute stressor.

Therefore, it is plausible that exposing non-stress controls to behavioural testing may have masked

some of the effects of sub-chronic stress exposure on cytokine concentrations.

An intriguing finding was that ovarian status impacted the neuroinflammatory consequences of

sub-chronic stress exposure, with overall more pronounced effects in sham-operated mice. Specifically,

in the frontal cortex and hippocampus of sham-operated mice only, stress exposure increased

concentrations of the pro-inflammatory IL-1β and resulted in a shift toward a pro-inflammatory

cytokine bias (IL-6:IL-10). The exaggerated neuroinflammatory response to stress in sham-operated

mice was observed in tandem with a greater behavioural susceptibility to sub-chronic stress exposure

(discussed below). Our findings in sham-operated females corroborate studies in male subjects, in

which IL-1β was increased by chronic stress exposure in the brain, and further found to be directly

implicated in the behavioural consequences of stress exposure (Goshen et al., 2008; Goshen and

Yirmiya, 2009; Wohleb et al., 2014). Further, the role of IL-6 in depression and stress-based models is

well-established (reviewed in Hodes et al., 2016), and a shift towards a higher IL-6: IL-10 ratio is

associated with passive-coping in response to social defeat stress in male rats (Wood et al., 2015).

Thus, our findings suggest that in middle-aged females, ovarian hormones may potentiate the

inflammatory consequences of stress exposure.

Ovarian hormones, especially 17β-estradiol, have well established anti-inflammatory properties

(Bruce-Keller et al., 2000; Vegeto et al., 2008). Consistent with this are the observed effects of long-

term ovariectomy to increase IL-6:IL-10 ratio in the frontal cortex and IL-6 concentrations in the

hippocampus under non-stress conditions. Given the role of IL-6 in MDD (Dowlati et al., 2010; Liu et

al., 2012), these observations may be related to the modest increase in depressive-like behaviour

induced by long-term ovariectomy under non-stress conditions in our middle-aged mice, which is in

line with findings in humans (Schmidt et al., 2015; Wilson et al., 2018) . However, in the frontal cortex

we also found that ovariectomy reduced concentrations of other proinflammatory cytokines including

IFN-γ and TNF-α. Further, in both the frontal cortex and hippocampus we observed an exaggerated

neuroinflammatory response to stress in sham-operated, rather than ovariectomized mice, highlighted

by increased IL-1β and increased IL-6:IL-10 ratio. Importantly, the anti-inflammatory and

neuroprotective effects of estrogens have been investigated mostly in younger females and in models of

brain injury or stroke, and in response to acute inflammatory challenges or *in vitro* (Selvamani and

Sohrabji, 2010; Suzuki et al., 2009; Vegeto et al., 2003). Age is a critical consideration here, as studies

using older females have observed neurotoxic effects of estrogens in similar models (Johnson and

Sohrabji, 2005; Nordell et al., 2003; Selvamani and Sohrabji, 2010), partially consistent with our

finding here that exposure to sub-chronic stress resulted in a larger neuroinflammatory response in

sham-operated middle-aged mice. Future studies need to consider age as a variable to further clarify the

role of ovarian hormones in the neuroinflammatory outcomes of stress exposure.

Long-term ovariectomy and sub-chronic stress exposure interact to affect ERK signaling

Long-term ovariectomy reduced phosphorylated ERK1/2 and its upstream activator MEK1/2 in the

frontal cortex under non-stress conditions, and stress exposure reduced pERK1/2 expression in sham-

operated mice only. The observed reduction with long-term ovariectomy may not be surprising, as this

group had significantly lower circulating concentrations of estradiol, which can rapidly activate

ERK1/2 (Bi et al., 2002; Fernandez et al., 2008). Perhaps more intriguing than baseline differences in

pERK1/2 expression were the differential effects of stress exposure, as abnormal ERK signaling has

been observed in post-mortem brains of individuals with MDD that died of suicide (Duric et al., 2010;

Dwivedi et al., 2006, 2001; Hsiung et al., 2003; Labonté et al., 2017). Notably, in human studies there

are reports of reduced ERK1/2 expression and activation in the prefrontal cortex and hippocampus of

males and females with MDD (Dwivedi et al., 2006, 2001; Hsiung et al., 2003), although data was not

stratified by sex. Our results contrast another study in which chronic stress exposure increased

pERK1/2 in the ventromedial PFC of female mice (Labonté et al., 2017), but this study used young

adult females whereas our subjects were middle-aged. Collectively, past studies indicate that

dysregulated ERK signaling is implicated in depression and the outcome of stress exposure and our

current findings extend previous work to suggest that ovarian hormones can influence the outcomes of

stress exposure on ERK signaling in females. In the present study, we only examined cell signaling

phosphoproteins in the frontal cortex, therefore future studies should examine these signaling pathways

in the hippocampus.

Long-term ovariectomy and stress did not significantly affect PSD-95 expression in the frontal

cortex in the present study. This finding is consistent with previous studies indicating that although

PSD-95 expression is sensitive to transient fluctuations in ovarian hormones (Spencer et al., 2010), it is

not modified two weeks post ovariectomy (Spencer-Segal et al., 2012; Waters et al., 2015, 2009; Zhang

et al., 2010). Several studies have found reductions in PSD-95 expression in response to chronic stress

exposure (Kallarackal et al., 2013; Kim and Leem, 2016; Pacheco et al., 2017) suggesting that the

shorter duration of stress exposure used in this study may not be sufficient to produce significant

modifications to PSD-95 expression.

Ovarian status influenced behaviour under non-stress conditions and in response to sub-chronic

stress exposure

Under non-stress conditions, long-term ovariectomy alone impacted some measures of depressive-like

behaviour, as seen by reduced latency to immobility in the forced swim test and a trend toward reduced

sucrose consumption. These behavioural differences may be linked to the reported effects of long-term

ovariectomy on cytokine concentrations in the brain. However, the depressogenic effects of long-term

ovariectomy were overall limited, as we did not observe significant differences between sham-operated

and ovariectomized mice in any other behavioural measure. This is partially consistent with other

studies in which the anxiogenic and depressogenic effects of ovariectomy (Estrada-Camarena et al.,

2011; Li et al., 2014) can recover after more prolong periods of ovarian hormone deprivation (de

Chaves et al., 2009; Estrada-Camarena et al., 2011).

Exposure to sub-chronic stress reduced latency to immobility in the forced swim test in sham-

operated mice only. This points to a greater susceptibility to stress-induced passive-coping behaviour,

which is in line with the exaggerated neuroinflammatory response to stress and the reduction in

pERK1/2 expression observed again in the sham-operated group. On the other hand, sub-chronic stress

exposure increased anhedonia-like behaviour and reduced body mass regardless of ovarian status,

suggesting similar susceptibilities in these domains. Interestingly, increased passive-coping behaviour

was associated with a pro-inflammatory cytokine bias (IL-6:IL-10) in the frontal cortex and

hippocampus suggesting that the greater susceptibility to stress-induced passive-coping behaviour in

sham-operated mice may be directly linked to the exaggerated neuroinflammatory response to stress.

Our current findings agree with a previous study in young adult female mice in which exposure

to a similar sub-chronic stress paradigm increased passive-coping behaviour in sham-operated, but not

ovariectomized mice (LaPlant et al., 2009). In contrast, other studies in rats and mice found that long-

term ovariectomy increased depressive-like behaviour after chronic stress exposure (Lagunas et al.,

2010; Mahmoud et al., 2016). These dissimilarities may be reconciled by comparing the duration of

exposure to stressors, which spanned 4-6 weeks in studies showing increased sensitivity to depressive-

like behaviour after long-term ovariectomy (Lagunas et al., 2010; Mahmoud et al., 2016), as opposed to

6 days in the current report and in LaPlant et al (2009), where increased sensitivity to stress is observed

in sham-operated females. Thus, together with previous studies our current data indicate that ovarian

hormones can confer either risk or resilience, depending on stress paradigm.

Conclusions

In conclusion, long-term ovarian hormone deprivation in middle-aged mice significantly altered the

neuroimmune environment in a brain region-specific manner, reduced expression of ERK pathway

phosphoproteins in the frontal cortex, and modestly increased depressive-like behaviour under non-

stress conditions. Importantly, ovarian status dictated some of the behavioural and neuroinflammatory

outcomes of sub-chronic stress exposure. Specifically, sham-operated mice had a greater

neuroinflammatory response to stress as seen by increased concentrations of IL-1β and increased IL-

6:IL-10 in the frontal cortex and hippocampus. We also observed stress-induced reductions in pERK1/2

expression in the frontal cortex of sham-operated mice only. This was coupled with an enhanced

behavioural susceptibility to sub-chronic stress exposure in sham-operated mice, as seen by increased

passive-coping behaviour in the forced swim test. These findings underscore the importance of

considering the immunomodulatory properties of ovarian hormones in stress and depression research.

Funding

This work was supported by a Canadian Institute of Health Research grant to LAMG (MOP 142308).

Declaration of interest

None.

References

Beagley, K.W., Gockel, C.M., 2003. Regulation of innate and adaptive immunity by the female sex
hormones oestradiol and progesterone. FEMS Immunol. Med. Microbiol. doi:10.1016/S0928-
8244(03)00202-5

Bi, R., Foy, M.R., Vouimba, R.-M., Thompson, R.F., Baudry, M., 2002. Cyclic changes in estradiol
regulate synaptic plasticity through the MAP kinase pathway. Proc. Natl. Acad. Sci.
doi:10.1073/pnas.241507698

Brechbuhl, J., Moine, F., Klaey, M., Nenniger-Tosato, M., Hurni, N., Sporkert, F., Giroud, C., Broillet,
M.-C., 2013. Mouse alarm pheromone shares structural similarity with predator scents. Proc. Natl.
Acad. Sci. doi:10.1073/pnas.1214249110

Bruce-Keller, A.J., Keeling, J.L., Keller, J.N., Huang, F.F., Camondola, S., Mattson, M.P., 2000.
Antiinflammatory effects of estrogen on microglial activation. Endocrinology 141, 3646–3656.
doi:10.1210/endo.141.10.7693

Brummelte, S., Galea, L.A.M., 2010. Depression during pregnancy and postpartum: Contribution of
stress and ovarian hormones. Prog. Neuro-Psychopharmacology Biol. Psychiatry.
doi:10.1016/j.pnpbp.2009.09.006

Burger, H.G., Dudley, E.C., Robertson, D.M., Dennerstein, L., 2002. Hormonal changes in the
menopause transition. Recent Prog. Horm. Res. 57, 257–75.

Can, A., Dao, D.T., Arad, M., Terrillion, C.E., Piantadosi, S.C., Gould, T.D., 2012. The mouse forced
swim test. J. Vis. Exp. e3638. doi:10.3791/3638

Cargnello, M., Roux, P.P., 2011. Activation and Function of the MAPKs and Their Substrates, the
MAPK-Activated Protein Kinases. Microbiol. Mol. Biol. Rev. doi:10.1128/MMBR.00031-10

Cohen, L.S., Soares, C.N., Vitonis, A.F., Otto, M.W., Harlow, B.L., 2006. Risk for New Onset of
Depression During the Menopausal Transition The Harvard Study of Moods and Cycles. Arch

Gen Psychiatry 63, 385–390. doi:10.1001/archpsyc.63.4.385

Cora, M.C., Kooistra, L., Travlos, G., 2015. Vaginal Cytology of the Laboratory Rat and

Mouse:Review and Criteria for the Staging of the Estrous Cycle Using Stained Vaginal Smears.

Toxicol. Pathol. 43, 776–793. doi:10.1177/0192623315570339

de Brito, R. de C.C.M., Lucena-Silva, N., Torres, L.C., Luna, C.F., Correia, J. de B., da Silva, G.A.P.,

2016. The balance between the serum levels of IL-6 and IL-10 cytokines discriminates mild and

severe acute pneumonia. BMC Pulm. Med. doi:10.1186/s12890-016-0324-z

de Chaves, G., Moretti, M., Castro, A.A., Dagostin, W., da Silva, G.G., Boeck, C.R., Quevedo, J.,

Gavioli, E.C., 2009. Effects of long-term ovariectomy on anxiety and behavioral despair in rats.

Physiol. Behav. doi:10.1016/j.physbeh.2009.03.016

Dowlati, Y., Herrmann, N., Swardfager, W., Liu, H., Sham, L., Reim, E.K., Lanctôt, K.L., 2010. A

Meta-Analysis of Cytokines in Major Depression. Biol. Psychiatry 67, 446–457.

doi:10.1016/j.biopsych.2009.09.033

Drevets, W.C., 2000. Functional anatomical abnormalities in limbic and prefrontal cortical structures in

major depression, in: Progress in Brain Research. doi:10.1016/S0079-6123(00)26027-5

Drevets, W.C., Price, J.L., Furey, M.L., 2008. Brain structural and functional abnormalities in mood

disorders: Implications for neurocircuitry models of depression. Brain Struct. Funct.

doi:10.1007/s00429-008-0189-x

Duman, R.S., Aghajanian, G.K., Sanacora, G., Krystal, J.H., 2016. Synaptic plasticity and depression:

New insights from stress and rapid-acting antidepressants. Nat. Med. doi:10.1038/nm.4050

Duric, V., Banasr, M., Licznerski, P., Schmidt, H.D., Stockmeier, C.A., Simen, A.A., Newton, S.S.,

Duman, R.S., 2010. A negative regulator of MAP kinase causes depressive behavior. Nat. Med.

16, 1328–1332. doi:10.1038/nm.2219

Dwivedi, Y., Rizavi, H.S., Conley, R.R., Pandey, G.N., 2006. ERK MAP kinase signaling in post-

mortem brain of suicide subjects: Differential regulation of upstream Raf kinases Raf-1 and B-

Raf. Mol. Psychiatry. doi:10.1038/sj.mp.4001744

Dwivedi, Y., Rizavi, H.S., Roberts, R.C., Conley, R.C., Pandey, G.N., 2001. Reduced activation and

expression of ERK1 / 2 MAP kinase in the post-mortem brain of depressed suicide subjects 916–

928.

Eid, R.S., Chaiton, J.A., Lieblich, S.E., Weinberg, J., Bodnar, T.S., Galea, L.A.M., 2019a. Early and

Late Effects of Maternal Experience on Hippocampal Neurogenesis, Microglia, and the

Circulating Cytokine Milieu. Neurobiol. Aging. doi:10.1016/j.neurobiolaging.2019.01.021

Eid, R.S., Gobinath, A.R., Galea, L.A.M., 2019b. Sex differences in depression: Insights from clinical
and preclinical studies. Prog. Neurobiol. doi:10.1016/J.PNEUROBIO.2019.01.006

Estrada-Camarena, E., López-Rubalcava, C., Hernández-Aragón, A., Mejía-Mauries, S., Picazo, O.,
2011. Long-term ovariectomy modulates the antidepressant-like action of estrogens, but not of
antidepressants. J. Psychopharmacol. doi:10.1177/0269881111408456

Felger, J.C., Haroon, E., Miller, A.H., 2015. Risk and resilience: Animal models shed light on the
pivotal role of inflammation in individual differences in stress-induced depression. Biol.
Psychiatry. doi:10.1016/j.biopsych.2015.04.017

Fernandez, S.M., Lewis, M.C., Pechenino, A.S., Harburger, L.L., Orr, P.T., Gresack, J.E., Schafe, G.E.,
Frick, K.M., 2008. Estradiol-Induced Enhancement of Object Memory Consolidation Involves
Hippocampal Extracellular Signal-Regulated Kinase Activation and Membrane-Bound Estrogen
Receptors. J. Neurosci. doi:10.1523/JNEUROSCI.1968-08.2008

Feyissa, A.M., Chandran, A., Stockmeier, C.A., Karolewicz, B., 2009. Reduced levels of NR2A and
NR2B subunits of NMDA receptor and PSD-95 in the prefrontal cortex in major depression. Prog.
Neuro-Psychopharmacology Biol. Psychiatry 33, 70–75. doi:10.1016/j.pnpbp.2008.10.005

Freeman, E.W., Sammel, M.D., Liu, L., Gracia, C.R., Nelson, D.B., Hollander, L., 2004. Hormones
and Menopausal Status as Predictors of Depression in Womenin Transition to Menopause. Arch.
Gen. Psychiatry 61, 62. doi:10.1001/archpsyc.61.1.62

Friedrich, M.J., 2017. Depression Is the Leading Cause of Disability Around the World. JAMA.
doi:10.1001/jama.2017.3826

Galea, L.A.M., McEwen, B.S., Tanapat, P., Deak, T., Spencer, R.L., Dhabhar, F.S., 1997. Sex
differences in dendritic atrophy of CA3 pyramidal neurons in response to chronic restraint stress.
Neuroscience 81, 689–697. doi:10.1016/S0306-4522(97)00233-9

Galea, L.A.M., Roes, M.M., Dimech, C.J., Chow, C., Mahmoud, R., Lieblich, S.E., Duarte-Guterman,
P., 2018. Premarin has opposing effects on spatial learning, neural activation, and serum cytokine
levels in middle-aged female rats depending on reproductive history. Neurobiol. Aging.
doi:10.1016/j.neurobiolaging.2018.06.030

Giefing-Kröll, C., Berger, P., Lepperdinger, G., Grubeck-Loebenstein, B., 2015. How sex and age
affect immune responses, susceptibility to infections, and response to vaccination. Aging Cell.
doi:10.1111/acel.12326

Goel, N., Workman, J.L., Lee, T.T., Innala, L., Viau, V., 2014. Sex differences in the HPA axis.
Compr. Physiol. 4, 1121–1155. doi:10.1002/cphy.c130054

Goshen, I., Kreisel, T., Ben-Menachem-Zidon, O., Licht, T., Weidenfeld, J., Ben-Hur, T., Yirmiya, R.,

2008. Brain interleukin-1 mediates chronic stress-induced depression in mice via adrenocortical

activation and hippocampal neurogenesis suppression. Mol. Psychiatry.

doi:10.1038/sj.mp.4002055

Goshen, I., Yirmiya, R., 2009. Interleukin-1 (IL-1): A central regulator of stress responses. Front.

Neuroendocrinol. doi:10.1016/j.yfrne.2008.10.001

Gouin, J.P., Glaser, R., Malarkey, W.B., Beversdorf, D., Kiecolt-Glaser, J., 2012. Chronic stress, daily

stressors, and circulating inflammatory markers. Heal. Psychol. doi:10.1037/a0025536

Gutiérrez-Lobos, K., Scherer, M., Anderer, P., Katschnig, H., 2002. The influence of age on the

female/male ratio of treated incidence rates in depression. BMC Psychiatry 2, 3.

doi:10.1186/1471-244X-2-3

Haapakoski, R., Mathieu, J., Ebmeier, K.P., Alenius, H., Kivimäki, M., 2015. Cumulative meta-

analysis of interleukins 6 and 1β, tumour necrosis factor α and C-reactive protein in patients with

major depressive disorder. Brain. Behav. Immun. 49, 206–215. doi:10.1016/j.bbi.2015.06.001

Hantsoo, L., Epperson, C.N., 2015. Premenstrual Dysphoric Disorder: Epidemiology and Treatment.

Curr. Psychiatry Rep. doi:10.1007/s11920-015-0628-3

Hendrick, V., Altshuler, L.L., Suri, R., 1998. Hormonal changes in the postpartum and implications for

postpartum depression. Psychosomatics 39, 93–101. doi:10.1016/S0033-3182(98)71355-6

Hodes, Georgia E, Kana, V., Menard, C., Merad, M., Russo, S.J., 2015. Neuroimmune mechanisms of

depression. Nat. Neurosci. 18, 1386–1393. doi:10.1038/nn.4113

Hodes, G.E., Ménard, C., Russo, S.J., 2016. Integrating Interleukin-6 into depression diagnosis and

treatment. Neurobiol. Stress. doi:10.1016/j.ynstr.2016.03.003

Hodes, G. E., Pfau, M.L., Purushothaman, I., Ahn, H.F., Golden, S.A., Christoffel, D.J., Magida, J.,

Brancato, A., Takahashi, A., Flanigan, M.E., Menard, C., Aleyasin, H., Koo, J.W., Lorsch, Z.S.,

Feng, J., Heshmati, M., Wang, M., Turecki, G., Neve, R., Zhang, B., Shen, L., Nestler, E.J.,

Russo, S.J., 2015. Sex Differences in Nucleus Accumbens Transcriptome Profiles Associated with

Susceptibility versus Resilience to Subchronic Variable Stress. J. Neurosci. 35, 16362–16376.

doi:10.1523/JNEUROSCI.1392-15.2015

Hollos, P., Marchisella, F., Coffey, E.T., 2018. JNK Regulation of Depression and Anxiety. Brain

Plast. 3, 145–155. doi:10.3233/bpl-170062

Hong, S., Beja-Glasser, V.F., Nfonoyim, B.M., Frouin, A., Li, S., Ramakrishnan, S., Merry, K.M., Shi,

Q., Rosenthal, A., Barres, B.A., Lemere, C.A., Selkoe, D.J., Stevens, B., 2016. Complement and

microglia mediate early synapse loss in Alzheimer mouse models. Science (80-.). doi:10.1126/science.aad8373

Howard, D.M., Adams, M.J., Clarke, T.-K., Hafferty, J.D., Gibson, J., Shirali, M., Coleman, J.R.I., Hagenaars, S.P., Ward, J., Wigmore, E.M., Alloza, C., Shen, X., Barbu, M.C., Xu, E.Y., Whalley, H.C., Marioni, R.E., Porteous, D.J., Davies, G., Deary, I.J., Hemani, G., Berger, K., Teismann, H., Rawal, R., Arolt, V., Baune, B.T., Dannlowski, U., Domschke, K., Tian, C., Hinds, D.A., Team, 23andMe Research, Consortium, M.D.D.W.G. of the P.G., Trzaskowski, M., Byrne, E.M., Ripke, S., Smith, D.J., Sullivan, P.F., Wray, N.R., Breen, G., Lewis, C.M., McIntosh, A.M., 2019. Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. bioRxiv 433367. doi:10.1101/433367

Hsiung, S.C., Adlersberg, M., Arango, V., Mann, J.J., Tamir, H., Liu, K.P., 2003. Attenuated 5-HT1A receptor signaling in brains of suicide victims: Involvement of adenylyl cyclase, phosphatidylinositol 3-kinase, Akt and mitogen-activated protein kinase. J. Neurochem. doi:10.1046/j.1471-4159.2003.01987.x

Isingrini, E., Camus, V., Le Guisquet, A.M., Pingaud, M., Devers, S., Belzung, C., 2010. Association between repeated unpredictable chronic mild stress (UCMS) procedures with a high fat diet: A model of fluoxetine resistance in mice. PLoS One. doi:10.1371/journal.pone.0010404

Johnson, A.B., Sohrabji, F., 2005. Estrogen's effects on central and circulating immune cells vary with reproductive age. Neurobiol. Aging. doi:10.1016/j.neurobiolaging.2004.12.006

Kallarackal, A.J., Kvarta, M.D., Cammarata, E., Jaberi, L., Cai, X., Bailey, A.M., Thompson, S.M., 2013. Chronic Stress Induces a Selective Decrease in AMPA Receptor-Mediated Synaptic Excitation at Hippocampal Temporoammonic-CA1 Synapses. J. Neurosci. doi:10.1523/JNEUROSCI.2588-13.2013

Kang, H.J., Voleti, B., Hajszan, T., Rajkowska, G., Stockmeier, C.A., Licznerski, P., Lepack, A., Majik, M.S., Jeong, L.S., Banasr, M., Son, H., Duman, R.S., 2012. Decreased expression of synapse-related genes and loss of synapses in major depressive disorder. Nat. Med. doi:10.1038/nm.2886

Kendler, K.S., Karkowski, L.M., Prescott, C.A., 1999. Causal Relationship Between Stressful Life Events and the Onset of Major Depression. Psychiatry Interpers. Biol. Process. 156, 837–841. doi:10.1176/ajp.156.6.837

Kessler, R.C., Bromet, E.J., 2013. The epidemiology of depression across cultures. Annu. Rev. Public Health 34, 119–38. doi:10.1146/annurev-publhealth-031912-114409

Kim, D.M., Leem, Y.H., 2016. Chronic stress-induced memory deficits are reversed by regular exercise
via AMPK-mediated BDNF induction. Neuroscience. doi:10.1016/j.neuroscience.2016.03.019

Klein, S.L., Flanagan, K.L., 2016. Sex differences in immune responses. Nat. Rev. Immunol.
doi:10.1038/nri.2016.90

Labonté, B., Engmann, O., Purushothaman, I., Menard, C., Wang, J., Tan, C., Scarpa, J.R., Moy, G.,
Loh, Y.-H.E., Cahill, M., Lorsch, Z.S., Hamilton, P.J., Calipari, E.S., Hodes, G.E., Issler, O.,
Kronman, H., Pfau, M., Obradovic, A.L.J., Dong, Y., Neve, R.L., Russo, S., Kazarskis, A.,
Tamminga, C., Mechawar, N., Turecki, G., Zhang, B., Shen, L., Nestler, E.J., 2017. Sex-specific
transcriptional signatures in human depression. Nat. Med. 23, 1102–1111. doi:10.1038/nm.4386

Lagunas, N., Calmarza-Font, I., Diz-Chaves, Y., Garcia-Segura, L.M., 2010. Long-term ovariectomy
enhances anxiety and depressive-like behaviors in mice submitted to chronic unpredictable stress.
Horm. Behav. 58, 786–791. doi:10.1016/j.yhbeh.2010.07.014

LaPlant, Q., Chakravarty, S., Vialou, V., Mukherjee, S., Koo, J.W., Kalahasti, G., Bradbury, K.R.,
Taylor, S. V., Maze, I., Kumar, A., Graham, A., Birnbaum, S.G., Krishnan, V., Truong, H.T.,
Neve, R.L., Nestler, E.J., Russo, S.J., 2009. Role of Nuclear Factor κB in Ovarian Hormone-
Mediated Stress Hypersensitivity in Female Mice. Biol. Psychiatry 65, 874–880.
doi:10.1016/j.biopsych.2009.01.024

Li, L.H., Wang, Z.C., Yu, J., Zhang, Y.Q., 2014. Ovariectomy results in variable changes in
nociception, mood and depression in adult female rats. PLoS One.
doi:10.1371/journal.pone.0094312

Liu, Y., Chun-Man Ho, R., Mak, A., 2012. Interleukin (IL)-6, tumour necrosis factor alpha (TNF-α)
and soluble interleukin-2 receptors (sIL-2R) are elevated in patients with major depressive
disorder: A meta-analysis and meta-regression. doi:10.1016/j.jad.2011.08.003

Mahmoud, Rand, Wainwright, S.R., Chaiton, J.A., Lieblich, S.E., Galea, L.A.M., 2016. Ovarian
hormones, but not fluoxetine, impart resilience within a chronic unpredictable stress model in
middle-aged female rats. Neuropharmacology 107, 278–293.
doi:10.1016/j.neuropharm.2016.01.033

Mahmoud, R., Wainwright, S.R., Chaiton, J.A., Lieblich, S.E., Galea, L.A.M., 2016. Ovarian
hormones, but not fluoxetine, impart resilience within a chronic unpredictable stress model in
middle-aged female rats. Neuropharmacology 107. doi:10.1016/j.neuropharm.2016.01.033

McEwen, B.S., Nasca, C., Gray, J.D., 2016. Stress Effects on Neuronal Structure: Hippocampus,
Amygdala, and Prefrontal Cortex. Neuropsychopharmacology. doi:10.1038/npp.2015.171

McKinnon, M.C., Yucel, K., Nazarov, A., MacQueen, G.M., 2009. A meta-analysis examining clinical
predictors of hippocampal volume in patients with major depressive disorder. J. Psychiatry
Neurosci. 34, 41–54.

Meziane, H., Ouagazzal, A.M., Aubert, L., Wietrzych, M., Krezel, W., 2007. Estrous cycle effects on
behavior of C57BL/6J and BALB/cByJ female mice: Implications for phenotyping strategies.
Genes, Brain Behav. doi:10.1111/j.1601-183X.2006.00249.x

Miller, A.H., Raison, C.L., 2016. The role of inflammation in depression: From evolutionary
imperative to modern treatment target. Nat. Rev. Immunol. doi:10.1038/nri.2015.5

Miller, A.H., Raison, C.L., 2006. Cytokines, p38 MAP Kinase and the Pathophysiology of Depression.
Neuropsychopharmacology. doi:10.1038/sj.npp.1301032

Nordell, V.L., Scarborough, M.M., Buchanan, A.K., Sohrabji, F., 2003. Differential effects of estrogen
in the injured forebrain of young adult and reproductive senescent animals. Neurobiol. Aging.
doi:10.1016/S0197-4580(02)00193-8

Orlowski, D., Elfving, B., Müller, H.K., Wegener, G., Bjarkam, C.R., 2012. Wistar rats subjected to
chronic restraint stress display increased hippocampal spine density paralleled by increased
expression levels of synaptic scaffolding proteins. Stress. doi:10.3109/10253890.2011.643516

Pacheco, A., Aguayo, F.I., Aliaga, E., Muñoz, M., García-Rojo, G., Olave, F.A., Parra-Fiedler, N.A.,
García-Pérez, A., Tejos-Bravo, M., Rojas, P.S., Parra, C.S., Fiedler, J.L., 2017. Chronic Stress
Triggers Expression of Immediate Early Genes and Differentially Affects the Expression of
AMPA and NMDA Subunits in Dorsal and Ventral Hippocampus of Rats. Front. Mol. Neurosci.
doi:10.3389/fnmol.2017.00244

Paolicelli, R.C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M., Ferreira,
T.A., Guiducci, E., Dumas, L., Ragozzino, D., Gross, C.T., 2011. Synaptic pruning by microglia is
necessary for normal brain development. Science (80-.). doi:10.1126/science.1202529

Rao, J.S., Harry, G.J., Rapoport, S.I., Kim, H.W., 2010. Increased excitotoxicity and
neuroinflammatory markers in postmortem frontal cortex from bipolar disorder patients. Mol.
Psychiatry 15, 384–392. doi:10.1038/mp.2009.47

Salk, R.H., Hyde, J.S., Abramson, L.Y., 2017. Gender differences in depression in representative
national samples: Meta-analyses of diagnoses and symptoms. Psychol. Bull. 143, 783–822.
doi:10.1037/bul0000102

Santarelli, L., Saxe, M., Gross, C., Surget, A., Battaglia, F., Dulawa, S., Weisstaub, N., Lee, J., Duman,
R., Arancio, O., Belzung, C., Hen, R., 2003. Requirement of hippocampal neurogenesis for the

behavioral effects of antidepressants. Science 301, 805–9. doi:10.1126/science.1083328

Schmidt, P.J., Ben Dor, R., Martinez, P.E., Guerrieri, G.M., Harsh, V.L., Thompson, K., Koziol, D.E.,

Nieman, L.K., Rubinow, D.R., 2015. Effects of Estradiol Withdrawal on Mood in Women With

Past Perimenopausal Depression. JAMA Psychiatry 72, 714.

doi:10.1001/jamapsychiatry.2015.0111

Selvamani, Amutha, Sohrabji, F., 2010. Reproductive age modulates the impact of focal ischemia on

the forebrain as well as the effects of estrogen treatment in female rats. Neurobiol. Aging.

doi:10.1016/j.neurobiolaging.2008.08.014

Selvamani, A., Sohrabji, F., 2010. The Neurotoxic Effects of Estrogen on Ischemic Stroke in Older

Female Rats Is Associated with Age-Dependent Loss of Insulin-Like Growth Factor-1. J.

Neurosci. doi:10.1523/jneurosci.0761-10.2010

Setiawan, E., Wilson, A.A., Mizrahi, R., Rusjan, P.M., Miler, L., Rajkowska, G., Suridjan, I., Kennedy,

J.L., Rekkas, P.V., Houle, S., Meyer, J.H., 2015. Role of Translocator Protein Density, a Marker

of Neuroinflammation, in the Brain During Major Depressive Episodes. JAMA Psychiatry 72,

268. doi:10.1001/jamapsychiatry.2014.2427

Shansky, R.M., Hamo, C., Hof, P.R., Lou, W., McEwen, B.S., Morrison, J.H., 2010. Estrogen

promotes stress sensitivity in a prefrontal cortex-amygdala pathway. Cereb. Cortex 20, 2560–

2567. doi:10.1093/cercor/bhq003

Shansky, R.M., Hamo, C., Hof, P.R., McEwen, B.S., Morrison, J.H., 2009. Stress-induced dendritic

remodeling in the prefrontal cortex is circuit specific. Cereb. Cortex 19, 2479–2484.

doi:10.1093/cercor/bhp003

Smith, J.A., Das, A., Butler, J.T., Ray, S.K., Banik, N.L., 2011. Estrogen or estrogen receptor agonist

inhibits lipopolysaccharide induced microglial activation and death. Neurochem. Res.

doi:10.1007/s11064-010-0336-7

Soares, C.N., 2014. Mood disorders in midlife women: Understanding the critical window and its

clinical implications. Menopause. doi:10.1097/GME.0000000000000193

Soares, C.N., 2013. Depression in peri- and postmenopausal women: Prevalence, pathophysiology and

pharmacological management. Drugs and Aging. doi:10.1007/s40266-013-0100-1

Spencer-Segal, J.L., Tsuda, M.C., Mattei, L., Waters, E.M., Romeo, R.D., Milner, T.A., McEwen, B.S.,

Ogawa, S., 2012. Estradiol acts via estrogen receptors alpha and beta on pathways important for

synaptic plasticity in the mouse hippocampal formation. Neuroscience.

doi:10.1016/j.neuroscience.2011.11.035

Spencer, J.L., Waters, E.M., Milner, T.A., Lee, F.S., McEwen, B.S., 2010. BDNF variant Val66Met

interacts with estrous cycle in the control of hippocampal function. Proc. Natl. Acad. Sci.

doi:10.1073/pnas.0915105107

Suuronen, T., Nuutinen, T., Huuskonen, J., Ojala, J., Thornell, A., Salminen, A., 2005. Anti-

inflammatory effect of selective estrogen receptor modulators (SERMs) in microglial cells.

Inflamm. Res. 54, 194–203. doi:10.1007/s00011-005-1343-z

Suzuki, S., Brown, C.M., Wise, P.M., 2009. Neuroprotective effects of estrogens following ischemic

stroke. Front. Neuroendocrinol. doi:10.1016/j.yfrne.2009.04.007

Syed, S.A., Beurel, E., Loewenstein, D.A., Lowell, J.A., Craighead, W.E., Dunlop, B.W., Mayberg,

H.S., Dhabhar, F., Dietrich, W.D., Keane, R.W., de Rivero Vaccari, J.P., Nemeroff, C.B., 2018.

Defective Inflammatory Pathways in Never-Treated Depressed Patients Are Associated with Poor

Treatment Response. Neuron. doi:10.1016/j.neuron.2018.08.001

Tapia-Gonzalez, S., Carrero, P., Pernia, O., Garcia-Segura, L.M., Diz-Chaves, Y., 2008. Selective

oestrogen receptor (ER) modulators reduce microglia reactivity in vivo after peripheral

inflammation: potential role of microglial ERs. J. Endocrinol. doi:10.1677/joe-07-0294

Vegeto, E., Belcredito, S., Etteri, S., Ghisletti, S., Brusadelli, A., Meda, C., Krust, A., Dupont, S.,

Ciana, P., Chambon, P., Maggi, A., 2003. Estrogen receptor-α mediates the brain

antiinflammatory activity of estradiol. Proc. Natl. Acad. Sci. doi:10.1073/pnas.1531957100

Vegeto, E., Benedusi, V., Maggi, A., 2008. Estrogen anti-inflammatory activity in brain: A therapeutic

opportunity for menopause and neurodegenerative diseases. Front. Neuroendocrinol.

doi:10.1016/j.yfrne.2008.04.001

Vegeto, E., Bonincontro, C., Pollio, G., Sala, a, Viappiani, S., Nardi, F., Brusadelli, a, Viviani, B.,

Ciana, P., Maggi, a, 2001. Estrogen prevents the lipopolysaccharide-induced inflammatory

response in microglia. J. Neurosci. 21, 1809–1818. doi:21/6/1809 [pii]

Wainwright, S.R., Barha, C.K., Hamson, D.K., Epp, J.R., Chow, C., Lieblich, S.E., Rutishauser, U.,

Galea, L.A.M., 2016. Enzymatic depletion of the polysialic acid moiety associated with the neural

cell adhesion molecule inhibits antidepressant efficacy. Neuropsychopharmacology 41, 1670–

1680. doi:10.1038/npp.2015.337

Wang, A.K., Miller, B.J., 2018. Meta-analysis of Cerebrospinal Fluid Cytokine and Tryptophan

Catabolite Alterations in Psychiatric Patients: Comparisons between Schizophrenia, Bipolar

Disorder, and Depression. Schizophr. Bull. doi:10.1093/schbul/sbx035

Wang, J.Q., Mao, L., 2019. The ERK Pathway: Molecular Mechanisms and Treatment of Depression.

Mol. Neurobiol. doi:10.1007/s12035-019-1524-3

Waters, E.M., Mitterling, K., Spencer, J.L., Mazid, S., McEwen, B.S., Milner, T.A., 2009. Estrogen

receptor alpha and beta specific agonists regulate expression of synaptic proteins in rat

hippocampus. Brain Res. doi:10.1016/j.brainres.2009.06.090

Waters, E.M., Thompson, L.I., Patel, P., Gonzales, A.D., Ye, H., Filardo, E.J., Clegg, D.J., Gorecka, J.,

Akama, K.T., McEwen, B.S., Milner, T.A., 2015. G-Protein-Coupled Estrogen Receptor 1 Is

Anatomically Positioned to Modulate Synaptic Plasticity in the Mouse Hippocampus. J. Neurosci.

doi:10.1523/jneurosci.1298-14.2015

Wilson, L., Pandeya, N., Byles, J., Mishra, G., 2018. Hysterectomy and incidence of depressive

symptoms in midlife women: The Australian Longitudinal Study on Women's Health. Epidemiol.

Psychiatr. Sci. doi:10.1017/S2045796016001220

Wohleb, E.S., Patterson, J.M., Sharma, V., Quan, N., Godbout, J.P., Sheridan, J.F., 2014. Knockdown

of Interleukin-1 Receptor Type-1 on Endothelial Cells Attenuated Stress-Induced

Neuroinflammation and Prevented Anxiety-Like Behavior. J. Neurosci.

doi:10.1523/jneurosci.3723-13.2014

Wood, S.K., Wood, C.S., Lombard, C.M., Lee, C.S., Zhang, X.Y., Finnell, J.E., Valentino, R.J., 2015.

Inflammatory factors mediate vulnerability to a social stress-induced depressive-like phenotype in

passive Coping Rats. Biol. Psychiatry. doi:10.1016/j.biopsych.2014.10.026

Woolley, C.S., Gould, E., Frankfurt, M., McEwen, B.S., 1990. Naturally occurring fluctuation in

dendritic spine density on adult hippocampal pyramidal neurons. J. Neurosci.

Workman, J.L., Brummelte, S., Galea, L.A.M., 2013. Postpartum Corticosterone Administration

Reduces Dendritic Complexity and Increases the Density of Mushroom Spines of Hippocampal

CA3 Arbours in Dams. J. Neuroendocrinol. 25, 119–130. doi:10.1111/j.1365-2826.2012.02380.x

Zhang, D., Zhang, J., Bian, C., Deng, Q., 2010. Postnatal and ovariectomic regulation of postsynaptic

density protein-95 in the hippocampus of female Sprague-Dawley rats. Synapse.

doi:10.1002/syn.20805