Crystal structure of *Sulfolobus solfataricus* topoisomerase III reveals a novel carboxyl-terminal zinc finger domain essential for decatenation activity

Hanqian Wang[1,+], Junhua Zhang[2,+], Zengqiang Gao[3], Xin Zheng[1], Keli Zhu[1,4], Zhenfeng Zhang[2], Zhiyong Zhang[5], Yuhui Dong[1,3], Li Huang[2,6,*] and Yong Gong[1,*]

[1] Multi-discipline Center, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, 100049, China

[2] State Key Laboratory of Microbial Resources, Institute of Microbiology, Chinese Academy of Sciences, Beijing, 100101, China

[3] Beijing Synchrotron Radiation Facility, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, 100049, China

[4] Institute of Physical Science and Information Technology, Anhui University, Anhui, 230601, China

[5] CAS Key Laboratory for Biomedical Effects of Nanomaterials and Nanosafety, CAS Key Laboratory of Nuclear Radiation and Nuclear Energy Technology, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, 100049, China

[6] College of Life Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China

* Correspondence to: Yong Gong, Multidiscipline Research Center, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China; Tel: +86 10 8823 5257; Fax: +86 1088 2352 57; Email: yonggong@ihep.ac.cn

to Li Huang, State Key Laboratory of Microbial Resources, Institute of Microbiology, Chinese Academy of Sciences, Beijing, 100101, Tel: +86 10 6480 7606; Fax: +86 10 6480 7429; Email: huangl@im.ac.cn

[+] These authors contributed equally to the paper as the first authors.

Running title: *Crystal structure of an archaeal DNA topoisomerase III*

ABSTRACT

DNA topoisomerases are essential enzymes for a variety of cellular processes

involved in DNA transactions. Mechanistic insights into type IA DNA

topoisomerases have come principally from studies on bacterial and

eukaryotic enzymes. A structural understanding of type IA topoisomerases in

Archaea is lacking. Here, we present a 2.1-angstrom crystal structure of

full-length *Sulfolobus solfataricus* topoisomerase III (*Sso* topo III), an archaeal

member of type IA topoisomerases. The structure shows that *Sso* topo III

adopts a characteristic torus-like architecture consisting of a four-domain core

region and a novel carboxyl-terminal zinc finger domain (domain V).

Structure-based mutation analyses reveal that a novel zinc-binding motif in

domain V is essential for the DNA decatenation activity of *Sso* topo III. Our

data indicate that *Sso* topo III represents a subclass of Type IA

topoisomerases capable of resolving DNA catenates using a domain

V-dependent mechanism.

58 **IMPORTANCE**

59 Type IA topoisomerases are omnipresent in all cellular life forms and serve

pivotal roles in cellular processes involved in DNA transactions. While

considerable insights have been gained into Type IA topoisomerases from

bacteria and eukaryotes, a structural understanding of type IA topoisomerases

in Archaea remains elusive. we first determined the crystal structure of

full-length *Sulfolobus solfataricus* topoisomerase III (*Sso* topo III), an archaeal

member of type IA topoisomerases. Our structure provides the first molecular

view of this archaeal topoisomerase, which removes negative supercoils and

decatenates DNA catenane. Our findings manifest that *Sso* topo III may serve

as an alternative prototype of type IA topoisomerases, whose decatenation

mechanism differs from that of known bacterial and eukaryotic topoisomerases

III such as *Escherichia coli* topoisomerase III (EcTOP3).

KEYWORDS

Archaea, DNA topoisomerase, crystal structure, zinc finger, decatenation

activity, enzyme mutation

INTRODUCTION

The overall topological state of DNA in cells is altered by DNA

topoisomerases. In all three domains of life, conserved DNA topoisomerases

are essential for a wide variety of cellular processes involved in DNA

transactions (1-4). There are two types of topoisomerases (1). Type I

topoisomerases cut only one strand of DNA, while type II topoisomerases

cleave both strands of the DNA duplex. Type I enzymes can be further

classified into three different families, types IA, IB and IC, on the basis of

transiently formed covalent intermediates formed (5'- or 3'- covalent

intermediate) (5). *Escherichia coli* topoisomerase I (EcTOP1) and

topoisomerase III (EcTOP3) are the best-characterized bacterial type IA

enzymes to date.

A conserved toroidal fold formed by four domains (I-IV) has been found in

the structures of type IA enzymes including EcTOP1 (6), EcTOP3 (7),

Thermotoga maritima topoisomerase I (TmTOP1) (8), *Mycobacterium*

tuberculosis topoisomerase I (MtTOP1) (9), *Mycobacterium smegmatis*

topoisomerase I (MsTOP1) (10), human topoisomerase IIIα (HsTop3α) (11),

human topoisomerase IIIβ (HsTop3β) (12) and a reverse gyrase (13) from

Archaeoglobus fulgidus, a hyperthermophilic archaeon. Structural studies

(14-16) as well as biochemical (17-20) and single-molecule (21-24)

experimental data have provided abundant evidences in support of an

enzyme-bridged strand passage model for type IA enzymes. According to this

model, a type IA topoisomerase nicks one DNA single-strand, and remains

attached to the two ends of the nick. That is, the enzyme is not only covalently

attached to one end of the nick, but also noncovalently bound to the other end

to generate a bridge through which another single-strand or the DNA duplex is

passed. The following step is the resealing of the nick and the release of the

enzyme. In this process, the OH group of the aromatic ring of the Tyr residue in

the enzyme nucleophilically attacks the scissile phosphodiester bond,

producing a transient 5' phosphotyrosine covalent bond.

Members of the type IA subfamily are further divided into two

mechanistically distinct subclasses based on their biochemical specificity for

substrates (2). EcTOP1 and EcTOP3 are the prototypes of these two

subclasses of type IA enzymes. EcTOP3 catalyzes the DNA catenates more

effectively than it relaxes negatively supercoiled DNA, whereas the converse is

true for EcTOP1 (23, 25-28). EcTOP3, but not EcTOP1, contains a unique

loop located close to the central hole (29). This positively charged loop, known

as the decatenation loop, is essential for the decatenation activity of EcTOP3

(24, 29).

Previous studies have characterized topoisomerase III (hereafter, *Sso* topo

III) from *Sulfolobus solfataricus*, a hyperthermophilic archaeon, as a member

of the type IA topoisomerase family (30, 31). A phylogenetic analysis of *Sso*

topo III, and the observation that *Sso* topo III inefficiently catalyzes the

relaxation of negatively supercoiled DNA, led to the suggestion that *Sso* topo

III is more similar to EcTOP3 than to EcTOP1 (30). More recently, Bizard et al.

provided convincing biochemical data in support of notion that *Sso* topo III is a

genuine decatenase that can efficiently unlink covalently closed catenanes

alone at temperatures up to 90℃ (32). However, little is known about the

structure of type IA topoisomerases (except for reverse gyrase) from the

Archaea. The structural basis for the decatenase activity of *Sso* topo III is

unclear.

Here, we have determined the crystal structure of full-length *Sso* topo III.

Sso topo III adopts a classical elongated toroidal-like shape with four domains

(I-IV) forming the protein core and a novel carboxyl-terminal zinc finger domain

(domain V) tethered to the core against domain II. Furthermore, domain V is

conserved among DNA topoisomerases III from the TACK superphylum of the

Archaea. In addition, structure-based mutational analyses show that, whereas

the zinc-binding motif of domain V was required for efficient DNA relaxation by

Sso topo III, its depletion nearly abolished the DNA decatenation activity of the

enzyme. These findings indicate that *Sso* topo III may represent as a novel

type of topoisomerases III which employ a decatenation mechanism distinctly

different from that of known bacterial and eukaryotic equivalent.

148
149

RESULTS

Overall structure of intact *Sso* topo III To obtain crystals of *Sso*

topo III, we overproduced and purified sufficient quantities of the mutant

enzyme (Y318F) in which the catalytic residue Tyr318 was replaced with Phe

(Fig. 1 A). The structure of the full-length enzyme was determined at 2.1 Å

resolution using molecular replacement (Table 1). The full-length protein

formed an asymmetric unit comprising four *Sso* topo III molecules. The four

molecules had essentially the same architecture. For convenience, one

molecule was chosen for the following structural analyses.

Our analysis showed that the overall structure of *Sso* topo III comprises four

domains (I–IV) (residues 1–593), forming the conserved core, and a novel

carboxyl-terminal domain V (residues 594–668) (Fig. 1A, 1B and S1). The

enzyme adopts a canonical toroidal-like elongated configuration with a large

hole (Fig. 1B and 1C), which is shared by all known type 1A topoisomerases

(33). The enzyme-bridged strand passage model proposes that the hole

accommodates single or double-stranded DNA (34).

Domain I (residues 1–219) exhibits a topoisomerase–primase fold (2), which

is found in type IA, IIA, and IIB topoisomerases, as well as in bacterial

primases. Within domain I, a disulphide bridge crosslinks Cys4 of a loop

(residues 1–10) and Cys34 of an α/β fold consisting of six parallel β strands

sandwiched with four α helices (Fig. S1). This connection results in tight

packing of domain I and stabilizes its architecture. In Domain II (residues

220–277, 408–474), the hinge region (residues 218–224, 471–480) is

composed of two loops linking domains I and II, and II and IV, respectively (Fig.

1B). The hinge is believed to serve as the principal pivot point for opening and

closing of the DNA gate in the strand passage reaction (6, 11, 35).

A long loop (residues 594–616) leads from helix α18 of domain IV to

domain V (residues 594–668), thus linking the core of the enzyme to domain V

(Fig. 1B and S1). Domain V attaches to the top of the topoisomerase (domain

II) against the hinge region (Fig. 1B and 1C). Three antiparallel α-helices,

including helix α19 (residues 616–636), helix α20 (residues 640–649), and

helix α21 (residues 655–666), are coiled around each other to constitute the

bulk of domain V (Fig. 1B). Interestingly, four residues (Cys602, Cys605,

Cys615 and His618) coordinate with a zinc atom to generate a novel

zinc-binding motif (Fig. 1D and S1). Among them, the three Cys residues come

from the long loop connecting domains IV/V and the His residue is from helix

α19. The presence of the znic-binding site was confirmed by an analysis of the

anomalous difference Fourier map (Fig. 1D). The recently characterized

mesophilic archaeal topoisomerase, *Methanosarcina acetivorans* topo IIIα

(*Mac* topo IIIα), also has one znic-binding site (36). However, the zinc-binding

motif of *Sso* topo III is the Cys3His1 type, whereas that of *Mac* topo IIIα is

tetraCys type. For *Sso* topo III, the zinc finger motif may contribute to the

stabilization of domain V. As revealed by sequence alignment, the

carboxyl-terminal zinc finger motif of the Cys3His1 type is highly conserved

among topoisomerases from the Crenarchaeota (Fig. S2A), suggesting that

this motif plays an important role. Intriguingly, small proteins homologous to

domain V are also found in Thaumarchaeota and Bathyarchaeota, both of

which are phyla within the TACK superphylum where Crenarchaeota also

belong, as well as Euryarchaeota and Thorarchaeota (Fig. S3). Homologues of

Sso topo III are present in Thaumarchaeota, Bathyarchaeota, Euryarchaeota

and Thorarchaeota, but these enzymes almost lack the carboxyl-terminal

domain similar to domain V in *Sso* topo III. It would be of interest to determine

if the domain V-like small proteins physically and functionally interact with the

type IA topoisomerases in these organisms. Notably, no structural elements

similar to domain V of *Sso* topo III is found in the Protein Data Bank, as

detected by DALI (37). These results suggest that the carboxyl-terminal fold of

Sso topo III, also found in several archaeal lineages, is unique.

Interface of domains II and V A distinguishing feature of the overall

structure of *Sso* topo III compared with that of other known available type IA

topoisomerases is the presence of three helices in domain V located close to

the hinge region of domain II (Fig. 1B). Two flexible loops, each about five

residues long, link these three helices. This arrangement causes the three

helices to pack against each other so that their movement is highly

coordinated, but they maintain a degree of flexibility. Additionally, within the

long linker loop (about 20 residues) connecting domains IV and V, there are

three successive turns, which maintain free space between the two domains.

The novel zinc-binding motif, located on the carboxyl-end of the long loop,

contributes to the stabilization of the loop (Fig. 1D). By analogy to the other

solved type IA topoisomerases (6), the interfaces between domain III and

domains IV and I in *Sso* topo III are conserved and far-ranging. However, the

interface between domains II and V within the enzyme is unique.

The longest one of the three helices in domain V, helix α19, acts as the

major binding platform for the interface between domains II and V. The

interface is mostly, but not exclusively, hydrophobic (Fig. 1E). Extensive

hydrophobic interactions mainly occur between the side chains of Leu219,

Phe220, Ile221, Pro222, Leu223, Pro224 and Phe226 in the hinge region,

which connects helix α6 of domain I with strand β9 of domain II, and the side

chains of Ala621, Leu625, Ala628, Leu631, Trp632 in helix α19 and Tyr656,

Val657 in helix α21 of domain V (Fig. 1E). Moreover, the main chain amides of

Gly654 and Lys655 in domain V form hydrophilic interactions with the

backbone carbonyl groups of Phe220 and Asn218 in the hinge region,

respectively, while the amide of Tyr656 makes hydrophilic interaction with the

backbone carbonyl groups of Asn218 (Fig. 1F). In addition, the NH group on

an indole group of Trp632 makes a hydrogen bond with the backbone carbonyl

group of Pro222. These interactions further stabilize the attachment of domain

V to domain II. Most importantly, the main chain carbonyl group and side chain

amide of Arg635 in the carboxyl-end of helix α19 in domain V form hydrogen

bonds with the main chain amide of Ile250 at the amino-end of helix α7 and the

carbonyl group of Pro224 at the amino-end of strand β9 in domain II,

respectively (Fig. 1F). Domain V and the main part of domain II are connected

by a single interaction, that is, double hydrogen bonds involving Arg635. This

may allow for independent and coordinated movements of domain V relative to

domain II at different stages during catalysis. Noticeably, Glu608 from domain

V forms salt bridge with Lys592 from domain IV on the other end of the

interface. Alignment of the sequences of type IA topoisomerases from the

Crenarchaeota shows that most of the above-mentioned residues contributing

to the interaction between domains II and V are highly conserved (Fig. S2B).

This implies that the interaction between domains II and V may be essential for

the functional role of domain V of the archaeal topoisomerases.

Comparison of the interactions around the hinge region

among type IA topoisomerases The studies on EcTOP3 (7, 14, 29)

indicated that the decatenation loop on domain IV is critical for efficient DNA

decatenation (Fig. 2A). A recent report on *Sso* topo III provides strong

biochemical evidence that the enzyme is a true decatenase (32). Surprisingly,

the structural equivalent of the decatenation loop is absent from *Sso* topo III

(Fig. 2A), suggesting that DNA decatenation by *Sso* topo III may differ

mechanistically from that by EcTOP3.

It is believed that the highly conserved hinge region joining domains II and IV

appears to be important for strand passage during catalysis (6). To understand

the decatenation mechanism of *Sso* topo III, we performed a structural

analysis of the potential interactions within the available type IA

topoisomerases between the hinge region and the carboxyl-terminal domain or

an additional subunit of the topoisomerase complex. No direct interactions

exist between the hinge regions and the C-terminus in TmTOP1 (8), MtTOP1

(9) and MsTOP1 (10). However, there are direct interactions between the

hinge region and parts of the carboxyl-terminal domain in several

topoisomerases or the other subunit of a topoisomerase complex (Fig. 2B).

The interactions may influence the opening state of their gates. The human

BLM helicase, topoisomerase IIIα (HsTop3α) and RMI1 proteins form a

minimal protein complex, which catalyses the dissolution of double-Holliday

junctions (dHJ) to produce non-crossover products (38). A structural analysis

of the Top3a–RMI1 complex showed that the hinge region makes intimate

contacts with the decatenation loop of the RMI1 subunit by hydrophobic

interaction, suggesting that this contact contributes to the modulation of the

opening-closing state of the gate in Top3α (11) (Fig. 2B). Similar to the

Top3a–RMI1 complex, Top3β forms a complex with the TDRD3 subunit. A

structural analysis of the complex indicated that the hinge region of Top3β is a

little bit apart from the insertion loop of TDRD3 (12) (Fig. 2B). In the case of

EcTOP1, the interaction between a unique α helix in domain VI of the

carboxyl-terminal domain and the hinge region mediates the major direct

contact between the carboxyl-terminal domain and the catalytic core (39) (Fig.

2B). The conformational changes of the carboxyl-terminal domain are

potentially transferred to the hinge region through the unique helix in domain

VI at some points in the catalytic cycle, thereby regulating the opening-closing

state of the gate. In Sso topo III, major contacts between the hinge region and

the longer helix α19 of domain V occur through stable and extensive

hydrophobic interactions (Fig. 2C), and several hydrogen bonds, located at the

two ends of the interface between domains II and V, are involved in the

interaction between domain V and the bulk of domain II and domain IV,

respectively (Fig. 1F). This network of interactions may allow domain V to

adhere to the hinge region mainly by the α19 helix during the strand passage

reaction, and possibly make the bulk of domain II and domain IV to move

relative to the hinge region at some points during the catalysis.

Deletion of the zinc-binding motif reduces the relaxation

activity of *Sso* topo III To explore the role of the carboxyl-terminal domain,

the zinc-binding motif in particular, in DNA relaxation by *Sso* topo III, a

carboxyl-terminally truncated mutant (597ΔC), which lacked whole domain V,

and a quadruple mutant (CCCH/4A), in which the four residues (Cys602,

Cys605, Cys615 and His618) coordinating with the zinc atom were all replaced

with Ala, were generated by mutagenesis (Table S1). In the CCCH/4A, the

mutation of the zinc-binding motif (3Cys1His type) resulted in the elimination of

the zinc atom (Fig. 1D). These two mutant proteins were overproduced in *E.*

coli and purified to homogeneity. The two mutant proteins showed the same

optimal temperature as the wild-type enzyme (75℃) in DNA relaxation (Fig.

S4). 597ΔC possessed about 50% of the wild-type activity (Fig. 3, compare top

and middle panel), whereas CCCH/4A retained about 15% of the wild-type

activity (Fig. 3, compare top and bottom panel). These results indicated that

the zinc-binding motif is necessary for efficient relaxation by *Sso* topo III.

To compare DNA relaxation reactions catalysed by wild-type and mutants

of *Sso* topo III, a time-course assay was performed (Fig. 4). The wild-type

protein displayed a profile of topoisomers indicative of a distribute mechanism

of relaxation (Fig. 4, top panel), as described recently (32). However, both

597ΔC and CCCH/4A exhibit a relaxation profile differently from the wild-type

Sso topo III (Fig. 4, compare top, middle, and bottom panel). The DNA

relaxation reaction catalysed by 597ΔC appeared to more distributive than that

by the wild-type protein. Interestingly, the relaxation reaction becomes more

distributive, when the unique zinc-binding motif was eliminated, than that as

the carboxyl-terminal domain of *Sso* topo III was wholly deleted (Fig. 4).

Together, these results indicated that domain V, especially the zinc-binding

motif, facilitates the relaxation of negatively supercoiled DNA by *Sso* topo III.

To probe the potential roles of domain V in *Sso*Topo III, we also

determined ssDNA binding by the domain V-deletion mutant protein 594ΔC

(residues 1-594). The mutant protein showed a ~ 5-fold lower affinity than

wild-type *Sso*Topo III for a 32-nt ssDNA containing the preferred cleavage site

(designated C32) than the wild-type protein (Fig. S5). Therefore, it appeared

that domain V contributes significantly to template binding by *Sso*Topo III.

The novel zinc-binding motif is essential for DNA

decatenation by *Sso* topo III To explore the role of the novel domain V

in *Sso* topo III-catalysed decatenation, wild-type and mutants of *Sso* topo III

were subjected to a decatenation assay using kinetoplast DNA (kDNA) (40) as

the substrate. Wild-type *Sso* topo III was optimally active in kDNA

decatenation at 90℃, in consistent with the observation made previously (32)

(Fig. S6A, B, C). The fast-migrating bands, corresponding to closed minicircles,

were present when the reactions were performed at very high temperatures

(90 °C) (Fig. 5A). These closed minicircles were released from the kDNA

network by *Sso* topo III, because the enzyme apparently was unable to relax

supercoiled DNA at 90 °C and above, but was still able to decatenate DNA

catenanes (32). Therefore, the presence of closed minicircles at very high

temperature was considered as a measure of the decatenation activity of *Sso*

topo III. The bands corresponding to closed minicircles were undetectable in

reactions containing 597ΔC (Fig. 5B), suggesting that the mutant protein had

no detectable decatenation activity at very high temperature. Similarly,

CCCH/4A was nearly inactive in decatenation (Fig. 5C). Taken together, we

concluded that domain V, the zinc-binding motif in particular, is essential for

the decatenation activity of *Sso* topo III.

DISCUSSION

Type IA topoisoerases are found in all three domains of life. Much of current

knowledge about the enzymes has come from structural and biochemical

studies on the ones from bacterial and eukaryotic origins (41). A structural

understanding of type IA topoisomerases (except for reverse gyrase) in the

Archaea remains elusive. In this work, we present the crystal structure of *Sso*

topo III, an archaeal type IA topoisomerase, at 2.1 Å resolution. The structure

provides the first molecular view of this archaeal topoisomerase, which

removes negative supercoils and decatenates DNA catenane (30, 32, 42). *Sso*

topo III is toroidal in shape with a four-domain protein core closely resembling

that found in other members of the topoisomerase IA family (6). A striking

feature of *Sso* topo III is the presence at its carboxyl-terminus of a novel zinc

finger-containing part that attaches to the hinge region of domain II. Intriguingly,

the enzyme lacks the decatenation loop found in some of the known type IA

topoisomerases capable of DNA decatenation. As revealed by mutational

analysis, the zinc-binding motif is critical for effective decatenation of *Sso* topo

III. Therefore, *Sso* topo III represents a novel type of topoisomerases which

employ a DNA decatenation mechanism different from those of the known

bacterial and eukaryotic topoisomerases III.

The carboxyl-terminal domain of a type IA topoisomerase varies widely in

composition, length and shape (41, 43), as well as in orientations relative to

the hinge region of the enzyme. Despite wide variations, the carboxyl-terminal

domain exhibits ssDNA-binding activity, and is responsible for the high

DNA-binding affinity of the enzyme. In the case of EcTOP1, domain IV extends

into the carboxyl-terminal domain consisting of three zinc ribbon domains

containing the tetracysteine motif and two zinc ribbon-like domains (39).

Among the five domains, four domains bind to ssDNA with primarily π-stacking

and cation-π interactions. Similarly, TmTOP1 has a zinc ribbon motif, but

without a bound zinc ion, at its C-terminus (8). The deletion of the

carboxyl-terminal domain resulted in a significant loss of the relaxation activity

of TmTOP1, possibly due to the reduction in DNA-binding by the mutant

protein (44, 45). By comparison, the truncation of the carboxyl-terminal domain

almost abolished the relaxation activity of EcTOP3 (46), whose C-terminus

displayed a partially disordered structure (7, 14). Unlike EcTOP1, MtTOP1 has

four tandem novel folds, but not the zinc-binding motif, at the C-terminus (9).

Similar to EcTOP3, the partial truncation of the carboxyl-terminal domain

abolished the relaxation activity of MtTOP1. It is generally believed that the

carboxyl-terminal domains of these two topoisomerases have ssDNA-binding

activity, although they lack a zinc-binding motif. Our structure of *Sso* topo III

provides the first molecular view of the carboxyl-terminal domain of an

archaeal type IA topoisomerase (Fig. 1B). The carboxyl-terminal domain of

Sso topo III is unique, and harbours a zinc-binding motif (3Cys1His type) (Fig.

1D) that differs from the tetracysteine zinc-binding motif conserved in

prokaryotic type IA topoisomerases such as EcTOP1 (39). A structural

analysis of the EcTOP1-DNA complex demonstrated that a zinc ribbon domain

or zinc ribbon-like domain, mainly consisting of β-sheets, is capable of binding

ssDNA, implying that these domains may not bind double-stranded DNA. In

some enzymes involved in DNA transactions, the zinc-finger domains, which

contain α-helixes, bind to double-stranded DNA (47). It will be interesting to

determine whether the carboxyl-terminal fold of *Sso* topo III can bind

double-stranded DNA. In *Sso* topo III, the zinc-binding motif is required for

efficient relaxation activity (Fig. 3 and 4). By comparison, TmTOP1 (44)

remained fully active in DNA relaxation when the zinc ribbon motif was deleted

while EcTOP1 (48) was inactive when the zinc ion was removed.

The first solved structure of a type IA topoisomerase indicates that domains

II and III may move away from domains I and IV in the strand passage reaction

(6). Further evidence for this movement has been provided by biochemical (19,

21), structural data (14, 15, 35) and single-molecule experiments (22-24).   A

structural comparison between EcTOP3 and EcTOP1 indicates that there are

two unique insertions in EcTOP3 in the vicinity of its central hole (29) (Fig. 2A).

One insertion (the decatenation loop) is rich in basic amino acids within

domain IV of EcTOP3, and the other (the acidic loop) is abundant in acidic

amino acids and is located within domain II (2) (Fig. 2A). Decatenation by the

enzyme is driven primarily by the interaction between the decatenation loop

and the acidic loop, both of which line the central hole of the topoisomerase,

during its gate dynamics. Compared with the interaction in EcTOP1, that in

EcTOP3 presumably stabilizes and holds the open state for longer, thus

allowing sufficient time for EcTOP3 to catch duplex DNA and pass it through

the break in a single strand into the gate (24). In the eukaryotic Top3a-RMI

complex, RMI1 coordinates the gate dynamics of Top3a through its unique

decatenation loop, which has a functionally equivalent role to that of the

decatenation loop in EcTOP3 (11). Hence, a unified loop-mediated

decatenation mechanism for prokaryotic and eukaryotic topoisomerases III

has been proposed (11).

Our results demonstrated that the zinc-binding motif is essential for the DNA

decatenation activity of *Sso* topo III (Fig. 5C). We provided the following

evidence in support of the notion that the decatenation mechanism of *Sso* topo

III is distinct from the current loop-mediated decatenation mechanism

proposed for type IA topoisomerase: (i) The decatenation domain (domain V)

of *Sso* topo III is a novel zinc finger fold (Fig. 1D), whereas the decatenation

loop is present in domain IV of EcTOP3 (29) (Fig. 2A). (ii) The decatenation

domain (domain V) of *Sso* topo III is on the outer edge of the topoisomerase

gate and attaches to domain II through the hinge region by extensive

hydrophobic interactions (Fig. 1D and 2C), whereas the decatenation loop in

EcTOP3 lines the DNA gate and is away from the hinge region of the catalytic

core (24) (Fig. 2A). (iii) The 594ΔC mutant lacking domain V showed an

approximately 5-fold reduction in ssDNA-binding compared with wild-type *Sso*

topo III (Fig. S5). In contrast, the deletion of the decatenation loop from

EcTOP3 did not reduce its ssDNA-binding activity (29). Obviously, further

biochemical studies are needed to understand the molecular details of the

unlinking of covalently closed catenanes catalysed by *Sso* topo III.

MATERIALS AND METHODS

438   **Plasmid and DNA constructs** Using the standard PCR cloning

439   strategy, *Sso* topo III gene from *S. solfataricus* genome was cloned into the

440   *Nde*I and *Xho*I sites of the modified version of vector pET28a (Novagene),

441   which harbored an N-terminal poly-histidine tag followed by a tobacco etch

442   virus (TEV) protease cleavage site (ENLYFQ/G) (where '/' donates the

443   cleavage site). However, the growth of *E.coli* cells harboring wild-type *Sso*

444   topo III gene were severely inhibited, possibly due to its toxicity. So, *Sso* topo

445   III gene was mutated to the *Sso* topo III Y318F by replacing residue Tyr318

446   with Phe using the QuickChange^{TM} Site-Directed Mutagenesis Kit (Stratagene)

447   according to the instructions of manufacturer. Site-directed mutagenesis of the

448   other *Sso* Topo III mutants used in this study was carried out according to

449   modified QuickChange^{TM} protocol (49). All the plasmid constructs and mutants

450   were confirmed by DNA double strands sequencing. Primers for *Sso* Topo III

451   site-directed mutagenesis are listed in Table S1.

452   **Protein Expression and Purification** The recombination *Sso*

453   topo III Y318F was overexpressed as a fusion protein with an N-terminal

454   poly-histidine tag in *E. coli* strain Rosetta2 (DE3) pLysS (Novagen). For each

455   purification procedure, 10-l *E. coli* cultures harbouring fusion proteins were

456   grown in the shaker at 37℃ at 185 r.p.m. When their density reached an

457   OD$_{600nm}$ of 1.2, sodium citrate was added into cultures to a final concentration

458   of 100 mM and temperature of the cultures dropped to 25℃, followed by the

459   addition of isopropylb-D-thiogalactoside to a final concentration of 0.5 mM.

After growth for further 6 h at 25℃, the cultures were harvested, and the

resulting cell sediments were suspended in a solubilization buffer containing

20 mM Tris-HCl (pH 8.0), 2 mM potassium citrate, 2 mM $MgCl_2$, 10 mM

imidazole and 500 mM NaCl supplemented with DNAase I (55 µg/ml,

Amersco), followed by disruption with a French press at 12,000 p.s.i. for three

cycles. To remove cell debris, cell lysates were centrifuged for 40 min at

16,000 r.p.m. The resultant supernatant was collected and loaded onto an

equilibrated Ni^{2+} Chelating Sepharose™ Fast Flow column (GE Healthcare).

Purification procedure with the nickel column were performed as described

before (50), except the insertion of an additional wash with high-salt of 1 M

NaCl. After several wash, *Sso* topo III Y318F protein was eluted with a buffer

containing 20 mM Tris-HCl (pH 8.0), 2 mM potassium citrate, and 300 mM

NaCl supplemented with 300 mM imidazole. The eluate was digested using

TEV protease to remove the poly-histidine tag, and dialyzed against buffer I

(20 mM Tris-HCl, pH 7.5, 10 mM potassium citrate, 150 mM NaCl) overnight at

4°C. The resultant eluate was filtered through a 0.45 µm filtration membrane

and then loaded onto a 5 ml SOURCE 15S column (GE Healthcare)

equilibrated with buffer I. The column was eluted with an 80 ml linear gradient

from 150 mM NaCl to 500 mM NaCl, both in buffer I. The fractions, eluted at

0.45-0.50 M NaCl, were pooled and concentrated to 4 mg ml^{-1} in a buffer

containing 20 mM Tris-HCl (pH7.5), 150 mM NaCl, 10 mM potassium citrate,

and 2 mM DTT for later usage. Purified protein was analysed by SDS-PAGE

for purity and quantified with Bradford assay using BSA as standards.

Expression and purification of the other *Sso* Topo III mutants (597ΔC and

CCCH/4A) were carried out according to the procedure described above, with

the exception of ion-exchange column chromatography in the final purification

step. The resultant nickel column eluate was loaded onto a 5 ml SOURCE 15Q

column (GE Healthcare), instead of 15S column, equilibrated with buffer I. The

column was eluted with an 80 ml linear gradient from 0.150 mM NaCl to 1.0 M

NaCl, both in buffer I. The fractions, eluted at 0.55-0.65 M NaCl, were pooled.

Crystallization *Sso* topo III Y318F was crystallized by the sitting-drop

vapor diffusion method, with 0.7µl drops of protein or protein-ssDNA complex

mixed with an equal volume of reservoir solution. Initial crystallization trials

screen clustered-needle-shaped crystals in reservoir solution containing 8%

(v/v) Tacsimate (pH 4.4), and 20% (w/v) Polyethyleneglycerol 3,350. Further

optimization of the crystallization condition led to the appearance of

rod-shaped crystals in the well in which 0.7µl protein solution was mixed with

an equal volume of reservoir solution consisting of 8% (v/v) Tacsimate (pH 4.4),

and 13.75% (w/v) Polyethyleneglycerol 3,350 at 12°C. The diffraction-level

crystals, by the microseeding method, appeared one week later and reached a

maximum size within 2 months. These crystals in the mother liquor were

exposed to air for 10 min, and transferred into a drop of paraffin and NVH mix

oil before being crycooled by plunging rapidly in liquid nitrogen.

Crystallographic Data Collection, Structure determination

and Refinement X-ray diffraction data were collected at the BL17U

beamline of the Shanghai Synchrotron Radiation Facility, and indexed,

integrated and scaled using the HKL2000 package (51). The initial phases

were determined by molecular replacement using the program PHASER (52),

with the structure of TmTOP1 (PDB ID: 2GAJ) as a searching model. The

structure refinement was carried out with phenix.refine (53) and Refmac5 (54).

Model building was carried out by iterative rounds of manual building with Coot

(55). MolProbity (56) was used to validate the structure. Data collection and

refinement statistics are listed in Table 1.

Assays of decatenation and relaxation activities DNA

relaxation assay was performed as described previously (30), except that the

final concentration of $MgCl_2$ is 20 mM, instead of 10 mM, in the standard

reaction mixture. Decatenation of kinetoplast DNA (kDNA) was carried out by a

similar protocol except that 400 ng kDNA (www.topogen.com) was used as

substrate in each standard reaction mixture. Reaction mixes were incubated at

90 °C for 30 min unless specified temperature otherwise, and the

enzyme-catalysed products were examined by electrophoresis in the indicated

agarose gels.

Data availability Atomic coordinates and structure factors for the

crystal structure of *Sso* topo (apo) have been deposited in the Protein Data

bank under accession codes 6K8N.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting

Information section.

ACKNOWLEDGEMENTS

We are grateful to Dr. Mei Li and the staff of beamline BL17U at the Shanghai

Synchrotron Radiation Facility and of beamline BL5A at the Photon Factory,

KEK (Tsukuba, Japan) for technical support. This study was financially

supported by National Natural Science Foundation of China [U1432241 to

Y.G. and Z.Z., 31730001 and 91751000 to L.H., 92051109 to Z.Z.]; Strategic

Priority Research Program of the Chinese Academy of Sciences

[XDB37040302 to Y.D.]; National Basic Research Program of China

[2017YFA0504900 to Y.D.]; and Beijing Municipal Science and Technology

Commission [Z191100007219007 to Y.D.]. Funding for open access charges

was provided by the National Natural Science Foundation of China.

We declare no conflict of interest.

REFERENCE

1.   Wang JC. 1996. DNA topoisomerases. Annu Rev Biochem 65:635-92.

2.   Champoux JJ. 2001. DNA topoisomerases: structure, function, and

mechanism. Annu Rev Biochem 70:369-413.

3. Vos SM, Tretter EM, Schmidt BH, Berger JM. 2011. All tangled up: how

cells direct, manage and exploit topoisomerase function. Nat Rev Mol

Cell Biol 12:827-41.

4. Pommier Y, Sun Y, Huang SN, Nitiss JL. 2016. Roles of eukaryotic

topoisomerases in transcription, replication and genomic stability. Nat

Rev Mol Cell Biol 17:703-721.

5. Forterre P. 2006. DNA topoisomerase V: a new fold of mysterious origin.

Trends Biotechnol 24:245-7.

6. Lima CD, Wang JC, Mondragon A. 1994. Three-dimensional structure

of the 67K N-terminal fragment of E. coli DNA topoisomerase I. Nature

367:138-46.

7. Mondragon A, DiGate R. 1999. The structure of Escherichia coli DNA

topoisomerase III. Structure 7:1373-83.

8. Hansen G, Harrenga A, Wieland B, Schomburg D, Reinemer P. 2006.

Crystal structure of full length topoisomerase I from Thermotoga

maritima. J Mol Biol 358:1328-40.

9. Tan K, Cao N, Cheng B, Joachimiak A, Tse-Dinh YC. 2016. Insights

from the Structure of Mycobacterium tuberculosis Topoisomerase I with

a Novel Protein Fold. J Mol Biol 428:182-93.

10. Cao N, Tan K, Zuo X, Annamalai T, Tse-Dinh Y-C. 2020. Mechanistic

insights from structure of Mycobacterium smegmatis topoisomerase I

with ssDNA bound to both N- and C-terminal domains. Nucleic acids

research 48:4448-4462.

11.   Bocquet N, Bizard AH, Abdulrahman W, Larsen NB, Faty M, Cavadini S,

Bunker RD, Kowalczykowski SC, Cejka P, Hickson ID, Thoma NH. 2014.

Structural and mechanistic insight into Holliday-junction dissolution by

topoisomerase IIIalpha and RMI1. Nat Struct Mol Biol 21:261-8.

12.   Goto-Ito S, Yamagata A, Takahashi TS, Sato Y, Fukai S. 2017.

Structural basis of the interaction between Topoisomerase IIIbeta and

the TDRD3 auxiliary factor. Sci Rep 7:42123.

13.   Rodriguez AC, Stock D. 2002. Crystal structure of reverse gyrase:

insights into the positive supercoiling of DNA. Embo j 21:418-26.

14.   Changela A, DiGate RJ, Mondragon A. 2001. Crystal structure of a

complex of a type IA DNA topoisomerase with a single-stranded DNA

molecule. Nature 411:1077-81.

15.   Zhang Z, Cheng B, Tse-Dinh YC. 2011. Crystal structure of a covalent

intermediate in DNA cleavage and rejoining by Escherichia coli DNA

topoisomerase I. Proc Natl Acad Sci U S A 108:6939-44.

16.   Berger JM, Fass D, Wang JC, Harrison SC. 1998. Structural similarities

between topoisomerases that cleave one or both DNA strands. Proc

Natl Acad Sci U S A 95:7876-81.

17.   Brown PO, Cozzarelli NR. 1981. Catenation and knotting of duplex DNA

by type 1 topoisomerases: a mechanistic parallel with type 2

topoisomerases. Proc Natl Acad Sci U S A 78:843-7.

18.   Kirkegaard K, Wang JC. 1985. Bacterial DNA topoisomerase I can relax

positively supercoiled DNA containing a single-stranded loop. J Mol Biol

185:625-37.

19.   Li Z, Mondragon A, DiGate RJ. 2001. The mechanism of type IA

topoisomerase-mediated DNA topological transformations. Mol Cell

7:301-7.

20.   Cejka P, Plank JL, Dombrowski CC, Kowalczykowski SC. 2012.

Decatenation of DNA by the S. cerevisiae Sgs1-Top3-Rmi1 and RPA

complex: a mechanism for disentangling chromosomes. Mol Cell

47:886-96.

21.   Leelaram MN, Bhat AG, Godbole AA, Bhat RS, Manjunath R, Nagaraja

V. 2013. Type IA topoisomerase inhibition by clamp closure. FASEB J

27:3030-8.

22.   Dekker NH, Rybenkov VV, Duguet M, Crisona NJ, Cozzarelli NR,

Bensimon D, Croquette V. 2002. The mechanism of type IA

topoisomerases. Proc Natl Acad Sci U S A 99:12126-31.

23.   Terekhova K, Marko JF, Mondragon A. 2014. Single-molecule analysis

uncovers the difference between the kinetics of DNA decatenation by

bacterial topoisomerases I and III. Nucleic Acids Res 42:11657-67.

24.   Mills M, Tse-Dinh YC, Neuman KC. 2018. Direct observation of

topoisomerase IA gate dynamics. Nat Struct Mol Biol 25:1111-1118.

25.   Liu LF, Wang JC. 1979. Interaction between DNA and Escherichia coli

DNA topoisomerase I. Formation of complexes between the protein and

superhelical and nonsuperhelical duplex DNAs. J Biol Chem

254:11082-8.

26.   Tse Y, Wang JC. 1980. E. coli and M. luteus DNA topoisomerase I can

catalyze catenation of decatenation of double-stranded DNA rings. Cell

22:269-76.

27.   DiGate RJ, Marians KJ. 1988. Identification of a potent decatenating

enzyme from Escherichia coli. J Biol Chem 263:13366-73.

28.   Terekhova K, Gunn KH, Marko JF, Mondragon A. 2012. Bacterial

topoisomerase I and topoisomerase III relax supercoiled DNA via

distinct pathways. Nucleic Acids Res 40:10432-40.

29.   Li Z, Mondragon A, Hiasa H, Marians KJ, DiGate RJ. 2000.

Identification of a unique domain essential for Escherichia coli DNA

topoisomerase III-catalysed decatenation of replication intermediates.

Mol Microbiol 35:888-95.

30.   Dai P, Wang Y, Ye R, Chen L, Huang L. 2003. DNA topoisomerase III

from the hyperthermophilic archaeon Sulfolobus solfataricus with

specific DNA cleavage activity. J Bacteriol 185:5500-7.

31.   Couturier M, Gadelle D, Forterre P, Nadal M, Garnier F. 2020. The

reverse gyrase TopR1 is responsible for the homeostatic control of DNA

supercoiling in the hyperthermophilic archaeon Sulfolobus solfataricus.

Molecular Microbiology 113:356-368.

32.   Bizard AH, Yang X, Debat H, Fogg JM, Zechiedrich L, Strick TR,

Garnier F, Nadal M. 2018. TopA, the Sulfolobus solfataricus

topoisomerase III, is a decatenase. Nucleic Acids Res 46:861-872.

33.   Capranico G, Marinello J, Chillemi G. 2017. Type I DNA

Topoisomerases. J Med Chem 60:2169-2192.

34.   Schoeffler AJ, Berger JM. 2008. DNA topoisomerases: harnessing and

constraining energy to govern chromosome topology. Q Rev Biophys

41:41-101.

35.   Feinberg H, Lima CD, Mondragon A. 1999. Conformational changes in

E. coli DNA topoisomerase I. Nat Struct Biol 6:918-22.

36.   Morales R, Sriratana P, Zhang J, Cann IK. 2011. Methanosarcina

acetivorans C2A topoisomerase IIIalpha, an archaeal enzyme with

promiscuity in divalent cation dependence. PLoS One 6:e26903.

37.   Holm L, Sander C. 1995. Dali: a network tool for protein structure

comparison. Trends Biochem Sci 20:478-80.

38.   Yang J, Bachrati CZ, Ou J, Hickson ID, Brown GW. 2010. Human

topoisomerase IIIalpha is a single-stranded DNA decatenase that is

stimulated by BLM and RMI1. J Biol Chem 285:21426-36.

39.   Tan K, Zhou Q, Cheng B, Zhang Z, Joachimiak A, Tse-Dinh YC. 2015.

Structural basis for suppression of hypernegative DNA supercoiling by

E. coli topoisomerase I. Nucleic Acids Res 43:11031-46.

40.   Englund PT. 2014. A passion for parasites. J Biol Chem 289:33712-29.

41.   Baker NM, Rajan R, Mondragon A. 2009. Structural studies of type I

topoisomerases. Nucleic Acids Res 37:693-701.

42.   Chen L, Huang L. 2006. Oligonucleotide cleavage and rejoining by

topoisomerase III from the hyperthermophilic archaeon Sulfolobus

solfataricus: temperature dependence and strand annealing-promoted

DNA religation. Mol Microbiol 60:783-94.

43.   Viard T, de la Tour CB. 2007. Type IA topoisomerases: a simple puzzle?

Biochimie 89:456-67.

44.   Viard T, Lamour V, Duguet M, Bouthier de la Tour C. 2001.

Hyperthermophilic topoisomerase I from Thermotoga maritima. A very

efficient enzyme that functions independently of zinc binding. J Biol

Chem 276:46495-503.

45.   Viard T, Cossard R, Duguet M, de La Tour CB. 2004. Thermotoga

maritima-Escherichia coli chimeric topoisomerases. Answers about

involvement of the carboxyl-terminal domain in DNA topoisomerase

I-mediated catalysis. J Biol Chem 279:30073-80.

46.   Zhang HL, DiGate RJ. 1994. The carboxyl-terminal residues of

Escherichia coli DNA topoisomerase III are involved in substrate binding.

J Biol Chem 269:9052-9.

47.   Krishna SS, Majumdar I, Grishin NV. 2003. Structural classification of

zinc fingers: survey and summary. Nucleic Acids Res 31:532-50.

48. Tse-Dinh YC. 1991. Zinc (II) coordination in Escherichia coli DNA

topoisomerase I is required for cleavable complex formation with DNA.

J Biol Chem 266:14317-20.

49. Liu H, Naismith JH. 2008. An efficient one-step site-directed deletion,

insertion, single and multiple-site plasmid mutagenesis protocol. BMC

Biotechnol 8:91.

50. Wang H, Liu X, Zhao J, Yue Q, Yan Y, Gao Z, Dong Y, Zhang Z, Fan Y,

Tian J, Wu N, Gong Y. 2018. Crystal structures of multicopper oxidase

CueO G304K mutant: structural basis of the increased laccase activity.

Scientific Reports 8:14252.

51. Otwinowski Z, Minor W. 1997. Processing of X-ray diffraction data

collected in oscillation mode. Methods Enzymol 276:307-326.

52. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC,

Read RJ. 2007. Phaser crystallographic software. J Appl Crystallogr

40:658-674.

53. Afonine PV, Grosse-Kunstleve RW, Adams PD. 2005. A robust

bulk-solvent correction and anisotropic scaling procedure. Acta

Crystallogr D Biol Crystallogr 61:850-5.

54. Murshudov GN, Skubak P, Lebedev AA, Pannu NS, Steiner RA,

Nicholls RA, Winn MD, Long F, Vagin AA. 2011. REFMAC5 for the

refinement of macromolecular crystal structures. Acta Crystallogr D Biol

Crystallogr 67:355-67.

55.   Emsley P, Cowtan K. 2004. Coot: model-building tools for molecular graphics. Acta Crystallogr D Biol Crystallogr 60:2126-32.

56.   Chen VB, Arendall WB, 3rd, Headd JJ, Keedy DA, Immormino RM, Kapral GJ, Murray LW, Richardson JS, Richardson DC. 2010. MolProbity: all-atom structure validation for macromolecular crystallography. Acta Crystallogr D Biol Crystallogr 66:12-21.

TABLE AND FIGURES LEGENDS

Table 1. Data collection and refinement statistics

	Native *Sso* topo III
Data collection	
Space group	P2$_1$
Cell dimensions	
a, b, c (Å)	110.7, 90.0, 156.2

α, β, γ (°)	90.0, 100.5, 90.0
Wavelength (Å)	0.98
Resolution (Å)	50-2.10 (2.14-2.10) [a]
Rmerge	0.057 (0.942)
⟨ I/σ(I)⟩	25.8 (2.0)
Completeness (%)	98.0 (94.5)
Redundancy	7.0
Refinement	
Resolution (Å)	50-2.10
No. reflections	171,737
Rwork/ Rfree	0.203/0.247
No. atoms	
Protein	21,559
DNA	-
ZN	4
Water	1020
B-factors	46.0

Rmsd bond length (Å)	0.008
Rmsd bond angle (°)	0.9
Ramachandran Plot	
Favoured (%)	97.8
Allowed (%)	2.2
Outliers (%)	0.0

a. The values in parenthesis mean those of the highest resolution shell.

FIGURES LEGENDS

FIG 1 Overall structure of full-length *Sso* topo III in apo state.

A. Domain organization of *Sso* topo III. *Sso* topo III is composed of N-terminal four domains (domain I-IV) and the unique carboxyl-terminal domain V. Domains I-V are colored hotpink, green, cyan, orange, red, respectively.

B. C. Two views of *Sso* topo III. Domains I-V are colored as the legend to Figure 1A. The close-up view (D) is highlighted by the solid box in B. Detailed

interactions (E and F) within the domains II/V interface is highlighted by the

solid box in C.

D. The close-up view shows the anomalous difference electron density map

(drawn in cyan mesh) around Zn (II) contoured at 15σ using the diffraction data

collected at the Zn absorption peak. The three Cys (C602, C605, C615) and

one His residue (H618) coordinating Zn (II) are shown in sticks. Zn (II) is

shown as a gray sphere.

E. Hydrophobic zipper between domain V residues from helix α19 and domain

II residues from the loop linking domain I and II. The hydrophobic interactions

are shown in space-filling presentations.

F. As in E but describing the corresponding hydrogen-bonding network. The

hydrogen bonds are shown as black dashed lines.

FIG 2 Comparison of structural elements around the hinge region in *Sso* topo

III, EcTOP1, EcTOP3, HsTop3α and HsTop3β.

A. Overall structural comparison among *Sso* topo III (hotpink), EcTOP3 (green)

and HsTop3α (cyan). For clarity, the RMI1 subunit of HsTop3α complex (PDB

entry 4CHT) is not shown. Domain V of *Sso* topo III is indicated. The

decatenation loop and the acidic loop of EcTOP3 (PDB entry 1D6M) are

shown, and their movement during the gate dynamics is shown with opposing

arrows.

B. The close-up views around the hinge regions of HsTop3α in Top3α-RMI1

complex (PDB entry 4CHT), HsTop3β in Top3β-TDRD3 complex (PDB entry

5GVE) complexes, and EcTOP1, mainly showing the hydrophobic interactions.

The hydrogen bonds are shown as black dashed lines. For clarity, the ssDNA

molecule of EcTOP1 complex (PDB entry 4RUL) is not shown.

C. The close-up view around the hinge region of *Sso* topo III, showing the

hydrophobic interactions. For clarity, helix α20 (residues 639-654) of *Sso* topo

III is not shown.

FIG 3 Relaxation activity of wild-type and mutant *Sso* topo III proteins.

pUC18 DNA was incubated at 75 °C for 30 min in relaxation reaction buffer (50

mM Tris-HCl (pH 8.8), 20 mM $MgCl_2$, 1 mM DTT, 0.1 mM EDTA, 90 mM

sodium chloride, 30 μgml^{-1} BSA, and 12% (vol/vol) ethylene glycol)) with

wild-type *Sso* topo III (top panel), 597ΔC (middle panel) and CCCH/4A (bottom

panel) at 500, 250, 62.5, 12, 6, 3, 1.5 and o fmol. Reactions were terminated;

the DNA products were separated in 1.4% agarose gels and visualized by

staining with ethidium bromide. Oc indicates open circle nicked or gapped

circular DNA; Sc denotes negatively supercoiled circular DNA; and Rel is the

relaxed topoisomers.

FIG 4 Time-course of DNA relaxation by wild-type and mutants *Sso* topo III

proteins.

DNA relaxation by wild-type *Sso* topo III (top panel), 597ΔC (middle panel) and

CCCH/4A mutant (bottom panel), were carried out over the indicated time

periods as described in the FIG 3. The molar ratio of protein to pUC18 DNA

was 1:1 in the reaction. Samples terminated at the indicated time-points.

Sample at o-min point indicated untreated substrate pUC18 DNA. The reaction

products were analyzed in 1.4% agarose gels and visualized by staining with

ethidium bromide. Oc indicates open circle nicked or gapped circular DNA; Sc

denotes negatively supercoiled circular DNA; and Rel is the relaxed

topoisomers.

FIG 5 kinetoplast DNA (kDNA) decatenation by wild-type and mutants *Sso*

topo III proteins.

Decatenation assays were carried out as described for DNA relaxation assay

except that kDNA (400 ng) was used as substrate with wild-type *Sso* topo III,

597ΔC and CCCH/4A mutant at 1000, 500, 250, 62.5 and 12 fmol. The

reactions were performed at 90°C for 30 min and were terminated. Samples

were electrophoresed through a 2% agarose gel, and the DNA products were

visualized by staining with ethidium bromide. Lane CK corresponds to the

kDNA control incubated at 90°C without the enzyme. Minicircles denote

minicircle DNA released by *Sso* topo III.

814
815

FIG 1

FIG 2

A

EcTOP3
Sso topo III
HsTop3α

Domain V

acidic loop

decatenation loop

C

Sso topo III

B

HsTop3α   RMI1

HsTop3β   TDRD3

EcTOPI

FIG 3

wildtype

597ΔC

CCCH/4A

FIG 4

wildtype

597ΔC

CCCH/4A

FIG 5

A

wildtype

CK   1000   500   250   62.5   12   fmol

← kDNA

← nicked or gapped circles

Rel

← supercoiled closed circles

B

597ΔC

CK   1000   500   250   62.5   12   fmol

→ kDNA

→ nicked or gapped circles

C

CCCH/4A

CK   1000   500   250   62.5   12   fmol

← kDNA

← nicked or gapped circles