Identification and evolution of avian endogenous foamy viruses

Yicong Chen[1,2,†], Xiaoman Wei[1,2,†], Guojie Zhang[3], Edward C. Holmes[4], Jie Cui[1,*]

[1]Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan, China; Institut Pasteur

of Shanghai, Chinese Academy of Sciences, Shanghai, China; Center for Biosafety

Mega-Science, Chinese Academy of Sciences.

[2]University of Chinese Academy of Sciences, Beijing, China.

[3]State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology,

Chinese Academy of Sciences, Kunming, China; China National GeneBank, BGI-Shenzhen,

Shenzhen, China; Section for Ecology and Evolution, Department of Biology, University of

Copenhagen, Copenhagen, Denmark; Center for Excellence in Animal Evolution and

Genetics, Chinese Academy of Sciences, Kunming, China.

[4]Marie Bashir Institute for Infectious Diseases and Biosecurity, Charles Perkins Centre,

School of Life and Environmental Sciences and Faculty of Medicine and Health, University

of Sydney, Sydney, NSW, Australia.

[*]Corresponding author: E-mail: jcui@ips.ac.cn

[†]These authors contributed equally to this work.

Abstract

A history of long-term co-divergence means that foamy viruses (family *Retroviridae*) provide

an ideal framework to understanding virus-host evolution over extended time periods.

Endogenous foamy viruses (EndFVs) are rare, and to date have only been described in a

limited number of mammals, amphibians, reptiles and fish genomes. By screening 414 avian

genomes we identified EndFVs in two bird species: the Maguari Stork (*Ciconia maguari*) and

the Oriental Stork (*C. boyciana*). Analyses of phylogenetic relationships, genome structures

and flanking sequences revealed a single origin of EndFVs in *Ciconia* species. In addition,

the marked incongruence between the virus and host phylogenies suggested that this

integration event occurred independently in birds. In sum, by providing evidence that birds

can be infected with foamy viruses, we fill the last major gap in the taxonomic distribution of

foamy viruses and their animal hosts.

Key words: endogenous foamy viruses; birds; incongruence; cross-species transmission

Introduction

Retroviruses (family *Retroviridae*) are viruses of substantial medical and economic

significance as some are associated with severe infectious disease or are oncogenic

(Hayward, et al. 2015; Aiewsakun and Katzourakis 2017; Xu, et al. 2018). Retroviruses are

also of evolutionary importance as they have occasionally invaded the host germ line, leading

to the generation of endogenous retroviruses (ERVs) and hence genomic 'fossils' (Stoye

2012; Johnson 2015, 2019). ERVs are widely distributed in vertebrates (Hayward, et al.

2013; Cui, et al. 2014; Hayward, et al. 2015; Xu, et al. 2018) and provide important insights

into the origin and long-term evolution of retroviruses. However, some complex retroviruses

such as lenti-, delta- and spuma viruses, only relatively rarely appear as endogenized forms.

As well leaving a litany of endogenous copies in host genomes, foamy viruses are of

particular importance because they exhibit a history of long-term co-divergence with their

vertebrate hosts (Switzer, et al. 2005). Endogenous foamy viruses (EndFVs) were first

discovered in sloths (Katzourakis, et al. 2009), and then found in several primate genomes

(Han and Worobey 2012b, 2014; Katzourakis, et al. 2014). The subsequent discovery of

EndFV and EndFV-like copies in fish genomes indicated that foamy viruses may have a deep

evolutionary history within the vertebrates (Han and Worobey 2012a; Ruboyianes and

Worobey 2016; Aiewsakun and Katzourakis 2017). Recently, three novel endogenous foamy

viruses were identified in reptile genomes, although there is disagreement over their origins

with some suggesting virus-host co-divergence over many millions of years (Wei et al. 2019),

and others favoring cross-species transmission events (Aiewsakun, et al. 2019; Wei, et al.

2019). To date, no EndFVs have been identified in avian genomes.

Materials and Methods

Genome screening and viral genome structure identification

All 147 avian genomes available in GenBank as of June 2019 (Supplementary Table S1) and

267 genomes from the 'Bird 10K' program were screened for endogenous foamy viruses

using the TBLASTN algorithm (Altschul, et al. 1990) and the protein sequences of

representative exogenous foamy viruses, EndFVs and endogenous foamy-like viruses

(Supplementary Table S2). A 35% sequence identity over a 30% region with an e-value set to

0.0001 was used to filter significant hits (Supplementary Table S3). Viral hits within large

scaffolds (>20 kb) were assumed to represent *bona fide* ERVs. We then extended the flanking

sequence of these hits to identify the viral long terminal repeats (LTRs) using BLASTN

(Altschul, et al. 1990), LTR Finder (Xu and Wang 2007) and LTRharvest (Ellinghaus, et al.

2008). In accordance with the nomenclature proposed for ERVs (Gifford, et al. 2018),

EndFVs were identified in the genomes of the Maguari Stork (*Ciconia maguari*) and the

Oriental stork (*C. boyciana*). These were termed 'ERV-Spuma.n-Cma'

and 'ERV-Spuma.n-Cbo', respectively (in which n represents the number of the viral

sequences extracted from host genome) (Supplementary Table S4). Putative genome

structures and conserved EndFV domains were identified using BLASTP, CD-search

(Marchler-Bauer and Bryant 2004; Marchler-Bauer, et al. 2017) and ORFfinder

(https://www.ncbi.nlm.nih.gov/orffinder/) at NCBI.

Molecular dating

ERV integration time can be approximately estimated using the relation T = (D/R)/2, in

which T is the integration time (million years, MY), D is the number of nucleotide

differences per site between the pairwise LTRs, and R is the genomic substitution rate

(nucleotide substitutions per site, per year). We used the previously estimated neutral

nucleotide substitution rate for birds (1.9×10^{-9} nucleotide substitutions per site, per year)

(Zhang, et al. 2014). Two full-length ERVs-Spuma-Cma containing a pairwise intact LTRs

were used to estimate integration time in this manner (Supplementary Table S5). We

excluded ERV-Spum.1-Cbo from this dating exercise due to its defective 5' LTR.

Phylogenetic analysis

To describe the evolutionary relationship of EndFVs to other representative retroviruses,

sequences of the Pol (Supplementary data set S1) and concatenated Gag-Pol-Env proteins

(Supplementary data set S2) were aligned using MAFFT 7.222 (Katoh and Standley 2013)

and confirmed manually in MEGA X (Kumar, et al. 2018). A phylogenetic tree of these data

was then inferred using the maximum likelihood (ML) method in PhyML 3.1 (Guindon, et al.

2010), incorporating 100 bootstrap replicates to assess node robustness. The best-fit

LG+Γ+I+F of amino acid substitution was selected for both Pol and concatenated

Gag-Pol-Env protein data sets using ProtTest (Abascal, et al. 2005).

Recombination analysis

To test for recombination in these data, we: (i) compared target site duplications (TSD)

flanking the ERVs, as it has previously been shown that ERVs not flanked by the same TSDs

likely arose by provirus recombination (Hughes and Coffin 2001), and (ii) screened for

recombination in the pol and gag-pol-env nucleotide sequences of mammalian, tuatara,

amphibian, lobe-finned fish and avian FVs/EndFVs using the Recombination Detection

Program 4 (RDP4) (Martin, et al. 2015)

Results and Discussion

Discovery and characterization of endogenous foamy viral elements in avian genomes

To identify potential foamy (-like) viral elements in birds, we collected 147 available bird

genomes from GenBank (Supplementary Table 1) and 267 genomes from the 'Bird 10K'

project (Zhang, et al. 2015) and performed *in silico* TBLASTN, using the amino acid

sequences of representative retroviruses as queries (Supplementary Table 2). This genomic

mining identified 16 significant hits in the Maguari Stork and the 12 in Oriental Stork

(Supplementary Table 3). We designated these ERV-Spuma.n-Cma and Spuma.n-Cbo,

respectively (Gifford, et al. 2018) (Supplementary Table S3, Table S4). We considered hits

within large scaffolds (>20 kilobases in length) to represent *bona fide* ERVs. We then

extended the flanking sequences of these EndFVs on both sides to search for LTRs, as these

define the boundary of the viral elements. Through this analysis we discovered two

full-length EndFVs in the Maguari stork genome and one in the Oriental stork genome. The

low copy number of EndFVs found in both two bird species accords with the observation that

avian genomes generally harbor small numbers of endogenous viruses (Cui, et al. 2014).

To further elucidate the relationship between these novel avian EndFVs and other

retroviruses, Pol gene sequences (>500 amino acid residue in length) were used in a

phylogenetic analysis (Fig. 1). Accordingly, our maximum likelihood phylogenetic tree

revealed that the EndFVs discovered in birds formed a close and well supported

monophyletic group within the foamy virus clade compatible with the idea that these avian

EndFVs might have a single origin. Notably, however, because they were most closely

related to the endogenous foamy viruses found in mammals rather than to those found in

reptiles, the phylogenetic position of the avian EndFVs described here was incongruent with

that of the host phylogeny (although the node associated with the tuatara EndFV was

relatively poorly supported) (Fig. 2). This, and the overall rarity of EndFVs in birds, suggests

that these avian EndFVs have an independent origin in birds and were not acquired through

virus-host co-divergence, such that a non-avian retrovirus jumped into birds at some point

during evolutionary history. No evidence for recombination was found in these data.

Genomic structure characterization

By searching for conserved domains against the Conserved Domains Database

(www.ncbi.nlm.nih.gov/Structure/cdd), we identified three typical foamy conserved domains

in the three full-length avian EndFVs: (1) the Spuma virus Gag domain (pfam03276)

(Winkler, et al. 1997), (2) the Spuma aspartic protease (A9) domain (pfam03539) that is

present in all mammalian foamy virus Pol proteins (Aiewsakun and Katzourakis 2017), and

the (3) foamy virus envelope protein domain (pfam03408) (Han and Worobey 2012a; Wei, et

al. 2019) (Supplementary Fig. S2). Furthermore, we identified an open reading frame 1

(ORF1) as an accessary gene in all three full-length avian EndFV genomes. These ORF1

sequences exhibited high nucleotide similarity (>88%) with each other, although not to any

known accessory genes from foamy viruses (and the ORF1 in ERV-Spuma.1-Cma was split

in two, due to two insertions and a nonsense mutation; Supplementary Fig. S1). Interestingly,

the internal promoter of this ORF was located between 3' end of the *env* gene and the 5' start

of the ORF gene. This is inconsistent with exogenous foamy viruses whose internal

promoters are located toward the 3' end of the *env* gene (Campbell, et al. 1994; Lochelt, et al.

1995).

In summary, the genomes of the new EndFVs documented here contained a pair of LTRs,

although with no sequence similarity to other EndFV LTRs, and exhibited a typical spuma

virus structure, with three main open reading frames (ORF) - gag, pol, and env - and one

putative additional accessory gene, ORF 1 (Fig. 3, Supplementary Fig. S1).

Vertical transmission of bird EndFVs

Surprisingly, ERV-Spuma.2-Cma and ERV-Spuma.1-Cbo shared 98% nucleotide sequence

identity and contained the same deletion in the pol gene (Supplementary Fig. S3). By

comparing the flanking sequences of these two EndFVs using BLASTN, we discovered that

two scaffolds containing EndFV (BDFF02011124.1 in the Oriental stork and scaffold3222 in

the Maguari Stork) shared 99% coverage with 98.66% sequence identity (e-value = 0.0).

Furthermore, upon the ERV insertion, the target DNA fragment is also duplicated, resulting

in target site duplication (TSDs) that differs among ERV insertions (Hughes and Coffin

2001). We were able to identify the same TSD flanking these two avian EndFVs

(Supplementary Fig. S3), confirming that they have been vertically inherited. As such, we

can infer that the most recent common ancestor of Maguari Stork and Oriental Stork,

estimated to have existed between 3.2-13.2 million years ago (Jetz, et al. 2012), carried these

EndFVs. However, neither EndFVs nor any solo LTRs were present in other bird species

from same order (Pelecaniformes), including the little egret (*Egretta garzetta*), crested ibis

(*Nipponia nippon*) and Yellow-throated sandgrouse (*Pterocles gutturalis*). Clearly, the study

of additional genomes sampled across the avian phylogeny is merited.

Estimation of ERV insertion times

To approximately estimate the insertion time of endogenous foamy viruses in two birds, we

utilized the LTR-divergence based on the degree of sequence divergence between the 5' and

3' LTRs with a known host nucleotide substitution rate (Dangel, et al. 1995; Johnson and

Coffin 1999). Accordingly, the two intact pairwise LTRs of ERVs-Spuma-Cma were selected

for time estimation (Supplementary Table S5). This analysis revealed insertion times of 3.15

and 13.95 MYA (million years ago), close to the estimated time of common ancestor of

Maguari Stork and Oriental Stork (Jetz, et al. 2012). However, the presence of multiple

premature stop codon suggests the invasion was ancient, and all estimates of integration times

should be treated with caution (Kijima and Innan 2010). Clearly, the discovery of additional

EndFVs will shed more light on the time-scale of these retroviral integration events.

In sum, we describe the presence and evolution of two novel avian endogenous foamy

viruses. This discovery fills the last major gap in our understanding of the taxonomic

distribution of the foamy viruses and helps reveal the evolutionary interactions between

retroviruses and their hosts over extended time-periods.

Acknowledgements

This work was supported by the Special Key Project of Biosafety Technologies

(2017YFC1200800) for the National Major Research & Development Program of China, and

Strategic Priority Research Program of the Chinese Academy of Sciences (XDA13010500).

J.C. is supported by National Natural Science Foundation of China (31671324, 31970176)

and CAS Pioneer Hundred Talents Program. E.C.H. is supported by an ARC Australian

Laureate Fellowship (FL170100022). G.Z. is supported by Strategic Priority Research

Program of the Chinese Academy of Sciences (XDB31020000) and Villum Foundation. We

thank B10K consortium for the early access of all bird genomes produced by B10K.

References

Abascal F, Zardoya R, Posada D. 2005. ProtTest: selection of best-fit models of protein

evolution. Bioinformatics 21:2104-2105.

Aiewsakun P, Katzourakis A. 2017. Marine origin of retroviruses in the early Palaeozoic Era.

Nat Commun 8:13954.

Aiewsakun P, Simmonds P, Katzourakis A. 2019. The First Co-Opted Endogenous Foamy

Viruses and the Evolutionary History of Reptilian Foamy Viruses. Viruses 11:641.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search

tool. J Mol Biol 215:403-410.

Campbell M, Renshaw-Gegg L, Renne R, Luciw PA. 1994. Characterization of the internal

promoter of simian foamy viruses. J Virol 68:4811-4820.

Cui J, Zhao W, Huang Z, Jarvis ED, Gilbert MT, Walker PJ, Holmes EC, Zhang G. 2014.

Low frequency of paleoviral infiltration across the avian phylogeny. Genome Biol 15:539.

Dangel AW, Baker BJ, Mendoza AR, Yu CY. 1995. Complement component C4 gene intron

9 as a phylogenetic marker for primates: long terminal repeats of the endogenous retrovirus

ERV-K(C4) are a molecular clock of evolution. Immunogenetics 42:41-52.

Ellinghaus D, Kurtz S, Willhoeft U. 2008. LTRharvest, an efficient and flexible software for

de novo detection of LTR retrotransposons. BMC Bioinformatics.

Gifford RJ, Blomberg J, Coffin JM, Fan H, Heidmann T, Mayer J, Stoye J, Tristem M,

Johnson WE. 2018. Nomenclature for endogenous retrovirus (ERV) loci. Retrovirology

15:59.

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing the

performance of PhyML 3.0. Syst Biol 59:307-321.

Han GZ, Worobey M. 2012a. An endogenous foamy-like viral element in the coelacanth

genome. PLoS Pathog 8:e1002790.

Han GZ, Worobey M. 2012b. An endogenous foamy virus in the aye-aye (Daubentonia

madagascariensis). J Virol 86:7696-7698.

Han GZ, Worobey M. 2014. Endogenous viral sequences from the Cape golden mole

(Chrysochloris asiatica) reveal the presence of foamy viruses in all major placental mammal

clades. PLoS One 9:e97931.

Hayward A, Cornwallis CK, Jern P. 2015. Pan-vertebrate comparative genomics unmasks

retrovirus macroevolution. Proc Natl Acad Sci U S A 112:464-469.

Hayward A, Grabherr M, Jern P. 2013. Broad-scale phylogenomics provides insights into

retrovirus-host evolution. Proc Natl Acad Sci U S A 110:20146-20151.

Hughes JF, Coffin JM. 2001. Evidence for genomic rearrangements mediated by human

endogenous retroviruses during primate evolution. Nat Genet 29:487-489.

Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. 2012. The global diversity of birds in

space and time. Nature 491:444.

Johnson WE. 2015. Endogenous Retroviruses in the Genomics Era. Annu Rev Virol

2:135-159.

Johnson WE. 2019. Origins and evolutionary consequences of ancient endogenous

retroviruses. Nat Rev Microbiol 17:355-370.

Johnson WE, Coffin JM. 1999. Constructing primate phylogenies from ancient retrovirus

sequences. Proc Natl Acad Sci U S A 96:10254-10260.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol 30:772-780.

Katzourakis A, Aiewsakun P, Jia H, Wolfe ND, LeBreton M, Yoder AD, Switzer WM. 2014.

Discovery of prosimian and afrotherian foamy viruses and potential cross species

transmissions amidst stable and ancient mammalian co-evolution. Retrovirology 11:61.

Katzourakis A, Gifford RJ, Tristem M, Gilbert MT, Pybus OG. 2009. Macroevolution of

complex retroviruses. Science 325:1512.

Kijima TE, Innan H. 2010. On the estimation of the insertion time of LTR retrotransposable

elements. Mol Biol Evol 27:896-904.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. Mol Biol Evol 35:1547-1549.

Lochelt M, Yu SF, Linial ML, Flugel RM. 1995. The human foamy virus internal promoter is

required for efficient gene expression and infectivity. Virology 206:601-610.

Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, Chitsaz F, Derbyshire MK, Geer

RC, Gonzales NR, et al. 2017. CDD/SPARCLE: functional classification of proteins via

subfamily domain architectures. Nucleic Acids Res 45:D200-d203.

Marchler-Bauer A, Bryant SH. 2004. CD-Search: protein domain annotations on the fly.

Nucleic Acids Res 32:W327-331.

Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. 2015. RDP4: Detection and analysis

of recombination patterns in virus genomes. Virus Evol 1:vev003.

Ruboyianes R, Worobey M. 2016. Foamy-like endogenous retroviruses are extensive and

abundant in teleosts. Virus Evol 2:vew032.

Stoye JP. 2012. Studies of endogenous retroviruses reveal a continuing evolutionary saga.

Nat Rev Microbiol 10:395-406.

Switzer WM, Salemi M, Shanmugam V, Gao F, Cong ME, Kuiken C, Bhullar V, Beer BE,

Vallet D, Gautier-Hion A, et al. 2005. Ancient co-speciation of simian foamy viruses and

primates. Nature 434:376-380.

Wei X, Chen Y, Duan G, Holmes EC, Cui J. 2019. A reptilian endogenous foamy virus sheds

light on the early evolution of retroviruses. Virus Evol 5:vez001.

Winkler I, Bodem J, Haas L, Zemba M, Delius H, Flower R, Flugel RM, Lochelt M. 1997.

Characterization of the genome of feline foamy virus and its proteins shows distinct features

different from those of primate spumaviruses. J Virol 71:6727-6741.

Xu X, Zhao H, Gong Z, Han GZ. 2018. Endogenous retroviruses of non-avian/mammalian

vertebrates illuminate diversity and deep history of retroviruses. PLoS Pathog 14:e1007072.

Xu Z, Wang H. 2007. LTR_FINDER: an efficient tool for the prediction of full-length LTR

retrotransposons. Nucleic Acids Res 35:W265-268.

Zhang G, Li C, Li Q, Li B, Larkin DM, Lee C, Storz JF, Antunes A, Greenwold MJ,

Meredith RW, et al. 2014. Comparative genomics reveals insights into avian genome

evolution and adaptation. Science 346:1311-1320.

Zhang G, Rahbek C, Graves GR, Lei F, Jarvis ED, Gilbert MT. 2015. Genomics: Bird

sequencing project takes off. Nature 522:34.

Fig. 1. Phylogenetic tree of retroviruses and endogenous retroviruses, including the

endogenous foamy viruses found in avian genomes. The tree was inferred using amino acid

sequences of the Pol gene, and rooted using the remaining retroviral taxa (excluding the

spumaviruses). The newly identified viral elements are labelled in red. * indicates the EndFV

found in the Maguari Stork genome, while † denotes the EndFV from the Oriental Stork

genome. The scale bar indicates the number of amino acid changes per site.

Fig. 2. Evolutionary history of foamy viruses (left) and their vertebrate hosts (right).

Associations between foamy viruses and their hosts are indicated by connecting lines. The

avian EndFV and its host are labelled in red. Note that the avian EndFVs are more closely

related to mammalian foamy viruses than the reptilian EndFV. Scale bars indicate the number

of amino acid changes per site in the viruses and the host divergence times (million years

ago, MYA).

Fig. 3. Genomic organization of exogenous foamy viruses (acc: NC_001364) and avian

endogenous foamy viruses. LTR, long terminal repeat; Pro, protease; RT, reverse

transcriptase; RH, ribonuclease H; IN, integrase.

FV Gag-Pol-Env phylogeny