**DNA METHYLATION BIOMARKERS OF MYOCARDIAL INFARCTION
AND CARDIOVASCULAR DISEASE**

Epigenetic biomarkers of myocardial infarction

Alba Fernández-Sanlés, PhD[a,b,c]; Sergi Sayols-Baixeras, PhD[a,d,e]; Isaac Subirana, PhD[a,f];
Mariano Sentí, MD, PhD[b]; Pérez-Fernández S, BSc, MSc[a,d]; Manuel Castro de Moura, PhD[g];
Manel Esteller, MD, PhD[g,h,i,j,k]; Jaume Marrugat, MD, PhD[a,d]; Roberto Elosua, MD, PhD[a,d,l].

a.-Cardiovascular Epidemiology and Genetics Research Group, REGICOR Study group,
IMIM (Hospital del Mar Medical Research Institute), Barcelona, Catalonia, Spain.
b.-Pompeu Fabra University (UPF), Barcelona, Catalonia, Spain.
c.-Medical Research Council (MRC) Integrative Epidemiology Unit at the University of Bristol,
Bristol, UK.
d.-CIBER Cardiovascular Diseases (CIBERCV), Spain.
e.-Department of Medical Sciences, Molecular Epidemiology, Uppsala University, Sweden.
f.-CIBER Epidemiology and Public Health (CIBERESP), Spain.
g.-Cancer Epigenetics and Biology Program (PEBC), Bellvitge Biomedical Research Institute
(IDIBELL), Barcelona, Catalonia, Spain.
h.-CIBER Cancer (CIBERONC), Spain.
i.-Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Catalonia,
Spain.
j.-Physiological Sciences Department, School of Medicine and Health Sciences, University of
Barcelona (UB), Barcelona, Catalonia, Spain.
k.-Josep Carreras Leukaemia Research Institute (IJC), Badalona, Catalonia, Spain.
l.-Medicine Department, Medical School, University of Vic-Central University of Catalonia
(UVic-UCC), Vic, Catalonia, Spain.

Corresponding author: Roberto Elosua, MD, PhD
IMIM, Hospital del Mar Medical Research Institute
Dr Aiguader 88, 08003 Barcelona, Catalonia, Spain
Telephone: (+34) 933 160800; Email: relosua@imim.es

Subject codes: Biomarkers; Epidemiology; Epigenetics; Myocardial Infarction.

Word count: 5,359

Total number of figures and tables: 5

TOC category: Clinical and population studies

TOC subcategory: Arteriosclerosis

ABSTRACT

Objective: To assess the association between DNA methylation and acute myocardial infarction, the predictive added value of the identified methylation marks, and the causality of those associations.

Approach and Results: We conducted a case-control, two-stage, epigenome-wide association study on acute myocardial infarction ($n_{discovery}$=391, $n_{validation}$=204). DNA methylation was assessed using the Infinium MethylationEPIC BeadChip (over 850,000 CpGs). DNA methylation was the exposure variable and myocardial infarction the outcome of interest. After a fixed-effects meta-analysis, 34 CpGs fulfilled Bonferroni significance. These findings were also analysed in two independent cohort studies (n~1,800 and n~2,500) with incident coronary (CHD) and cardiovascular disease (CVD). The Infinium HumanMethylation450 BeadChip was used in these two studies (over 480,000 CpGs) and only 12 of the 34 CpGs were available in those samples. Finally, we validated four of them in association with incident CHD: *AHRR*-mapping cg05575921, *PTCD2*-mapping cg25769469, intergenic cg21566642 and *MPO*-mapping cg04988978. The four CpGs were also associated with classical cardiovascular risk factors. A methylation risk score based on those CpGs did not improve the predictive capacity of the Framingham risk function. To assess the causal effects of those CpGs we performed Mendelian randomization analysis but only one metQTL could be identified and the results were not conclusive.

Conclusions: We have identified 34 CpGs related to acute myocardial infarction. These loci highlight the relevance of smoking, lipid metabolism, and inflammation in the biological mechanisms related to myocardial infarction. Four were additionally associated with incident CHD and CVD but did not provide additional predictive information.

Keywords: DNA methylation; Epigenome-wide association study; Predictive biomarkers; Myocardial infarction; Cardiovascular disease

Abbreviations

450k: Infinium HumanMethylation450 BeadChip

MI: Myocardial Infarction

CHD: Coronary Heart Disease

CVD: Cardiovascular Disease

CVR: Cardiovascular Risk

CVRF: Cardiovascular Risk Factor

EPIC: Infinium MethylationEPIC BeadChip

EWAS: Epigenome-Wide Association Study

FOS: Framingham Offspring Study

MRS: Methylation Risk Score

REGICOR: REgistre GIroní del COR

WHI: Women's Health Iniciative

INTRODUCTION

Cardiovascular disease (CVD) and more specifically coronary heart disease (CHD) remains the number one cause of death and disease burden worldwide.[1,2] At the individual level, prevention is based on the estimation of cardiovascular risk.[3] However, the sensitivity of cardiovascular risk estimation is low and a significant proportion of CHD events occurs in individuals classified as having moderate or low risk.[4] Additionally, the use of currently available drugs to control classical cardiovascular risk factors (CVRFs) does not prevent all CHD events, underlining the need to identify new strategies for reducing this residual cardiovascular risk.[5] Thus, information encoded in biological mechanisms should be unravelled to find new predictive biomarkers and potential therapeutic targets. Among these biomarkers, DNA methylation marks arise as emerging candidates.

DNA methylation is an epigenetic mechanism consisting on chemical modifications of cytosines, mostly followed by guanines (CpGs).[6] Epigenome-wide association studies (EWASs) make it possible to find DNA methylation biomarkers of different traits and outcomes. In fact, DNA methylation pattern is associated with multiple chronic diseases,[7] including CVD and CHD.[8–13] However, the clinical value of the identified biomarkers is still limited, and the epigenetic landscape underlying CVD is not completely understood.

The most common technology to assess DNA methylation is based on commercial arrays, which do not cover the whole methylome. Moreover, most current knowledge on the relation between DNA methylation and cardiovascular risk comes from studies based on the Infinium HumanMethylation450 BeadChip (Illumina, CA, USA; from now on, 450k)[14] – which has been replaced by the Infinium MethylationEPIC BeadChip (Illumina, CA, USA; from now on, EPIC). Compared to the 450k, EPIC interrogates 413,745 more methylation sites (but excludes 42,859) increasing the genomic coverage. Moreover, EPIC is enriched with functional sites analyses such as enhancers, DNase hypersensitive sites, and miRNA promoter regions.[15] Thus, the new chip has the potential to identify novel DNA methylation-based biomarkers of cardiovascular events.

We hypothesized that DNA methylation is associated with MI risk, and that some of these

epigenetic marks could be predictive of future risk, and have causal effects on cardiovascular

outcomes. Thus, this study had three aims: 1) to unravel genomic methylation loci associated

with myocardial infarction (MI), 2) to assess their predictive capacity of cardiovascular risk,

and 3) to decipher the causality of those associations.

MATERIALS AND METHODS

Study design and populations

We designed an epigenome-wide association study (EWAS) using three populations: the

Girona Heart Registry (REGICOR, REgistre GIroní del COR), the Women's Health Initiative

(WHI), and the Framingham Offspring Study (FOS). We first performed a two-stage EWAS

on acute MI using two independent age- and sex-matched case-control studies designed in

REGICOR. Then, we validated the significant results in the other two populations with

incident cases of CHD and CVD.

Case-control studies of acute MI in REGICOR

The sample used in the discovery stage (REGICOR-1) involved 416 individuals (208 MI

cases and 208 controls). The sample in the validation stage (REGICOR-2) comprised 208

individuals (104 cases and 104 controls). Cases were selected from patients who were

consecutively attended for a first acute MI in the reference hospital of the monitored area, in

the province of Girona, in the northeast of Spain. Women were overrepresented to achieve

their inclusion as 50% of our sample. Controls were participants in a population-based

survey performed in the same monitored area. They were randomly selected from those

attending the 2009-2013 follow-up visit (n=4,980), and matched by age and sex with the MI

cases.[16] All participants were of European descent and provided informed written consent.

The study was approved by the local ethics committee (2015/6199/I; 2018/7855/I) and meets

the principles expressed in the Declaration of Helsinki and the relevant Spanish legislation.

Samples with incident cases of CHD and CVD

The WHI sample is a case-control study nested in a cohort. The FOS sample is a

prospective cohort study. Both samples were available in the database of Genotypes and

Phenotypes (http://dbgap.ncbi.nlm.nih.gov; project number #9047). The graphical abstract

shows the design and flow-chart of this study.

Assessment of cardiovascular outcomes

The outcomes assessed were acute MI in REGICOR, and incident CHD and CVD in the WHI

and FOS samples. Additional details are provided in the Supplementary material.

Assessment of DNA methylation

DNA methylation was assessed genome-wide from peripheral blood with commercial arrays

from Illumina (CA, USA). The Infinium MethylationEPIC BeadChip, covering over 850,000

CpGs, was used in the REGICOR samples. The Infinium HumanMethylation450 BeadChip,

covering over 480,000 CpGs, was used in the WHI and FOS samples. A detailed quality

control pipeline for the methylation data is available in the Supplementary material.

Methylation status at each CpG was reported by β-values.[17]

Covariates

In the REGICOR case-control studies the following covariates were considered: self-reported

smoking, diabetes, hypercholesterolemia and hypertension (Supplementary material). In the

WHI and FOS studies self-reported smoking and glycaemia, total and HDL cholesterol, and

blood pressure measurement were considered. Moreover, we inferred the peripheral blood

cell counts with the *FlowSorted.Blood.450k* R package.[18] We also estimated two surrogate

variables for unknown sources of potential technical or biological confounding using the *sva*

R package.[19]

Statistical analysis

All statistical analyses were performed using R version 3.4.0. The codes of the Singularity

images are available in the repositories at https://github.com/regicor/methylation_ami/. A

detailed description of the statistical methods is provided in the Supplementary material.

Association between DNA methylation and cardiovascular outcomes

Logistic regression was used in the analyses in the REGICOR and WHI samples, while Cox

regression was used in the FOS sample. We considered the cardiovascular event (acute MI,

CHD or CVD) as the outcome and DNA methylation as the exposure.

We defined three models. Model 1 was adjusted for estimated cell counts and two surrogate

variables (plus age and ethnicity in the WHI sample, plus age and sex in the FOS samples).

Model 2 was additionally adjusted for smoking. Model 3 was further adjusted for diabetes,

hypercholesterolemia and hypertension.

In order to reduce epigenomic inflation, we corrected the coefficients, the standard errors and

the *p* values using the *bacon* R package if necessary.[20] We used coefficients and standard

errors from the regression models as the input data and we set a random seed at 123.

We selected those associations from the discovery stage (REGICOR-1) with a corrected *p-*

value$<10^{-5}$ for assessment in the validation stage (REGICOR-2). Moreover, we performed a

fixed-effect meta-analysis of the corrected effect sizes observed in both stages, weighted by

the inverse of the variance. We defined as statistically significant any association fulfilling the

Bonferroni correction for multiple comparisons (0.05 divided by the number of probes

analyzed). Thereafter, we studied the association of the identified CpGs with incident CHD

and with CVD events in the WHI and the FOS samples, separately. The results from both

samples were meta-analysed (for CHD and CVD, separately), and significance was defined

accordingly to the number of probes analyzed.

Association between the identified CpGs and CVRFs

We analysed whether the methylation levels of the identified CpGs were associated with

individual CVRFs in the four samples using multiple linear regression, and then meta-

analysed the results. Significance was also defined accordingly to Bonferroni criteria.

Methylation risk score (MRS) and predictive capacity

We developed a weighted MRS based on the CpGs identified. We evaluated the association

between this score and CVD and CHD incidence in the FOS sample, using Cox regression.

All analyses were adjusted for age, sex, diabetes, smoking, systolic blood pressure,

hypertensive treatment, and levels of total cholesterol and cholesterol in high-density

lipoproteins (HDL-C).[21] We also assessed the potential added predictive value of including

the MRS in the Framingham risk function. We evaluated the increase in the discrimination

and the reclassification.

***In silico* functional analysis**

We performed a pathway and network analysis of the genes found as differentially

methylated in association with MI, using the Ingenuity Pathway Analysis (IPA) software

(http://www.ingenuity.com/, QIAGEN, Redwood City, CA, USA). We uploaded the Gene

symbol identifiers of those genes. The reference database underlying IPA was the Ingenuity

Knowledge Base (Genes only). Only human annotations were considered. Pathway analyses

were performed with IPA's Core Analysis module. "Canonical pathways" and "Diseases and

functions" terms with a $p<0.05$ after Benjamini-Hochberg multiple testing correction were

defined as a statistically significant overrepresentation of input genes in a given process.

Causality of associations between DNA methylation and cardiovascular outcomes

We performed Mendelian Randomization studies using the MR-Base platform.[22] We selected

methylation-level quantitative trait loci (meQTL) from a public database[23] as instrumental

variables, and interrogated their association with CHD using summary statistic data from a

meta-analysis of GWAS assessing CHD.[24] A more detailed description of the analysis is

included in the Supplementary material.

RESULTS

Quality control of DNA methylation data, cardiovascular outcomes and covariates

We finally included 391 individuals (196 cases and 195 controls) in the REGICOR-1 sample,

204 individuals (101 cases and 103 controls) in the REGICOR-2 sample, 1,863 women in the

WHI sample, and 2,540 participants in the FOS sample. The main sociodemographic and

clinical characteristics of the three populations are shown in Tables 1 and 2. Regarding the

number of CpGs, we analyzed 811,610 CpGs in the REGICOR-1 sample, 820,183 CpGs in

the REGICOR-2 sample, 478,369 CpGs in the WHI sample, and 483,656 CpGs in the FOS

sample.

Association between DNA methylation and cardiovascular outcomes

A.- Two-stage EWAS on acute myocardial infarction

A.1. Discovery stage:

The associations from the discovery stage (REGICOR-1) that were taken to the subsequent

validation (*p-value*<10^{-5}), and their Manhattan and Q-Q plots are shown in the

Supplementary Table 1, and Supplementary Figures 1-2. In total, we identified 68

suggestively significant CpGs (Supplementary Figure 3). Model 1 provided 56 CpGs, of

which three were also found in both model 2 and 3, and 13 in model 2. One additional CpG

was found in both model 2 and 3, two in model 2 and nine in model 3.

A.2. Validation and meta-analysis:

The association studies performed in the validation stage included the 68 suggestively

significant CpGs. We meta-analyzed the results of those 68 associations from both stages.

We identified 34 differentially methylated CpGs related to MI, with similar effect sizes in all

three models and both stages for most of the CpGs. Specifically, all of them were statistically

significant (*p-value*$<6.17 \cdot 10^{-8}$) in model 1, 17 in model 2, and three in model 3 (Table 3, and

Supplementary Table 2). The 34 CpGs were located in 25 different loci (26 genes, with one

CpG mapping to two genes) and nine intergenic regions. Fourteen of these CpGs should be

interpreted with caution, as the *p-value* in the validation population was not significant

considering the 68 CpGs analyzed (*p-value*$>0.05/68=7.35 \cdot 10^{-4}$) (Table 3, and Supplementary

Table 2).

A.3. *In silico* functional analysis:

We performed an IPA analysis of the 26 identified genes. Among the 10 most significant

molecular and cellular functions, this set of genes was enriched on lipid metabolism. The

enriched physiological system development and functions included the connective tissue,

hematological system, hematopoiesis, and the endocrine system. Inflammatory response

and disease, metabolic disease, and cardiovascular disease were also among the 20 most

significant diseases/disorders (Supplementary Figure 4).

B.- Follow-up association studies on incident CHD and CVD events

Out of the 34 identified CpGs associated with MI, only 12 were available in the samples with

incident cases (whose DNA methylation was profiled with the array 450k). In total, we

validated four CpGs after the meta-analysis of the separate association studies in the WHI

and the FOS samples (*p-value*$<0.05/12=4.17 \cdot 10^{-3}$): *AHRR*-mapping cg05575921, *PTCD2*-

mapping cg25769469, intergenic cg21566642 and *MPO*-mapping cg04988978. The four

CpGs were associated with CHD but cg25769469 was not related to CVD (Table 4,

Supplementary Table 3).

Association between the identified CpGs and CVRFs

Table 5 shows the associations observed between the identified CpGs and classical CVRFs.

The four identified CpGs were related to some CVRF [*p-value*<0.05/(4 CpGs*8

CVRF)=$1.56 \cdot 10^{-3}$].

Association between MRS and incidence of CHD and CVD

The associations between MRS and the incidence of coronary (n=94) and cardiovascular

(n=222) events in the FOS population are shown in Supplementary Table 4. The median of

the follow-up periods for CVD and CHD incidence were 7.67 and 7.87 years, respectively.

The MRS was not associated with higher cardiovascular risk independently of the classical

CVRFs. Consistently, the addition of the MRS to the Framingham risk function did not

improve its predictive capacity in the FOS cohort (Supplementary Table 4).

Causality of the associations between DNA methylation and cardiovascular outcomes

We identified one genetic variant (rs72617176) associated with cg21566642 methylation

level. Only the Wald ratio method could be performed, and the association was not

significant (Supplementary Table 5). We couldn't create genetic instrumental variables to

assess the causality of the association of the other three CpGs with CHD.

DISCUSSION

We have identified 34 methylation loci associated with acute MI in a two-stage EWAS,

analyzing ~850,000 CpGs. All but two of these MI-associated sites (one located in *AHRR*

and the other one intergenic) are newly reported. Of those, 12 CpGs could be studied in

association with incident cases of CHD and CVD, and we identified four of them associated

with incident CHD (three of them also with incident CVD). All four were also related to

traditional CVRFs, supporting their role in the development of these diseases. However, their

clinical utility as predictive biomarkers or drug targets was not proven.

Recently, two EWASs on incident CHD were published providing different findings from ours.

Ward-Caviness *et al* found nine CpGs associated with incident acute MI.[9] Agha *et al* reported

52 CpGs related to incident CHD.[8] None of them was replicated in our study. This lack of

concordance could be related to methodological differences (incident vs prevalent cases;

considered confounder variables), and highlights the complexity of the study of these

diseases.

CpG sites associated with acute MI events

The 34 identified CpGs showed similar effect sizes in the two REGICOR samples. Although

14 CpGs were non-significant in the validation sample, we considered them potentially

relevant because the magnitude of their association with MI was comparable across the two

samples and the statistical power of the validation sample was lower. Similarly, all but three

CpGs (*AHRR*-mapping cg05575921, *F2RL3*-mapping cg03636183, and the intergenic

cg21566642) showed consistent effect sizes in the three models. The effect size of those

three was reduced by half when adjusted for smoking, which highlights the important role of

this risk factor in the MI context. In fact, all three sites are widely described to be related to

smoking.[16,25,26]

Differentially methylated genes were enriched in diverse molecular and physiological

pathways, including lipid metabolism and metabolic and inflammatory diseases, underlining

their relevance on the pathogenesis of CHD. Considering the Bonferroni correction, three

different CpGs were associated with MI independently of traditional CVRFs (smoking,

diabetes, hypertension and hypercholesterolemia), suggesting that other biological pathways

are triggered in the MI context. To our knowledge, no EWAS on any trait has previously

found any of these three CpGs. For instance, cg12066594 is located within a regulatory

region of *PSMB7*, which encodes a subunit of the proteasome, a complex essential for

protein degradation and involved in inflammation.[27] A polymorphism in a gene encoding

another proteasome subunit (*PSMA6*) has been associated with CHD and MI.[28–31] According

to the EPIC manifest, the probes of cg12066594 and cg09967176 contain both of them three

different genetic variants (rs114673478, rs569238855, rs539730482 in that of cg12066594;

and rs570492126, rs534702669, rs181647444 in that of cg09967176). rs539730482, located

in the single base extension site (one basepair upstream of the CpG site), and rs181647444,

in the CpG site, are variants in an African population and their frequency is very low (Minor

Allele Frequency <1%).[32,33] Thus, as the individuals in this study are of European ancestry

and these variants showed very low frequency in this population, the identified associations

most likely are not confounded by the genetic variants.

Nonetheless, we cannot infer the biological sequence of the epigenetic marks, the related

biological mechanisms, and the clinical event. One possible scenario could be that the

identified DNA methylation marks occurred before the acute event, as potential biological

mechanisms involved in MI pathogenesis. This may be the case of the three CpGs that were

related to smoking. Conversely, as blood samples of MI cases were collected within the initial

24 h after hospitalization, the other possibility could be that methylation at the identified

CpGs had changed as a consequence of the acute event or the therapeutic procedures. If

the first scenario can be proven in further studies, these DNA methylation marks could be

potential predictive biomarkers of MI or new therapeutic targets. If they are found to be post-

MI marks, further studies could evaluate their potential as biomarkers of prognosis.

CpG sites consistently related to prevalent and incident CVD events

Twelve of the 34 identified CpGs could be evaluated in prospective samples and four of them

were also significantly related to incident cases of CHD. cg21566642 maps to an intergenic

region, and cg05575921, cg04988978 and cg25769469 annotate to *AHRR, MPO* and

PTCD2, respectively. To our knowledge, these CpGs were not associated with

cardiovascular events in previous EWAS reports.

cg21566642 and cg05575921 were highly and inversely associated with smoking, which is

supported by previous EWAS.[25,26] We have also previously reported both CpGs as related to

age-independent cardiovascular risk,[13] and they have been related to all-cause mortality in

an EWAS.[34] cg05575921 was further associated directly with HDL-C and inversely with

cholesterol in low-density lipoproteins (LDL-C) and triglyceride levels in our study. This CpG

has been related to both CHD prevalence and incidence in a candidate gene study.[35]

cg04988978 and cg25769469 annotate to *MPO* and *PTCD2*, respectively. Both CpGs were

associated directly with HDL-C and inversely with triglyceride and glucose levels. *MPO*

encodes the myeloperoxidase, which promotes atherosclerotic lesions by enhancing APOB

oxidation within low-density lipoproteins[36] and was causally associated with incident

cardiovascular outcomes.[37] One CpG located within *PTCD2* was previously identified to be

associated with hypertension in obstructive sleep apnea patients.[38]

MRSs as predictive CVD biomarkers

To assess the value of the four identified CpGs as predictive biomarkers, we followed the

AHA recommendations.[39] However, neither we observed an association between the MRS

and the incidence of CVD events in the FOS, nor we observed an improvement in the

predictive capacity of the Framingham risk function when including this score. This highlights

the challenge of novel biomarkers to improve cardiovascular risk prediction.

Causality of the associations between methylation loci and cardiovascular outcomes

MR studies could only be performed on the association of cg21566642 with acute MI and

CHD. Although a non-causal relationship was suggested, this must be interpreted with

caution as only one genetic variable was used as the genetic instrument. In fact, the four

CpGs associated not only with acute MI, but also incident CHD, may suggest that DNA

methylation changes at those loci occur prior to the event.

Strengths and limitations

The main strength of our study is that it is the first two-stage EWAS on MI to be based on

more than 800,000 CpGs across the genome. Moreover, we aimed to validate our findings in

prospective samples of CHD and CVD as a proxy of MI. Also, we aimed to prove the clinical

relevance of our findings. However, some limitations should be acknowledged. First, two

thirds of the CpGs identified in the initial case-control study could not be assessed in the

incident studies as the methylation arrays differed in the number of CpGs (EPIC VS 450k,

respectively). Second, we used self-reported information about cardiovascular risk factors in

the case-control study, as an event such as MI modifies risk factor levels during the acute

phase. Third, we cannot infer causality since changes in methylation could have occurred as

a consequence of the acute phase and disease management of the MI event. We aimed to

perform MR studies of the association between the identified CpGs and cardiovascular

events, but available meQTL datasets are still limited. Last, our study is based on

populations of European origin and the results cannot be extrapolated to other populations.

CONCLUSIONS

Our study provides 34 novel DNA methylation loci related to MI. The results shed some light

on the molecular landscape of MI, highlighting the importance of traditional CVRFs and

inflammation in the development of CHD. Our results question the relevance of DNA

methylation as a predictive biomarker.

Acknowledgements

a) Acknowledgements: We thank Elaine M. Lilly, PhD, for her critical reading and

revision of the English text.

b) Sources of funding: This project was funded by the Carlos III Health Institute–

European Regional Development Fund (FIS PI18/00017, FIS PI15/00051,

PI12/00232, CIBERCV, CIBERESP, CIBERONC), PERIS from Agència de Gestió

d'Ajuts Universitaris i de Recerca (SLT002/16/00088) and the Government of

Catalonia through the Agency for Management of University and Research Grants

(2014SGR240, 2017SGR946). S. Sayols-Baixeras was funded by the Instituto de

Salud Carlos III Fondos FEDER (IFI14/00007). A. Fernández-Sanlés was funded by

the Spanish Ministry of Economy and Competitiveness (BES-2014–069718).

The Framingham Heart Study (FHS) is conducted and supported by the National

Heart, Lung, and Blood Institute (NHLBI) in collaboration with Boston University

(Contract No. N01-HC-25195 and HHSN268201500001I). The Women's Health

Initiative (WHI) program is funded by the NHLBI (Contracts N01WH22110, 24152,

32100-2, 32105-6, 32108-9, 32111-13, 32115, 32118-32119, 32122, 42107-26,

42129-32, and 44221). This manuscript was not prepared in collaboration with

investigators of the FHS/ WHI, has not been reviewed and/or approved by the

FHS/WHI, and does not necessarily reflect the opinions or views of the FHS and WHI

investigators or the NHLBI.

Disclosures: The authors declare that they have no competing interests.

REFERENCES
437

1.   Roth GA, Abate D, Abate KH, et al. Global, regional, and national age-sex-specific
mortality for 282 causes of death in 195 countries and territories, 1980-2017: a
systematic analysis for the Global Burden of Disease Study 2017. *Lancet.*
2018;392(10159):1736-88.
2.   James SL, Abate D, Abate KH, et al. Global, regional, and national incidence,
prevalence, and years lived with disability for 354 diseases and injuries for 195 countries
and territories, 1990–2017: a systematic analysis for the Global Burden of Disease
Study 2017. *The Lancet.* 2018;392(10159):1789-858.
3.   Piepoli MF, Hoes AW, Agewall S, et al. 2016 European Guidelines on cardiovascular
disease prevention in clinical practice. *European Heart Journal.* 2016;37(29):2315-81.
4.   Marrugat J, Vila J, Baena-Díez JM, Grau M, Sala J, Ramos R, Subirana I, Fitó M, Elosua
R. Relative Validity of the 10-Year Cardiovascular Risk Estimate in a Population Cohort
of the REGICOR Study. *Revista Española de Cardiología (English Edition).*
2011;64(5):385-94.
5.   Patrono C. Fighting residual cardiovascular risk in stable patients with atherosclerotic
vascular disease: COMPASS in context. *Cardiovascular research.* 2017;113(14):e61-3.
6.   Petronis A. Epigenetics as a unifying principle in the aetiology of complex traits and
diseases. *Nature.* 2010;465(7299):721-7.
7.   Jin Z, Liu Y. DNA methylation in human diseases. *Genes & Diseases.* 2018;5(1):1–8.
8.   Agha G, Mendelson MM, Ward-Caviness CK, et al. Blood Leukocyte DNA Methylation
Predicts Risk of Future Myocardial Infarction and Coronary Heart Disease. *Circulation.*

2019;140(8):645-57.
9.   Ward-Caviness CK, Agha G, Chen BH, et al. Analysis of repeated leukocyte DNA
methylation assessments reveals persistent epigenetic alterations after an incident
myocardial infarction. *Clinical Epigenetics*. 2018;10(1):161.
10. Westerman K, Sebastiani P, Jacques P, Liu S, Demeo D, Ordovás JM. DNA methylation
modules associate with incident cardiovascular disease and cumulative risk factor
exposure. *Clinical Epigenetics*. 2019;11(1):142.
11. Westerman K, Fernández-Sanlés A, Patil P, Sebastiani P, Jacques P, Starr JM, J. Deary
I, Liu Q, Liu S, Elosua R, DeMeo DL, Ordovás JM. Epigenomic Assessment of
Cardiovascular Disease Risk and Interactions With Traditional Risk Metrics. *Journal of
the American Heart Association*. 2020;9(8):e015299.
12. Fernández-Sanlés A, Sayols-Baixeras S, Subirana I, Degano IRIR, Elosua R.
Association between DNA methylation and coronary heart disease or other
atherosclerotic events: A systematic review. *Atherosclerosis*. 2017;263:325-33.
13. Fernández-Sanlés A, Sayols-Baixeras S, Curcio S, Subirana I, Marrugat J, Elosua R.
DNA methylation and age-Independent cardiovascular risk, an epigenome-Wide
approach the REGICOR study (REgistre GIroní del COR). *Arteriosclerosis, Thrombosis,
and Vascular Biology*. 2018;38(3):645-52.
14. Sandoval J, Heyn H, Moran S, Serra-Musach J, Pujana MA, Bibikova M, Esteller M.
Validation of a DNA methylation microarray for 450,000 CpG sites in the human
genome. *Epigenetics*. 2011;6(6):692–702.
15. Moran S, Arribas C, Esteller M. Validation of a DNA methylation microarray for 850,000
CpG sites of the human genome enriched in enhancer sequences. *Epigenomics*.
2016;8(3):389-99.
16. Sayols-Baixeras S, Lluís-Ganella C, Subirana I, et al. Identification of a new locus and
validation of previously reported loci showing differential methylation associated with
smoking. The REGICOR study. *Epigenetics*. 2015;10(12):1156-65.
17. Du P, Zhang X, Huang C-C, Jafari N, Kibbe WA, Hou L, Lin SM. Comparison of Beta-
value and M-value methods for quantifying methylation levels by microarray analysis.
BMC Bioinformatics. 2010;11(1):587.
18. Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH,
Wiencke JK, Kelsey KT. DNA methylation arrays as surrogate measures of cell mixture
distribution. *BMC bioinformatics*. 2012;13(1):86.
19. Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing
batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics. 2012;28(6):882-3.
20. van Iterson M, van Zwet EW, BIOS Consortium BT, Heijmans BT. Controlling bias and
inflation in epigenome- and transcriptome-wide association studies using the empirical
null distribution. *Genome biology*. 2017;18(1):19.
21. D'Agostino RB, Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, Kannel WB.
General Cardiovascular Risk Profile for Use in Primary Care. *Circulation*. 2008;117(6).
22. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports systematic
causal inference across the human phenome. *eLife*. 2018;7:e34408.
23. Gaunt TR, Shihab HA, Hemani G, Min JL, Woodward G, Lyttleton O, Zheng J, Duggirala
A, McArdle WL, Ho K, Ring SM, Evans DM, Davey Smith G, Relton CL. Systematic
identification of genetic influences on methylation across the human life course. *Genome
biology*. 2016;17(1):61.
24. Nikpay M, Goel A, Won H-H, et al. A comprehensive 1000 Genomes–based genome-
wide association meta-analysis of coronary artery disease. *Nature Genetics*.
2015;47(10):1121–1130.
25. Shenker NS, Polidoro S, van Veldhoven K, Sacerdote C, Ricceri F, Birrell MA, Belvisi
MG, Brown R, Vineis P, Flanagan JM. Epigenome-wide association study in the
European Prospective Investigation into Cancer and Nutrition (EPIC-Turin) identifies
novel genetic loci associated with smoking. *Human Molecular Genetics*.
2013;22(5):843–851.

26. Gao X, Jia M, Zhang Y, Breitling LP, Brenner H. DNA methylation changes of whole
blood cells in response to active smoking exposure in adults: a systematic review of
DNA methylation studies. *Clinical Epigenetics*. 2015;7(1):113.
27. Deshmukh FK, Yaffe D, Olshina MA, Ben-Nissan G, Sharon M. The contribution of the
20s proteasome to proteostasis. *Biomolecules*. 2019;9(5).
28. Wang H, Jiang M, Zhu H, Chen Q, Gong P, Lin J, Lu J, Qiu J. Quantitative assessment
of the influence of *PSMA6* variant (rs1048990) on coronary artery disease risk.
Molecular Biology Reports. 2013;40(2):1035-41.
29. Heckman MG, Soto-Ortolaza AI, Diehl NN, Rayaprolu S, Brott TG, Wszolek ZK, Meschia
JF, Ross OA. Genetic variants associated with myocardial infarction in the *PSMA6* gene
and Chr9p21 are also associated with ischaemic stroke. *European Journal of Neurology*.
2013;20(2):300–308.
30. Liu X, Wang X, Shen Y, Wu L, Ruan X, Lindpaintner K, Yusuf S, Engert JC, Anand S,
Tan X, Liu L. The functional variant rs1048990 in PSMA6 is associated with
susceptibility to myocardial infarction in a Chinese population. *Atherosclerosis*.
2009;206(1):199–203.
31. Ozaki K, Sato H, Iida A, Mizuno H, Nakamura T, Miyamoto Y, Takahashi A, Tsunoda T,
Ikegawa S, Kamatani N, Hori M, Nakamura Y, Tanaka T. A functional SNP in *PSMA6*
confers risk of myocardial infarction in the Japanese population. *Nature Genetics*.
2006;38(8):921–925.
32. Pidsley R, Zotenko E, Peters TJ, Lawrence MG, Risbridger GP, Molloy P, Van Djik S,
Muhlhausler B, Stirzaker C, Clark SJ. Critical evaluation of the Illumina MethylationEPIC
BeadChip microarray for whole-genome DNA methylation profiling. *Genome biology*.
2016;17(1):208.
33. McCartney DL, Walker RM, Morris SW, Mcintosh AM, Porteous DJ, Evans KL.
Identification of polymorphic and off-target probe binding sites on the Illumina Infinium
MethylationEPIC BeadChip. *Genom Data*. 2016;9:22-4.
34. Svane A, Soerensen M, Lund J, Tan Q, Jylhävä J, Wang Y, Pedersen N, Hägg S,
Debrabant B, Deary I, Christensen K, Christiansen L, Hjelmborg J. DNA Methylation and
All-Cause Mortality in Middle-Aged and Elderly Danish Twins. *Genes*. 2018;9(2):78.
35. Ligthart S, Marzi C, Aslibekyan S, et al. DNA methylation signatures of chronic low-
grade inflammation are associated with complex diseases. *Genome biology*.
2016;17(1):255.
36. Teng N, Maghzal GJ, Talib J, Rashid I, Lau AK, Stocker R. The roles of
myeloperoxidase in coronary artery disease and its potential implication in plaque
rupture. *Redox Report*. 2017;22(2):51–73.
37. Yao C, Chen G, Song C, et al. Genome-wide mapping of plasma protein QTLs identifies
putatively causal genes and pathways for cardiovascular disease. *Nature
Communications*. 2018;9(1):3268.
38. Chen Y-C, Chen T-W, Su M-C, et al. Whole Genome DNA Methylation Analysis of
Obstructive Sleep Apnea: IL1R2, NPR2 , AR , SP140 Methylation and Clinical
Phenotype. *Sleep*. 2016;39(4):743–755.
39. Hlatky MA, Greenland P, Arnett DK, et al. Criteria for Evaluation of Novel Markers of
Cardiovascular Risk. *Circulation*. 2009;119(17):2408-16.

558
559
560
HIGHLIGHTS
- 34 novel loci differentially methylated in relation to acute myocardial infarction were
identified.
- Four of those methylation loci were further associated with incident cases of coronary
heart disease.
- The identified loci highlight the relevance of traditional cardiovascular risk factors and
biological mechanisms such as lipid metabolism and inflammation underlying
myocardial infarction.

TABLES

572 **Table 1. Descriptive characteristics of the populations used in the two-stage EWAS on**
573 **acute myocardial infarction (AMI): REGICOR-1 and REGICOR-2.**

Variables		Discovery: REGICOR-1		Validation: REGICOR-2	
		N=391	NA‡	N=204	NA‡
AMI cases/controls, n (%)‡					
	AMI cases	*196 (50.1)*	0	*101 (49.5)*	0
	Controls	*195 (49.9)*	0	*103 (50.5)*	0
Age		63.2 (6.94)	0	61.7 (6.90)	0
	AMI cases	*63.0 (6.96)*	0	*61.6 (6.83)*	0
	Controls	*63.3 (6.94)*	0	*61.7 (7.01)*	0
Sex, female, n (%)		201 (51.4)	0	100 (49.0)	0
	AMI cases	*100 (51.0)*	0	*49 (48.5)*	0
	Controls	*101 (51.8)*	0	*51 (49.5)*	0
Smokers, n (%)		93 (24.4)	10	67 (33.3)	3
	AMI cases	*76 (40.6)*	9	*45 (45.0)*	1
	Controls	*17 (8.76)*	1	*22 (21.8)*	2
BMI, kg/m2*‡		28.5 (4.75)	52	27.5 (4.87)	23
	AMI cases	*28.1 (4.34)*	52	*28.2 (5.87)*	23
	Controls	*28.8 (5.02)*	0	*26.9 (3.90)*	0
Hypercholesterolaemia, n (%)‡		179 (53.0)	53	93 (51.4)	23
	AMI cases	*93 (64.1)*	51	*51 (65.4)*	23
	Controls	*86 (44.6)*	2	*42 (40.8)*	0
HTN, n (%)‡		212 (57.1)	20	105 (54.1)	10
	AMI cases	*120 (68.2)*	20	*62 (68.1)*	10
	Controls	*92 (47.2)*	0	*43 (41.7)*	0
Diabetes, n (%)		87 (24.7)	39	44 (23.8)	19
	AMI cases	*56 (35.7)*	39	*27 (32.9)*	19
	Controls	*31 (15.9)*	0	*17 (16.5)*	0

*Mean (Standard deviation).
†Median (Interquartile Range).
‡ NA, missing information; AMI, Acute myocardial infarction; BMI, Body mass index;
Hypercholesterolaemia, defined as self-reported high cholesterol levels or treatment; HTN,
Hypertension, defined as self-reported high blood pressure or treatment; Diabetes, defined
as self-reported diabetes or treatment.
580

Table 2. Descriptive characteristics of the populations used in the follow-up association studies on incident cardiovascular (CVD) and coronary heart disease (CHD) events: Women's Health Initiative (WHI) and Framingham Offspring Study (FOS).

Variables	WHI N=1863	NA	FOS CHD N=2266	NA	FOS CVD N=2123	NA
Age	64.2 (7.03)	0	65.7 (8.82)	0	65.3 (8.66)	0
Sex, female, n (%)	1863 (100)	0	1284 (56.7)	0	1206 (56.8)	0
CHD, n (%)‡	914 (49.1)	0	106 (4.68)	0	_	_
CVD, n (%)‡	984 (52.8)	0	_	_	222 (10.5)	0
Smokers, n (%)	174 (9.46)	24	217 (9.62)	11	202 (9.56)	11
BMI, kg/m2*‡	29.8 (5.97)	11	28.1 (5.39)	8	28.0 (5.33)	6
Total cholesterol, mg/dl*	233 (42.3)	1	189 (36.2)	1	190 (35.8)	1
LDL-C, mg/dl*‡	152 (38.7)	31	107 (30.8)	2	108 (30.5)	2
HDL-C, mg/dl*‡	52.2 (13.2)	1	58.3 (18.3)	2	58.7 (18.3)	2
TG, mg/dl†	126 [93.0;176]	1	101 [73.0;140]	1	100 [73.0;139]	1
Cholesterol treatment, n (%)	166 (11.4)	403	871 (38.5)	3	787 (37.1)	3
Hypercholesterolaemia, n (%)‡	913 (55.8%)	227	1033 (45.7)	4	943 (44.5)	4
SBP, mmHg*‡	131 (17.5)	0	126 (16.9)	1	125 (16.7)	1
DBP, mmHg*‡	76.4 (9.27)	0	72.1 (9.99)	3	72.3 (9.92)	3
HTN, n (%)‡	952 (61.2)	308	1217 (53.9)	8	1110 (52.5)	7
HTN treatment, n (%)‡	672 (45.9)	400	1025 (45.4)	7	927 (43.8)	6
Glucose, mg/dl†	107 (38.3)	1	106 (21.8)	2	105 (21.5)	2
Diabetes, n (%)	319 (17.1)	0	307 (14.3)	126	266 (13.3)	119
Diabetes treatment, n (%)	178 (9.57)	3	87 (4.04)	110	72 (3.56)	102

*Mean (Standard deviation). †Median (Interquartile range). ‡ CHD, Coronary heart disease; CVD, Cardiovascular disease; BMI, Body mass index; LDL-C, Cholesterol in low-density lipoprotein; HDL-C, Cholesterol in high-density lipoprotein; TG, Triglycerides; Hypercholesterolaemia, defined as previous treatment or total cholesterol ≥240 mmHg or LDL-C ≥160 mmHg; SBP, Systolic blood pressure; DBP, Diastolic blood pressure; HTN, Hypertension, defined as previous treatment or SBP ≥140 mmHg or DBP ≥90 mmHg; Diabetes, defined as previous treatment or glycaemia ≥126mg/dl.

Table 3. Significant CpGs differentially methylated in association with prevalent myocardial infarction in the fixed-effects meta-analyses of the REGICOR case-control samples. Model 1 was adjusted for estimated cell counts and two surrogate variables. Model 2 was further adjusted for smoking status. Model 3 was additionally adjusted for diabetes, hypercholesterolemia, and hypertension. The Bonferroni-corrected p threshold was $1.67 \cdot 10^{-8}$.

	Genomic information			Model 1			Model 2			Model 3		
CpG	Chr*	Position (bp)	Gene	Coef*	SE*	*P*	Coef*	SE*	*P*	Coef*	SE*	*P*
A) CpGs significant in both the meta-analysis and the validation stage												
cg10636246	1	159,046,973	AIM2	$-9.62 \cdot 10^{-1}$	$1.41 \cdot 10^{-1}$	$1.03 \cdot 10^{-11}$	$-9.82 \cdot 10^{-1}$	$1.58 \cdot 10^{-1}$	$5.21 \cdot 10^{-10}$	$-8.56 \cdot 10^{-1}$	$1.68 \cdot 10^{-1}$	$3.68 \cdot 10^{-7}$
cg08345526	1	226,916,320	ITPKB	1.79	$2.93 \cdot 10^{-1}$	$1.02 \cdot 10^{-9}$	1.85	$3.29 \cdot 10^{-1}$	$1.71 \cdot 10^{-8}$	1.60	$3.45 \cdot 10^{-1}$	$3.42 \cdot 10^{-6}$
cg01552966	2	25,528,330	DNMT3A	1.15	$1.93 \cdot 10^{-1}$	$2.43 \cdot 10^{-9}$	1.25	$2.22 \cdot 10^{-1}$	$1.78 \cdot 10^{-8}$	1.04	$2.33 \cdot 10^{-1}$	$7.66 \cdot 10^{-6}$
cg25420477	2	70,319,121	Intergenic	1.54	$2.52 \cdot 10^{-1}$	$1.08 \cdot 10^{-9}$	1.63	$2.84 \cdot 10^{-1}$	$9.01 \cdot 10^{-9}$	1.47	$3.01 \cdot 10^{-1}$	$9.45 \cdot 10^{-7}$
cg06528771	2	102,513,985	Intergenic	-1.71	$2.79 \cdot 10^{-1}$	$9.88 \cdot 10^{-10}$	-1.89	$3.16 \cdot 10^{-1}$	$2.26 \cdot 10^{-9}$	-1.81	$3.48 \cdot 10^{-1}$	$1.81 \cdot 10^{-7}$
cg12206840	2	136,826,451	Intergenic	1.12	$1.98 \cdot 10^{-1}$	$1.31 \cdot 10^{-8}$	1.12	$2.23 \cdot 10^{-1}$	$4.89 \cdot 10^{-7}$	1.03	$2.33 \cdot 10^{-1}$	$9.08 \cdot 10^{-6}$
cg10650214	3	45,914,188	LZTFL1	1.28	$1.97 \cdot 10^{-1}$	$8.01 \cdot 10^{-11}$	1.26	$2.19 \cdot 10^{-1}$	$8.05 \cdot 10^{-9}$	1.18	$2.31 \cdot 10^{-1}$	$2.99 \cdot 10^{-7}$
cg13373048	3	71,892,575	Intergenic	-2.76	$3.57 \cdot 10^{-1}$	$1.02 \cdot 10^{-14}$	-3.24	$4.23 \cdot 10^{-1}$	$1.93 \cdot 10^{-14}$	-3.03	$4.29 \cdot 10^{-1}$	$1.77 \cdot 10^{-12}$
cg03733470	3	196,613,114	SENP5	-1.20	$2.07 \cdot 10^{-1}$	$5.94 \cdot 10^{-9}$	-1.28	$2.30 \cdot 10^{-1}$	$2.70 \cdot 10^{-8}$	-1.17	$2.46 \cdot 10^{-1}$	$2.01 \cdot 10^{-6}$
cg25769469	5	71,643,841	PTCD2	-1.33	$2.20 \cdot 10^{-1}$	$1.33 \cdot 10^{-9}$	-1.38	$2.53 \cdot 10^{-1}$	$5.40 \cdot 10^{-8}$	-1.41	$2.80 \cdot 10^{-1}$	$4.92 \cdot 10^{-7}$
cg12066594	9	127,162,066	PSMB7	-1.89	$3.37 \cdot 10^{-1}$	$2.00 \cdot 10^{-8}$	-2.57	$4.07 \cdot 10^{-1}$	$2.67 \cdot 10^{-10}$	-2.36	$4.25 \cdot 10^{-1}$	$2.83 \cdot 10^{-8}$
cg24501073	11	34,210,305	ABTB2	$9.89 \cdot 10^{-1}$	$1.73 \cdot 10^{-1}$	$1.07 \cdot 10^{-8}$	$9.76 \cdot 10^{-1}$	$1.96 \cdot 10^{-1}$	$6.11 \cdot 10^{-7}$	$8.23 \cdot 10^{-1}$	$1.98 \cdot 10^{-1}$	$3.07 \cdot 10^{-5}$
cg02042026	11	113,936,782	ZBTB16	1.04	$1.70 \cdot 10^{-1}$	$8.50 \cdot 10^{-10}$	$9.88 \cdot 10^{-1}$	$1.88 \cdot 10^{-1}$	$1.51 \cdot 10^{-7}$	$9.82 \cdot 10^{-1}$	$2.04 \cdot 10^{-1}$	$1.50 \cdot 10^{-6}$
cg15867698	14	69,438,267	ACTN1	1.13	$1.86 \cdot 10^{-1}$	$1.58 \cdot 10^{-9}$	1.26	$2.14 \cdot 10^{-1}$	$3.33 \cdot 10^{-9}$	1.14	$2.25 \cdot 10^{-1}$	$4.04 \cdot 10^{-7}$
cg07252680	14	94,857,224	SERPINA1	-1.56	$2.41 \cdot 10^{-1}$	$1.17 \cdot 10^{-10}$	-1.43	$2.65 \cdot 10^{-1}$	$6.57 \cdot 10^{-8}$	-1.41	$2.86 \cdot 10^{-1}$	$7.93 \cdot 10^{-7}$
cg09967176	15	77,280,925	Intergenic	-1.28	$1.78 \cdot 10^{-1}$	$6.93 \cdot 10^{-13}$	-1.40	$2.03 \cdot 10^{-1}$	$5.21 \cdot 10^{-12}$	-1.16	$2.10 \cdot 10^{-1}$	$3.82 \cdot 10^{-8}$
cg17261469	16	87,420,879	FBXO31	$9.93 \cdot 10^{-1}$	$1.73 \cdot 10^{-1}$	$9.49 \cdot 10^{-9}$	$9.96 \cdot 10^{-1}$	$1.94 \cdot 10^{-1}$	$3.01 \cdot 10^{-7}$	$9.98 \cdot 10^{-1}$	$2.04 \cdot 10^{-1}$	$9.92 \cdot 10^{-7}$
cg22146772	17	26,793,673	Intergenic	1.23	$1.97 \cdot 10^{-1}$	$5.38 \cdot 10^{-10}$	1.18	$2.22 \cdot 10^{-1}$	$8.98 \cdot 10^{-8}$	1.06	$2.28 \cdot 10^{-1}$	$3.31 \cdot 10^{-6}$
cg04988978	17	56,359,578	MPO	-1.31	$2.07 \cdot 10^{-1}$	$2.33 \cdot 10^{-10}$	-1.45	$2.37 \cdot 10^{-1}$	$9.71 \cdot 10^{-10}$	-1.22	$2.43 \cdot 10^{-1}$	$5.41 \cdot 10^{-7}$
cg19448292	20	35,504,064	C20orf118	1.24	$1.93 \cdot 10^{-1}$	$1.17 \cdot 10^{-10}$	1.38	$2.25 \cdot 10^{-1}$	$7.89 \cdot 10^{-10}$	1.15	$2.30 \cdot 10^{-1}$	$6.05 \cdot 10^{-7}$
B) CpGs significant in the meta-analysis†												
cg15724772	1	6,373,534	ACOT7	1.17	$1.94 \cdot 10^{-1}$	$1.76 \cdot 10^{-9}$	1.16	$2.17 \cdot 10^{-1}$	$8.14 \cdot 10^{-8}$	1.00	$2.30 \cdot 10^{-1}$	$1.38 \cdot 10^{-5}$

CpG	Chr	Position	Gene	Coef	SE	p	Coef	SE	p	Coef	SE	p
cg11919692	1	26,343,868	Intergenic	1.08	$1.98 \cdot 10^{-1}$	$4.94 \cdot 10^{-8}$	1.11	$2.19 \cdot 10^{-1}$	$4.00 \cdot 10^{-7}$	1.04	$2.34 \cdot 10^{-1}$	$9.29 \cdot 10^{-6}$
cg21566642	2	233,284,661	Intergenic	$-6.53 \cdot 10^{-1}$	$1.13 \cdot 10^{-1}$	$7.70 \cdot 10^{-9}$	$-3.03 \cdot 10^{-1}$	$1.56 \cdot 10^{-1}$	$5.26 \cdot 10^{-2}$	$-4.14 \cdot 10^{-1}$	$1.70 \cdot 10^{-1}$	$1.50 \cdot 10^{-2}$
cg07386061	3	52,492,874	*NISCH*	$9.94 \cdot 10^{-1}$	$1.79 \cdot 10^{-1}$	$2.86 \cdot 10^{-8}$	1.03	$2.01 \cdot 10^{-1}$	$3.27 \cdot 10^{-7}$	$8.58 \cdot 10^{-1}$	$2.09 \cdot 10^{-1}$	$4.01 \cdot 10^{-5}$
cg05575921	5	373,378	*AHRR*	$-7.51 \cdot 10^{-1}$	$1.25 \cdot 10^{-1}$	$2.05 \cdot 10^{-9}$	$-4.04 \cdot 10^{-1}$	$1.91 \cdot 10^{-1}$	$3.48 \cdot 10^{-2}$	$-4.46 \cdot 10^{-1}$	$2.04 \cdot 10^{-1}$	$2.87 \cdot 10^{-2}$
cg04725636	8	67,004,538	*DNAJC5B*	$-7.99 \cdot 10^{-1}$	$1.45 \cdot 10^{-1}$	$3.33 \cdot 10^{-8}$	$-7.21 \cdot 10^{-1}$	$1.59 \cdot 10^{-1}$	$5.47 \cdot 10^{-6}$	$-7.46 \cdot 10^{-1}$	$1.74 \cdot 10^{-1}$	$1.76 \cdot 10^{-5}$
cg01381203	12	2,985,630	*FOXM1;RHNO1*	$-8.83 \cdot 10^{-1}$	$1.62 \cdot 10^{-1}$	$5.17 \cdot 10^{-8}$	$-8.39 \cdot 10^{-1}$	$1.76 \cdot 10^{-1}$	$1.88 \cdot 10^{-6}$	$-9.17 \cdot 10^{-1}$	$1.95 \cdot 10^{-1}$	$2.44 \cdot 10^{-6}$
cg02175213	12	69,352,407	*CPM*	-1.08	$1.85 \cdot 10^{-1}$	$5.15 \cdot 10^{-9}$	-1.10	$2.06 \cdot 10^{-1}$	$9.76 \cdot 10^{-8}$	$-8.20 \cdot 10^{-1}$	$2.14 \cdot 10^{-1}$	$1.27 \cdot 10^{-4}$
cg06084585	13	50,797,125	*DLEU1*	-1.60	$2.87 \cdot 10^{-1}$	$2.37 \cdot 10^{-8}$	-1.47	$3.14 \cdot 10^{-1}$	$2.96 \cdot 10^{-6}$	-1.36	$3.30 \cdot 10^{-1}$	$3.54 \cdot 10^{-5}$
cg16125725	15	70,101,302	Intergenic	-2.03	$3.45 \cdot 10^{-1}$	$4.02 \cdot 10^{-9}$	-2.30	$3.97 \cdot 10^{-1}$	$6.82 \cdot 10^{-9}$	-2.20	$4.18 \cdot 10^{-1}$	$1.39 \cdot 10^{-7}$
cg04983687	16	88,558,223	*ZFPM1*	$7.21 \cdot 10^{-1}$	$1.29 \cdot 10^{-1}$	$2.61 \cdot 10^{-8}$	$6.77 \cdot 10^{-1}$	$1.42 \cdot 10^{-1}$	$1.79 \cdot 10^{-6}$	$4.42 \cdot 10^{-1}$	$1.46 \cdot 10^{-1}$	$2.49 \cdot 10^{-3}$
cg22508829	17	17,843,585	*TOM1L2*	1.28	$1.96 \cdot 10^{-1}$	$7.97 \cdot 10^{-11}$	1.35	$2.26 \cdot 10^{-1}$	$2.41 \cdot 10^{-9}$	1.10	$2.28 \cdot 10^{-1}$	$1.23 \cdot 10^{-6}$
cg13064873	17	48,360,726	*TMEM92-AS1*	-1.08	$1.85 \cdot 10^{-1}$	$5.31 \cdot 10^{-9}$	-1.19	$2.12 \cdot 10^{-1}$	$1.82 \cdot 10^{-8}$	-1.04	$2.23 \cdot 10^{-1}$	$3.19 \cdot 10^{-6}$
cg03636183	19	17,000,585	*F2RL3*	$-7.44 \cdot 10^{-1}$	$1.36 \cdot 10^{-1}$	$4.56 \cdot 10^{-8}$	$-3.50 \cdot 10^{-1}$	$1.80 \cdot 10^{-1}$	$5.24 \cdot 10^{-2}$	$-3.88 \cdot 10^{-1}$	$1.90 \cdot 10^{-1}$	$4.12 \cdot 10^{-2}$

596 **Chr, Chromosome; Coef; Coefficient of association; SE, Standard error.*

597 *†These CpGs should be interpreted with caution, as they were not significant in the validation stage (Supplementary Table 2).*

Table 4. Significant CpGs differentially methylated in association with incident cardiovascular/coronary disease in the fixed-effects meta-analyses of the samples from the Women's Health Initiative (WHI) and the Framingham Offspring Study (FOS). Model 1 was adjusted for age, estimated cell counts and two surrogate variables (plus ancestry in WHI, plus sex in FOS). Model 2 was further adjusted for smoking status. Model 3 was additionally adjusted for diabetes, hypercholesterolemia, and hypertension. The Bonferroni-corrected p threshold was $4.17 \cdot 10^{-3}$ (those significant are highlighted in bold).

Genomic information				Model 1			Model 2			Model 3		
CpG	Chr*	Position (bp)	Gene	Coef*	SE*	*P*	Coef*	SE*	*P*	Coef*	SE*	*P*
A) *Incident coronary heart disease*												
cg21566642	2	233,284,661	Intergenic	**$-1.34 \cdot 10^{-1}$**	**$4.33 \cdot 10^{-2}$**	**$2.00 \cdot 10^{-3}$**	$-9.63 \cdot 10^{-2}$	$4.83 \cdot 10^{-2}$	$4.62 \cdot 10^{-2}$	$-9.66 \cdot 10^{-2}$	$5.39 \cdot 10^{-2}$	$7.32 \cdot 10^{-2}$
cg05575921	5	373,378	*AHRR*	**$-1.80 \cdot 10^{-1}$**	**$4.73 \cdot 10^{-2}$**	**$1.38 \cdot 10^{-4}$**	$-1.33 \cdot 10^{-1}$	$5.81 \cdot 10^{-2}$	$2.21 \cdot 10^{-2}$	$-1.22 \cdot 10^{-1}$	$6.52 \cdot 10^{-2}$	$6.05 \cdot 10^{-2}$
cg25769469	5	71,643,841	*PTCD2*	**$-2.62 \cdot 10^{-1}$**	**$7.94 \cdot 10^{-2}$**	**$9.67 \cdot 10^{-4}$**	**$-2.52 \cdot 10^{-1}$**	**$8.02 \cdot 10^{-2}$**	**$1.70 \cdot 10^{-3}$**	$-2.14 \cdot 10^{-1}$	$8.89 \cdot 10^{-2}$	$1.59 \cdot 10^{-2}$
cg04988978	17	56,359,578	*MPO*	**$-2.28 \cdot 10^{-1}$**	**$7.49 \cdot 10^{-2}$**	**$2.29 \cdot 10^{-3}$**	$-2.07 \cdot 10^{-1}$	$7.58 \cdot 10^{-2}$	$6.19 \cdot 10^{-3}$	$-1.06 \cdot 10^{-1}$	$8.35 \cdot 10^{-2}$	$2.03 \cdot 10^{-1}$
B) *Incident cardiovascular disease*												
cg21566642	2	233,284,661	Intergenic	**$-1.68 \cdot 10^{-1}$**	**$3.97 \cdot 10^{-2}$**	**$2.34 \cdot 10^{-5}$**	**$-1.31 \cdot 10^{-1}$**	**$4.40 \cdot 10^{-2}$**	**$2.94 \cdot 10^{-3}$**	$-1.37 \cdot 10^{-1}$	$4.85 \cdot 10^{-2}$	$4.87 \cdot 10^{-3}$
cg05575921	5	373,378	*AHRR*	**$-2.08 \cdot 10^{-1}$**	**$4.30 \cdot 10^{-2}$**	**$1.30 \cdot 10^{-6}$**	**$-1.67 \cdot 10^{-1}$**	**$5.24 \cdot 10^{-2}$**	**$1.42 \cdot 10^{-3}$**	$-1.61 \cdot 10^{-1}$	$5.79 \cdot 10^{-2}$	$5.54 \cdot 10^{-3}$
cg25769469	5	71,643,841	*PTCD2*	$-1.92 \cdot 10^{-1}$	$7.27 \cdot 10^{-2}$	$8.40 \cdot 10^{-3}$	$-1.77 \cdot 10^{-1}$	$7.35 \cdot 10^{-2}$	$1.58 \cdot 10^{-2}$	$-1.26 \cdot 10^{-1}$	$8.09 \cdot 10^{-2}$	$1.20 \cdot 10^{-1}$
cg04988978	17	56,359,578	*MPO*	**$-2.04 \cdot 10^{-1}$**	**$6.72 \cdot 10^{-2}$**	**$2.42 \cdot 10^{-3}$**	$-1.83 \cdot 10^{-1}$	$6.79 \cdot 10^{-2}$	$6.99 \cdot 10^{-3}$	$-9.08 \cdot 10^{-2}$	$7.36 \cdot 10^{-2}$	$2.18 \cdot 10^{-1}$

*Chr, Chromosome; Coef; Coefficient of association; SE, Standard error.

Table 5. Significant associations between the identified CpGs and classical cardiovascular risk factors (CVRFs) in the fixed-effects meta-analyses of the four samples. The significant associations are highlighted in bold ($p < 1.56 \cdot 10^{-3}$ †).

CpG	Smoking	BMI*	HDL-C*	LDL-C*	TG*	Glucose	SBP	DBP*	
cg21566642	-1.61	$9.55 \cdot 10^{-3}$	$7.69 \cdot 10^{-4}$	$-1.10 \cdot 10^{-3}$	$-4.17 \cdot 10^{-4}$	$2.32 \cdot 10^{-4}$	$2.30 \cdot 10^{-3}$	$7.97 \cdot 10^{-3}$	**Coefficient**
	$4.22 \cdot 10^{-2}$	$2.65 \cdot 10^{-3}$	$1.21 \cdot 10^{-3}$	$4.09 \cdot 10^{-4}$	$1.95 \cdot 10^{-4}$	$4.20 \cdot 10^{-4}$	$9.02 \cdot 10^{-4}$	$1.67 \cdot 10^{-3}$	**SE***
	$<2.2 \cdot 10^{-16}$	**$3.21 \cdot 10^{-4}$**	$5.24 \cdot 10^{-1}$	$7.11 \cdot 10^{-3}$	$3.22 \cdot 10^{-2}$	$5.81 \cdot 10^{-1}$	$1.08 \cdot 10^{-2}$	**$1.78 \cdot 10^{-6}$**	***P***
cg05575921	-1.61	$1.17 \cdot 10^{-2}$	$4.36 \cdot 10^{-3}$	$-1.35 \cdot 10^{-3}$	$-7.09 \cdot 10^{-4}$	$1.86 \cdot 10^{-4}$	$2.15 \cdot 10^{-3}$	$7.54 \cdot 10^{-3}$	**Coefficient**
	$4.22 \cdot 10^{-2}$	$2.46 \cdot 10^{-3}$	$1.11 \cdot 10^{-3}$	$3.77 \cdot 10^{-4}$	$1.79 \cdot 10^{-4}$	$3.87 \cdot 10^{-4}$	$8.23 \cdot 10^{-4}$	$1.52 \cdot 10^{-3}$	**SE***
	$<2.2 \cdot 10^{-16}$	**$2.22 \cdot 10^{-6}$**	**$8.22 \cdot 10^{-5}$**	**$3.39 \cdot 10^{-4}$**	**$7.63 \cdot 10^{-5}$**	$6.30 \cdot 10^{-1}$	$9.06 \cdot 10^{-3}$	**$7.69 \cdot 10^{-7}$**	***P***
cg25769469	$-9.01 \cdot 10^{-2}$	$-1.12 \cdot 10^{-2}$	$6.76 \cdot 10^{-3}$	$2.47 \cdot 10^{-4}$	$-1.38 \cdot 10^{-3}$	$-1.52 \cdot 10^{-3}$	$-2.35 \cdot 10^{-3}$	$-4.17 \cdot 10^{-3}$	**Coefficient**
	$5.08 \cdot 10^{-2}$	$2.64 \cdot 10^{-3}$	$1.20 \cdot 10^{-3}$	$4.11 \cdot 10^{-4}$	$1.93 \cdot 10^{-4}$	$4.17 \cdot 10^{-4}$	$9.10 \cdot 10^{-4}$	$1.69 \cdot 10^{-3}$	**SE***
	$7.65 \cdot 10^{-2}$	**$2.32 \cdot 10^{-5}$**	**$1.92 \cdot 10^{-8}$**	$5.48 \cdot 10^{-1}$	**$8.75 \cdot 10^{-13}$**	**$2.69 \cdot 10^{-4}$**	$9.97 \cdot 10^{-3}$	$1.36 \cdot 10^{-2}$	***P***
cg04988978	$-7.29 \cdot 10^{-2}$	$-1.15 \cdot 10^{-2}$	$5.28 \cdot 10^{-3}$	$3.59 \cdot 10^{-4}$	$-1.19 \cdot 10^{-3}$	$-1.88 \cdot 10^{-3}$	$-2.88 \cdot 10^{-3}$	$-3.54 \cdot 10^{-3}$	**Coefficient**
	$5.15 \cdot 10^{-2}$	$2.64 \cdot 10^{-3}$	$1.20 \cdot 10^{-3}$	$4.08 \cdot 10^{-4}$	$1.92 \cdot 10^{-4}$	$4.15 \cdot 10^{-4}$	$9.02 \cdot 10^{-4}$	$1.67 \cdot 10^{-3}$	**SE***
	$1.57 \cdot 10^{-1}$	**$1.26 \cdot 10^{-5}$**	**$1.08 \cdot 10^{-5}$**	$3.80 \cdot 10^{-1}$	**$5.79 \cdot 10^{-10}$**	**$5.97 \cdot 10^{-6}$**	**$1.41 \cdot 10^{-3}$**	$3.47 \cdot 10^{-2}$	***P***

*BMI, Body mass index; HDL-C, Cholesterol in high-density lipoprotein; LDL-C, Cholesterol in low-density lipoprotein; TG, Triglycerides; SBP, Systolic blood pressure; DBP, Diastolic blood pressure; SE, Standard Error.

†0.05/(number of CpGs*8 CVRFs)