Apple ripening is controlled by a NAC transcription factor

Running title: NAC18.1 controls apple ripening

Zoë Migicovsky[1], Trevor H. Yeats[2,6], Sophie Watts[1], Jun Song[3], Charles F. Forney[3], Karen Burgher-MacLellan[3], Daryl J. Somers[4], Yihi Gong[5,#], Zhaoqi Zhang[5], Julia Vrebalov[2,6], James G. Giovannoni[6,7], Jocelyn K. C. Rose[2]*, Sean Myles[1]*

1. Department of Plant, Food and Environmental Sciences, Dalhousie University, Faculty of Agriculture, Agricultural Campus, Truro, NS, Canada.

2. Plant Biology Section, School of Integrative Plant Science, Cornell University, Ithaca, NY 14853 USA

3. Agriculture and Agri-Food Canada, Kentville, Nova Scotia, B4N 1J5, Canada

4. Vineland Research and Innovation Centre, Vineland Station, ON, L0R 2E0, Canada

5. College of Horticulture, South China Agriculture University, Guangzhou, China.

6. Boyce Thompson Institute, Cornell University, Ithaca, NY 14853 USA.

7. United States Department of Agriculture, Robert W. Holley Center, Cornell University, Ithaca, NY 14853 USA

[#]Present address: College of Agriculture and Biotechnology, Hunan University of Humanities, Science and Technology, Loudi 417000, China

* Co-corresponding authors: Sean Myles (sean.myles@dal.ca) and Jocelyn K. C. Rose (jr286@cornell.edu).

Abstract

Softening is a hallmark of ripening in fleshy fruits, and has both desirable and undesirable implications for texture and postharvest stability. Accordingly, the timing and extent of ripening and associated textural changes are key targets for improving fruit quality through breeding. Previously, we identified a large effect locus associated with harvest date and firmness in apple (*Malus domestica*) using genome-wide association studies (GWAS). Here, we present additional evidence that polymorphisms in or around a transcription factor gene, *NAC18.1*, may cause variation in these traits. First, we confirmed our previous findings with new phenotype and genotype data from ~800 apple accessions. In this population, we compared a genetic marker within *NAC18.1* to markers targeting three other firmness-related genes currently used by breeders (*ACS1*, *ACO1*, and *PG1*), and found that the *NAC18.1* marker was the strongest predictor of both firmness at harvest and firmness after three months of cold storage. By sequencing *NAC18.1* across 18 accessions, we revealed two predominant haplotypes containing the single nucleotide polymorphism (SNP) previously identified using GWAS, as well as dozens of additional SNPs and indels in both the coding and promoter sequences. *NAC18.1* encodes a protein with high similarity to the NON-RIPENING (NOR) transcription factor, a regulator of ripening in tomato (*Solanum lycopersicum*). To test whether these genes are functionally orthologous, we introduced both *NAC18.1* transgene haplotypes into the tomato *nor* mutant and showed that both haplotypes complement the *nor* ripening deficiency. Taken together, these results indicate that polymorphisms in *NAC18.1* may underlie substantial variation in apple firmness through modulation of a conserved ripening program.

Keywords:

Apple, fruit ripening, NAC-domain transcription factor

Introduction

Despite their diverse structure, ontogeny, and biochemical composition, fleshy fruits from a taxonomically broad range of species undergo coordinated ripening processes that have many features in common. Ripening involves numerous physiological and biochemical changes that render the fruit attractive and nutritious for consumption by seed-dispersing animals, or human consumers in the case of crops. These changes include the accumulation of sugars, pigments, and flavor or aroma compounds, as well as a loss of flesh firmness due in large part to the controlled modification and depolymerization of cell wall polysaccharides (Wang et al., 2018). Processes involved in ripening are regulated by conserved and convergently evolved networks of transcription factors and hormones, such as ethylene in climacteric fruit where a respiratory burst occurs at the beginning of ripening (Lü et al., 2018).

Various aspects of fruit ripening have been particularly well-studied in tomato (*Solanum lycopersicum* L.) and the characterization of tomato ripening mutants has revealed a regulatory network consisting of transcription factors, hormones, and epigenetic modifications (Giovannoni et al., 2017). Among the best studied ripening-related transcription factors in tomato is NON-RIPENING (NOR), a NAC [No apical meristem (NAM), *Arabidopsis* transcription activation factor (ATAF), Cup-shaped cotyledon (CUC)] domain transcription factor (Giovannoni et al., 2004) expressed early in ripening (Shinozaki et al., 2018). NAC genes comprise one of the largest plant-specific families of transcription factors, with specific members regulating development, defense, and senescence (Mathew and Agarwal, 2018). While all NAC genes share a conserved DNA-binding (NAC) domain, specific functional clades are defined in terms of their

more variable domains, particularly the C-terminal transcriptional regulatory region. These domains can act directly as transcriptional activators, or can facilitate interaction with other transcription factors in order to fine-tune transcriptional control. NAC transcription factors have been implicated in ripening phenotypes in diverse species including tomato (Kumar et al., 2018), melon (Ríos et al., 2017), banana (Shan et al., 2012), peach (Pirona et al., 2013) and apricot (García-Gómez et al., 2019).

Apple (*Malus* x *domestica* Borkh.) fruit exhibit extensive variation in the extent and timing of ripening and softening. In previous genome-wide association studies (GWAS) of 689 apple accessions, we found a single nucleotide polymorphism (SNP) on chromosome 3 within the coding sequence of a NAC transcription factor, *NAC18.1* (MD03G1222600 in the GDDH13 v1.1 reference genome (Daccord et al., 2017)), associated with harvest date and firmness (Migicovsky et al., 2016). This SNP results in an aspartate (D) to tyrosine (Y) mutation at a highly conserved position of the *NAC18.1* amino acid sequence and we refer to this putatively causal SNP as D5Y. Subsequently, GWAS of three additional germplasm collections confirmed the association between the genomic region containing *NAC18.1* and ripening time (Urrestarazu et al., 2017; Larsen et al., 2019; Jung et al., 2020). Thus, the *NAC18.1* gene is a strong candidate for mediating apple ripening time and firmness, and the D5Y SNP may be of utility for marker-assisted breeding.

Due to a prolonged juvenile phase, it is particularly challenging for apple breeders to evaluate fruit quality traits. As a result, in recent years considerable effort has been invested in developing molecular markers that can be used to select for fruit quality traits at the seedling stage. In particular, markers in three genes are currently used by apple breeders to select for desirable fruit texture. The first is *1-AMINOCYCLOPROPANE-1-CARBOXYLATE SYNTHASE 1*

(*ACS1*), which encodes the ripening-associated isoform of an ethylene biosynthesis gene. The *ACS1-2* allele contains a retrotransposon insertion thought to confer low ethylene production and longer shelf life in 'Fuji' and other apple cultivars homozygous for this allele (Sunako et al., 1999; Harada et al., 2000). The second gene corresponds to another enzyme involved in ethylene biosynthesis, *AMINOCYCLOPROPANE-1-CARBOXYLATE OXIDASE 1* (*ACO1*), which has a similar reduced-functionality allele (Costa et al., 2005). Finally, *POLYGALACTURONASE 1* (*PG1*) encodes an enzyme that hydrolyzes pectin polysaccharides in the cell wall and middle lamella and thus has been implicated in affecting apple fruit firmness (Atkinson et al., 2012). Several apple *PG* alleles have been identified (Costa et al. 2010). Apple firmness was identified as one of the five most important traits for genomics-assisted breeding (Laurens et al., 2018) and markers for the desirable alleles of *ACS1*, *ACO1*, and *PG1* are widely available (Baumgartner et al., 2016).

The ripening of climacteric fruits, like apples, is regulated by the plant hormone ethylene (Grierson, 2013). All three firmness-related genes described above are ethylene-dependent: their expression is mediated by ethylene after the initiation of the climacteric process and is repressed by exposure to the ethylene inhibitor 1-methylcyclopropene (1-MCP) (Brummell and Harpster, 2001; Dandekari et al., 2004; Costa et al., 2005, 2010; Tadiello et al., 2016; Di Guardo et al., 2017). While *PG1* has been associated with apple firmness using GWAS (Kumar et al., 2013; Di Guardo et al., 2017), the discovery of *ACS1*, *ACO1* and *PG1* markers was driven largely by linkage mapping studies of bi-parental families (Costa et al., 2010; Longhi et al., 2012, 2013; Bink et al., 2014; Sadok et al., 2015). These studies did not consistently detect all three loci, which may have been due to limited sample sizes, lack of segregation between parents, and/or differences in phenotyping methods. Subsequent testing has revealed either low, or no power for

these markers to predict firmness phenotypes, raising doubt about their utility in marker-assisted breeding (Costa et al., 2013; Nybom et al., 2013; McClure et al., 2018; Chagné et al., 2019).

Here we extend our previous work and report an evaluation of the role of *NAC18.1* in apple ripening and softening. First, we tested the utility of the three established firmness-related markers (*ACO1*, *ACS1* and *PG1*) and the *NAC18.1* D5Y marker to predict harvest date, firmness and softening during storage across more than 800 diverse apple accessions. Second, we determined the effect of ethylene and 1-MCP on the expression levels of all four genes. Third, we sequenced the *NAC18.1* gene from a subset of apple cultivars to discover potentially causal alleles in linkage disequilibrium (LD) with D5Y. Finally, we heterologously expressed the *NAC18.1* gene in the tomato *non-ripening (nor)* mutant to test whether it functions as a component of a conserved fruit ripening program via complementation.

Materials and Methods

Germplasm sources

Apple samples taken for this study included three different sources of germplasm. The majority of the samples were from Canada's Apple Biodiversity Collection (ABC), an orchard located in Kentville, Nova Scotia, which contains 1,113 apple accessions. A comprehensive description of the ABC is provided in Watts et al. (submitted). Briefly, the ABC is a diverse germplasm collection planted in duplicate in an incomplete block design, which includes 1 of 3 standards per grid, allowing for correction of positional effects using a REstricted Maximum Likelihood (REML) model, described in Migicovsky et al. (2017). Samples from the ABC were used for phenotyping of harvest date and firmness, as well as genotyping of texture-related genetic markers.

Additional apple samples were taken from the United States Department of Agriculture (USDA) apple germplasm collection and the Cornell apple breeding program in Geneva, New York. Samples from Geneva were used for sequencing of *NAC18.1*.

Lastly, 'Golden Delicious' apples from a commercial orchard in Nova Scotia were harvested before the climacteric stage from 2004-2007, as described previously (Yang et al., 2013, 2016). Samples from this previous work were used to test the expression levels of genes of interest.

Apple phenotyping

In 2017, we evaluated harvest date for 1,348 trees and fruit firmness for 1,328 trees within the ABC. Due to the diversity of apples within the collection, a variety of methods were used to determine the appropriate time to harvest. First, we observed if the tree had dropped fruit or, for red apples, if the fruit were a deep red color. Next, a sample apple was taken from each tree and touched to assess firmness, tasted to assess starch and sweetness, cut in half to check browning of seeds, and then sprayed with iodine solution to assess starch content (Blanpied and Silsby, 1992a). Fruit were deemed mature and ready to harvest at a starch-iodine index of 6 (Blanpied and Silsby, 1992b).

Once harvested, the fruit were evaluated for firmness. We recorded the firmness (kg/cm^2) of 5 fruit per tree using a penetrometer with a 1 cm diameter (Fruit Texture Analyzer, GS-14, Güss Manufacturing). A small section of skin was removed using a vegetable peeler, and each fruit was placed on the penetrometer platform so that the piston entered the middle of the apple where the skin had been removed. The data were automatically recorded into a spreadsheet. After these measurements were taken, 5 additional fruit from each accession (combined across trees/replicates) were placed in cold storage (5°C). Fruit was removed from storage after 3

months and was left at room temperature for 24 hours before being evaluated again for firmness using the same method. Firmness after storage and the percent change in firmness from harvest to post-storage were both calculated.

Harvesting fruit from the ABC orchard often lasted more than one day and so differences in harvest date within a week reflect the time required to harvest the orchard, rather than meaningful biological differences. As a result, we recorded harvest dates as the Monday of each week for all trees harvested throughout the week. We used the "lmer" function in the R package lme4 (Bates et al., 2015) to fit a REML model for harvest date and firmness at harvest. Next, we calculated the least squares mean using the "lsmeans" function in the lsmeans R package (Lenth, 2016), resulting in one value per accession. After running the REML model, we had 862 unique accessions with harvest dates and 859 accessions with firmness at harvest measurements. Due to the number of fruits available and storage capacity, replicates were combined from multiple trees prior to storage and so it was not possible to fit a REML model for firmness after storage or change in firmness measurements, which both had sample sizes of 535 unique accessions. We calculated the correlation between each of the phenotypes using Pearson correlation tests and the "ggpairs" function in the GGally R package (Schloerke et al., 2020).

Texture-related genetic markers

DNA was extracted from leaf tissue collected from the ABC orchard using silica columns, quantified using PicoGreen (Thermo) and normalized to a concentration of 20 ng μL^{-1}. Genotyping was conducted using PCR and high resolution melting (HRM) on a LightScanner HR384 (BioFire). Primers are listed in Supplemental Table S1.

Genotyping of the *PG1* SNP marker was based on the GenBank sequence L27743.1, where the T allele is favorable at position 437 and the G allele is unfavorable and leads to increased softening during storage (Costa et al., 2010). Three allelic combinations of the observed indel, *ACS1-1/1*, *1-1/2* and *1-2/2*, have been associated with high, medium, and low ethylene production, respectively (Sunako et al., 1999; Harada et al., 2000; Oraguzie et al., 2004; Costa et al., 2005). The exact position of the indel from *ACS1-2* (GenBank: AB010102.1) is 1,320 to 1,483 bp (163 bp), and from *ACS1-1* (GenBank: AY062129.1) is 4,500 to 4,525 bp (25 bp). The size difference between alleles is 138 bp. The *ACO1* marker involves an unfavorable 62 bp insertion in the third intron of *ACO1* (Costa et al., 2005). The third intron spans from 1,083 bp to 1,300 bp of the GenBank sequence Y14005.1 and the indel is found from position 1,297 bp to 1,358 bp. The D5Y mutation in *NAC18.1* is a nonsynonymous SNP at position 30,698,039 on chromosome 3, according to reference genome version GDDH13 v1.1 (Daccord et al. 2017) and is associated with both harvest date and firmness (Migicovsky et al., 2016). The desirable C allele encodes an aspartic acid (D) at the fifth amino acid position of the NAC18.1 protein, while the undesirable A allele encodes a tyrosine (Y). The names and IDs of samples, their phenotypes and their genotypes are provided in Table S2.

Phenotypic variance explained by texture-related markers

To determine the proportion of phenotypic variance explained by each marker of interest (ACO1, ACS1, PG1, and NAC18.1) we used a type 2 ANOVA from the 'car' package in R (Fox and Weisberg, 2018) with the markers encoded as co-dominant. To determine the phenotypic variance explained by each marker after accounting for harvest date, we also performed a type 2 ANOVA including the four markers and harvest date as factors. The results of the models were

visualized using the "geom_tile" function in the ggplot2 R package (Wickham, 2016). We determined the association between each marker and each phenotype using Spearman's rank correlation test. We visualized the results using the "geom_boxplot" function in ggplot2 in R (Wickham, 2016).

Sequencing of NAC18.1

DNA was isolated from leaves sampled from 18 apple accessions growing in Geneva, NY. A 2.3 kb amplicon including the *NAC18.1* gene and ~800 bp of upstream sequence was amplified by PCR using primers NAC18F2 and NAC18R2 (Table S1) and Phusion® High-Fidelity PCR Master Mix with HF Buffer (NEB). PCR product size and purity was confirmed by agarose gel electrophoresis, and the remaining product was purified using a DNA Clean & Concentrator kit (Zymo Research). The resulting DNA fragment was cloned into the plasmid pMiniT 2.0 and transformed into *E. coli* using the NEB® PCR Cloning Kit (NEB).

Individual colonies were selected for complete sequencing of the cloned amplicon using the primers NAC18F2, NAC18F3, NAC18F4, NAC18R1, and NAC18R2 (Table S1). For accessions homozygous for the D5Y SNP, the NAC18.1 amplicon from a single clone was sequenced. For heterozygous accessions, two clones representing each D5Y allele were selected based on partial sequencing of the D5Y region, followed by complete sequencing of the 2.3 kb amplicon, as described above. The nucleotide sequences were aligned using MUSCLE (Edgar, 2004) and used to construct a maximum-likelihood phylogenetic tree in MEGA7 (Kumar et al., 2016).

Gene expression analyses of apple fruit

Gene expression levels of *ACO1*, *ACS1*, *PG1*, and *NAC18.1* were evaluated using q-PCR

with and without treatment of ethylene and 1-methylcyclopropene (1-MCP), using methods

described previously in (Yang et al., 2013, 2016). Briefly, in 2004 and 2005, 'Golden Delicious'

fruit were harvested from a commercial orchard before the climacteric stage. Ethylene gas (36

μL L^{-1}) was applied for 24 hours at 20°C to initiate ripening while control fruits were stored for

24 hours at 20°C without ethylene. All fruit were then stored at 20°C for 21 days, with sampling

occurring at day 0, 7, 13, and 21 of each year. In 2006 and 2007, 'Golden Delicious' were once

again harvested at the pre-climacteric stage. Apples were either treated with 1-MCP (1 μL L^{-1} of

EthylBloc, 0.14%, Rohm and Haas Company) for 12 hours at 20°C in a sealed container, or

stored at 20°C for 12 hours without 1-MCP. Fruits were then stored at 20°C, with sampling

occurring on day 0, 7, 14, and 22.

Total RNA was extracted from frozen apple tissues using a hot borate method with some

modification in the extraction buffer, as described in (Yang et al., 2016). RNA extracts were

treated with DNase I using a DNA-free Kit following the manufacturer's recommendations

(Applied Biosystems). First-strand cDNA synthesis was performed using 2 μg DNase I-treated

total RNA. The oligonucleotide primers used for real-time quantitative qRT-PCR analysis were

designed from sequence information in NCBI (Table S1). Conditions for all PCR reactions were

optimized as previously described (Yang et al., 2013) and efficiency values for each gene are

shown in Table S3. Two reference genes, *MdActin* and *MdUBI*, were used in the real-time qPCR

analysis to normalize the expression patterns. Samples from day 0 (assigned an arbitrary quantity

of "1") were used as a calibrator to calculate relative quantities (Yang et al., 2016). The

experimental design was a balanced randomized block design with random effects of year, plate,

and row (on plate). The fixed effects were treatments for both ethylene (for example, control

(Day 7) and ethylene (Day 7)) and 1-MCP experiments (such as control (Day 7) and 1-MCP

(Day 7)).

Molecular cloning and plant transformation

The hypothetical coding sequences (CDSs) corresponding to the consensus sequence of

the "A" and "C" haplotypes of *NAC18.1* were synthesized as a double-stranded DNA (gBlock)

by Integrated DNA Technologies (IDT), with 20 bp of flanking sequences added to both ends to

facilitate Gibson Assembly of the *NAC18.1* sequences between the *Asc*I and *Pac*I restriction sites

of pMDC32 (Curtis and Grossniklaus, 2003). Constructs were assembled using the NEBuilder

HiFi DNA Assembly Cloning Kit (NEB) and their integrity verified by Sanger sequencing.

Plasmids were transformed into *Agrobacterium tumefaciens*, which was used to transform

tomato (*S. lycopersicum*) cotyledon explants (Van Eck et al., 2019) derived from the tomato *nor*

mutant in the cv. Ailsa Craig background (LA3770, Tomato Genetics Resource Center,

https://tgrc.ucdavis.edu/). A total of 10 T0 plants were recovered per construct and plants derived

from two independent transformation events per construct were selected for further

characterization in the T1 generation.

Transgenic tomato characterization

Fruit were harvested when visually ripe, or in the case of the *nor* mutant, at the equivalent

age as ripe cv. Ailsa Craig fruit (the near isogenic wild-type control), as determined by tagging

of flowers at anthesis. The color of the fruit surface was measured using a CR-400 Chroma

Meter (Konica Minolta), and fruit were weighed, photographed, and dissected. Pericarp tissue

was frozen in liquid nitrogen and stored at -80˚C. Frozen tissue was ground to a fine powder and RNA extracted using a modified version of the protocol described in Chang et al. (1993). Briefly, approximately 400 mg of tissue was added to a preheated (80˚C) two-phase system consisting of 500 µL of water-saturated phenol and 500 µL of extraction buffer (100 mM Tris [pH 8.0], 25 mM EDTA, 2 M NaCl, 2% CTAB, 2% PVP, and 2% [v/v] beta-mercaptoethanol). The mixture was vortexed and incubated for 5 min at 65˚C and then cooled to room temperature before extracting and precipitating RNA, as previously described (Chang et al., 1993).

RNA was treated with RNase-free DNase (Promega) and used for cDNA synthesis with RNA to cDNA EcoDry™ Premix with Oligo dT primer (Takara Bio). The cDNA was used as a template for quantitative PCR using Luna Universal qPCR Master Mix (NEB) and a Viia7 real-time PCR instrument (Life Technologies/ABI). Gene-specific primers are listed in Table S1. Quantification used the ΔC_t method with *RIBOSOMAL PROTEIN L2* (*RPL2*) as a reference gene, and statistical significance of the ΔC_t values was tested by a one-way ANOVA followed by Tukey's HSD test.

Results

Evaluation of texture-related markers

Fruit were harvested over a 65 day period and their firmness at harvest (N = 859) and after 3 months of cold storage (N = 535) was found to differ by 7-fold across the apple accessions from the ABC. We observed a strong relationship between harvest date and firmness: late-harvested apples were firmer both at harvest (R^2 = 0.25, p < 1 x 10^{-15}) and after storage (R^2 = 0.24, p < 1 x 10^{-15}), and they also softened less during storage (R^2 = 0.086, p = 4.53 x 10^{-12})

(Figure S1). Firmness at harvest was also significantly correlated with firmness after storage (R^2 = 0.54, p < 1 x 10^{-15}; Figure S1). Post-harvest storage resulted in significant softening: on average, apples lost 40 percent of their firmness during 3 months of cold storage (Table S2).

We tested the utility of four genetic markers to predict firmness-related phenotypes and found that the *NAC18.1* marker outperformed the other three for both firmness at harvest and firmness after storage. However, softening (the loss in firmness during storage) was best predicted by the *PG1* marker. Our results suggest that the markers in *ACO1* and *ACS1* have little to no predictive power for firmness-related traits across diverse apple germplasm (Figure 1; Figure S2). When we performed the same analysis with harvest date as a factor, harvest date accounted for 10.9% and 14.6% of the variation in firmness at harvest and firmness after storage, respectively. In this model, the amount of phenotypic variance in firmness at harvest explained by *NAC18.1* was reduced from 18.19%, when harvest date was not included in the model, to 2.64% (Figure 1; Figure S3).

Figure 1: The utility of four genetic markers for predicting harvest date and firmness-related apple phenotypes. For each trait, the percent variance explained for each of the four markers is shown in cases where a significant effect was detected from a type 2 ANOVA. A type 2 ANOVA was also performed with the four genetic markers and harvest date as a co-factor (Figure S3).

The genotypes of the four texture-related genetic markers across the nine most popular apple cultivars sold in the USA in 2018 (Home - USApple 2021) are presented in Figure 2. All nine cultivars were homozygous for the desirable (firm) C allele of *NAC18.1*, while only 2 to 4 of the cultivars had homozygous desirable genotypes for the other markers. Among the top cultivars, only 'Fuji', released in 1962, was homozygous for desirable alleles at all markers, while 'McIntosh', initially discovered in 1811 and commercially released in 1870, was homozygous for undesirable alleles at all markers except for *NAC18.1*. 'Empire', an offspring of 'McIntosh' and 'Red Delicious' released in 1966, was homozygous for the desirable *NAC18.1* alleles like its parents, but inherited undesirable alleles from 'McIntosh' and thus was heterozygous for *ACS1* and *ACO1*, and homozygous for undesirable *PG1* alleles.

Release Date	Cultivar	ACS1	ACO1	PG1	NAC18.1
1868	'Granny Smith'	Heterozygous	Homozygous (Undesirable)	Homozygous (Desirable)	Homozygous (Desirable)
1870	'McIntosh'	Homozygous (Undesirable)	Homozygous (Undesirable)	Homozygous (Undesirable)	Homozygous (Desirable)
1895	'Red Delicious'	Heterozygous	Homozygous (Desirable)	Heterozygous	Homozygous (Desirable)
1914	'Golden Delicious'	Heterozygous	Homozygous (Undesirable)	Heterozygous	Homozygous (Desirable)
1962	'Fuji'	Homozygous (Desirable)	Homozygous (Desirable)	Homozygous (Desirable)	Homozygous (Desirable)
1966	'Empire'	Heterozygous	Heterozygous	Homozygous (Undesirable)	Homozygous (Desirable)
1974	'Gala'	Homozygous (Desirable)	Heterozygous	Heterozygous	Homozygous (Desirable)
1990	'Pink Lady®'	Heterozygous	Homozygous (Undesirable)	Homozygous (Desirable)	Homozygous (Desirable)
1991	'Honeycrisp'	Heterozygous	Homozygous (Undesirable)	Homozygous (Desirable)	Homozygous (Desirable)

Genotypes ■ Homozygous (Desirable)   ■ Heterozygous   ■ Homozygous (Undesirable)

Figure 2: Genotypes of four texture-related genetic markers across the top 9 apple cultivars sold in the United States (U.S. Apple Association). The "desirable" allele for each marker is defined as the allele that has been reported to lead to firmer apple texture.

Using q-PCR, we evaluated the expression of the four candidate firmness genes across a three week period and observed that transcript levels of *ACO1*, *ACS1* and *PG1* genes were up-regulated in response to treatment with ethylene and suppressed following exposure to the ethylene-inhibitor, 1-MCP. *NAC18.1* transcript levels, however, remained unaffected by exposure to ethylene and 1-MCP (Figures S4 and S5).

Resequencing of NAC18.1

DNA sequences from 24 *NAC18.1* haplotypes confirmed the expected D5Y genotype in all individuals and revealed a number of additional SNPs and indels within both coding and non-

coding regions of *NAC18.1* (Table S4). A multiple sequence alignment and subsequent

phylogenetic analysis indicated two major clades, corresponding to the D5Y A and C alleles

(Figure 3A). In addition to the amino acid change resulting from D5Y, several additional amino

acid changing polymorphisms were observed between sequences from D5Y genotypes and the

reference sequence of *NAC18.1*. For example, near the site of the D5Y polymorphism, all "A"

haplotypes also had a 12 nucleotide insertion that introduced the amino acid sequence QPQP

(Figure 3B, Figure S6).

Figure 3: Polymorphisms within the *NAC18.1* gene. (A) Maximum likelihood phylogenetic tree

of 24 *NAC18.1* sequences from 18 apple cultivars and the reference genome (GDDH13 v1.1)

sequences of *NAC18.1* (Md03g1222600) and its closest homolog (Md11g1239900). A graphical

representation of the 2.3 kb multiple sequence alignment is also shown. (B) Amino acid

sequence alignment of the N-terminal region of *NAC18.1,* illustrating additional variation in the

coding sequence in strong LD with the D5Y variant. The complete amino acid sequence

alignment is provided in Figure S6.

Transgenic complementation of the tomato non-ripening (nor) mutant

A BLAST search of predicted proteins from the ITAG 3.20 tomato genome (https://solgenomics.net/) using *NAC18.1* as a query identified only two matching sequences: *Solyc10g006880* (*NOR*) and *Solyc07g063420* (*NOR-LIKE 1*). To test the functional conservation of *NAC18.1* and *NOR*, we introduced constructs individually conferring constitutive expression of each of the *NAC18.1* haplotype CDSs into the tomato *nor* mutant. Two independent lines for each construct were characterized in the T1 generation with respect to their ability to rescue the ripening deficiency of the *nor* mutant. In contrast to *nor*, fruit from all four lines changed color at maturity, although internal fruit color change did not occur to the same extent as observed in a WT control (Figure 4A). To complement this qualitative phenotypic assessment, we also conducted quantitative colorimetry of the surface of the fruit (Figure 4B). Fruit from all the transgenic lines exhibited a significant increase in the *a** (green-red) component of color space relative to the *nor* mutant, although only *NAC18.1^A #6* achieved similar *a** levels to WT.

Figure 4: Transgenic complementation of the tomato *nor* mutant using *NAC18.1* transgene

restores ripening. **(**A) Mature fruit of the tomato *nor* mutant and four independent T1 transgenic

lines constitutively expressing either of two alleles of the *NAC18.1* transgene, *NAC18.1^C* and

NAC18.1^A and isogenic WT control (cv. Ailsa Craig). (B) Quantitative colorimetry of the fruit

surface of *nor,* the transgenic fruit and WT fruit. The *a** component (green-red axis) is shown, with the mean ± SE superimposed in black over the raw values ($n = 5$) in a color approximating the external color of the fruit. Genotypes not sharing a letter (a-e) are statistically distinct by one-way ANOVA and Tukey's HSD test ($p < 0.05$).

The different degrees to which ripe fruit color was restored in the transgenic plants might be a consequence of either the different alleles of *NAC18.1*, or of different levels of transgene expression in each line. To address this, we analyzed the expression level of *NAC18.1* using qRT-PCR primers designed to target both *NAC18.1* alleles. Expression levels of the *NAC18.1* transgene were not statistically different between each independent line ($p > 0.5$) (Figure S7A). Next, we measured the expression level of several genes associated with tomato ripening physiology: *PHYTOENE SYNTHASE 1 (PSY1)* encodes an enzyme in an early stage of the carotenoid synthesis pathway, which is responsible for the production of red pigments during fruit ripening (Fray and Grierson, 1993), and its expression is impaired in the *nor* mutant (Osorio *et al.*, 2011); *POLYGALACTURONASE 2* (*PG2*) encodes an enzyme catalyzing pectin depolymerization associated with fruit softening (Biggs and Handa, 1989); and *1-AMINOCYCLOPROPANE-1-CARBOXYLATE SYNTHASE 2* (*ACS2*) encodes an enzyme that synthesizes 1-aminocyclopropane-1-carboxylate, the immediate precursor of ethylene (Nakatsuka et al., 1998).

Expression of all three marker genes was enhanced in the *NAC18.1* transgenic lines relative to the *nor* mutant control, although not to the same extent as observed in WT ripe fruit (Figure S7B-D). Visual color change was consistent with the expression of the carotenoid biosynthetic gene *PSY1* (Figure S7B). Gene expression analysis further confirmed the induction

of genes involved in ripening-associated cell wall remodeling (Figure S7C) and ethylene synthesis (Figure S7D). In contrast to the consistent level of *NAC18.1* expression observed in each line, the marker genes were more variable in their expression levels between lines. A similar pattern was observed for all marker genes, with the *NAC18.1^C* #9 line showing the smallest induction of marker gene expression relative to *nor*. In the case of *PG2* and *ACS2*, the difference in expression in *NAC18.1^C* #9 was not statistically significant relative to *nor* ($p = 0.09$ and 0.18, respectively). Consistent with these results, fruit from this line also exhibited the lowest amount of red color development (Figure S7B). Taken together, these results indicate that a canonical ripening program can be induced in the tomato *nor* mutant through the heterologous expression of either apple *NAC18.1* allele.

Discussion

Genomics-assisted breeding has tremendous potential in perennial crops, such as apple, where a lengthy juvenile phase and large plant size make phenotyping at the adult stage time-consuming and expensive. However, the genetic markers used for culling progeny at the seedling stage must accurately predict the trait of interest at the adult stage in order for genomics-assisted breeding to be effective and cost-efficient (Luby and Shaw, 2001). Apple texture has been repeatedly identified as a key breeding target because of consumer demand for crisp, firm apples that retain their desirable texture during storage (Harker et al., 2003; Yue et al., 2013; Laurens et al., 2018). Markers for three genes (*ACS1*, *ACO1*, and *PG1*) are widely used in genomics-assisted apple breeding programs to predict firmness. However, GWAS for firmness suggests a single major effect locus for firmness at the *NAC18.1* gene (Migicovsky et al., 2016; Urrestarazu et al., 2017; McClure et al., 2018; Larsen et al., 2019). Here we have presented further evidence

that whereas *ACS1*, *ACO1* and, *PG1* are weak predictors of firmness at harvest in genetically diverse apples, *NAC18.1* is a strong predictor and may serve a functional role in the fruit ripening pathway.

Single-marker correlation tests between each of the four markers and four phenotypes revealed statistically significant ($P < 0.05$) associations in every case except one (Figure S2). This result was not surprising: many genome-wide markers are expected to be correlated with firmness-related phenotypes since the genetic structure of our apple population is strongly correlated with these traits. For example, without correcting for the effects of population structure, 39% and 17% of genome-wide SNPs were significantly associated ($P < 0.05$) with harvest date and firmness, respectively, in the USDA collection, which is genetically identical to most of the population studied here (Migicovsky et al., 2016). Single-marker association tests do not account for population structure using genome-wide markers, as is customarily done when performing GWAS. Thus, we are only able to compare the relative power of each marker to predict these phenotypes. We found that the D5Y marker in *NAC18.1* had a 3 to 14 times greater effect on firmness at harvest and harvest date than the markers in *ACO1*, *ACS1* and *PG1* (Figure S2). We assessed the combined effects of the markers on these phenotypes using type 2 ANOVA and demonstrated that the D5Y marker in the *NAC18.1* gene is a far stronger predictor of harvest date and firmness at harvest than the markers for *ACS1*, *ACO1* and *PG1*. For predicting firmness after storage, however, *NAC18.1* performs only slightly better than *PG1*, while softening during three months of cold storage is best predicted by the marker in *PG1* (Figure 1). It is worth noting that a recent GWAS found that variation in firmness loss may not be due to variation within *PG1* but rather at a neighboring ethylene response factor (ERF) gene, *ERF* (MDP0000855671) (McClure et al., 2018). *ERF* may have been missed by previous linkage mapping studies due to a

lack of mapping resolution compared to GWAS. Future fine mapping is required to determine if markers at *ERF* serve as superior predictors of softening during storage compared to the widely used *PG1* marker.

In our study, by far the strongest association observed for any marker-phenotype combination was between harvest date and the D5Y marker in *NAC18.1*: 41% of the variance in harvest date is accounted for by *NAC18.1* (Figure 1). On average, accessions homozygous for the late-ripening *NAC18.1* allele are harvested 29 days later than accessions homozygous for the early-ripening allele, which represents nearly half of the 65-day harvest season (Figure S2). This observation is consistent with strong GWAS signals for harvest date and ripening period discovered in and around the *NAC18.1* gene (Migicovsky et al., 2016; Urrestarazu et al., 2017; Larsen et al., 2019; Jung et al., 2020). The D5Y marker was also the strongest predictor of firmness at harvest and firmness after storage (Figure 1). We observed strong positive correlations between harvest date and both firmness at harvest ($R^2 = 0.25$, $p < 1 \times 10^{-15}$) and firmness after storage ($R^2 = 0.24$, $p < 1 \times 10^{-15}$; Figure S1), consistent with previous work showing that early-ripening apples tend to be softer (Vincent, 1989; Johnston et al., 2002; Oraguzie et al., 2004, 2007; Nybom et al., 2013; Chagné et al., 2014). When harvest date is included as a factor in the type 2 ANOVA instead of a response variable, it is the strongest predictor of firmness both at harvest and after storage (Figure S3). Thus, an apple's firmness is largely determined by its harvest date. Therefore, markers that predict harvest date, such as the D5Y marker in *NAC18.1*, will be more effective for breeding than the other firmness-related markers tested here. This conclusion is in agreement with previous work suggesting that screening for *ACO1*, *ACS1* and *PG1* is not cost effective and that a marker for harvest date is of greater value to improving firmness via marker-assisted breeding (Nybom et al., 2013).

The best predictor of softening during storage was the marker in *PG1*, which accounted for 8% of the variance. The *PG1* marker was also significantly associated with firmness after storage, accounting for 7% of the variance in addition to the 9% accounted for by *NAC18.1* (Figure 1). Most apples are consumed after being stored. Thus, the most relevant phenotype for apple quality from the perspectives of both consumers and breeders is firmness after storage. The firmness of an apple after storage is a consequence of how firm it was when harvested and how much firmness loss it experienced during storage. Firmness loss, or softening, in climacteric fruit like apple is regulated by ethylene, and previous studies have shown that *ACO1*, *ACS1* and *PG1* transcripts increase when exposed to ethylene (Atkinson et al., 1998; Wakasa et al., 2006; Mann et al., 2008; Zhu and Barritt, 2008; Costa et al., 2010; Longhi et al., 2012; Di Guardo et al., 2017). We confirmed this result by showing that expression of these three genes is up-regulated by ethylene and down-regulated by the ethylene inhibitor, 1-MCP (Figures S4 and S5). The expression of *NAC18.1*, however, was unaffected by treatments with ethylene and 1-MCP. NAC transcription factors in apple are primarily involved in the regulation of growth and development (Li et al., 2020) and previous work has suggested that ethylene may not be required for on-tree apple ripening (Lau et al., 1986; Blankenship and Unrath, 1988). Our results suggest that genetic variation at the *NAC18.1* locus affects fruit firmness before harvest via a ripening pathway that is independent of ethylene, while softening during storage is ethylene-dependent and influenced by variation in or near *PG1*.

In the context of a marker-assisted breeding program, our results suggest that apple texture could be improved by selecting for the D5Y marker in *NAC18.1*. However, the potential for this marker to improve apple texture should be considered in light of several factors. First,

we measured firmness with a penetrometer, but we recognize that our measures are only a proxy for the texture desired by consumers, which may be better captured using consumer panels and/or other mechanical devices that better mimic the chewing process (Costa et al., 2011). Second, all 9 of the most popular cultivars in the USA tested here are homozygous for the *NAC18.1* allele associated with late-harvested, firm apples (Figure 2). Out of 1,056 accessions genotyped, 696 (66%) are homozygous for the desirable *NAC18.1* allele, and its high frequency suggests this allele may already be under selection by breeders. Indeed, a recent population genetic analysis found evidence of positive selection for the desirable *NAC18.1* allele (Migicovsky et al., 2021). As a result, this allele may not segregate in many breeding populations. This may be the reason that *NAC18.1* was only identified as a firmness locus via GWAS in diverse populations, and not in numerous bi-parental breeding populations, emphasizing the need for germplasm collections that maintain diverse populations (Migicovsky et al. 2019). Third, highly aromatic apple cultivars are generally less firm, and selection for firmness using the D5Y marker may therefore result in lower production of volatile organic compounds, and thus apples with less pronounced aromas (Song and Bangerth, 1996; Farneti et al., 2017). Finally, selection for firmness using the D5Y marker will also select for late-harvested cultivars, which could result in a future excess of commercial cultivars with a compressed harvest window near the end of the harvest season, risking fruit loss to late season weather and reducing the opportunity to pick and sell fruit throughout the season. A more thorough understanding of the relationship between harvest date and firmness may lead to novel ways to break the correlation between these two phenotypes and to thereby enable the development of new apple cultivars with desirable firmness attributes, whose harvest dates are spread throughout the harvest season.

While the D5Y mutation in *NAC18.1* is a strong functional candidate variant for apple

firmness, we discovered numerous DNA sequence variants in *NAC18.1*, often in perfect LD with

D5Y, that are also putatively functional (Figure 3). The magnitude of DNA polymorphism at this

locus, and the multitude of putatively causal variants, is consistent with the results of Larsen et

al. (2019), who found 18 SNPs and 2 indels within *NAC18.1* across 11 apple accessions. Our

analysis of *NAC18.1* sequences revealed a number of mutations that are candidates for causal

association with ripening phenotypes. For example, we observed an insertion of four amino acids

(QPQP) 11 amino acids downstream of the D5Y mutation in all A haplotypes upstream of the

NAC DNA binding domain. Glutamine-rich sequences are common in eukaryotic transcription

factors, and polymorphisms in these motifs have been shown to alter the activity of

transcriptional activators (Atanesyan et al., 2012). On the other hand, we also observed several

SNPs resulting in amino acid changes within the DNA binding domain and C-terminal

transcriptional activator domain of *NAC18.1*. Given the pattern of LD we observed across

NAC18.1 (Figure 3), there remains the possibility that the GWAS signals at *NAC18.1* are driven

by a causal variant outside of the *NAC18.1* coding sequence that acts independently of the

NAC18.1 gene. The identification of a causal polymorphism within the coding sequence or

promoter of *NAC18.1* will require additional high-resolution genetic mapping and/or

transcriptome analysis efforts.

Since *NAC18.1* is a homolog of the well-known ripening gene *NOR* in tomato, we used

tomato as a model to explore the role of *NAC18.1* in the ripening process. We observed enhanced

ripening in the tomato *nor* mutant following heterologous expression of either of the *NAC18.1*

alleles (Figure 4). Although the expression level of the *NAC18.1* transgene was comparable

across transgenic lines (Figure S7A), the level of ripening marker induction in the distinct lines

was significantly different (Figure S7B-D). We had hypothesized that an allele-specific trend in our transgenic experiments would indicate that coding sequence variants are responsible for the association of *NAC18.1* with variation in apple ripening. While our results cannot confirm an allele-specific trend, they do verify that both alleles of *NAC18.1* encode functional proteins that can promote ripening.

In tomato, *nor* was first identified as a spontaneous mutation in an heirloom cultivar (Tigchelaar et al., 1973). While this *nor* allele exhibits recessive behavior and has been assumed to confer a complete loss-of-function, due to a 2 nucleotide deletion resulting in a truncated protein (Giovannoni et al., 2004), it was recently demonstrated that it is actually a dominant negative allele (Gao et al., 2019; Wang et al., 2019). These studies used CRISPR/Cas9 to generate *bona fide* null alleles of *nor*, which showed evidence of ripening relative to the spontaneous *nor* allele, although to a lesser extent than WT fruit. In light of these studies, we speculate that the action of the *NAC18.1* transgenes was likely attenuated by the dominant-negative activity of the spontaneous *nor* allele used in our work. Further heterologous characterization of *NAC18.1* would likely benefit from the use of null *nor* mutants, or double mutants of *nor* and *nor-like 1* (Gao et al., 2018). This may allow for more precise quantitative comparisons between alleles in order to resolve whether differences in the coding sequence of *NAC18.1* confer different degrees of ripening.

Although we are unable to conclude whether polymorphisms in coding or regulatory sequences affect the activity of *NAC18.1* in apple, an increasing body of evidence generated using tomato as a model system indicates that coding sequence polymorphisms of *NOR* can influence firmness and timing of ripening. For example, the 'Alcobaça' tomato cultivar has firm fruit, delayed ripening, and long shelf life that is conferred by the *alcobaça* mutation, and a

recent study has shown that *ALCOBACA* is allelic with *NOR*, and that the *alcobaça* allele of *NOR* contains a valine to aspartate mutation at position 106 within the NAC domain (Kumar et al., 2018). An additional complete loss of function allele of *NOR* was also found in another long shelf life tomato cultivar (Kumar et al., 2018).

In conclusion, the results presented here provide evidence that *NAC18.1* is involved in apple ripening via an ethylene-independent mechanism, and that genetic variation in and/or near the *NAC18.1* gene influences apple ripening. The present study lays the groundwork for future efforts to compare the effect of different *NAC18.1* haplotypes and determine the precise causal variant(s) underlying this agriculturally important gene. Ultimately, we envision that the identification of precise causal variants for ripening in apple could be exploited using gene editing for fast and efficient introgression of desirable texture phenotypes.

Data availability

The authors declare that all data supporting the findings of this study are available within the paper and its supplementary information files.

Acknowledgments

We thank Susan Brown (Cornell University), Benjamín Gutiérrez (USDA-ARS Plant Genetics Resource Unit), and C. Thomas Chao (USDA-ARS Plant Genetics Resource Unit) for apple leaf tissue samples from orchards of the Cornell apple breeding program and the USDA apple germplasm collection. Transgenic tomato plants were generated by the Boyce Thompson Institute Biotechnology Center (https://btiscience.org/). ZM was supported by National Science Foundation Plant Genome Research Program 1546869. TY and JR were supported by a grant

from the USDA-ARS Multistate (NE-1336) project, NYC-184821. This research was supported

in part by funding the National Sciences and Engineering Research Council of Canada (SM) and

A-Base funding (NOI-1238) from Agriculture and Agri-Food Canada (JS). The authors

gratefully acknowledge the farm services team at the Kentville Research and Development

Centre for maintaining the apple collection studied here.

Conflicts of interest

The authors declare no conflicts of interest.

Supplemental Figures

Figure S1: Correlations among phenotypes. The distributions of each phenotype are shown as well as dot plots of comparisons between each pair of phenotypes. The results of a Pearson correlation test are provided for each pairwise comparison.

Figure S2: Genotype-phenotype correlations. Box plots display the variation in phenotypes across genotypic classes for four different firmness-related genetic markers. The phenotypes include (A) harvest date, (B) firmness at harvest, (C) firmness after storage, and (D) change in firmness (softening). Sample sizes are shown in parentheses under each genotypic class. The desirable genotype is on the right, while the undesirable genotype is on the left in every case. The P value and rho value are shown from performing a Spearman rank correlation test. The difference in phenotype values between genotypic classes are shown across the top of each plot.

Figure S3: Prediction of firmness-related apple phenotypes. Four genetic markers and harvest date were included as factors in a type 2 ANOVA with three different phenotypes as outcomes. The proportion of the variance explained is shown in cases where a statistically significant result ($P < 0.05$) was observed.

Figure S4: Relative expression of four firmness-related genes with and without exposure to ethylene over 3 weeks of storage. The ANOVAs testing for the effects of ethylene were significant (p < 0.05) for ACO1 (a), ACS1 (b) and PG1 (c), but not for NAC18.1 (d). Due to the balanced design, one standard error of the means (SEM) bar is used to represent the SEM for each population.

Figure S5: Relative expression of four firmness-related genes with and without exposure to the ethylene-inhibitor 1-MCP over 3 weeks of storage. The ANOVAs testing for the effects of 1-MCP were significant ($p < 0.05$) for ACO1 (a), ACS1 (b) and PG1 (c), but not for NAC18.1 (d). Due to the balanced design, one standard error of the means (SEM) bar is used to represent the SEM for each population.

Figure S6: Amino acid sequence alignment of *NAC18.1*. Polymorphisms are highlighted in colour. Raw sequencing data is available in Table S4 in fasta format.

Figure S7: Ripening marker gene expression in transgenic tomatoes. Gene expression was evaluated as the difference in threshold cycle (ΔC_t, $\log_2$ scale) by qRT-PCR using *RPL2* as a reference gene. The mean ± SE is superimposed in black over the raw values in gray ($n = 5$). Genotypes not sharing a letter (a-d) are statistically distinct by one-way ANOVA and Tukey's HSD test ($p < 0.05$). (A) The *NAC18.1* transgene. (B) *PSY1*, encoding the phytoene synthase

carotenoid biosynthetic gene. (C) *PG2*, encoding the major ripening-associated polygalacturonase involved in pectin hydrolysis. (D) *ACS2*, encoding 1-aminocyclopropane-1-carboxylate synthase, a ripening-associated isoform of the ethylene biosynthesis enzyme.

Supplemental Table Legends

Table S1: Primer names and sequences.

Table S2: A list of sample IDs, sample names, genotypes and phenotypes.

Table S3: Efficiency values for gene expression assays of apple genes ACO1, ACS1, PG1 and NAC18.1.

Table S4: DNA sequences from 24 *NAC18.1* haplotypes in fasta format. Amino acid sequence alignment is visualized in Figure S6.

References

Atanesyan, L., Günther, V., Dichtl, B., Georgiev, O., and Schaffner, W. (2012). Polyglutamine tracts as modulators of transcriptional activation from yeast to mammals. *Biological Chemistry* 393, 63–70. doi:10.1515/bc-2011-252.

Atkinson, R. G., Bolitho, K. M., Wright, M. A., Iturriagagoitia-Bueno, T., Reid, S. J., and Ross, G. S. (1998). Apple ACC-oxidase and polygalacturonase: ripening-specific gene expression and promoter analysis in transgenic tomato. *Plant Mol. Biol.* 38, 449–460.

Atkinson, R. G., Sutherland, P. W., Johnston, S. L., Gunaseelan, K., Hallett, I. C., Mitra, D., et al. (2012). Down-regulation of POLYGALACTURONASE1 alters firmness, tensile strength and water loss in apple (Malus x domestica) fruit. *BMC Plant Biol.* 12, 129.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects Models Usinglme4. *Journal of Statistical Software* 67. doi:10.18637/jss.v067.i01.

Baumgartner, I. O., Kellerhals, M., Costa, F., Dondini, L., Pagliarani, G., Gregori, R., et al. (2016). Development of SNP-based assays for disease resistance and fruit quality traits in apple (Malus × domestica Borkh.) and validation in breeding pilot studies. *Tree Genet. Genomes* 12, 35.

Bink, M. C. A. M., Jansen, J., Madduri, M., Voorrips, R. E., Durel, C.-E., Kouassi, A. B., et al. (2014). Bayesian QTL analyses using pedigreed families of an outcrossing species, with application to fruit firmness in apple. *Theor. Appl. Genet.* 127, 1073–1090.

Blankenship, S. M., and Unrath, C. R. (1988). Internal ethylene levels and maturity of «Delicious» and «Golden Delicious» apples destined for prompt consumption. *J. Am. Soc. Hortic. Sci.* 113, 88–91.

Blanpied, G. D., and Silsby, K. J. (1992a). Predicting Harvest Date Windows for Apples. Available at: https://ecommons.cornell.edu/handle/1813/3299 [Accessed August 28, 2020].

Blanpied, G. D., and Silsby, K. J. (1992b). Predicting Harvest Date Windows for Apples. *Cornell University Library*. Available at: https://ecommons.cornell.edu/handle/1813/3299 [Accessed October 26, 2020].

Brummell, D. A., and Harpster, M. H. (2001). Cell wall metabolism in fruit softening and quality and its manipulation in transgenic plants. *Plant Mol. Biol.* 47, 311–340.

Chagné, D., Dayatilake, D., Diack, R., Oliver, M., Ireland, H., Watson, A., et al. (2014). Genetic and environmental control of fruit maturation, dry matter and firmness in apple (Malus × domestica Borkh.). *Hortic Res* 1, 14046.

Chagné, D., Vanderzande, S., Kirk, C., Profitt, N., Weskett, R., Gardiner, S. E., et al. (2019). Validation of SNP markers for fruit quality and disease resistance loci in apple (Malus × domestica Borkh.) using the OpenArray® platform. *Horticulture Research* 6. doi:10.1038/s41438-018-0114-2.

Chang, S., Puryear, J., and Cairney, J. (1993). A simple and efficient method for isolating RNA from pine trees. *Plant Molecular Biology Reporter* 11, 113–116. doi:10.1007/bf02670468.

Costa, F., Cappellin, L., Longhi, S., Guerra, W., Magnago, P., Porro, D., et al. (2011). Assessment of apple (Malus×domestica Borkh.) fruit texture by a combined acoustic-mechanical profiling strategy. *Postharvest Biol. Technol.* 61, 21–28.

Costa, F., Cappellin, L., Zini, E., Patocchi, A., Kellerhals, M., Komjanc, M., et al. (2013). QTL validation and stability for volatile organic compounds (VOCs) in apple. *Plant Sci.* 211, 1–7.

Costa, F., Peace, C. P., Stella, S., Serra, S., Musacchi, S., Bazzani, M., et al. (2010). QTL dynamics for fruit firmness and softening around an ethylene-dependent polygalacturonase gene in apple (Malusxdomestica Borkh.). *Journal of Experimental Botany* 61, 3029–3039. doi:10.1093/jxb/erq130.

Costa, F., Stella, S., Van de Weg, W. E., Guerra, W., Cecchinel, M., Dallavia, J., et al. (2005). Role of the genes Md-ACO1 and Md-ACS1 in ethylene production and shelf life of apple (Malus domestica Borkh). *Euphytica* 141, 181–190. doi:10.1007/s10681-005-6805-4.

Daccord, N., Celton, J.-M., Linsmith, G., Becker, C., Choisne, N., Schijlen, E., et al. (2017). High-quality de novo assembly of the apple genome and methylome dynamics of early fruit development. *Nat. Genet.* 49, 1099–1106.

Dandekari, A. M., Teo, G., Defilippi, B. G., Uratsu, S. L., Passey, A. J., Kader, A. A., et al. (2004). Effect of down-regulation of ethylene biosynthesis on fruit flavor complex in apple fruit. *Transgenic Res.* 13, 373–384.

Di Guardo, M., Bink, M. C. A. M., Guerra, W., Letschka, T., Lozano, L., Busatto, N., et al. (2017). Deciphering the genetic control of fruit texture in apple by multiple family-based analysis and

genome-wide association. *J. Exp. Bot.* 68, 1451–1466.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32, 1792–1797. doi:10.1093/nar/gkh340.

Farneti, B., Di Guardo, M., Khomenko, I., Cappellin, L., Biasioli, F., Velasco, R., et al. (2017). Genome-wide association study unravels the genetic control of the apple volatilome and its interplay with fruit texture. *J. Exp. Bot.* 68, 1467–1478.

Fox, J., and Weisberg, S. (2018). *An R Companion to Applied Regression*. SAGE Publications.

Gao, Y., Wei, W., Zhao, X., Tan, X., Fan, Z., Zhang, Y., et al. (2018). A NAC transcription factor, NOR-like1, is a new positive regulator of tomato fruit ripening. *Horticulture Research* 5. doi:10.1038/s41438-018-0111-5.

Gao, Y., Zhu, N., Zhu, X., Wu, M., Jiang, C.-Z., Grierson, D., et al. (2019). Diversity and redundancy of the ripening regulatory networks revealed by the fruitENCODE and the new CRISPR/Cas9 CNR and NOR mutants. *Horticulture Research* 6. doi:10.1038/s41438-019-0122-x.

García-Gómez, B. E., Salazar, J. A., Dondini, L., Martínez-Gómez, P., and Ruiz, D. (2019). Identification of QTLs linked to fruit quality traits in apricot (Prunus armeniaca L.) and biological validation through gene expression analysis using qPCR. *Molecular Breeding* 39. doi:10.1007/s11032-018-0926-7.

Giovannoni, J., Nguyen, C., Ampofo, B., Zhong, S., and Fei, Z. (2017). The Epigenome and Transcriptional Dynamics of Fruit Ripening. *Annual Review of Plant Biology* 68, 61–84. doi:10.1146/annurev-arplant-042916-040906.

Giovannoni, J., Tanksley, S., Vrebalov, J., and Noensie, F. (2004). Nor gene compositions and methods for use thereof. *US Patent*.

Grierson, D. (2013). "Ethylene and the Control of Fruit Ripening," in *The Molecular Biology and Biochemistry of Fruit Ripening* The Plant Hormone Ethylene., eds. G. B. Seymour, M. Poole, J. J. Giovannoni, and G. A. Tucker (Oxford, UK: Blackwell Publishing Ltd.), 43–73.

Harada, T., Sunako, T., Wakasa, Y., Soejima, J., Satoh, T., and Niizeki, M. (2000). An allele of the 1-aminocyclopropane-1-carboxylate synthase gene (Md-ACS1) accounts for the low level of ethylene production in climacteric fruits of some apple cultivars. *Theoretical and Applied Genetics* 101, 742–746. doi:10.1007/s001220051539.

Harker, F. R., Gunson, F. A., and Jaeger, S. R. (2003). The case for fruit quality: an interpretive review of consumer attitudes, and preferences for apples. *Postharvest Biol. Technol.* 28, 333–347.

Johnston, J. W., Hewett, E. W., Hertog, M. L. A. T. M., and Harker, F. R. (2002). Harvest date and fruit size affect postharvest softening of apple fruit. *J. Hortic. Sci. Biotechnol.* 77, 355–360.

Jung, M., Roth, M., Aranzana, M. J., Auwerkerken, A., Bink, M., Denancé, C., et al. (2020). The apple REFPOP—a reference population for genomics-assisted breeding in apple. *Horticulture Research* 7, 189.

Kumar, R., Tamboli, V., Sharma, R., and Sreelakshmi, Y. (2018). NAC-NOR mutations in tomato Penjar accessions attenuate multiple metabolic processes and prolong the fruit shelf life. *Food Chem.* 259, 234–244.

Kumar, S., Garrick, D. J., Bink, M. C., Whitworth, C., Chagné, D., and Volz, R. K. (2013). Novel genomic approaches unravel genetic architecture of complex traits in apple. *BMC Genomics* 14, 393.

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* 33, 1870–1874.

Larsen, B., Migicovsky, Z., Jeppesen, A. A., Gardner, K. M., Toldam-Andersen, T. B., Myles, S., et al. (2019). Genome-Wide Association Studies in Apple Reveal Loci for Aroma Volatiles, Sugar Composition, and Harvest Date. *The Plant Genome* 12, 180104. doi:10.3835/plantgenome2018.12.0104.

Lau, O. L., Liu, Y., and Yang, S. F. (1986). Effects of fruit detachment on ethylene biosynthesis and loss of flesh firmness, skin color, and starch in ripening Golden Delicious apples. *J. Am. Soc. Hortic. Sci.* 111, 731–734.

Laurens, F., Aranzana, M. J., Arus, P., Bassi, D., Bink, M., Bonany, J., et al. (2018). An integrated approach for increasing breeding efficiency in apple and peach in Europe. *Hortic Res* 5, 11.

Lenth, R. V. (2016). Least-Squares Means: TheRPackagelsmeans. *Journal of Statistical Software* 69. doi:10.18637/jss.v069.i01.

Li, H., Ran, K., Dong, Q., Zhao, Q., and Shi, S. (2020). Cloning, sequencing, and expression analysis of 32 NAC transcription factors (MdNAC) in apple. *PeerJ* 8, e8249.

Longhi, S., Hamblin, M. T., Trainotti, L., Peace, C. P., Velasco, R., and Costa, F. (2013). A candidate gene based approach validates Md-PG1 as the main responsible for a QTL impacting fruit texture in apple (Malus x domestica Borkh). *BMC Plant Biol.* 13, 37.

Longhi, S., Moretto, M., Viola, R., Velasco, R., and Costa, F. (2012). Comprehensive QTL mapping survey dissects the complex fruit texture physiology in apple (Malus x domestica Borkh.). *J. Exp. Bot.* 63, 1107–1121.

Luby, J. J., and Shaw, D. V. (2001). Does Marker-assisted Selection Make Dollars and Sense in a Fruit Breeding Program? *HortScience* 36, 872–879. doi:10.21273/hortsci.36.5.872.

Lü, P., Yu, S., Zhu, N., Chen, Y.-R., Zhou, B., Pan, Y., et al. (2018). Genome encode analyses reveal the basis of convergent evolution of fleshy fruit ripening. *Nat Plants* 4, 784–791.

Mann, H. S., Alton, J. J., Kim, S., and Tong, C. B. S. (2008). Differential expression of cell-wall–modifying genes and novel cDNAs in apple fruit during storage. *J. Am. Soc. Hortic. Sci.* 133, 152–157.

Mathew, I. E., and Agarwal, P. (2018). May the Fittest Protein Evolve: Favoring the Plant-Specific Origin and Expansion of NAC Transcription Factors. *BioEssays* 40, 1800018. doi:10.1002/bies.201800018.

McClure, K. A., Gardner, K. M., Douglas, G. M., Song, J., Forney, C. F., DeLong, J., et al. (2018). A Genome Wide Association Study of Apple Quality and Scab Resistance. *The Plant Genome* 11, 1–14. doi:10.3835/plantgenome2017.08.0075.

Migicovsky, Z., Gardner, K. M., Money, D., Sawler, J., Bloom, J. S., Moffett, P., et al. (2016). Genome to Phenome Mapping in Apple Using Historical Data. *Plant Genome* 9. doi:10.3835/plantgenome2015.11.0113.

Migicovsky, Z., Gardner, K. M., Richards, C., Thomas Chao, C., Schwaninger, H. R., Fazio, G., et al. (2021). Genomic consequences of apple improvement. *Hortic Res* 8, 9.

Migicovsky, Z., Li, M., Chitwood, D. H., and Myles, S. (2017). Morphometrics Reveals Complex and Heritable Apple Leaf Shapes. *Front. Plant Sci.* 8, 2185.

Nakatsuka, A., Murachi, S., Okunishi, H., Shiomi, S., Nakano, R., Kubo, Y., et al. (1998). Differential Expression and Internal Feedback Regulation of 1-Aminocyclopropane-1-Carboxylate Synthase, 1-Aminocyclopropane-1-Carboxylate Oxidase, and Ethylene Receptor Genes in Tomato Fruit during Development and Ripening. *Plant Physiology* 118, 1295–1305. doi:10.1104/pp.118.4.1295.

Nybom, H., Ahmadi-Afzadi, M., Sehic, J., and Hertog, M. (2013). DNA marker-assisted evaluation of fruit firmness at harvest and post-harvest fruit softening in a diverse apple germplasm. *Tree Genetics & Genomes* 9, 279–290. doi:10.1007/s11295-012-0554-z.

Oraguzie, N. C., Iwanami, H., Soejima, J., Harada, T., and Hall, A. (2004). Inheritance of the Md-ACS1 gene and its relationship to fruit softening in apple (Malus × domestica Borkh.). *Theoretical and Applied Genetics* 108, 1526–1533. doi:10.1007/s00122-003-1574-8.

Oraguzie, N. C., Volz, R. K., Whitworth, C. J., Bassett, H. C. M., Hall, A. J., and Gardiner, S. E. (2007). Influence of Md-ACS1 allelotype and harvest season within an apple germplasm collection on fruit softening during cold air storage. *Postharvest Biol. Technol.* 44, 212–219.

Pirona, R., Eduardo, I., Pacheco, I., Da Silva Linge, C., Miculan, M., Verde, I., et al. (2013). Fine mapping and identification of a candidate gene for a major locus controlling maturity date in peach. *BMC Plant Biology* 13, 166. doi:10.1186/1471-2229-13-166.

Ríos, P., Argyris, J., Vegas, J., Leida, C., Kenigswald, M., Tzuri, G., et al. (2017). ETHQV6.3 is involved in melon climacteric fruit ripening and is encoded by a NAC domain transcription factor. *Plant J.* 91, 671–683.

Sadok, I. B., Ben Sadok, I., Tiecher, A., Galvez-Lopez, D., Lahaye, M., Lasserre-Zuber, P., et al. (2015). Apple fruit texture QTLs: year and cold storage effects on sensory and instrumental traits. *Tree Genetics & Genomes* 11. doi:10.1007/s11295-015-0947-x.

Schloerke, B., Crowley, J., Cook, D., Briatte, F., Marbach, M., Thoen, E., et al. (2020). *GGally: Extension to "ggplot2."* Available at: https://CRAN. R-project.org/package=GGally.

Shan, W., Kuang, J.-F., Chen, L., Xie, H., Peng, H.-H., Xiao, Y.-Y., et al. (2012). Molecular characterization of banana NAC transcription factors and their interactions with ethylene signalling component EIL during fruit ripening. *J. Exp. Bot.* 63, 5171–5187.

Shinozaki, Y., Nicolas, P., Fernandez-Pozo, N., Ma, Q., Evanich, D. J., Shi, Y., et al. (2018). High-resolution spatiotemporal transcriptome mapping of tomato fruit development and ripening. *Nat. Commun.* 9, 364.

Song, J., and Bangerth, F. (1996). The effect of harvest date on aroma compound production from "Golden Delicious" apple fruit and relationship to respiration and ethylene production. *Postharvest Biol. Technol.* 8, 259–269.

Sunako, T., Sakuraba, W., Senda, M., Akada, S., Ishikawa, R., Niizeki, M., et al. (1999). An Allele of the Ripening-Specific 1-Aminocyclopropane-1-Carboxylic Acid Synthase Gene (ACS1) in Apple Fruit

with a Long Storage Life. *Plant Physiology* 119, 1297–1304. doi:10.1104/pp.119.4.1297.

Tadiello, A., Longhi, S., Moretto, M., Ferrarini, A., Tononi, P., Farneti, B., et al. (2016). Interference with ethylene perception at receptor level sheds light on auxin and transcriptional circuits associated with the climacteric ripening of apple fruit (Malus x domestica Borkh.). *Plant J.* 88, 963–975.

Tigchelaar, E. C., Tomes, M. L., Kerr, E. A., and Barman, R. J. (1973). A new fruit ripening mutant, non-ripening (nor). *Rep. Tomato Genet. Coop. Tomato Genet. Coop.* 23, 33.

Urrestarazu, J., Muranty, H., Denancé, C., Leforestier, D., Ravon, E., Guyader, A., et al. (2017). Genome-Wide Association Mapping of Flowering and Ripening Periods in Apple. *Front. Plant Sci.* 8, 1923.

U.S. Apple Association *Apple Industry At-a-Glance*. Available at: http://usapple.org/the-industry/apple-industry-at-a-glance/ [Accessed February 11, 2018].

Van Eck, J., Keen, P., and Tjahjadi, M. (2019). Agrobacterium tumefaciens-Mediated Transformation of Tomato. *Methods in Molecular Biology*, 225–234. doi:10.1007/978-1-4939-8778-8_16.

Vincent, J. F. V. (1989). Relationship between density and stiffness of apple flesh. *J. Sci. Food Agric.* 47, 443–462.

Wakasa, Y., Kudo, H., Ishikawa, R., Akada, S., Senda, M., Niizeki, M., et al. (2006). Low expression of an endopolygalacturonase gene in apple fruit with long-term storage potential. *Postharvest Biology and Technology* 39, 193–198. doi:10.1016/j.postharvbio.2005.10.005.

Wang, D., Yeats, T. H., Uluisik, S., Rose, J. K. C., and Seymour, G. B. (2018). Fruit Softening: Revisiting the Role of Pectin. *Trends in Plant Science* 23, 302–310. doi:10.1016/j.tplants.2018.01.006.

Wang, R., da Rocha Tavano, E. C., Lammers, M., Martinelli, A. P., Angenent, G. C., and de Maagd, R. A. (2019). Re-evaluation of transcription factor function in tomato fruit development and ripening with CRISPR/Cas9-mutagenesis. *Scientific Reports* 9. doi:10.1038/s41598-018-38170-6.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer Science & Business Media.

Yang, X., Song, J., Campbell-Palmer, L., Fillmore, S., and Zhang, Z. (2013). Effect of ethylene and 1-MCP on expression of genes involved in ethylene biosynthesis and perception during ripening of apple fruit. *Postharvest Biol. Technol.* 78, 55–66.

Yang, X., Song, J., Du, L., Forney, C., Campbell-Palmer, L., Fillmore, S., et al. (2016). Ethylene and 1-MCP regulate major volatile biosynthetic pathways in apple fruit. *Food Chemistry* 194, 325–336. doi:10.1016/j.foodchem.2015.08.018.

Yue, C., Karina Gallardo, R., Luby, J., Rihn, A., McFerson, J. R., McCracken, V., et al. (2013). An Investigation of U.S. Apple Producers' Trait Prioritization—Evidence from Audience Surveys. *HortScience* 48, 1378–1384.

Zhu, Y., and Barritt, B. H. (2008). Md-ACS1 and Md-ACO1 genotyping of apple (Malus x domestica Borkh.) breeding parents and suitability for marker-assisted selection. *Tree Genet. Genomes* 4, 555–562.