Allelic diversity of *NAC18.1* is a major determinant of fruit firmness and harvest date in

apple (*Malus domestica*)

Trevor H. Yeats[1,6,7], Zoë Migicovsky[2], Sophie Watts[2], Jun Song[3], Charles F. Forney[3], Karen

Burgher-MacLellan[3], Daryl J. Somers[4], Julia Vrebalov[5], James G. Giovannoni[5,6], Jocelyn K. C.

Rose[1]*, Sean Myles[2]*

1. Plant Biology Section, School of Integrative Plant Science, Cornell University, Ithaca, NY

14853 USA

2. Department of Plant, Food and Environmental Sciences, Dalhousie University, Faculty of

Agriculture, Agricultural Campus, Truro, NS, Canada.

3. Agriculture and Agri-Food Canada, AFHRC, Kentville, Nova Scotia, B4N 1J5, Canada

4. Vineland Research and Innovation Centre, Vineland Station, ON, L0R 2E0, Canada

5. Boyce Thompson Institute, Cornell University, Ithaca, NY 14853 USA.

6. United States Department of Agriculture, Robert W. Holley Center, Cornell University, Ithaca,

NY 14853 USA

7. Present address: Plant Breeding & Genetics Section, School of Integrative Plant Science,

Cornell University, Ithaca, NY 14853 USA

* Co-corresponding authors: Sean Myles (seanmichaelmyles@dal.ca) and Jocelyn K. C. Rose

(jr286@cornell.edu).

Highlight:

NAC18.1 is a member of a family of conserved transcriptional regulators of ripening that underlies variation in fruit firmness and harvest date in diverse apple accessions.

Abstract:

Softening is a hallmark of ripening in fleshy fruits, and has both desirable and undesirable implications for texture and postharvest stability. Accordingly, the timing and extent of ripening and associated textural changes are key targets for improving fruit quality through breeding. Previously, we identified a large effect locus associated with harvest date and firmness in apple (*Malus domestica*) using genome-wide association studies (GWAS). Here, we present additional evidence that polymorphisms in or around a transcription factor gene, *NAC18.1*, cause variation in these traits. First, we confirmed our previous findings with new phenotype and genotype data from ~800 apple accessions. In this population, we compared *NAC18.1* to three other ripening-related markers currently used by breeders (*ACS1*, *ACO1*, and *PG1*), and found that the effect of the *NAC18.1* genotype on both traits greatly exceeded that observed for the other markers. By sequencing *NAC18.1* across 18 accessions, we revealed two predominant haplotypes containing the SNP previously identified using GWAS, as well as dozens of additional SNPs and indels in both the coding and promoter sequences. *NAC18.1* encodes a protein with high similarity to the NON-RIPENING (NOR) transcription factor, an early regulator of ripening in tomato (*Solanum lycopersicum*). To test whether these genes are functionally orthologous, we introduced *NAC18.1* transgenes into the tomato *nor* mutant and showed that both haplotypes complement the *nor* ripening deficiency. Taken together, these

results indicate that polymorphisms in *NAC18.1* underlie substantial variation in apple firmness

and harvest time through modulation of a conserved ripening program.

Keywords:

Apple, fruit ripening, NAC-domain transcription factor

Abbreviations:

ABC   Apple Biodiversity Collection

ACS1 1-AMINOCYCLOPROPANE-1-CARBOXYLATE SYNTHASE 1

ACS2 1-AMINOCYCLOPROPANE-1-CARBOXYLATE SYNTHASE 2

ACO1 AMINOCYCLOPROPANE-1-CARBOXYLATE OXIDASE 1

CIELAB   International Commission on Illumination 1976 color space

HRM   High Resolution Melting

LD   Linkage disequilibrium

PG1   POLYGALACTURONASE 1

PG2   POLYGALACTURONASE 2

PSY1 PHYTOENE SYNTHASE 1

REML Restricted Maximum Likelihood

RPL2 RIBOSOMAL PROTEIN L2

USDA United States Department of Agriculture

Introduction

Despite their diverse structure, ontogeny, and biochemical composition, fleshy fruits from a taxonomically broad range of species undergo coordinated ripening programs that have many features in common. Ripening involves numerous physiological and biochemical changes that render the fruit attractive and nutritious for consumption by seed-dispersing animals, or human consumers in the case of cultivated crops. These include the ripening-associated accumulation of sugars, pigments, and flavor or aroma compounds, as well as a loss of flesh firmness due in large part to the controlled modification and depolymerization of cell wall polysaccharides (Wang *et al.*, 2018). These processes are regulated by conserved and convergently evolved networks of transcription factors and hormones, such as ethylene in so-called climacteric fruit (Lü *et al.*, 2018).

Various aspects of fruit ripening have been particularly well studied in tomato (*Solanum lycopersicum*) and the characterization of tomato ripening mutants has revealed a regulatory network, consisting of transcription factors, hormones, and epigenetic modifications (Giovannoni *et al.*, 2017). Of the genes that regulate ripening, the *NON-RIPENING (NOR)* gene appears to exhibit the earliest expression preceding ripening (Shinozaki *et al.*, 2018). *NOR* encodes a NAC domain transcription factor (Giovannoni *et al.*, 2004). This family of transcription factors is one of the largest plant-specific families of transcription factors, with specific members regulating development, defense, and senescence (Mathew and Agarwal, 2018). While all members share a conserved DNA-binding (NAC) domain, specific functional clades are defined in terms of their more variable domains, particularly the C-terminal transcriptional regulatory region. These domains can act directly as transcriptional activators, or can facilitate interaction with other transcription factors in order to fine-tune transcriptional control.

Apple (*Malus domestica*) fruit exhibit extensive variation in the extent and timing of

ripening and ripening-associated softening. This is reflected in three related agronomic traits:

harvest date, firmness, and storability. Harvest date is typically determined by the semi-

quantitative measurement of starch, based on iodine-staining patterns (Blanpied and Silsby,

1992). Firmness can also be used as a measure of ripening, although apples soften less than many

other fleshy fruits, and are more noted for their characteristic and desirable "crisp" texture.

Importantly, postharvest storability is highly correlated with fruit firmness (McClure *et al.*, 2018;

Ornelas-Paz *et al.*, 2018). Thus, firmness is typically a desirable fruit quality trait in apple,

although it is often associated with later harvest dates that can be undesirable if the growing

season is short (Migicovsky *et al.*, 2016).

Due to a prolonged juvenile phase, it is particularly challenging to evaluate fruit quality

traits in the context of an apple breeding program. As a result, in recent years considerable effort

has been invested in developing molecular markers that can be used to select for fruit quality

traits at the seedling stage. In particular, three markers have been proposed for predicting

firmness. The first is *1-AMINOCYCLOPROPANE-1-CARBOXYLATE SYNTHASE 1* (*ACS1*),

which encodes the ripening-associated isoform of an ethylene biosynthesis gene. The *ACS1-2*

allele contains a retrotransposon insertion thought to confer low ethylene production and longer

shelf life in Fuji and other apple cultivars homozygous for this allele (Sunako *et al.*, 1999;

Harada *et al.*, 2000). The second is a marker for a gene corresponding to another enzyme

involved in ethylene biosynthesis, *AMINOCYCLOPROPANE-1-CARBOXYLATE OXIDASE 1*

(*ACO1*; Costa *et al.*, 2005). Finally, a marker has been developed for alleles of

POLYGALACTURONASE 1 (PG1; Costa *et al.*, 2010). This gene encodes an enzyme that

hydrolyzes pectin polysaccharides and has been implicated in apple fruit firmness by RNAi-

based gene silencing experiments (Atkinson *et al.*, 2012).

In a previous study, we conducted GWAS on a range of traits in apple using genotype

data and historical phenotype records for 689 accessions of the United States Department of

Agriculture (USDA) apple germplasm collection (Migicovsky *et al.*, 2016). Our analysis

revealed a single nucleotide polymorphism (SNP) on chromosome 3 within the coding sequence

of the *NAC18.1* gene that was significantly associated with firmness and harvest date. This SNP

results in an aspartate (D) to tyrosine (Y) mutation at position 5 of the NAC18.1 amino acid

sequence and we refer to this SNP as D5Y (Migicovsky *et al.*, 2016). Subsequently, GWAS of

two European apple germplasm collections confirmed the association between *NAC18.1* and

ripening time (Urrestarazu *et al.*, 2017; Larsen *et al.*, 2019). NAC transcription factors have also

been linked to harvest date in other Rosaceae species, including peach (*Prunus persica;* Pirona *et*

al., 2013) and apricot (*P. armeniaca;* García-Gómez *et al.*, 2019).

Here we report an extension of our previous findings using three approaches. First, we

conducted additional phenotyping and genotyping of apple germplasm in order to compare the

predictive power of the *NAC18.1* D5Y SNP with other proposed firmness/harvest date markers.

Secondly, we sequenced the *NAC18.1* gene from a subset of apple cultivars in order to discover

additional polymorphisms in linkage disequilibrium (LD) with D5Y. Finally, we used

heterologous expression of the *NAC18.1* gene in the tomato *non-ripening (nor)* mutant to test

whether it functions as a component of a conserved fruit ripening program.

Materials and Methods

Apple Firmness and Harvest Date

The Apple Biodiversity Collection (ABC) is an apple orchard located in Kentville, Nova

Scotia, which contains 1,113 accessions. It was first established in spring 2012, when trees were

budded onto M.9 rootstocks. In fall 2012, the trees were uprooted and kept in cold storage,

before planting in spring 2013. The trees were planted in an incomplete block design, which

includes 1 of 3 standards per grid, allowing for correction of positional effects using a REstricted

Maximum Likelihood (REML) model, described in Migicovsky *et al.* (2018).

In 2017, we evaluated harvest date for 1,348 trees and fruit firmness for 1,328 trees

within the ABC orchard. Due to the diversity of apples within the collection, a variety of

methods were used to determine the appropriate time to harvest. First, we observed if the tree

had dropped fruit or, for red apples, if the fruit were a deep red color. Next, a sample apple was

taken from each tree and touched to assess firmness, tasted to assess starch and sweetness, cut in

half to check browning of seeds, and then sprayed with iodine solution to assess starch content

(Blanpied and Silsby, 1992).

When fruit were determined to be mature, they were harvested and evaluated for

firmness. We recorded the firmness ($kg\ cm^{-2}$) of 5 fruit per tree using a penetrometer with a 1 cm

diameter (Fruit Texture Analyzer, GS-14, Güss Manufacturing). A small section of skin was

removed using a vegetable peeler, and each fruit was placed on the penetrometer platform so that

the piston entered the middle of the apple where the skin had been removed. Data were

automatically recorded into a spreadsheet.

Due to the number of trees, harvesting the ABC orchard often lasted more than one day.

Therefore, differences in harvest date within a week were simply due to the time required to

harvest, and the harvest date for each tree was recorded as the Monday of that week. We used the

"*lmer*" function in the R package *lme4* (Bates *et al.*, 2015) to fit a REML model for harvest and

firmness data. Next, we calculated the least squares mean using the "*lsmeans*" function in the

lsmeans R package (Lenth, 2016), resulting in one value per unique accession. We fit this model

for 866 unique accessions with harvest dates and 863 accessions with firmness measurements.

High Resolution Melting (HRM) Genotyping

DNA was extracted using silica columns from fresh leaf tissue collected from the ABC

orchard, quantified using PicoGreen (Thermo) and normalized to a concentration of 20 ng μL^{-1}.

Genotyping was conducted using PCR and high resolution melting (HRM) on a LightScanner

HR384 (BioFire). Primers are listed in Supplemental Table S1.

Genotyping of the PG1 SNP marker was based on the GenBank sequence L27743.1,

where the G allele at position 437 is unfavorable and the T allele is favorable (Costa *et al.*,

2010). Three allelic combinations of the observed indel, ACS1-1/1, 1-1/2 and 1-2/2, have been

associated with high, medium, and low ethylene production, respectively (Sunako *et al.*, 1999;

Harada *et al.*, 2000; Oraguzie *et al.*, 2004; Costa *et al.*, 2005). The exact position of the indel is

from ACS1-2 (GenBank: AB010102.1) 1,320 to 1,483 bp (163 bp), and from ACS1-1

(GenBank: AY062129.1) 4,500 to 4,525 bp (25 bp). The size difference between alleles is 138

bp. The ACO marker involves an unfavorable 62 bp insertion in the third intron of ACO1 (Costa

et al., 2005). The third intron spans from 1,083 bp to 1,300 bp of the GenBank sequence

Y14005.1 and the indel is found from position 1,297 bp to position 1,358 bp. The D5Y mutation

in *NAC18.1* is a nonsynonymous SNP at position 30,698,039 on chromosome 3, according to

reference genome version GDDH13 v1.1 (https://iris.angers.inra.fr/gddh13/) and is associated

with both harvest date and firmness (Migicovsky *et al.*, 2016). The desirable C allele encodes an

aspartic acid (D) at the fifth amino acid position of the NAC18.1 protein, while the undesirable A

allele encodes a tyrosine (Y).

Marker-phenotype associations

We evaluated the ability of markers at NAC18.1, ACO1, ACS1, and PG1 to predict both

harvest date and firmness of accessions in the ABC orchard. The number of accessions with

phenotype and genotype information ranged from 754 to 852 depending on the trait/marker

combination. The association test was conducted using Spearman's rank correlation test. We

visualized results using the "*geom_boxplot*" function in *ggplot2* in R (Wickham, 2016). Lastly,

we tested which mode of inheritance best fit the data using SNPstats (Solé *et al.*, 2006), and

selected the model with the lowest Akaike Information Criterion value.

***Sequencing of* NAC18.1**

DNA was isolated from leaves sampled from 18 apple accessions growing in Geneva,

NY. A 2.3 kb amplicon including the *NAC18.1* gene and ~800 bp of upstream sequence was

amplified by PCR using primers NAC18F2 and NAC18R2 (Supplemental Table S1) and

Phusion® High-Fidelity PCR Master Mix with HF Buffer (NEB). PCR product size and purity

was confirmed by agarose gel electrophoresis, and the remaining product was purified using

DNA Clean & Concentrator kit (Zymo Research). The resulting DNA fragment was cloned into

the plasmid pMiniT 2.0 and transformed into *E. coli* using the NEB® PCR Cloning Kit (NEB).

Individual colonies were selected for complete sequencing of the cloned amplicon using

the primers NAC18F2, NAC18F3, NAC18F4, NAC18R1, and NAC18R2 (Supplemental Table

S1). For accessions homozygous for the D5Y SNP, the NAC18.1 amplicon from a single clone

was sequenced. For heterozygous accessions, two clones representing each D5Y allele were

selected based on partial sequencing of the D5Y region, followed by complete sequencing of the

2.3 kb amplicon, as described above. The nucleotide sequences were aligned by MUSCLE

(Edgar, 2004) and used to construct a maximum-likelihood phylogenetic tree in MEGA7 (Kumar

et al., 2016).

Molecular cloning and plant transformation

The hypothetical coding sequences (CDSs) corresponding to the consensus sequence of

the "A" and "C" haplotypes of *NAC18.1* were synthesized as a double-stranded DNA (gBlock)

by IDT, with 20 bp of flanking sequence added to facilitate Gibson Assembly of the *NAC18.1*

sequences between the *Asc*I and *Pac*I restriction sites of pMDC32 (Curtis and Grossniklaus,

2003). Constructs were assembled using the NEBuilder HiFi DNA Assembly Cloning Kit (NEB)

and their integrity verified by Sanger sequencing. Plasmids were transformed into

Agrobacterium tumefaciens, which was used to transform tomato cotyledon explants (Van Eck *et*

al., 2019) derived from the tomato *nor* mutant in the cv. Ailsa Craig background (LA3770,

Tomato Genetics Resource Center, https://tgrc.ucdavis.edu/). A total of 10 T0 plants were

recovered per construct and plants derived from two independent transformation events per

construct were selected for further characterization in the T1 generation.

Transgenic tomato characterization

Fruit were harvested when visually ripe, or in the case of the *nor* mutant, at the equivalent

age as ripe cv. Ailsa Craig fruit (the near isogenic wild-type control), as determined by tagging

of flowers at anthesis. The color of the fruit surface was measured using a CR-400 Chroma

Meter (Konica Minolta), and fruit were weighed, photographed, and dissected. Pericarp tissue

was frozen in liquid nitrogen and stored at -80˚C. Frozen tissue was ground to a fine powder and

RNA extracted by a modified version of Chang et al. (1993). Briefly, approximately 400 mg of

tissue was added to a preheated (80˚C) two-phase system consisting of 500 µL of water-saturated

phenol and 500 µL of extraction buffer (100 mM Tris [pH 8.0], 25 mM EDTA, 2 M NaCl, 2%

CTAB, 2% PVP, and 2% [v/v] beta-mercaptoethanol). The mixture was vortexed and incubated

for 5 min at 65˚C and then cooled to room temperature before extracting and precipitating RNA,

as previously described (Chang *et al.*, 1993).

RNA was treated with RNase-free DNase (Promega) and used for cDNA synthesis with

RNA to cDNA EcoDry™ Premix with Oligo dT primer (Takara Bio). The cDNA was used as

template for quantitative PCR using Luna Universal qPCR Master Mix (NEB) and a Viia7 real-

time PCR instrument (Life Technologies/ABI). Gene-specific primers are listed in Supplemental

Table S1. Quantification used the ΔC_t method with *RIBOSOMAL PROTEIN L2* (*RPL2*) as a

reference gene, and statistical significance of the ΔC_t values was tested by a one-way ANOVA

followed by Tukey's HSD test.

Results

Evaluation of markers for firmness and harvest date in apple

We sought to re-evaluate the ability of D5Y and other published markers to predict fruit

firmness and harvest date using new phenotype and genotype data: specifically, we aimed to

address some experimental limitations of our previous work. First, the GWAS that identified

D5Y, and failed to find associations for previously published markers, made use of only 8,000

SNPs (Migicovsky *et al.*, 2016). Given the rapid LD decay in apple, our low SNP density was

insufficient to conclude that we had exhaustively searched the apple genome for loci involved in

firmness and harvest date (Migicovsky *et al.*, 2016). Secondly, the historic phenotype data we

used were imprecise. For example, firmness was recorded as either "firm" or "soft" rather than

as a biomechanical measurement, and harvest date was recorded as simply "early", "mid", or

"late". These factors would have limited the detection of additional loci that underlie firmness

and/or harvest date.

In order to directly compare the NAC18.1 D5Y marker that we identified with previously

published markers, we developed HRM genotyping assays for the D5Y SNP, as well as markers

at *ACS1, ACO1,* and *PG1*. We used these assays to genotype accessions from the ABC orchard

(Supplementary Table S2). While this orchard largely consists of clones of individuals present in

the population used in our previous GWAS (i.e. the USDA germplasm collection), it also

contains a number of additional accessions. More importantly, for the purposes of phenotyping,

the ABC orchard is planted in an incomplete block design that allows for modeling of location

effects using REML methods (Migicovsky *et al.*, 2018). Using this approach, we generated

firmness and harvest date measures for over 800 unique accessions for which HRM genotype

data were available.

We evaluated the mode of inheritance for each genetic marker and phenotype

combination and found that, while the effects of the markers at *ACS1, ACO1,* and *PG1* were all

dominant, the D5Y marker at *NAC18.1* had a codominant effect. As a result, accessions that

were heterozygous for D5Y had firmer fruit (an increase of 1.2 kg cm^{-2}) and a later harvest date

(20.13 days) than those that were homozygous for the 'soft' A allele. The combined effect of two

C alleles was notably stronger than the effect of a single C allele, increasing firmness by 2.24 kg

cm^{-2} and harvest date by a month (30.52 days), when compared to AA genotypes. The difference

between firm/late and soft/early genotypes for D5Y was at least 4 times higher for firmness and

3 times higher for harvest date than for the markers at *ACS1, ACO1,* and *PG1* (Figure 1).

Figure 1. Distribution of phenotypic variation across genotypic classes for markers at
*NAC18.1***,** *PG1***,** *ACO1***, and** *ACS1***.** A. Boxplots of variation in firmness across each marker B.
Boxplots of variation in harvest date across each marker. The lower and upper hinges of the

boxplots correspond to the first and third quartiles. The putative inheritance model, allele effect,

correlation (rho) and *p*-value determined using a Spearman's rank correlation test, and the

number of accessions within each genotypic class are listed.

To evaluate the distribution of firmness/harvest date markers across commercial

cultivars, we present the genotypes for all markers genotyped here in nine of the top ten apple

cultivars sold in the USA (Table 1). All nine cultivars were homozygous for the favorable (firm)

C allele of *NAC18.1*, while only 2 to 4 of the cultivars had homozygous favorable genotypes for

the other markers. Among the top cultivars, only 'Fuji', released in 1962, was homozygous for

favorable alleles at all markers, while 'McIntosh', initially discovered in 1811 and commercially

released in 1870, was homozygous for undesirable alleles at all markers except for the *NAC18.1*

marker.

Table 1. Genotypes of markers for 9 of the 10 most widely sold apple cultivars in the United States.

		Genotypes[a]			
Release Date	Cultivar	NAC18.1	ACS1	ACO1	PG1
1868	Granny Smith	+	±	−	+
1870	McIntosh	+	−	−	−
1895	Red Delicious	+	±	+	±
1914	Golden Delicious	+	±	-	±
1952	Braeburn	+	±	n.d.	±
1962	Fuji	+	+	+	+
1974	Gala	+	+	±	±
1990	Cripps Pink	+	±	−	+
1991	Honeycrisp	+	±	−	+

[a] Genotypes are coded as + for homozygous, favorable; − for homozygous, unfavorable; ± for heterozygous.

Resequencing of NAC18.1

Since the D5Y SNP is predicted to result in an amino acid change in the NAC18.1 protein and LD decay is generally high in apple (r^2 decayed to < 0.2 within 100 bp), it is possible that D5Y is causative for the early/firm phenotype (Migicovsky *et al.*, 2016). However, GBS sequence data are too sparse to rule out additional polymorphisms in LD with D5Y contributing to the phenotype. To better understand allelic diversity of *NAC18.1*, we selected 6 cultivars each of the D5Y A/A, A/C, and C/C genotypes, based on our HRM genotyping data. For homozygous

cultivars, single alleles were sequenced, while for the heterozygous samples we resequenced

both alleles, resulting in *NAC18.1* sequences from 24 haplotypes. These were compared to two

reference sequences from the GDDH13 v1.1 apple genome (Daccord *et al.*, 2017):

Md03g1222600 and *Md11g1239900*, representing *NAC18.1* and its closest paralog, respectively.

In addition to confirming the expected D5Y genotype in all individuals, a number of

additional SNPs and indels were revealed within both coding and non-coding regions of the

NAC18.1 allele. Multiple sequence alignment and subsequent phylogenetic analysis indicated

two major clades corresponding to the D5Y A and C genotypes (Figure 2A). SNPs and small

indels were observed throughout the sequenced region. Notably, in addition to the D5Y amino

acid change, several additional amino acid changes were observed between sequences from D5Y

genotypes and the reference sequence of NAC18.1. For example, near the site of the D5Y

polymorphism, all "A" haplotypes also had a 12 nucleotide insertion that introduced the amino

acid sequence QPQP (Figure 2B).

Taken together, these results indicate that the *NAC18.1* locus exhibits a number of

polymorphisms that are in strong LD with each other across the 2.3 kb that we sequenced. Many

of these are candidates for causality, as they affect either the coding sequence or promoter of the

gene. On the other hand, it is possible the signal at D5Y is driven by a causal variant outside of

the 2.3kb region sequenced here that acts completely independently from *NAC18.1*. This led us

to look for further evidence of the involvement of NAC18.1 in the ripening process in apple.

Figure 2. Polymorphisms in LD with the D5Y SNP. A. Maximum likelihood phylogenetic tree

of 24 *NAC18.1* sequences from 18 apple cultivars and the reference genome (GDDH13 v1.1)

sequences of *NAC18.1* (*Md03g1222600*) and its closest homolog (*Md11g1239900*). A graphical

representation of the 2.3 kb multiple sequence alignment is also shown. B. Amino acid sequence

alignment of the N-terminal region of NAC18.1, illustrating additional variation in the coding

sequence strongly correlated with the D5Y variant. The complete amino acid sequence alignment

is given in Supplemental Figure S1.

Transgenic complementation of the tomato* non-ripening (nor) *mutant

A BLAST search of predicted proteins from the ITAG 3.20 tomato genome

(https://solgenomics.net/) using NAC18.1 as a query identified only two sequences matching the

full length of the query: *Solyc10g006880* (*NOR*) and *Solyc07g063420* (*NOR-LIKE 1*). To test the

functional conservation of *NAC18.1* and *NOR*, we introduced transgenic constructs individually

conferring constitutive expression of each of the *NAC18.1* haplotype CDSs into the tomato *nor*

mutant. Two independent lines for each construct were characterized in the T1 generation with

respect to their ability to rescue the ripening deficiency of the *nor* mutant. In contrast to *nor*, all

four lines changed color at maturity, although internal color change did not occur to the same

extent as observed in a WT control (Figure 3A). To complement this qualitative phenotypic

assessment, we also conducted quantitative colorimetry of the surface of fruits (Figure 3B). Fruit

from all the transgenic lines exhibited a significant increase in the *a** (green-red) component of

CIELAB (International Commission on Illumination 1976) color space relative to the *nor*

mutant, although only *NAC18.1^A #6* achieved similar *a** levels to WT.

The different degrees to which ripe fruit color was restored in the transgenic plants might

be a consequence of either the different alleles of *NAC18.1*, or of different levels of transgene

expression in each line. To address this, we analyzed the expression level of *NAC18.1* using

qRT-PCR primers designed to target both *NAC18.1* alleles. Expression levels of the NAC18.1

transgene were not statistically different ($p > 0.5$) between each independent line (Figure 4A).

Next, we measured the expression level of several genes associated with tomato ripening

physiology: *PHYTOENE SYNTHASE 1 (PSY1)* encodes an early enzyme of carotenoid synthesis

responsible for the production of red pigments during fruit ripening (Fray and Grierson, 1993).

Consistent with the green color of mature *nor* fruit*,* expression of *PSY1* is impaired in the *nor*

mutant (Osorio *et al.*, 2011); *POLYGALACTURONASE 2* (*PG2*) encodes an enzyme catalyzing

pectin depolymerization associated with fruit softening (Biggs and Handa, 1989); and *1-*

AMINOCYCLOPROPANE-1-CARBOXYLATE SYNTHASE 2 (*ACS2*) encodes an enzyme that

synthesizes 1-aminocyclopropane-1-carboxylate, the immediate precursor of ethylene

(Nakatsuka *et al.*, 1998).

Expression of all three marker genes was enhanced in the *NAC18.1* transgenic lines

relative to the *nor* mutant control, although not to the same extent as observed in WT ripe fruit

(Figure 3B-D). In contrast to the consistent level of NAC18 expression observed in each line, the

marker genes were more variable in their expression level between lines. A similar pattern was

observed for all marker genes, with the *NAC18.1^C* #9 line showing the smallest induction of

marker gene expression relative to *nor*. In the case of *PG2* and *ACS2*, the difference in

expression in *NAC18.1^C* #9 was not statistically significant relative to *nor* (p = 0.09 and 0.18,

respectively). Consistent with these results, fruit from this line also exhibited the lowest amount

of color development (Figure 3B). Taken together, these results indicate that a canonical ripening

program can be induced in the tomato *nor* mutant through the heterologous expression of either

apple *NAC18.1* allele.

Figure 3. Transgenic complementation of the tomato *nor* mutant. A. Mature fruits of the tomato *nor* mutant and four independent T1 transgenic lines constitutively expressing either of two alleles of the *NAC18.1* transgene, *NAC18.1^C* and *NAC18.1^A* and isogenic WT control (cv.

Ailsa Craig). B. Quantitative colorimetry of the fruit surface of *nor,* the transgenic fruit and WT

fruit. The *a** component (green-red axis) is shown, with the mean ± SE superimposed in black

over the raw values in gray ($n = 5$). Genotypes not sharing a letter (a-e) are statistically distinct

by one-way ANOVA and Tukey's HSD test ($p < 0.05$).

Discussion

Genomics-assisted breeding has tremendous potential in perennial crops like apple,

where a lengthy juvenile phase and large plant size make phenotyping at the adult stage time-

consuming and expensive. However, the genetic markers used for culling progeny at the seedling

stage must accurately predict the trait of interest at the adult stage in order for genomics-assisted

breeding to be effective (Luby and Shaw, 2001). Firmness and harvest date, which are highly

correlated, are key breeding targets in apple, and markers in three genes (*ACS1*, *ACO1*, and *PG1*)

are widely used in genomics-assisted apple breeding programs to predict firmness. However,

recent GWAS analyses for firmness and harvest date have failed to identify significant

associations at any of these loci, but rather have uncovered a single major effect locus at the

NAC18.1 gene (Migicovsky et al. 2016; McClure et al. 2018; Larsen et al. 2019). Based on the

results of this current study, we propose several possible explanations for these findings.

First, the markers in *ACS1*, *ACO1*, and *PG1* were identified using linkage mapping in bi-

parental crosses, which only captured variation segregating in the progeny of those crosses.

These mapping studies generally made use of commercially successful cultivars, and here we

show that the most commercially successful apple cultivars in the USA are all homozygous for

the desirable *NAC18.1* alleles (Table 1). Thus, the *NAC18.1* alleles we identified via GWAS may

not have segregated in the progeny of the crosses used to discover the markers at *ACS1, ACO1*

and *PG1*. This may be the primary reason *NAC18.1* was not identified using traditional linkage

mapping approaches. Second, the most extensive GWAS we have performed to date may have

failed to detect association signals at *ACS1, ACO1* and *PG1* since the SNP density used (~1 SNP

every 100kb) provided poor genome-wide coverage, given the rapid LD decay observed in our

GWAS population (r^2 decayed to < 0.2 within 100 bp). For example, the SNP closest to *ACO1* in

our GWAS was >100 kb away and was thus unlikely to be in LD with genetic variants within

ACO1 (Migicovsky et al. 2016). Given these limitations, we examined the value of each of these

markers to predict firmness and harvest date in a diverse collection of over 750 unique apple

accessions.

The correlation test between each of the markers and the two phenotypes, harvest date

and firmness, revealed statistically significant ($P < 0.05$) associations in every case except one

(Figure 1). ACO1 was not correlated with firmness in our data, which is consistent with a recent

study that found that a marker for ACO1 did not explain any of the variation in crispness or

firmness across 207 accessions from New Zealand's apple breeding program (Chagne et al.,

2019). Our observation that most of the marker-trait tests that we performed resulted in

significant associations is not surprising: many, if not most, markers genome-wide are expected

to be correlated with harvest date and firmness since the genetic structure of our apple population

is strongly correlated with these traits. For example, without correcting for the effects of

population structure, 39% and 17% of genome-wide SNPs were significantly associated (P <

0.05) with harvest date and firmness, respectively, in the USDA collection, which is genetically

identical to most of the population studied here (Migicovsky et al, 2016). Because we performed

single-marker association tests in the present study, we were unable to account for population

structure using genome-wide markers as is customarily done when performing GWAS. Thus, we

were only able to compare the relative power of each marker to predict firmness and harvest

date. We found that the D5Y marker in *NAC18.1* had a 3 to 14 times greater effect on firmness

and harvest than the markers in *ACO1*, *ACS1* and *PG1* (Figure 1). The superior predictive power

of the NAC18.1 marker suggests that it has more potential to advance genomics-assisted

breeding for firmness and harvest date compared to markers currently in use. Results from our

forthcoming GWAS that include all of these markers and properly account for the effects of

relatedness and population structure will be necessary to definitively conclude which, if any, of

these traditionally used markers predict firmness and/or harvest date in diverse collections of

apple.

While the D5Y mutation in *NAC18.1* is a strong functional candidate variant, we

discovered numerous DNA sequence variants in NAC18.1, often in perfect LD with D5Y, that

are also putatively functional (Figure 2). The magnitude of DNA polymorphism at this locus,

and the multitude of putatively causal variants, are consistent with the results of Larsen et al.

(2019), who found 18 SNPs and 2 indels within *NAC18.1* across 11 apple accessions. Given the

pattern of LD we observed across *NAC18.1* (Figure 2), we considered the possibility that the

GWAS signals at *NAC18.1* are in fact driven by a causal variant outside of the *NAC18.1* coding

sequence that acts independently of the *NAC18.1* gene. Since *NAC18.1* is a homolog of the well-

known ripening gene *NOR* in tomato, we used tomato as a model to explore the role of *NAC18.1*

in the ripening process.

We observed enhanced ripening in the *nor* mutant following heterologous expression of

either of the *NAC18.1* alleles. Notably, the synthesis of carotenoids was restored in the

transgenic fruit, resulting in visual color change that approached the level of WT at the fruit

surface (Figure 3B), although dissection of fruit revealed incomplete internal color change in all

lines (Figure 3A). Visual color change was consistent with the expression of the carotenoid

biosynthetic gene *PSY1* (Figure 4B). Gene expression analysis further confirmed the induction of

genes involved in ripening-associated cell wall remodeling (Figure 4C) and ethylene synthesis

(Figure 4D). Although the expression level of the *NAC18.1* transgene was comparable across

transgenic lines (Figure 4A), the level of ripening marker induction in the distinct lines was

significantly different (Figure 4B-D). We had hypothesized that an allele-specific trend in our

transgenic experiments would indicate that coding sequence variants are responsible for the

association of *NAC18.1* with variation in apple ripening. While our results do not indicate such a

trend, they do confirm that both alleles of *NAC18.1* encode functional proteins that can promote

ripening.

Figure 4. Ripening marker gene expression in transgenic tomatoes. Gene expression was evaluated as the difference in threshold cycle (ΔC_t, $\log_2$ scale) by qRT-PCR using *RPL2* as a reference gene. The mean ± SE is superimposed in black over the raw values in gray ($n = 5$). Genotypes not sharing a letter (a-d) are statistically distinct by one-way ANOVA and Tukey's HSD test ($p < 0.05$). A. The *NAC18.1* transgene. B. *PSY1*, encoding the phytoene synthase carotenoid biosynthetic gene. C. *PG2*, encoding the major ripening-associated polygalacturonase involved in pectin hydrolysis. D. *ACS2*, encoding 1-aminocyclopropane-1-carboxylate synthase, a ripening-associated isoform of the ethylene biosynthesis pathway.

In tomato, *nor* was first identified as a spontaneous mutation in an heirloom cultivar (Tigchelaar *et al.*, 1973). While this *nor* allele exhibits recessive behavior and has been assumed to be a complete loss-of-function, due to a 2 nt deletion resulting in a truncated protein (Giovannoni *et al.*, 2004), it was recently demonstrated that it is actually a dominant negative allele (Wang *et al.*, 2019; Gao *et al.*, 2019). These studies used CRISPR/Cas9 to generate *bona fide* null alleles of *nor* , which showed evidence of ripening relative to the spontaneous *nor* allele, although to a lesser extent than WT fruit. In light of these studies, we speculate that the action of the *NAC18.1* transgenes was likely attenuated by the dominant-negative activity of the spontaneous *nor* allele used in our work. Further heterologous characterization of *NAC18.1* would likely benefit from the use of null *nor* mutants, or double mutants of *nor* and *nor-like 1* (Gao *et al.*, 2018). This may allow for more precise quantitative comparisons between alleles in order to resolve whether differences in the coding sequence of *NAC18.1* confer different degrees of ripening.

Although we are unable to conclude whether polymorphisms in coding or regulatory

sequences affect the activity of *NAC18.1* in apple, an increasing body of evidence generated

using tomato as a model system indicates that coding sequence polymorphisms of *NOR* can

influence firmness and timing of ripening. For example, the Portuguese Alcobaça cultivar of

tomato has firm fruit, delayed ripening, and long shelf life that is conferred by the *alcobaça*

mutation, and a recent study has shown that *ALCOBACA* is allelic with *NOR*, and that the

alcobaça allele of *NOR* contains a valine to aspartate mutation at position 106 within the NAC

domain (Kumar *et al.*, 2018). An additional complete loss of function allele of *NOR* was also

found in another long shelf life tomato cultivar (Kumar *et al.*, 2018).

Our analysis of *NAC18.1* sequences revealed a number of mutations that are candidates

for causal association with ripening phenotypes. For example, we observed an insertion of four

amino acids (QPQP) 11 amino acids downstream of the D5Y mutation in all A haplotypes

upstream of the NAC DNA binding domain. Glutamine-rich sequences are common in

eukaryotic transcription factors, and polymorphisms in these motifs have been shown to alter the

activity of transcriptional activators (Atanesyan *et al.*, 2012). On the other hand, we also

observed several SNPs resulting in amino acid changes within the DNA binding domain and C-

terminal transcriptional activator domain of NAC18.1. Clearly, resolution of a causal

polymorphism within the coding sequence or promoter of *NAC18.1* will require substantial

additional effort. In this regard, we are resequencing the *NAC18.1* locus in our complete

diversity panel in order to identify intragenic recombination that is informative in combination

with our long-term phenotyping of the collection.

In conclusion, the results presented here provide evidence of the superiority of GWAS for

the identification of genetic markers associated with firmness and harvest date in diverse

germplasm. By validating NAC18.1 as a functional candidate for apple fruit ripening, we lay the

groundwork for future efforts to compare the effect of different NAC18.1 haplotypes and

determine the precise causal variant(s) underlying this agriculturally important gene. Ultimately,

we envision that the identification of precise causal variants for ripening in apple could be

exploited using gene editing for fast and efficient introgression of this desirable trait.

Acknowledgments

We thank Susan Brown (Cornell University), Benjamín Gutiérrez (USDA-ARS Plant

Genetics Resource Unit), and C. Thomas Chao (USDA-ARS Plant Genetics Resource Unit) for

apple leaf tissue samples from orchards of the Cornell apple breeding program and the USDA

apple germplasm collection. Transgenic tomato plants were generated by the Boyce Thompson

Institute Biotechnology Center (https://btiscience.org/). ZM was supported by National Science

Foundation Plant Genome Research Program 1546869. TY and JR were supported by a grant

from the USDA-ARS Multistate (NE-1336) project, NYC-184821. This research was supported

in part by funding the National Sciences and Engineering Research Council of Canada (SM) and

A-Base funding (NOI-1238) from Agriculture and Agri-Food Canada (JS). The authors

gratefully acknowledge the farm services team at the Kentville Research and Development

Centre for maintaining the apple collection studied here.

Supplemental Figure Legends

Supplemental Figure S1. Complete amino acid sequence alignment of 24 NAC18.1

haplotypes.

Supplemental Table Legends

Supplemental Table S1. PCR primers used in this work.

Supplemental Table S2. Genotypes of *NAC18.1* (D5Y), *ACS1*, and *ACO1* across 866 apple

accessions.

References

Atanesyan L, Günther V, Dichtl B, Georgiev O, Schaffner W. 2012. Polyglutamine tracts as

modulators of transcriptional activation from yeast to mammals. Biological Chemistry **393**, 63–

70.

Atkinson RG, Sutherland PW, Johnston SL, Gunaseelan K, Hallett IC, Mitra D, Brummell

DA, Schröder R, Johnston JW, Schaffer RJ. 2012. Down-regulation of

POLYGALACTURONASE1 alters firmness, tensile strength and water loss in apple (Malus x

domestica) fruit. BMC Plant Biology **12**, 129.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using

lme4. Journal of Statistical Software **67**, 1–48.

Biggs MS, Handa AK. 1989. Temporal Regulation of Polygalacturonase Gene Expression in

Fruits of Normal, Mutant, and Heterozygous Tomato Genotypes. Plant Physiology **89**, 117–125.

Blanpied GD, Silsby KJ. 1992. Predicting Harvest Date Windows for Apples.

Chang S, Puryear J, Cairney J. 1993. A simple and efficient method for isolating RNA from

pine trees. Plant Molecular Biology Reporter **11**, 113–116.

Costa F, Peace CP, Stella S, Serra S, Musacchi S, Bazzani M, Sansavini S, Van de Weg

WE. 2010. QTL dynamics for fruit firmness and softening around an ethylene-dependent

polygalacturonase gene in apple (Malus×domestica Borkh.). Journal of Experimental Botany **61**,

3029–3039.

Costa F, Stella S, Van de Weg WE, Guerra W, Cecchinel M, Dallavia J, Koller B, Sansavini

S. 2005. Role of the genes Md-ACO1 and Md-ACS1 in ethylene production and shelf life of

apple (Malus domestica Borkh). Euphytica **141**, 181–190.

Curtis MD, Grossniklaus U. 2003. A Gateway Cloning Vector Set for High-Throughput

Functional Analysis of Genes in Planta. Plant Physiology **133**, 462–469.

Daccord N, Celton J-M, Linsmith G, *et al.* 2017. High-quality *de novo* assembly of the apple

genome and methylome dynamics of early fruit development. Nature Genetics **49**, 1099–1106.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Research **32**, 1792–1797.

Fray RG, Grierson D. 1993. Identification and genetic analysis of normal and mutant phytoene

synthase genes of tomato by sequencing, complementation and co-suppression. Plant Molecular

Biology **22**, 589–602.

Gao Y, Wei W, Zhao X, *et al.* 2018. A NAC transcription factor, NOR-like1, is a new positive

regulator of tomato fruit ripening. Horticulture Research **5**, 75.

Gao Y, Zhu N, Zhu X, *et al.* 2019. Diversity and redundancy of the ripening regulatory

networks revealed by the fruitENCODE and the new CRISPR/Cas9 CNR and NOR mutants.

Horticulture Research **6**, 39.

García-Gómez BE, Salazar JA, Dondini L, Martínez-Gómez P, Ruiz D. 2019. Identification

of QTLs linked to fruit quality traits in apricot (Prunus armeniaca L.) and biological validation

through gene expression analysis using qPCR. Molecular Breeding **39**, 28.

Giovannoni J, Nguyen C, Ampofo B, Zhong S, Fei Z. 2017. The Epigenome and

Transcriptional Dynamics of Fruit Ripening. Annual Review of Plant Biology **68**, 61–84.

Giovannoni J, Tanksley S, Vrebalov J, Noensie F. 2004. NOR gene compositions and

methods for use thereof.

Harada T, Sunako T, Wakasa Y, Soejima J, Satoh T, Niizeki M. 2000. An allele of the 1-

aminocyclopropane-1-carboxylate synthase gene (Md-ACS1) accounts for the low level of

ethylene production in climacteric fruits of some apple cultivars. Theoretical and Applied

Genetics **101**, 742–746.

Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis

Version 7.0 for Bigger Datasets. Molecular Biology and Evolution **33**, 1870–1874.

Kumar R, Tamboli V, Sharma R, Sreelakshmi Y. 2018. NAC-NOR mutations in tomato

Penjar accessions attenuate multiple metabolic processes and prolong the fruit shelf life. Food

Chemistry **259**, 234–244.

Larsen B, Migicovsky Z, Jappesen AA, Gardner KM, Toldam-Andersen TB, Myles S,

Ørgaard M, Petersen MA, Pedersen C. 2019. Genome-Wide Association Studies in Apple

Reveal Loci for Aroma Volatiles, Sugar Composition and Harvest Date. The Plant Genome **in**

press.

Lenth RV. 2016. Least-Squares Means: The R Package lsmeans. Journal of Statistical Software

69, 1–33.

Lü P, Yu S, Zhu N, *et al.* 2018. Genome encode analyses reveal the basis of convergent

evolution of fleshy fruit ripening. Nature Plants **4**, 784.

Luby JJ, Shaw DV. 2001. Does Marker-assisted Selection Make Dollars and Sense in a Fruit

Breeding Program? HortScience **36**, 872–879.

Mathew IE, Agarwal P. 2018. May the Fittest Protein Evolve: Favoring the Plant-Specific

Origin and Expansion of NAC Transcription Factors. BioEssays **40**, 1800018.

McClure KA, Gardner KM, Douglas GM, *et al.* 2018. A Genome-Wide Association Study of

Apple Quality and Scab Resistance. The Plant Genome **11**, 0.

Migicovsky Z, Gardner KM, Money D, *et al.* 2016. Genome to Phenome Mapping in Apple

Using Historical Data. The Plant Genome **9**, 0.

Migicovsky Z, Li M, Chitwood DH, Myles S. 2018. Morphometrics Reveals Complex and

Heritable Apple Leaf Shapes. Frontiers in Plant Science **8**.

Nakatsuka A, Murachi S, Okunishi H, Shiomi S, Nakano R, Kubo Y, Inaba A. 1998.

Differential Expression and Internal Feedback Regulation of 1-Aminocyclopropane-1-

Carboxylate Synthase, 1-Aminocyclopropane-1-Carboxylate Oxidase, and Ethylene Receptor

Genes in Tomato Fruit during Development and Ripening. Plant Physiology **118**, 1295–1305.

Oraguzie NC, Iwanami H, Soejima J, Harada T, Hall A. 2004. Inheritance of the Md-ACS1

gene and its relationship to fruit softening in apple (Malus × domestica Borkh.). Theoretical and

Applied Genetics **108**, 1526–1533.

Ornelas-Paz J de J, Quintana-Gallegos BM, Escalante-Minakata P, Reyes-Hernández J,

Pérez-Martínez JD, Rios-Velasco C, Ruiz-Cruz S. 2018. Relationship between the firmness of

Golden Delicious apples and the physicochemical characteristics of the fruits and their pectin

during development and ripening. Journal of Food Science and Technology **55**, 33–41.

Osorio S, Alba R, Damasceno CMB, *et al*. 2011. Systems Biology of Tomato Fruit

Development: Combined Transcript, Protein, and Metabolite Analysis of Tomato Transcription

Factor (nor, rin) and Ethylene Receptor (Nr) Mutants Reveals Novel Regulatory Interactions.

Plant Physiology **157**, 405–425.

Pirona R, Eduardo I, Pacheco I, *et al*. 2013. Fine mapping and identification of a candidate

gene for a major locus controlling maturity date in peach. BMC Plant Biology **13**, 166.

Shinozaki Y, Nicolas P, Fernandez-Pozo N, *et al*. 2018. High-resolution spatiotemporal

transcriptome mapping of tomato fruit development and ripening. Nature Communications **9**,

364.

Solé X, Guinó E, Valls J, Iniesta R, Moreno V. 2006. SNPStats: a web tool for the analysis of

association studies. Bioinformatics **22**, 1928–1929.

Sunako T, Sakuraba W, Senda M, Akada S, Ishikawa R, Niizeki M, Harada T. 1999. An

Allele of the Ripening-Specific 1-Aminocyclopropane-1-Carboxylic Acid Synthase Gene

(ACS1) in Apple Fruit with a Long Storage Life. Plant Physiology **119**, 1297–1304.

Tigchelaar EC, Tomes ML, Kerr EA, Barman RJ. 1973. A new fruit ripening mutant, non-

ripening (nor). Rep. Tomato Genet. Coop **23**, 33.

Urrestarazu J, Muranty H, Denancé C, *et al.* 2017. Genome-Wide Association Mapping of

Flowering and Ripening Periods in Apple. Frontiers in Plant Science **8**.

Van Eck J, Keen P, Tjahjadi M. 2019. Agrobacterium tumefaciens-Mediated Transformation

of Tomato. In: Kumar S, Barone P, Smith M, eds. Transgenic Plants: Methods and Protocols.

New York, NY: Springer New York, 225–234.

Wang R, Tavano EC da R, Lammers M, Martinelli AP, Angenent GC, Maagd RA de. 2019.

Re-evaluation of transcription factor function in tomato fruit development and ripening with

CRISPR/Cas9-mutagenesis. Scientific Reports **9**, 1696.

Wang D, Yeats TH, Uluisik S, Rose JKC, Seymour GB. 2018. Fruit Softening: Revisiting the

Role of Pectin. Trends in Plant Science **23**, 302–310.

Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.