1 **Towards Improved Molecular Identification Tools in Fine Fescue (*Festuca* L.,**
2 **Poaceae) Turfgrasses: Nuclear Genome Size, Ploidy, and Chloroplast Genome**
3 **Sequencing**

4 **Yinjie Qiu**[1*]**, Cory D. Hirsch**[2]**, Ya Yang**[3] **and Eric Watkins**[1*]

5 ¹ Department of Horticultural Science, University of Minnesota, St. Paul, MN, 55108, USA;
6   qiuxx221@umn.edu (Y.Q.); ewatkins@umn.edu (E.W.)
7 2 Department of Plant Pathology, University of Minnesota, St. Paul, MN, 55108, USA; cdhirsch@umn.edu (C.
8   H.)
9 3 Department of Plant and Microbial Biology, University of Minnesota, St. Paul, MN, 55108, USA;
10   yayang@umn.edu (Y.Y.)
11 * Correspondence: qiuxx221@umn.edu (Y.Q.); ewatkins@umn.edu (E.W.); Tel.: +1- 612-624-7496 (E.W.)

12 **Abstract:** Fine fescues (*Festuca* L., Poaceae) are turfgrass species that perform well in low-
13 input environments. Based on morphological characteristics, the most commonly-utilized
14 fine fescues are divided into five taxa: three are subspecies within *F. rubra* L. and the
15 remaining two are treated as species within the *F. ovina* L. complex. Morphologically, these
16 five taxa are very similar, both identification and classification of fine fescues remain
17 challenging. In an effort to develop identification methods for fescues, we used flow
18 cytometry to estimate genome size, ploidy level, and sequenced the chloroplast genome of
19 all five taxa. Fine fescue chloroplast genome sizes ranged from 133,331 to 133,841 bp and
20 contained 113 to 114 genes. Phylogenetic relationship reconstruction using whole
21 chloroplast genome sequences agreed with previous work based on morphology.
22 Comparative genomics suggested unique repeat signatures for each fine fescue taxon that
23 could potentially be used for marker development for taxon identification.

24 **Keywords:** Fine fescue, chloroplast genome, phylogeny, comparative genomics
25

26 **1. Introduction**

27   With ca. 450 species, Fescues (*Festuca* L., Poaceae) is a large and diverse genus of
28 perennial grasses (Clayton and Renvoize 1986). Fescue species are distributed mostly in
29 temperate zones of both the northern and southern hemispheres, but most commonly found
30 in the northern hemisphere (Jenkin 1959). Several of the fescue species have been commonly
31 used as turfgrass. Based on both leaf morphology and nuclear ITS sequences, fescue species
32 can be divided into two groups: broad-leaved fescues and fine-leaved fescues (Torrecilla
33 and Catalán 2002). Broad-leaved fescues commonly used as turfgrass include tall fescue
34 (*Festuca arundinacea* Schreb.) and meadow fescue (*Festuca pratensis* Huds.). Fine-leaved
35 fescues are a group of cool-season grasses that include five commonly used taxa called fine
36 fescues. Fine fescues include hard fescue (*Festuca brevipila* Tracey, 2n=6x=42), sheep fescue
37 (*Festuca ovina* L., 2n=4x=28), strong creeping red fescue (*Festuca rubra* ssp. *rubra* 2n=8x=56),
38 slender creeping red fescue (*Festuca rubra* ssp. *litoralis* (G. Mey.) Auquier 2n=6x=42), and
39 Chewings fescue (*Festuca rubra* ssp. *fallax* (Thuill.) Nyman 2n=6x=42) (Ruemmele et al. 1995).
40 All five taxa share very fine and narrow leaves and have been used for forage, turf, and
41 ornamental purposes. They are highly tolerant to shade and drought, prefer low pH (5.5-
42 6.5) and low fertility soils (Beard 1972). Additionally, fine fescues grow well in the shade or
43 sun, have reduced mowing requirements, and do not need additional fertilizer or
44 supplemental irrigation (Ruemmele et al. 1995).
45   Based on morphological and cytological features, fine fescues are currently divided into
46 two groups referred to as the *F. rubra* complex (includes *F. rubra* ssp. *litoralis*, *F. rubra* ssp.

rubra, *F. rubra* ssp. *fallax*) and the non- rhizomatous *F. ovina* complex (includes *F. brevipila*
and *F. ovina*) (Ruemmele et al. 1995). While it is relatively easy to separate fine fescue taxa
into their proper complex based on the presence and absence of rhizome, it is challenging
to identify taxon within the same complex. In the *F. rubra* complex, both ssp. *litoralis* and
ssp. *rubra* are rhizomatous while ssp. *fallax* is non-rhizomatous. However, the separation of
ssp. *litoralis* from ssp. *rubra* using rhizome length is challenging. Taxon identification within
the *F. ovina* complex relies heavily on leaf characters; however, abundant morphological and
ecotype diversity within *F. ovina* makes taxa identification difficult (Piper 1906). This is
further complicated by inconsistent identification methods between different continents.
For example, in the United States, sheep fescue is described as having a bluish gray leaf color
and hard fescue leaf blade color is considered green (Beard 1972), while in Europe, it is the
opposite (Hubbard 1968). Because the ploidy level of the five taxa varies from tetraploid to
octoploid, beyond morphological classifications, laser flow cytometry has been used to
determine ploidy level of fine fescues and some other fescue species (Huff and Palazzo
1998). A wide range of DNA contents within each complex suggests that the evolutionary
history of each named species is complicated, and interspecific hybridization might interfere
with species determination using this approach. Plant breeders have been working to
improve fine fescues for turf use for several decades, with germplasm improvement efforts
focused on disease resistance, traffic tolerance, and ability to perform well under heat stress
(Casler 2003). Turfgrass breeders have utilized germplasm collections from old turf areas as
a source of germplasm (Bonos and Huff 2013); however, confirming the taxon identity in
these collections has been challenging. A combination of molecular markers and flow
cytometry could be a valuable tool for breeders to identify fine fescue germplasm (Hebert
et al. 2003).
Due to the complex polyploidy history of fine fescues, sequencing plastid genomes
provides a more cost-effective tool for taxon identification than the nuclear genome because
it is often maternally inherited, lacks of heterozygosity, is present in high copies and usable
even in partially degraded material (Bryan et al. 1999, Provan et al. 2001). Previous studies
have developed universal polymerase chain reaction (PCR) primers to amplify non-coding
polymorphic regions for DNA barcoding in plants for species identification (Baldwin et al.
1995, Demesure et al. 1995). However, the polymorphisms discovered from these regions
are often single nucleotide polymorphisms that are difficult to apply using PCR screening
methods. For these reasons, it would be helpful to assemble chloroplast genomes and
identify simple sequence repeat (SSR) and tandem repeats polymorphisms. Chloroplast
genome sequencing has been simplified due to improved sequencing technology. In
turfgrass species, high throughput sequencing has been used to assemble the chloroplast
genomes of perennial ryegrass (*Lolium perenne* cv. Cashel) (Diekmann et al. 2009), tall fescue
(*Lolium arundinacea* cv. Schreb) (Cahoon et al. 2010), diploid *Festuca ovina*, *Festuca pratensis*,
Festuca altissima (Hand et al. 2013), and bermudagrass (*Cynodon dactylon*) (Huang et al. 2017).
To date, there is limited molecular biology information on fine fescue taxon identification
and their phylogenetic position among other turfgrass species (Hand et al. 2013, Cheng et
al. 2016). In this study, we used flow cytometry to confirm the ploidy level of five fine fescue
cultivars, each representing one of the five commonly utilized fine fescue taxon. We then
reported the complete chloroplast genome sequences of these five taxa, carried out
comparative genomics and phylogenetic inference. Based on the genome sequence we
identified unique genome features among fine fescue taxa and predicted taxon specific SSR
and tandem repeat loci for molecular marker development.

2. Results

2.1 Species Ploidy Level Confirmation

We used flow cytometry to estimate the ploidy levels of five fine fescue taxa by measuring the DNA content in each nucleus. DNA content was reflected by the flow cytometry mean PI-A value. Overall, fine fescue taxa had mean PI-A values roughly from 110 to 180 (**Figure 1 and Figure S1**). *Festuca rubra* ssp. *rubra* cv. Navigator II (2n=8x=56) had the highest mean PI-A value (181.434, %rCV 4.4). *Festuca rubra* ssp. *litoralis* cv. Shoreline (2n=6x=42) and *F. rubra* ssp. *fallax* cv. Treazure II (2n=6x=42) had similar mean PI-A values of 137.852, %rCV 3.7 and 145.864, %rCV 3.5, respectively. *Festuca brevipila* cv. Beacon (2n=6x=42) had a mean PI-A of 165.25, %rCV 1.9, while *F. ovina* cv. Quatro (2n=4x=28) had a mean PI-A of 108.43, %rCV 2.9. Standard reference *L. perenne* cv. Artic Green (2n=2x=14) had a G1 phase mean PI-A of 63.91, %rCV 3.0. USDA *F. ovina* PI 230246 (2n=2x=14) had a G1 mean PI-A of 52.73 (histogram not shown). The estimated genome size of USDA PI 230246 was 4.67 pg/2C. Estimated ploidy level of *F. brevipila* cv. Beacon was 6.3, *F. ovina* cv. Quatro was 4.11, *F. rubra* ssp. *rubra* cv. Navigator II was 6.9, *F. rubra* ssp. *litoralis* cv. Shoreline was 5.2, and *F. rubra* ssp. *fallax* cv. Treazure II was 5.5 (**Table 1**). All newly estimated ploidy levels roughly correspond to previously reported ploidy levels based on chromosome counts.

Figure 1. Flow cytometry results for the five fine fescue taxa. *Lolium perenne* (2n=2x=14) was used as the reference. Flow cytometry was able to separate *F. rubra* ssp. *rubra* from the other two subspecies

115 in the *F. rubra* complex. The mean PI-A values of *F. rubra* ssp. *fallax* and *F. rubra* ssp. *litoralis* were
116 similar (145.86 to 137.85).

Table 1. Summary of flow cytometry statistics, genome size estimation, and ploidy level estimation
of fine fescue species. *Lolium perenne* 2C DNA content was used to calculate fine fescue and USDA *F. ovina* PI 230246 genome size, calculated PI 230246 DNA content was used as reference to infer fine fescue ploidy level

Species name	Chromosome count	Cultivar name	Mean PI-A	%rCV *	Estimated Genome Size (pg/Nuclei)	Estimated Ploidy Level
F. brevipila	2n=6x=42	Beacon	165.3	1.9	14.6	6.3
F. ovina	2n=4x=28	Quatro	108.4	2.9	9.6	4.1
F. ovina PI 230246	2n=2x=14	NA	52.7	3.1	4.7	1.7
F. rubra ssp. rubra	2n=8x=56	Navigator II	181.4	4.4	16.1	6.9
F. rubra ssp. litoralis	2n=6x=42	Shoreline	137.9	3.7	12.2	5.2
F. rubra ssp. fallax	2n=6x=42	Treazure II	145.9	3.5	12.9	5.5
L. perenne	2n=2x=14	Artic Green	63.9	3.0	5.7	2.0

* %rCV: Quality of laser alignment. Low %rCV suggests high resolution sensitivity.

2.2 Plastid Genome Assembly and Annotation of Five Fescue Taxa

A total of 47,843,878 reads were produced from the five fine fescue taxa. After Illumina
adaptor removal, we obtained 47,837,438 reads. The assembled chloroplast genomes ranged
from 133,331 to 133,841 bp. The large single copy (LSC) and small single copy (SSC) regions
were similar in size between the sequenced fine fescue accessions (78 kb and 12 kb,
respectively). *Festuca ovina* and *F. brevipila* in the *F. ovina* complex had exactly the same size
inverted repeat (IR) region (42,476 bp). In the *F. rubra* complex, *F. rubra* ssp. *rubra* and *F.*
rubra ssp. *litoralis* had the same IR size (21,235 bp). Species in the *F. rubra* complex had a
larger chloroplast genome size compared to species in the *F. ovina* complex. All chloroplast
genomes shared similar GC content (38.4%) (**Figure 2**, **Table 2**). The fine fescue chloroplast
genomes encoded for 113-114 genes, including 37 transfer RNAs (tRNA), 4 ribosomal RNAs
(rRNA), and 72 protein-coding genes (**Table 2**). Genome structures were similar among all
five fine fescue taxa sequenced, except that the pseudogene *accD* was annotated in all three
subspecies of *F. rubra*, but not in the *F. ovina* complex (**Table S1**).

(a)   (b)

Figure 2. Whole chloroplast genome structure of *F. ovina* complex (a) and *F. rubra* complex (b). Genes
inside the circle are transcribed clockwise, genes outside are transcribed counter-clockwise. Genes
belong to different functional groups are color coded. GC content is represented by the dark gray
inner circle, the light gray corresponded to the AT content. IRA(B), inverted repeat A(B); LSC, large
single copy region; SSC, small single copy region.

Table 2. Characteristics of fine fescue chloroplast genomes.

	F. brevipila cv. Beacon	*F. ovina* cv. Quatro	*F. rubra* ssp. *rubra* cv. Navigator II	*F. rubra* ssp. *litoralis* cv. Shoreline	*F. rubra* ssp. *fallax* cv. Treazure II
NCBI GenBank ID	MN309822	MN309824	MN309825	MN309823	MN309826
Total Genome Size (bp)	133,331	133,508	133,804	133,814	133,841
Large Single Copy (bp)	78,462	78,632	78,888	78,909	78,882
Small Single Copy (bp)	12,393	12,400	12,446	12,435	12,451
Inverted Repeat (bp)	42,476	42,476	42,470	42,470	42,508
Ratio of LSC (%)	58.85	58.9	58.96	58.97	58.94
Ratio of SSC (%)	9.29	9.29	9.3	9.29	9.3
Ratio of IRs (%)	31.86	31.82	31.74	31.74	31.76
GC content (%)	38.4	38.4	38.4	38.4	38.4

2.3 Chloroplast Genome IR Expansion and Contraction

Contraction and expansion of the IR regions resulted in the size variation of chloroplast
genomes. We examined the four junctions in the chloroplast genomes, LSC/IRa, LSC/IRb,
SSC/IRa, and SSC/IRb of the fine fescue and the model turfgrass species *L. perenne*. Although
the chloroplast genome of fine fescue taxa were highly similar, some structural variations

were still found in the IR/LSC and IR/SSC boundary. Similar to *L. perenne*, fine fescue taxa
chloroplast genes *rpl22-rps19, rps19-psbA* were located in the junction of IR and LSC; *rps15-*
ndhF and *ndhH-rps15* were located in the junction of IR/SSC. In the *F. ovina* complex, the
rps19 gene was 37 bp into the LSC/IRb boundary while in the *F. rubra* complex and *L. perenne*,
the *rps19* gene was 36 bp into the LSC/IRb boundary (**Figure 3**). The *rsp15* gene was 308 bp
from the IRb/SSC boundary in *F. ovina* complex, 307 bp in *F. rubra* complex, and 302 bp in *L.*
perenne. Both the *ndhH* and the pseudogene fragment of the *ndhH* (*ψndhH*) genes spanned
the junction of the IR/SSC. The *ψndhH* gene crossed the IRb/SSC boundary with 32 bp into
SSC in *F. brevipila* and *F. ovina*, 9 bp in *F. rubra* ssp. *rubra* and *F. rubra* ssp. *litoralis*, 10 bp in
F. rubra ssp. *fallax,* and 7 bp in *L. perenne*. The *ndhF* gene was 88 bp from the IRb/SSC
boundary in *F. brevipila* and *F. ovina*, 91 bp in *F. rubra* ssp. *rubra*, 84 bp in *F. rubra* ssp. *litoralis*,
77 bp in *F. rubra* ssp. *fallax*, and 102 bp in *L. perenne*. Finally, the *psbA* gene was 87 bp apart
from the IRa/LSC boundary into the LSC in *L. perenne* and *F. ovina* complex taxa but 83 bp
in the *F. rubra* complex taxa.

Figure 3. Comparison for border positions of LSC, SSC and IR regions among five fine fescues and *L.*
perenne. Genes are denoted by boxes, and the gap between the genes and the boundaries are indicated
by the number of bases unless the gene coincides with the boundary. Extensions of genes are also
indicated above the boxes.

2.4 Whole Chloroplast Genome Comparison and Repetitive Element Identification

Genome-wide comparison among five fine fescue taxa showed high sequence similarity
with most variations located in intergenic regions (**Figure 4**). To develop markers for species
screening, we predicted a total of 217 SSR markers for fine fescue taxa sequenced (*F. brevipila*
39; *F. ovina* 45; *F. rubra* ssp. *rubra* 45; *F. rubra* ssp. *litoralis* 46; *F. rubra* ssp. *fallax* 42) that
included 17 different repeat types for the fine fescue species (**Figure 5a, Table S2**). The most
frequent repeat type was A/T repeats, followed by AT/AT. The pentamer AAATT/AATTT
repeat was only presented in the rhizomatous *F. rubra* ssp. *litoralis* and *F. rubra* ssp. *rubra*,
while ACCAT/ATGGT was only found in *F. ovina* complex species *F. brevipila* and *F. ovina*.
Similar to SSR loci prediction, we also predicted long repeats for the fine fescue species and
identified a total of 171 repeated elements ranging in size from 20 to 51 bp (**Figure 5b, Table**

S3). Complementary (C) matches were only identified in *F. brevipila* and *F. ovina*. *F. rubra*
species had more palindromic (P) and reverse (R) matches. Number of forward (F) matches
were similar between five taxa. Selected polymorphic regions were validated by PCR and
gel electrophoresis assay (**Figure S2**).

183
Figure 4. Sequence identity plot of fine fescues chloroplast genome sequences with *F. ovina* (2x) as the
reference using mVISTA. A cut-off of 70% identify was used for the plots, and the percent of identity
varies from 50% to 100% as noted on the y-axis. Most of the sequence variations between fine fescues
were in intergenic regions. Taxa in the *F. ovina* complex, *F. brevipila* and *F. ovina* showed high sequence
similarity. Similarly, subspecies within *F. rubra* complex also showed high sequence similarity.

Figure 5. (a) SSR repeat type and numbers in the five fine fescue taxa sequenced. Single nucleotide
repeat type has the highest frequency. No hexanucleotide repeats were identified in the fine fescue

chloroplast genomes sequenced. One penta-nucleotide repeat type (AAATT/AATTT) is unique to *F.*
rubra ssp. *rubra* and *F. rubra* ssp. *litoralis*; One penta-nucleotide repeat type (ACCAT/ATGGT) is
unique to *F. brevipila* and *F. ovina*. (b) Long repeats sequences in fine fescue chloroplast genomes.
Complement (C) match was only identified in the *F. ovina* complex; Reverse (R) match has the most
number variation in fine fescues.

Figure S2. Examples of PCR validation of predicted repeat regions based on fine fescue chloroplast
genomes. PCR primers were developed using Primer3 module(Untergasser, Cutcutache et al. 2012).
Primers used for the PCR assays (1) Forward primer 5'-GTCGTGGAATGCTTTTGTTCTC-3'; Reverse
primer 5'-AGTGGATTCATCAGATGATACA-3'; (2) Forward primer 5'-TTCCTCTTTTCATTG-
CAAAGTGGT AT-3'; Reverse primer 5'-TACTCGGAGGTTCGAATCCTTCC-3'. PCR products were
examined on 1% agarose gel and gel images showed fragment size separation between different
taxa(a). Figure (b) and (c) showed partial sequence alignment of regions amplified by primer sets (1
and 2).

2.5 SNP and InDel Distribution in the Coding Sequence of Five Fine Fescue Species

To identify single nucleotide polymorphisms (SNPs, non-reference allele in this
content), we used the diploid *F. ovina* chloroplast genome (JX871940.1) as the reference for
the mapping and used the genome annotation file to identify genic and non-genic SNPs.
The total genic and non-genic sequence of (JX871940.1) were 60,582 and 72,583 bp,
respectively. We found SNP polymorphisms were over-present within intergenic regions in
the *F. ovina* complex (~0.3 SNP/Kbp more), while were under-present in the *F. rubra* complex
(~0.5 SNP/Kbp less). Most InDels were located in intergenic regions of the fine fescue species
(**Table 3**). Between *F. ovina* and the *F. rubra* complex, *the ropC2* gene had the most SNPs (4
vs 31). The *rbcL* gene also has a high level of variation (1 vs 14.3). In addition, *rpoB, ccsA,*
NADH dehydrogenase subunit genes (*ndhH, ndhF, ndhA*), and ATPase subunit genes (*atpA,*
atpB, aptF) also showed variation between *F. ovina* and *F. rubra* complexes. Less SNP and
InDel variation were found within each complex (**Table 3, Table S4 and S5**).

Table 3. Distribution of SNPs and InDels for the five fine fescue taxa sequenced in this study.

Species	F. brevipila	F. ovina	F. rubra ssp. rubra	F. rubra ssp. litoralis	F. rubra ssp. fallax
Total number of SNPs	98	134	638	615	624
SNPs in the coding region	35	52	306	301	300
SNPs in intergenic region	63	82	332	314	324
SNPs per Kbp in genic region	0.5777	0.8583	5.0510	4.9685	4.9520
SNPs per Kbp in non-genic region	0.8680	1.1297	4.5741	4.3261	4.4639
Total number of InDels	112	102	149	156	149
InDels in the coding region	22	17	27	26	27
InDels in intergenic region	90	85	122	130	122
Percentage of InDels in the intergenic region	80.36	83.33	81.88	83.33	81.88
Average sequencing depth	171.61	86.81	101.58	77.04	50.94

2.6 Nucleotide Diversity Calculation

A sliding window analysis successfully detected highly variable regions in the fine
fescue chloroplast genomes (**Figure 6, Table S6**). The average nucleotide diversity (Pi)
among fine fescue taxa was relatively low (0.00318). The IR region showed lower variability
than the LSC and SSC region. There were several divergent loci having a Pi value greater
than 0.01 (*psbK-psbI*, *trnfM-trnE*, *trnC-rpoB*, *psbH-petB*, *trnL-trnF*, *trnS-rps4*, *aptB-rbcL-psaI*,
and *rpl32-trnL*), but mostly within intergenic regions. The *rbcL-psaI* region contained a
highly variable *accD-like* region in some fine fescue taxa, so we looked at the structural
variation of 10 taxa in the *Festuca - Lolium* complex. We found taxa in broad-leaved fescue
and *F. rubra* complex had similar structure, while *F. ovina* (2x, 4x) and *F. brevipila* had a 276
bp deletion in the *rbcL-psaI* intergenic region (**Figure 7**).

Figure 6. Sliding window analysis of fine fescue whole chloroplast genomes. Window size: 600 bp, step size: 200 bp. X-axis: the position of the midpoint of a window (kb). Y-axis: nucleotide diversity of each window. Inverted repeat regions are highlighted in grey. *rpl32-trnL* region has the most nucleotide diversity followed by *psbH- petB-trnL-trnF-trnS-rps4* region.

Figure 7. The alignment of *rbcL-psaI* intergenic sequence shows that the pseudogene *accD* is missing in both *F. ovina* (2x, 4x) and *F. brevipila* but present in the *F. rubra* complex and other species examined in this study. Species were ordered by complexes.

2.7 Phylogenetic Reconstruction of Fine Fescue Species

We reconstructed the phylogenetic relationships of taxa within the *Festuca - Lolium* complex using the chloroplast genomes sequenced in our study and eight publicly available complete chloroplast genomes including six taxa within the *Festuca-Lolium* complex (**Figure 8**). The dataset included 125,824 aligned characters, of which 3,923 were parsimony-informative and 91.11% characters are constant. The five fine fescue taxa were split into two clades ([ML]BS=100). In the *F. ovina* complex, two *F. ovina* accessions included in the

phylogenetic analysis, a diploid one from GenBank, and a tetraploid one newly sequenced in this study are paraphyletic to *F. brevipila* ([ML]BS=100). All three subspecies of *F. rubra* formed a strongly supported clade ([ML]BS=100). Together they are sisters to the *F. ovina* complex ([ML]BS=100).

Figure 8. Maximum likelihood (ML) phylogram of the *Festuca - Lolium* complex with 1,000 bootstrap replicates. Fine fescues were grouped into previous named complexes (*F. ovina* and *F. rubra*), sister to broad leaved fescues in the *Festuca – Lolium* complex.

3. Discussion

In this study, we used flow cytometry to determine the ploidy level of five fine fescue cultivars, assembled the chloroplast genomes for each, and identified structural variation and mutation hotspots. We also identified candidate loci for marker development to facilitate fine fescue species identification. Additionally, we reconstructed the phylogenetic relationships of the *Festuca-Lolium* complex using plastid genome information generated in this study along with other publicly-available plastid genomes.

While most crop plants are highly distinctive from their close relatives, *Festuca* is a species-rich genus that contains species with highly similar morphology and different ploidy level. Consequently, it is difficult for researchers to interpret species identity. In our case, flow cytometry was able to successfully separate fine fescue taxa *F. brevipila* cv. Beacon, *F. ovina* cv. Quatro and *F. rubra* ssp. *rubra* cv. Navigator II based on the estimated ploidy levels. However, it is difficult to distinguish between *F. rubra* ssp. *litoralis* cv. Shoreline and *F. rubra* ssp. *fallax* cv. Treazure II as they had similar PI-A values based on flow cytometry. We noticed that the average mean PI-A of the diploid *L. perenne* (63.91) was higher than the mean PI-A of diploid *F. ovina* (52.73), suggesting that *F. ovina* taxa have smaller genome size than *L. perenne*. The ploidy estimation in the *F. ovina* complex are fairly consistent while the estimations of genome sizes in the *F. rubra* complex are smaller than we expected, even though these two complexes are closely related. Indeed, a similar finding was reported by Huff et al (Huff and Palazzo 1998) who reported that *F. brevipila* has a larger genome size than *F. rubra* ssp. *litoralis* and *F. rubra* ssp. *fallax*, both of which have the same ploidy level as *F. brevipila*. The range of variation in DNA content within each complex suggest a complicated evolutionary history in addition to polyploidization (Huff and Palazzo 1998).

When we cannot identify taxon based on the ploidy level, we need different approaches to identify them. The presence and absence of rhizome formation could be taken into consideration; for example, *F. rubra* ssp. *fallax* cv. Treazure II is a bunch type turfgrass, while *F. rubra* ssp. *litoralis* cv. Shoreline forms short and slender rhizomes (Meyer and Funk 1989). This method may not be effective because rhizome formation can be greatly affected by environmental conditions (Yang et al. 2015, Ma and Huang 2016).

To further develop molecular tools to facilitate species identification, we carried out chloroplast genome sequencing. We assembled the complete chloroplast genomes of five low-input turfgrass fine fescues using Illumina sequencing. Overall, the chloroplast genomes had high sequence and structure similarity among all five fine fescue taxa sequenced, especially within each complex. All five chloroplast genomes share similar gene content except for the three species in the *F. rubra* complex that have a pseudogene Acetyl-coenzyme A carboxylase carboxyl transferase subunit (*accD*). The *accD* pseudogene is either partially or completely absent in some monocots. Instead, a nuclear-encoded ACC enzyme has been found to replace the plastic *accD* gene function in some angiosperm linage (Rousseau-Gueutin et al. 2013). Indeed, even though the *accD* pseudogene is missing in the *F. brevipila* chloroplast genome, the gene transcript was identified in a transcriptome sequencing dataset (unpublished data), suggesting that this gene has been translocated to nucleus genome. Previous studies have shown that broad-leaf fescues, *L. perenne*, *O. sativa*, and *H. vulgare* all had the pseudogene *accD* gene, while it was absent in diploid *F. ovina*, *Z. mays*, *S. bicolor*, *T. aestivum,* and *B. distachyon* (Hand et al. 2013). Broad-leaf and fine-leaf fescues diverged around 9 Mya ago (Fjellheim et al. 2006), which raises an interesting question about the mechanisms of the relocation of *accD* among closely related taxa in the *Festuca-Lolium* complex and even within fine fescue species.

In plants, chloroplast genomes are generally considered "single copy" and lack recombination due to maternal inheritance (Ebert and Peakall 2009). For this reason, chloroplast genomes are convenient for developing genetic markers for distinguishing species and subspecies. We have identified a number of repeat signatures that are unique to a single species or species complex in fine fescue. For example, complement match is only identified in *F. ovina* complex, and *F. rubra* complex has more reversed matches. We also identified two SSR repeats unique to each of the two complexes. The first one consists of AAATT/AATTT repeat units is unique to *F. rubra* ssp. *litoralis* and *F. rubra* ssp. *rubra*, and the second one consists of ACCAT/ATGGT repeat units is unique to *F. brevipila* and *F. ovina*. In cases like the identification of hexaploids *F. brevipila*, *F. rubra* ssp. *fallax*, and *Festuca rubra* ssp. *litoralis*, it is critical to have these diagnostic repeats given all three taxa share similar PI-A values from flow cytometry. Taxon-specific tandem repeats could also aid the SSR repeats for species identification. We used chloroplast sequence developed candidate primer sets to solve the problem. Primer set (1) provided a clear separation of *F. rubra* ssp. *litoralis* cv. Shoreline and *F. rubra* ssp. *fallax* cv. Treazure II when flow cytometry was not able to separate them. Primer set (2) provided clear separation of *F. brevipila* cv. Beacon and *F. ovina* cv. Quatro, which provided an alternative method for *F. ovina* complex taxa identification. By combining both flow cytometry and candidate primer sets developed in this study, researchers will be able to identify fine fescue taxa within and between two complexes.

Nucleotide diversity analysis suggested that several variable genome regions exist among the five fine fescue taxa sequenced in this study. These variable regions included previously known highly variable chloroplast regions such as *trnL-trnF* and *rpl32-trnL* (Demesure et al. 1995, Dong et al. 2012). These regions have undergone rapid nucleotide

substitution and are potentially informative molecular markers for characterization of fine
fescue species.

Phylogeny inferred from DNA sequence is valuable for understanding the evolutionary
context of a species. The phylogenetic relationship of fine fescue using whole plastid
genome sequences agrees with previous classification based on genome size estimation and
morphology (Huff and Palazzo 1998, Cheng et al. 2016). The *F. ovina* complex includes *F.
ovina* and *F. brevipila* and the *F. rubra* complex includes *F. rubra* ssp. *rubra*, *F. rubra* ssp. *litoralis*
and *F. rubra* ssp. *fallax*, with the two rhizomatous subspecies (ssp. *rubra* and ssp. *literalis*)
being sister to each other. Within the *Festuca – Lolium* complex, fine fescues are monophyletic
and together sister to a clade consists of broad-leaved fescues and *Lolium*. In our analysis, *F.
brevipila* (6x) is nested within *F. ovina* and sister to the diploid *F. ovina*. It is likely that *F.
brevipila* arose from the hybridization between *F. ovina* (2x) and *F. ovina* (4x). Considering
the complex evolutionary history of this genus, further research using nuclear loci sequences
are needed to provide a more accurate phylogeny tree and validate this hypothesis.

The diversity of fine fescue provides valuable genetic diversity for breeding and
cultivar development. Breeding fine fescue cultivars for better disease resistance, heat
tolerance, and traffic tolerance could be achieved through screening wild accessions and by
introgressing desired alleles into elite cultivars. Some work has been done using *Festuca*
accessions in the USDA Germplasm Resources Information Network (GRIN)
(https://www.ars-grin.gov) to breed for improved forage production in fescue species
(Robbins et al. 2016). To date, there are 229 *F. ovina* and 486 *F. rubra* accessions in the USDA
GRIN. To integrate this germplasm into breeding programs, plant breeders and other
researchers need to confirm the ploidy level using flow cytometry and further identify them
using molecular markers. Resources developed in this study could provide the tools to
screen the germplasm accessions and refine the species identification so breeders can
efficiently use these materials for breeding and genetics improvement of fine fescue species.

4. Materials and Methods

Plant Material

Seeds from the fine fescue cultivars were obtained from the 2014 National Turfgrass
Evaluation Program (www.ntep.org, USA) and planted in the Plant Growth Facility at the
University of Minnesota, St. Paul campus under 16 hours daylight (25 °C) and 8 hours dark
(16 °C) with weekly fertilization. Single genotypes of *F. brevipila* cv. Beacon, *F. rubra* ssp.
litoralis cv. Shoreline, *F. rubra* ssp. *rubra* cv. Navigator II, *F. rubra* ssp. *fallax* cv. Treazure II,
and *F. ovina* cv. Quatro were selected and used for chloroplast genome sequencing.

Flow Cytometry

To determine the ploidy level of the cultivars used for sequencing and compare them
to previous work (2n=4x=28: *F. ovina*; 2n=6x=42: *F. rubra* ssp. *litoralis*, *F. rubra* ssp. *fallax*, and
F. brevipila; 2n=8x=56: *F. rubra* ssp. *rubra*), flow cytometry was carried out using *Lolium
perenne* cv. Artic Green (2n=2x=14) as the reference. Samples were prepared using CyStain
PI Absolute P (Sysmex, product number 05-5022). Briefly, to prepare the staining solution
for each sample, 12 μL propidium iodide (PI) was added to 12 mL of Cystain UV Precise P
staining buffer with 6 μL RNase A. To prepare plant tissue, a total size of 0.5 cm x 0.5 cm
leaf sample of the selected fine fescue was excised into small pieces using a razor blade in
500 μL CyStain UV Precise P extraction buffer and passed through a 50 μm size filter
(Sysmex, product number 04-004-2327). The staining solution was added to the flow-

through to stain nuclei in each sample. Samples were stored on ice before loading the flow
cytometer. Flow cytometry was carried out using the BD LSRII H4760 (LSRII) instrument
with PI laser detector using 480V with 2,000 events at the University of Minnesota Flow
Cytometry Resource (UCRF). Data was visualized and analyzed on BD FACSDiva 8.0.1
software. To estimate the genome size, *L. perenne* DNA (5.66 pg/2C) was used as standard
(Arumuganathan, Tallury et al. 1999), USDA PI 230246 (2n=2x=14) was used as diploid fine
fescue relative (unpublished data). To infer fine fescues ploidy, estimation was done using
equations (1) and (2) (Doležel et al. 2007).
382

$$\text{Sample 2C DNA Content} = \text{Standard 2C DNA Content (pg DNA)} \times \frac{\text{(Sample G1 Peak Mean)}}{\text{(Standard G1 Peak Mean)}} \quad (1)$$

$$\text{Sample Ploidy} = \frac{2n \times \text{Sample pg/Nucleus}}{\text{Diploid Relative pg/Nucleus}} \quad (2)$$

DNA Extraction and Sequencing

To extract DNA for chloroplast genome sequencing, 1 g of fresh leaves were collected
from each genotype and DNA was extracted using the Wizard Genomic DNA Purification
Kit (Promega, USA) following manufacturer instructions. DNA quality was examined on
0.8% agarose gel and quantified via PicoGreen (Thermo Fisher, Catalog number: P11496).
Sequencing libraries were constructed by NovoGene, Inc. (Davis, CA) using Nextera XT
DNA Library Preparation Kit (Illumina) and sequenced in 150 bp paired-end mode, using
the HiSeq X Ten platform (Illumina Inc., San Diego, CA, USA) with an average of 10 million
reads per sample. All reads used in this study were deposited in the NCBI Sequence Read
Archive (SRA) under BioProject PRJNA512126.
396

Chloroplast Genome Assembly and Annotation
Raw reads were trimmed of Illumina adaptor sequences using Trimmomatic (v. 0.32)
(Bolger et al. 2014). Chloroplast genomes were *de novo* assembled using NovoPlasty v. 2.0
(Dierckxsens et al. 2016). Briefly, *rbcL* gene sequence from diploid *F. ovina* (NCBI accession
number: JX871940) was extracted and used as the seed to initiate the assembly. NovoPlasty
assembler configuration was set as follows: *k-mer* size = 39; insert size = auto; insert range =
1.8; and insert range strict 1.3. Reads with quality score above 25 were used to complete the
guided assembly using *F. ovina* (NCBI accession number: JX871940) as the reference.
Assembled plastid genomes for each taxon were manually corrected by inspecting the
alignments of reads used in the assembly. The assembled chloroplast genomes were
deposited under BioProject PRJNA512126, GenBank accession numbers MN309822-
MN309826.
The assembled chloroplast genomes were annotated using the GeSeq pipeline (Tillich,
Lehwark et al. 2017) and corrected using DOGMA online interface
(https://dogma.ccbb.utexas.edu) (Wyman, Jansen et al. 2004). BLAT (a BLAST-like
alignment tool (Kent 2002) protein, tRNA, rRNA, and DNA search identity threshold was
set at 80% in the GeSeq pipeline using the default reference database with the generate
codon-based alignments option turned on. tRNAs were also predicted via tRNAscan-SE
v2.0 and ARAGORN v 1.2.38 using the bacterial/plant chloroplast genetic code (Lowe and
Eddy 1997, Laslett and Canback 2004). The final annotation was manually inspected and
corrected using results from both pipelines. The circular chloroplast map was drawn by the
OrganellarGenomeDRAW tool (OGDRAW) (Lohse et al. 2007).

 Nucleotide Polymorphism of Fine Fescue Species

To identify genes with the most single nucleotide polymorphism, quality trimmed
sequencing reads of the five fine fescues were mapped to the diploid *Festuca ovina*
chloroplast genome (NCBI accession number: JX871940) using BWA v.0.7.17 (Li and Durbin
2009). SNPs and short indels were identified using bcftools v.1.9 with the setting "mpileup
-Ou" and called via bcftools using the -mv function (Quinlan and Hall 2010). Raw SNPs
were filtered using bcftools filter -s option to filter out SNPs with low quality (Phred score
cutoff 20, coverage cutoff 20). The subsequent number of SNPs per gene and InDel number
per gene was calculated using a custom perl script SNP_vcf_from_gene_gff.pl
(https://github.com/qiuxx221/fine-fescue-).

To identify simple sequence repeat (SSR) markers for plant identification,
MIcroSAtellite identification tool (MISA v 1.0) was used with a threshold of 10, 5, 4, 3, 3, 3
repeat units for mono-, di-, tri-, tetra-, penta-, and hexanucleotide SSRs, respectively (Thiel
et al. 2003). The identification of repetitive sequences and structure of whole chloroplast
genome was done via REPuter program online server (https://bibiserv.cebitec.uni-
bielefeld.de/reputer) (Kurtz et al. 2001). Program configuration was set with minimal repeat
size set as 20 bp and with sequence identify above 90%. Data was visualized using ggplot2
in R (v 3.5.3). Finally, the sliding window analysis was performed using DnaSP (v 5) with a
window size of 600 bp, step size 200 bp to detected highly variable regions in the fine fescue
chloroplast genome (Librado and Rozas 2009).

Comparative Chloroplast Genomics Analysis

To compare fine fescue species chloroplast genome sequence variations, the five
complete chloroplast genomes were aligned and visualized using mVISTA, an online suite
of computation tools with LAGAN mode (Brudno et al. 2003, Frazer et al. 2004). The diploid
Festuca ovina (NCBI accession number: JX871940) chloroplast genome and annotation were
used as the template for the alignment.

Phylogenetic Analysis of Fine Fescues and Related Festuca species

To construct the phylogenetic tree of the fine fescues using the whole chloroplast
genome sequence, chloroplast genomes of 8 species were downloaded from GenBank. Of
the 8 downloaded genomes, perennial ryegrass (*Lolium perenne*, AM777385), Italian ryegrass
(*Lolium multiflorum*, JX871942), diploid *Festuca ovina* (JX871940), tall fescue (*Festuca
arundiancea*, FJ466687), meadow fescue (*Festuca pratensis*, JX871941), and wood fescue
(*Festuca altissima*, JX871939) were within the *Festuca- Lolium* complex. Turfgrass species
outside of *Festuca-Lolium* complex including creeping bentgrass (*Agrostis stolonifera* L.,
EF115543) and *Cynodon dactylon* (KY024482.1) were used as an outgroup. All chloroplast
genomes were aligned using the MAFFT program (v 7) (Katoh and Standley 2013);
alignments were inspected and manually adjusted. Maximum likelihood (ML) analyses was
performed using the RAxML program (v 8.2.12) under GTR+G model with 1,000 bootstrap
(Stamatakis 2006). The phylogenetic tree was visualized using FigTree (v 1.4.3)
(https://github.com/rambaut/figtree) (Rambaut 2012).

5. Conclusions

Five newly-sequenced complete chloroplast genomes of fine fescue taxa were reported
in this study. Chloroplast genome structure and gene contents are both conserved, with the
presence and absence of *accD* pseudogene being the biggest structural variation between the

F. ovina and the *F. rubra* complexes. We identified SSR repeats and long sequence repeats of fine fescues and discovered several unique repeats for marker development. The phylogenetic constructions of fine fescue species in the *Festuca - Lolium* complex suggested a robust and consistent relationship compared to the previous identification using flow cytometry. This information provided a reference for future fine fescue taxa identification.

Supplementary Materials:

1. **Figure S1**. Flow cytometry nuclei population distribution of *L. perenne*, fine fescues, and diploid USDA PI accession. G1 populations were gated in red, G2 population was only gated in *L. perenne* in green.
2. **Figure S2.** Examples of PCR validation of predicted repeat regions based on fine fescue chloroplast genomes.
3. **Table S1.** Fine fescue chloroplast genomes gene content by gene category.
4. **Table S2**. SSR loci types and number distributions of fine fescue species predicted using MISA program.
5. **Table S3**. Tandem repeat loci and repeat types predicted using PEPuter program in fine fescue species.
6. **Table S4.** SNPs number per gene distribution for fine fescue species. *rpoC2* gene has the most SNPs (31) in *F. rubra* complex comparing to *F. ovina* species.
7. **Table S5.** InDel number per gene distribution for fine fescue species. *ndhA* gene had the most InDels *F. rubra* species. *atpI* gene had the most InDesl in *F. ovina* species.
8. **Table S6**. Sliding window analysis using DnaSP at window size of 600 bp, step size 200 bp to detect highly variable regions in the fine fescue chloroplast genome

Author Contributions: Y. Q. performed the experiments, analyzed the data, and wrote the manuscript; C. H. helped analyze data, wrote perl scripts; Y. Y. helped with phylogenetic analysis; E. W. secured funding for this project, supervised this research, provided suggestions, and comments. All authors contributed to the revision of the manuscript and approved the final version.

Funding: This research is funded by the National Institute of Food and Agriculture, U.S. Department of Agriculture, Specialty Crop Research Initiative under award number 2017-51181-27222.

Acknowledgments: The authors would like to thank Minnesota Supercomputing Institute for the high-performance computing clusters. We would also like to thank Jill Ekar, Jason Motl, and Therese Martin for providing instruction on operating the flow cytometer.

This manuscript has been released as a pre-print at bioRxiv, (Qiu, et al. 2019)

Conflicts of Interest: The authors declare no conflict of interest.

References

Arumuganathan, K., S. Tallury, M. Fraser, A. Bruneau and R. Qu (1999). "Nuclear DNA content of thirteen

turfgrass species by flow cytometry." <u>Crop science</u> **39**(5): 1518-1521.

Baldwin, B. G., M. J. Sanderson, J. M. Porter, M. F. Wojciechowski, C. S. Campbell and M. J. Donoghue (1995).

"The ITS region of nuclear ribosomal DNA: a valuable source of evidence on angiosperm phylogeny."

<u>Annals of the Missouri botanical garden</u>: 247-277.

Beard, J. B. (1972). "Turfgrass: Science and culture."

Bolger, A. M., M. Lohse and B. Usadel (2014). "Trimmomatic: a flexible trimmer for Illumina sequence data."

<u>Bioinformatics</u> **30**(15): 2114-2120.

Bonos, S. A. and D. R. Huff (2013). "Cool-season grasses: Biology and breeding." <u>Turfgrass: Biology, use, and</u>

<u>management</u>(turfgrassbiolog): 591-660.

Brudno, M., C. B. Do, G. M. Cooper, M. F. Kim, E. Davydov, E. D. Green, A. Sidow, S. Batzoglou and N. C. S.

Program (2003). "LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic

DNA." <u>Genome research</u> **13**(4): 721-731.

Bryan, G., J. McNicoll, G. Ramsay, R. Meyer and W. De Jong (1999). "Polymorphic simple sequence repeat

markers in chloroplast genomes of Solanaceous plants." <u>Theoretical and Applied Genetics</u> **99**(5): 859-867.

Cahoon, A. B., R. M. Sharpe, C. Mysayphonh, E. J. Thompson, A. D. Ward and A. Lin (2010). "The complete

chloroplast genome of tall fescue (Lolium arundinaceum; Poaceae) and comparison of whole plastomes

from the family Poaceae." <u>American Journal of botany</u> **97**(1): 49-58.

Casler, M. D. (2003). <u>Turfgrass biology, genetics, and breeding</u>, John Wiley & Sons.

Cheng, T., C. Xu, L. Lei, C. Li, Y. Zhang and S. Zhou (2016). "Barcoding the kingdom Plantae: new PCR primers

for ITS regions of plants with improved universality and specificity." <u>Molecular Ecology Resources</u> **16**(1):

138-149.

Clayton, W. D. and S. A. Renvoize (1986). "Genera graminum. Grasses of the world." <u>Genera graminum.</u>

<u>Grasses of the World.</u> **13**.

Demesure, B., N. Sodzi and R. Petit (1995). "A set of universal primers for amplification of polymorphic non-

coding regions of mitochondrial and chloroplast DNA in plants." <u>Molecular ecology</u> **4**(1): 129-134.

Diekmann, K., T. R. Hodkinson, K. H. Wolfe, R. van den Bekerom, P. J. Dix and S. Barth (2009). "Complete

chloroplast genome sequence of a major allogamous forage species, perennial ryegrass (Lolium perenne

L.)." <u>DNA research</u> **16**(3): 165-176.

Dierckxsens, N., P. Mardulyn and G. Smits (2016). "NOVOPlasty: de novo assembly of organelle genomes from

whole genome data." <u>Nucleic acids research</u> **45**(4): e18-e18.

Doležel, J., J. Greilhuber and J. Suda (2007). "Estimation of nuclear DNA content in plants using flow

cytometry." <u>Nature protocols</u> **2**(9): 2233.

Dong, W., J. Liu, J. Yu, L. Wang and S. Zhou (2012). "Highly variable chloroplast markers for evaluating plant

phylogeny at low taxonomic levels and for DNA barcoding." <u>PloS one</u> **7**(4): e35071.

Ebert, D. and R. Peakall (2009). "Chloroplast simple sequence repeats (cpSSRs): technical resources and

recommendations for expanding cpSSR discovery and applications to a wide array of plant species."

<u>Molecular Ecology Resources</u> **9**(3): 673-690.

Fjellheim, S., O. A. Rognli, K. Fosnes and C. Brochmann (2006). "Phylogeographical history of the widespread

meadow fescue (Festuca pratensis Huds.) inferred from chloroplast DNA sequences." <u>Journal of</u>

<u>Biogeography</u> **33**(8): 1470-1478.

Frazer, K. A., L. Pachter, A. Poliakov, E. M. Rubin and I. Dubchak (2004). "VISTA: computational tools for

comparative genomics." <u>Nucleic acids research</u> **32**(suppl_2): W273-W279.

Hand, M. L., G. C. Spangenberg, J. W. Forster and N. O. Cogan (2013). "Plastome sequence determination and
comparative analysis for members of the Lolium-Festuca grass species complex." <u>G3: Genes, Genomes,</u>
<u>Genetics</u>: g3. 112.005264.
Hebert, P. D., A. Cywinska, S. L. Ball and J. R. Dewaard (2003). "Biological identifications through DNA
barcodes." <u>Proceedings of the Royal Society of London. Series B: Biological Sciences</u> **270**(1512): 313-321.
Huang, Y.-Y., S.-T. Cho, M. Haryono and C.-H. Kuo (2017). "Complete chloroplast genome sequence of
common bermudagrass (Cynodon dactylon (L.) Pers.) and comparative analysis within the family
Poaceae." <u>PloS one</u> **12**(6): e0179055.
Hubbard, C. E. (1968). "Grasses. A guide to their structure, identification, uses, and distribution in the British
Isles." <u>Grasses. A guide to their structure, identification, uses, and distribution in the British Isles.</u>
Huff, D. R. and A. J. Palazzo (1998). "Fine fescue species determination by laser flow cytometry." <u>Crop science</u>
38(2): 445-450.
Jenkin, T. J. (1959). "Fescue Species (Festuca L.)." <u>In: Roemer, T. & W. Rudorf. Handbuch der</u>
<u>Pflanzenzüchtung</u>(4): 418–434.
Katoh, K. and D. M. Standley (2013). "MAFFT multiple sequence alignment software version 7: improvements
in performance and usability." <u>Molecular biology and evolution</u> **30**(4): 772-780.
Kent, W. J. (2002). "BLAT—the BLAST-like alignment tool." <u>Genome research</u> **12**(4): 656-664.
Kurtz, S., J. V. Choudhuri, E. Ohlebusch, C. Schleiermacher, J. Stoye and R. Giegerich (2001). "REPuter: the
manifold applications of repeat analysis on a genomic scale." <u>Nucleic acids research</u> **29**(22): 4633-4642.
Laslett, D. and B. Canback (2004). "ARAGORN, a program to detect tRNA genes and tmRNA genes in
nucleotide sequences." <u>Nucleic acids research</u> **32**(1): 11-16.
Li, H. and R. Durbin (2009). "Fast and accurate short read alignment with Burrows–Wheeler transform."
<u>bioinformatics</u> **25**(14): 1754-1760.
Librado, P. and J. Rozas (2009). "DnaSP v5: a software for comprehensive analysis of DNA polymorphism
data." <u>Bioinformatics</u> **25**(11): 1451-1452.
Lohse, M., O. Drechsel and R. Bock (2007). "OrganellarGenomeDRAW (OGDRAW): a tool for the easy
generation of high-quality custom graphical maps of plastid and mitochondrial genomes." <u>Current genetics</u>
52(5-6): 267-274.
Lowe, T. M. and S. R. Eddy (1997). "tRNAscan-SE: a program for improved detection of transfer RNA genes in
genomic sequence." <u>Nucleic acids research</u> **25**(5): 955-964.
Ma, X. and B. Huang (2016). "Gibberellin-stimulation of rhizome elongation and differential GA-responsive
proteomic changes in two grass species." <u>Frontiers in plant science</u> **7**: 905.
Meyer, W. A. and C. R. Funk (1989). "Progress and Benefits to Humanity from Breeding Cool-Season Grasses
for Turf 1." <u>Contributions from breeding forage and turf grasses</u>(contributionsfr): 31-48.
Piper, C. V. (1906). <u>North American species of Festuca</u>, US Government Printing Office.
Provan, J., W. Powell and P. M. Hollingsworth (2001). "Chloroplast microsatellites: new tools for studies in
plant ecology and evolution." <u>Trends in ecology & evolution</u> **16**(3): 142-147.
Qiu, Y., C. D. Hirsch, Y. Yang and E. Watkins (2019). "Chloroplast Genome Sequencing and Comparative
Analysis for Fine Fescue (Festuca L., Poaceae) Turfgrasses." <u>bioRxiv</u>: 708149.
Quinlan, A. R. and I. M. Hall (2010). "BEDTools: a flexible suite of utilities for comparing genomic features."
<u>Bioinformatics</u> **26**(6): 841-842.
Rambaut, A. (2012). FigTree v1. 4.

Robbins, M. D., J. E. Staub and B. S. Bushman (2016). "Development of fine-leaved Festuca grass populations

identifies genetic resources having improved forage production with potential for wildfire control in the

western United States." Euphytica **209**(2): 377-393.

Rousseau-Gueutin, M., X. Huang, E. Higginson, M. Ayliffe, A. Day and J. N. Timmis (2013). "Potential

functional replacement of the plastidic acetyl-CoA carboxylase subunit (accD) gene by recent transfers to

the nucleus in some angiosperm lineages." Plant physiology **161**(4): 1918-1929.

Ruemmele, B., L. Brilman and D. Huff (1995). "Fine fescue germplasm diversity and vulnerability." Crop

science **35**(2): 313-316.

Stamatakis, A. (2006). "RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of

taxa and mixed models." Bioinformatics **22**(21): 2688-2690.

Thiel, T., W. Michalek, R. Varshney and A. Graner (2003). "Exploiting EST databases for the development and

characterization of gene-derived SSR-markers in barley (Hordeum vulgare L.)." Theoretical and applied

genetics **106**(3): 411-422.

Tillich, M., P. Lehwark, T. Pellizzer, E. S. Ulbricht-Jones, A. Fischer, R. Bock and S. Greiner (2017). "GeSeq–

versatile and accurate annotation of organelle genomes." Nucleic acids research **45**(W1): W6-W11.

Torrecilla, P. and P. Catalán (2002). "Phylogeny of broad-leaved and fine-leaved Festuca lineages (Poaceae)

based on nuclear ITS sequences." Systematic Botany **27**(2): 241-252.

Untergasser, A., I. Cutcutache, T. Koressaar, J. Ye, B. C. Faircloth, M. Remm and S. G. Rozen (2012). "Primer3—

new capabilities and interfaces." Nucleic acids research **40**(15): e115-e115.

Wyman, S. K., R. K. Jansen and J. L. Boore (2004). "Automatic annotation of organellar genomes with

DOGMA." Bioinformatics **20**(17): 3252-3255.

Yang, M., L. Zhu, C. Pan, L. Xu, Y. Liu, W. Ke and P. Yang (2015). "Transcriptomic analysis of the regulation of

rhizome formation in temperate and tropical lotus (Nelumbo nucifera)." Scientific reports **5**: 13059.

606