The Gut Microbiome of Exudivorous Wild and Captive Marmosets

Joanna Malukiewicz (corresponding author)
Primate Genetics Laboratory, German Primate Research Center, Leibniz Institute for Primate Research, Goettingen, Germany

Reed A. Cartwright
School of Life Sciences and The Biodesign Institute, Arizona State University, Tempe AZ, USA

Jorge A. Dergam
Department of Animal Biology, Federal University of Vicosa, Vicosa, MG, Brazil

Claudia S. Igayara
Guarulhos Municipal Zoo, Guarulhos, SP, Brazil

Sharon Kessler
Department of Psychology, Faculty of Natural Sciences, University of Stirling, Stirling, , Scotland
Department of Anthropology, Durham University, Durham DH1 3LE, UK

Silvia B. Moreira
Centro de Primatologia do Rio de Janeiro, Guapimirim, RJ, Brazil

Leanne T. Nash
School of Human Evolution and Social Change, Arizona State University, Tempe, AZ, USA

Patricia A. Nicola
Programa de Pos-Graduacao Ciências da Saúde e Biológicas, Universidade Federal do Vale do São Francisco, Petrolina, PE

Luiz C.M. Pereira
Centro de Conservação e Manejo de Fauna da Caatinga, Universidade Federal do Vale do Sao Francisco, Petrolina PE, Brazil

Alcides Pissinatti
Centro de Primatologia do Rio de Janeiro, Guapimirim, RJ, Brazil

Carlos R. Ruiz-Miranda
Laboratorio das Ciencias Ambientais, Centro de Biociencias e Biotecnologia, Universidade Estadual do Norte Fluminense, Campos dos Goytacazes RJ, Brazil

Andrew T. Ozga
Center for Evolution and Medicine, Arizona State University, Tempe, Arizona, USA
Halmos College of Arts and Sciences, Nova Southeastern University, Fort Lauderdale, FL, USA

Christian Roos
Primate Genetics Laboratory, German Primate Research Center, Leibniz Institute for Primate Research, Göttingen, Germany
Gene Bank of Primates, German Primate Research Center, Leibniz Institute for Primate Research, Göttingen, Germany

Daniel L. Silva
Núcleo de Pesquisas em Ciências Biológicas - NUPEB, Federal University of Ouro Preto, Ouro Preto, MG, Brazil

Anne C. Stone
School of Human Evolution and Social Change, Arizona State University, Tempe, AZ, USA
Institute of Human Origins, Arizona State University, Tempe, Arizona, USA
Center for Evolution and Medicine, Arizona State University, Tempe, Arizona, USA

Adriana D. Grativol
Laboratorio das Ciencias Ambientais, Centro de Biociencias e Biotecnologia, Universidade Estadual do Norte Fluminense, Campos dos Goytacazes RJ, Brazil

Abstract

Among mammals, captive dietary specialists like primate folivores are prone to gastrointestinal distress and suffer the greatest gut microbiome diversity losses relative to wild individuals. Less is known about the gut microbiome of mammalian exudivores, which exploit tree gums and sap, and increased knowledge could improve management of wild exudivores and facilitate captive exudivore welfare. A number of primates, including *Callithrix* marmosets, represent key mammalian exudivores. Captive marmosets commonly develop gastrointestinal distress symptoms that in human diseases are linked to microbiome dysbiosis. Thus, we studied wild and captive *Callithrix* gut microbiome composition and predictive function through 16S V4 ribosomal subunit sequencing of 59 wild, translocated, and captive pure and hybrid *Callithrix*. We found that host environment had a stronger effect on the gut microbiome than host taxon. Captive marmosets showed relatively reduced gut microbiome diversity. Wild *Callithrix* gut microbiomes were enriched for carbohydrate function and *Bifidobacterium,* which process host-indigestible carbohydrates. Captive marmosets had the highest relative abundance of Enterobacteriaceae, a family containing several pathogenic bacteria. Captive marmosets also showed gut microbiome composition aspects seen in human gastrointestinal diseases. Thus, captivity may perturb the exudivore gut microbiome, which raises implications for captive exudivore welfare and calls for modified husbandry practices.

Keywords

exudivore, non-human primate, microbiome, Neotropical, microbiome, *Bifidobacterium*

Introduction

The mammalian gut microbiome plays an important role in host physiology [1-2], and microbiome dysbiosis may negatively impact host health [2-7]. Closely related mammals generally share similar gut microbiome composition (GMC), which is usually enriched for bacteria associated with the main macronutrients of a host's feeding strategy [8-10]. Yet, the environment significantly alters individual host microbiomes, as evidenced by differences in microbiome composition between wild and captive conspecifics [9, 11-14]. Gut microbiome studies of captive and wild mammals show that non-human primates (NHPs) experience relatively large losses of native gut microbiome diversity in captivity compared to the wild [6,11]. Additionally, folivorous NHPs are especially prone to GI problems in captivity [15], and among humans and non-human primates (NHPs) dysbiosis in GMC has been tied to gastro-intestinal (GI) diseases [7,15,16].

A number of mammals are exudivorous [16], particularly primates [18], but their gut microibome remains less studied relative to other mammals. However, the Brazilian *Callithrix* marmosets, obligate NHP exudivores, are excellent models to improve our understanding of exudivore gut microbiome. *Callithrix* species are endemic to Brazil and in the wild subsist on hard to digest oligosaccharides of tree gums or, if they gouge, hardened saps, that require fermentation for digestion [19, 20]. Marmosets are regularly maintained in captivity as biomedical research models [21], captive breeding of *C. aurita* (one of the 25 most endangered primates in the world [22]), and due to illegal pet trafficking [23, 24]. In captivity, marmosets commonly develop symptoms of GI distress referred to by some authors as marmoset wasting syndrome) like chronic diarrhea, chronic enteritis, and chronic colitis without clear pathogenesis

[25-28]. As mammalian dietary specialists, it is likely that captivity causes gut microbiome dysbiosis in *Callithrix,* which in turn may be implicated in the pathogenesis of GI distress.

Given these problems, further investigation of the gut microbiome of mammalian exudivores is necessary and will provide baseline data to improve their welfare in captivity. A necessary first step towards understanding diseased GMC profiles is defining baseline GMC variation of non-diseased individuals [2,29]. Thus, comparing GMC of wild and captive conspecifics is an important step for such approaches. Up to now, *Callithrix* gut microbiome studies have focused on captive *C. jacchus* to identify specific bacterial strains [30-33] and on how life history, social, or laboratory conditions affect the GMC [34-36]. Here, we determine gut microbiome profiles of wild and translocated/captive *Callithrix* sampled throughout Brazil. We applied 16S ribosomal subunit (rRNA, V4 region) gene amplicon sequencing of *Callithrix* anal microbiota, and investigated GMC and gut microbiome predictive functional profiles. Anal swabs were sampled from 59 individuals of four species and three hybrid types (Table 1) that were sampled in the wild and Brazilian captive facilities (Figure 1). We hypothesize that: (1) *Callithrix* taxa share similar bacterial taxa as part of host GMC; (2) captive *Callithrix* have less diverse GMC than wild *Callithrix*; and (3) *Callithrix* GMC and predictive functional profiles are strongly biased toward carbohydrate metabolism.

Figure 1. Natural *Callithrix* ranges and sampling locations in Brazil. Sampling locations are represented by different colored shapes and species names are written next to their respective ranges. Legend abbreviations are as follows: MG: Minas Gerais, Rio de Janeiro: RJ, Pernambuco: PE, São Paulo: SP; CPRJ: Centro de Primatologia do Rio de Janeiro; CEMAFAUNA: Centro de Conservação e Manejo da Fauna da Caatinga.

Table 1. Host count distribution by taxon, hybrid status, and environment categories

Classification	Species	Hybrid	Captive	Translocated	Wild
C. aurita	10	0	5	3	2
C. geoffroyi	4	0	3	0	1
C. jacchus	9	0	9	0	0
C. penicillata	7	0	6	1	0
C. jacchus x *C. penicillata*	0	23	1	22	5
C. penicillata x *C. geoffroyi*	0	5	0	0	0
C. aurita x *Callithrix* sp.	0	1	0	1	0
Total	**30**	**29**	**24**	**27**	**8**

Materials and Methods

Sample Collection

We collected anal swabs between 2015 and 2016 from 59 adult individuals (Figure 1, Table 1 and Supplementary Table 1). Marmoset sampling was authorized by the Brazilian Environmental Ministry (SISBIO protocol #47964-2), the Arizona State University IACUC (protocol# 15-144R). Wild animals were captured with Tomahawk style traps baited with bananas. All sampled animals were immobilized by injection of ketamine (10 mg/kg of body

weight) into the inner thigh intramuscular region. All were photographed, weighed, measured, examined clinically by veterinarians, and deemed healthy upon examination. Copan FLOQSwab tips were inserted and rotated in the anal region and then each swab was submerged in a storage buffer (50 mM Tris pH 8.0, 50 mM EDTA, 50 mM Sucrose, 100 mM NaCl, 1% SDS) for several seconds and then discarded. After processing, animals were returned to cages, and allowed to recover. Wild marmosets were released at original capture sites. Specimens taxon identification followed phenotype descriptions in [37-40], and confirmed by whole genome genotyping for some individuals (unpublished data). Specimens were also classified by their environment as wild (captured as free-range individuals), translocated (born wild but later put into captivity), or captive (born and raised in captivity).

Sample Processing and Sequencing

Bacterial DNA extraction from *Callithrix* anal swabs was carried out by adding 0.3g zirconia beads to tubes containing storage buffer and swabbed material following a modified phenol-chloroform protocol [41]. Modifications included beating the samples on a vortex fitted with a horizontal vortex adaptor (#13000-V1-24, Mo Bio, Carlsbad, CA, USA) for 10 minutes at step "2Aiii," precipitating samples in 100% ethanol in step "2Axvi," and rehydrating DNA pellets in 25 uL low TE buffer at step "2Axxii." Extracted DNA was quantified on a Qubit3 (Life Technologies, Carlsbad, CA, USA) with a dsDNA HS Assay Kit (Life Technologies). Then the V4 region of the bacterial 16S rRNA gene was amplified in triplicate using the barcoded primer set 515f/806r [42]. Genetic samples have been registered in the Brazilian CGen SISGEN database (Supplementary Table 2). Amplicon triplicates were combined for each individual and then pooled in equimolar amounts into a multiplexed Illumina sequencing library. The library was purified with a Zymo DNA Concentrator and Cleaner-5 (#D4013, Zymo Research, Irving,

CA, USA) and size selected for 375-380 base pairs with Agencourt Ampure XP (#A63880, Beckman Coulter, Indianapolis, IN, USA) magnetic beads. Following international regulations, DNA extraction and library preparation was carried out at the Federal University of Viçosa, Brazil and then sequenced at Arizona State University, USA on an Illumina MiSeq for 2x250 cycles. The dataset supporting the conclusions of this article is available in the NCBI SRA repository under Bioproject SUB6302859.

Bioinformatics and Statistical Analysis

Code for bioinformatics analysis described below for Qiime2-2018.6 and Qiime2-2020.2 [43] is available at https://github.com/Callithrix-omics/callithrix_microbiome. Data were demultiplexed using default parameters in Qiime2-2018.6. The DADA2 Qiime2-2018.6 plug-in [44] was used to quality-filter and trim sequences and join paired-end reads. Upon trimming, the first 10 and last 30 nucleotides were removed from reverse reads due to low base quality. These steps resulted in feature tables of DNA sequences and their per-sample counts, termed operational taxonomic units (OTUs). MAAFT and FastTree Qiime2-2018.6 plug-ins [45,46] aligned and produced a phylogenetic tree of feature sequences, which was then mid-pointed rooted.

Qiime2-2018.6's q2-diversity plug-in was used to estimate alpha diversity measures (Shannon's diversity index, observed OTUs, and Pielou's Evenness) and calculate beta diversity weighted UniFrac distance [47]. Based on rarefaction, we set the minimum sampling depth to 45000 reads for alpha and beta diversity analyses. Box-plots were generated for each alpha diversity measure according to host taxon and environment. Kruskal-Wallis group and pairwise tests determined statistical significance for each category. Statistical differences between host

Figure 2. Boxplots of gut microbiome species alpha diversity by host taxon showing Shannon diversity index (A), number of observed OTUs (B), and Pielou E evenness (C). Classifications along the x-axis are A/Red=*C. aurita*, AH=*C. aurita* hybrid, G/Orange= *C. geoffroyi*, J/Green= *C. jacchus*, P/Yellow= *C. penicillata*, JP/Purple= *C. jacchus* x *C. penicillata* hybrids, and PG/ Blue= *C. penicillata* x *C. geoffroyi* hybrids.

Figure 3. Boxplots of gut microbiome species alpha diversity by host environment showing Shannon diversity index (A), number of observed OTUs (B), and Pielou E evenness (C). Classifications along the x-axis are C/Red=captive, T/Dark Blue=translocated, and W/Orange=wild.

sex, taxon, and environmental classifications were tested using adonis PERMANOVA test for beta group significance based on weighted UniFrac distances with the Qiime2-2020.2 adonis plug-in [48,49]. Weighted UniFrac principle coordinate analysis (PCoA) plots were generated according to host taxon and environment. Statistical significance of UniFrac PCoA plots within each category was determined by PERMANOVA group and pairwise tests [50]. All pairwise comparisons were corrected with the Benjamini–Hochberg procedure [51] (Benjamini and Hochberg, 1995). The corrected P-value was considered significant at p<0.05.

Taxonomic composition of samples was determined with the Qiime2-2018.6 q2-feature-classifier plug-in, which was trained on the Greengenes OTU database (version 13_8) [52] at the 99% level. Reads were limited to 500 nucleotides from the portion of the 16S V4 region bounded by the 515F/806R primer pair. Afterward, Qiime2-2018.6 was used to generate bar plots of bacterial taxonomic abundance based on host taxon and environment. The core gut OTUs that were present in at least 50% of all sampled hosts was determined with the Qiime2-2018.6 feature-table core-features visualizer and OTU taxonomic classifications were based on the q2-feature-classifier classifications. We used linear discriminant analysis coupled with effect size (LEfSe) on the Huttenhower lab Galaxy server (http://huttenhower.sph.harvard.edu/galaxy/), with LDA score>2.5 and P-value<0.05, to determine which OTUs have differential abundance between wild, captive, and translocated marmosets.

PICRUST 1.1.3 [53] was used for predictive functional profiling of bacterial communities via a phylogeny-based approach to correlate sampled bacterial species to the Greengenes 13_5 99% OTU reference dataset. Initially, the Greengenes reference data were used in QIIME2 to make a closed-reference OTU table from our sequence reads and converted to biom format. The OTU table was normalized by 16s rRNA copy number in PICRUST and the

extrapolated metagenome results then were input into HUMAnN 0.99 [54]. This program

reconstructs microbial pathway abundances from metagenomic data using the KEGG Orthology

catalogue [55-57]. The HUMAnN KEGG pathway module (functional units of genes) output was

used in LefSe [58] on the Huttenhower lab Galaxy server (http://huttenhower.sph.harvard.edu/

galaxy/) to identify how abundance of modules differ between marmosets categorized by taxon

and environment with linear discriminant analysis (LDA) coupled with effect size. These

differences were then plotted in GraPhlAn 0.9 [59].

Results

A total of 11,123,817 sequence reads was obtained with an average of 188,539 (129,170

+/- SD) reads per sample. After quality filtering, 9,063,712 reads remained with an average

153,622 (103,648 +/- SD) reads per sample. Afterward, merging of paired-end sequences

produced 843,615 reads, with an average of 142,891 (96,076 +/- SD) reads per sample. This

information is detailed in Supplementary Table 3.

Gut Microbiome Alpha Diversity based on Host Taxon and Environment

Rarefaction excluded a total of seven individuals from alpha diversity analysis

(Supplementary Table 1). Although taxa differed significantly in Shannon's diversity index

(Figure 2A; Kruskal-Wallis group test, H=16.88, p=0.01), Kruskal-Wallis pairwise tests were not

significant (Supplementary Table 4). The number of observed OTUs among taxa ranged from 61

to 571 (Supplementary Table 1), and median values varied significantly among taxa (Figure 2B;

Kruskal-Wallis group test, H=28.16, p=0.00). For pairwise comparisons, the median observed

Figure 4. PCoA of weighted Unifrac distances by host taxon (A) and environment (B). Abbreviations and color codes in plot legends are the same as in Figures 2-3.

OTUs in *C. jacchus* x *C. penicillata* hybrids were significantly different from that of *C. aurita* and *C. penicillata* x *C. geoffroyi* hybrids (Supplementary Table 5). Although taxa differed significantly in species evenness (Figure 2C; Kruskal-Wallis group test, H=16.41, p=0.01), subsequent pairwise tests were not significant (Supplementary Table 6). Shannon's index differed significantly among wild, translocated, and captive marmosets (Figure 3A; Kruskal-Wallis group test, H=9.85, p=0.007), and pairwise differences were significant between captive vs. translocated and captive vs. wild groups (Supplementary Table 7). Differences in median observed OTUs were significant among host environment categories (Figure 3B; Kruskal-Wallis group test, H=15.95, p=0.00), specifically between translocated vs. captive and translocated vs. wild (Supplementary Table 8). Species evenness differences were significant among host environments (Figure 3C; Kruskal-Wallis group test, H=10.03, p=0.007), with captive vs. translocated and captive vs. wild comparisons showing significant pairwise differences (Supplementary Table 9).

Gut Microbiome Beta Diversity based on Host Taxon and Environment

No differences in gut microbiome composition were observed between marmoset male and female hosts (adonis, F=1.03, df=2, R^2=0.04, p-value= 0.4). Gut microbiome composition between host taxa was significant when hybrids were included (adonis, F=2.66, df=2, R2=0.26, p-value=0.001), but no differences were detected between *Callithrix* species (adonis, F=0.76, df=3, R2=0.10, p-value=0.75). Lastly, 27% of variation in marmoset gut microbiome composition could be significantly attributed to differences between host environments (adonis,

Figure 5. Relative abundances of gut bacterial classes by environment. Each bar represents a marmoset host, and hosts are classified as C (captive), T (translocated), and W (wild). Colors within each bar represent the top 11 most abundant bacterial classes. Black represents all other bacterial classes. Top colors represent greater bacterial relative abundance than those taxa at the bottom of each bar.

F=8.88, df=6, R²=0.26, p-value=0.001). PCoA plots of weighted Unifrac distances also suggested that environment more strongly effects host differences in gut microbial composition than taxon affiliation (Figures 4A-B). The first PCoA axis explained 33.02% of gut microbial variation and the second axis explained 19.49%. In separating the samples by taxon (Figure 4A), some taxa showed loose clustering, but most of them showed interdispersion in the upper and lower right quadrants. Distances among taxa were significant (PERMANOVA, pseudo-F= 2.67, p-value=0.002), but there were few significant pairwise comparisons among them (Supplementary Table 10). Separation of hosts according to environment resulted in the most prominent clustering patterns between wild, translocated, and captive samples, particularly along the first axis (Figure 4B). Weighted Unifrac distances between samples by environment were significant at both the group level (PERMANOVA, pseudo-F=8.88, p-value=0.001), as well as for all pairwise comparisons (Supplementary Table 11). When only captive and translocated pure species hosts were analyzed together most samples clustered together along both the first and second components (Supplementary Figure 1A-B), regardless of species category. Group level (PERMANOVA, pseudo-F=1.20, p-value=0.257) nor pairwise weighted Unifrac distances were significant between host species (Supplementary Table 12). There were also no significant differences between captive and translocated hosts (PERMANOVA, pseudo-F=1.89, p-value=0.128).

Differential OTU Abundance based on Host Taxon and Environment

The core bacterial classes among hosts (Figure 5, Supplementary Figure 2; Supplementary Table 13) included Gammaproteobacteria, Epsilonproteobacteria, Bacteroidia,

Table 2. Core predictive functional bacterial KEGG modules present among all sampled marmoset hosts at a coverage of >90%

Module ID	KEGG Module Definition	Level 3 KEGG Brite Category
M00001	Glycolysis (Embden-Meyerhof pathway), glucose => pyruvate	Central carbohydrate metabolism
M00002	Glycolysis, core module involving three-carbon compounds	Central carbohydrate metabolism
M00003	Gluconeogenesis, oxaloacetate => fructose-6P	Central carbohydrate metabolism
M00004	Pentose phosphate pathway (Pentose phosphate cycle)	Central carbohydrate metabolism
M00006	Pentose phosphate pathway, oxidative phase, glucose 6P => ribulose 5P	Central carbohydrate metabolism
M00007	Pentose phosphate pathway, non-oxidative phase, fructose 6P => ribose 5P	Central carbohydrate metabolism
M00049	Adenine nucleotide biosynthesis, IMP => ADP/dADP,ATP/dATP	Purine Metabolism<
M00050	Guanine ribonucleotide biosynthesis IMP => GDPGTP	Purine Metabolism<
M00051	Uridine_monophosphate biosynthesis glutamine + PRPP => UMP	Pyrimidine Metabolism
M00115	NAD biosynthesis aspartate => NAD	Cofactor and vitamin biosynthesis
M00119	Pantothenate biosynthesis valine/L-aspartate => pantothenate	Cofactor and vitamin biosynthesis
M00123	Biotin biosynthesis, pimeloyl-CoA => biotin	Cofactor and vitamin biosynthesis
M00125	Riboflavin biosynthesis, GTP => riboflavin/FMN/FAD	Cofactor and vitamin biosynthesis
M00126	Tetrahydrofolate biosynthesis, GTP => THF	Cofactor and vitamin biosynthesis
M00157	F-type ATPase, bacteria	ATP Synthesis

M00164	ATP synthase	Photosynthesis
M00222	Phosphate transport system	Phosphate and Amino Acid Transport System
M00229	Arginine transport system	Phosphate and Amino Acid Transport System
M00299	Spermidine/putrescine transport system	Mineral and Organic Ion Transport System
M00360	Aminoacyl-tRNA_biosynthesis prokaryotes	Aminoacyl tRNA

Actinobacteria, Clostridia, and Alphaproteobacteria among others. Hosts showed variation in both bacterial class and abundance among individuals (Supplementary Figure 2), particularly in *C. aurita*, *C. geoffroyi*, and *C. penicillata*- taxa that were sampled across different locations and environments. Then, a strong pattern of bacterial taxonomic abundance (limited to class for brevity) emerges upon classifying hosts by environment (Figure 5). At the family level these patterns are dominated by: (1) Enterobacteriaceae (class Gammaproteobacteria) in captive hosts; (2) families Helicobacteraceae and Campylobacteraceae (class Epsilonproteobacteria), Bacteroidaceae (class Bacterioidia), and Sphingomonadaceae (class Alphaproteobacteria) in translocated hosts; and (3) Helicobacteraceae and Campylobacteraceae, Bacterioidia (family undetermined), Bifidobacteriaceae (class Actinobacteria), and Veillonellaceae (class Clostridia) in wild hosts.

LEfSe differential gut microbiome bacteria abundance testing between hosts found several microbe genera to differ between wild, captive, and translocated hosts (Supplementary Figure 3). The wild marmoset gut microbiome was enriched for 8 bacterial genera, including *Megasphaera*, *Campylobacter*, *Bifidobacterium* with the former having the highest LDA score. For captivity, 12 bacteria taxa were enriched for this host environmental category, with the Enterobacteriaceae family

having the highest LDA score followed by *Providencia.* For translocated hosts, 41 bacterial taxa were enriched for this host environment, with *Helicobacter* having the highest LDA score.

Predictive Metagenomic and Metabolic Characterization of Marmoset Gut Microbiome

Metagenome prediction by PICRUSt gave an average weighed NSTI value across hosts of 0.035 and host scores ranged from 0.008 to 0.147 (Supplementary Table 14). The 16S rRNA gene amplicon data analyzed in QIIME2 resulted in predicted proteins classified as KEGG orthologos (KOs) with a total of 6909 KOs across all hosts. Using these KOs with HUMAnN for metabolic reconstruction detected 113 KEGG modules (functional units of genes) collectively present in all samples (Supplementary Table 15). Among these modules, 20 were present across all samples at a coverage of >90% and were identified as core modules (Table 2). The core modules were found to be involved with the following Level 3 KEGG BRITE hierarchies: central carbohydrate metabolism (6 modules), purine metabolism (2 modules), pyrimidine metabolism (1 module), cofactor and vitamin biosynthesis (5 modules), ATP synthesis (1 module), photosynthesis (1 module), phosphate and amino acid transport system (2 modules), mineral and organic ion transport system (1 module), and aminoacyl tRNA (1 module).

Several KEGG modules were differently abundant for host taxonomy and environment (Figures 6-7, Supplementary Figures 4-5). A total of 57 modules differed for taxa (Supplementary Figure 4 and 5), with *C. jacchus x C. penicillata* hybrids having the largest number of differently abundant modules and *C. jacchus* possessing the second highest number. Differences among the three groups of host environment were not particularly concentrated around certain categories but rather spread out across clades of modules.

A:M00178 Ribosome bacteria
AA:M00051 Uridine_monophosphate biosynthesis glutamine + PRPP => UMP
BB:M00064 ADP-L-glycero-D-manno-heptose_biosynthesis
C:M00335 Sec secretion system
CC:M00060 Lipopolysaccharide biosynthesis KDO2-lipid A
D:M00331 Type II general secretion system
DD:M00061 Uronic acid metabolism
E:M00326 RTX toxin transport system
EE:M00001 Glycolysis Embden-Meyerhof pathway glucose => pyruvate
F: M00336   Twin-arginine translocation Tat system
G:M00229 Arginine transport system
GG:M00007 Pentose phosphate pathway oxidative phase fructose 6P => ribulose 5P
GH:M00006 Pentose phosphate pathway oxidative phase glucose 6P => ribulose 5P
H:M00222 Phosphate transport system
HH:M00009 Citrate_cycle_TCA_cycle_Krebs_cycle
I:M00237 Branched-chain amino acid transport system
II:M00002 Glycolysis core module involving three-carbon compound
J:M00277 PTS system N-acetylgalactosamine-specific II componentl
JJ:M00003 Gluconeogenesis oxaloacetate =>fructose-6P
K:M00230 Glutamate/aspartate transport system
KK:M00004 Pentose phosphate pathway Pentose phosphate cycle
L:M00323 Urea transport system
LL:M00166 Reductive pentose phosphate cycle RuBP + CO2 => glyceraldehyde-3P M00166
M:M00200 Histidine transport system
MM:M00165 Reductive pentose phosphate cycle Calvin cycle
N:M00322 Neutral amino acid transport system
NN:M00167 Reductive pentose phosphate cycle glyceraldehyde-3P => RuBP
O:M00219 AI-2 transport system
OO:M00174 Methane oxidation methylotroph methane => CO2
P:M00197 Putative fructooligosaccharide transport system
PP:M00360 Aminoacyl-tRNA_biosynthesis_prokaryotes
Q:M00194 Maltose/maltodextrin transport system
QQ:M00359 Aminoacyl-tRNA_biosynthesis_eukaryotes
R:M00202 Oligogalacturonide transport system
RR:M00311 2-oxoglutarate:ferredoxin oxidoreductas
S:M00207 Putative multiple sugar transport system
SS:M00149 Succinate_dehydrogenase_prokaryotes
T:M00200 Putative sorbitol/mannitol transport system
TT:M00150 Fumarate_reductase_prokaryotes
U:M00198 Putative sn-glycerol-phosphate transport system
UU:M00157 F-type ATPase bacteria
V:M00220 Rhamnose_transport_system
VV:M00156 Cytochrome_c_oxidase_cbb3-type
W:M00302 2-Aminoethylphosphonate transport system
WW:M00144 NADH quinone_oxidoreductase_prokaryotes
X:M00321 Bicarbonate transport system
XX:M00161 Photosystem_II
Y:M00185 Sulfate transport system
YY:M00164 ATP synthase
Z:M00300 Putrescine transport system
ZZ:M00163 Photosystem I
a:M00034 Methionine salvage pathway
aa:M00324 Dipeptide transport system
b:M00017 Methionine biosynthesis apartate => homoserine => methionine
bb:M00239 Peptides/nickel transport system
cccc:M0001lCitrate cycle second carbon oxidation 2-oxoglutarate => oxaloacetate
c:M00035 Methionine degradation
cc:M00349 Microcin C transport system
d: M0028   Ornithine biosynthesis glutamate_=> ornithine
dd:M00348 Glutathione transport system
e:M00025 Tyrosine biosynthesis chorismate => tyrosine
ee:M00245 Cobalt transport system
f: M0133   Polyamine biosynthesis arginine => agmatine =>_putrescine => spermidine
ff:M00317 Manganese/iron_transport_system
g:M00136 GABA biosynthesis prokaryotes putrescine => GABA
gg:M00276 PTS system mannose-specific II component
h:M00016 Lysine biosynthesis succinyl-DAP pathway aspartate => lysine
hh:M00278 PTS system sorbose-specific II component
i:M00019 Valine/isoleucine biosynthesis pyruvate => valine/2-oxobutanoate => isoleucine
ii:M00277 PTS system N-acetylgalactosamine-specific II component
j:M00026 Histidine biosynthesis PRPP => histidine M00026
k:M00124 Pyridoxal biosynthesis erythrose-4P => pyridoxal-5P
l:M00125 Riboflavin biosynthesis GTP => riboflavin/FMN/FAD
m:M00127 PTS system N-acetylgalactosamine-specific II componentl
n:M00119 Pantothenate biosynthesis valine/L-aspartate => pantothenate
o:M00123 Biotin biosynthesis pimeloyl-ACP/CoA => biotin
p:M00115 NAD biosynthesis aspartate => NAD
q:M00116 Menaquinone biosynthesis chorismate => menaquinone
u:M00117 Ubiquinone biosynthesis prokaryotes chorismate => ubiquinone
v:M00126 Tetrahydrofolate biosynthesis GTP => THF
w:M00048 Inosine monophosphate biosynthesis PRPP + glutamine => IMP
x:M00050 Guanine ribonucleotide biosynthesis IMP => GDPGTP
y:M00049 Adenine ribonucleotide biosynthesis IMP => ADPATP
z:M00053 Pyrimidine deoxyribonucleotide biosynthesis CDP/CTP => dCDP/dCTPdTDP/dTTP

Figure 6. HUMAnN metabolic reconstruction of marmoset fecal microbiome based on host environment. The cladogram shows 3 KEGG BRITE hierarchical levels with an outer, generalized level, and ending with the inner-level annotated by letters that are detailed further in the accompanying legend. Circles not differentially abundant in any species are clear. Environment classifications and colors follow the same abbreviations as Figures 3.

Figure 7. Differentially abundant KEGG metabolic modules inferred by LEfSe are colored by their associated environment classification, which follow the same abbreviations as Figures 3. The plot shows LDA scores resulting from LeFSe analysis for differentially abundant KEGG modules across host environment.

Discussion

Our results revealed that in marmosets as other animals, both host environment and taxonomy shape the gut microbiome [60-63]. However, host environment in this study showed stronger influence on marmoset GMC than host taxonomy, perhaps due to how closely related *Callithrix* species are as a recent primate radiation [64,65]. As host environment had the strongest effect on marmoset GMC within our study, the below discussion focuses largely on this factor.

The Wild *Callithrix* Gut Microbiome is Associated with Carbohydrate Metabolism

As exudivores, wild *Callithrix* regularly exploit tree gums or hardened saps [19,20,66,67]. Morphological adaptations of marmosets for exudivory include a well-developed caecum for fermentation of plant exudates [68] and a lower-anterior dentition "tooth scraper" to perforate tree bark and lianas [69]. In line with such morphological adaptations to exudate exploitation, our data show that the wild *Callithrix* gut microbiome is geared towards carbohydrate metabolism.

While wild and captive/translocated marmosets possessed similar classes of bacterial phyla (Table 3), bacterial abundance differed starkly between host environment. Unlike non-captive marmosets, wild marmosets had a GMC enriched for the Bifidobacteriaceae family (Actionobacterium), which includes genera like *Bifibacterium* that are common in animal gastrointestinal tracts and metabolize host-indigestible carbohydrates [70,71]. Wild marmosets gut microbiomes were also enriched for other bacteria involved in carbohydrate metabolism like *Megasphaera* and *Pectinatus* (both Firmicutes), which are actually known as important beer spoilage bacteria [72].

Table 3. Percentages of most abundant bacterial taxa present in marmoset hosts according to host environment

Phylum	Captive (%)	Translocated (%)	Wild (%)
Actinobacteria	6	6	32
Bacteroidetes	8	20	17
Firmicutes	6	7	14
Proteobacteria	76	63	35

The principle core predictive functional KEGG modules expressed in all sampled marmosets were related to central carbohydrate metabolism as well as purine metabolism, pyrimidine metabolism, cofactor and vitamin biosynthesis. Core predictive modules in both wild and non-wild hosts for carbohydrate metabolism were involved with glycolysis and the pentose phosphate pathway, pathways essential for the production of ATP and NADPH. However, the gut microbiome of wild marmosets was particularly enriched for predictive modules involved in carbohydrate metabolism including that of glycolysis and pentose phosphate biosynthesis, perhaps due to enrichment of gut microbes like Actinobacteria in wild hosts.

Captive Marmoset GMC Mirrors Aspects of Human GI Distress

While methodological differences between studies need to be taken into consideration, certain trends arise in GMC of captive marmosets in our study and captive *C. jacchus* from previous studies. Actinobacteria was not the most abundant gut bacterial taxa for most captive *Callithrix* and there is increased abundance of gut Proteobacteria among captive marmosets here and other studies [36,70]. One study of captive *C. jacchus* uniquely compared gut microbiome composition between a specific pathogen free colony, housed under conditions that minimized outside contamination, with a

conventionally housed group [34]. Many barrier marmosets had relatively high abundance of *Bifibacterium* (approximately 1% to 80%)*,* while *Fusobacterium* B (phylum Fusobacteria) was dominant in conventionally housed hosts [34].

The GMC of captive and translocated marmosets shows similarity to certain aspects of the GMC of human GI diseases associated with gut microbiome dysbiosis [7,18,73,74]. Crohn's Disease (CD) generally affects the terminal ileum and colon [75,76] and GMC of CD sufferers is less diverse relative to healthy humans [18,77,78]. The gut microbiome of CD patients is enriched for bacterial taxa that include Enterobacteriaceae and Veillonellaceae, and decreased in Bifidobacteriaceae [18,79]. We also observed similar patterns in non-wild marmosets, relative to wild counterparts. Enterobacteriaceae is frequently associated with intestinal diseases and contains a number of pathogenic bacterial strains of *Salmonella*, *Escherichia*, and *Shigella* [80,81] and potentially pathogenic Enterobacteriaceae were previously found in captive and urban *Callithrix* [82]. Translocated marmosets GMC showed higher abundance of Bacteroidetes and Clostridia, a pattern also seen in ulcerative colitis [79], a GI disease which usually involves rectal and colon inflammation [75,76].

Importance of *Bifidobacterium* and Natural Diets for the Exudivore Gut

Data from this and previously published studies suggest that *Bifidobacterium* is an important gut microbe for animal hosts that nutritionally exploit plant exudates. Further, high abundance of *Bifidobacterium* in gut microbiomes of exudivore specialists may be a biomarker for host gut microbiome eubiosis. In the specialist exudivore the pygmy loris (*Nycticebus pygmaeus*), two wild individuals possessed Actinobacteria (10.98%) as major gut microbiome bacterial phyla [83], and gut microbiome function was enriched for carbohydrate metabolism. *Bifidobacterium* was also the most abundant and enriched gut bacterial taxa in wild individuals of *Nycticebus javanicus,* another primate

exudivore [84]. The genomes of three isolates of *Bifidobacterium callitrichos* from a captive *C. jacchus* fecal sample contained predicted genes associated with galactose and arabinose metabolism, which are major constituents of tree gums eaten by *C. jacchus* [70]. In 3 US captive facilities, *C. jacchus* collectively shared four species of *Bifidobacterium* which possessed genes encoding ATP-binding cassette proteins, which are important for nutrient transport that may be specific to marmoset gut [32].

One concern for a lack of tree gums in the diet of captive specialist exudivores is the development of health issues as well as a negative impact on breeding and survivability [85,86]. Current captive *Callithrix* husbandry, often omits gums from the marmoset diet or uses it as an enrichment ingredient (Table 4) [86]. In humans suffering from GI diseases, increasing plant-based foods and dietary fiber, resulted in increasing microbiome diversity, remission of GI symptoms, and decreasing risk of GI [7,87]. Such diets may increase gut abundance of bacteria such as *Bifibacterium* that produce short chain fatty acids like butyrate, which may guard against proliferation of pathogenic bacteria in the gut and decrease chronic inflammation [7,87]. Lack of access to a natural-wild diet for marmosets and other exudivory specialists may also promotes loss of native gut microbe *Bifidobacterium* and enrichment of potentially pathogenic bacterial strains of Enterobacteriaceae.

Table 4. Diet collectively fed to marmoset hosts in sampled captive facilities

Fruits	Papaya, Orange, Banana, Apple, Pear, Avacado, Kiwi, Melon, Mango
Carbs	Sweet Potato, Potato, Beets
Veggies	Cucumber, Eggplant, Pumpkin, Chuchu, Cauliflower, Carrots
Proteins	Cooked Chicken, Cooked Egg

Animal studies increasingly recognize the need to include host-microbiome associations in management of captive populations [2,17,88]. It has also been demonstrated, as in mice and wood-rats,

that feeding more natural diets to individuals in captivity helped maintain host GMC profiles closer to wild, free-ranging hosts [88,89]. Plausibly, GI distress in captive *Callithrix* may be related to unnatural diets that lower abundance of *Bifidobacterium* and increase presence of potentially pathogenic bacteria Enterobacteriacea thereby driving dysbiosis in the gut microbiome. Thus, based on such observations, captive facilities should consider provisioning of gum in the diet of specialized exudivores to improve host welfare by maintaining gut microbiomes closer to that of wild populations.

Conclusion

The results of this study have given us an initial genus-level look into the gut microbiome of specialized exudivorous NHPs. Our major findings indicate that, just as in other primates, the gut microbiome of *Callithrix* is sensitive to environmental factors, and these findings likely apply more broadly to other animal exudivores. Our findings also suggest that carbohydrate metabolism is a core function of the *Callithrix* gut microbiome. Yet, although marmosets are specialized exudivores, there is a wide gap between the nutritional intake of marmosets in the wild and captivity, where the marmoset physiology for gummivory is generally not considered. From an evolutionary and health point of view, this disconnect between wild and captive *Callithrix* diets leaves open several questions concerning maladaptation of the marmoset digestive system, perturbation of gut microbiota, malabsorption of nutrients, and long term health effects. More studies are needed to understand better the health and reproductive consequences of omitting as well as increasing gum intake by specialized exudivores in captivity. Also, systematic studies are needed to better understand whether *Bifidobacterium* plays a central role in maintaining euobisis in the gut microbiomes of *Callithrix* and other exudivory specialists. Overall, such information will expand baseline gut microbiome data available for wild and non-wild

exudivores to allow for the development of new tools to improve the management, welfare, and

conservation of animal exudivores.

Funding

This work was supported by a Brazilian CNPq Jovens Talentos Postdoctoral Fellowship, an American Society of Primatologists Conversation Small Grant, and an International Primatological Society Research Grant.

Acknowledgements

We thank Vanner Boere, Ita de Oliveira e Veronica Souza, Rodrigo S. Carvalho, the Guarulhos Zoo staff, the CEMAFAUNA staff, the CPRJ staff, SERCAS staff, and AMLD staff for assistance with wild and captive populations. We are very grateful to Cecil M. Lewis and Tanvi Honap and the LMAMR staff at University of Oklahoma for donation of PCR primers. We thank Dietmar Zinner, Corinna Ross, and two anonymous reviewers for comments on prior versions of this work.

References

1. Human Microbiome Project Consortium. Structure, function and diversity of the healthy human microbiome. Nature 2012;486(7402):207-14.
2. West AG, Waite DW, Deines P, Bournec DG, Digby A, Kenziee VJ, Taylor MW. The microbiome in threatened species conservation. Biological Conservation. 2019; 229:85–98. (doi:10.1016/j.biocon.2018.11.016f).
3. Cho I, Blaser MJ. The human microbiome: at the interface of health and disease. Nat Rev Genet. 13;13(4):260–70. (doi.org/10.1038/nrg3182).
4. Nicholson JK, Holmes E, Kinross J, Burcelin R, Gibson G, Jia W, Pettersson S. Host-gut microbiota metabolic interactions. Science. 2012;336;1262-1267. (doi:10.1126/science.1223813).
5. Amato KR, Yeoman CJ, Kent A, Righini N, Carbonero F, Estrada A, Gaskins HR, Stumpf RM, Yildirim S, Torralba M, et al. Habitat degradation impacts black howler monkey (*Alouatta pigra*) gastrointestinal microbiomes. ISME J. 2013; 7:1344–1353. (doi:10.1038/ismej.2013.16).
6. Clayton JB, Vangay P, Huang H, Ward T, Hillmann BM, Al-Ghalith GA, Travis DA, Long HL, Tuan BV, Minh VV, et al. Captivity humanizes the primate microbiome. Proc Natl Acad Sci USA. 2016;113(37):10376–81. (doi:10.1073/pnas.1521835113).
7. DeGruttola AK, Low D, Mizoguchi A, Mizoguchi E. Current understanding of dysbiosis in disease in human and animal models. Inflammatory Bowel Diseases. 2016;22(5):1137–50. (doi:10.1097/MIB.0000000000000750).
8. Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, Schlegel ML, Tucker TA, Schrenzel MD, Rob Knight R, et al. Evolution of mammals and their gut microbe. Science. 2008; 320;1647–1651. (doi: 10.1126/science.1155725).
9. Guo W, Mishra S, Wang C, Zhang H, Ning R, Kong F, Zeng B, Zhao J, Li Y. Comparative study of gut microbiota in wild and captive giant pandas (*Ailuropoda melanoleuca*). Genes. 2019; 10(10):827.

10. Youngblut ND, Reischer GH, Walters W, Schuster N, Walzer C, Stalder G, RLey RE, Andreas H. Farnleitner AH. Host diet and evolutionary history explain different aspects of gut microbiome diversity among vertebrate clades. Nat Commun. 2019;10:2200. (doi:10.1038/s41467-019-10191-3).

11. McKenzie VJ, Song SJ, Delsuc F, Prest TL, Oliverio AM, Korpita TM, Alexiev A, Amato KR, Metcalf JL, Kowalewski M,et al. The Effects of Captivity on the Mammalian Gut Microbiome. Integrative and Comparative Biology. 2017;57(4):690–704. (doi:10.1093/icb/icx090).

12. Gao H, Chi X, Qin W, Wang L, Song P, Cai Z, Zhang J, Zhang T. Comparison of the gut microbiota composition between the wild and captive Tibetan wild ass (*Equus kiang*). J Appl Microbiol. 2019;126(6):1869–78. (doi.org/10.1111/jam.14240)

13. Gibson KM, Nguyen BN, Neumann LM, Miller M, Buss P, Daniels S, Ahn MJ, Crandall KA, Pukazhenthi B. Gut microbiome differences between wild and captive black rhinoceros – implications for rhino health. Scientific Reports. 2019;9(1):7570. (doi:10.1038/s41598-019-43875-3).

14. Sun CH, Liu HY, Liu B, Yuan BD, Lu CH. Analysis of the gut microbiome of wild and captive Père David's deer. Front Microbiol. 2019;10. (doi:10.3389/fmicb.2019.02331).

15. Frankle JS, Mallott EK, Hopper LM, Ross SR, Amato KR. 2019. The effect of captivity on the primate gut microbiome varies with host dietary niche. *American Journal of Primatology.* 81:e23061. (doi:10.1002/ajp.23061).

16. Gevers D, Kugathasan S, Denson LA, Vázquez-Baeza Y, Van Treuren W, Ren B, Schwager E, Knights D, Song SJ, Moran Yassour M, et al. The treatment-naive microbiome in new-onset Crohn's disease. Cell Host and Microbe. 2014;15(3):382–92. (doi:10.1016/j.chom.2014.02.005).

17. Pineda-Munoz S, Alroy J. Dietary characterization of terrestrial mammals. Proc. R. Soc. B 2014;281: 20141173.

18. Cabana F, Dierenfeld ES, Donati G, Nekaris K. Exploiting a readily available but hard to digest resource: a review of exudativorous mammals identified thus far and how they cope in captivity. Integr Zool. 2018;13:94–111.

19. Nash LT. Dietary, behavioral, and morphological aspects of gummivory in primates. Yearb Phys Anthropol. 1986;29:113-137.

20. Smith A. Exudativory in primates: interspecific patterns. In: Burrow AM, Nash LT, editors. The Evolution of Exudativory in Primates. New York: Springer-Verlag. 2010. pp. 45-87.

21. Marini R, Wachtman L, Tardif S, Mansfield K, Fox J, editors. The common marmoset in captivity and biomedical research. London:Academic Press. 2018.

22. Carvalho RS, Fransen SJ, Valença-Montenegro MM, Dunn NJ, Igayara-Souza CA, Port-Carvalho M, Dominic Wormell, Melo FR, Silva A, Lacerda WR, Jerusalinsky L. *Callithrix aurita*. In: Schwitzer C, Mittermeier RA, Rylands AB, Chiozza F, Williamson EA, Byler D, Wich S, Humle T, Johnson C, Mynott H, and G. McCabe G, editors. Primates in Peril: The World's 25 Most Endangered Primates. Washington, DC:IUCN SSC Primate Specialist Group, International Primatological Society, Global Wildlife Conservation, and Bristol Zoological Society. 2018–2020; p. 78-81.

23. Silva FFR, Malukiewicz J, Silva LC, Carvalho RS, Ruiz-Miranda, CR, Coelho FAS, Figueira MP, Boere V, Silva IO. A survey of wild and introduced marmosets (*Callithrix*: Callitrichidae) in the southern and eastern portions of the state of Minas Gerais, Brazil. Primate Conservation. 2018;32:1–18.

24. Malukiewicz J. A Review of experimental, natural, and anthropogenic Hybridization in *Callithrix* Marmosets. Int J Primatol. 2018; 40(1):72–98. (doi.org/10.1007/s10764-018-0068-0).

25. Ialeggio DM, Baker AJ. Results of a preliminary survey into wasting marmoset syndrome in callitrichid collections. Toronto, Ontario, Canada.1995;148–158.

26. Ludlage E, Mansfield K. Clinical care and diseases of the common marmoset (*Callithrix jacchus*). Comparative Medicine. 2003;53:369–382.

27. Nakashima E, Okano Y, Niimi K, Takahashi E. Detection of calprotectin and apoptotic activity in the colon of marmosets with chronic diarrhea. The Journal of Veterinary Medical Science.2013;75:1633–1636. (doi: 10.1292/jvms.13-0257).

28. Yoshimoto T, Niimi K, Takahashi, E. Tranexamic acid and supportive measures to treat wasting marmoset syndrome. Comparative Medicine. 2016;66:468–473.

29. Jiménez RR, Sommer S. The amphibian microbiome: natural range of variation, pathogenic dysbiosis, and role in conservation. Biodivers Conserv. 2017;26(4):763–86. (doi:10.1007/s10531-016-1272-x).

30. Endo A, Futagawa-Endo Y, Schumann P, Pukall R, Dicks LMT. *Bifidobacterium reuteri* sp. nov., *Bifidobacterium callitrichos* sp. nov., *Bifidobacterium saguini* sp. nov., *Bifidobacterium stellenboschense* sp. nov. and *Bifidobacterium biavatii* sp. nov. isolated from faeces of common marmoset (*Callithrix jacchus*) and red-handed tamarin (*Saguinus midas*). Systematic and Applied Microbiology. 2012;(2):92–7. (doi: 10.1016/j.syapm.2011.11.006).

31. Modesto M, Watanabe K, Arita M, Satti M, Oki K, Sciavilla P, Patavino C, Cammà C, Michelin S, Sgorbati B, Mattarelli P. *Bifidobacterium jacchi* sp. nov., isolated from the faeces of a baby common marmoset (*Callithrix jacchus*). International Journal of Systematic and Evolutionary Microbiology. 2019;69(8):2477–85. (doi: 10.1099/ijsem.0.003518).

32. Brown CJ, Mtui D, Oswald BP, Van Leuven JT, Vallender EJ, Schultz-Darken N, Ross CN, Tardif SD, Austad SN, Forney LJ. Comparative genomics of *Bifidobacterium* species isolated from marmosets and humans. Am J Primatol. 2019;81:10–11. (doi:10.1002/ajp.22983).

33. Lugli GA, Milani C, Duranti S, Alessandri G, Turroni F, Mancabelli L, Tatoni D, Ossiprandi MC, van Sinderen D, Ventura M. Isolation of novel gut bifidobacteria using a combination of metagenomic and cultivation approaches. Genome Biol. 2019;20(1). (doi: 10.1186/s13059-019-1711-6).

34. Ross CN, Austad S, Brasky K, Brown CJ, Forney LJ, Gelfond JA, Landord R, Richardson A, Tardif SD. The development of a specific pathogen free (SPF) barrier colony of marmosets (*Callithrix jacchus*) for ageing research. Aging (Albany NY). 2017;9(12):2544–58.

35. Reveles KR, Patel S, Forney L, Ross CN. Age-related changes in the marmoset gut microbiome. Am J Primatol. 2019;81(2):e22960. (doi.org/10.1002/ajp.22960).

36. Zhu L, Clayton JB, Haute MJSV, Yang Q, Hassenstab HR, Mustoe AC, Knights D, Benson AK, French JA. Sex bias in gut microbiome transmission in newly paired marmosets (*Callithrix jacchus*). mSystems. 2020;5(2). (doi:10.1128/mSystems.00910-19).

37. Coimbra-Filho AF, Pissinatti A, Rylands AB. Experimental multiple hybridism and natural hybrids among *Callithrix* species from eastern Brazil. In: Rylands A, editor. (Marmosets and tamarins: Systematics, behaviour, and ecology. New York: Oxford University Press.1993; p 95–120.

38. Fuzessy LF, Silva I de O, Malukiewicz J, Silva FFR, Pônzio M do C, Boere V, Ackermann RR. Morphological variation in wild marmosets (*Callithrix penicillata* and *C. geoffroyi*) and their hybrids. Evol Biol. 2014;41(3):480–93. (doi: 10.1007/s11692-014-9284-5).

39. Malukiewicz J, Boere V, Fuzessy LF, Grativol AD, French JA, Silva I de O, Pereira LCM, Ruiz-Miranda CR, Valença YM, Stone AC. Hybridization effects and genetic diversity of the common and black-tufted marmoset (*Callithrix jacchus* and *Callithrix penicillata*) mitochondrial control region. Am J Phys Anthropol. 2014;155(4):522–36. (doi:10.1002/ajpa.22605).

40. Carvalho RS. Conservação do saguis-da-serra-escuro (*Callithrix aurita* (Primates)) – Analise molecular e colormetrica de populações do gênero *Callithrix* e seus híbridos. PhD dissertation 2015, Universidade do Estado do Rio de Janeiro.

41. Zoetendal EG, Heilig HG, Klaassens ES, Booijink CC, Kleerebezem M, Smidt H, de Vos WM. Isolation of DNA from bacterial samples of the human gastrointestinal tract. J Nat Protoc. 2006;1:870–873. (doi: 10.1038/nprot.2006.142).

42. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J 2012;6(8):1621-4.

43. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, Huttley GA, et al., QIIME allows analysis of high-throughput community sequencing data. Nat Methods. 2010;7;335–336. (doi:10.1038/nmeth.f.303).

44. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13(7):581–3. (doi.org/10.1038/nmeth.3869).

45. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately maximum-likelihood trees for large alignments. PLOS ONE. 2010;5:e9490. (doi: 10.1371/journal.pone.0009490).

46. Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7: improvements in performance and usability. Molecular Biology and Evolution. 2013;30:772–780. (doi:10.1093/molbev/mst010).

47. Lozupone CA, Hamady M, Kelley ST, Knight R. Quantitative and Qualitative β diversity measures lead to different insights into factors that structure microbial communities. AEM. 2007;73(5):1576–85. (doi.org/10.1128/AEM.01996-06).

48. Anderson MJ. A new method for non-parametric multivariate analysis of variance. Austral ecology. 26:32–46, 2001. (doi:10.1111/j.1442-9993.2001.01070.pp.x).

49. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, M. Stevens MHH, Szoecs E, Helene Wagner H. Vegan: community ecology package. 2018. R package version 2.5-3.

50. Anderson MJ. Permutational multivariate analysis of variance (PERMANOVA). Wiley StatsRef: Statistics Reference Online 2017. (doi:10.1002/9781118445112.stat07841).

51. Benjamini, Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing, J R Stat Soc B. Series B (Methodological).1995;57:289–300.

52. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen GL. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. Appl Environ Microb. 2006;72(7): 5069-5072.

53. Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, Clemente JC, Burkepile DE, Thurber RLV, Knight R, et al. Predictive functional profiling of microbial communities using 16S rRNA marker gene sequences. Nature Biotechnology. 2013;31(9):814–21. (doi: 10.1038/nbt.2676).

54. Abubucker S, Segata N, Goll J, Schubert AM, Izard J, Cantarel BL, Rodriguez-Mueller B, Zucker J, Thiagarajan M, Henrissat B, et al. Metabolic reconstruction for metagenomic data and

its application to the Human Microbiome. PLoS Comput Biol. 2012;8(6):e1002358. (doi:10.1371 journal.pcbi.1002358).

55. Kanehisa M, Goto S. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 2002;**28(1):**27-30.

56. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Res. 2017;45(D1):D353–61. (doi:10.1093/nar/gkw1092).

57. Kanehisa M, Sato Y, Furumichi M, Morishima K, Tanabe M. New approach for understanding genome variations in KEGG. Nucleic Acids Res. 2019;47(D1):D590–5. (doi:10.1093/nar/gky962).

58. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C.Metagenomic biomarker discovery and explanation. Genome Biology. 2011.12:R60. (doi:10.1186/gb-2011-12-6-r60).

59. Asnicar F, Weingart G, Tickle TL, Huttenhower C, Segata N. Compact graphical representation of phylogenetic data and metadata with GraPhlAn. PeerJ. 2015;3:e1029. (doi: 10.7717/peerj.1029).

60. Colman DR, Toolson EC, Takacs-Vesbach CD. Do diet and taxonomy influence insect gut bacterial communities? Mol Ecol. 2012;21(20):5124–37. (doi:10.1111/j.1365-294X.2012.05752.x).

61. Sullam KE, Essinger SD, Lozupone CA, O'Connor MP, Rosen GL, Knight R, Kilham SS, Russell JA. Environmental and ecological factors that shape the gut bacterial communities of fish:a meta-analysis. Mol Ecol. 2012;21(13):3363-78. (doi:10.1111/j.1365-294X.2012.05552.x).

62. Kohl KD, Skopec MM, Dearing MD. Captivity results in disparate loss of gut microbial diversity in closely related hosts. Conserv Physiol. 2014;2(1). (doi: 10.1093/conphys/cou009).

63. Hale VL, Tan CL, Niu K, Yang Y, Knight R, Zhang Q, Cui D, Amato KR. Diet versus phylogeny: a comparison of gut microbiota in captive colobine monkey species. Microb Ecol. 2018;75(2):515–27. (doi: 10.1007/s00248-017-1041-8).

64. Malukiewicz J, Hepp CM, Guschanski K, Stone AC. Phylogeny of the I group of *Callithrix* marmosets based on complete mitochondrial genomes. Am J Phys Anthropol. 2016 20;162(1):157–69. (doi.org/10.1002/ajpa.23105 doi: 10.1002/ajpa.23105).

65. Malukiewicz J, Cartwright RA, Curi NHA, Dergam JA, Igayara CS, Moreira SB, Molina CV, Nicola PA, Noll A, Passamani M, Pereira LCM, Pissinatti A, Ruiz-Miranda CR, Silva DL, Stone AC, Dietmar Zinner D, Roos C. Mitogenomic phylogeny of *Callithrix* with special focus on human transferred taxa. BMC Genomics. (in review).

66. Martins MM, Setz EZF. Diet of buffy tufted-eared marmosets (*Callithrix aurita*) in a forest fragment in Southeastern Brazil. International Journal of Primatology. 2000,;21:467–476. (doi:10.1023/A:1005491903220).

67. Castro CSS, Araujo A. Diet and feeding behavior of marmoset, *Callithrix jacchus*. Braz. J. Ecol. 2007; 7:14–19.

68. Caton JM, Hill DM, Hume ID, Crook GA. The digestive strategy of the common marmoset, *Callithrix jacchus*. Comparative Biochemistry and Physiology Part A: Physiology. 1996;114(1):1–8. doi:10.1016/0300-9629(95)02013-6.

69. Coimbra-Filho AF, Mittermeier RA (1976). Tree-gouging, exudate-eating, and the 'short-tusked' condition in *Callithrix* and *Cebuella*. In Kleinman DG, editor. The biology and conservation of the Callitrichidae. Washington, DC: Smithsonian Institution Press. 1976; p.105-15.

70. Albert K, Rani A, Sela DA. The comparative genomics of *Bifidobacterium callitrichos* reflects dietary carbohydrate utilization within the common marmoset gut. Microbial Genomics. 2018;4(6):e000183. (doi:10.1099/mgen.0.000183).

71. O'Callaghan A, van Sinderen D. *Bifidobacteria* and their role as members of the human gut microbiota. Front Microbiol. 2016;7. (doi:10.3389/fmicb.2016.00925).

72. Paradh AD. Gram-negative spoilage bacteria in brewing. In: Hill AE, editor.Woodhead Publishing Series in Food Science, Technology and Nutrition,Brewing Microbiology,Woodhead Publishing. 2015. p. 175-194.

73. Ott SJ, Musfeldt M, Wenderoth DF, Hampe J, Brant O, Fölsch UR, Timmis KN, Schreiber S. Reduction in diversity of the colonic mucosa associated bacterial microflora in patients with active inflammatory bowel disease. Gut. 2004;53(5):685–93. (doi:10.1136/gut.2003.025403).

74. Manichanh C, Rigottier-Gois L, Bonnaud E, Gloux K, Pelletier E, Frangeul L, Nalin R, Jarrin C, Chardon P, Marteau P, Roca J, Dore J.Reduced diversity of faecal microbiota in Crohn's disease revealed by a metagenomic approach. Gut. 2006;55(2):205–11. (doi:10.1136/gut.2005.073817).

75. Abraham C, Cho JH. Inflammatory Bowel Disease. N Engl J Med. 2009; 361(21):2066–78. (doi:10.1056/NEJMra0804647).

76. Baumgart DC, Sandborn, WJ. (2012) Crohn's disease. The Lancet. 2012;380:1590–1605. (doi: 10.1016/S0140-6736(12)60026-9).

77. Pérez-Brocal V, García-López R, Nos P, Beltrán B, Moret I, Moya A. Metagenomic Analysis of Crohn's Disease Patients Identifies Changes in the Virome and Microbiome Related to Disease Status and Therapy, and Detects Potential Interactions and Biomarkers. Inflamm Bowel Dis. 2015(11):2515–32. (doi: 10.1097/MIB.0000000000000549).

78. Pascal V, Pozuelo M, Borruel N, Casellas F, Campos D, Santiago A, Martinez X, Varela E, Sarrabayrouse G, Machiels K, et al. A microbial signature for Crohn's disease. Gut. 2017;66(5):813–22. (doi: 10.1136/gutjnl-2016-313235).

79. Alam, MT, Amos GCA, Murphy ARJ, Murch S, Wellington EMH, Arasaradnam RP. Microbial imbalance in inflammatory bowel disease patients at different taxonomic levels. Gut Pathog. 2020;12:1. (doi:10.1186/s13099-019-0341-6).

80. Paterson DL. Resistance in gram-negative bacteria: enterobacteriaceae. Am J Med. 2006;119:S20-28. (doi:10.1016/j.amjmed.2006.03.013).

81. Kang E, Crouse A, Chevallier L, Pontier SM, Alzahrani A, Silué N, Campbell-Valois FX, Montaguelli X, Guehneid S, Malo D. Enterobacteria and host resistance to infection. Mamm Genome. 2018;29(7):558–76. (doi:10.1007/s00335-018-9749-4).

82. Verona CES. Parasitos em sagüi-de-tufo-branco (*Callithrix jacchus*) no Rio de Janeiro. PhD Dissertation 2015. Fundação Oswaldo Cruz.

83. Xu B, Xu W, Yang F, Li J, Yang Y, Tang X, Mu Y, Zhou J, Huang Z. Metagenomic analysis of the pygmy loris fecal microbiome reveals unique functional capacity related to metabolism of aromatic compounds. PLoS ONE. 2013;8(2):e56565. (doi:10.1371/journal.pone.0056565).

84. Cabana F, Clayton JB, Nekaris KAI, Wirdateti W, Knights D, Seedorf H. Nutrient-based diet modifications impact on the gut microbiome of the Javan slow loris (*Nycticebus javanicus*). Sci Rep. 2019;9(1). (doi:10.1038/s41598-019-40911-0).

85. Cabana F, Nekaris KAI. Diets high in fruits and low in gum exudates promote the occurrence and development of dental disease in pygmy slow loris (*Nycticebus pygmaeus*). Zoo Biol. 2015; 34:547–553. (doi: 10.1002/zoo.21245).

86. Cabana F, Maguire R, Hsu CD, Plowman A. Identification of possible nutritional and stress risk factors in the development of marmoset wasting syndrome. Zoo Biol. 2018b;37(2):98-106.

87. Grosse CSJ, Christophersen CT, Devine A, Lawrance IC. The role of a plant-based diet in the pathogenesis, etiology and management of the inflammatory bowel diseases. Expert Review of Gastroenterology & Hepatology. 2020;14(3):137–45. (doi: 10.1080/17474124.2020.1733413)

88. Leeuwen P van, Mykytczuk N, Mastromonaco GF, Schulte-Hostedde AI. Effects of captivity, diet, and relocation on the gut bacterial communities of white-footed mice. Ecology and Evolution. 2020;10(11):4677–90. (doi: 10.1002/ece3.6221).

89. Martínez-Mota R, Kohl KD, Orr TJ, Dearing MD. Natural diets promote retention of the native gut microbiota in captive rodents. The ISME Journal. 2020;14:67-78. (doi: 10.1038/s41396-019-0497-6).

Supplementary Figure 1. PCoA of weighted Unifrac distances by host taxon (A) and environment (B) for non-hybrid individuals only. Classifications along the x-axis are A/Red=*C. aurita*, G/blue= *C. geoffroyi*, J/Orange= *C. jacchus,* P/blue= *C. penicillata.* In panel A, classifications along the x-axis are C/Red=captive, T/Dark Blue=translocated, and W/Orange=wild. In panel B, classifications along the x-axis are C/Red=captive, T/Dark Blue=translocated.

Supplementary Figure 2. Relative abundances of gut bacterial classes by taxon. Each bar represents a marmoset host, and hosts are classified as A=*C. aurita*, AH=*C. aurita* hybrid, G=*C. geoffroyi*, J= *C. jacchus*, P=*C. penicillata*, JP=*C. jacchus* x *C. penicillata* hybrids, and PG=*C. penicillata* x *C. geoffroyi* hybrids. Top colors represent greater bacterial relative abundance than those taxa at the bottom of each bar.

Supplementary Figure 3. Differentially abundant bacterial abundances inferred by LEfSe are colored by their associated environment classification, C/Red=Captive, T/Green=Translocated, W/Blue=Wild.

Supplementary Figure 4. HUMAnN metabolic reconstruction of marmoset fecal microbiome based on host taxon. The cladogram shows 3 KEGG BRITE hierarchical levels with an outer, generalized level, and ending with the inner-level annotated by letters that are detailed further in the accompanying legend. Circles not differentially abundant in any species are clear. Taxon classifications and colors are as AH/ Red=*C. aurita* hybrid, J/Green=*C. jacchus*, P/Purple=*C. penicillata*, JP/Blue=*C. jacchus* x *C. penicillata* hybrids, PG/sea foam green=*C. penicillata* x *C. geoffroyi.*

Supplementary Figure 5. Differentially abundant KEGG metabolic modules inferred by LEfSe are colored by their associated host taxon classification, which follow the same abbreviations as Supplementary Figure 4. The plot shows LDA scores resulting from LeFSe analysis for differentially abundant KEGG modules across host taxa.