Common neural and transcriptional correlates of inhibitory control across emotion, memory, and response inhibition

Wei Liu[1], Nancy Peeters[1], Guillén Fernández[1], Nils Kohn[1]

1. Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Centre, The Netherlands

Correspondence:

Wei Liu

Department of Cognitive Neuroscience

Donders Institute for Brain, Cognition and Behaviour

Radboud University Medical Centre

Trigon Building, Kapittelweg 29

6525 EN Nijmegen, The Netherlands

Tel: +31 (0)24 36 53276

E-mail: W.Liu@donders.ru.nl

Abstract

Inhibitory control is crucial for regulating emotions and it may potentially also control memory retrieval.
Here, we provide neurobiological and transcriptional evidence to link emotional and memory control, and
thereby we identify neural networks and molecular mechanisms as potential targets for interventions.
Based on MRI coordinates from 95 published fMRI studies, meta-analytic approaches revealed a common
role of the right inferior parietal lobule embedded in a frontal-parietal-insular network during emotion and
memory control, which is similarly recruited during response inhibition tasks. These co-activation patterns
do also overlap with the neural networks associated with "inhibition", "cognitive control", and "working
memory". Linking neural commonalities to transcriptional commonalities using the Allen Human Brain
Atlas, we demonstrate that emotion regulation and memory control-related brain activity patterns are
associated with transcriptional profiles of a specific set of genes. As a control, we confirmed that
transcriptional correlates are more similar between networks defined by inhibition-related tasks compared
to transcriptional overlap between this inhibition network and the default mode network. Gene ontology
enrichment analysis of "inhibition-related" genes revealed associations with glutamatergic synapse,
neuropeptide-related activity and risk for major psychiatric disorders as well as Parkinson's disease. In
summary, this study identified a neural network and a set of genes associated with inhibitory control
across emotion regulation, memory control, and response inhibition. These findings facilitate our
understanding of neurobiological correlates of inhibitory control and may contribute to novel brain
stimulation and pharmacological interventions in individuals with such deficit.

                                      MAIN TEXT

Introduction

One of the most influential cognitive theories of emotion regulation proposes that it is primarily supported
by inhibitory control (*1*, *2*). A potentially related cognitive process, memory control, is defined as an
ability to actively reduce the accessibility of memories. Memory control is also thought to be supported by
inhibitory control (*3–5*). Recently, Engen and Anderson summarized the conceptual link between emotion
regulation and memory control, and raised the question whether there is a common neurobiological
mechanism supporting these two processes (*6*). Here, we set out to empirically demonstrate the
neurobiological commonalities between the two processes by first analyzing fMRI data and then linking
neuroimaging findings with post mortem gene expression data.

The idea that inhibitory control, as the fundamental cognitive process, supports other higher-level
processes such as emotion regulation and memory control is supported by behavioral and neural evidence.
Behaviorally, a positive correlation between emotion regulation and memory control performances has
been found (*7*). Human neuroimaging results showed overlapping recruitment of right superior medial
frontal gyrus (rSFG), right inferior frontal gyrus (rIFG) in both emotion regulation (*8–10*) and memory
control (*5*, *11*). Beyond these frontal regions, the parietal cortex is another candidate region that may play
a critical role in both emotion regulation and memory control. Starting from the idea of multiple-demand
system in the brain (*12*), both the evidence from task-fMRI (*13*) and resting-state fMRI (*14*, *15*) suggest
that a fronto-parietal network supports the initiation of top-down control and control adjustment in
response to task-goals and feedbacks.

Common neural correlates between emotion regulation or memory control with response inhibition tasks
(e.g. stop-signal task) have been investigated in two recent meta-analyses (*11*, *16*). However, no previous
formal meta-analysis has investigated the neural commonalities between emotion regulation and memory
control to pinpoint the overlap in inhibitory control. Using Activation likelihood estimation (ALE), we
aim to demonstrate that emotion regulation and memory control evoke activation of similar brain regions
with a spatial pattern that is similar to activation of typical response inhibition paradigms including stop-
signal and go/no-go tasks. Moreover, beyond overlapping regional brain activity, we also are interested in
overlapping co-activation patterns of associated brain regions. We used meta-analytic connectivity
modeling (MACM) to test whether these brain regions acts as an interconnected functional network. We
expected to find a similar set of co-activated brain regions that are associated with both emotion regulation
and memory control, defining a core "inhibition-related" network.

Identifying an underlying neural network is relevant for potentially brain stimulation interventions
including but not limited to Transcranial Magnetic Stimulation (TMS) and Transcranial Direct Current
Stimulation (tDCS). However, further dissection of molecular underpinnings is necessary to expand our
understanding of inhibition control, which in turn may pave the way for pharmacological interventions.
Therefore, we reasoned whether emotion regulation and memory control are not only related to a common
neural network defined by activity and connectivity, but also by associated similarities in spatial
transcriptional profiles. Conventional imaging-genetic methods including candidate gene methods and
genome-wide association approaches cannot reveal the relationship between localized gene transcription
and task-related brain activity, even though they already showed the association between multiple
common gene variants and neuroimaging measures (*17*, *18*). Thus, to explore the relationship between
spatial transcriptional profiles and emotion regulation or memory control-related brain activity, we
adopted a recently developed approach to associate spatial maps of gene expression in post-mortem brain

tissues with brain activation measures (*19*, *20*). Spatial pattern analysis of gene expression maps of the
Allen Human Brain Atlas (AHBA) together with neuroimaging revealed fundamental features of
transcriptional regulation (See review by Fornito, Arnatkevičiūtė, & Fulcher, 2018) and related
disruptions in brain disorders (*22–25*). However, the correspondence between spatial transcriptional
profiles and neural functionality has not yet provided full insight into how genetic correlates are linked to
core cognitive abilities (e.g. inhibitory control in this study). Based on the assumption that spatial
transcriptional profiles not only co-vary with the connectional architecture but also support the task-
evoked, synchronous brain activity (*19*, *20*, *26*, *27*) we expected to find similar spatial transcriptional
profiles between emotion regulation, memory control, and response inhibition.

Results
101
Summary of the analytical pipeline

To investigate the neural and transcriptional commonality of emotion regulation and memory control, we
combined task fMRI data, neuroimaging meta-analytic approaches, and postmortem gene-expression data.
First, we used the ALE method to generate the task activation map for each paradigm of interest (e.g.
emotion regulation or memory control) based on 95 published task fMRI studies with in total 1857 healthy
participants (Figure 1A). Then, we used the brain regions identified by the ALE as seed regions and
estimated the meta-analytic connectivity map (or co-activation map), using information from BrainMap
(Figure 1B). And to explore the behavioral relevance of these findings, we examined the associations of
these co-activation patterns in a data-driven way via Neurosynth with behavioral domains (Figure 1C).
Next, we calculated the spatial association between task activation maps and human gene expression maps
to identify their common spatial transcriptional profiles (Figure 1D). Finally, we performed a systematic
and integrative analysis of the resulting gene list to gain an in-depth understanding of putative biological
functions and disease associations of the identified "inhibition-related "gene set (Figure 1E).

Regional brain activity associated with emotion regulation and memory control

Taking data from 95 published studies including in total 1857 subjects, we used 15 Emotion Regulation
studies to represent emotion regulation, 15 Think/No-think studies to represent memory control, and 27
Go/No-go studies as well as 38 Stop Signal studies to represent response inhibition. (A list of studies and
coordinates are available via the Open Science Framework; Search and inclusion criteria in *Materials and*
Methods).

Regional brain activity of emotion regulation

The meta-analysis of the emotion regulation studies revealed six brain regions that are active during
"regulation" compared to a "passive viewing" condition (FWE-cluster level corrected p<.05, uncorrected
p<.001, threshold permutations=1000). Emotion regulation task consistently led to activation in right
insular/ inferior frontal gyrus (IFG), left IFG, left insular, middle cingulate gyrus, right inferior parietal
lobule (IPL) and left supplementary motor area (SMA) (Table 1, Figure 2A).

Regional brain activity of memory control

Memory control studies revealed five brain regions during the "No-Think" condition compared to the
"Think" condition (FWE-cluster level corrected p<.05, uncorrected p<.001, threshold permutations=1000):
left insular, right dorsolateral prefrontal cortex (DLPFC), right IPL, left precuneus, and SMA (Table 1,
Figure 2B).

Regional brain activity of response inhibition

We used the same meta-analytical approach for all Go/No-go and Stop-signal studies and found similar
significant clusters during the "control" condition compared to "baseline" condition (No-go vs go; Stop vs
Go) in insular, IFG, middle cingulate gyrus, SMA, IPL, and DLPFC (Figure 2C and Figure 2D, Table S1
and Table S2).

Analyses of convergence and divergence

To examine the spatial convergence and divergence between emotion regulation and memory control
activations, we performed a formal conjunction and contrast analysis (FWE-cluster level corrected p<.05,
uncorrected p<.001, threshold permutations=1000). The conjunction analysis revealed two shared clusters
within the IPL (cluster 1 peak MNI:60/-42/42; cluster 2 peak MNI:58/-46/40). We did not find any
reliable clusters in the contrast analysis.

Co-activation maps of regions associated with emotion regulation and memory control

To gain deeper insight into the co-activation profiles of brain regions associated with emotion regulation
and memory control, we used a large database of fMRI studies (*BrainMap*: http://www.brainmap.org/)
and applied meta-analytic connectivity modelling (MACM). This approach reveals brain regions that are
consistently activated together with the seed regions resulting from the ALE meta-analyses. To control for
specific modelling methods and sizes of seed regions, we also performed a similar co-activation analyses
using *Neurosynth* (http://neurosynth.org/) based on peak voxel coordinates of each seed region instead of
the entire clusters. The two approaches yielded similar co-activation maps for each ROI. In the main test,
we only report results from the *BrainMap* analysis (*Results from the Neurosynth can be found in*
Supplemental Materials, Figure S1 and S2.)

In total, we analyzed six ROIs from the meta-analysis of emotion regulation studies. MACM analyses for
the left IFG and right IPL revealed co-activation patterns in the direct vicinity of the seed regions. Co-
activation analyses of other seed regions including bilateral insula, SMA, MACC revealed co-activation
between these ROIs and with cerebellum, IFG, IPL, thalamus, and DLPFC (Figure 3A). Similarly, we
analyzed five seed regions from the meta-analysis of memory control studies. ROIs including insula,
MACC, SMA, DLPFC, right IPL tended to be co-activated. Only the MACM analysis for the precuneus
ROI identified co-activation with additional regions (left inferior occipital cortex, right middle occipital
cortex, left precentral sulcus) (Figure 3B). In addition, we investigated the co-activation profiles of
response inhibition tasks. The co-activation patterns of 11 ROIs from Go/No-go meta-analysis and ten
ROIs from Stop-signal meta-analysis were estimated using the same method. We found a brain network
including bilateral DLPFC, insular, IPL, thalamus, SMA, and MACC across the co-activation patterns of
ROIs from response inhibition (*See Supplemental Materials*).

Behavioral profiles of the co-activation maps

To identify the cognition domains most strongly associated with these co-activation maps, we created the
behavioral profiles of each co-activation maps using the *Neurosynth* database.

First, we quantified the behavioral profiles of co-activation maps of emotion regulation and memory
control. As expected, these two sets of co-activation maps were both strongly characterized by terms such
as "inhibition", "cognitive control", and "working memory". Then, the behavioral profiles of Stop-signal
and Go/no-go co-activation maps were estimated in the same way. Finally, we investigated the
commonalities of behavioral characteristics of all co-activation maps across four paradigms and revealed

that these co-activation maps have the highest correlations to terms including "inhibition", "cognitive
control", and "working memory", compared to other items (Figure 4).

Transcriptional signatures underlying the Emotion Regulation and Memory control

Next, to understand the transcriptional correlates that may be associated with brain activity elicited by
each task, we combined the Allen Human Brain Atlas (AHBA), a brain-wide atlas of gene expression of
post mortem brains and spatial pattern correlation. More specifically, we aimed to identify the genes for
which the spatial transcriptional patterns are similar to the spatial pattern of brain activity during a given
task.

Transcriptional correlates of brain activity

Emotion regulation-related brain activity patterns were correlated with spatial transcriptional maps of
1529 genes across cortical regions ($p<0.05$, FDR corrected; 7% of the total). The activation-gene
expression association analysis showed the correlation between memory control-related brain activity
patterns and the expression pattern of 2531 genes ($p<0.05$, FDR corrected; 12% of the total). The top ten
genes with the highest spatial similarities are represented in Figure 5A and 5B. (*Complete gene lists in*
Open Science Framework(OSF) repository; Details of OSF in Methods and Supplemental Materials)

Common transcriptional profiles and "inhibition-related" genes

We found a substantial overlap (Figure 5A) between the two identified gene lists for the emotion
regulation and memory control activation networks: there are in total 1130 genes (55% of the memory
control associated genes; 73% of the emotion regulation associated genes) whose expression pattern
correlated with the brain activity pattern of both tasks. We further evaluated the significance of the overlap
by generating 5000 null distributions of the overlapping genes. Specifically, two gene lists with identical
sizes (list1=2531; list2=1529) were randomly selected from all the genes (N=20787), and then
overlapping genes between the two random lists were identified. This procedure was repeated 5000 times,
and we found that the amount of the overlapping genes between the memory control and the emotion
regulation was significantly larger than the amount of randomly sampled overlapping genes lists
($p<0.001$). Furthermore, we defined genes of which the expression patterns are correlated with the
activation maps of the four tasks at issue (memory control, emotion regulation, stop-signal, and go/no-go)
as "inhibition-related" genes, which are 1061 genes (*See OSF repository*), of which 93.89% were also
found in genes overlapping with memory control and emotion regulation activation patterns.

Specificity of "inhibition-related" genes

To test the specificity of the "inhibition-related" genes versus genes related to neuronal activity and brain
function in general, we first generated, also using an ALE meta-analysis, the activation map of "Default
Mode Network" (DMN), or "Task-Negative Network" (See Materials and Method; Figure S3). Next, we
identified a list of genes associated with the DMN network via the AHBA and the same activation-gene
expression association analysis. Given the fact that the DMN has been associated with homeostasis and
undirected thought or mind-wandering, we expected that DMN-associated genes are different from
inhibition-association genes. To test this, we calculated the number of overlapping genes between all
possible combinations of two out of the five reported gene lists (memory control, emotion regulation,
stop-signal, and go/no-go, and DMN) across different thresholds. As depicted in Figure 5C, genes
associated with inhibitory-control tasks are more similar to each other than the associated genes between
the inhibitory-control tasks and the DMN (mean Within Inhibitory =41.75%, mean Inhibitory & DMN
=35.60%, t=7.21, $p=7.81 \times 10^{-13}$).

Biological functions and disease associations of the "inhibition-related" genes

Using gene ontology (GO) (*28*), a widely-used literature-based gene-to-function annotation analysis, we generated a list of biological functions related to the "inhibition-related" genes. Furthermore, due to common finding of impairment in inhibitory control in neurological and psychiatric disorders, we explored the human disease association of the identified gene list.

Enrichment for biological functions

Gene Ontology Enrichment Analysis (GOEA) of "inhibition-related" genes identified 19 related biological functions (Table S3). We further visualized all significant items separately (Figure S4-S6) according to their categories (Biological Processes (BP), Molecular Function (MF), and Cellular Component (CC)) to reveal the hierarchical and network structure of significant items and related, but not significant GO items. Considering the frequently seen words across BP, MF and CC items, we grouped the 19 significant biological functions into two clusters: "synaptic/synapse" and "neuropeptide". More specifically, within the cluster "synaptic/synapse", 30 genes within the list are associated with chemical synaptic transmission, ten genes with regulation of synaptic plasticity, and 39 genes with the glutamatergic synapse. For the cluster of "neuropeptide", 17 genes within the list are associated with neuropeptide signaling pathways and eight genes with neuropeptide hormone activity.

Enrichment for human diseases

Additionally, we used *ToppGene* (https://toppgene.cchmc.org/) to perform the gene list enrichment analysis of "inhibition-related" genes for human diseases. Using *DisGeNET* (www.disgenet.org), a comprehensive database on the relationship between human diseases and genes, we found significant associations between our gene list and 13 disease terms (Figure 5D, Table S4). After clustering disease terms describing the same disease, we found that the "inhibition-related" genes are associated with bipolar disorder, schizophrenia, alcoholic intoxication, depression, autism spectrum disorders, neuroblastoma, and Parkinson disease.

Discussion

Inhibitory control is a fundamental cognitive function supporting other processes like emotion regulation and response inhibition. Here, we provide neurobiological evidence from human neuroimaging and transcriptional mapping to support the concept that there is one generic neural network of inhibitory control with a set of "inhibition-related" genes modulating not only response inhibition but also emotion regulation and memory retrieval. Our meta-analysis of 95 neuroimaging studies revealed a common role of the right inferior parietal lobule and related regions in a frontal-parietal-insular network during emotion regulation and memory control. These co-activation patterns were also similar to the meta-analysis results of response inhibition tasks, and "inhibition-related" network reported in the literature. Additionally, we used the Allen Human Brain Atlas as an avenue to link this neural network to common transcriptional profiles and identified "inhibition-related" genes, which are associated with glutamatergic synapse function, neuropeptide-related activity and the risk for several psychiatric disorders and Parkinson's disease.

The idea that inhibitory control is the underlying core cognitive function in emotion regulation was already suggested before (*1*, *8*, *29–31*). Similarly, it was also suggested that inhibitory control plays a fundamental role in memory control (*32*). A unified theory proposed a central role of inhibitory control in various psychological domains (e.g. motor inhibition, emotional response, and memory retrieval)

depending on external task requirements (*7*, *33*, *34*). Engen and Anderson recently reviewed behavioral
and neuroimaging studies in this field and proposed the conceptual link between emotion regulation and
memory control (*6*). However, there has been little empirical support for this link. Our multimodal
analysis provides rich evidence beyond neuroimaging supporting a conceptual link, and suggests that
inhibitory control as well as its underlying neural and transcriptional correlates modulates both emotion
regulation and memory control.

Brain activation patterns of emotion regulation and memory control found here are in accordance with
previous meta-analyses (emotion regulation: e.g. Buhle et al., 2014; Kohn et al., 2014; memory control:
e.g. Guo et al., 2018). We find one overlapping region between emotion regulation and memory control,
the right inferior parietal lobule. Although, considering the central role of inhibitory control, only one
overlapping brain region seems surprising at first glance. However, meta-analytic connectivity modelling
revealed that other regions (including IFG, insula, preSMA/MACC, IPL) that lacked significantly
overlapping activations across emotion regulation and memory control, form a tightly integrated network.
Our results suggest that although these regions do not overlap strictly, they belong to the same functional
network. Our behavioral profile analysis corroborated this interpretation: both co-activation maps of
emotion regulation and memory control, as well as stop-signal and go/no-go paradigms, have comparable
behavioral profiles and were characterized by terms like "inhibition", "cognitive control", and "working
memory".

We have demonstrated neural commonality of emotion regulation and memory control. Next, we
proceeded to investigate if transcriptional profiles overlap with activity patterns in a similar way.
Critically, this study adopted an imaging-genetic approach to investigate the common transcriptional
signatures across neural networks of emotion regulation and memory control. Activation-gene expression
association analysis revealed a largely overlapping gene list whose expression patterns were similar to the
activation patterns. Furthermore, we identified a list of "inhibition-related" genes and characterized their
biological functions and diseases associations. "Inhibition-related" genes were primarily associated with
synaptic transmission and plasticity, in particular at the glutamatergic synapse. Glutamate is an excitatory
neurotransmitter, and glutamatergic abnormalities have long served as a major hypothesis for
understanding schizophrenia, depression, and Parkinson's disease, and it's associated emotional and
cognitive deficits (*35–39*). "Neuropeptide" is another frequently reported key word, which is known to be
involved in a wide range of behaviors and brain function (e.g. role of oxytocin in social cognition) (*40*).
Even though, our current method lacks specificity to link a given neural network to any specific type of
neuropeptide, both glutamate and neuropeptides are involved in neural information transfer, which may
relate to molecular mechanisms underlying synchronous brain activity during inhibitory control.
Inhibition-related" genes were also associated with risks for several psychiatric disorders (e.g. depression,
bipolar disorder, schizophrenia, and autism) and Parkinson's disease, a disorder within impaired inhibition
(*41–43*).

Recently, neuroimaging (*44*, *45*) and genetic studies (*46–48*) collectively demonstrated the potential
common biological roots across psychiatric disorders. However, common phenotypes across disorders are
less well understood. Thus, the National Institute of Mental Health's Research Domain Criteria (NIMH's
RDoC) (https://www.nimh.nih.gov/research/research-funded-by-nimh/rdoc/index.shtml) summarized
several domains of phenotypes, where inhibitory control is a central aspect. Here, our results highlighted
the critical role of inhibitory control and its biological underpinning across psychiatric disorders. Firstly,
dysfunctional inhibitory control (e.g. impaired response inhibition, lack of emotion regulation, and
compromised memory control) is evident across different psychiatric disorders (*49*, *50*, *59*, *60*, *51–58*).
Also, inhibitory control deficits can be further linked to some disorder-specific symptoms (e.g. lack of

inhibition of negative thoughts (or rumination) in depression, lack of fear control in anxiety, and failure to
avoid retrieval of traumatic memories in PTSD). Secondly, our results suggest an overlap between brain
regions (frontal-parietal-insular network) involved in inhibitory control and regions whose structural
abnormalities were observed consistently in a variety of psychiatric diagnoses (*44*). Thirdly, "inhibition-
related" genes which we identified by spatial transcriptional profiles that overlap with activation patterns
of inhibitory control tasks, were also associated with the risks for a variety of psychiatric disorders.

Our study has two limitations that should be mentioned. First, our neural commonality analyses were
based on fMRI studies only. However, overlap in fMRI activation or co-activation patterns lacks temporal
information of the underlying cognitive processes. Electroencephalography (EEG) or
Magnetoencephalography (MEG) in humans could provide further confirmatory evidence for the idea of
common cognitive process. Recently, Castiglione and colleagues reported that memory control elicited an
electrophysiological signature, increased right frontal beta, which was seen in the Stop Signal task (*61*).
Follow-up electrophysiological studies or even a meta-analysis of them might confirm the idea of a
common electrophysiological signature. Second, the current activation-gene expression association
analysis is still preliminary (e.g. low sample size and spatial resolution of the post-mortem data), and
without the possibility of causal interference. Nevertheless, the method already showed great potential
when helping to understand basic molecular principles of both the structural and functional connectome
(*See review by Fornito et al., 2018*) and it identified molecular mechanisms underlying changes in brain
structure or function associated with brain disorders (e.g. autism spectrum disorder (*22*), Huntington's
disease (*23*), schizophrenia (*25*), and Alzheimer's disease (*24*)). Results from these studies are consistent
with the genetics of neuropsychiatric disorders using conventional methods like genome-wide association
studies or animal models. Taken together, although preliminary, neuroimaging-gene expression
association analysis has demonstrated its potential to bridge brain structure-function associations and to
reveal its underlying molecular processes. To detect more specific associations between spatial
transcriptional profiles and neuroimaging data, large sets of postmortem gene-expression data with higher
spatial resolution need to be collected, and more dedicated analytical methods need to be developed and
validated (*62*). Also, new methods may better control for the effect of domain-general genes that are
supporting brain function in general.

In summary, our multimodal analysis identified a frontal-parietal-insular neural network and a set of genes
associated with inhibitory control across emotion regulation, memory control and response inhibition. The
integrative approach established here bridges between cognitive, neural, and molecular correlates of
inhibitory control and can be used to study other higher-level cognitive process. Our findings may
deepening our understanding of emotion regulation and memory control in health, and pave the way for
better emotion regulation and memory control by targeting the core inhibitory-related network or related
molecular targets in patients with such deficit at issue.

Materials and Methods

Literature searches, selection and coordinates extraction

In total, we perform literature searches for four task paradigms (think/no-think, emotion regulation, stop-signal, and go/no-go) and one network ("default mode network"). To avoid biases, we use the same inclusion criteria during the search.

(1) We only included data from studies on healthy adults with no prior report of neurological, medical, or psychiatric disorders in the current meta-analysis, while results of patients or specific sub-group effects (e.g., gender differences) were not included. Articles including patients were only selected if they reported results for a control group separately, and only the control group was included here.

(2) Only neuroimaging studies, which used whole-brain fMRI and reported coordinates for brain activation or deactivation in standard anatomical reference space (Talairach/Tournoux; Montreal Neurological Institute (MNI)) were considered. Coordinates originally published in Talairach space were converted to MNI space using the algorithm implemented in GingerALE 2.3.5 (http://www.brainmap.org/ale/)

(3) Only studies reporting results of General Linear Model (GLM) contrasts were included, while studies focusing on functional connectivity, structural, resting-state or brain–behavior correlations were excluded.

(4) Only studies reporting whole brain analyses were included, while studies based on partial coverage or employing only region-of-interest analyses were excluded.

Detailed search and extraction procedures were as following for each paradigm:

Think/No-Think studies

A step-wise procedure was used to search articles, published before February 2018, using functional MRI to investigate brain activity during Think/No-Think paradigm. First, we used standard search in PubMed and ISI Web of Science to perform the search. More specifically, we used the combination of following key words during the search: "memory regulation", "memory control", "memory suppression regulation", "memory inhibition", "think/no-think", "fMRI", "neuroimaging", "functional magnetic resonance imaging", or "functional MRI". At the same time, we carefully exclude studies using the "directed forgetting" paradigm, which targets at the memory control during the encoding. Next, two lab members compared the search results with a recent review article (*3*) to find additional relevant studies. The same two lab members independently extracted the coordinates and other essential information (e.g. sample size, type of stimulus) extraction based on the identified think/no-think literatures and then cross validated the coordinates. In summary, this search and inclusion/ exclusion criteria led to 15 think/no-think studies (353 subjects and 249 foci).

Emotional Regulation Studies

We take the databases of previously published meta-analyses on emotion regulation (*9*, *63*). We use the keywords:"emotion regulation", "affective regulation", "implicit emotion regulation", "explicit emotion regulation", "interpersonal emotion regulation", "extrinsic emotion regulation, "intrinsic emotion regulation", "reappraisal", "suppression", "distraction", "detachment", "labelling", "affective labelling", "reinterpretation", "rumination", "fMRI", "neuroimaging", "functional magnetic resonance imaging", or "functional MRI". In the case that a study did not report the contrast of interest for this meta-analysis, the corresponding authors were contacted and asked to provide more information on their data. In the following the term "experiment" refers to any single contrast analysis, while the term "study" refers to a

scientific publication, usually reporting several contrasts, i.e. experiments. This search and the employed
inclusion/exclusion criteria led to a total inclusion of 107 studies from peer-reviewed journals by July 31st,
2017 (385 experiments, 3204 participants).

Each experiment was manually coded by the authors of previous meta-analysis (C.M. and N.K) with terms
that described the experimental design with respect to contrast, stimulus type utilized, emotion regulation
strategy, goal of the strategy, valence of the stimuli, tactics of the strategy and the task nature. To achieve
a more appropriate comparison between think/no-think and emotion regulation, we restrict inclusion to ER
studies that used the "suppression" or "distraction" strategy. We excluded all the studies used "cognitive
reappraisal" strategy due to high-level cognitive processes instead of inhibitory control involved and the
potential imbalance in study sample size. These criteria led to inclusion of 15 emotion regulation studies
(387 subjects and 165 foci).

Go/No-go and Stop-signal studies

A similar procedure was used to search published whole-brain functional MRI studies using the "Go/No-
go" and "stop-signal" paradigm. To confirm the completeness of our search, we compared our results with
used studies in a recent meta-analysis of motor inhibitory and memory control (*11*). Again, two lab
members (W.L and N.P) performed the coordinates and study information extraction for the Go/No-go
and Stop-signal studies.

Default Mode Network (DMN) (task-negative network) identification

We also performed a coordinated-based meta-analysis to identify the DMN. Instead of manually searching
related studies and extracting coordinates, we used the BrainMap database (*64*, *65*) to find peak
coordinates of task-independent deactivation reported in neuroimaging literatures. This method was used
before by Laird and colleagues (*66*) to identify the core regions in DMN. More specifically, we searched
the BrainMap for all contrasts that were labelled as "deactivation" and "low-level control" during
submission. "Deactivation" refers to contrasts in which stronger signal was observed during a baseline
condition than during task condition (e.g. Control-Task); "low-level control" are conditions in which
either fixation or resting was defined as baseline. Our search was further limited to "normal mapping",
which means that participants who diagnosed with disease or disorders were excluded. In total, 105
studies (1588 foci) matched our search criteria, and were used in the following analyses.

Activation Likelihood Estimation (ALE) analyses

The ALE analyses were based on the revised ALE algorithm (*67*) in GingerALE 2.3. Firstly, two separate
meta-analyses were conducted for the Think/No-Think (contrast: No-Think vs. Think) and emotion
regulation (contrast: Regulation vs. View) tasks using cluster-level inference (FWE cluster-level
correction p<0.05, uncorrected cluster-forming threshold p<0.001, threshold permutations=1000).
Secondly, contrast analyses (*68*) were conducted between the Think/No-Think and emotion regulation
tasks. In these contrast analyses, the thresholded activation maps from the two separate analyses, as well
as the pooled results from both tasks were used as inputs. Conjunction and contrast maps between the
conditions were given as output. For the output images, the same cluster-level threshold correction was
used (FWE cluster-level correction p<0.05, uncorrected cluster-forming threshold p<0.001, threshold
permutations=1000).

Additionally, meta-analyses of published fMRI studies using Stop-signal and Go/No-go paradigm were
also performed. The same software (GingerALE 2.3) and threshold (cluster-level p<0.05) was used to
perform the analysis. Due to the unbalanced number of studies and subjects included (*emotion*

regulation=387; think/no-think=353; stop-signal=672; go/no-go=445), no conjunction or contrast
analyses were performed for four tasks to identify the overlap.

Co-activation analyses using BrainMap

We conducted the Meta-Analytic Connectivity Modeling (MACM) analyses on the regions from the ALE
meta-analysis. More specifically, for each ROI, we used the BrainMap database (*69, 70*) to search for
experiments that also activated the particular ROI. Next, we retrieved all foci reported in the identified
experiments. Finally, ALE analyses were performed over these foci to identify regions of significant
convergence. Sequentially, raw co-activation maps were corrected for multiple comparisons (Voxel-wise
False Discovery Rate(FDR)<0.05. All clusters size> 200mm^3)

Functional profiles of the co-activation maps

To assess associated functional terms of the co-activation maps generated by MACM, we used the
NeuroSynth meta-analytic database (www.neurosynth.org) (*71*). We followed the methodology (See
Meta-analytic Functional Gradients section) used in a previous study to assess topic terms associated with
the principal connectivity gradient in the human brain (*72*). More specifically, we conducted a term-based
meta-analysis for the same list of NeuroSynth topic terms as Margulies and colleagues did. This list
covered well-studied functional terms from low level cognition (e.g., visual perception, and auditory) to
high level cognition (e.g., language, and rewarding). Sequentially, we examined the association between
these term-based activation maps with our four sets of co-activation maps (emotion regulation, think/no-
think, stop-signal, go/no-go). For each co-activation map, a spatial similarity index (r statistic) between
each co-activation map and each meta map of functional term was provided. The terms were then ordered
based on the average correlation for the interpretation and visualization.

Activation-gene expression association analysis

We used a recently developed activation-gene expression association analysis to link the task-related brain
activity to gene expression in postmortem human brains. This analysis can identify a list of associated
genes based on the MRI-space statistical map(s). This analysis presupposes that if certain gene(s) are
associated with the cognitive task of interest, then spatial distributions of their expression values and task-
related activation pattern measured by functional MRI should be similar.

The Allen Human Brain Atlas (http://www.brainmap.org) was used in the gene expression decoding
analysis. The atlas provided genome-wide microarray-based gene expression data based on six
postmortem human brains (Gene expression level of over 62000 gene probes for around 1000 sampling
sites across the whole brain were recorded) (*73*). Additionally, structural brain imaging data of each donor
was collected and provided, which enable users to visualize gene expression in its naive space and
perform the registration to the standard MRI MNI space.

Previous studies used slightly different statistical methods to associate task-independent MRI-based brain
measures (e.g. cortical thickness, functional or structural networks,) with the gene expression data (*74–76*).
We used the method developed by Gorgolewski and colleagues implemented in the alleninfo tool (*19*)
(https://github.com/chrisfilo/alleninf). This method was, originally designed for the association analysis
between voxel-vise statistical maps and gene expression maps. This method was also by default
implemented in Neurovault (*77*) (https://neurovault.org/). The method has two important features: (1)
nonlinear coregistration of the donor brain with MNI space was allowed and (2) the ability to use random
effects model make it possible to generalize the results to the whole population. The activation-gene
expression association analysis works as following: (1) data from gene probes was aggregated for each
gene, resulting in 20787 gene expression maps. (2) For each gene expression map, MNI coordinates of

each sampling location (the locations which brain tissues were analyzed for the gene expression data)
were extracted to draw a spherical ROI (r=4mm). We used these ROIs to extract the average values of the
ALE statistical map within each ROI. Next the gene expression and meta-analysis vectors were correlated.
(3) This extraction and correlation procedure was repeated for each gene expression map to quantify the
spatial pattern similarity between the statistical maps and gene expression map. (4) Multiple correction
was implemented using the threshold $P_{FDR}<0.05$ to generate the significant associated gene list(s) among
the 20787 genes. Since negative correlation between brain measures and gene expression is difficult to
explain, we only considered the positively correlated genes. Additionally, because of fundamental
differences in gene expression between cortical and subcortical regions, we only performed our analyses
within the cortical regions.

The described association analysis takes MRI statistical map(s) as input(s) and will output a list of
significantly associated genes. To investigate the common transcriptional signatures (associated genes) of
emotion regulation and memory control, we used the unthresholded statistical maps from the ALE
analyses to the identify the task-associated gene list via the association algorithm. Next, the common gene
list was generated by overlapping the emotion regulation-associated gene list and memory control-
associated gene list. To further investigate whether the two gene lists significantly overlapped with each
other, we generated the null distribution of the number of overlapped genes by randomly creating two
gene lists (with the same size as the real gene lists) from the 20787 genes (5000 times) and calculated the
number of overlapped genes. Then the p-value of the overlap can be estimated by comparing the number
of the overlapped genes to the null distribution. To identify the "inhibition-related" genes, we first used
the same activation-gene expression association analysis to identify the stop-signal and go/no-go-
associated gene list and then defined the overlap between four gene lists (emotion regulation, think/no-
think, stop-signal, and go/no-go) as "inhibitory-related" genes.

We aim to improve the specificity of our activation-gene expression association analysis and safeguard the
possibility that these identified genes may only support the general functional brain network architecture
instead of particular cognitive functions. To rule out this possibility, we performed a dedicated control
analysis. An unthreholded DMN ALE map was used as a comparison with four inhibitory control tasks
because control processes hardly take place in the DMN and the DMN has been associated with
homeostasis and undirected thought or mind-wandering. Rather than arbitrarily using the q<0.05 (p_{FDR}
<0.05) as the threshold, we used a range of different numbers of genes (x from 1 to 2000, with step size 10)
as thresholds to identify the top x most similar genes. We calculated two kinds of overlap under different
thresholds: the first one is "within control" overlap, which is the overlap in gene expression association
between two inhibitory tasks (e.g. think/no-think and emotion regulation, or stop-signal and go/no-go); the
second one is "inhibitory & DMN" overlap, which is the transcriptional overlap between one of the
control-related tasks and DMN (e.g. think/no-think and DMN, or emotion regulation and DMN). Finally,
we average the percentage of overlap within each kind of overlap for certain threshold x, and compare the
average percentages across all the thresholds.

Gene Ontology Enrichment Analysis

The Gene Ontology(GO) is a widely used bioinformatics tool to interpret the complex gene list based on
the knowledge regarding functions of genes and gene products (*28*, *78*). To systematically investigate the
biological meaning of the "inhibitory- related" genes, we use GO to perform the overrepresentation test
for Biological Processes (BP), Molecular Function (MF), and Cellular Component (CC). Currently,
experimental findings from over 140 000 published papers are represented as over 600 000
experimentally-supported GO annotations in the GO knowledgebase. For the input gene list, "Gene
Ontology enrichment analysis" can identify relevant groups of genes that function together, and associate

and statistically test the relationship between these groups and Go annotations. In this way, it can reduce a
big list of genes to a much smaller number of biological functions (BP, MF, or CC), and make the input
list more easily comprehendible. This method has been applied successfully to understand the output of
many gene expression studies including the study that also used the Allen Human Brain Atlas (*75*). More
specifically, we used GOATOOLS (*79*) to perform the GO analyses based on Ontologies and Associations
downloaded on 15th November 2018. All of the significant GO items were corrected by FDR correction
(p<0.05). We further identified the frequently seen words within all the significant GO items by counting
the frequency of the words after removal of not meaningful words (e.g. "of" and "the").

Disease-related gene set enrichment

Although GO analysis can provide insights into the biological functions (BP, MF or CC) of the overlapped
gene list, the approach does not provide sufficient information to identify disease associations of the gene
list. We leveraged *ToppGene* (*80*) to explore the gene-diseases associations. The ToppGene platform can
cluster groups of genes together according to their disease associations, and perform a statistical test as
well as the multiple testing error corrections. The latest version of DisGeNET included associations
between 17,549 genes and 24,166 diseases, disorders, or abnormal human phenotypes. In this study, our
analysis was based on one of the sub-database of DisGeNET, the DisGeNET Curated (expert-curated
associations obtained from UniProt and CTD human datasets). All of the significant diseases items were
corrected by FDR correction (p<0.05).

Data and code availability.

All the data (excluding neuroimaging data) are stored in Open Science Framework (OSF). Open access
data includes study summary, extracted coordinates for ALE, significant associated genes for each task
paradigm, and overlapped gene list(s). (OSF link: https://osf.io/6wz2j/). Neurovault was used to store all
the neuroimaging data (e.g. results of ALE from Figure2, MACM from Figure3, and "Top Genes"
expression maps from Figure 4) and provide 3D visualization of all the statistical maps. (Neurovault link:
https://neurovault.org/collections/4845/). Other research data supporting reported findings are available
from the authors upon request.

For the neuroimaging meta-analysis, two software (GingerALE and Sleuth)
(http://www.brainmap.org/software.html) were used. The Neurosynth Tool
(https://github.com/neurosynth/neurosynth), a Python package which covered most of the functions on
Neurosynth (http://www.neurosynth.org/) website, was used in functional profile analyses and co-
activation analyses (*see Supplementary Materials*). Activation-gene expression association analyses were
based on alleninfo tool (https://github.com/chrisfilo/alleninf) and web application of gene decoder within
the Neurovault (https://neurovault.org/). Nilearn (https://nilearn.github.io/) was used to load, manipulate,
and visualize MRI statistical maps.

GOATOOLS (https://github.com/tanghaibao/goatools), a Python library, was used for Gene Ontology
analyses and related visualization. The Ontology data was downloaded from Gene Ontology website
(http://geneontology.org/ontology/) and the Association data was downloaded from National Center for
Biotechnology Information (ftp://ftp.ncbi.nlm.nih.gov/gene/DATA/). ToppGene Suite
(https://toppgene.cchmc.org/) was used for disease-related gene set enrichment. The gene-disease
associations database can be downloaded from http://www.disgenet.org/.

Anaconda (https://www.anaconda.com/) Python 3.6 version for Win10 was used as the platform for all the
programming and statistical analyses. Custom python scripts were written to perform the all analyses

described based on the mentioned Python packages; all code are available from the authors upon request
and will be released via the OSF as soon as possible.

References

1. K. N. Ochsner, J. J. Gross, The cognitive control of emotion. *Trends Cogn. Sci.* **9**, 242–249 (2005).

2. J. J. Gross, R. A. Thompson, Emotion regulation: Conceptual foundations. (2007).

3. M. C. Anderson, S. Hanslmayr, Neural mechanism of motivated forgetting. *Trends Cogn. Sci.* **18**, **Issue:**, 1–14 (2014).

4. M. C. Anderson, C. Green, Suppressing unwanted memories by executive control. *Nature*. **410**, 366 (2001).

5. M. C. Anderson, K. N. Ochsner, B. Kuhl, J. Cooper, E. Robertson, S. W. Gabrieli, G. H. Glover, J. D. E. Gabrieli, Neural systems underlying the suppression of unwanted memories. *Science (80-.).* **303**, 232–235 (2004).

6. H. G. Engen, M. C. Anderson, Memory Control: A Fundamental Mechanism of Emotion Regulation. *Trends Cogn. Sci.* **22**, 982–995 (2018).

7. B. E. Depue, J. M. Orr, H. R. Smolker, F. Naaz, M. T. Banich, The Organization of Right Prefrontal Networks Reveals Common Mechanisms of Inhibitory Regulation Across Cognitive, Emotional, and Motor Processes. *Cereb. Cortex*. **26**, 1634–1646 (2016).

8. K. N. Ochsner, S. A. Bunge, J. J. Gross, J. D. E. Gabrieli, Rethinking feelings: an FMRI study of the cognitive regulation of emotion. *J. Cogn. Neurosci.* **14**, 1215–1229 (2002).

9. N. Kohn, S. B. Eickhoff, M. Scheller, A. R. Laird, P. T. Fox, U. Habel, Neural network of cognitive emotion regulation—an ALE meta-analysis and MACM analysis. *Neuroimage*. **87**, 345–355 (2014).

10. J. T. Buhle, J. A. Silvers, T. D. Wager, R. Lopez, C. Onyemekwu, H. Kober, J. Weber, K. N. Ochsner, Cognitive reappraisal of emotion: a meta-analysis of human neuroimaging studies. *Cereb. cortex*. **24**, 2981–2990 (2014).

11. Y. Guo, T. W. Schmitz, M. Mur, C. S. Ferreira, M. C. Anderson, A supramodal role of the basal ganglia in memory and motor inhibition: Meta-analytic evidence. *Neuropsychologia*. **108**, 117–134 (2018).

12. J. Duncan, The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends Cogn. Sci.* **14**, 172–179 (2010).

13. E. Fedorenko, J. Duncan, N. Kanwisher, Broad domain generality in focal regions of frontal and parietal cortex. *Proc. Natl. Acad. Sci.* **110**, 16616–16621 (2013).

14. J. D. Power, A. L. Cohen, S. M. Nelson, G. S. Wig, K. A. Barnes, J. A. Church, A. C. Vogel, T. O. Laumann, F. M. Miezin, B. L. Schlaggar, others, Functional network organization of the human brain. *Neuron*. **72**, 665–678 (2011).

15. N. U. F. Dosenbach, D. A. Fair, F. M. Miezin, A. L. Cohen, K. K. Wenger, R. A. T. Dosenbach, M. D. Fox, A. Z. Snyder, J. L. Vincent, M. E. Raichle, others, Distinct brain networks for adaptive and stable task control in humans. *Proc. Natl. Acad. Sci.* **104**, 11073–11078 (2007).

16. R. Langner, S. Leiberg, F. Hoffstaedter, S. B. Eickhoff, Towards a human self-regulation system: Common and distinct neural signatures of emotional and behavioural control. *Neurosci. Biobehav. Rev.* (2018).

17. L. T. Elliott, K. Sharp, F. Alfaro-Almagro, S. Shi, K. L. Miller, G. Douaud, J. Marchini, S. M. Smith, Genome-wide association studies of brain imaging phenotypes in UK Biobank. *Nature*. **562**, 210 (2018).

18. A. R. Hariri, V. S. Mattay, A. Tessitore, B. Kolachana, F. Fera, D. Goldman, M. F. Egan, D. R. Weinberger, Serotonin transporter genetic variation and the response of the human amygdala. *Science (80-.).* **297**, 400–403 (2002).

19. K. Gorgolewski, A. Fox, L. Chang, A. Schäfer, K. Arélin, I. Burmann, J. Sacher, D. Margulies, Tight fitting genes: Finding relations between statistical maps and gene expression patterns. *Organ. Hum. Brain Mapping. Hamburg, Ger.* (2014).

20. X. Kong, Y. Song, Z. Zhen, J. Liu, Genetic Variation in S100B Modulates Neural Processing of Visual
Scenes in Han Chinese. *Cereb. Cortex*, bhv322 (2016).

21. A. Fornito, A. Arnatkevičiūtė, B. D. Fulcher, Bridging the Gap between Connectome and Transcriptome.
Trends Cogn. Sci. (2018), doi:10.1016/j.tics.2018.10.005.

22. R. Romero-Garcia, V. Warrier, E. T. Bullmore, S. Baron-Cohen, R. A. I. I. Bethlehem, Synaptic and
transcriptionally downregulated genes are associated with cortical thickness differences in autism. *Mol.*
Psychiatry, 1–12 (2018).

23. P. McColgan, S. Gregory, K. K. Seunarine, A. Razi, M. Papoutsi, E. Johnson, A. Durr, R. A. C. Roos, B. R.
Leavitt, P. Holmans, others, Brain regions showing white matter loss in Huntington's disease are enriched
for synaptic and metabolic genes. *Biol. Psychiatry*. **83**, 456–465 (2018).

24. M. J. Grothe, J. Sepulcre, G. Gonzalez-Escamilla, I. Jelistratova, M. Schöll, O. Hansson, S. J. Teipel, A. D.
N. Initiative, Molecular properties underlying regional vulnerability to Alzheimer's disease pathology. *Brain*.
141, 2755–2771 (2018).

25. I. A. C. C. Romme, M. A. de Reus, R. A. Ophoff, R. S. Kahn, M. P. van den Heuvel, Connectome
disconnectivity and cortical gene expression in patients with schizophrenia. *Biol. Psychiatry*. **81**, 495–502
(2017).

26. S. Berto, G.-Z. Wang, J. Germi, B. C. Lega, G. Konopka, Human Genomic Signatures of Brain Oscillations
During Memory Encoding. *Cereb. Cortex*. **28**, 1733–1748 (2017).

27. J. M. Shine, M. Breakspear, P. T. Bell, K. Ehgoetz Martens, R. Shine, O. Koyejo, O. Sporns, R. A. Poldrack,
Human cognition involves the dynamic integration of neural activity and neuromodulatory systems. *Nat.*
Neurosci. (2019), doi:10.1038/s41593-018-0312-0.

28. M. Ashburner, C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis, K. Dolinski, S. S.
Dwight, J. T. Eppig, others, Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25 (2000).

29. T. D. Wager, M. L. Davidson, B. L. Hughes, M. A. Lindquist, K. N. Ochsner, Prefrontal-subcortical
pathways mediating successful emotion regulation. *Neuron*. **59**, 1037–1050 (2008).

30. K. McRae, S. E. Jacobs, R. D. Ray, O. P. John, J. J. Gross, Individual differences in reappraisal ability: Links
to reappraisal frequency, well-being, and cognitive control. *J. Res. Pers.* **46**, 2–7 (2012).

31. B. J. Schmeichel, R. N. Volokhov, H. A. Demaree, Working memory capacity and the self-regulation of
emotional expression and experience. *J. Pers. Soc. Psychol.* **95**, 1526 (2008).

32. B. J. Levy, M. C. Anderson, Inhibitory processes and the control of memory retrieval. *Trends Cogn. Sci.* **6**,
299–305 (2002).

33. A. R. Aron, T. W. Robbins, R. A. Poldrack, Inhibition and the right inferior frontal cortex: one decade on.
Trends Cogn. Sci. **18**, 177–185 (2014).

34. A. R. Aron, T. W. Robbins, R. A. Poldrack, Inhibition and the right inferior frontal cortex. *Trends Cogn. Sci.*
8, 170–177 (2004).

35. M. Laruelle, L. S. Kegeles, A. ABI-DARGHAM, Glutamate, dopamine, and schizophrenia: from
pathophysiology to treatment. *Ann. N. Y. Acad. Sci.* **1003**, 138–158 (2003).

36. M. Amalric, Targeting metabotropic glutamate receptors (mGluRs) in Parkinson's disease. *Curr. Opin.*
Pharmacol. **20**, 29–34 (2015).

37. J. W. Murrough, C. G. Abdallah, S. J. Mathew, Targeting glutamate signalling in depression: progress and
prospects. *Nat. Rev. Drug Discov.* **16**, 472 (2017).

38. A. L. Gray, T. M. Hyde, A. Deep-Soboslay, J. E. Kleinman, M. S. Sodhi, Sex differences in glutamate
receptor gene expression in major depression and suicide. *Mol. Psychiatry*. **20**, 1057 (2015).

672 39. G. Sanacora, G. Treccani, M. Popoli, Towards a glutamate hypothesis of depression: an emerging frontier of
673    neuropsychopharmacology for mood disorders. *Neuropharmacology*. **62**, 63–77 (2012).

674 40. T. W. Robbins, Neurochemistry of cognition. *Oxford Textb. Cogn. Neurol. Dement.*, 91 (2016).

675 41. D. J. Zgaljardic, J. C. Borod, N. S. Foldi, P. Mattis, A review of the cognitive and behavioral sequelae of
676    Parkinson's disease: relationship to frontostriatal circuitry. *Cogn. Behav. Neurol.* **16**, 193–210 (2003).

677 42. H. Bokura, S. Yamaguchi, S. Kobayashi, Event-related potentials for response inhibition in Parkinson's
678    disease. *Neuropsychologia*. **43**, 967–975 (2005).

679 43. S. Gauggel, M. Rieger, T. A. Feghoff, Inhibition of ongoing responses in patients with Parkinson's disease. *J.*
680    *Neurol. Neurosurg. Psychiatry*. **75**, 539–544 (2004).

681 44. M. Goodkind, S. B. Eickhoff, D. J. Oathes, Y. Jiang, A. Chang, L. B. Jones-Hagata, B. N. Ortega, Y. V
682    Zaiko, E. L. Roach, M. S. Korgaonkar, others, Identification of a common neurobiological substrate for
683    mental illness. *JAMA psychiatry*. **72**, 305–315 (2015).

684 45. Z. Sha, T. D. Wager, A. Mechelli, Y. He, Common Dysfunction of Large-Scale Neurocognitive Networks
685    across Psychiatric Disorders. *Biol. Psychiatry* (2018).

686 46. A. J. Schork, H. Won, V. Appadurai, R. Nudel, M. Gandal, O. Delaneau, M. Revsbech Christiansen, D. M.
687    Hougaard, M. Bækved-Hansen, J. Bybjerg-Grauholm, M. Giørtz Pedersen, E. Agerbo, C. Bøcker Pedersen,
688    B. M. Neale, M. J. Daly, N. R. Wray, M. Nordentoft, O. Mors, A. D. Børglum, P. Bo Mortensen, A. Buil, W.
689    K. Thompson, D. H. Geschwind, T. Werge, A genome-wide association study of shared risk across
690    psychiatric disorders implicates gene regulation during fetal neurodevelopment. *Nat. Neurosci.*, 240911
691    (2019).

692 47. Cross-Disorder Group of the Psychiatric Genomics Consortium., Identification of risk loci with shared
693    effects on five major psychiatric disorders: a genome-wide analysis. *Lancet*. **381**, 1371–1379 (2013).

694 48. V. Anttila, B. Bulik-Sullivan, H. K. Finucane, R. K. Walters, J. Bras, L. Duncan, V. Escott-Price, G. J.
695    Falcone, P. Gormley, R. Malik, others, N. Patsopoulos, others, Analysis of shared heritability in common
696    disorders of the brain. *Science (80-.)*. **360**, eaap8757 (2018).

697 49. S. Erk, A. Mikschl, S. Stier, A. Ciaramidaro, V. Gapp, B. Weber, H. Walter, Acute and sustained effects of
698    cognitive emotion regulation in major depression. *J. Neurosci.* **30**, 15726–15734 (2010).

699 50. J. Joormann, I. H. Gotlib, Emotion regulation in depression: relation to cognitive inhibition. *Cogn. Emot.* **24**,
700    281–298 (2010).

701 51. J. C. Magee, R. E. Zinbarg, Suppressing and focusing on a negative memory in social anxiety: Effects on
702    unwanted thoughts and mood. *Behav. Res. Ther.* **45**, 2836–2849 (2007).

703 52. J. Lipszyc, R. Schachar, Inhibitory control and psychopathology: a meta-analysis of studies using the stop
704    signal task. *J. Int. Neuropsychol. Soc.* **16**, 1064–1076 (2010).

705 53. R. B. Price, J. Mohlman, Inhibitory control and symptom severity in late life generalized anxiety disorder.
706    *Behav. Res. Ther.* **45**, 2628–2639 (2007).

707 54. E. Falconer, R. Bryant, K. L. Felmingham, A. H. Kemp, E. Gordon, A. Peduto, G. Olivieri, L. M. Williams,
708    The neural networks of inhibitory control in posttraumatic stress disorder. *J. psychiatry Neurosci. JPN*. **33**,
709    413 (2008).

710 55. M. T. Tull, H. M. Barrett, E. S. McMillan, L. Roemer, A preliminary investigation of the relationship
711    between emotion regulation difficulties and posttraumatic stress symptoms. *Behav. Ther.* **38**, 303–313 (2007).

712 56. A. Amstadter, Emotion regulation and anxiety disorders. *J. Anxiety Disord.* **22**, 211–221 (2008).

713 57. T. Ehring, D. Quack, Emotion regulation difficulties in trauma survivors: The role of trauma type and PTSD
714    symptom severity. *Behav. Ther.* **41**, 587–598 (2010).

715   58.   A. Catarino, C. S. Küpper, A. Werner-Seidler, T. Dalgleish, M. C. Anderson, Failing to forget: Inhibitory-
716   control deficits compromise memory suppression in posttraumatic stress disorder. *Psychol. Sci.* **26**, 604–616
717   (2015).

718   59.   W. Yang, Q. Chen, P. Liu, H. Cheng, Q. Cui, D. Wei, Q. Zhang, J. Qiu, Abnormal brain activation during
719   directed forgetting of negative memory in depressed patients. *J. Affect. Disord.* **190**, 880–888 (2016).

720   60.   M. D. Sacchet, B. J. Levy, J. P. Hamilton, A. Maksimovskiy, P. T. Hertel, J. Joormann, M. C. Anderson, A.
721   D. Wagner, I. H. Gotlib, Cognitive and neural consequences of memory suppression in major depressive
722   disorder. *Cogn. Affect. Behav. Neurosci.* **17**, 77–93 (2017).

723   61.   A. Castiglione, J. Wagner, M. Anderson, A. R. Aron, Preventing a Thought from Coming to Mind Elicits
724   Increased Right Frontal Beta Just as Stopping Action Does. *Cereb. Cortex*, 1–13 (2019).

725   62.   A. Arnatkeviciūtė, B. D. Fulcher, A. Fornito, A practical guide to linking brain-wide gene expression and
726   neuroimaging data. *Neuroimage.* **189**, 353–367 (2019).

727   63.   C. Morawetz, S. Bode, B. Derntl, H. R. Heekeren, The effect of strategies, goals and stimulus material on the
728   neural mechanisms of emotion regulation: A meta-analysis of fMRI studies. *Neurosci. Biobehav. Rev.* **72**,
729   111–128 (2017).

730   64.   A. R. Laird, J. J. Lancaster, P. T. Fox, Brainmap. *Neuroinformatics.* **3**, 65–77 (2005).

731   65.   P. T. Fox, J. L. Lancaster, Mapping context and content: the BrainMap model. *Nat. Rev. Neurosci.* **3**, 319
732   (2002).

733   66.   A. R. Laird, S. B. Eickhoff, K. Li, D. A. Robin, D. C. Glahn, P. T. Fox, Investigating the Functional
734   Heterogeneity of the Default Mode Network Using Coordinate-Based Meta-Analytic Modeling. *J. Neurosci.*
735   **29**, 14496–14505 (2009).

736   67.   S. B. Eickhoff, A. R. Laird, C. Grefkes, L. E. Wang, K. Zilles, P. T. Fox, Coordinate-based activation
737   likelihood estimation meta-analysis of neuroimaging data: A random-effects approach based on empirical
738   estimates of spatial uncertainty. *Hum. Brain Mapp.* **30**, 2907–2926 (2009).

739   68.   S. B. Eickhoff, D. Bzdok, A. R. Laird, C. Roski, S. Caspers, K. Zilles, P. T. Fox, Co-activation patterns
740   distinguish cortical modules, their connectivity and functional differentiation. *Neuroimage.* **57**, 938–949
741   (2011).

742   69.   A. R. Laird, S. B. Eickhoff, P. M. Fox, A. M. Uecker, K. L. Ray, J. J. Saenz, D. R. McKay, D. Bzdok, R. W.
743   Laird, J. L. Robinson, others, The BrainMap strategy for standardization, sharing, and meta-analysis of
744   neuroimaging data. *BMC Res. Notes.* **4**, 349 (2011).

745   70.   A. R. Laird, S. B. Eickhoff, F. Kurth, P. M. Fox, A. M. Uecker, J. A. Turner, J. L. Robinson, J. L. Lancaster,
746   P. T. Fox, ALE meta-analysis workflows via the brainmap database: progress towards a probabilistic
747   functional brain atlas. *Front. Neuroinform.* **3**, 23 (2009).

748   71.   T. Yarkoni, R. A. Poldrack, T. E. Nichols, D. C. Van Essen, T. D. Wager, Large-scale automated synthesis
749   of human functional neuroimaging data. *Nat. Methods.* **8**, 665 (2011).

750   72.   D. S. Margulies, S. S. Ghosh, A. Goulas, M. Falkiewicz, J. M. Huntenburg, G. Langs, G. Bezgin, S. B.
751   Eickhoff, F. X. Castellanos, M. Petrides, E. Jefferies, J. Smallwood, Situating the default-mode network
752   along a principal gradient of macroscale cortical organization. *Proc. Natl. Acad. Sci.* **113**, 12574–12579
753   (2016).

754   73.   M. J. Hawrylycz, E. S. Lein, A. L. Guillozet-Bongaarts, E. H. Shen, L. Ng, J. A. Miller, L. N. Van De
755   Lagemaat, K. A. Smith, A. Ebbert, Z. L. Riley, C. Abajian, C. F. Beckmann, A. Bernard, D. Bertagnolli, A. F.
756   Boe, P. M. Cartagena, M. Mallar Chakravarty, M. Chapin, J. Chong, R. A. Dalley, B. D. Daly, C. Dang, S.
757   Datta, N. Dee, T. A. Dolbeare, V. Faber, D. Feng, D. R. Fowler, J. Goldy, B. W. Gregor, Z. Haradon, D. R.
758   Haynor, J. G. Hohmann, S. Horvath, R. E. Howard, A. Jeromin, J. M. Jochim, M. Kinnunen, C. Lau, E. T.
759   Lazarz, C. Lee, T. A. Lemon, L. Li, Y. Li, J. A. Morris, C. C. Overly, P. D. Parker, S. E. Parry, M. Reding, J.
760   J. Royall, J. Schulkin, P. A. Sequeira, C. R. Slaughterbeck, S. C. Smith, A. J. Sodt, S. M. Sunkin, B. E.

Swanson, M. P. Vawter, D. Williams, P. Wohnoutka, H. Ronald Zielke, D. H. Geschwind, P. R. Hof, S. M.
Smith, C. Koch, S. G. N. Grant, A. R. Jones, An anatomically comprehensive atlas of the adult human brain
transcriptome. *Nature*. **489**, 391–399 (2012).

74.   G. Z. Wang, T. G. Belgard, D. Mao, L. Chen, S. Berto, T. M. Preuss, H. Lu, D. H. Geschwind, G. Konopka,
Correspondence between Resting-State Activity and Brain Gene Expression. *Neuron*. **88**, 659–666 (2015).

75.   J. Richiardi, A. Altmann, A.-C. Milazzo, C. Chang, M. M. Chakravarty, T. Banaschewski, G. J. Barker, A. L.
W. Bokde, U. Bromberg, C. Buchel, P. Conrod, M. Fauth-Buhler, H. Flor, V. Frouin, J. Gallinat, H. Garavan,
P. Gowland, A. Heinz, H. Lemaitre, K. F. Mann, J.-L. Martinot, F. Nees, T. Paus, Z. Pausova, M. Rietschel,
T. W. Robbins, M. N. Smolka, R. Spanagel, A. Strohle, G. Schumann, M. Hawrylycz, J.-B. Poline, M. D.
Greicius, L. Albrecht, C. Andrew, M. Arroyo, E. Artiges, S. Aydin, C. Bach, T. Banaschewski, A. Barbot, G.
Barker, N. Boddaert, A. Bokde, Z. Bricaud, U. Bromberg, R. Bruehl, C. Buchel, A. Cachia, A. Cattrell, P.
Conrod, P. Constant, J. Dalley, B. Decideur, S. Desrivieres, T. Fadai, H. Flor, V. Frouin, J. Gallinat, H.
Garavan, F. G. Briand, P. Gowland, B. Heinrichs, A. Heinz, N. Heym, T. Hubner, J. Ireland, B. Ittermann, T.
Jia, M. Lathrop, D. Lanzerath, C. Lawrence, H. Lemaitre, K. Ludemann, C. Macare, C. Mallik, J.-F. Mangin,
K. Mann, J.-L. Martinot, E. Mennigen, F. Mesquita de Carvahlo, X. Mignon, R. Miranda, K. Muller, F. Nees,
C. Nymberg, M.-L. Paillere, T. Paus, Z. Pausova, J.-B. Poline, L. Poustka, M. Rapp, G. Robert, J. Reuter, M.
Rietschel, S. Ripke, T. Robbins, S. Rodehacke, J. Rogers, A. Romanowski, B. Ruggeri, C. Schmal, D.
Schmidt, S. Schneider, M. Schumann, F. Schubert, Y. Schwartz, M. Smolka, W. Sommer, R. Spanagel, C.
Speiser, T. Spranger, A. Stedman, S. Steiner, D. Stephens, N. Strache, A. Strohle, M. Struve, N.
Subramaniam, L. Topper, R. Whelan, S. Williams, J. Yacubian, M. Zilbovicius, C. P. Wong, S. Lubbe, L.
Martinez-Medina, A. Fernandes, A. Tahmasebi, Correlated gene expression supports synchronous activity in
brain networks. *Science (80-.)*. **348**, 1241–1244 (2015).

76.   J. Seidlitz, F. Váša, M. Shinn, R. Romero-Garcia, K. J. Whitaker, P. E. Vértes, K. Wagstyl, P. Kirkpatrick
Reardon, L. Clasen, S. Liu, A. Messinger, D. A. Leopold, P. Fonagy, R. J. Dolan, P. B. Jones, I. M. Goodyer,
A. Raznahan, E. T. Bullmore, Morphometric Similarity Networks Detect Microscale Cortical Organization
and Predict Inter-Individual Cognitive Variation. *Neuron*. **97**, 231-247.e7 (2018).

77.   K. J. Gorgolewski, G. Varoquaux, G. Rivera, Y. Schwarz, S. S. Ghosh, C. Maumet, V. V Sochat, T. E.
Nichols, R. A. Poldrack, J.-B. Poline, others, NeuroVault. org: a web-based repository for collecting and
sharing unthresholded statistical maps of the human brain. *Front. Neuroinform*. **9**, 8 (2015).

78.   D. W. Huang, B. T. Sherman, R. A. Lempicki, Systematic and integrative analysis of large gene lists using
DAVID bioinformatics resources. *Nat. Protoc*. **4**, 44–57 (2009).

79.   D. V Klopfenstein, L. Zhang, B. S. Pedersen, F. Ram\'\irez, A. W. Vesztrocy, A. Naldi, C. J. Mungall, J. M.
Yunes, O. Botvinnik, M. Weigel, others, GOATOOLS: A Python library for Gene Ontology analyses. *Sci.*
Rep. **8**, 10872 (2018).

80.   J. Chen, E. E. Bardes, B. J. Aronow, A. G. Jegga, ToppGene Suite for gene list enrichment analysis and
candidate gene prioritization. *Nucleic Acids Res*. **37**, W305--W311 (2009).

Acknowledgments: We thank Prof. Hans van Bokhoven for comments on the early results of this study,
Dr. Yuhua Guo for providing brain activation coordinates of unpublished studies. **Funding:** W.L. is
funded by the PhD scholarship (201606990020) of the Chinese Scholarship Council. **Author**
contributions: The experiment was designed by W.L., G.F., and N.K. Data were collected by W.L., N.P.,
and N.K. Analyses were performed by W.L. and W.L., N.P. wrote the manuscript. W.L, G.F., and N.K.
revised and edited the manuscript. All authors provided essential feedback on the manuscript at different
stages, contributed substantially to, and approved the final version of the manuscript. **Competing**
interests: All authors declare that they have no competing interests. **Data and materials availability**: All
data reported in this study was public available via our Open Science Framework and Neurovault
repository. All custom code is available from the authors upon request and will be released via the
personal GitHub as soon as possible.

Figures and Tables

Figure 1 Schematic of the pipeline for the investigation of neural and transcriptional commonality. (A) Activation likelihood estimation (ALE) meta-analyses of functional MRI studies. MRI coordinates of reported brain activations from relevant studies were used to generate the activation map for each tasks of interest. (B) Meta-analytic connectivity modeling of seed regions resulting from the ALE analyses was conducted to search co-activated regions over all studies in BrainMap. Coactivation maps were estimated via ALE. (C) To generate behavioral profiles of co-activation maps, we used Neurosynth to acquire a list of meta-maps including the behavioral terms from low-level to high level cognitive processes (total Number of terms=23). Next, each co-activation map was compared to all meta-maps (e.g. inhibition, decision making, and social cognition) to compute the similarity index for each meta-map. (D) The activation patterns were associated to gene expression maps from Allen Institute for Brain Science (AIBS). Each gene expression map was vectorized based on the expression measures within the three dimension maps. The ALE values at the corresponding brain regions on the activation map were extracted and vectorized. To identify the genes whose spatial patterns are similar to specific activation map, similarity between the all vector pairs were quantified. "Inhibitory-related" genes were defined by identifying genes of which the expression patterns are correlated with the activation maps of four paradigms (ER, Emotion Regulation; TNT, Think/No-Think; SS, Stop-signal; GN, Go/No-go). (E) "Inhibition-related" genes were associated with biological functions (GO terms) and diseases terms for the interpretation using Gene Ontology Enrichment Analysis (GOEA).

Figure 2 Brain activity underlying four inhibitory control tasks revealed by Activation likelihood estimation (ALE) meta-analyses. (A) Brain regions significantly more activated in the "Regulation" compared to the "Baseline" condition during the Emotion Regulation task. (B) Brain regions significantly more activated in the "Control" compared to the "non-control" condition during the memory control task. (C) Brain regions significantly more activated in the "No-Go" compared to the "Go" condition during the Go/No-go task. (D) Brain regions significantly more activated in the "Stop" compared to the "Go" condition during the Stop-signal task.

Figure 3. Co-activation maps of Emotion Regulation and Memory Control revealed by Meta-analytic connectivity modeling (MACM). (A) Co-activation maps for regions of interests (ROIs) resulting from ALE analysis of the emotion regulation. (B) Co-activation maps for ROIs resulting from ALE analysis of the memory control. ALE-based ROIs are projected onto glass brains (left column), and co-activation patterns are rendered on an MNI template second to ninth column (right columns)

Figure 4 Behavioral profiles of co-activation maps across four inhibitory control tasks. Spatial similarity between Neurosynth meta-maps and co-activation maps across 23 topic terms. Terms are ordered by the mean r-values across the row(s) (all co-activation maps). "Working memory", "cognitive control", and "inhibition" are located at the top, suggesting the common stronger association. Domain-specific cognitive functions (e.g. autobiographical memory, emotion) are located at the bottom, suggesting a limited association.

Figure 5 Common transcriptional profiles revealed by activation-gene expression association analysis. (A) Emotion Regulation-related brain activation patterns are correlated with the expression pattern of 1529 genes. Memory control-related brain activation patterns are correlated with the expression pattern of 2531 genes. 1130 genes correlated with both the emotion regulation and memory control-related brain activity patterns. (B) Visualization and comparison of brain activation patterns and expression patterns of "Top Genes". "Top Genes" are 10 genes with the most similar expression patterns as the brain activation. (C) The gene lists of two inhibitory control tasks were more overlap with each other (Within Inhibitory) compared to one of the inhibitory control and DMN (Inhibitory & DMN). (D) Disease associations of "inhibitory-related" genes. "Inhibition-related" genes are associated with genetic risks for major psychiatric disorders and Parkinson disease. Terms are ordered by the percent (Left Green) of hit count in query list to hit count in genome. Hit count in query list is the number of genes belongs to the "inhibitory-related" genes (Right Dark Blue). Hit count in genome is the total number of genes associated with risk for certain disease terms based on the DisGeNET (Right Light Blue).

895 **Table1** Significant activated clusters during emotion regulation and think/no-think task

Task	Brain region	Hemisphere	MNI coordinates	Cluster size	Peak voxel Value
Emotion Regulation	Insula/IFG	R	36 16 6	209	0.02056
	IFG	L	-46 44 -4	138	0.02051
	Insula	L	-34 16 12	156	0.02089
	Middle Cingulate Gyrus	L/R	10 22 44	215	0.0189
	Inferior parietal lobule	R	60 -42 46	90	0.01688
	SMA	L	-4 6 62	110	0.0217
Memory Control	Insula	L	-36 20 -4	127	0.017665
	IFG	R	34 38 30	303	0.019
	Inferior parietal lobule	R	64 -48 28	126	0.0155
	Precuneus	L	-16 -64 56	159	0.017254
	SMA	R	12 12 60	169	0.022948

896   IFG: inferior frontal gyrus; SMA: supplementary motor area; DLPFC: dorsolateral prefrontal cortex