HD-tDCS over mIPS causally modulates online reach correction

Alexander Goettker[1], Brandon Caie[2], Jerrold Jeyachandra[2], Sisi Xu[2], Jason Gallivan[2], Jacek Dmochowski[3], Katja Fiehler[1,4] & Gunnar Blohm[2]

[1] Justus-Liebig University Giessen, 35394 Giessen, Germany

[2] Center of Neuroscience Studies, Queen's University, Kingston, ON, Canada

[3] Department of Biomedical Engineering, City College of New York, New York NY 1003

[4] Center for Mind, Brain and Behavior (CMBB), Universities Marburg and Giessen, Germany

Abstract

Brain lesion and stimulation studies have suggested posterior parietal cortex and the medial intraparietal sulcus in particular as a crucial hub for online movement error corrections. However, causal evidence for this is still sparse. Indeed, lesion studies are potentially confounded by compensatory reorganization mechanisms while brain stimulation studies have produced heterogeneous results when employing transcranial magnetic stimulation. Here we employed a complementary approach using fMRI-guided high-definition transcranial direct current stimulation (HD-tDCS) of the left medial intraparietal sulcus (mIPS) to re-examine the role of mIPS in online reach corrections to jumping targets. We obtained two independent measures of stimulation-induced changes in brain activity by modeling current flow in the brain and through EEG recordings before and after the stimulation. To quantify behavioral effects we computed movement curvature as a measure of online correction. We demonstrate that both of our measurements of brain activity were consistent with a polarity-specific modulation of the online correction for targets jumping to the contralateral side of the stimulation. Importantly, using a mediation analysis of the relationship between stimulation current and movement curvature, we provide causal evidence that the induced current modifies brain activity, which then leads to the observed behavioral changes. This unique combination of methods and analysis thus provides complementary evidence for the crucial role of the posterior parietal cortex in online error correction, while at the same time setting a new methodological standard with respect to the causal influence of transcranial direct current stimulation.

Keywords

Motor control, arm movements, parietal cortex, transcranial electrical stimulation, EEG

Correspondence

Alexander Goettker, Justus Liebig University Giessen, Otto-Behagel Strasse 10F, 35394 Giessen, Germany; Contact: Alexander.Goettker@psychol.uni-giessen.de

Introduction

In order to interact with objects in our environment we often bring our hands to the object to manually explore and manipulate them. Those movements can be inaccurate due to sensory-motor noise (Harris & Wolpert, 1998; van Beers, Baraduc & Wolpert, 2002; Faisal, Selen & Wolpert, 2008) or sudden changes in the environment. To compensate for these internal and external factors, our online-control system can correct for errors that arise during the movement (see Gaveau et al. 2014 for a recent review). The online-control system reacts fast and automatically (Pisella et al., 2000) and occurs without the awareness of an error (Pelisson, Prablanc, Goodale & Jeannerod, 1986). Online-corrections in response to changes in the environment appear to be based on a comparison of a continuously updated representation of the target position and an internal estimate of the state of the hand (Prablanc & Martin, 1992; Desmurget et al. 2001). While the target position is usually determined by visual information (Sarlegna et al., 2003), the internal estimate of the hand state relies more on a combination of somatosensory feedback and an efference copy of the motor command (Miall & Wolpert, 1996; Scott, 2012), as corrections are possible without the vision of the hand (Pelisson et al., 1986; Sarlegna et al., 2015).

There is a vast literature on online-correction behavior (Prablanc, Desmurget & Grea, 2003; Wijdenes, Brenner & Smeets, 2011; Gaveau et al. 2014, Sarlegna & Mutha, 2015); however, much less is known about the underlying neural mechanisms during online control (see Battaglia-Mayer et al. 2014). One of the important neural structures implicated in the online-correction process is the posterior parietal cortex (PPC); situated between visual and motor areas representing hand and target locations respectively (see Battaglia-Mayer et al. 2014; Archambault et al., 2015). There is evidence for the activation of the PPC during visually guided movements (Ogawa et al. 2006; Limanowski et al., 2017) and TMS over the PPC impairs online-corrections contralateral to the side of stimulation (Desmurget et al.1999; Tunik et al. 2005). In addition, lesions in the PPC result in an absent online correction response while leaving reaching to static targets unimpaired (Pisella et al.,2000; Grea et al.,2002), indicating a specific causal role of the PPC in online-corrections. More specifically, lesions in the intraparietal sulcus have been related to optic ataxia (Rossetti et al., 2003; Anderson et al. 2015), which manifests in impaired reaching to peripheral targets (Hwang & Anderson, 2012) and impaired online control (Grea et al. 2002)

Despite this evidence for a causal role of the PPC there are some caveats. A recent study

which used TMS at four different locations in the PPC was not able to replicate an effect on

online control of hand or foot movements (Marigold et al., 2019), suggesting that some parts

of the PPC might be more involved in movement planning than in online control.

Additionally, due to their complexity TMS studies typically only have a limited sample of

participants which often show inconsistent results with a general increase in the variability of

the online correction behavior (Desmurget et al. 1999). Lesion studies (see Cavina-Pratesi,

Connolly & Milner, 2013; Buiatti, Skrap & Shallice, 2013) provide evidence for some causal

involvement of the PPC, but cannot point to the exact mechanisms underlying behavioral

differences (Jonas & Kording, 2017) and suffer the confound of compensatory brain

reorganization mechanisms (see Rorden & Karnath, 2004).

Therefore, we aimed at reinvestigating the causal role of the PPC in online control by using

transcranial direct current stimulation (tDCS). tDCS is a noninvasive method to increase or

decrease cortical excitability in humans. In contrast to TMS, which disrupts the cortical

output of an area through the generation or inhibition of spikes (Siebner, Hartwigsen,

Kassuba, & Rothwell, 2009, see Liu et al. 2018), tDCS is capable of bidirectionally

influencing dendritic input through modulation of pre-synaptic activity (Bączyk &

Jankowska, 2014). As such, tDCS offers a valuable complementary approach to previous

lesion or TMS studies, as tDCS allows for a continuous and long-lasting stimulation.

Importantly, through recent advances it is possible to model the flow of the induced current

(Dmochowski et al, 2011; Saturnio et al, 2015, 2018). Modeling current flow provides the

opportunity to estimate the orientation and the magnitude of the current that impinges on the

target area to predict changes in the brain activity and inter-individual differences on the

behavioral level. Using this approach, we targeted the middle intraparietal sulcus (mIPS), a

region in the PPC, which represents a reach and eye movement map and is involved in

movement planning (Beurze et al., 2007; Gertz & Fiehler, 2015; Vesia et al., 2010; Gallivan

et al., 2011).

To independently measure and verify the electrophysiological changes induced by tDCS, we

used EEG recordings. EEG has been used to investigate the effects of tDCS on cortical

excitability in humans (Mangia et al, 2014; Lauro et al, 2014), but the sources of variability

underlying the relationship between behavioral and neural changes following tDCS are not

well established. Here, we address this relationship by measuring EEG before and after

stimulation and relate these recordings to the individual sources of anatomical variability

through current modeling. In addition, we use the EEG recordings to test if changes in brain activity are directly related to changes in the online correction behavior. Of special interest, are particular brain oscillations that are thought to be a mechanism for cortical communication during motor planning and execution (Rubino, Robbins, & Hatsopoulos, 2006; see Battaglia Mayer, Ferrari-Roniolo & Visco-Comandini, 2015). Specifically, osciallations in the beta-band (15-30 Hz) have been implicated in the communication between the PPC and premotor cortex (Pani et al. 2014; Stetson & Anderson, 2014), somatosensory cortex (Zhang et al. 2008; Tsujimoto et al., 2009) and primary motor cortex (Menzer et al., 2014). In addition, the beta-band power is thought to be inversely related with movement preparation (Pani et al. 2014; Stetson & Anderson, 2014) which is crucial especially for online correction.

Thus far, previous research has investigated the relationship between tDCS and EEG (e.g. Mangia et al, 2014; Lauro et al, 2014), tDCS and behavior (e.g. Jacobsen et al., 2012) and EEG and behavior (e.g. Pani et al. 2014; Stetson & Anderson, 2014), however to our knowledge our study is the first one that combines all these relevant measures. We designed a mediation analysis (Baron & Kenny, 1986) based on the modeled current, changes in brain oscillations and the changes in behavior to reconstruct the causal structure of the stimulation. By leveraging the double dissociation in current polarity afforded by tDCS and our unique set of measurements, we demonstrate that the induced current is related to the behavioral changes, but importantly this relationship is mediated by altered brain activity. By applying such an interventional methodology (Marinescu, Lawlor & Kording, 2018), our results provide causal evidence for the role of mIPS in online reach correction, while also establishing a new methodological approach with respect to the causal influence of transcranial direct current stimulation.

Methods

Overview. By stimulating the left mIPS with HD-tDCS we aimed to provide complementary evidence for a causal role of the PPC during the online control of right-arm reaching. By combining detailed psychophysics, current forward modeling and independently measured stimulation-induced EEG changes, we sought to establish a causal relationship between the induced current, changes in brain activity and the altered behavior. We made use of the bidirectional effect of tDCS by stimulating all participants with anodal and cathodal

stimulation in randomized order on two different testing days, separated by at least one week.
To achieve anatomically accurate stimulation of the targeted area, we used task-based fMRI
to localize the left mIPS in each participant and registered the location on the skull (see fMRI
localizer task and **preparation of stimulation**) to place the electrodes for the stimulation
accordingly (see **HD-tDCS**). Based on the placement of the electrodes and the participant's
brain scan we were able to model the current flow and quantify the expected effect of the
stimulation at the targeted site on a subject-by-subject level (see **Current Forward**
Modeling). For the behavioral experiment, each day was divided into three different
consecutive sessions (baseline, stimulation, and post-stimulation). To independently measure
changes in brain activity caused by the stimulation we used the same electrodes as during
HD-tDCS stimulation to record the EEG response in the baseline and post-stimulation
sessions (see **EEG Recording**). During the baseline session participants performed four
blocks without any stimulation to obtain a representative measurement of their performance.
The task was to reach to the location of visually presented dots, and some of them jumped to a
new location at hand movement onset (see **Experimental setup and paradigm**). During the
stimulation session participants also completed four blocks, but now with constant direct
current stimulation. The stimulation duration was 20 minutes, and all of the participants
finished their four blocks during the time of stimulation. After stimulation, we measured post-
stimulation trials, where participants again completed four blocks without stimulation**.** We
quantified the online correction performance by measuring the curvature of the respective
movements (see **Behavioral Data**). This combination of measurements allowed us to perform
a mediation analysis to establish a relationship between the induced current and behavioral
changes, while showing that the behavioral changes are mediated by underlying changes in
brain activity as measured through EEG.

Participants. Ten healthy right-handed volunteers participated (age range: 20 – 39 years; 7
females) in the experiment. All participants had normal or corrected-to-normal vision. All
participants gave informed written consent to procedures that were approved by the Queen's
University General Board of Ethics. Participants received financial compensation for their
participation. One participant was excluded from the analysis as he did not exhibit online
corrections during his baseline testing session.

Figure 1. Overview of experiment. (**A**) Sketch of experimental setup. Participants sat in a
chair in front of the tilted surface, which they viewed through a semi-silvered mirror. In this
way, visual feedback about their hand position was occluded. (**B**) General overview of the
different sessions. After they completed the functional localizer task in the MRI scanner,
participants completed the main experiment on two testing days, separated by one week. (**C**).
Depiction of the different trial types during the experiment. In static trials participants had to
perform reaching movements to one of three possible target locations. In the jumping trials
participants started their hand movement to the central target but it jumped either to the left or
the right at hand movement onset.

fMRI localizer task. To localize the left mIPS in each participant, all participants took part in

a localizer testing session in an MRI-scanner prior to the main experiment (see figure 1B).

MR imaging was performed at the Queen's MRI facility using a 3-Tesla Magnetom Trio

scanner (Siemens Medical Systems, Erlangen, Germany). The left mIPS was functionally

localized on an individual basis using an interleaved center-out reaching task (Vesia et al.,

2010; Vingerhoets, 2014).

Inside the MRI, the participant rested supine with a 32-channel head coil worn on the head.

All stimuli were rear-projected using an LCD projector (NEC LT 265 DLP projector;

resolution, 1024 x 768; 60 Hz refresh rate) onto a screen mounted behind the participant, and

viewed by the participant via a mirror mounted to the head coil directly above the eyes. In

each task block, the participants were presented with a circular arrangement of 8 white circles (~15 degrees of visual angle) surrounding a central red or green square. When the central square was red, participants were instructed to make self-paced center-out reaching movements (small pivots about the wrist and elbow with their index finger) while fixating on the central fixation square; when the central square was green participants were instructed to make self-paced center-out saccadic eye movements while keeping their hand in a relaxed home position on their chest. The reach-saccade localizer included 8 reaching blocks (16 s per block), 8 saccade blocks (16 s per block), and 3 fixation/baseline blocks (10 s per block), which were placed at the beginning, middle and end of the run. The localizer run lasted a total of 5 min 22 s (157 brain volumes).

Functional MRI volumes based on the blood oxygenation level-dependent (BOLD) signal were collected using T2*-weighted gradient echo planar imaging with 240 mm x 240 mm field of view with 80 x 80 acquisition matrix size for an in-plane resolution of 3mm, repetition time (TR) = 2 s; time to echo (TE) = 30 ms, and flip angle (FA) = 90°. Each volume comprised of 35 contiguous axial slices (slice thickness 4mm) acquired in an interleaved order along the anterior commissure-posterior commissure (AC-PC) line. A 176 slice, high-resolution T1-weighted structural volume was acquired along the same orientation as the functional images using a 3D MP-RAGE sequence (single shot, ascending sequence in the sagittal plane with TR = 1760 ms, TE = 2.2 ms, FoV = 256 mm, flip angle = 90°, and voxel size = 1 mm^3).

Data from the localizer run were registered to the corresponding structural image, aligned on the plane between the anterior commissure and posterior commissure. All preprocessing and univariate analyses were performed using Brain Voyager QX version 2.6 (Brain Innovation). Preprocessing for the localizer data included slice scan-time correction, 3D motion correction (alignment of the first volume of the functional scan, which was closest in time, to the anatomical scan), high-pass temporal filtering of 3 cycles/run, functional-to-anatomical co-registration, and spatial smoothing (trilinear-sinc interpolation performed during realignment, sinc interpolation performed during reorientation). We carried out subject-level analyses using a general linear model (GLM) with condition predictors created from boxcar functions convolved with a double-gamma hemodynamic response function (HRF). These were aligned to the onset of each reach and saccade block, with its duration based on block length. In a whole-brain analysis for each, the BOLD signal was contrasted between the reaching condition and the fixation condition to identify the hotspots of the left mIPS. The location of

the left mIPS was defined by selecting the voxel peak medial to the intraparietal sulcus in the left hemisphere (Vingerhoets, 2014).

Preparation of stimulation. During the main experiment, we used neuronavigation (Brainsight, Rogue Research Inc., Montreal, Quebec) to place the central electrode on the scalp over the fMRI-identified location of the left mIPS, so that the peak electric field would be directed to the mIPS. The participant-image registration used the landmarks of the nose tip, the nasion and the left and right intertragal notches. After the registration, the target location on the scalp was identified as the inline projection of the selected target region mapped onto the scalp that minimized the distance between target region and the target location on the skull.

After the registration of the target location on the scalp with Brainsight, a 10-20 system EEG mesh cap was fastened on the head of the participant. Embedded plastic holders in the mesh cap were used as holders for the electrodes. The central electrode was put directly above the target location on the scalp (by adjusting the cap), and additionally 4 pick-up electrodes were placed in a 3 cm radius surrounding the central electrode, leading to a 4x1 electrode configuration for HD-tDCS (Datta et al., 2009).

Experimental setup and paradigm. Participants sat in a dark room at a tilted reaching setup (see figure 1A). Their head was immobilized by a personalized dental impression, attached to the setup. The tilted reaching setup consisted of an overhead monitor (ViewPixx 1920*1200 Pixel; 52 *29 cm, VPixx Technologies Inc., Saint-Bruno, QC Canada) for the presentation of the visual stimuli, a tilted reaching surface below the screen and a semi-silvered mirror positioned exactly in between the reach surface and the screen so that stimuli presented on the screen were projected onto the reach surface. The mirror allowed the participants to see the visual stimuli, but occluded the sight of their hand. All surfaces were tilted by 30°, which allowed for comfortable reaching and facilitated eye-tracking. The eye-tracker was attached to the frame of the semi silvered-mirror. The eye movements were tracked with an EyeLink 1000 eye tracking system (SR Research; sampling rate, 500 Hz; accuracy, 0.5°). Hand position during the reaching was recorded by localizing the 3D position of infrared light-emitting diodes taped to the tip of the right index finger (Optotrak Certus; Northern Digital; sampling rate 400 Hz; accuracy, 0.1 mm).

Participants were asked to perform reaching movements from an initial starting location to a visual target at one of three possible target locations. In some of the trials, the visual target

changed position at the onset of hand movement. At the beginning of each trial, a white dot (30 pixels diameter) indicated the starting location. The starting location was presented on the horizontal midline slightly above the bottom end of the screen. Participants had to place their right hand at the starting location and were told to also look there. Although participants could not see their own hand in the experiment, during the beginning of the block or after the end of the previous trial a red marker indicated their hand position within 5cm of the start position to guide them back to the starting location. When the hand was kept at the starting location for 500 ms (defined as a frame of 50 pixels around the starting location) the starting location disappeared and a dot representing the target locations appeared. The central target was presented in a distance of 23 cm above the starting location, at the midline of the screen. The left and the right target locations were placed at the same height, but displaced laterally by 7.5 cm relative to the center target. As soon as the target appeared, participants were instructed to make a reaching movement to the target location and were allowed to move their eyes. The target stayed on for 1000 ms and afterwards the starting location appeared to signal the start of the next trial (see figure 1C).

Each block had a total of 80 trials, lasting approximately 4.5 minutes. Across trials the different target locations were randomized. Of the 80 trials, 20 trials changed the target position at hand movement onset. This small amount of target jump trials mitigated participants ability to predict jump trials. For the static targets the target was equally likely to appear at each of the target locations (left, center, right). The 20 trials jumping trials initially appeared at the central location, but jumped with equal probability to the left or right at hand movement (jumping left, jumping right). Hand movement onset was calculated online as the moment where the hand was outside of the initial frame around the starting location. In total, participants completed 1920 trials (2 polarities * 3 sessions * 4 blocks * 80 trials) including 480 jumping trials per participant.

EEG Recording. During the baseline and post-stimulation session we used the central and four surrounding electrodes to record the EEG-response. This allowed us to assess the influence of HD-tDCS on oscillatory signatures of reaching. EEG data was acquired from the same electrodes delivering the stimulation using a 16 channel USB amplifier (V-AMP, Brain Products). Reference and ground electrodes were placed on the seventh cervical vertebrate and the olecranon respectively to minimize noise from muscular activity. EEG data was sampled at 2 kHz and converted from analog to digital with a resolution of 24 bits at 0.0489

µV/bit. Event timing was tracked by a trigger shared between the EEG amplifier and the
Eyelink 1000, allowing for behavioral events to be time-locked to the EEG signal.

HD-tDCS. The electrodes were sintered Ag/AgCl ring electrodes (HD-target package, Soterix
Medial Inc., New York, NY, USA) with a diameter of 1 cm. To improve the conductance, the
hair was separated under the scalp using cotton swabs and the plastic holders were filled with
conducting gel. An impedance check was performed prior to the experiment to ensure
appropriate impedance limits (< 20 kΩ).

The stimulation was administered for 21 minutes. During the first 30 seconds the current
strength ramped up and during the last 30 seconds ramped down again, leading to 20 minutes
of constant stimulation with a current strength of 2mA. The stimulation was controlled by
microprocessor-controlled constant current source (DC-STIMULATOR MC 8, NeuroConn
GmbH, Ilmenau, Germany) to deliver continuous direct current. The direction of the current
led to either anodal or cathodal stimulation under the central electrode.

Data Analysis

Behavioral Data. First, the data obtained by the Optotrak and Eyelink were low-pass filtered
by an autoregressive forward backward filter with 50Hz cut-off frequency. Then, velocity and
acceleration were computed using a central difference algorithm. The offline analysis was
performed in MATLAB (The Mathworks, Natick, MA). The hand movement onset was
defined as the point where the hand was moved outside of the initial frame around the start
position. The end position was defined as the time when the velocity of the hand was lower
than 1 cm/s.

The onset of online correction was defined by the interpolation method of Oustwoud-Widjens
et al. (2014). The average x-position of a movement to the central target was calculated for
each session. Afterwards the difference between the x-position of each single jumping trial
and the average for the central location was computed. This gave us a difference function,
where we performed a linear regression between the points of 25% and 75% from the
absolute maximum difference. The point where this line crosses zero was used as the onset of
online correction. Based on the detected movement endpoints we computed the horizontal-
and vertical-error and the precision of the movements. Horizontal- and vertical-error were
based on the distance between the endpoint and the target location and the precision of the

movements was defined as the area of an error ellipse which contained 95% of the movement endpoints.

The main variable that we used to quantify the online correction performance was the normalized curvature. The normalized curvature was defined as the maximum distance between the movement trajectory and a straight line which connected the start and the endpoint of the hand movement, divided by the length of the trajectory of the movement. Additionally, we did not only look at the absolute curvature of the trajectory, but also analyzed the development of curvature over time. Therefore, we aligned all trials on the calculated onset of the online correction and calculated the curvature in the same way with a sliding window of 100 ms in steps of 10 ms. We normalized the curvature over time based on the maximum curvature for the respective baseline session and then compared the peak curvature depending on the session and stimulation type. To quantify the effect of the stimulation on behavior we computed the difference between the stimulation and the baseline session for anodal and cathodal stimulation for our two measurements of curvature, absolute curvature and the peak curvature in the time course after correction onset. We collapsed the data over the four blocks of each session (except for the baseline where we only used the last 3 blocks, see exclusion of trials) and calculated the mean and the standard deviation for each participant for each of the sessions for the different variables. To test for systematic influences on the movement endpoints we ran a repeated measures ANOVA with the factors polarity (anodal, cathodal), session (baseline, stimulation, post-stimulation) and target type (static vs online corrected) separated for targets on the left and right. To test for effects of the stimulation on online correction we conducted repeated measures ANOVAs with the factors polarity (anodal, cathodal), target location (jumping left, jumping right) and session (baseline, stimulation, post-stimulation). As post-hoc analysis we conducted one-way ANOVAs with the factor session for either the anodal or cathodal values or post-hoc t-tests with Bonferroni correction.

Current Forward Modeling. We simulated the induced electric field at the site of stimulation by constructing a tetrahedral mesh from the T1 and T2-weighted MRI of each participant and numerically solving a forward model for Maxwell's equations across tissue boundaries (see Figure 2). In tDCS, the current flows through a heterogeneous medium with a complex spatial distribution that depends on both head geometry and the electrical conductivities of the various tissues. Thus, it is important to consider the influence of

individual differences in anatomical properties such as cortical surface folding and skull
thickness on the current path from scalp to target.

Figure 2. Source reconstruction of HD-tDCS over mIPS. The left medial intraparietal sulcus
(mIPS) was localized during a separate functional MRI scan (Methods: Localizer task), and
superimposed over a segmented head model (A-E). T1-weighted structural scans were
segmented into tissue categories for finite element modeling of the electric field induced by
HD-tDCS (Methods: Current Forward Modeling). Visualized is the electric field density across
the scalp (A), skull (B), pial layer (C), gray matter (D), and white matter (E), showing a
progressive decrease in the strength of the resulting electric field closer to neural tissue. Current
models were performed in SimNIBS, with tissue conductivity values given in Table 1.

The MR images were segmented into tissue categories (scalp, skull, cerebrospinal fluid, gray
matter, white matter) using an automated procedure in Freesurfer (Dale et al, 1999). This
generated a tetrahedral mesh that was used as the model for simulating volume conduction in
SimNIBS (Saturnino et al., 2018). A detailed explanation of the pipeline can be found in
Saturnino et al (2018). Tissue conductance values used are reported in Table 1. The
localization of mIPS was transformed from the Brainvoyager System Coordinates to
SimNIBS space to guide the central electrode placement in the model with a custom Matlab
function. Electrodes are modelled as elliptical nodes of dimensions 1cm x 1cm x 1mm on the
tetrahedral surface. A ring mesh was generated with a 3cm radius to automatically place the 4

surrounding electrodes. Center electrodes were set to apply +/- 2mA, and each surrounding electrode -/+ 0.5mA. The system of partial differential electromagnetic equations was solved using a conjugate gradient method to compute the vector field.

The resulting electric field is a vector field with a magnitude and orientation at the target site. We thus computed the magnitude of the radial and tangential (orthogonal complement of radial) component of the induced current at the localized mIPS site for a regression analysis with EEG and behavior. Electric fields were visualized with Gmsh (Geuzaine & Remacle, 2009) and MATLAB .

Table 1: Tissue Conductivity Values based on Wagner, 2004

White matter	**0.126 S/m**
Gray Matter	**0.275 S/m**
CSF Volume	**1.654 S/m**
Bone	**0.01 S/m**
Scalp	**0.465 S/m**

EEG Analysis. Data was band-pass filtered between 1 and 45 Hz using a zero-phase digital filter. A notch filter excluded frequencies between 55 and 65 Hz to filter noise from the power line frequency. Each electrode was filtered independently. Trials were then aligned in separate analyses to the hand movement onset. A surface Laplacian was computed with respect to the central electrode in order to topologically localize the filtered data to the point on the scalp closest to mIPS (Babiloni et al, 2001). As we were interested in the online correction performance we used only the trials where a target jump occurred and due to the lateralization of the stimulation we also separated the analysis for trials jumping to the left or right.

For the time frequency analysis, data was analyzed in a 750 ms window covering [-250,500] ms relative to the target jump. Time-frequency energy representations were computed using a complex Morlet wavelet (BrainWave Toolbox; Jobst et al., 2018) with a width parameter of 7 centered on each time (0-500ms) and frequency (1-45 Hz) following the method of Tallon-Baudry et al (1997). The data was baseline variance-normalized using the z score with respect to each time vector. We computed such a representation for each baseline and post-stimulation session for the anodal and cathodal stimulation and again separated targets

jumping to the left and right. Then we again quantified the effect of the stimulation by subtracting the baseline representation from that in the post-stimulation session.

Mediation analysis. To examine the relationship between the behavioral changes, the modeled current flow, and the EEG response, we performed a mediation analysis (Baron & Kenny, 1986; see MacKinnon et al., 2007). Due to the lateralization of the stimulation. this was completed separately for trials jumping to the left and to the right. First, we established a relationship between the induced current based on the forward modeling and the behavioral changes. In a second step we correlated the induced current with the changes in the EEG to establish a relationship between stimulation and the potential mediator. To do so, we correlated the current vector with each point in the time frequency representation (TFR). By doing so we found two spots in the TFR response that were correlated with the induced current (first 100 ms after target jump for 11 to 14 Hz and 100 to 200 ms from 20 to 26 Hz). To be able to represent that correlation and use it for the mediation analysis, we averaged the TFR differences in these time windows and took it as a measure of induced changes in the EEG. Following this, we correlated the changes in EEG with the behavioral changes. After these individual steps, the crucial step was now to compute the partial correlation between the induced current and behavior while controlling for the changes in EEG, allowing us to establish the causal structure of our stimulation.

Exclusion of trials. We had to exclude some trials from our analysis due to a missing signal of the Optotrak marker or where we could not determine hand movement onset or offset (~6.5% of trials). Additionally, we excluded trials with hand movement onset latency faster than 100 ms or slower than 700 ms, indicating that participants probably did not conform to the task (~5.9% of trials). For the online correction targets, we excluded trials where the maximum deviation between the trajectory to the central target and the jumped target was less than 3 cm and the calculated online correction latency was outside of 100 and 700 ms, indicating that no appropriate correction had been made (<0.1% of trials). To calculate the average value for the baseline session we excluded trials from the first block, in order to eliminate contamination from any practice effects.

This led to a total of 13850 out of 15840 trials (9 Participants *1920 trials - 9 Participants * 2 Days* 80 trials for the exclusion of the first block). For our EEG analysis we had to exclude the data from one participant as due to technical problems with determining the timing of the trials in the EEG-recording.

Data and analysis code for this project is available at: https://osf.io/kbyxc/

Results

We used HD-tDCS over the left mIPS to elucidate the causal role of the PPC for online-control during reaching movements. In our task, participants had to perform reach-to-point movements to three different target locations. Occasionally the central targets would jump to the left or right at hand movement onset, thereby eliciting an online adjustment of the movement trajectory. We hypothesized that, depending on the polarity of the stimulation, there would be a differential effect on the online correction: cathodal stimulation, which should lead to a reduction of activity similar to artificial or real lesions, should lead to worse online correction performance. This would be reflected in a higher curvature of the movement. Anodal stimulation, which should lead to an increase in activity, should lead to an improved online correction performance, thus a lower curvature of the movement, In a next step, we used two independent measures of the changes in brain activity, one based on reconstruction of the current flow and one based on changes in EEG activity, and related these to the behavioral changes. The combination of our measurements allowed us to unravel the causal structure of the stimulation effect by linking them in a mediation analysis.

Figure 3. **Depiction of hand movements** (**A**) Example reach trajectories of one baseline session for one representative participant. Grey paths show movements to static targets, black lines show movements to targets that jumped at hand movement onset. The red circles show the starting position and the possible target locations. (**B**) Corresponding end points of the reaches in (A). Again grey circles show the end points of movements to static targets, the black circles for the targets that jumped at hand movement onset. The red circles depict the target location. Separate error ellipses were fitted to each of the possible conditions.

Behavioral Analysis

Movement accuracy and precision. To verify that participants were able to adjust their movements online we examined the trajectories of the individual arm movements (see Figure 3A). Participants were able to correct their movement in flight and land close to the same position as for the static movements. In order to assess the performance of the reaches we analyzed at the endpoints of the reaching movements. We calculated the horizontal- and vertical-position error as measurements of the accuracy and fitted an error ellipse to the endpoints of each condition as an indicator of the precision (see Figure 3B). We did not observe any systematic modulations of the horizontal error or precision of the movements and the endpoints for the corrected movements only had a higher vertical error than the comparable static movements across the different sessions ($F(1,8) = 15.061$, $p = .005$ for jumping left; $F(1,8) = 11.900$, $p = .009$ for jumping right). Importantly, there was no polarity specific effect on the endpoints for any of the measures. Based on this result, we continued with analyzing online control of the reaching movement with confidence that any polarity-specific effect we observe there cannot be explained by a systematic variation of the endpoint based on the stimulation.

Movement curvature. To measure the effectiveness of the online correction we looked at the normalized curvature of the reaching trajectories. As mentioned earlier, a lower curvature indicates a smoother integration of the updated movement plan. A hypothetical maximal curvature value would be obtained if the movement was not corrected inflight but if the trajectory would be based on two completely separate movements. We used a repeated measurement ANOVA with three factors (polarity, target location, session) to test for possible effects on the online correction behavior. We found a significant main effect of session (F $(2,16) = 4.584$, $p = .027$, $\eta^2 = .364$) and two interactions. The main effect was driven by a decrease in curvature across each session (Base: $M = 0.160$ +/- 0.031; Stimulation: $M = 0.154$ +/- 0.030; Post-Stim: $M = 0.149$ +/- 0.028), suggesting that participants generally improved during the course of the experiment.

The interactions showed that there was an additional effect caused by the type of stimulation. We found a significant interaction between polarity and session (F $(2,16) = 5.216$, $p = .018$, $\eta^2 = .395$) and between target location and polarity (F $(1,8) = 6.514$, $p = .034$, $\eta^2 = .449$). While the curvature was slightly higher for the anodal stimulation blocks during the baseline, there was a reduction in curvature during anodal stimulation, which did not occur with cathodal

stimulation (see Figure 4 A). We investigated this effect further by running two repeated measures ANOVAs only with the factor Session for the anodal and the cathodal values, respectively. While we found a significant reduction over the sessions for the anodal stimulation (F (2,16) = 8.002, p = .004, η^2 = .500), there was no reduction of the curvature for the cathodal stimulation (F (2,16) = 0.474, p = .631, η^2 = .056). The interaction between target location and polarity was based on a higher curvature for cathodal than for anodal stimulation for rightward (contralateral) targets (see Figure 4 B). There is a marginal reduction of the curvature for anodal stimulation for targets corrected to the right in comparison to the left (t (8) = 2.16, p = .063), but no difference depending on the direction with cathodal stimulation (t (8) = 0.16, p = .878). Thus, taken together there seems to be a reduction of curvature with anodal stimulation, whereas an increase in curvature with cathodal stimulation. This effect was most prominent for targets jumping to the right, i.e., the contralateral side of the stimulation.

Figure 4. Effects on the curvature of the trajectories. (**A**) Curvature of the hand movements averaged across participants and shown separated for the different sessions and polarities (circle = anodal stimulation, triangle = cathodal). (**B**) Average curvature across participants for targets jumping to the left or right for both polarities. Error bars depict the standard error of the mean.

To investigate the integration of the new target location into the movement we also examined the development of curvature over time (see Figure 5 A). We aligned all trials to the calculated onset of the online correction (see Methods) and calculated the curvature with a sliding window of 100 ms in steps of 10 ms. Interestingly, the latency of the online correction also decreased across the different sessions (F (2,16) = 10.093, p = .001, η^2 = .558: Base: M =

291.70 +/-63.42 ms; Stim: M = 280.43+/-64.54 ms; Post-Stim: M = 274.13+/-57.55 ms), but it was not affected by the polarity of the stimulation ($F(1,8) = 2.539$, $p = .150$, $\eta^2 = .241$). To quantify the possible effect of the more effective integration of the updated movement plan, we normalized the curvature over time for each participant based on their peak of the time course in the baseline condition and computed the difference between the peak curvature for the stimulation session and the baseline value. We did so for each of the polarities as well as for targets both jumping to the right and left (both polarities are shown in Figure 5 A, but note that only trials jumping to the right are shown). We found a significant difference between the anodal and cathodal stimulation only during the stimulation for targets jumping to the right (t (8) = 3.725, $p = .006$), which was not present on the left side (t (8) = 1.37, $p = .21$). The fact that the contralateral side is significant but the ipsilateral is not, is not enough to show that both sides are actually different. Thus, we compared the difference between anodal and cathodal stimulation on both sides directly and again obtained a significant result ($t(8) = 2.687$, $p = .028$), indicating a hemifield specific effect of the stimulation. Furthermore, anodal stimulation led to a reduction in curvature, whereas cathodal stimulation led to an increase. Thus, in contrast to the latency of the online control system, we found a significant effect of the stimulation on the curvature of the reaching trajectories. Anodal stimulation seems to lead to a more effective updating and correction of the ongoing movement for targets that jumped into the contralateral field from the stimulation site, whereas the opposite was the case for cathodal stimulation.

Figure 5. Development of curvature over time. (**A**) shows the average curvature over time aligned to online correction onset for the different sessions (Baseline = black, Stimulation = red, Post-Stim = blue). The left panel shows the curvature for targets jumping to the right during the sessions with anodal stimulation, the right panel with cathodal stimulation. Curvature was normalized so that the maximum of the baseline was assigned a 1. The shaded area shows the standard error of the mean. (**B**) depicts the average differences between the peak curvature during baseline and the stimulation. Circles show the effect of anodal stimulation, triangles of cathodal stimulation. Error bars depict the standard error of the mean. The star indicates significance at the 1% level.

Causal relation between induced current and behavioral changes

Thus far our results indicate a relation between the polarity of the stimulation that we induced and participants' online correction behavior. To test for a causal relationship, we next sought to provide evidence that the amount of current delivered to the mIPS, quantified based on an individual model of current flow for each participant (see current forward modeling in methods), is directly linked to their behavior and that this relationship is mediated by a change in brain activity. To establish such a relationship our data need to fulfill the following three criteria: (1) There needs to be a correlation between the induced current and the change in behavior. (2) There needs to be a relationship between the induced current and changes in brain activity, quantified in our case via EEG measurement before and after the stimulation. (3) The altered brain activity needs to be related to the changes in behavior and crucially this relationship must be the determining factor for the correlation between the induced current and the behavior.

(1) Relationship between induced current and behavior. To test this relationship, we first quantified the induced current. We estimated the current that affected the left mIPS based on the electrode placement and the structural MRI scan of each participant (see current forward modeling in methods). The current was decomposed in its radial and tangential component.

By design, anodal stimulation led to positive values, whereas cathodal stimulation led to
negative values. We correlated the reconstructed current vectors with the effect of the
stimulation on curvature and online curvature (Stimulation-Baseline for both measures).
While there was no systematic relationship between the behavior and the radial component of
the current (see Fig 6A), we observed a significant relationship between the tangential
component of the current reconstruction and the curvature measurements for targets jumping
to the right (see Figure 6B for online curvature). Importantly, this relationship was again only
present on the contralateral side of the stimulation, thus for targets jumping to the right.

Figure 6. Relationship between induced current and behavior. **(A)** Relationship between the
radial component of the induced current and the difference in peak reach curvature. Targets that
jumped to the left are depicted by the black triangles pointing to the left. Targets jumping to the
right are depicted by the open triangles pointing to the right. The solid line shows a regression
for the targets jumping to the left, the dotted line shows the regression to the data jumping to the
right. **(B)** Similar as A, but here the relationship with the tangential current is plotted. The red
data points show the significant relationship between the tangential current and change in peak
curvature for targets jumping to the right.

(2) Relationship between current and EEG. We now have established the correlation
between the induced current and the behavioral changes. To investigate whether the current
also changed the brain activity, we analyzed the EEG response for trials that jumped either to
the left or to the right. We measured the EEG in the baseline and the post-stimulation session,
which allowed us to compute the effect of the stimulation on the brain activity. Recall that,
during the stimulation, the sensors are being used for stimulation and thus cannot be used to
record EEG activity. We computed TFR response for each session and direction of target
jump separately and then computed the difference between the post-stimulation and baseline.

Figure 7 shows the difference in power across the different frequencies for anodal and cathodal stimulation respectively for targets jumping to the right. By looking at the figure it seems that anodal stimulation is slightly decreasing the power of alpha and beta frequencies shortly after the target jump, whereas cathodal stimulation seems to increase the power in a similar range (compare Figure 7 A+B). To relate these changes with the induced current we wanted to establish areas in the TFR that were related to the induced current. Similar to the relation of the behavior, we found that the tangential current was related to the TFR responses for targets jumping to the right. Significant correlations were found quite localized at 11-14 Hz (alpha) in the first 100 ms after target jump and 20-26 Hz (beta) between 100 and 200 ms after target jump (see Figure 7C). As a representation of the observed correlation and in order to quantify the effect of the stimulation we extracted the average TFR response for targets jumping to the right for each participant in these time windows (see Figure 7D). For these time windows we found a significant difference between the effect of anodal and cathodal stimulation for targets jumping to the right ($t(7) = 3.23$, $p = .015$). Anodal stimulation led to a reduction of the power ($M = -0.42$, $SD = 0.59$), while cathodal stimulation increased the power ($M = 0.22$, $SD = 0.28$). This was true for the alpha as well as the beta activity, as the change in power in those two time windows was highly correlated ($r(16) = .81$, $p < .001$). If we computed the same difference for targets to the left we observed no significant correlation with the current and also no difference in the TFR representation ($t(7) = 0.44$, $p = .67$). Thus, we can establish a systematic influence of the induced current on the TFR response for targets jumping to the contralateral side of the stimulation. Note that this analysis was irrespective of any behavioral changes and only considered the lateralization of the brain with respect to coding of target jump direction.

Figure 7. Relationship between induced current and EEG response. (**A+B**) Differences between Post-Stimulation session and Baseline in the TFR response aligned to the moment of the target jump. (**A**) shows the effect of anodal stimulation and (**B**) the effect of cathodal stimulation, both for targets jumping to the right. Blueish colors depict reduction of the power, reddish color increase in power. (**C**) depicts significant parts of a point-by-point correlation between the TFR response for targets jumping to the right and the tangential current. The p-value of these correlation is color-coded and to improve visibility only significant parts are shown. (**D**) Correlation between the average TFR values in the significant portions and the induced tangential current. Targets that jumped to the left are depicted by the black triangles pointing to the left. The significant relationship between tangential current and changes in the TFR for targets jumping to the right are depicted by the red open triangles pointing to the right. The solid line shows the regression for the targets jumping to the left, the dotted line shows the regression for the targets jumping to the right.

(3) Mediation of the relationship between current and behavior via brain activity. Based on a causal interpretation of the effects, the induced current influences the brain response and this changed brain response should then affect the behavior. We now ask whether the independently established changes in EEG serve as a mediator for the relationship between current and behavior.

We tested for a relationship between the changes in EEG and behavior and observed a
significant relationship between the changes in the TFR and curvature (r(16) = .62, p = .01;
see Figure 8) as well as the curvature over time for targets jumping to the right (r(16) = .73, p
=.001). Crucially, when we controlled for the changes in the TFR response by controlling for
these changes in a partial correlation between the tangential current with the behavior, the
correlations were significantly decreased (see Figure 8 for curvature; r (16) = -.54, p = .035
for online curvature).

Thus, when taken together we found a polarity-specific modulation of the online correction
for targets jumping to the contralateral side of the stimulation. Anodal stimulation slightly
improved the correction, reflected in a lower curvature, whereas cathodal stimulation
produced the opposite effect. We found a significant relationship between the tangential
current and the observed behavioral changes and could successfully establish the changes in
the EEG response as a mediator of this effect. These results suggest that the induced current
adjusted the early alpha and beta oscillations after the target jump and these changes then
affected the online correction performance.

Figure 8. Mediation analysis. Relationship between the induced tangential current, the
changes in the TFR response of the EEG and the change in curvature in the behavioral response.
Solid arrows depict significant correlations. The partial correlation against the EEG response is
reported beneath the relationship of induced current and behavior.

Discussion

The goal of our study was to establish a causal role of the PPC for online reach control. We were able to show not only the effects on behavior, but more importantly, reconstruct the causal structure of the stimulation with a mediation analysis. This was achieved by using a unique set of measurements starting from a model of current flow for each individual participant, pre- and post stimulation EEG recordings, and the analysis of behavioral changes. Our results provide interesting insights into the role of the mIPS for online control while establishing a new methodological approach to provide causality and explain some of the variability of tDCS stimulation.

Online control behavior

Our results fit well into the literature on behavioral effects on online movement control (Gaveau et al., 2014) and optic ataxia (Rossetti et al., 2003; Anderson et al. 2015). Lesions in the PPC lead to severe impairments of online corrections (Pisella et al.,2000; Grea et al., 2002; Buiatti et al., 2013). Similarly, TMS stimulation over the PPC impedes online reach control (Desmurget et al., 1999), while keeping reaches to static movements intact (Desmurget et al., 1999; Grea et al., 2002). Here, we show that anodal stimulation, related to an increase in activity, leads to improved online correction performance, while cathodal stimulation, related to a decrease in activity, did not (Figure 4). Importantly, this effect was observed contralateral to the hemisphere of stimulation (Figure 5), providing evidence for the lateral specificity of stimulation. We did not observe any systematic changes in some more general behavioral measurements such as accuracy or precision. This finding rules out the possibility that our observed effects on online control are based on a change in the representation of the target position, suggesting that the stimulation indeed mainly affected the efficiency of the online correction and not the endpoint of the movement. It is important to note here that participants were allowed to freely move their eyes, so they performed reach-to-point movements towards targets in the fovea: reaches to targets in the periphery are increasing after lesions of the intraparietal sulcus (Hwang et al., 2012).

The exact behavioral and neural mechanisms of how movements can be adjusted in flight are still unknown, however recent optimal feedback control models provide a potential framework for understanding how the brain might achieve this (Todorov & Martin, 2002; Scott, 2004; Scott 2012). These models propose that after establishing a behavior goal, the

state of the body is estimated based on a combination of efferent and delayed afferent information and then the motor action is implemented by a control policy which will continuously control the movement (Scott 2016; Heed et al., 2016; Scott, 2017). One key advantage of the optimal control theory framework is that it can explain why despite some variability in the individual trajectories (see Figure 3), the endpoints of a movement are still comparable and accurate.

To probe the online correction system studies typically use two different manipulations: perturbations of the ongoing movement to change the estimation of the hand position or perturbations of the target position to change the estimate of the task goal (for reviews see Scott, 2012; 2016). Participants are faster in reacting to a mechanical perturbation than to a change in the visual target location (Dimitriou et al. 2013), suggesting that there are possible differences in the system when changing either the state estimate or redefining the task goal. In line with this interpretation, the faster but less reliable somatosensory information is mainly used to estimate the state of the body even when visual information is available (Crevecoeur, Munoz & Scott, 2016). In contrast, a target step like in our experiment can only be signaled by visual information, requiring an updating of the reach goal. Based on the long latency (~ 250-300 ms) of the online correction in our task, the large target step in our case seems to reflect a higher-level corrective process redefining the task (Scott, 2016) instead of an already present internal representation of different motor plans (Nashed, Crevecour, Scott, 2014).

The PPC could serve as a state estimator, due to its multiple afferent connections and role in sensorimotor integration (Sirigu et al., 1996; Mulliken et al., 2008; Pilacinski et al., 2019; Heed & Medendorp, 2019). However, although our results provide additional evidence for the causal role of the PPC in online movement control, it is not a uniform cortical area and different parts of the PPC might fulfill different subtasks (Heed & Medendorp, 2019). A recent study by Marigold et al. (2019) used TMS over four different areas of the PPC, which were all shifted laterally to the IPS, and did not observe any effects on online control. The authors suggested that these areas might be more involved in reach planning instead of online control. With respect to our stimulation site, the mIPS, it seems that it can be functionally separated from the more anterior part of the IPS (Reichenbach et al. 2014). TMS stimulation of the aIPS was shown to disrupt grip size, while keeping reaching intact (Tunik et al., 2005) and seems to be more related to integrating visual information used to guide a movement (Reichenbach et al., 2014). In contrast, the mIPS seems to be more involved in integrating the incoming somatosensory information like proprioceptive feedback (Reichenbach et al., 2014).

In our study no visual information about the position of the hand was present, making the
corrections only possible by comparing an internal estimate of the hand position with the
updated target location.

Thus, the integration of visual versus somatosensory information is mirrored between the
behavioral effects and the different brain areas. There seems to be a functional gradient
implemented in the PPC: from the estimates of the environment, including possible changes
in task goal, in its rostral pole, to the state estimate of the body, in its caudal pole (Heed &
Medendorp, 2019). Our results show that stimulation of the mIPS, which is related to the
integration of the incoming somatosensory information (Reichenbach et al., 2014), possibly
changes the internal state estimate which is then reflected in changes in the efficiency of
online control.

Brain oscillations as communication mechanism

It is important to mention here, that although the PPC is causally involved in online
movement control, it is not the only relevant area. Active perturbations of the hand activate a
broad network ranging from the somatosensory cortex, primary motor cortex, dorsal premotor
cortex (PMd) to more parietal areas. Changes in firing rates in the somatosensory and motor
cortex are observed as early as 25 ms after the perturbation (Omrani et al. 2016). Besides the
PPC, the dorsal premotor cortex (PMd) plays a crucial role in updating target location (see
Battaglia-Mayer et al. 2014; Archambault 2015). It provides a dynamic error signal indicating
the new goal (Lee & van Donkelaar, 2005), which then is passed to the PPC (Archambault et
al., 2015). Interestingly, a lesion in the PMd mainly affects the latency of the movement, but
not its accuracy (Buiatti et al. 2013), which suggests that the movement correction is
computed in the PPC (Battaglia-Mayer et al. 2014).

In order to adjust ongoing movements there are two crucial steps needed. First one needs to
inhibit the old goal and afterwards integrate the state estimate of the hand with the new goal.
These two steps seem to be related to specific frequencies of brain oscillations. The initial
inhibition arriving from the PMd is presumably related to alpha oscillations (Pani et al.,
2014), while the integration of the new goal and preparation of the movement may be related
to beta frequencies (Pani et al., 2014; Stetson & Anderson, 2014; Engel & Fries, 2010). In
line with previous studies, we observed changes in these two frequency bands.

Our results show that stimulation with tDCS influenced the power of alpha and beta
frequency bands. Anodal stimulation significantly reduced alpha and beta power (Figure 7),
suggesting that a more efficient online correction is based on a combination of less inhibition
of the old goal and a better integration of the state estimate and the new goal. Cathodal
stimulation led to an increase in power in both frequency bands, suggesting a stronger
inhibition of the previous movement goal with less integration of the state estimate with the
new movement goal, leading to an even higher curvature than without stimulation.

Remarkably, we could reconstruct the two relevant frequency bands based on a correlation
between our two independent measurements of changes in the brain activity, without taking
into account the behavioral changes (Figure 7C). The correlation with the induced current also
provided the same temporal structure: an early alpha response after the target jump followed
by a slightly later beta activity. However, it is important to note, that despite their potential
different functions, alpha and beta frequencies seems to be coupled (Grabot et al., 2019) as
changes in their respective power were highly correlated in our study.

After the decision to adjust the ongoing movement there are two possibilities for how this can
be achieved computationally: Either one simply replaces the old movement by the updated
solution, or the old and new movement goals exist in parallel and there is continuous change
from the old towards the new movement. Neuronal recordings in the PPC suggest that the two
possibilities are combined. A third of the neurons behave in line with the replacing
hypothesis, while the rest of the neurons behave according to the superposition hypothesis,
(Dickey et al., 2013). Therefore, online corrections lead to a decrease in the correlation
between neural responses and behavior due to overlap of separate motor plans across the PPC,
as well as the PMd (Georgopulos, 1983; Archambault et al., 2009; 2011; 2015; Dickey et al.,
2013). Thus, the influence of tDCS could either act on the reliability of the state estimate,
which influences the integration of the state with the updated movement goal, or the
stimulation could change the balance between these two sets of neurons; both effects could
lead to a smoother online correction.

Caveats of results and strengths of our multimodal approach

While our study provides important insights into mechanisms behind online control during
reaching and provides a new methodology and experimental design for tDCS-stimulation
studies, there are certain caveats with respect to our results. Stimulation studies typically rely
on the assumption that behavior won't change in the absence of stimulation, but, trial-by-trial

effects and learning in behavioral paradigms are the norm, not the exception (Fecteau et al, 2003). To statistically account for learning, manipulations of neural mechanisms should thus take into account the potential for ongoing changes in behavior. Our baseline/stimulation/post design allowed us to assess the improvement over time as well as the bidirectional effect of tDCS depending on the polarity (see Figure 4). But as participants improved in the task across the different sessions, it could be the case that the learning effect, in addition to a lack of statistical power, could hide additional effects, for example an influence on the latency of online correction. Based on the learning effect one possibility is that our results could also be interpreted based on an increase or impairment of trial-by-trial learning, while anodal stimulation increases the learning and therefore leads to smoother corrections. To rule that explanation out, in future work one could add an additional sham condition to measure the change in behavior without any stimulation. However, based on the fact that people got more comfortable with the task with continuous practice throughout the task and the fact that we did not observe any polarity specific effect on other parameters like the accuracy or latency of online correction, we propose that there was a direct influence of the tDCS on the online control during reaching.

While the EEG recordings provide good temporal resolution, its spatial resolution is limited and cannot directly show the effect of tDCS on the neuronal level. However, EEG is a non-invasive method that can image the relationship between ongoing neural oscillations and behavior. Importantly, given that evoked cortical potentials from repetitive sensory and magnetic stimulation are subject to potentiation (Teyler et al, 2005; Esser et al, 2006; Clapp et al, 2012; Ferreri et al, 2013), it follows that the EEG can index long-term changes to neural activity induced by tDCS.

Delivering current to humans noninvasively has been met with significant methodological challenges; much of the current delivered transcranially is shunted by the scalp, and the rest spreads diffusely across the cortex (Vöröslakos et al, 2018). Recent attempts to model the induced electric field has provided the opportunity to design electrode montages to increase the specificity of the induced current; like the HD-tDCS stimulation that we used in our experiment (Dmochowski et al, 2011; Saturnio et al, 2015, 2018). Additionally, the current flow depends on the individual anatomy (Miranda et al., 2006), which may explain some of the variability and conflicting results in tDCS studies (Jacobsen et al., 2012; Hill et al., 2016; Kang et al., 2016).

Despite these caveats, our experimental design contrasting anodal and cathodal stimulation
combined with our multimodal approach (event-related EEG), fMRI-guided localization of
stimulation target, and individualized current modeling work allow us to account for most of
the potential problems. Specifically, modeled current source strength correlations with
stimulation-induced EEG changes produced an independent subject-by-subject predictor for
expected tDCS-induced behavioral changes. Although there might be potential differences in
individual values for tissue and bone conductance (McCann et al., 2019), the reconstruction of
the current flow for each individual participant allows us to model and account for the
individual variability of the induced current across participants. By independently measuring
the change in brain activity via EEG, we could validate our current reconstructions due to the
relationship between the induced current and changes in relevant frequency bands in the EEG.
Thus, our study design together with the applied mediation analysis can overcome some of
the caveats of the individual measurements and has one big advantage: it is possible to
establish the causal structure of the stimulation while at the same time gain insight into
potential mechanisms due to the continuous change in brain activity and behavior, which is
not possible with real or virtual lesions who simply disrupt the activity. It also allows to
integrate and make use of the variability that is typically found in the results of tDCS studies
(Jacobsen et al., 2012) by the use of a regression-based mediation analysis, as a step to make
tDCS a more useful tool for research and clinical applications.

Establishing causality for functional reorganization

One goal in neuroscience is to uncover how specific areas of the brain mediate specific
functions. This understanding requires causal intervention (Marinescu & Kording, 2018), and
is built on the methods with which we can change the activity of neurons, such as stimulation
or lesions (Fellows et al 2005). This approach permits to identify clinically relevant regions of
the brain that lead to behavioral deficits when damaged. However, information in the brain is
at least partially redundant (Price & Friston, 2002; Cisek, 2012; Pitkow & Angelaki, 2017),
functionality is shared across different behaviors (Dehaene, 2016; Krakauer et al, 2017),
interventions can cause indirect effects in downstream circuits (Kumar et al, 2013; Otchy,
2015), and there is at least some degree of functional reorganization following damage
(O'Shea et al, 2007). Thus, intervention studies necessarily provide information about how
the brain responds to a change, but it may or may not answer a causal question about the
function of a brain region in isolation (Young et al, 2000).

By combining current reconstructions, EEG analysis and behavioral changes, we provided a new methodological approach to address causal questions about how the brain responds to HD-tDCS over time. We used this methodology to gain evidence for the causal role of an externally applied electric field centered around mIPS (stimulation period) on future EEG activity and reaching behavior during the stimulation. Since the EEG recorded at the stimulation site reflects a spatial summation of neural activity in the brain, and not just that of mIPS, our results implicate the underlying network with which it participates. Future work could leverage our causal methodologies to probe how multiple brain regions interact following brain stimulation, and the combination of our measurements can be adapted to almost any behavioral task and stimulation of different areas. This is an important step towards translating knowledge of the adaptive brain to our understanding of functional localization of behavior.

Acknowledgements

AG received funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—project number 222641018—SFB/TRR 135 Project A1, and by the DFG International Research Training Group 1901. **BC**, **SX**, **JJ** & **GB** were supported by a NSERC Discovery Grant (Canada), a NSERC CREATE International Research and Training grant (Canada), and the Canadian Foundation for Innovation (CFI). **JG** received funding from NSERC Discovery Grant (Canada) and CFI award. **JD** received funding from the Army Research Laboratory (W911-NF-10-2-0022) and the National Institutes of Health (NIH–NIDA K18DA045437), USA. **KF** received funding from the Deutsche Forschungs-gemeinschaft (DFG, German Research Foundation)—project number 222641018—SFB/TRR 135 Project A4, and by the DFG International Research Training Group IRTG 1901.

Author contributions

AG Conception and Design of experiment, Performed Experiments, Analyzed behavioral and EEG data and performed the mediation analysis, Interpreted the results of the experiments, Prepared figures, Drafted manuscript, Edited and revised manuscript, Approved final version of the experiment.

BC Performed current reconstructions and EEG data analysis, Interpreted the results of the experiment, Prepared figures, Drafted manuscript, Edited and revised manuscript, Approved final version of the manuscript.

JJ Performed current reconstructions and EEG data analysis, Edited and revised manuscript, Approved final version of the experiment.

SX Performed experiments, Analyzed fMRI data, Prepared figures, Drafted parts of the method section, Edited and revised manuscript, Approved final version of the experiment.

JG Designed fMRI task and fMRI analysis, Edited and revised manuscript, Approved final
version of the experiment.
JD Designed current reconstruction pipeline, Edited and revised manuscript, Approved final
version of the experiment.
KF Conception and Design of experiment, Interpreted the results of the experiments, Edited
and revised manuscript, Approved final version of the experiment.
GB Conception and Design of research, conception and design of analysis methods,
Interpreted the results of the experiment, Edited and revised manuscript, Approved final
version of the manuscript.
862

References

Andersen, R. A., Andersen, K. N., Hwang, E. J., & Hauschild, M. (2014). Optic ataxia: from
Balint's syndrome to the parietal reach region. *Neuron*, *81*(5), 967-983.
Archambault, P. S., Caminiti, R., & Battaglia-Mayer, A. (2009). Cortical mechanisms for
online control of hand movement trajectory: the role of the posterior parietal cortex.
Cerebral Cortex, *19*(12), 2848-2864.
Archambault, P. S., Ferrari-Toniolo, S., & Battaglia-Mayer, A. (2011). Online control of hand
trajectory and evolution of motor intention in the parietofrontal system. *Journal of
Neuroscience*, *31*(2), 742-752.
Archambault, P. S., Ferrari-Toniolo, S., Caminiti, R., & Battaglia-Mayer, A. (2015). Visually-
guided correction of hand reaching movements: the neurophysiological bases in the
cerebral cortex. *Vision research*, *110*, 244-256.
Bączyk, M., & Jankowska, E. (2014). Presynaptic actions of transcranial and local direct
current stimulation in the red nucleus. *The Journal of physiology*, *592*(19), 4313-4328.
Baron, R. M., & Kenny, D. A. (1986). The moderator–mediator variable distinction in social
psychological research: Conceptual, strategic, and statistical considerations. *Journal of
personality and social psychology*, *51*(6), 1173.
Battaglia-Mayer, A., Buiatti, T., Caminiti, R., Ferraina, S., Lacquaniti, F., & Shallice, T.
(2014). Correction and suppression of reaching movements in the cerebral cortex:
physiological and neuropsychological aspects. *Neuroscience & Biobehavioral Reviews*,
42, 232-251.
Battaglia-Mayer, A., Ferrari-Toniolo, S., & Visco-Comandini, F. (2015). Timing and
communication of parietal cortex for visuomotor control. *Current opinion in
neurobiology*, *33*, 103-109.
Beurze, S. M., De Lange, F. P., Toni, I., & Medendorp, W. P. (2007). Integration of target and
effector information in the human brain during reach planning. *Journal of
neurophysiology*, *97*(1), 188-199.
Buiatti, T., Skrap, M., & Shallice, T. (2013). Reaching a moveable visual target: Dissociations
in brain tumour patients. *Brain and cognition*, *82*(1), 6-17.
Buzsáki, G., Anastassiou, C. A., & Koch, C. (2012). The origin of extracellular fields and
currents—EEG, ECoG, LFP and spikes. *Nature reviews neuroscience*, *13*(6), 407.
Cavina-Pratesi, C., Connolly, J. D., & Milner, A. D. (2013). Optic ataxia as a model to
investigate the role of the posterior parietal cortex in visually guided action: evidence from
studies of patient MH. *Frontiers in human neuroscience*, *7*, 336.

Clapp, W. C., Hamm, J. P., Kirk, I. J., & Teyler, T. J. (2012). Translating long-term
potentiation from animals to humans: a novel method for noninvasive assessment of
cortical plasticity. *Biological psychiatry*, *71*(6), 496-502.
Cisek, P. (2012). Making decisions through a distributed consensus. *Current opinion in
neurobiology*, *22*(6), 927-936.
Crevecoeur, F., Munoz, D. P., & Scott, S. H. (2016). Dynamic multisensory integration:
somatosensory speed trumps visual accuracy during feedback control. *Journal of
Neuroscience*, *36*(33), 8598-8611.
Datta, A., Bansal, V., Diaz, J., Patel, J., Reato, D., & Bikson, M. (2009). Gyri-precise head
model of transcranial direct current stimulation: improved spatial focality using a ring
electrode versus conventional rectangular pad. *Brain stimulation*, *2*(4), 201-207.
Dehaene, S., & Cohen, L. (2007). Cultural recycling of cortical maps. *Neuron*, *56*(2), 384-
398.
Desmurget, M., Epstein, C. M., Turner, R. S., Prablanc, C., Alexander, G. E., & Grafton, S. T.
(1999). Role of the posterior parietal cortex in updating reaching movements to a visual
target. *Nature neuroscience*, *2*(6), 563.
Desmurget, M., Gréa, H., Grethe, J. S., Prablanc, C., Alexander, G. E., & Grafton, S. T.
(2001). Functional anatomy of nonvisual feedback loops during reaching: a positron
emission tomography study. *Journal of Neuroscience*, *21*(8), 2919-2928.
Dickey, A. S., Amit, Y., & Hatsopoulos, N. G. (2013). Heterogeneous neural coding of
corrective movements in motor cortex. *Frontiers in neural circuits*, *7*, 51.
Dmochowski, J. P., Datta, A., Bikson, M., Su, Y., & Parra, L. C. (2011). Optimized multi-
electrode stimulation increases focality and intensity at target. *Journal of neural
engineering*, *8*(4), 046011.
Engel, A. K., & Fries, P. (2010). Beta-band oscillations—signalling the status quo?. Current
opinion in neurobiology, 20(2), 156-165.
Esser, S. K., Huber, R., Massimini, M., Peterson, M. J., Ferrarelli, F., & Tononi, G. (2006). A
direct demonstration of cortical LTP in humans: a combined TMS/EEG study. Brain
research bulletin, *69*(1), 86-94.
Faisal, A. A., Selen, L. P., & Wolpert, D. M. (2008). Noise in the nervous system. *Nature
reviews neuroscience*, *9*(4), 292.
Fecteau, J. H., & Munoz, D. P. (2003). Exploring the consequences of the previous trial.
Nature Reviews Neuroscience, *4*(6), 435.
Fellows, L. K., Heberlein, A. S., Morales, D. A., Shivde, G., Waller, S., & Wu, D. H. (2005).
Method matters: an empirical study of impact in cognitive neuroscience. *Journal of
cognitive neuroscience*, *17*(6), 850-858.
Ferreri, F., & Rossini, P. M. (2013). TMS and TMS-EEG techniques in the study of the
excitability, connectivity, and plasticity of the human motor cortex. *Reviews in the
Neurosciences*, *24*(4), 431-442.
Gallivan, J. P., McLean, D. A., Smith, F. W., & Culham, J. C. (2011). Decoding effector-
dependent and effector-independent movement intentions from human parieto-frontal
brain activity. *Journal of Neuroscience*, *31*(47), 17149-17168.
Gaveau, V., Pisella, L., Priot, A. E., Fukui, T., Rossetti, Y., Pélisson, D., & Prablanc, C.
(2014). Automatic online control of motor adjustments in reaching and grasping.
Neuropsychologia, *55*, 25-40.
Georgopoulos, A. P., Kalaska, J. F., Caminiti, R., & Massey, J. T. (1983). Interruption of
motor cortical discharge subserving aimed arm movements. *Experimental Brain Research*,
49(3), 327-340.
Gertz, H., & Fiehler, K. (2015). Human posterior parietal cortex encodes the movement goal
in a pro-/anti-reach task. *Journal of neurophysiology*, *114*(1), 170-183.

Goettker, A., Caie, B., Jeyachandra, J., Xu, S., Gallivan, J., Dmochowski, J., … Blohm, G. (2019, July 30). HD-tDCS over mIPS causally modulates online reach correction. Retrieved from osf.io/kbyxc

Grabot, L., Kononowicz, T. W., La Tour, T. D., Gramfort, A., Doyère, V., & Van Wassenhove, V. (2019). The Strength of Alpha–Beta Oscillatory Coupling Predicts Motor Timing Precision. *Journal of Neuroscience*, *39*(17), 3277-3291.

Gréa, H., Pisella, L., Rossetti, Y., Desmurget, M., Tilikete, C., Grafton, S., ... & Vighetto, A. (2002). A lesion of the posterior parietal cortex disrupts on-line adjustments during aiming movements. *Neuropsychologia*, *40*(13), 2471-2480.

Geuzaine, C., & Remacle, J. F. (2009). Gmsh: A 3-D finite element mesh generator with built-in pre-and post-processing facilities. *International journal for numerical methods in engineering*, *79*(11), 1309-1331.

Harris, C. M., & Wolpert, D. M. (1998). Signal-dependent noise determines motor planning. *Nature*, *394*(6695), 780.

Hill, A. T., Fitzgerald, P. B., & Hoy, K. E. (2016). Effects of anodal transcranial direct current stimulation on working memory: a systematic review and meta-analysis of findings from healthy and neuropsychiatric populations. *Brain stimulation*, *9*(2), 197-208.

Hwang, E. J., Hauschild, M., Wilke, M., & Andersen, R. A. (2012). Inactivation of the parietal reach region causes optic ataxia, impairing reaches but not saccades. *Neuron*, *76*(5), 1021-1029.

Jacobson, L., Koslowsky, M., & Lavidor, M. (2012). tDCS polarity effects in motor and cognitive domains: a meta-analytical review. *Experimental brain research*, *216*(1), 1-10.

Jobst, C., Ferrari, P., Isabella, S., & Cheyne, D. (2018). BrainWave: A Matlab Toolbox for Beamformer Source Analysis of MEG Data. *Frontiers in neuroscience*, *12*.

Jonas, E., & Kording, K. P. (2017). Could a neuroscientist understand a microprocessor?. *PLoS computational biology*, *13*(1), e1005268.

Kang, N., Summers, J. J., & Cauraugh, J. H. (2016). Transcranial direct current stimulation facilitates motor learning post-stroke: a systematic review and meta-analysis. *J Neurol Neurosurg Psychiatry*, *87*(4), 345-355.

Kirby, M. (1972). On acting and not-acting. *The drama review: TDR*, 3-15.

Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A., & Poeppel, D. (2017). Neuroscience needs behavior: correcting a reductionist bias. *Neuron*, *93*(3), 480-490.

Kumar, A., Vlachos, I., Aertsen, A., & Boucsein, C. (2013). Challenges of understanding brain function by selective modulation of neuronal subpopulations. *Trends in neurosciences*, *36*(10), 579-586.

Lauro, L. J. R., Rosanova, M., Mattavelli, G., Convento, S., Pisoni, A., Opitz, A., ... & Vallar, G. (2014). TDCS increases cortical excitability: direct evidence from TMS–EEG. *Cortex*, *58*, 99-111.

Limanowski, J., Kirilina, E., & Blankenburg, F. (2017). Neuronal correlates of continuous manual tracking under varying visual movement feedback in a virtual reality environment. *Neuroimage*, *146*, 81-89.

Liu, A., Vöröslakos, M., Kronberg, G., Henin, S., Krause, M. R., Huang, Y., ... & Berényi, A. (2018). Immediate neurophysiological effects of transcranial electrical stimulation. *Nature communications*, *9*(1), 5092.

MacKinnon, D. P., Fairchild, A. J., & Fritz, M. S. (2007). Mediation analysis. *Annu. Rev. Psychol.*, *58*, 593-614.

Mangia, A. L., Pirini, M., & Cappello, A. (2014). Transcranial direct current stimulation and power spectral parameters: a tDCS/EEG co-registration study. *Frontiers in human neuroscience*, *8*, 601.

Marigold, D. S., Lajoie, K., & Heed, T. (2019). No effect of triple-pulse TMS medial to intraparietal sulcus on online correction for target perturbations during goal-directed hand and foot reaches. *bioRxiv*, 645853.

McCann, H., Pisano, G., & Beltrachini, L. (2019). Variation in reported human head tissue electrical conductivity values. *bioRxiv*, 511006.

Medendorp, W. P., & Heed, T. (2019, April 28). State Estimation in Posterior Parietal Cortex: Distinct Poles of Environmental and Bodily States. https://doi.org/10.31219/osf.io/wbcru

Menzer, D. L., Rao, N. G., Bondy, A., Truccolo, W., & Donoghue, J. P. (2014). Population interactions between parietal and primary motor cortices during reach. *American Journal of Physiology-Heart and Circulatory Physiology*.

Miall, R. C., & Wolpert, D. M. (1996). Forward models for physiological motor control. *Neural networks*, *9*(8), 1265-1279.

Miranda, P. C., Lomarev, M., & Hallett, M. (2006). Modeling the current distribution during transcranial direct current stimulation. *Clinical neurophysiology*, *117*(7), 1623-1629.

Ogawa, K., Inui, T., & Sugio, T. (2006). Separating brain regions involved in internally guided and visual feedback control of moving effectors: an event-related fMRI study. *Neuroimage*, *32*(4), 1760-1770.

Omrani, M., Murnaghan, C. D., Pruszynski, J. A., & Scott, S. H. (2016). Distributed task-specific processing of somatosensory feedback for voluntary motor control. *Elife*, *5*, e13141.

Otchy, T. M., Wolff, S. B., Rhee, J. Y., Pehlevan, C., Kawai, R., Kempf, A., ... & Ölveczky, B. P. (2015). Acute off-target effects of neural circuit manipulations. *Nature*, *528*(7582), 358.

O'Shea, J., Johansen-Berg, H., Trief, D., Göbel, S., & Rushworth, M. F. (2007). Functionally specific reorganization in human premotor cortex. *Neuron*, *54*(3), 479-490.

Pani, P., Di Bello, F., Brunamonti, E., D'Andrea, V., Papazachariadis, O., & Ferraina, S. (2014). Alpha-and beta-band oscillations subserve different processes in reactive control of limb movements. *Frontiers in behavioral neuroscience*, *8*, 383.

Pisella, L., Grea H, Tilikete C, Vighetto A, Desmurget M, Rode G, Boisson D, & Rossetti Y. (2000) An "automatic pilot" for the hand in human posterior parietal cortex: toward reinterpreting optic ataxia. *Nat Neurosci*, *3*, 729-736.

Pélisson, D., Prablanc, C., Goodale, M. A., & Jeannerod, M. (1986). Visual control of reaching movements without vision of the limb. *Experimental Brain Research*, *62*(2), 303-311.

Pitkow, X., & Angelaki, D. E. (2017). Inference in the brain: statistics flowing in redundant population codes. *Neuron*, *94*(5), 943-953.

Prablanc, C., & Martin, O. (1992). Automatic control during hand reaching at undetected two-dimensional target displacements. *Journal of neurophysiology*, *67*(2), 455-469.

Prablanc, C., Desmurget, M., & Gréa, H. (2003). Neural control of on-line guidance of hand reaching movements. *In Progress in brain research* (Vol. 142, pp. 155-170). Elsevier.

Price, C. J., & Friston, K. J. (2002). Degeneracy and cognitive anatomy. *Trends in cognitive sciences*, *6*(10), 416-421.

Rorden, C., & Karnath, H. O. (2004). Using human brain lesions to infer function: a relic from a past era in the fMRI age?. *Nature Reviews Neuroscience*, *5*(10), 812.

Rossetti, Y., Pisella, L., & Vighetto, A. (2003). Optic ataxia revisited. *Experimental Brain Research*, *153*(2), 171-179.

Rubino, D., Robbins, K. A., & Hatsopoulos, N. G. (2006). Propagating waves mediate information transfer in the motor cortex. *Nature neuroscience*, *9*(12), 1549.

Sarlegna, F., Blouin, J., Bresciani, J. P., Bourdin, C., Vercher, J. L., & Gauthier, G. M. (2003). Target and hand position information in the online control of goal-directed arm movements. *Experimental Brain Research*, *151*(4), 524-535.

Sarlegna, F. R., & Mutha, P. K. (2015). The influence of visual target information on the online control of movements. *Vision research*, *110*, 144-154.

Saturnino, G. B., Antunes, A., & Thielscher, A. (2015). On the importance of electrode parameters for shaping electric field patterns generated by tDCS. *Neuroimage*, *120*, 25-35.

Saturnino, G. B., Puonti, O., Nielsen, J. D., Antonenko, D., Madsen, K. H. H., & Thielscher, A. (2018). SimNIBS 2.1: A Comprehensive Pipeline for Individualized Electric Field Modelling for Transcranial Brain Stimulation. *bioRxiv*, 500314.

Scott, S. H. (2004). Optimal feedback control and the neural basis of volitional motor control. *Nature Reviews Neuroscience*, *5*(7), 532.

Scott, S. H. (2012). The computational and neural basis of voluntary motor control and planning. *Trends in cognitive sciences*, *16*(11), 541-549.

Scott, S. H. (2016). A functional taxonomy of bottom-up sensory feedback processing for motor actions. *Trends in neurosciences*, *39*(8), 512-526.

Siebner, H. R., Hartwigsen, G., Kassuba, T., & Rothwell, J. C. (2009). How does transcranial magnetic stimulation modify neuronal activity in the brain? Implications for studies of cognition. *Cortex*, *45*(9), 1035-1042.

Scott, S. H. (2012). The computational and neural basis of voluntary motor control and planning. *Trends in cognitive sciences*, *16*(11), 541-549.

Stetson, C., & Andersen, R. A. (2014). The parietal reach region selectively anti-synchronizes with dorsal premotor cortex during planning. *Journal of Neuroscience*, *34*(36), 11948-11958.

Teyler, T. J., Hamm, J. P., Clapp, W. C., Johnson, B. W., Corballis, M. C., & Kirk, I. J. (2005). Long-term potentiation of human visual evoked responses. *European Journal of Neuroscience*, *21*(7), 2045-2050.

Todorov, E., & Jordan, M. I. (2002). Optimal feedback control as a theory of motor coordination. *Nature neuroscience*, *5*(11), 1226.

Tsujimoto, T., Mima, T., Shimazu, H., & Isomura, Y. (2009). Directional organization of sensorimotor oscillatory activity related to the electromyogram in the monkey. *Clinical Neurophysiology*, *120*(6), 1168-1173.

Tunik, E., Frey, S. H., & Grafton, S. T. (2005). Virtual lesions of the anterior intraparietal area disrupt goal-dependent on-line adjustments of grasp. Nature neuroscience, 8(4), 505.

van Beers, R. J., Baraduc, P., & Wolpert, D. M. (2002). Role of uncertainty in sensorimotor control. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *357*(1424), 1137.

Vesia, M., Prime, S. L., Yan, X., Sergio, L. E., & Crawford, J. D. (2010). Specificity of human parietal saccade and reach regions during transcranial magnetic stimulation. *Journal of Neuroscience*, *30*(39), 13053-13065.

Vingerhoets, G. (2014). Contribution of the posterior parietal cortex in reaching, grasping, and using objects and tools. *Frontiers in psychology*, *5*, 151.

Vöröslakos, M., Takeuchi, Y., Brinyiczki, K., Zombori, T., Oliva, A., Fernández-Ruiz, A., ... & Berényi, A. (2018). Direct effects of transcranial electric stimulation on brain circuits in rats and humans. *Nature communications*, *9*(1), 483.

Wagner, T. A., Zahn, M., Grodzinsky, A. J., & Pascual-Leone, A. (2004). Three-dimensional head model simulation of transcranial magnetic stimulation. *IEEE Transactions on Biomedical Engineering*, *51*(9), 1586-1598.

Wijdenes, L. O., Brenner, E., & Smeets, J. B. (2011). Fast and fine-tuned corrections when
the target of a hand movement is displaced. *Experimental Brain Research*, *214*(3), 453-
462.
Wijdenes, L. O., Brenner, E., & Smeets, J. B. (2014). Analysis of methods to determine the
latency of online movement adjustments. *Behavior research methods*, *46*(1), 131-139.
Young, M. P., Hilgetag, C. C., & Scannell, J. W. (2000). On imputing function to structure
from the behavioural effects of brain lesions. *Philosophical Transactions of the Royal*
Society of London. Series B: Biological Sciences, *355*(1393), 147-161.
Zhang, Y., Chen, Y., Bressler, S. L., & Ding, M. (2008). Response preparation and inhibition:
the role of the cortical sensorimotor beta rhythm. *Neuroscience*, *156*(1), 238-246.