TWO-SIGMA: a novel TWO-component SInGle cell Model-based Association method for single-cell RNA-seq data

Eric Van Buren [1], Ming Hu [2], Chen Weng [3], Fulai Jin [3], Yan Li [3], Di Wu [1,4*], Yun Li [1,5,6*]

[1]Department of Biostatistics, The University of North Carolina at Chapel Hill
[2]Department of Quantitative Health Sciences, Lerner Research Institute, Cleveland Clinic Foundation
[3]Department of Genetics, School of Medicine, Case Western Reserve University
[4]Department of Oral and Craniofacial Health Sciences, Adams School of Dentistry, The University of North Carolina at Chapel Hill
[5]Department of Genetics, The University of North Carolina at Chapel Hill
[6]Department of Computer Science, The University of North Carolina at Chapel Hill

* To whom correspondence should be addressed: Email: yun_li@med.unc.edu; did@email.unc.edu

July 22, 2019

Abstract

Two key challenges in the analysis of single cell RNA-seq (scRNA-seq) data are excess zeros due to "drop-out" events and substantial overdispersion due to stochastic and systematic differences. Association analysis of scRNA-seq data is further confronted with the possible dependency introduced by measuring multiple single cells from the same biological sample. To address these three challenges, we propose TWO-SIGMA: a TWO-component SInGle cell Model-based Association method. The first component models the drop-out probability with a mixed-effects logistic regression, and the second component models the (conditional) mean read count with a mixed-effects negative binomial regression. Our approach simultaneously allows for overdispersion and accommodates dependency in both drop-out probability and mean mRNA abundance at the gene level, leading to improved statistical power while still providing highly interpretable coefficient estimates. Simulation studies and real data analysis show advantages in type-I error control, power enhancement, and parameter estimation over alternative approaches including MAST and a zero-inflated negative binomial model without random effects. TWO-SIGMA is implemented in the R package "twosigma" available at https://github.com/edvanburen/twosigma.

1 Introduction

Advancements in single-cell sequencing technologies have created many exciting opportunities to researchers yet have also posed many challenges relating to data analysis. Expression profiles can now be analyzed at the single-cell level, providing new insights into the cellular heterogeneity of gene expression. Three characteristic features of single-cell transcriptome sequencing data include excess zero counts, overdispersion of observed counts, and a large number of cells that are sequenced from a relatively small number of samples [1]. Technological limitations including low capture rate and amplification failure lead to "drop-out" events, in which the data may capture only a fraction of the transcriptome of a given cell and mistakenly generate zero measurements for expressed genes. The presence of these zeros creates a dataset with an excess of zeros (often called "zero-inflation") beyond those that occur due to biological factors; these excess zeros often

necessitate special modelling approaches such as a two-component model [2, 3]. Overdispersion, in which the variance in expression exceeds the mean expression, is commonly observed in count-based quantitative sequencing due to large variance in expression and within gene variability over time or across samples. The ideal method should also account for the correlation present because multiple cells are sequenced from the same biological sample. These challenges motivated us to develop the new statistical method TWO-SIGMA. It is designed for association analysis where the primary interest is in performing statistical inference on covariates of interest, such as a treatment effect.

Previous methods for scRNA-seq association analysis do not simultaneously model the excess zero counts, overdispersion, and within-sample correlation. A two-component negative binomial mixture method to perform single-cell differential expression (DE) analysis for only a two-group comparison was developed in [4]. Others have discussed log-linear mixed effects negative binomial regression models in general [5] and as specifically applied to microbiome data [6]. The two-component zero-inflated model without random effects has a long history in the analysis of count and microbiome data [7, 8, 9], however its application in scRNA-seq data is limited. Furthermore, a zero-inflated negative binomial mixed effects model has previously been proposed for modelling zero-inflated count data [10]. The focus of that work was on typical repeated measures applications where the number of repeated measures per individual is small, in contrast to genomic applications which tend to include more repeated measures than samples. Because attention is split between zero-inflated and a similar approach called a hurdle model, and between the Poisson and negative binomial distributions, details regarding the performance and robustness of the zero-inflated negative binomial mixed-effects model are not discussed in as much detail as we can here. Finally, DESingle was introduced as a zero-inflated negative binomial regression to detect differential expression (DE) in scRNA-seq data [11]. Their method is designed for DE detection with a two-level grouping variable and does not employ a regression modeling framework to control for other covariates or account for within-sample correlation.

Previously, the MAST method was introduced as a log-normal hurdle model for the analysis of scRNA-seq data [2]. MAST is similar to TWO-SIGMA in that it also fits a two-component model to account for excess zeros. The two methods differ, however, in that we fit a zero-inflated model on the observed counts while MAST fits a hurdle model (described in more detail in the next section) on the log scale. More importantly, although the ability to include random effects in either component of MAST is mentioned, they do not prioritize their inclusion for scRNA-seq data and do not evaluate the impact of random effects on the model's performance. We will revisit the comparison with MAST in the methods section. Several unsupervised methods, such as ZINB-WaVE [12] and ZIFA [13] have also been proposed for scRNA-seq analysis. Although ZINB-WaVE is similar to our method in that it uses a zero-inflated negative binomial model, both ZINB-WaVE and ZIFA are primarily designed for unsupervised settings in which dimension reduction is the primary goal.

Others use variations of the zero-inflated negative binomial distribution to construct observation-level weights that can be incorporated into the popular bulk RNA-seq pipelines found in the DESeq2 [14] or edgeR [15] Bioconductor packages [16, 17]. These pipelines do not allow for random effects and can involve some transformation of the data in processing or analysis. Thus, one advantage of employing TWO-SIGMA over using the weighting approaches of [16] and [17] is the ability to use random effects to investigate the (gene-specific) within-individual correlation without the need to transform the data.

In this paper, we develop TWO-SIGMA, a TWO-component SInGle cell Model-based Association (TWO-SIGMA) method for single-cell gene expression data. The first component models the probability of "drop-out" with a mixed-effects logistic regression model, while the second component models the (conditional) negative binomial mean parameter with a mixed-effects negative binomial regression model. TWO-SIGMA is extremely flexible in that it does not require a log-transformation of the outcome and allows for (i) overdispersed and zero-inflated counts, (ii) a correlation structure between cells from the same individual, (iii) unbalanced designs (in which the number of cells does not need to be identical for all samples), and (iv) covariates that may differ across the two components.

TWO-SIGMA therefore combines the advantages of random effect modeling and zero-inflated regression models. TWO-SIGMA is the only method to our knowledge for analyzing single-cell gene expression count data that simultaneously allows for zero-inflation, overdispersion, and correlation between cells from the same sample/individual. Simulations studies show that TWO-SIGMA outperforms alternative approaches in both type-I error and power when the data contains even moderate within-sample correlation.

The rest of the article proceeds as follows: first, we specify the TWO-SIGMA model, discuss implications of its parameterization, and provide details on parameter estimation. Next we describe both traditional methods and a new *ad hoc* method to decide whether random effects should be included in our zero-inflated negative binomial model. Then we show simulation results and an application to a dataset of pancreatic islet single-cells, respectively. Finally, we conclude with a discussion.

2 Materials and Methods

2.1 Zero-inflated negative binomial distribution

For a given gene, let i index the samples sequenced and j index the n_i single cells from sample i. Consider the following parameterization of the negative binomial probability mass function (p.m.f.) at a non-negative integer y_{ij} corresponding to the observed read count:

$$\Pr(Y_{ij} = y_{ij}) = f(y_{ij}; \mu_{ij}, \phi)$$
$$= \frac{\Gamma\left(y_{ij} + \phi\right)}{\Gamma(y_{ij} + 1)\Gamma\left(\phi\right)} \left(\frac{1}{1 + \frac{1}{\phi}\mu_{ij}}\right)^{\phi} \left(\frac{\frac{1}{\phi}\mu_{ij}}{1 + \frac{1}{\phi}\mu_{ij}}\right)^{y_{ij}}$$
$$, \ y_{ij} = 0, 1, 2, \ldots$$

With this parameterization, $\mathrm{E}(Y_{ij}) = \mu_{ij}$ and
$\mathrm{Var}(Y_{ij}) = \mu_{ij} + \frac{1}{\phi}\mu_{ij}^2$, such that ϕ is the overdispersion parameter ($\phi > 0$). This parameterization is appealing for interpretability because as $\frac{1}{\phi} \to 0^+$, the density above approaches the Poisson density with mean μ_{ij}. Thus, the model can accommodate data from a Poisson distribution using a very large value of ϕ. A zero-inflated Poisson model is thus nested within this parameterization as mentioned further in the discussion.

To accommodate the excess zeros often observed in scRNA-seq data, we employ the zero-inflated negative binomial distribution (ZINB). This distribution mixes a point mass at zero (from which observations are considered "drop-out") with the negative binomial distribution. Let p_{ij} and μ_{ij} be the probability of drop-out and the mean read count conditional on not being dropped-out for cell j from sample i, respectively. The p.m.f. of one observation Y_{ij} under the ZINB distribution is given by:

$$\mathrm{P}(Y_{ij} = 0) = p_{ij} + (1 - p_{ij})f(0; \mu_{ij}, \phi)$$
$$\mathrm{P}(Y_{ij} = y_{ij}) = (1 - p_{ij})f(y_{ij}; \mu_{ij}, \phi), y_{ij} = 1, 2, 3, \ldots \tag{1}$$

The ZINB distribution thus assumes that there are two sources of zeros in the data: the first source is the process that governs drop-out and the second source is from the negative binomial process for genes that are not dropped-out. This differs from hurdle models used for gene expression data [2] which use a left-truncated distribution for the positive expression component—meaning zeros can only be generated from the drop-out process. Thus, the hurdle model does not allow zero expression measurements due to biological variation.

Interpretations from the zero-inflated model are quite natural for single-cell gene expression data because it is reasonable to believe that some observed zeros are "structural zeros" with *bona fide* zero expression due to stochastic biological factors (e.g. transcriptional bursting, cell cycle) and not due to technical drop-out [1]. Although semantic, the distinction regarding the source of zeros affects the interpretation of model coefficients and is important because these models are often misinterpreted by researchers [18].

Table 1: Type-I error evaluations in simulated data: Shows type-I error using the LRT to test the joint null hypothesis of a simulated binary disease status indicator, $H_0 : \alpha_1 = 0$, $\beta_1 = 0$ versus $H_a : \alpha_1 \neq 0$ or $\beta_1 \neq 0$, with a significance level of 0.05. "2-S" refers to TWO-SIGMA, ZINB refers to a zero-inflated negative binomial model without random effects and MAST refers to the model described in [2]. 10,000 genes were simulated.

				Case 1 in Supplement: 50 Cells per 1000 Ind.			Case 3 in Supplement: 500 Cells per 100 Ind.			Case 4 in Supplement: 2000 Cells per 25 Ind.		
	Sim Params											
	ϕ	σ_a	σ_b	2-S	ZINB	MAST	2-S	ZINB	MAST	2-S	ZINB	MAST
No	10	0	0	**0.049**	0.051	0.089	**0.042**	0.050	0.090	**0.041**	0.052	0.090
R.E.	2	0	0	**0.048**	0.051	0.080	**0.038**	0.044	0.079	**0.041**	0.052	0.086
	1	0	0	**0.048**	0.052	0.081	**0.044**	0.051	0.087	**0.042**	0.051	0.090
Small	10	0.1	0.1	**0.051**	0.132	0.144	**0.056**	0.534	0.313	**0.077**	0.795	0.487
R.E.	2	0.1	0.1	**0.051**	0.078	0.089	**0.057**	0.323	0.176	**0.072**	0.643	0.361
	1	0.1	0.1	**0.049**	0.066	0.095	**0.053**	0.224	0.174	**0.075**	0.548	0.361
Large	10	0.5	0.5	**0.051**	0.621	0.290	**0.055**	0.941	0.716	**0.076**	0.984	0.875
R.E.	2	0.5	0.5	**0.053**	0.505	0.275	**0.056**	0.909	0.685	**0.076**	0.974	0.857
	1	0.5	0.5	**0.050**	0.404	0.247	**0.053**	0.873	0.649	**0.074**	0.964	0.827

2.2 TWO-SIGMA

We can now provide the full TWO-SIGMA specification:

$$\text{logit}(p_{ij}) = \boldsymbol{z_{ij}^T}\boldsymbol{\alpha} + a_i, a_i \sim N(0, \sigma_a^2)$$
$$\log(\mu_{ij}) = \boldsymbol{x_{ij}^T}\boldsymbol{\beta} + b_i, b_i \sim N(0, \sigma_b^2), \text{ assume } a_i \perp\!\!\!\perp b_i \tag{2}$$

The model is fit for each gene individually, so all parameters are gene-specific. $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ are fixed effect coefficient vectors and the corresponding vectors of covariates $\boldsymbol{z_{ij}}$ and $\boldsymbol{x_{ij}}$ can be different. a_i and b_i are sample-specific intercepts (discussed more in the next section). Prediction of sample-specific intercepts and estimation of the variance components σ_a^2 and σ_b^2 allow us to investigate heterogeneity among individuals, and tests of whether the variance components equal zero allow us to separately (or jointly) evaluate the need for random effects. Separate variance components are estimated because the different link functions in the two components correspond to linear predictors with different scales. Including the random effects terms also helps control for any within-sample correlation, providing more accurate estimates and standard errors of fixed effect parameters.

As part of our **twosigma** R package, we employ the **glmmTMB** package [19] to fit the model specified in equation (2). This package is well-suited to fit generalized linear mixed models (GLMMs) because the user can easily specify an arbitrarily complex model composed of fixed and random effects. More details regarding computational considerations can be found in section 5 of the supplement.

To summarize, TWO-SIGMA can control for different covariates in each component, incorporate random effects to accommodate within-sample dependency, analyze unbalanced data, and allow for zero-inflated and overdispersed counts. The implementation of the model strikes a balance between computational accuracy and efficiency, even as the number of random effects (number of samples in the context of the scRNA-seq data) or the number of single cells per sample increases.

2.3 Evaluating the need for random effects

One primary methodological contribution of TWO-SIGMA for scRNA-seq data analysis is the inclusion of random effect terms in each of the two components, which is a well-established technique to account for within-sample correlation. Ignoring random effects in TWO-SIGMA is equivalent to assuming that cells from the same sample/individual are independent. This assumption can lead to underestimated standard

errors and thus inflated type-I errors. An example is given in table 3 in the real data analysis section.

Evaluating the need for a random effect term involves a hypothesis test of whether the corresponding variance component(s) equal zero. For example, consider testing whether random effects are needed in either component of TWO-SIGMA with $H_0 : \sigma_a = \sigma_b = 0$ versus $H_a : \sigma_a > 0$ or $\sigma_b > 0$ using the Likelihood Ratio Test (LRT). This procedure requires fitting the model under both the null and the alternative hypotheses, but is a preferred method to determine whether the random effect terms significantly improve model fit. Other less desirable post-fitting options to compare models with and without random effects include information criteria like AIC and BIC or Wald tests of the variance components [20]. Critically, all options discussed require fitting the "full" model including the random effect terms. The scRNA-seq application is distinct from typical repeated measures analyses in that the number of repeated measures (cells) typically far exceeds the number of samples. Such designs can entail more extensive computational time for each gene over scenarios involving a smaller number of repeated measures from a modest number of individuals. These computational burdens are especially relevant given that scRNA-seq data typically include thousands of genes, each of which is fit separately using TWO-SIGMA. It would therefore be useful to identify the genes that are most likely to need the random effect terms without having to fit the full model to each gene.

2.4 ad hoc approach

We utilize the following *ad hoc* approach to determine whether random effects are needed: using a one-way ANOVA, we regress the Pearson residuals from a zero-inflated negative binomial regression model without random effects on the sample label and take the p-value from the overall F test. This p-value serves as a rudimentary measure of whether the residuals tend to differ across samples. If they do, this is evidence that residuals are not exchangeable across samples. The full TWO-SIGMA specification including random effects will then be fit to more formally evaluate the need for random effect terms. In contrast, when the residuals show no tendency of differing across samples, we do not have evidence to believe that they are structured/clustered within samples and thus will not fit the full model with random effects. Through simulations we found that this procedure is very effective in identifying the need for random effects. Results from applying this proposed method to a real dataset of pancreatic islet cells are given in the data analysis section. In simulations, computation runtime was the longest for models attempting to fit random effects when variance components were truly zero (see supplementary tables 1-4). Therefore, as discussed more in section 3 of the supplementary file, the *ad hoc* method can dramatically reduce overall computation time over many genes in addition to increasing model parsimony where most appropriate.

2.5 Simulation studies

To evaluate the performance of TWO-SIGMA, we simulated data in a variety of scenarios. Competing methods for comparison include a zero-inflated negative binomial model without random effects (ZINB model) and MAST using the $\log_2$ transformation of the counts in a hurdle model. Simulated additional covariates included age and the cellular detection rate (CDR, see the real data analysis section and [2] for more details) to mimic our real data analysis. Values of $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ were designed to mimic realistic parameter values observed in our pancreatic data analysis. Models were evaluated using the likelihood ratio test on the joint null hypothesis that a binary disease status indicator is not associated with expression through either drop-out probability or the conditional mean, $H_0 : \alpha_1 = \beta_1 = 0$. We consider two different ways of simulating data: one in which the number of samples far exceeds the number of cells per sample, as is typical in most repeated measures contexts, and the other in which the number of cells far exceeds the number of samples, as is the case in scRNA-seq data. In each scenario we simulated 10,000 genes and used 0.05 as the nominal significance rate to evaluate type-I error and power.

3 Results

3.1 Type-I error control

Table 1 shows results from simulations in which the true values of the overdispersion parameter ϕ and the variance components σ_a, σ_b vary. Type-I error is well-controlled for TWO-SIGMA in the scenarios involving

Figure 1: Power evaluations in simulated data: Shows the power to test $H_0 : \alpha_1 = \beta_1 = 0$ by varying the effect size with 500 cells from each of 100 individuals. Values of ϕ, σ_a, and σ_b were all set to 0.1 to mimic the "Small R.E." section of table 1 and 10,000 genes were simulated. Because of the type-I error inflation from the ZINB model and MAST seen in table 1, true power was calculated and plotted for these methods using the empirical significance threshold from the corresponding setting under the null. TWO-SIGMA can bypass the need for computationally expensive resampling procedures needed to generate true power because it preserves the type-I error as seen in table 1. See the discussion at the beginning of section 4 of the supplement for more details about computing true power and discussion regarding power trends across the different methods.

more individuals than cells. When the number of cells increases, type-I errors from TWO-SIGMA are slightly inflated over the nominal rate of 5%, but consistently remain superior to the results from the ZINB model or MAST in the presence of ignored non-zero variance components. For example, the last row of table 1 shows that, when $\phi = 1$ and $\sigma_a = \sigma_b = 0.5$, type-I error for TWO-SIGMA increases from 0.05 to 0.053 to 0.074 as the number of individuals decreases from 1000 to 100 to 25. In contrast, the ZINB model and MAST have inflated type-I errors in every scenario that increase to nearly 1 as the number of individuals decreases. This is not surprising because both of the latter methods cannot account for any within-sample dependency structure among the single cells from the same sample. Ignoring the dependency introduced by even a moderate random effect size can thus have a drastic impact on the type-I error. When true variance components are zero, both TWO-SIGMA and the ZINB model preserve type-I error while MAST consistently has higher type-I error, as seen in the first three rows of table 1. Coverage of confidence intervals for α, β, and ϕ always approaches the nominal level (Supplementary tables S1-S4). The reason for the slightly inflated type-I error for TWO-SIGMA observed in the scenario with 25 individuals is worth mentioning briefly. The smaller number of individuals (25) provides less information to estimate the sample-specific variance components σ_a and σ_b and few unique values of the simulated binary disease status indicator. The slightly lower coverage for variance components in the last 6 sets of supplementary table 4 is one illustration of the (relative) difficulty in getting precise variance component estimates. TWO-SIGMA outperforms MAST or the ZINB model in preserving type-I error and estimating parameters under a variety of sample size breakdowns and with a variety of true parameter values. See supplementary figures S4-S5 for type-I error across more stringent significance thresholds for representative scenarios.

3.2 TWO-SIGMA retains high power under a variety of scenarios

Because the ZINB model and MAST both have heavily inflated type-I errors in many cases, using raw (or "apparent") power does not provide a fair comparison for these two methods. For each method and each simulation setting under the null, we therefore calculate the empirical significance threshold, defined as the test statistic value at the quantile associated with 1 minus the significance level. A percentage of statistics equal to the nominal significance level will then be larger than this threshold. For various alternative hypotheses, we calculate "true" power for MAST and the ZINB model by using the empirical significance threshold from the corresponding setting under the null as the rejection threshold instead of a usual theoretical threshold (e.g. 5.9915 from χ_2^2 at the .05 level). Figure 1 plots raw power for TWO-SIGMA and true power for MAST in the ZINB model in the following four scenarios: effect in both components, in either the same or opposite directions, and effects in only one of the two components. In the first three scenarios, MAST consistently has the lowest power, while TWO-SIGMA and the ZINB model have very similar power in the first two scenarios, beginning at around 20% and increasing to nearly 100%. The ZINB model has higher power than TWO-SIGMA in the third scenario but the lowest power in the fourth scenario. In simulation, computing the empirical significance thresholds and true power is straightforward and computationally included. In real data settings, however, computationally intensive resampling approaches are needed for reliable estimates of the empirical significance thresholds. Because TWO-SIGMA preserves type-I error, we can rely on raw power and can therefore bypass the need for any resampling approach for valid inference. This is a key advantage and shows that TWO-SIGMA is more robust and flexible than the ZINB model while both preserving the type-I error and having high power without any additional computation. When the effect is only in the zero-inflation component, power is lower for all methods than in the first three scenarios. Such effects present only in the zero-inflation component are known to be more difficult to detect, as seen in [8]. For full power results, including more detailed comparisons to the ZINB model with additional discussion, see section 4, figures S6-S8 and tables S5-S12 of the supplement.

3.3 Sensitivity of the *ad hoc* method for random effect screening in simulated data

We evaluated the *ad hoc* screening procedure used to select genes possibly in need of random effect terms using the simulated data. We found it to be effective as a screening procedure: for the data used in Table 1, most genes with non-zero variance components had p-values less than 0.05 and were flagged as in need of estimating variance components using TWO-SIGMA. When the variance components are zero, however,

Figure 2: Presence of overdispersion in real data: Shows the need of a non-linear mean-variance relationship in the pancreatic islet data. Each point represents the mean-variance relationship for one gene. In the legend ϕ represents the overdispersion parameter of the negative binomial distribution and p represents the drop-out probability.

the p-values from the ad hoc method are close to uniformly distributed, indicating that the procedure is not too liberal under the null. See section 2 and figures S2 and S3 of the supplement for more details and some examples.

3.4   Pancreas real data analysis

For illustrative purposes we applied TWO-SIGMA to a dataset of pancreatic islet cells isolated from nine individuals (see section 1 of the supplement and [21] for full details on the data processing and generation steps). To focus on the most informative cells and genes, we applied rather aggressive filtering of the data to keep the top 2,000 genes by number of transcripts observed and only keep cells with more than 1000 transcripts across these genes. After merging across all nine individuals, we were left with 1,290 genes and 10,269 single cells of which we used only the 7,774 for which cell type information was available based on the expression of signature genes. Here we focus our attention on alpha and beta cells, which compose the majority (55% and 34%, respectively) of the cells in our dataset. Type-II diabetes (T2D) status and age were used as sample-specific covariates in all analyses. The cellular detection rate (CDR) is defined in [2] as the percentage of genes expressed over some background level of expression (often chosen to be zero). The CDR therefore has a biological interpretation as a cellular scaling factor and is a surrogate for both technical and biological variation. This confirms the conclusions of [22] that the CDR can explain a substantial proportion of observed expression variability and should be included in any association analysis of scRNA-seq data. As such, we include CDR in all analyses performed and stratified by cell type. For more details about the pancreas data processing, see section 1 and figure S1 of the supplementary file.

Figure 2 plots the relationship of mean versus variance for the 1,290 genes we used in our analysis. It shows that the Poisson and zero-inflated Poisson models cannot adequately account for the overdispersion observed in many genes. In contrast, TWO-SIGMA can accommodate these mean-variance pairs in a quadratic relationship via the overdispersion parameter ϕ. Because we have only nine individuals, we chose to focus on analyses excluding the zero-inflation random effect terms a_i to improve convergence and overall model fit. Some genes still showed convergence issues–partly indicative of a misspecified or overparameterized model and partly due to the small number of cells and samples in the dataset. As a general guideline, we recommend that users with concerns or limited computational resources begin including random effects in the mean component, and scale upwards to include random effects in the zero-inflation component if

Table 2: Rejection summaries from the pancreas data: Shows the proportion of genes in the pancreatic islet data with rejected nulls for various hypotheses related to T2D. The TWO-SIGMA model as specified in equation (2) was fit with no zero-inflation variance component (no ZIVC).

Hypothesis	Alpha Cells No ZIVC	Beta Cells No ZIVC
Overall Disease Status	0.161	0.111
Overall R.E. Test	0.738	0.724
NB vs. Poisson	0.627	0.555

performance is satisfactory.

Table 2 shows the proportions of genes showing statistically significant results at the .05 level for three types of hypothesis tests: the joint test of significance for the binary disease indicator $H_0 : \alpha_1 = \beta_1 = 0$, the test of the mean model variance component $H_0 : \sigma_b = 0$, and the test for the presence of overdispersion $H_0 : \frac{1}{\phi} = 0$. For example, when fitting the TWO-SIGMA model without the zero-inflation variance component to alpha cells, 73.8% of genes had statistically significant variance components in the mean model.

3.5 Cell-type specific genes often show a need for random effect inclusion

We matched 234 and 120 genes in our data that were identified in previous studies as cell-type specific in alpha or beta cells, respectively. ([23], supplementary table 10). After stratifying the data by cell type and removing genes with more than 90% or less than 10% zeros, we fit TWO-SIGMA (excluding a_i as mentioned previously) to the remaining 222 alpha cell-specific and 111 beta cell-specific genes to alpha cells and beta cells, respectively. Of these, 93 alpha cell-specific genes and 85 beta cell-specific genes had statistically significant variance components σ_b. This suggests that non-negligible between-sample variation—not attributable to cell-type—is present for these genes.

3.6 Impact of ignoring within-sample correlation

Table 3: Influence of failing to include needed random effects: Gives mean component estimates for gene *RPS29* with (top panel) and without (bottom panel) random effects

Effect	Estimate	Std. Error	z value	p-value
Intercept	0.521	0.207	2.515	0.012
T2D	-0.349	0.292	-1.197	0.231
age	-0.284	0.256	-1.109	0.267
CDR	0.394	0.011	36.284	<.001
σ_b	0.490			
Intercept	0.833	0.021	40.094	<.001
T2D	-0.605	0.032	-19.090	<.001
age	-0.057	0.017	-3.324	<.001
CDR	0.390	0.015	26.611	<.001

Models for genes that mistakenly exclude the b_i random effect term often show highly significant results for covariates; this significance can disappear when including the random intercept term—possibly indicative of a false positive due to failing to account for within-sample correlation. For example, gene *RPS29* demonstrates this pattern in alpha cells. Table 3 shows that failing to include random effects—and thereby assuming independence of all single cells—can lead to vastly underestimated standard errors. T2D status and age change from highly significant to insignificant when including a random intercept term. The standard error for the coefficient of T2D increases by a factor of 9 from 0.032 to 0.292, and the magnitude of the point estimate is halved from -0.605 to -0.349. Individual covariates such as T2D can thus exhibit dramatically

Table 4: Agreement between TWO-SIGMA and MAST: Shows the agreement in rejecting the omnibus null hypothesis of an association between T2D status and gene expression in alpha cells using a Bonferroni adjusted significance level of 5×10^{-5}.

TWO-SIGMA	MAST	
	No Reject	Reject
No Reject	1013	273
Reject	1	3

increased type-I error when random effects are incorrectly ignored. In contrast, the coefficient and associated standard error for the cellular detection rate (CDR) are nearly identical in the two models. This result is expected given that CDR is a cell-level covariate and shows that including sample-specific random effects leads to very minor changes in the estimation of any covariates that are not sample-specific. Our emphasis in this section is not to draw conclusions about any association between *RPS29* and T2D, but rather to illustrate that ignoring random effects has the potential to alter scientific conclusions.

We also used alpha cells to test the overall effect of T2D using both TWO-SIGMA to MAST. Table 4 shows that MAST rejects in many more instances than TWO-SIGMA. Of the 273 genes that were rejected with MAST but not with TWO-SIGMA, 234 have statistically significant variance components in TWO-SIGMA. This further illustrates the possibility that fixed effect coefficients can be mistakenly deemed significant in the presence of within-sample correlation.

3.7 The *ad hoc* method successfully separates genes that need random effects

Finally, we used all 1,290 genes from the islet dataset to demonstrate the usefulness of the *ad hoc* method to determine the need for the random effects terms b_i. Figure 3 shows that likelihood ratio statistics from formal testing of b_i are consistently larger for genes selected by the procedure than those not selected. This pattern suggests that the *ad hoc* procedure described earlier can effectively identify genes that will exhibit non-zero variance components in real data.

4 Discussion

We have developed a two-component zero-inflated negative binomial model with random effects for association analysis of scRNA-seq data. The model builds on the well-established literature in both zero-inflated models and generalized linear mixed models. It keeps the data on the original scale while simultaneously allowing for zero-inflation, overdispersion, and within-sample correlation.

Including random effects explicitly controls for within-sample correlation, and can improve mean parameter and standard error estimates. Given that many scRNA-seq studies have few samples, it would be reasonable to consider controlling for sample as a fixed effect rather than a random effect. However, there are two reasons to prefer incorporating a random effect to a fixed effect approach. First, we are interested in estimating a variance component that can apply to all samples in the population. Second, the random intercepts explicitly control for the within-sample correlation, rather than only providing adjusted parameter estimates for included covariates. An alternative approach to accounting for such sample-level repeated measures would be to fit a marginal model with the generalized estimating equations (GEE) approach instead of a mixed effects model [24]. We chose not to do so for two main reasons: first, we retain the flexibility for sample-level prediction. Second, given that many scRNA-seq experiments are conducted over a small number of samples, it is likely that the empirical (sandwich) covariance estimate would underestimate the true standard errors [24].

Incorrectly excluding random effects and assuming independence of cells can lead to underestimated standard errors of fixed effects and can therefore increase the type-I error of hypothesis tests relating to fixed effects parameters. See table 3 for an example. If the random effect terms do not contribute to the model fit, as judged by a statistical test or practical significance, they can be removed easily within the general framework of TWO-SIGMA. Random intercepts can also be useful even when they are not of direct

Figure 3: Ability of the ad hoc method to identify genes in need of random effects: Shows boxplots of the LR statistics from the joint test of the need for random effects, $H_0 : \sigma_a = \sigma_b = 0$, using TWO-SIGMA. Genes that our *ad hoc* procedure suggests need random effects ("Need RE") and genes the procedure suggests do not ("Don't Need RE") are compared. Both panels were created using TWO-SIGMA as specified in equation (2) but with no zero-inflation variance component (no ZIVC).

interest: they often capture the effects of omitted sample-specific covariates, and can limit the bias of fixed effect coefficients caused by misspecification. For example, if cell-type information is missing, and varies between individuals, a random intercept term can limit the resulting bias and p-value inflation observed in fixed effects parameters. Our *ad hoc* method proves to be a useful tool to both: (1) select genes that could benefit from including random effect terms and (2) reduce overall computation time by suggesting genes that do not need to be fit including random effect terms.

Because we expect *a priori* that zero-inflation will occur in scRNA-seq, it is beneficial to include a component dedicated to it. The zero-inflation component in TWO-SIGMA is flexible in that it allows for a different set of covariates from the mean model, or no covariates at all. For example, one might be interested in using zero-inflation only to improve mean parameter estimation. In this scenario, a constant probability of drop-out could be assumed via an intercept-only regression model. This would prevent coefficient estimates in the mean model from being overly shrunk towards zero, as would occur if drop-out was not accounted for, but would also limit the total number of parameters estimated and maximize model parsimony. Even if the data are not truly generated from a zero-inflated process, or if drop-out is viewed as a "nuisance," using the two component model in equation (2) can be a convenient choice to improve model fit and fixed effect parameter estimation. See section 6 of the supplement and supplementary figure S9 for more discussion regarding the zero-inflation component.

Fitting the TWO-SIGMA model also provides a way to choose between the zero-inflated negative binomial and zero-inflated Poisson distributions that may be useful in a standard data analysis; if the overdispersion parameter estimate is small, one can justifiably reduce model complexity and fit a Poisson model. Specifically, one can test $H_0 : \frac{1}{\phi} = 0$ versus $H_a : \frac{1}{\phi} > 0$ using the likelihood ratio test (p-values come from a 50:50 mixture of χ_1^2 and a point mass at zero). As seen in table 2, the p-value from such a test will often suggest a significant deviation from the Poisson model in scRNA-seq data.

Finally, our experience suggests that variance component estimates are often much smaller in the zero-inflation component than in the mean component. Therefore, as we did in the real data analysis, it might be a pragmatic choice to exclude random effects from the zero-inflation component of TWO-SIGMA. A key strength of TWO-SIGMA is the flexibility to easily customize the model within the general framework either

a priori or via iterative removal based on statistical hypothesis tests of features such as random effects, overdispersion, or the drop-out component.

5 Data Availability

The pancreas Drop-seq data is available at GEO: GSE101207.

6 Funding

This work was supported by the National Institute of Health [R01HL129132 to Yun Li and E.V.B, U54DK107977 to M.H., R01CA215347 to D.W., R01HG009658 to F.J., R01DK113185 to Yan Li].

Conflict of Interest Statement. None declared.

References

[1] Wills, Q. F. et al. (2014) Single-cell gene expression analysis reveals genetic associations masked in whole-tissue experiments. *Nature Biotechnology,* **31**, 748–753.

[2] Finak, G., McDavid, A., Yajima, M., Deng, J., Gersuk, V., Shalek, A. K., Slichter, C. K., Miller, H. W., McElrath, M. J., Prlic, M., Linsley, P. S., and Gottardo, R. (Dec, 2015) MAST: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biology,* **16**(1), 278.

[3] Hicks, Stephanie C, T. F. W., Teng, M., and Irizarry, R. A. (11, 2017) Missing data and technical variability in single-cell RNA-sequencing experiments. *Biostatistics,* **19**(4), 562–578.

[4] Kharchenko, P. V., Silberstein, L., and Scadden, D. Y. (2014) Bayesian approach to single-cell differential expression analysis. *Nature Methods,* **11**, 740–744.

[5] Booth, J. G. et al. (2003) Negative binomial loglinear mixed models. *Statistical Modelling,* **3**, 179–191.

[6] Zhang, X. et al. (2017) Negative binomial mixed models for analyzing microbiome count data. *BMC Bioinformatics,* **18**.

[7] Lambert, D. (1992) Zero-Inflated Poisson Regression, With an Application to Defects in Manufacturing. *Tecnometrics,* **34**.

[8] Chen, E. Z. and Li, H. (2016) A two-part mixed-effects model for analyzing longitudinal microbiome compositional data. *Bioinformatics,* **32**, 2611–2617.

[9] Hall, D. B. (2000) Zero-Inflated Poisson and Binomial Regression with Random Effects: A Case Study. *Biometrics,* **56**.

[10] Min, Y. and Agresti, A. (2005) Random effect models for repeated measures of zero-inflated count data. *Statistical Modelling,* **5**, 1–19.

[11] Miao, Z., Deng, K., Wang, X., and Zhang, X. (2018) DEsingle for detecting three types of differential expression in single-cell RNA-seq data. *Bioinformatics,* **34**(18), 3223–3224.

[12] Risso, D., Perraudeau, F., Gribkova, S., Dudoit, S., and Vert, J.-P. (2017) A general and flexible method for signal extraction from single-cell RNA-seq data. *Nature Communications,* **9**.

[13] Pierson, E. and Yau, C. (2015) ZIFA: Dimensionality reduction for zero-inflated single-cell gene expression analysis. *Genome Biology,* **16**.

[14] Love, M. I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology,* **15**, 550.

[15] McCarthy, J., D., Chen, Yunshun, Smyth, and K., G. (2012) Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research,* **40**(10), 4288–4297.

[16] Van den Berge, K., Perraudeau, F., Soneson, C., Love, M. I., Risso, D., Vert, J.-P., Robinson, M. D., Dudoit, S., and Clement, L. (Feb, 2018) Observation weights unlock bulk RNA-seq tools for zero inflation and single-cell applications. *Genome Biology,* **19**(1), 24.

[17] Van den Berge, K., Soneson, C., Love, M. I., Robinson, M. D., and Clement, L. (2017) zingeR: unlocking RNA-seq tools for zero-inflation and single cell applications. *bioRxiv,*.

[18] Preisser, J. et al. (2012) Review and recommendations for zero-inflated count regression modeling of dental caries indices in epidemiological studies. *Caries Research,* **46**, 413–423.

[19] Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Maechler, M., and Bolker, B. M. (2017) glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal,* **9**(2), 378–400.

[20] Fitzmaurice, G. M. et al. (2003) Applied Longitudinal Analysis, Second Edition, John Wiley & Sons, Inc., Hoboken, NJ.

[21] Fang, Z., Weng, C., Li, H., Tao, R., Mai, W., Liu, X., Lu, L., Lai, S., Duan, Q., Alvarez, C., Arvan, P., Wynshaw-Boris, A., Li, Y., Pei, Y., Jin, F., and Li, Y. (2019) Single-Cell Heterogeneity Analysis and CRISPR Screen Identify Key -Cell-Specific Disease Genes. *Cell Reports,* **26**(11), 3132 – 3144.e7.

[22] Hicks, S. C. et al. (2015) On the widespread and critical impact of systematic bias and batch effects in single-cell RNA-Seq data. *bioRxiv,* **025528**.

[23] Lawlor, N. et al. (2017) Single-cell transcriptomes identify human islet cell signatures and reveal cell-typespecific expression changes in type 2 diabetes. *Genome Research,* **27**, 208–222.

[24] Agresti, A. (2013) Categorical Data Analysis, Third Edition, John Wiley & Sons, Inc., Hoboken, NJ.