Restriction Enzyme Based Enriched L1Hs sequencing (REBELseq)

Benjamin C. Reiner, PhD[1,†,*]; Glenn A. Doyle, PhD[1,†]; Andrew E. Weller, MD[1]; Rachel N.

Levinson, BA[1]; Esin Namoglu, BS[1]; Alicia Pigeon, BS[1]; Gabriella Arauco-Shapiro, MS[1]; Emilie

Dávila Perea[1]; Cyndi Shannon Weickert, PhD[2]; Gustavo Turecki, MD, PhD[3]; Deborah C. Mash,

PhD[4]; Richard C. Crist, PhD[1] and Wade H. Berrettini, MD, PhD[1]

[1] Center for Neurobiology and Behavior, Department of Psychiatry, Perelman School of

Medicine, University of Pennsylvania; [2] Department of Neuroscience and Physiology, Upstate

Medical University; [3] McGill Group for Suicide Studies, Douglas Mental Health University

Institute, McGill University; [4] Dr. Kiran C. Patel College of Allopathic Medicine, NOVA

Southeastern University, [†] These authors contributed equally to this work, [*] corresponding

author: bcreiner@pennmedicine.upenn.edu

Corresponding author information

Benjamin C. Reiner, PhD

125 S. 31st St., room 2208-1

Philadelphia, PA 19104

Office: (215)-573-0278

Fax: (215)-573-2041

Email: bcreiner@pennmedicine.upenn.edu

Abstract

Long interspersed element-1 retrotransposons (LINE-1 or L1) are ~6 kb mobile DNA

elements implicated in the origins of many Mendelian and complex diseases. The actively

retrotransposing L1s are mostly limited to the L1 human specific Ta subfamily. In this

manuscript, we present REBELseq as a method for the construction of differentially amplified

next-generation sequencing libraries and bioinformatic identification of Ta subfamily long

interspersed element-1 human specific elements. REBELseq was performed on DNA isolated

from NeuN+ neuronal nuclei from postmortem brain samples of 177 individuals and empirically-

driven bioinformatic and experimental cutoffs were established. REBELseq reliably identified

both known and novel Ta subfamily L1 insertions distributed throughout the genome.

Differences in the proportion of individuals possessing a given reference or non-reference

retrotransposon insertion were identified. We conclude that REBELseq is an unbiased, whole

genome approach to the amplification and detection of Ta subfamily L1 retrotransposons.

Introduction

Retrotransposons are a class of mobile DNA elements capable of replicating and

inserting copies of themselves elsewhere in the genome using an RNA intermediate (1). Long

interspersed element-1 (L1), a type of retrotransposon that is currently active in the human

genome, is estimated to constitute approximately 17% of the human genome (2). Despite their

abundance, most L1s are truncated or mutated to the point of no longer being able to

retrotranspose (3). There are ~100 L1s in each human genome that are competent, meaning

they are full length and capable of replicating, the vast majority of which belong to the L1 human

specific (L1Hs) Ta subfamily (4). Alternatively, in agreement with Kazazian *et al.* (5), Beck *et al.*

described two competent L1s of the pre-Ta subfamily (2). A competent L1Hs is ~6 kb in length

and contains a promoter, 5' and 3' untranslated regions, and two open reading frames (ORF):

ORF1, encoding an RNA binding protein (6) , and ORF2, encoding a fusion protein that acts as

both a reverse transcriptase (7) and endonuclease (8).

Whereas competent retrotransposons are mostly limited to full length L1Hs of the Ta

subfamily, the remnants of vast numbers of retrotransposons, from both evolutionarily older and

Ta subfamily L1Hs can be found throughout the human genome. Notably, Ta subfamily L1Hs

contain identifying nucleotides in the 3' UTR that can be utilized for their differential amplification

by polymerase chain reaction (PCR) (4). Retrotransposition of competent L1Hs elements

frequently results in 5' truncation of the new L1Hs element insertions. Therefore, to reliably

amplify and detect full length and truncated L1Hs elements, including germline polymorphic and

individual somatic mutations, from the Ta subfamily, PCR primers for amplification must be

targeted near the 3' end of the L1Hs Ta subfamily sequence.

In this manuscript, we present REBELseq (Restriction Enzyme Based Enriched L1Hs

sequencing), a scalable technique for the differential amplification of both full length and 5'

truncated L1s. As described below, we specifically target the Ta subfamily of L1Hs and

summarize the results of the application of REBELseq to DNA samples from 177 individuals.

Materials and Methods

<u>Brain samples and purification of genomic DNA from NeuN+ nuclei</u>

177 fresh frozen human postmortem prefrontal cortex brain samples (Brodmann's Area

9 or 10) were provided by the Douglas-Bell Canada Brain Bank at McGill University, the Human

Brain and Spinal Fluid Resource Center at UCLA, the Sydney Brain Bank at Neuroscience

Research Australia, or the University of Miami Brain Endowment Bank. All work was approved

by the University of Pennsylvania Institutional Review Board as category four exempt human

subject research. All reported age, sex and ethnicity data are based on associated medical

records. Sample set contained 138 males (Age 45.6 ± 15.5; 93 European, 25 African, 15

Hispanic, 2 Asian and 3 of unknown ethnic origin) and 39 females (Age 55.5 ± 19.2; 33

European, 1 African, 3 Hispanic and 2 of unknown ethnic origin). See Supplemental Methods for

full details.

Brain samples were homogenized and co-stained with DAPI and an α-NeuN-AF488

antibody (Millipore # MAB377X) using a modification of a previously described method (9), and

NeuN+ and NeuN- nuclei were isolated using a FACSAria II (Beckman-Coulter). Isolated nuclei

were lysed overnight in digestion buffer in a 56 °C water bath, and the following morning

genomic DNA (gDNA) was isolated using the Zymo Research genomic DNA clean and

concentrator kit (#D4011). The concentration of gDNA was quantified using a Qubit 3

fluorometer and high-sensitivity double-stranded DNA assay (ThermoFisher #Q32854). See

Supplemental Methods for full details.

Ta subfamily L1Hs-enriched library construction and sequencing

Ta subfamily L1Hs-enriched next generation sequencing libraries were constructed

utilizing the REBELseq technique (Fig. 1). 33 ng of gDNA extracted from NeuN+ neuronal nuclei

was digested with HaeIII, in the presence of shrimp alkaline phosphatase. A single primer

extension utilizing a 3' diagnostic 'A' nucleotide (L1HsACA primer, see Supplemental Oligomers

for all primer sequences; all oligos from IDT, Iowa, USA), a more stringent variation of a primer

originally designed by Ewing and Kazazian (10), extends only Ta subfamily L1 3' UTR sequence

(4) and the adjacent downstream genomic DNA, leaving a terminal 3' A-overhang. The A-

overhang products were ligated to a double stranded T-linker molecule, originally designed for a

different technique (11), and the ligated products were amplified using the L1HsACA primer and

a T-linker specific primer to enrich the number of copies of each unique Ta subfamily L1Hs

insertion (Primary PCR). The Primary PCR product was diluted and used as a template for a

hemi-nested Secondary PCR reaction using the T-linker primer and the Seq2-L1HsG primer.

The purpose of the hemi-nested secondary PCR reactions is three-fold: to reduce the length of

3' L1 sequence carried forward, to add an Illumina sequencing primer, and, most importantly, to

use the L1Hs diagnostic 3' G nucleotide of the Seq2-L1HsG primer to further enrich the L1Hs

Ta subfamily. Secondary PCR products were cleaned and size selected using KAPA Pure

Beads (Roche #KK8002) according to the manufacturer's protocol for a 0.55X-0.75X double-

sided size selection, generating a purified library of 200-1,000 bp fragments. A Tertiary PCR

using 5' overhang primers then added Illumina flow cell adapters, containing a single

multiplexing index, to both ends of the amplicon. Sequencing libraries then underwent a final

cleanup and size selection using KAPA Pure Beads 0.55X-0.75X double-sided size selection

and the average fragment size of the library was determined using the 2100 Bioanalyzer and

high sensitivity DNA kit (Agilent #5067-4626). The quantity of sequenceable library for each

sample was determined using the KAPA Library Quantification Kit (Roche #KK4835) on the

7900HT real-time PCR (Applied Biosystems). Barcoded samples were pooled in groups of 6,

and pooled libraries were sequenced by the University of Pennsylvania Next-Generation

Sequencing Core, using one pool per lane, on an Illumina HiSeq 4000 utilizing 150 bp paired-

end sequencing. See Supplemental Methods for full details.

Bioinformatics

Demultiplexed sequencing data was cleaned and quality trimmed to a Phred quality

score of Q ≥ 20 using BBTools bbduk (https://sourceforge.net/projects/bbmap), and trimmed

read 1 data was aligned to the hg19 build of the human genome with Bowtie2-2.1.0 (12) using

end-to-end, very-sensitive alignment to generate a sequence alignment map (SAM) file for each

individual. ~10% of unaligned reads contained a poly(T) sequence after a genomic sequence

which made it unalignable. Assuming these poly(T) stretches likely corresponded to the 3'

poly(A) tail of the L1Hs sequence, they were trimmed, and the trimmed reads were aligned to

hg19 with Bowtie2-2.1.0 using end-to-end, very-sensitive alignment parameters to generate a

second SAM alignment file for each individual. SAM files were converted to BAM format, with

reads having alignment MapQ scores < 30 being discarded, using samtools-0.1.19 (13). Using

samtools-0.1.19, the two BAM files for each individual were combined. The resulting single BAM

files from all individuals were then merged using RG tags, a tag that allows for the combination

of reads from multiple BAM files, while still retaining the source (i.e. the individual) from which

each aligned read was generated, and the merged file was sorted and indexed. The merged

BAM file was stripped into BAM files for individual chromosomes using samtools-0.1.19, and the

chromosome specific BAM files were used as input for the custom python script

REBELseq_v1.0. REBELseq_v1.0 utilizes the sorted and indexed reads in the chromosome

specific BAM files to generate peaks of overlapping reads within a 150 base pair sliding window,

and does so with respect to DNA strand. Each peak corresponds to a putative L1

retrotransposon insertion, for which the peak's genomic coordinates, number of unique reads

per peak, number of reads per individual sample and average read alignment quality (mean

MapQ) are determined. The REBELseq_v1.0 output file was then further annotated using the

custom python script REBELannotate_v1.0 for L1Hs annotated in hg19 repeat masker and

L1Hs identified in the 1000 genomes data (14). REBELannotate_v1.0 utilizes a browser

extensible data (BED) formatted file to annotate genomic features of interest that overlap with or

occur within 500 base pairs downstream, with respect to strand, of the peak being annotated.

The REBELseq_v1.0 and REBELannotate_v1.0 custom scripts are based on work originally

described by Ewing and Kazazian (10), and all python scripts and necessary reference files

discussed in this manuscript are available online (https://github.com/BenReiner/REBELseq).

The raw sequencing data that was analyzed to support the findings of this manuscript are

available from the corresponding author upon request. See Supplemental Methods for full

details.

PCR Validations

Primers used in PCR experiments for method validation were designed using Primer3-2.2.3 in a

Perl script (makeprimers.pl), originally written by Adam Ewing (10), with an optimal Tm setting of

58°C (minimum 56°C and max 63°C), an optimal primer length of 24 nucleotides (minimum 21

and max 27) and the GC-clamp option set to 1. 25 µL reactions were constructed using 1x Go-

Taq colorless hotstart master mix (#M5133, Promega), 1 ng of gDNA for both an individual

predicted to have a particular L1Hs insertion or an individual predicted not to have the insertion,

and either 0.2 µM of the filled site and empty site (empty allele detection) or the filled site and

L1HsACA primer (filled allele detection). Samples were thermally cycled as 95 °C 2:30 min, then

10 cycles of touchdown PCR at 95 °C 0:30 min, 69-60 °C 0:30 min, 72 °C 1:30 min, then 25

cycles of 95 °C 0:30 min, 60 °C 0:30 min, 72 °C 1:30 min, then 72 °C 5:00 min, 4 °C hold. Go-

Taq green flexi buffer (#M8911, Promega) was added to all samples after amplification, and

samples were separated by 1.2% gel electrophoresis and visualized on a GelDoc XR+ (Bio-

Rad).

Statistics

Data are presented as mean ± standard deviation.

Results

REBELseq identifies L1Hs retrotransposon insertions

We performed REBELseq using DNA isolated from postmortem cortex NeuN+ nuclei

samples of 177 individuals and identified a total of 157,178 independent putative Ta subfamily

L1Hs insertions, with 1,050 annotating to hg19 known reference L1Hs elements (ref L1Hs) and

156,128 not annotating to hg19 known reference L1Hs (non-ref L1Hs). The ref L1Hs we

detected represented ~68% of all L1Hs annotated in hg19 repeat masker (1,050/1,544; see

Discussion) and were distributed across all chromosomes (Fig. 2a). Of the non-ref L1Hs, 432

annotated to known L1s in the 1000 genomes data (known non-ref L1Hs), while the other

155,696 are putative novel non-ref L1Hs. The ref L1Hs had an average mean MapQ of 38.82 ±

4.10. Having high confidence that the detected ref L1Hs were real, we used the ref L1Hs

average mean MapQ minus two standard deviations (MapQ ≥ 30.62) as a bioinformatic cutoff of

our data, after which we retained 974 ref L1Hs (92.76% of the 1,050 ref L1s), 430 known non-

ref L1Hs (99.54% of 432) and 127,976 putative novel non-ref L1Hs (82.20% of 155,696).

In the remaining known (ref L1Hs and known non-ref L1Hs) and putative novel L1Hs, we

next examined the average number of sequencing reads per person for a given L1Hs insertion

(Fig. 2b), and a clear difference between the known and novel L1Hs emerged, with 92.6% of

putative novel L1Hs having average reads < 10, and 87.7% of known L1Hs having average

reads ≥ 10 and 74.2% having average reads ≥ 100. Having confidence in the known L1Hs pool,

an average of ≥ 100 sequencing reads per person was used as a bioinformatic cutoff, which

retained 699 ref L1Hs (66.57%), 332 known non-ref L1Hs (76.85%) and 1,842 putative novel

non-ref L1Hs (1.18%).

<u>L1Hs insertion validation by PCR</u>

Having established bioinformatic cutoffs for our data, we next sought to experimentally

determine the proportion of bioinformatically-detected putative L1Hs insertions that were real

using PCR of the 3' insertion junction of the allele containing the insertion (filled allele) and the

corresponding genomic region on the 'empty' allele (see Fig. 3 for example image). Having

confidence that the known L1Hs insertions (ref L1Hs and known non-ref L1Hs) are real because

they were previously reported, we focused on the validation of the putative novel non-ref L1Hs

insertions predicted by REBELseq. The percentages of putative novel non-ref L1Hs insertions

that could be validated by average read number bin were experimentally determined, and these

data were used to calculate the number of detected putative novel non-ref L1Hs insertions that

we would expect to be true positives in each read bin. Using the number of putative novel non-

ref L1Hs insertions predicted to be true positives and the numbers of known L1Hs detected, we

calculated the probability of validating any detected L1Hs insertion per read bin and the

cumulative probability of validating any detected L1Hs insertions using the lower value of a read

bin as a minimum cutoff for the data (Table 1).

<u>Distribution of detected L1Hs</u>

The genomic distribution of known and novel L1Hs, per 10 MB genomic window, of our

highest quality L1Hs detection data was ascertained (≥ 1,000 average reads, Fig. 4a), and the

distribution of the number of individuals having a given known (Fig. 4b) or novel (Fig. 4c) L1Hs

for these data was assessed.

Discussion

While other methods for preparation of L1-targeted next generation sequencing libraries

exist (10, 11, 15-22), some are labor-intensive when scaled to a large number of individual

genomes, because of reliance on gel purification (10, 11) or because they require preparation of

multiple amplicon libraries per individual sample, due to the use of multiple hemi-degenerate

primers (10, 19). Some methods rely on the random shearing of DNA (18, 22, 23), a process

that requires specialized equipment and somewhat abundant DNA quantity, while others may

have reduced specificity for the Ta subfamily of L1Hs, which constitutes the majority of actively

replicating L1 retrotransposons in the human genome (16, 20, 21, 23). REBELseq was

designed as a high throughput alternative method to reliably detect the 3' flanking region of Ta

subfamily L1Hs elements with limited gDNA input. It should be noted that REBELseq was not

intended to determine whether an L1Hs insertion is full length or truncated or if ORF1 and ORF2

are intact, but merely to determine the presence or absence of the Ta subfamily L1Hs. After

identification of insertions of interest, these other aspects can be determined by long range PCR

and Sanger sequencing.

Other methods to produce L1-enriched libraries employing restriction enzyme digests of

gDNA as a starting point have been described (15). There is no set requirement for which

restriction enzyme to use in REBELseq other than that it generates blunt ends and cuts 5' of the

L1HsACA primer that identifies the 3' end of the Ta subfamily L1Hs elements. REBELseq

begins with restriction enzyme digestion of gDNA with HaeIII. This restriction enzyme was

chosen for the following reasons: 1) it creates blunt ends at 5'-GG|CC-3' sequences such that

polishing of cleaved ends is unnecessary; 2) it cuts the human genome to an average size of

342 ± 478 base pairs (NEBcutter) which is similar to the range of fragments generated by

sonication; 3) it does not cut within the 3' end of the L1Hs sequence targeted by our L1Hs Ta

subfamily-specific primers. One possible concern of using a single restriction enzyme digestion

in REBELseq library construction is the possibility that the restriction site for the enzyme occurs

immediately downstream from a L1 insertion, thus preventing the single primer extension into

the downstream gDNA (see Fig. 1). While this raises the possibility that using a single restriction

enzyme digestion during REBELseq library construction may not detect all L1 insertions, it does

not reduce the validity of L1 insertion that are detected. One possible solution to this concern

would be to perform restriction enzyme digestions with two or more enzymes that meet the

above criteria and pool the fragments before the single primer extension step.

Our method is specifically designed to leverage diagnostic nucleotides specific to the Ta

subfamily of L1Hs elements for differential amplification. It can easily be adapted to detect the

5' flanking region of full length L1Hs elements in the human genome, although diagnostic

nucleotides are limited within the 5' end of L1Hs for the Ta subfamily. Reliable primers targeting

the 5' end of Ta subfamily L1Hs and an enzyme other than HaeIII would be required for analysis

of full length L1Hs elements, as HaeIII makes a single digestion in the L1Hs sequence near the

5' end. REBELseq could also be utilized to differentially amplify and identify evolutionarily older

pre-Ta subfamily L1Hs insertions (by changing the last three nucleotides of the L1HsACA

primer to 'ACG') or any type of L1Hs element (by eliminating the last nucleotide of the L1HsACA

primer to end in 'AC'). Additionally, this method could be utilized to amplify and identify almost

any other genomic element with a specific pair of nested primers (to replace L1HsACA and

L1HsG).

The REBELseq technique is compatible with human gDNA purified from any tissue

source, including gDNA purified from leukocytes in saliva and blood. In the application of

REBELseq presented in this manuscript, 177 fresh frozen human postmortem prefrontal cortex

brain samples were utilized for purification of gDNA from NeuN+ neuronal nuclei. Utilizing gDNA

from NeuN+ neuronal nuclei allows for the identification of germline L1s, similar to using gDNA

from a peripheral source, and individual neuronally relevant somatic mutations occurring: only in

the brain (an L1 that retrotransposed early in neuronal development), in a limited number of

neurons (an L1 that retrotransposed in a neuronal progenitor cell, producing a cluster of

neurons harboring the insertion), or in a single neuron (an L1 that retrotransposed in the neuron

post-mitotically), results which could be verified by comparing the REBELseq results from

NeuN+ nuclei to those of a peripheral tissue. Using REBELseq to compare Ta subfamily L1Hs

from multiple brain regions would also allow for the identification of regionally specific L1Hs

insertions, possibly relevant in the context of neurological and psychiatric disease.

Using REBELseq, we identified 157,178 independent putative Ta subfamily L1Hs

insertions, with 1,050 annotating to reference L1Hs in hg19 repeat masker. The ref L1Hs

detected represent ~68% of all hg19 repeat masker L1Hs (Fig. 2a). The L1Hs elements

annotated in hg19 repeat masker represent both Ta subfamily L1Hs, which were specifically

targeted, and pre-Ta subfamily L1Hs. Thus, we did not expect to detect all L1Hs annotated in

hg19 repeat masker. Previous work using the genomes of 15 individuals aligned to hg18

showed that the pre-Ta subfamily of L1Hs represented ~36% of L1Hs insertions, with Ta

subfamily L1Hs elements composing the remainder (~64%) which closely approximates the

number we detected (10).

The bioinformatic cutoffs described in this manuscript were empirically derived by the

average mean MapQ score for the ref L1Hs, the average number of sequencing reads per

individual per insertion (Fig. 2b), and experimental confirmation of putative novel non-ref L1Hs

insertions (Table 1). We were unable to experimentally validate all putative novel L1Hs in any of

our average read bins. This is possibly due to the detected L1Hs insertion being sequencing

reads of a chimeric PCR product, or possibly because Ta subfamily L1Hs frequently

retrotranspose into repetitive sequences in the genome (e.g. the remnants of older repetitive

elements), making both their genomic alignment and PCR validation difficult. We believe this

suggests the proportion we were able to validate likely represents a minimum value, with

additional detected insertions being confirmable with more complex PCR methods. ~70% of all

known L1Hs averaged at least 100 sequencing reads per person, with our experimentally

determined probability of validating any detected L1Hs above 100 reads being ~59%.

Examining the data with ≥1,000 reads per person, we still see ~47% of all detected known L1Hs

and a probability of validating any detected L1Hs of almost 90%. These data suggest that novel

non-ref L1Hs having ≥1,000 reads per person include what are likely real polymorphic and

somatic L1Hs insertions.

When examining the genomic distribution of known and novel L1Hs having an average

of ≥1,000 sequencing reads per person (Fig. 4a), we see that both the known and novel L1Hs

are dispersed throughout the genome, suggesting that REBELseq is an unbiased whole

genome approach. Assessing the distribution of the number of individuals having either a known

(Fig. 4b) or novel (Fig. 4c) L1Hs insertion, we observe that both groups have L1 insertions

occurring throughout the range of possible values (i.e. number of individuals). With respect to

the known L1Hs, there appears to be a trimodal distribution with ~37% of L1Hs occurring in 45-

85 individuals and ~9% occurring in ≤ 3 and ≥ 170, meaning ~55% of insertions occur in only

~29% of the possible numbers of individuals. Our finding that a large proportion of known L1Hs

insertions occur in a small fraction of individuals disagrees with a previous report (10), but we

believe this may be due to their small sample size, thereby diminishing the likelihood that an

individual would not have a given insertion. ~70% of novel L1Hs occur in ≤ 7 people, while ~8%

of novel L1Hs occur in more than two thirds of individuals, suggesting that these L1Hs insertions

likely represent uncatalogued ref L1Hs occurring at common minor allele frequencies.

Taken together, we believe these data demonstrate that REBELseq reliably detects both

known and novel Ta subfamily L1Hs insertions, and that this technique could be a powerful

method for examining the association of the prevalence of L1Hs insertions in a broad range of

human disease.

Acknowledgements

The authors wish to thank those providing support for this work. BR is supported, in part, by a

2017 NARSAD Young Investigator Grant (#26634) from the Brain and Behavior Research

Foundation as the Patrick A. Coffer Investigator, funding for which was generously provided by

Ronald and Kathy Chandonais. BR was supported during part of this work by T32 MH014654

(PI is WB). This work was funded by National Institute of Health grants to WB (R21NS095756,

R01DA040972 & R01MH109260). RC was supported by K01DA036751. Additionally, the

authors wish to thank all the participants, and their families, who donated their brain tissue to

research for making these studies possible.

Conflict of Interests

The authors report no conflicts of interest.

1.   Richardson SR, Morell S, Faulkner GJ. L1 retrotransposons and somatic mosaicism in the brain.
Annual review of genetics. 2014;48:1-27.
2.   Beck CR, Collier P, Macfarlane C, Malig M, Kidd JM, Eichler EE, et al. LINE-1 retrotransposition
activity in human genomes. Cell. 2010;141(7):1159-70.
3.   Ostertag EM, Kazazian HH, Jr. Biology of mammalian L1 retrotransposons. Annual review of
genetics. 2001;35:501-38.
4.   Boissinot S, Chevret P, Furano AV. L1 (LINE-1) retrotransposon evolution and amplification in
recent human history. Molecular biology and evolution. 2000;17(6):915-28.
5.   Kazazian HH, Jr., Wong C, Youssoufian H, Scott AF, Phillips DG, Antonarakis SE. Haemophilia A
resulting from de novo insertion of L1 sequences represents a novel mechanism for mutation in man.
Nature. 1988;332(6160):164-6.
6.   Naufer MN, Callahan KE, Cook PR, Perez-Gonzalez CE, Williams MC, Furano AV. L1
retrotransposition requires rapid ORF1p oligomerization, a novel coiled coil-dependent property
conserved despite extensive remodeling. Nucleic acids research. 2016;44(1):281-93.
7.   Mathias SL, Scott AF, Kazazian HH, Jr., Boeke JD, Gabriel A. Reverse transcriptase encoded by a
human transposable element. Science. 1991;254(5039):1808-10.
8.   Feng Q, Moran JV, Kazazian HH, Jr., Boeke JD. Human L1 retrotransposon encodes a conserved
endonuclease required for retrotransposition. Cell. 1996;87(5):905-16.
9.   Jiang Y, Matevossian A, Huang HS, Straubhaar J, Akbarian S. Isolation of neuronal chromatin
from brain tissue. BMC Neurosci. 2008;9:42.
10.   Ewing AD, Kazazian HH, Jr. Whole-genome resequencing allows detection of many rare LINE-1
insertion alleles in humans. Genome research. 2011;21(6):985-90.
11.   Streva VA, Jordan VE, Linker S, Hedges DJ, Batzer MA, Deininger PL. Sequencing, identification
and mapping of primed L1 elements (SIMPLE) reveals significant variation in full length L1 elements
between individuals. BMC Genomics. 2015;16:220.
12.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9(4):357-9.
13.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
14.   Sudmant PH, Rausch T, Gardner EJ, Handsaker RE, Abyzov A, Huddleston J, et al. An integrated
map of structural variation in 2,504 human genomes. Nature. 2015;526(7571):75-81.
15.   Badge RM, Alisch RS, Moran JV. ATLAS: a system to selectively identify human-specific L1
insertions. Am J Hum Genet. 2003;72(4):823-38.
16.   Baillie JK, Barnett MW, Upton KR, Gerhardt DJ, Richmond TA, De Sapio F, et al. Somatic
retrotransposition alters the genetic landscape of the human brain. Nature. 2011;479(7374):534-7.
17.   Costello M, Fleharty M, Abreu J, Farjoun Y, Ferriera S, Holmes L, et al. Characterization and
remediation of sample index swaps by non-redundant dual indexing on massively parallel sequencing
platforms. BMC Genomics. 2018;19(1):332.
18.   Erwin JA, Paquola AC, Singer T, Gallina I, Novotny M, Quayle C, et al. L1-associated genomic
regions are deleted in somatic cells of the healthy human brain. Nat Neurosci. 2016;19(12):1583-91.
19.   Evrony GD, Cai X, Lee E, Hills LB, Elhosary PC, Lehmann HS, et al. Single-neuron sequencing
analysis of L1 retrotransposition and somatic mutation in the human brain. Cell. 2012;151(3):483-96.
20.   Iskow RC, McCabe MT, Mills RE, Torene S, Pittard WS, Neuwald AF, et al. Natural mutagenesis of
human genomes by endogenous retrotransposons. Cell. 2010;141(7):1253-61.
21.   Kvikstad EM, Piazza P, Taylor JC, Lunter G. A high throughput screen for active human
transposable elements. BMC Genomics. 2018;19(1):115.
22.   Upton KR, Gerhardt DJ, Jesuadian JS, Richardson SR, Sanchez-Luque FJ, Bodea GO, et al.
Ubiquitous L1 mosaicism in hippocampal neurons. Cell. 2015;161(2):228-39.

23.   Zhao B, Wu Q, Ye AY, Guo J, Zheng X, Yang X, et al. Somatic LINE-1 retrotransposition in cortical
neurons and non-brain tissues of Rett patients and healthy individuals. PLoS Genet.
2019;15(4):e1008043.

Figure legends

Figure 1: Schematic of the construction of Ta subfamily enriched L1Hs sequencing libraries.

gDNA isolated from NeuN+ nuclei was enzymatically digested with HaeIII to fragment the

genome. A single primer extension using Ta subfamily specific L1HsACA primer extends the 3'

end of the L1 sequence into the downstream gDNA. The 3' 'A' overhang from the single primer

extension is ligated to a custom T-linker, and primary PCR amplifies the construct using

L1HsACA and T-linker specific primers. Hemi-nested secondary PCR using the L1Hs specific

L1HsG primer and T-linker primer reduces the length of the L1 sequence carried forward and

adds a sequencing adapter to the L1 end. Tertiary PCR uses 5' overhang primers to add

barcodes to the L1 end and Illumina flow cell adapters to both ends of the amplicon.

Figure 2: Detection levels of Ta subfamily L1Hs. **a** The number of ref L1Hs detected per

chromosome versus the number of L1Hs annotated in hg19 repeat masker. Overall, we

detected ~68% of L1Hs annotated in hg19 repeat masker, which aligns with our expectations

based on the literature (see discussion). **b** Average number of sequencing reads per person for

a given L1 insertion. Data was binned to show contrast between the average number of

sequencing reads per person seen for known and putative novel L1Hs insertions. An average

number of sequencing reads per person ≥ 100 was used as a bioinformatic cutoff.

Figure 3: Example image of a successful and unsuccessful confirmatory PCR. PCR

experiments were conducted to determine the proportion of putative novel non-ref L1Hs

insertions detected by REBELseq that could be independently validated. For each insertion, a

person predicted to have the L1Hs insertion (+) and a person not predicted to have the insertion

(-) were used to amplify the genomic region purported to contain the L1Hs insertion (Filled site,

F) and the same genomic region if it did not contain the insertion (Empty site, E). Insertion #1 is

a positive confirmation of the results predicted by REBELseq, while Insertion #2 is a negative

confirmation.

<u>Figure 4:</u> Distributions of high confidence L1Hs insertions. **a** The genomic distribution of known

L1Hs (above the chromosome numbers) and novel L1Hs (below the chromosome numbers) per

10 MB window of each chromosome. Alternating color pattern and labeled central blocks

represent the different chromosomes. Known and novel L1Hs are distributed throughout the

genome, suggesting REBELseq is an unbiased whole genome approach. **b** Number of

individuals sharing a known L1Hs insertion. The number of known L1Hs insertions shared by

different numbers of individuals shows a trimodal distribution. While some ref L1Hs occur with a

rate of affected individuals approaching 1.0, the other local maxima are focused at few or less

than half of surveyed individuals. This demonstrates that known L1Hs should be considered

polymorphic in nature, rather than ubiquitous, in the human genome. **c** Number of individuals

sharing a novel L1Hs insertion. The number of novel L1Hs insertions shared by different

numbers of individuals shows a right skewed distribution. While some novel L1Hs were detected

in most individuals, most novel L1Hs were detected in one or a few individuals.

<u>Table 1:</u> Table describing the rates of putative L1Hs insertion validation by average number of

reads. The percentage of successfully validated L1Hs per read bin was used to calculate the

number of real novel non-ref L1Hs. The percent validated per bin was calculated as the sum of

real novel non-ref L1Hs, number of detected known non-ref L1Hs and number of detected ref

L1Hs, divided by the sum of the number of detected novel non-ref L1Hs, number of detected

known non-ref L1Hs and number of detected ref L1Hs. The cumulative probability of being

validated was calculated similarly, we the bottom of the lowest included read bin serving as a

lower cutoff.

440   **Figure 1**

Human gDNA extracted from NeuN+ neuronal nuclei

Digestion of gDNA with HaeIII in presence of rSAP

Single primer extension of L1

Ligation to double stranded T-linker

Primary PCR

Secondary PCR

Tertiary PCR

■ Human DNA   ■ T_a subfamily L1Hs
■ Double stranded T-linker
■ L1 targeted sequencing adapter
■ T-linker targeted sequencing adapter
■ Illumina flow cell adapters   ■ Barcode

Figure 2

	chr 1	chr 2	chr 3	chr 4	chr 5	chr 6	chr 7	chr 8	chr 9	chr 10	chr 11	chr 12	chr 13	chr 14	chr 15	chr 16	chr 17	chr 18	chr 19	chr 20	chr 21	chr 22	chr X	chr Y
hg19 ref L1Hs	119	104	124	151	109	105	83	69	57	48	69	55	38	46	30	31	13	60	12	22	13	6	129	35
Detected ref L1Hs	84	81	82	92	73	66	58	47	37	39	44	44	29	35	24	23	9	46	6	13	9	5	85	19
Percent detected	71	78	66	61	67	63	70	68	65	81	64	80	76	76	80	74	69	77	50	59	69	83	66	54

	1-9	10-24	25-49	50-99	100-249	250-499	500-999	≥1000
% known L1Hs	12.26	4.47	4.61	4.47	7.86	6.85	9.59	49.89
% novel L1Hs	92.59	3.35	1.63	0.99	0.69	0.25	0.15	0.36

Average reads per person

458 **Figure 3**

	Insertion #1				Insertion #2			
L1 predicted	**+**		**-**		**+**		**-**	
Allele	E	F	E	F	E	F	E	F

476   **Figure 4**

486 **Table 1**

Avg. # of reads	novel non-ref L1Hs					known non-ref L1Hs	ref L1Hs	% validated per read bin	cumulative probability of being validated (%)
	# Successful	# Attempted	% Successful	# detected	# real	# detected	# detected		
≥1,000	32	44	72.7	460	334	227	465	89.1	89.1
500-999	16	41	39.0	186	73	43	90	64.6	83.8
250-499	13	40	32.5	320	104	28	67	48.0	75.9
100-249	6	36	16.7	876	146	34	77	26.0	58.8

490