Eukaryotic virus composition can predict the efficiency of carbon export in the

global ocean

Hiroto Kaneko[1,+], Romain Blanc-Mathieu[1,2,+], Hisashi Endo[1], Samuel Chaffron[3,4],

Tom O. Delmont[4,5], Morgan Gaia[4,5], Nicolas Henry[6], Rodrigo Hernández-Velázquez[1],

Canh Hao Nguyen[1], Hiroshi Mamitsuka[1], Patrick Forterre[7], Olivier Jaillon[4,5],

Colomban de Vargas[6], Matthew B. Sullivan[8], Curtis A. Suttle[9], Lionel Guidi[10] and

Hiroyuki Ogata[1,*]

+ Equal contribution

* Corresponding author

Affiliations:

1: Bioinformatics Center, Institute for Chemical Research, Kyoto University,

Gokasho, Uji, Kyoto 611-0011, Japan

2: Laboratoire de Physiologie Cellulaire & Végétale, CEA, Univ. Grenoble Alpes,

CNRS, INRA, IRIG, Grenoble, France.

3: Université de Nantes, CNRS UMR 6004, LS2N, F-44000 Nantes, France.

4: Research Federation (FR2022) *Tara* Oceans GO-SEE, Paris, France

5: Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, 91000

Evry, France

6: Sorbonne Universités, CNRS, Laboratoire Adaptation et Diversité en Milieu Marin,

Station Biologique de Roscoff, 29680 Roscoff, France

7: Institut Pasteur, Department of Microbiology, 25 rue du Docteur Roux, 75015,

Paris, France

8: Department of Microbiology and Department of Civil, Environmental and Geodetic

Engineering, Ohio State University, Columbus, OH, United States of America

9: Departments of Earth, Ocean & Atmospheric Sciences, Microbiology &

Immunology, and Botany, and the Institute for the Oceans and Fisheries, University

of British Columbia, Vancouver, BC, V6T 1Z4, Canada

10: Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefanche, LOV,

F-06230 Villefranche-sur-mer, France

Summary

The biological carbon pump, in which carbon fixed by photosynthesis is exported to the deep ocean through sinking, is a major process in Earth's carbon cycle. The proportion of primary production that is exported is termed the carbon export efficiency (CEE). Based on in-lab or regional scale observations, viruses were previously suggested to affect the CEE (i.e., viral "shunt" and "shuttle"). In this study, we tested associations between viral community composition and CEE measured at a global scale. A regression model based on relative abundance of viral marker genes explained 67% of the variation in CEE. Viruses with high importance in the model were predicted to infect ecologically important hosts. These results are consistent with the view that the viral shunt and shuttle functions at a large scale and further imply that viruses likely act in this process in a way dependent on their hosts and ecosystem dynamics.

Introduction

A major process in the global cycling of carbon is the oceanic biological carbon pump (BCP), an organism-driven process by which atmospheric carbon (*i.e.,* CO_2) is transferred and sequestered to the ocean interior and seafloor for periods ranging from centuries to hundreds of millions of years. Between 15% and 20% of net primary production (NPP) is exported out of the euphotic zone, with 0.3% of fixed carbon reaching the seafloor annually (Zhang et al., 2018). However, there is wide variation in estimates of the proportion of primary production in the surface ocean that is exported to depth, ranging from 1% in the tropical Pacific to 35-45% during the North Atlantic bloom (Buesseler and Boyd, 2009). As outlined below, many factors affect the BCP.

Of planktonic organisms living in the upper layer of the ocean, diatoms (Tréguer et al., 2018) and zooplankton (Turner, 2015) have been identified as important contributors to the BCP in nutrient-replete oceanic regions. In the oligotrophic ocean, cyanobacteria, collodarians (Lomas and Moran, 2011), diatoms (Agusti et al., 2015; Karl et al., 2012; Leblanc et al., 2018), and other small (pico- to nano-) plankton (Lomas and Moran, 2011) have been implicated in the BCP. Sediment trap studies suggest that ballasted aggregates of plankton with biogenic minerals contribute to carbon export to the deep sea (Iversen and Ploug, 2010; Klaas and Archer, 2002). The BCP comprises three processes: carbon fixation, export, and remineralization. As these processes are governed by complex interactions between numerous members of planktonic communities (Zhang et al., 2018), the BCP is expected to involve various organisms, including viruses (Zimmerman et al., 2019a).

Viruses have been suggested to regulate the efficiency of the BCP. Lysis of host cells by viruses releases cellular material in the form of dissolved organic matter

(DOM), which fuels the microbial loop and enhances respiration and secondary

production (Gobler et al., 1997; Weitz et al., 2015). This process, coined "viral shunt

(Wilhelm and Suttle, 1999)", can reduce the carbon export efficiency (CEE) because

it increases the retention of nutrients and carbon in the euphotic zone and prevents

their transfer to higher trophic levels as well as their export from the euphotic zone to

the deep sea (Fuhrman, 1999; Weitz et al., 2015). However, an alternative process is

also considered, in which viruses contribute to the vertical carbon export (Weinbauer,

2004). For instance, a theoretical study proposed that the CEE increases if viral lysis

augments the ratio of exported carbon relative to the primary production-limiting

nutrients (nitrogen and phosphorous) (Suttle, 2007). Laboratory experimental studies

reported that cells infected with viruses form larger particles (Peduzzi and Weinbauer,

1993; Yamada et al., 2018), can sink faster (Lawrence and Suttle, 2004), and can lead

to preferential grazing by heterotrophic protists (Evans and Wilson, 2008) and/or to

higher growth of grazers (Goode et al., 2019). This process termed "viral shuttle

(Sullivan et al., 2017)" is supported by several field studies that reported association

of viruses with sinking material. Viruses were observed in sinking material in the

North Atlantic Ocean (Proctor and Fuhrman, 1991) and sediment of coastal waters

where algal blooms occur (Lawrence et al., 2002; Tomaru et al., 2007, 2011). In

addition, vertical transport of bacterial viruses between photic and aphotic zones was

observed in the Pacific Ocean (Hurwitz et al., 2015) and in *Tara* Oceans virome data

(Brum et al., 2015). A systematic analysis of large-scale omics data from oligotrophic

oceanic regions revealed a strong positive association between carbon flux and

bacterial dsDNA viruses (*i.e.*, cyanophages), which were previously unrecognized as

possible contributors to the BCP (Guidi et al., 2016). More recently, viral infection of

blooms of the photosynthetic eukaryote *Emiliania huxleyi* in the North Atlantic were

found to be accompanied by particle aggregation and greater downward vertical flux

of carbon, with the highest export during the early stage of viral infection (Laber et

al., 2018). These studies raise the question of the overall impact of viruses infecting

eukaryotes on oceanic carbon cycling and export. Given the significant contributions

of eukaryotic plankton to ocean biomass and net production (Hirata et al., 2011; Li,

1995) and their observed predominance over prokaryotes in sinking materials of

Sargasso Sea oligotrophic surface waters (Fawcett et al., 2011; Lomas and Moran,

2011), various lineages of eukaryotic viruses may be responsible for a substantial part

of the variation in carbon export across oceanic regions.

If the "viral shunt" and "shuttle" processes function at a global scale and if

these involve specific viruses, we expect to detect a statistical association between

viral community composition and CEE in a large scale omics data. To our knowledge,

such an association has never been investigated. Although this test per se does not

prove that viruses regulate CEE, we consider the association is worth being tested

because such an association is a necessary condition for the global model of viral

shunt and shuttle and, under its absence, we will have to reconsider the model. Deep

sequencing of planktonic community DNA and RNA, as carried out in *Tara* Oceans,

has enabled the identification of marker genes of major viral groups infecting

eukaryotes (Hingamp et al., 2013; Carradec et al., 2018; Culley, 2018). To examine

the association between viral community composition and CEE, we thus used the

comprehensive organismal dataset from the *Tara* Oceans expedition (Carradec et al.,

2018; Sunagawa et al., 2015), as well as related measurements of carbon export

estimated from particle concentrations and size distributions observed *in situ* (Guidi et

al., 2016).

In the present study, we identified several hundred marker-gene sequences of

nucleocytoplasmic large DNA viruses (NCLDVs) in metagenomes of 0.2–3 μm size

fraction. We also identified RNA and ssDNA viruses in metatranscriptomes of four

eukaryotic size fractions spanning 0.8 to 2,000 μm. The resulting profiles of viral

distributions were compared with an image-based measure of carbon export efficiency

(CEE), which is defined as the ratio of the carbon flux at depth to the carbon flux at

surface.

Results and Discussion

Detection of diverse eukaryotic viruses in *Tara* Oceans gene catalogs

We used profile hidden Markov model-based homology searches to identify marker-

gene sequences of eukaryotic viruses in two ocean gene catalogs. These catalogs were

previously constructed from environmental shotgun sequence data of samples

collected during the *Tara* Oceans expedition. The first catalog, the Ocean Microbial

Reference Gene Catalog (OM-RGC), contains 40 million non-redundant genes

predicted from the assemblies of *Tara* Oceans viral and microbial metagenomes

(Sunagawa et al., 2015). We searched this catalog for NCLDV DNA polymerase

family B (PolB) genes, as dsDNA viruses may be present in microbial metagenomes

because large virions (> 0.2 μm) have been retained on the filter or because viral

genomes actively replicating or latent within picoeukaryotic cells have been captured.

The second gene catalog, the Marine Atlas of *Tara* Oceans Unigenes (MATOU),

contains 116 million non-redundant genes derived from metatranscriptomes of single-

cell microeukaryotes and small multicellular zooplankton (Carradec et al., 2018). We

searched this catalog for NCLDV PolB genes, RNA-dependent RNA polymerase

(RdRP) genes of RNA viruses, and replication-associated protein (Rep) genes of

ssDNA viruses, since transcripts of viruses actively infecting their hosts, as well as

genomes of RNA viruses, have been captured in this catalog.

We identified 3,874 NCLDV PolB sequences (3,486 in metagenomes and 388

in metatranscriptomes), 975 RNA virus RdRP sequences, and 299 ssDNA virus Rep

sequences (Table 1). These sequences correspond to operational taxonomic units

(OTUs) at a 95% identity threshold. All except 17 of the NCLDV PolBs from

metagenomes were assigned to the families *Mimiviridae* ($n = 2{,}923$),

Phycodnaviridae ($n = 348$), and *Iridoviridae* ($n = 198$) (Table 1). The larger numbers

of PolB sequences assigned to *Mimiviridae* and *Phycodnaviridae* compared with other

NCLDV families are consistent with a previous observation based on a smaller

dataset (Hingamp et al., 2013). The divergence between these environmental

sequences and reference sequences from known viral genomes was greater in

Mimiviridae than in *Phycodnaviridae* (Figure 1a, S1a and S2). Within *Mimiviridae,*

83% of the sequences were most similar to those from algae-infecting *Mimivirus*

relatives. Among the sequences classified in *Phycodnaviridae,* 93% were most similar

to those in *Prasinovirus*, whereas 6% were closest to *Yellowstone lake phycodnavirus*,

which is closely related to *Prasinovirus*. Prasinoviruses are possibly over-represented

in the metagenomes because the 0.2 to 3 μm size fraction selects their picoeukaryotic

hosts. RdRP sequences were assigned mostly to the order *Picornavirales* ($n = 325$),

followed by the families *Partitiviridae* ($n = 131$), *Narnaviridae* ($n = 95$),

Tombusviridae ($n = 45$), and *Virgaviridae* ($n = 33$) (Table 1), with most sequences

being distant (30% to 40% amino acid identity) from reference viruses (Figures 1b,

S1b and S3). These results are consistent with previous studies on the diversity of

marine RNA viruses, in which RNA virus sequences were found to correspond to

diverse positive-polarity ssRNA and dsRNA viruses distantly related to well-characterized viruses (Culley, 2018). *Picornavirales* may be over-represented in the metatranscriptomes because of the polyadenylated RNA selection. The majority ($n = 201$) of Rep sequences were annotated as *Circoviridae*, known to infect animals, which is consistent with a previous report (Wang et al., 2018). Only eight were annotated as plant ssDNA viruses (families *Nanovoridae* and *Gemniviridae*) (Table 1). Most of these environmental sequences are distant (40% to 50% amino acid identity) from reference sequences (Figures 1c, S1c and S4). Additional 388 NCLDV PolBs were detected in the metranscriptomes. The average cosmopolitanism (number of samples where an OTU was observed by at least two reads) for PolBs in metagenomes was 23 samples against 2.9 for metranscriptome-derived PolB sequences, 5.5 for Reps, and 5.8 for RdRPs. Within metatranscriptomes, the average gene-length normalized read counts for PolBs were respectively ten and three times lower than those of RdRPs and Reps. Therefore, PolBs from metatranscriptomes were not further used in our study.

Composition of eukaryotic viruses can explain the variation of carbon export efficiency

Among the PolB, RdRP, and Rep sequences identified in the *Tara* Oceans gene catalogs, 38%, 18%, and 11% (total = 1,523 sequences), respectively, were present in at least five samples and had matching carbon export measurement data (Table 1). We used the relative abundance (defined as the centered log-ratio transformed gene-length normalized read count) profiles of these 1,523 marker-gene sequences at 59 sampling sites in the photic zone of 39 *Tara* Oceans stations (Figure 2) to test for association between their composition and a measure of carbon export efficiency (CEE, see Transparent Methods, Figure S5). A partial least squares (PLS) regression model

explained 67% (coefficient of determination $R^2 = 67\%$) of the variation in CEE with a

Pearson correlation coefficient of 0.84 between observed and predicted values. This

correlation was confirmed to be statistically significant by permutation test ($P < 1 \times$

10^{-4}) (Figure 3a).

We also tested for their association with estimates of carbon export flux at 150

meters (CE_{150}) and NPP. PLS regressions explained 54% and 64% of the variation in

CE_{150} and NPP with Pearson correlation coefficients between observed and predicted

values of 0.74 (permutation test, $P < 1 \times 10^{-4}$) and 0.80 (permutation test, $P < 1 \times$

10^{-4}), respectively (Figure S6). In these three PLS regression models, 83, 86, and 97

viruses were considered to be key predictors (*i.e.*, Variable Importance in the

Projection [VIP] score > 2) of CEE, CE_{150}, and NPP, respectively. PLS models for

NPP and CE_{150} shared a larger number of predictors (52 viruses) compared to the PLS

models for NPP and CEE (seven viruses) (two proportion Z-test, $P = 4.14 \times 10^{-12}$).

Consistent with this observation, CE_{150} was correlated with NPP (Pearson's $r = 0.77$;

parametric test, $P < 1 \times 10^{-12}$). This result implies that the magnitude of export in the

analyzed samples was partly constrained by primary productivity. However, CEE was

not correlated with NPP ($r = 0.16$; parametric test, $P = 0.2$) or CE_{150} ($r = 0.002$;

parametric test, $P = 0.99$). Thus, as expected, primary productivity was not a major

driver for the efficiency of carbon export.

The 83 viruses (5% of the viruses included in our analysis) that were

associated with CEE with a VIP score > 2 are considered to be important predictors of

CEE in the PLS regression (Figure 3b, Supplemental Data 1), and these viruses are

hereafter referred to as VIPs (Viruses Important in the Prediction). Fifty-eight VIPs

had positive regression coefficient, and 25 had negative regression coefficient in the

prediction (Figure 3b). Most of the positively associated VIPs showed high relative

abundance in the Mediterranean Sea and in the Indian Ocean where CEE tends to be

high compared with other oceanic regions (Figure 4). Among them, 15 (red labels in

Figure 4) also had high relative abundance in samples from other oceanic regions,

showing that these viruses are associated with CEE at a global scale. In contrast,

negatively associated VIPs tend to have higher relative abundance in the Atlantic

Ocean and the Southern Pacific Ocean where CEE is comparatively lower. In the

following sections, we investigate potential hosts of the VIPs in order to interpret the

statistical association between viral community composition and CEE in the light of

previous observations in the literature.

Viruses correlated with CEE infect ecologically important hosts

Most of the VIPs (77 of 83) belong to *Mimiviridae* ($n = 34$ with 25 positive

VIPs and nine negative VIPs), *Phycodnaviridae* ($n = 24$ with 18 positive VIPs and six

negative VIPs), and ssRNA viruses of the order *Picornavirales* ($n = 19$ with 13

positive VIPs and six negative VIPs) (Figure 3b, Table S1). All the phycodnavirus

VIPs were most closely related to prasinoviruses infecting Mamiellales, with amino

acid sequence percent identities to reference sequences ranging between 35% and

95%. The six remaining VIPs were two NCLDVs of the family *Iridoviridae*

negatively associated with CEE, three RNA viruses (two ssRNA viruses of the family

Hepeviridae negatively associated with CEE and one dsRNA virus of the family

Partitiviridae positively associated with CEE), and one ssDNA virus of the family

Circoviridae positively associated with CEE.

Host information may help understand the relationship between these VIPs

and CEE. We performed genomic context analysis for PolB VIPs and phylogeny-

guided network-based host prediction for PolB and RdRP to infer putative virus–host

relationships (see Transparent Methods).

Taxonomic analysis of genes predicted in 10 metagenome-assembled genomes

(MAGs) from the eukaryotic size fractions and 65 genome fragments (contigs)

assembled from the prokaryotic size fraction encoding VIP PolBs further confirmed

their identity as *Mimiviridae* or *Phycodnaviridae* (Figure S7). The size of MAGs

ranged between 30 kbp and 440 kbp with an average of 210 kbp (Table S2). The

presence of genes with high sequence similarities to cellular genes in a viral genome

is suggestive of a virus–host relationship (Monier et al., 2009; Yoshikawa et al.,

2019). Two closely related *Mimiviridae* VIPs, PolB 000079111 (positively associated

with CEE) and PolB 000079078 (negatively associated with CEE), were

phylogenetically close to the pelagophyte virus *Aureococcus anophagefferens virus*

(AaV). One MAG (268 kbp in size) corresponding to PolB 000079111 encoded seven

genes showing high similarities to genes from Pelagophyceae, and another MAG (382

kbp in size), corresponding to PolB 000079078, encoded five genes similar to genes

from Pelagophyceae. All but one of these 12 genes was encoded on a genome

fragment containing genes annotated as viral, including five NCLDV core genes

(Supplemental Data 2), excluding the possibility of contamination in these MAGs.

Two closely related *Phycodnaviridae* VIPs, PolB 001064263 and 010288541, were

positively associated with CEE. Both of these PolBs correspond to a MAG (134 kbp

in size) encoding one gene likely derived from Mamiellales. The genomic fragment

harboring this cellular gene was found to encode 10 genes annotated as viral

(Supplemental Data 2).

We conducted a phylogeny-guided, network-based host prediction analysis for

Mimiviridae, *Phycodnaviridae*, and *Picornavirales* (Figures S8 and S9). Only a subset

of the VIPs was included in this analysis because we kept the most reliable sequences

to obtain a well-resolved tree topology. Within the *Prasinovirus* clade, which

contained thirteen VIPs (nine positive and four negative), seven different eukaryotic

orders were detected as predicted host groups for 10 nodes in the tree. Mamiellales,

the only known host group of prasinoviruses, was detected at eight nodes (five of

them had no parent-to-child relationships), whereas the other six eukaryotic orders

were found at only one node (or two in the case of Eutreptiales) (Figure S8). The

order Mamiellales includes three genera (*Micromonas*, *Ostreococcus*, and

Bathycoccus), which are bacterial-sized green microalgae common in coastal and

oceanic environments and are considered to be influential actors in oceanic systems

(Monier et al., 2016). Various prasinoviruses (fourteen with available genome

sequences) have been isolated from the three genera.

Within the family *Mimiviridae*, which contains fifteen VIPs (10 positive and

five negative), twelve different orders were predicted as putative host groups (Figure

S8). Collodaria was detected at 15 nodes (two of them had no parent-to-child

relationships), and Prymnesiales at six nodes (three of them had no parent-to-child

relationships), whereas all other orders were present at a maximum of one node each

with no parent-to-child relationships. The nodes enriched for Prymnesiales and

Collodaria fell within a monophyletic clade (marked by a red arrow in Figure S8)

containing four reference haptophyte viruses infecting Prymnesiales and two

reference haptophyte viruses infecting Phaeocystales. Therefore, the environmental

PolB sequences in this *Mimiviridae* clade (including five positive VIPs and one

negative VIP) are predicted to infect Prymnesiales or related haptophytes. The

detection of Collodaria may be the result of indirect associations that reflect a

symbiotic relationship with Prymnesiales, as some acantharians, evolutionarily related

to the Collodaria, are known to host Prymnesiales species (Mars Brisbin et al., 2018).

Known species of Prymnesiales and Phaeocystales have organic scales, except one

Prymnesiales species, *Prymnesium neolepis*, which bears siliceous scales (Yoshida et

al., 2006). Some species can form blooms and colonies. Previous studies revealed the

existence of diverse and abundant noncalcifying picohaptophytes in open oceans

(Endo et al., 2018; Liu et al., 2009). Haptophytes as a whole have been estimated to

contribute from 30% to 50% of the total photosynthetic standing stock across the

world ocean (Hirata et al., 2011; Liu et al., 2009). They constitute an important

mixotrophic group in oligotrophic waters (Endo et al., 2018), and mixotrophy is

proposed to increase vertical carbon flux by enabling the uptake of organic forms of

nutrients (Ward and Follows, 2016). Clear host prediction was not made for the other

nine *Mimiviridae* VIPs shown in the phylogenetic tree. Three VIPs (two positive and

one negative) in the tree were relatives of AaV. One negatively associated VIP was a

relative of *Cafeteria roenbergensis virus* infecting a heterotrophic protist. The five

remaining *Mimiviridae* VIPs are very distant from any known *Mimiviridae*.

Sixteen *Picornavirales* VIPs (eleven positive and five negative) were included

in the phylogeny-guided, network-based host prediction analysis (Figure 9). Nine

(seven positive and two negative) were grouped within *Dicistroviridae* (known to

infect insects) and may therefore infect marine arthropods such as copepods, the most

ubiquitous and abundant mesozooplankton groups involved in carbon export (Turner,

2015). Three other *Picornavirales* VIPs were placed within a clade containing known

bacillarnaviruses. Two of them (35179764 and 33049404) were positively associated

with CEE and had diatoms of the order Chaetocerotales as a predicted host group. The

third one (107558617) was negatively associated with CEE and distant from other

bacillarnaviruses, and had no host prediction. Diatoms have been globally observed in

the deep sea (Agusti et al., 2015; Leblanc et al., 2018) and identified as important

contributors of the biological carbon pump (Tréguer et al., 2018). One positively

associated VIP (32150309) was in a clade containing *Aurantiochytrium single-stranded RNA virus* (AsRNAV), infecting a marine fungoid protist thought to be an important decomposer (Takao et al., 2005). The last three *Picornavirales* VIPs (59731273, 49554577, and 36496887) had no predicted host and were too distant from known *Picornavirales* to speculate about their putative host group.

Outside *Picornavirales*, three RNA virus VIPs (two *Hepeviridae*, negatively associated, and one *Partitiviridae*, positively associated) were identified, for which no reliable host inferences were made by sequence similarity. Known *Hepeviridae* infect metazoans, and known *Partitiviridae* infect fungi and plants. The two *Hepeviridae*-like viruses were most closely related to viruses identified in the transcriptomes of mollusks (amino acid identities of 48% for 42335229 and 43% for 77677770) (Shi et al., 2016). The *Partitiviridae*-like VIP (35713768) was most closely related to a fungal virus, *Penicillium stoloniferum virus S* (49% amino acid identity).

One ssDNA virus VIP (38177659) was positively associated with CEE. It was annotated as a *Circoviridae*, although it groups with other environmental sequences as an outgroup of known *Circoviridae*. This VIP was connected with copepod, mollusk, and Collodaria OTUs in the co-occurrence network but no enrichment of predicted host groups was detected for its clade. *Circoviridae*-like viruses are known to infect copepods (Dunlap et al., 2013) and have been reported to associate with mollusks (Dayaram et al., 2015), but none have been reported for Collodaria.

Overall, we could infer hosts for 36 VIPs (Tables S3 and S4). Most of the predicted hosts are known to be ecologically important as primary producers (Mamiellales, Prymnesiales, Pelagophyceae, and diatoms) or grazers (copepods). Of these, diatoms and copepods are well known as important contributors to the BCP but others (*i.e.*, Mamiellales, Prymnesiales, Pelagophyceae) have not been recognized as

341 major contributors to the BCP. Our analysis also revealed that positive and negative

342 VIPs are not separated in either the viral or host phylogenies.

Viruses positively correlated with CEE tend to interact with silicified organisms

The phylogeny-guided, network-based host prediction analysis correctly predicted

known virus–host relationships (for viruses infecting Mamiellales, Prymnesiales, and

Chaetocerotales) using our large dataset, despite the reported limitations of these co-

occurrence network-based approaches (Coenen and Weitz, 2018). This result

prompted us to further exploit the species co-occurrence networks (Table S5) to

investigate functional differences between the eukaryotic organisms predicted to

interact with positive VIPs, negative VIPs, and viruses less important for prediction of

CEE (VIP score < 2) (non-VIPs). Positive VIPs had a greater proportion of

connections with silicified eukaryotes ($Q = 0.001$), but not with chloroplast-bearing

eukaryotes ($Q = 0.16$) nor calcifying eukaryotes ($Q = 1$), compared to non-VIPs

(Table S6). No functional differences were observed between negative VIPs and non-

VIPs viruses (Table S6) or positive VIPs (Table S7).

Multifarious ways viruses affect the fate of carbon

Our analysis revealed that eukaryotic virus composition was able to predict CEE in

the global sunlit ocean and 83 out of the 1,523 viruses had a high importance in the

predictive model. This association is not a proof that the viruses are the cause of the

variation of CEE. For example, a virus may be found to be associated with CEE if its

host affects CEE regardless of viral infection. This would be the case especially if

latent/persistent viruses are widespread and abundant in phytoplankton (Goic and

Saleh, 2012). Organisms that preferentially grow in marine snow (Bochdansky et al.,

2017) may bring associations between viruses infecting those organisms and CEE;

this could be the case for the AsRNAV-related VIP that we identified. Alternatively,

the observed associations between VIPs and CEE may reflect a more direct causal

relationship, which we attempt to explore in light of the large body of literature on the

mechanisms by which viruses impact the fate of carbon in the oceans.

370   Among the 83 VIPs, 58 were positively associated with CEE. Such a positive

association is expected from the "viral shuttle" model, which states that viral activity

could facilitate carbon export to the deep ocean (Fuhrman, 1999; Sullivan et al., 2017;

Weinbauer, 2004) because viral infection can facilitate cell sinking (Lawrence and

Suttle, 2004) and increase the sizes of particles (Peduzzi and Weinbauer, 1993;

Yamada et al., 2018); for instance, a virus may induce secretion of sticky material that

contributes to cell/particle aggregation, such as transparent exopolymeric particles

(TEP) (Nissimov et al., 2018). The data we used to estimate carbon export are based

on the particle size distribution and concentration, and do not convey information

regarding the aggregation status of particles. Therefore, we cannot directly test for a

relation between viruses and aggregation at the sampling sites. Nonetheless, we found

that CEE (*i.e.*, $CE_{deep}/CE_{surface}$) increased with the change of particles size from

surface to deep ($\rho = 0.42$, $P = 8 \times 10^{-9}$) (Figure S10). This positive correlation may

reflect an elevated level of aggregation (either enhanced by viral activity or not) in

places where CEE is high, although it could be also due to the presence of large

organisms at depth.

386   Greater aggregate sinking along with higher particulate carbon fluxes was

observed in North Atlantic blooms of *Emiliania huxleyi* that were infected early by

the virus EhV, compared with late-infected blooms (Laber et al., 2018). In the same

bloom, viral infection stage was found to proceed with water column depth (Sheyn et

al., 2018). Enhanced TEP production for these same early infected calcifying

populations was observed over a three-day period in deck-board bottle incubations

(Laber et al., 2018). Laboratory observations also exist for enhanced TEP production

and aggregate formation during the early phase of EhV infection of a calcifying *E.*

huxleyi strain (Laber et al., 2018; Nissimov et al., 2018). These observations strongly

suggest that infection-induced TEP production in organisms containing dense material

(*e.g.*, calcite scales for *E. huxleyi*) can facilitate carbon export. No EhV-like PolB

sequences were detected in our dataset, which was probably due to sampled areas and

seasons.

Laboratory experiments suggest that viruses closely related to positive VIPs,

such as prasinoviruses, have infectious properties that may drive carbon export.

Cultures of *Micromonas pusilla* infected with prasinoviruses showed increased TEP

production compared with non-infected cultures (Lønborg et al., 2013), although it is

not known if this increase leads to aggregation. The hosts of prasinoviruses have been

proposed to contribute to carbon export because they were observed in abyssopelagic

zone at sampling sites dominated by Mamiellales in their surface waters in the

western subtropical North Pacific (Shiozaki et al., 2019). Some prasinoviruses encode

glycosyltransferases (GTs) of the GT2 family. Similar to the a098r gene (GT2) in

Paramecium bursaria Chlorella virus 1, the expression of GT2 family members

during infection possibly leads to the production of a dense fibrous hyaluronan

network at the surface of infected cells. Such a network may trigger the aggregation

of host cells, facilitate viral propagation (Van Etten et al., 2017), and increase the cell

wall C:N ratio. We detected one GT2 in a MAG of two *Phycodnaviridae*-like positive

VIPs (000200745 and 002503270) predicted to infect Mamiellales, one in a MAG

corresponding to the putative pelagophyte positive VIP 000079111 related to AaV

and six in two MAGs (three each) corresponding to two *Mimiviridae*-like positive

VIPs (000328966 and 001175669). *Phaeocystis globosa virus* (PgV), closely related

to the positive VIP PolB 000912507 (Figure S8), has been linked with increased TEP

production and aggregate formation during the termination of a *Phaeocystis* bloom

(Brussaard et al., 2007). Two closely related bacillarnavirus VIPs were positively

associated with CEE and predicted to infect Chaetocerales. A previous study revealed

an increase in abundance of viruses infecting diatoms of *Chaetoceros* in both the

water columns and the sediments during the bloom of their hosts in a coastal area

(Tomaru et al., 2011), suggesting sinking of cells caused by viruses. Furthermore, the

diatom *Chaetoceros tenuissimus* infected with a DNA virus (CtenDNAV type II) has

been shown to produce higher levels of large-sized particles (50 to 400 μm) compared

with non-infected cultures (Yamada et al., 2018).

   The other 25 VIPs were negatively associated with CEE. This association is

compatible with the "viral shunt," which increases the amount of DOC (Wilhelm and

Suttle, 1999) and reduces the transfer of carbon to higher trophic levels and to the

deep ocean (Fuhrman, 1999; Weitz et al., 2015). Increased DOC has been observed in

culture of Mamiellales lysed by prasinoviruses (Lønborg et al., 2013). Although this

culture-based observation may be difficult to extrapolate to natural conditions, where

the cell concentration and thus the contact rate with viruses are probably lower,

Mamiellales species are known to form blooms during which cell densities may be

comparable with cultures (Zhu et al., 2005). A field study reported that PgV, to which

the negative VIP PolB 000054135 is closely related (Figure S8), can be responsible

for up to 35% of cell lysis per day during bloom of its host (Baudoux et al., 2006),

which is likely accompanied by consequent DOC release. Similarly, the decline of a

bloom of the pelagophyte *Aureococcus anophagefferens* has been associated with

active infection by AaV (to which one negative VIP is closely related)

(Moniruzzaman et al., 2017). Among RNA viruses, eight were negative VIPs (six

Picornavirales and two *Hepeviridae*). The higher representation of *Picornavirales* in

the virioplankton (Culley, 2018) than within cells (Urayama et al., 2018) suggests that

they are predominantly lytic, although no information exists regarding the effect of

Picornavirales on DOC release.

It is likely that the "viral shunt" and "viral shuttle" simultaneously affect and

modulate CEE in the global ocean (Zimmerman et al., 2019a). The relative

importance of these two phenomena must fluctuate considerably depending on the

host traits, viral effects on metabolism, and environmental conditions. Reflecting this

complexity, viruses of a same host group could be found to be either positively or

negatively associated with CEE. For example, among prasinoviruses most likely

infecting Mamiellales, 18 were positive VIPs and six were negative VIPs. Two

closely related prasinoviruses (sharing 97.5% genome-wide identity) are known to

exhibit different ecological strategies with notably distinct molecular signatures on the

organic matter released upon infection of the same host (Zimmerman et al., 2019b).

We found that even two very closely related *Mimiviridae* viruses (PolBs 000079111

and 000079078 sharing 94% nucleotide identity over their full gene lengths) most

likely infecting pelagophyte algae were positively and negatively associated with

CEE. Furthermore, it is known that an early-infected *E. huxleyi* system was linked

with both higher aggregation "at surface" and higher remineralization "at deep"

compared to late-infected blooms (Laber et al., 2018). Therefore, the viral effect on

carbon cycle may vary also with depth.

Five percent of the tested viruses were associated with CEE in our study.

Similarly, four percent of bacterial virus populations were found to be associated with

the magnitude of carbon export at 150 meters (Guidi et al., 2016). These results

suggest that viruses affecting carbon export are rather uncommon. It is plausible that

such viruses affect CEE by infecting organisms that are functionally important

(abundant or keystone species), as we observed in host prediction. The vast majority

(95%) of non-VIPs may not have a significant impact on CEE, because they do not

strongly impact the host population, for instance, by stably coexisting with their hosts.

It is worth noting that experimental studies have reported cultures of algae with

viruses that reach a stable co-existence state after a few generations (Yau et al., 2020).

It is also possible that some of these non-VIPs can impact carbon export but were not

captured in the infection stage affecting the export process. Viruses captured in our

samples can represent active viruses in different infectious stages (early, mid or late)

for metagenomes and metatranscriptomes or at the post-lysis stage for metagenomes.

Potential effects of global climate changes on viral shunt/shuttle and CEE

Increasing evidence suggests that the biological carbon pump is highly dependent on

the planktonic community composition, and as discussed above, viruses represent a

possible key parameter that determines the efficiency of carbon export. In the photic

layer of the oceans, the composition of planktonic communities is strongly affected by

sea surface temperature (Salazar et al., 2019; Sunagawa et al., 2015), and CEE may

therefore be affected by ocean warming. Our result indicated that viruses infecting

small phytoplankton such as Mamiellales and haptophytes are likely associated with

CEE. Interestingly, many studies showed that high temperature and/or CO_2 levels are

associated with an increased contribution of small sized phytoplankton to the total

biomass (Hare et al., 2007; Mousing et al., 2014; Sugie et al., 2020).

An increase in CO_2 level in the surface seawater also causes a decrease in pH

(i.e., ocean acidification). Previous studies demonstrated that the decrease in seawater

pH negatively affect the growth of calcified and silicified phytoplankton cells (Doney

et al., 2009; Endo et al., 2016; Petrou et al., 2019). The biogenic minerals such as

calcium carbonate and silica act as ballasts in sinking particles (Iversen and Ploug,

2010). Given the statistical association that we detected between the viruses positively

correlated with CEE and the silicified predicted host planktons, the ocean

acidification may decrease the viral shuttle and thus CEE globally in the future.

The increased sea surface temperature will decrease the nutrient supply at the

surface of the oligotrophic ocean by preventing the vertical mixing. The decrease in

nutrient availability of surface seawaters possibly diminishes the net primary

production (NPP) and the magnitude of carbon export (corresponded to CE_{150} in our

study) (Riebesell et al., 2009). Consistently, a downward trend of global

phytoplankton abundance has been observed by satellite (Boyce et al., 2010). In such

a scenario of the global decrease of NPP, the efficiency of export would be an

important factor for a precise estimation of carbon export in the future ocean. In this

regard, the role of marine viruses in the carbon cycle and export should be further

investigated and eventually be integrated into prospective models for the climate

change.

Conclusions

Eukaryotic virus community composition was able to predict CEE at 59 sampling

sites in the photic zone of the world ocean. This statistical association was detected

based on a large omics dataset collected throughout the oceans and processed with

standardized protocols. The predictability of CEE by viral composition is consistent

with the hypothesis that "viral shuttle" and "shunt" are functioning at a global scale.

Among 83 viruses with a high importance in the prediction of CEE, 58 viruses were

positively and 25 negatively correlated with carbon export efficiency. Most of these

viruses belong to *Prasinovirus*, *Mimiviridae*, and *Picornavirales* and are either new to science or with no known roles in carbon export efficiency. Thirty-six of these "select" viruses were predicted to infect ecologically important hosts such as green algae of the order Mamiellales, haptophytes, diatoms, and copepods. Positively associated viruses had more predicted interactions with silicified eukaryotes than non-associated viruses did. Overall, these results imply that the effect of viruses on the "shuttle" and "shunt" processes could be dependent on viral hosts and ecosystem dynamics.

Limitations of the study

The observed statistical associations between viral compositions and examined parameters (*i.e.*, CEE, CE and NPP) do not convey the information about the direction of their potential causality relationships, and they could even result from indirect relationships as discussed above. Certain groups of viruses detected in samples may be over- or under-represented because of the technical limitations in size fractionation, DNA/RNA extraction and sequencing.

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by Lead Contact, Hiroyuki Ogata (ogata@kuicr.kyoto-u.ac.jp).

Materials Availability

This study did not generate unique reagent.

Data and Code Availability

The authors declare that the data supporting the findings of this study are available within the paper and its supplemental files, as well as at the GenomeNet FTP: ftp://ftp.genome.jp/pub/db/community/tara/Cpump/Supplementary_material/.

Our custom R script used to test for association between viruses and environmental variables (CEE, CE_{150} and NPP) is available along with input data at the GenomeNet FTP: ftp://ftp.genome.jp/pub/db/community/tara/Cpump/Supplementary_material/PLSreg/.

The Taxon Interaction Mapper (TIM) tool developed for this study and used for virus host prediction is available at https://github.com/RomainBlancMathieu/TIM.

Supplemental Files

- Supplemental_Information.pdf: supplemental figures and tables, and transparent methods
- Supplemental_Data_1_2.xlsx

Acknowledgements

We thank the *Tara* Oceans consortium, the projects Oceanomics and France Genomique (grants ANR-11-BTBR-0008 and ANR-10-INBS-09), and the people and sponsors who supported the *Tara* Oceans Expedition (http://www.embl.de/tara-oceans/) for making the data accessible. This is contribution number XXX of the *Tara* Oceans Expedition 2009–2013. Computational time was provided by the SuperComputer System, Institute for Chemical Research, Kyoto University. We thank Barbara Goodson, Ph.D., and Sara J. Mason, M.Sc., from Edanz Group (https://en-author-services.edanzgroup.com/) for editing a draft of this manuscript. This work

was supported by JSPS/KAKENHI (Nos. 26430184, 18H02279, and 19H05667 to

H.O. and Nos. 19K15895 and 19H04263 to H.E.), Scientific Research on Innovative

Areas from the Ministry of Education, Culture, Science, Sports and Technology

(MEXT) of Japan (Nos. 16H06429, 16K21723, and 16H06437 to H.O.), the

Collaborative Research Program of the Institute for Chemical Research, Kyoto

University (2019-29 to S.C.), the Future Development Funding Program of the Kyoto

University Research Coordination Alliance (to R.B.M.), the ICR-KU International

Short-term Exchange Program for Young Researchers (to S.C.), and the Research

Unit for Development of Global Sustainability (to H.O. and T.O.D.).

Author contributions

H.O. and R.B.M. conceived the study. R.B.M and H.K. performed most of the

analyses. H.E. and L.G. designed carbon export analysis. R.H.V and S.C. performed

network analysis. N.H. and C.d.V. analyzed eukaryotic sequences. T.O.D., M.G., P.F.

and O.J. analyzed viral MAGs. C.H.N. and H.M. contributed to statistical analysis.

M.B.S. and C.A.S. contributed to interpretations. All authors edited and approved the

final version of the manuscript.

Declaration of Interests

576 The authors declare no competing interests.

References

Agusti, S., González-Gordillo, J.I., Vaqué, D., Estrada, M., Cerezo, M.I., Salazar, G.,
Gasol, J.M., and Duarte, C.M. (2015). Ubiquitous healthy diatoms in the deep sea
confirm deep carbon injection by the biological pump. Nat. Commun. *6*, 7608.

Baudoux, A., Noordeloos, A., Veldhuis, M., and Brussaard, C. (2006). Virally
induced mortality of Phaeocystis globosa during two spring blooms in temperate
coastal waters. Aquat. Microb. Ecol. *44*, 207–217.

Bochdansky, A.B., Clouse, M.A., and Herndl, G.J. (2017). Eukaryotic microbes,
principally fungi and labyrinthulomycetes, dominate biomass on bathypelagic marine
snow. ISME J. *11*, 362–373.

Boyce, D.G., Lewis, M.R., and Worm, B. (2010). Global phytoplankton decline over
the past century. Nature *466*, 591–596.

Brum, J.R., Ignacio-Espinoza, J.C., Roux, S., Doulcier, G., Acinas, S.G., Alberti, A.,
Chaffron, S., Cruaud, C., Vargas, C. de, Gasol, J.M., et al. (2015). Patterns and
ecological drivers of ocean viral communities. Science *348*, 1261498.

Brussaard, C.P.D., Bratbak, G., Baudoux, A.-C., and Ruardij, P. (2007). Phaeocystis
and its interaction with viruses. Biogeochemistry *83*, 201–215.

Buesseler, K.O., and Boyd, P.W. (2009). Shedding light on processes that control
particle export and flux attenuation in the twilight zone of the open ocean. Limnol.
Oceanogr. *54*, 1210–1232.

Carradec, Q., Pelletier, E., Silva, C.D., Alberti, A., Seeleuthner, Y., Blanc-Mathieu,
R., Lima-Mendez, G., Rocha, F., Tirichine, L., Labadie, K., et al. (2018). A global
ocean atlas of eukaryotic genes. Nat. Commun. *9*, 373.

Coenen, A.R., and Weitz, J.S. (2018). Limitations of Correlation-Based Inference in
Complex Virus-Microbe Communities. MSystems *3*, e00084-18.

Culley, A. (2018). New insight into the RNA aquatic virosphere via viromics. Virus
Res. *244*, 84–89.

Dayaram, A., Goldstien, S., Argüello-Astorga, G.R., Zawar-Reza, P., Gomez, C.,
Harding, J.S., and Varsani, A. (2015). Diverse small circular DNA viruses circulating
amongst estuarine molluscs. Infect. Genet. Evol. J. Mol. Epidemiol. Evol. Genet.
Infect. Dis. *31*, 284–295.

Doney, S.C., Fabry, V.J., Feely, R.A., and Kleypas, J.A. (2009). Ocean Acidification:
The Other CO_2 Problem. Annu. Rev. Mar. Sci. *1*, 169–192.

Dunlap, D.S., Ng, T.F.F., Rosario, K., Barbosa, J.G., Greco, A.M., Breitbart, M., and
Hewson, I. (2013). Molecular and microscopic evidence of viruses in marine
copepods. Proc. Natl. Acad. Sci. *110*, 1375–1380.

Endo, H., Sugie, K., Yoshimura, T., and Suzuki, K. (2016). Response of Spring
Diatoms to CO2 Availability in the Western North Pacific as Determined by Next-
Generation Sequencing. PLOS ONE *11*, e0154291.

Endo, H., Ogata, H., and Suzuki, K. (2018). Contrasting biogeography and diversity
patterns between diatoms and haptophytes in the central Pacific Ocean. Sci. Rep. *8*,
10916.

Evans, C., and Wilson, W.H. (2008). Preferential grazing of Oxyrrhis marina on virus
infected Emiliania huxleyi. Limnol. Oceanogr. *53*, 2035–2040.

Fawcett, S.E., Lomas, M.W., Casey, J.R., Ward, B.B., and Sigman, D.M. (2011).
Assimilation of upwelled nitrate by small eukaryotes in the Sargasso Sea. Nat.
Geosci. *4*, 717–722.

Fuhrman, J.A. (1999). Marine viruses and their biogeochemical and ecological
effects. Nature *399*, 541–548.

Gobler, C.J., Hutchins, D.A., Fisher, N.S., Cosper, E.M., and Saňudo- Wilhelmy,
S.A. (1997). Release and bioavailability of C, N, P Se, and Fe following viral lysis of
a marine chrysophyte. Limnol. Oceanogr. *42*, 1492–1504.

Goic, B., and Saleh, M.-C. (2012). Living with the enemy: viral persistent infections
from a friendly viewpoint. Curr. Opin. Microbiol. *15*, 531–537.

Goode, A.G., Fields, D.M., Archer, S.D., and Martínez, J.M. (2019). Physiological
responses of Oxyrrhis marina to a diet of virally infected Emiliania huxleyi. PeerJ *7*,
e6722.

Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlimi, A., Roux, S., Darzi, Y.,
Audic, S., Berline, L., Brum, J.R., et al. (2016). Plankton networks driving carbon
export in the oligotrophic ocean. Nature *532*, 465.

Hare, C., Leblanc, K., DiTullio, G., Kudela, R., Zhang, Y., Lee, P., Riseman, S., and
Hutchins, D. (2007). Consequences of increased temperature and CO2 for
phytoplankton community structure in the Bering Sea. Mar. Ecol. Prog. Ser. *352*, 9–
16.

Hingamp, P., Grimsley, N., Acinas, S.G., Clerissi, C., Subirana, L., Poulain, J.,
Ferrera, I., Sarmento, H., Villar, E., Lima-Mendez, G., et al. (2013). Exploring
nucleo-cytoplasmic large DNA viruses in Tara Oceans microbial metagenomes. ISME
J. *7*, 1678–1695.

Hirata, T., Hardman-Mountford, N.J., Brewin, R.J.W., Aiken, J., Barlow, R., Suzuki,
K., Isada, T., Howell, E., Hashioka, T., Noguchi-Aita, M., et al. (2011). Synoptic
relationships between surface Chlorophyll-*a* and diagnostic pigments specific to
phytoplankton functional types. Biogeosciences *8*, 311–327.

Hurwitz, B.L., Brum, J.R., and Sullivan, M.B. (2015). Depth-stratified functional and
taxonomic niche specialization in the "core" and "flexible" Pacific Ocean Virome.
ISME J. *9*, 472–484.

Iversen, M.H., and Ploug, H. (2010). Ballast minerals and the sinking carbon flux in
the ocean: carbon-specific respiration rates and sinking velocity of marine snow
aggregates. Biogeosciences *7*, 2613–2624.

Karl, D.M., Church, M.J., Dore, J.E., Letelier, R.M., and Mahaffey, C. (2012).
Predictable and efficient carbon sequestration in the North Pacific Ocean supported
by symbiotic nitrogen fixation. Proc. Natl. Acad. Sci. *109*, 1842–1849.

Klaas, C., and Archer, D.E. (2002). Association of sinking organic matter with
various types of mineral ballast in the deep sea: Implications for the rain ratio. Glob.
Biogeochem. Cycles *16*, 63-1-63–14.

Laber, C.P., Hunter, J.E., Carvalho, F., Collins, J.R., Hunter, E.J., Schieler, B.M.,
Boss, E., More, K., Frada, M., Thamatrakoln, K., et al. (2018). Coccolithovirus
facilitation of carbon export in the North Atlantic. Nat. Microbiol. *3*, 537–547.

Lawrence, J.E., and Suttle, C.A. (2004). Effect of viral infection on sinking rates of
Heterosigma akashiwo and its implications for bloom termination. Aquat. Microb.
Ecol. *37*, 1–7.

Lawrence, J.E., Chan, A.M., and Suttle, C.A. (2002). Viruses causing lysis of the
toxic bloom-forming alga Heterosigma akashiwo (Raphidophyceae) are widespread in
coastal sediments of British Columbia, Canada. Limnol. Oceanogr. *47*, 545–550.

Leblanc, K., Quéguiner, B., Diaz, F., Cornet, V., Michel-Rodriguez, M., Durrieu de
Madron, X., Bowler, C., Malviya, S., Thyssen, M., Grégori, G., et al. (2018).
Nanoplanktonic diatoms are globally overlooked but play a role in spring blooms and
carbon export. Nat. Commun. *9*, 953.

Li, W. (1995). Composition of Ultraphytoplankton in the Central North-Atlantic. Mar.
Ecol. Prog. Ser. *122*, 1–8.

Liu, H., Probert, I., Uitz, J., Claustre, H., Aris-Brosou, S., Frada, M., Not, F., and de
Vargas, C. (2009). Extreme diversity in noncalcifying haptophytes explains a major
pigment paradox in open oceans. Proc. Natl. Acad. Sci. U. S. A. *106*, 12803–12808.

Lomas, M.W., and Moran, S.B. (2011). Evidence for aggregation and export of
cyanobacteria and nano-eukaryotes from the Sargasso Sea euphotic zone.
Biogeosciences *8*, 203–216.

Lønborg, C., Middelboe, M., and Brussaard, C.P.D. (2013). Viral lysis of
Micromonas pusilla: impacts on dissolved organic matter production and
composition. Biogeochemistry *116*, 231–240.

Mars Brisbin, M., Mesrop, L.Y., Grossmann, M.M., and Mitarai, S. (2018). Intra-host
Symbiont Diversity and Extended Symbiont Maintenance in Photosymbiotic
Acantharea (Clade F). Front. Microbiol. *9*.

Monier, A., Pagarete, A., de Vargas, C., Allen, M.J., Read, B., Claverie, J.-M., and
Ogata, H. (2009). Horizontal gene transfer of an entire metabolic pathway between a
eukaryotic alga and its DNA virus. Genome Res. *19*, 1441–1449.

Monier, A., Worden, A.Z., and Richards, T.A. (2016). Phylogenetic diversity and
biogeography of the Mamiellophyceae lineage of eukaryotic phytoplankton across the
oceans. Environ. Microbiol. Rep. *8*, 461–469.

Moniruzzaman, M., Wurch, L.L., Alexander, H., Dyhrman, S.T., Gobler, C.J., and
Wilhelm, S.W. (2017). Virus-host relationships of marine single-celled eukaryotes
resolved from metatranscriptomics. Nat. Commun. *8*, 16054.

Mousing, E., Ellegaard, M., and Richardson, K. (2014). Global patterns in
phytoplankton community size structure—evidence for a direct temperature effect.
Mar. Ecol. Prog. Ser. *497*, 25–38.

Nissimov, J.I., Vandzura, R., Johns, C.T., Natale, F., Haramaty, L., and Bidle, K.D.
(2018). Dynamics of transparent exopolymer particle production and aggregation
during viral infection of the coccolithophore, Emiliania huxleyi. Environ. Microbiol.
20, 2880–2897.

Peduzzi, P., and Weinbauer, M.G. (1993). Effect of concentrating the virus-rich 2-
2nm size fraction of seawater on the formation of algal flocs (marine snow). Limnol.
Oceanogr. *38*, 1562–1565.

Petrou, K., Baker, K.G., Nielsen, D.A., Hancock, A.M., Schulz, K.G., and Davidson,
A.T. (2019). Acidification diminishes diatom silica production in the Southern Ocean.
Nat. Clim. Change *9*, 781–786.

Proctor, L.M., and Fuhrman, J.A. (1991). Roles of viral infection in organic particle
flux. Mar. Ecol. Prog. Ser. *69*, 133–142.

Riebesell, U., Kortzinger, A., and Oschlies, A. (2009). Sensitivities of marine carbon
fluxes to ocean change. Proc. Natl. Acad. Sci. *106*, 20602–20609.

Salazar, G., Paoli, L., Alberti, A., Huerta-Cepas, J., Ruscheweyh, H.-J., Cuenca, M.,
Field, C.M., Coelho, L.P., Cruaud, C., Engelen, S., et al. (2019). Gene Expression
Changes and Community Turnover Differentially Shape the Global Ocean
Metatranscriptome. Cell *179*, 1068-1083.e21.

Sheyn, U., Rosenwasser, S., Lehahn, Y., Barak-Gavish, N., Rotkopf, R., Bidle, K.D.,
Koren, I., Schatz, D., and Vardi, A. (2018). Expression profiling of host and virus
during a coccolithophore bloom provides insights into the role of viral infection in
promoting carbon export. ISME J. 1.

Shi, M., Lin, X.-D., Tian, J.-H., Chen, L.-J., Chen, X., Li, C.-X., Qin, X.-C., Li, J.,
Cao, J.-P., Eden, J.-S., et al. (2016). Redefining the invertebrate RNA virosphere.
Nature *540*, 539–543.

Shiozaki, T., Hirose, Y., Hamasaki, K., Kaneko, R., Ishikawa, K., and Harada, N.
(2019). Eukaryotic Phytoplankton Contributing to a Seasonal Bloom and Carbon
Export Revealed by Tracking Sequence Variants in the Western North Pacific. Front.
Microbiol. *10*.

Sugie, K., Fujiwara, A., Nishino, S., Kameyama, S., and Harada, N. (2020). Impacts
of Temperature, CO2, and Salinity on Phytoplankton Community Composition in the
Western Arctic Ocean. Front. Mar. Sci. *6*, 821.

Sullivan, M.B., Weitz, J.S., and Wilhelm, S. (2017). Viral ecology comes of age.
Environ. Microbiol. Rep. *9*, 33–35.

Sunagawa, S., Coelho, L.P., Chaffron, S., Kultima, J.R., Labadie, K., Salazar, G.,
Djahanschiri, B., Zeller, G., Mende, D.R., Alberti, A., et al. (2015). Ocean plankton.
Structure and function of the global ocean microbiome. Science *348*, 1261359.

Suttle, C.A. (2007). Marine viruses--major players in the global ecosystem. Nat. Rev.
Microbiol. *5*, 801–812.

Takao, Y., Nagasaki, K., Mise, K., Okuno, T., and Honda, D. (2005). Isolation and
characterization of a novel single-stranded RNA Virus infectious to a marine fungoid
protist, Schizochytrium sp. (Thraustochytriaceae, Labyrinthulea). Appl. Environ.
Microbiol. *71*, 4516–4522.

Tomaru, Y., Hata, N., Masuda, T., Tsuji, M., Igata, K., Masuda, Y., Yamatogi, T.,
Sakaguchi, M., and Nagasaki, K. (2007). Ecological dynamics of the bivalve-killing
dinoflagellate Heterocapsa circularisquama and its infectious viruses in different
locations of western Japan. Environ. Microbiol. *9*, 1376–1383.

Tomaru, Y., Fujii, N., Oda, S., Toyoda, K., and Nagasaki, K. (2011). Dynamics of
diatom viruses on the western coast of Japan. Aquat. Microb. Ecol. *63*, 223–230.

Tréguer, P., Bowler, C., Moriceau, B., Dutkiewicz, S., Gehlen, M., Aumont, O.,
Bittner, L., Dugdale, R., Finkel, Z., Iudicone, D., et al. (2018). Influence of diatom
diversity on the ocean biological carbon pump. Nat. Geosci. *11*, 27–37.

Turner, J.T. (2015). Zooplankton fecal pellets, marine snow, phytodetritus and the
ocean's biological pump. Prog. Oceanogr. *130*, 205–248.

Urayama, S., Takaki, Y., Nishi, S., Yoshida- Takashima, Y., Deguchi, S., Takai, K.,
and Nunoura, T. (2018). Unveiling the RNA virosphere associated with marine
microorganisms. Mol. Ecol. Resour. *18*, 1444–1455.

Van Etten, J.L., Agarkova, I., Dunigan, D.D., Tonetti, M., De Castro, C., and Duncan,
G.A. (2017). Chloroviruses Have a Sweet Tooth. Viruses *9*.

Wang, H., Wu, S., Li, K., Pan, Y., Yan, S., and Wang, Y. (2018). Metagenomic
analysis of ssDNA viruses in surface seawater of Yangshan Deep-Water Harbor,
Shanghai, China. Mar. Genomics *41*, 50–53.

Ward, B.A., and Follows, M.J. (2016). Marine mixotrophy increases trophic transfer
efficiency, mean organism size, and vertical carbon flux. Proc. Natl. Acad. Sci. *113*,
2958–2963.

Weinbauer, M.G. (2004). Ecology of prokaryotic viruses. FEMS Microbiol. Rev. *28*,
127–181.

Weitz, J.S., Stock, C.A., Wilhelm, S.W., Bourouiba, L., Coleman, M.L., Buchan, A.,
Follows, M.J., Fuhrman, J.A., Jover, L.F., Lennon, J.T., et al. (2015). A multitrophic
model to quantify the effects of marine viruses on microbial food webs and ecosystem
processes. ISME J. *9*, 1352–1364.

Wilhelm, S.W., and Suttle, C.A. (1999). Viruses and Nutrient Cycles in the
SeaViruses play critical roles in the structure and function of aquatic food webs.
BioScience *49*, 781–788.

Yamada, Y., Tomaru, Y., Fukuda, H., and Nagata, T. (2018). Aggregate Formation
During the Viral Lysis of a Marine Diatom. Front. Mar. Sci. *5*.

Yau, S., Krasovec, M., Benites, L.F., Rombauts, S., Groussin, M., Vancaester, E.,
Aury, J.-M., Derelle, E., Desdevises, Y., Escande, M.-L., et al. (2020). Virus-host
coexistence in phytoplankton through the genomic lens. Sci. Adv. *6*, eaay2587.

Yoshida, M., Noël, M.-H., Nakayama, T., Naganuma, T., and Inouye, I. (2006). A
haptophyte bearing siliceous scales: ultrastructure and phylogenetic position of
Hyalolithus neolepis gen. et sp. nov. (Prymnesiophyceae, Haptophyta). Protist *157*,
213–234.

Yoshikawa, G., Blanc-Mathieu, R., Song, C., Kayama, Y., Mochizuki, T., Murata, K.,
Ogata, H., and Takemura, M. (2019). Medusavirus, a novel large DNA virus
discovered from hot spring water. J. Virol. JVI.02130-18.

Zhang, C., Dang, H., Azam, F., Benner, R., Legendre, L., Passow, U., Polimene, L.,
Robinson, C., Suttle, C.A., and Jiao, N. (2018). Evolving paradigms in biological
carbon cycling in the ocean. Natl. Sci. Rev. *5*, 481–499.

Zhu, F., Massana, R., Not, F., Marie, D., and Vaulot, D. (2005). Mapping of
picoeucaryotes in marine ecosystems with quantitative PCR of the 18S rRNA gene.
FEMS Microbiol. Ecol. *52*, 79–92.

Zimmerman, A.E., Howard-Varona, C., Needham, D.M., John, S.G., Worden, A.Z.,
Sullivan, M.B., Waldbauer, J.R., and Coleman, M.L. (2019a). Metabolic and
biogeochemical consequences of viral infection in aquatic ecosystems. Nat. Rev.
Microbiol. 1–14.

Zimmerman, A.E., Bachy, C., Ma, X., Roux, S., Jang, H.B., Sullivan, M.B.,
Waldbauer, J.R., and Worden, A.Z. (2019b). Closely related viruses of the marine
picoeukaryotic alga Ostreococcus lucimarinus exhibit different ecological strategies.
Environ. Microbiol. *21*, 2148–2170.

Figure legends

Figure 1: Viruses of eukaryotic plankton identified in *Tara* Oceans samples are distantly related to characterized viruses. Unrooted maximum likelihood phylogenetic trees containing environmental (black) and reference (red) viral sequences for NCLDV DNA polymerase family B (**a**), RNA virus RNA-dependent RNA polymerase (**b**), and ssDNA virus replication-associated protein (**c**). A rectangular representation of these trees with branch support values is provided in Figure S2–S4.

Figure 2: Carbon export efficiency and relative marker-gene occurrence of eukaryotic plankton viruses along the sampling route. a Carbon export efficiency estimated at 39 *Tara* Oceans stations where surface and DCM layers were sampled for prokaryote-enriched metagenomes and eukaryotic metatranscriptomes. **b and c** Relative marker-gene occurrence of major groups of viruses of eukaryotic plankton for NCLDVs in metagenomes (**b**) and for RNA and ssDNA viruses in metatranscriptomes (**c**) at 59 sampling sites.

Figure 3: Relative abundance of eukaryotic plankton viruses associated with carbon export efficiency in the global ocean. a Bivariate plot between predicted and observed values in a leave-one-out cross-validation test for carbon export efficiency. The PLS regression model was constructed using occurrence profiles of 1,523 marker-gene sequences (1,309 PolBs, 180 RdRPs and 34 Reps) derived from environmental samples. r, Pearson correlation coefficient; R^2, the coefficient of determination between measured response values and predicted response values. R^2, which was calculated as $1 - SSE/SST$ (sum of squares due to error and total)

measures how successful the fit is in explaining the variance of the response values.

The significance of the association was assessed using a permutation test ($n = 10,000$)

(grey histogram in **a**). The red diagonal line shows the theoretical curve for perfect

prediction. **b** Pearson correlation coefficients between CEE and occurrence profiles of

83 viruses that have VIP scores > 2 (VIPs) with the first two components in the PLS

regression model using all samples. PLS components 1 and 2 explained 83% and 11%

of the variance of CEE, respectively. Fifty-eight VIPs had positive regression

coefficients in the model (shown with circles), and 25 had negative regression

coefficients (shown with triangles).

Figure 4: Biogeography of viruses associated with carbon export efficiency. The

upper panel shows carbon export efficiency (CEE = $CE_{deep}/CE_{surface}$) for 59 sampling

sites. The bottom panel is a map reflecting relative abundances, expressed as centered

log-ratio transformed, gene-length normalized read counts of viruses positively and

negatively associated with CEE that have VIP scores > 2 (VIPs). MS, Mediterranean

Sea; IO, Indian Ocean; SAO, South Atlantic Ocean; SPO, South Pacific Ocean; NPO,

North Pacific Ocean; NAO, North Atlantic Ocean. The bottom horizontal axis is

labeled with *Tara* Oceans station numbers, sampling depth (SRF, surface; DCM, deep

chlorophyll maximum), and abbreviations of biogeographic provinces. Viruses

labeled in red correspond to positive VIPs that tend more represented in more than

one biogeographic province.

Tables

Table 1: Taxonomic breakdown of viral marker genes

Viruses		Identified	Used in PLS regression*
NCLDVs	Mimiviridae	2,923	1,148
	Phycodnaviridae	348	99
	Iridoviridae	198	59
	Other NCLDVs **	17	3
	Total	3,486	1,309
RNA viruses	Picornavirales (ssRNA+)	325	80
	Partitiviridae (dsRNA)	131	22
	Narnaviridae (ssRNA+)	95	6
	Other families	289	53
	Unclassified	78	9
	RNA viruses	57	10
	Total	975	180
ssDNA viruses	Circoviridae	201	22
	Geminiviridae	4	0
	Nanoviridae	4	0
	Unclassified	39	2
	ssDNA viruses	51	10
	Total	299	34
	All	4,760	1,523

* The marker genes had to occurred in at least five samples and harbor a Spearman correlation coefficient > |0.2| with carbon export efficiency.
** There was no unclassified NCLDV.

Figure 1

Figure 2

a

b

Relative Occurrence (%)

■ *Mimiviridae* ■ *Phycodnaviridae* ■ *Iridoviridae* □ Other NCLDVs

c

Relative Occurrence (%)

■ *Picornavirales* ■ *Partitiviridae* ■ *Narnaviridae* □ Other RNA viruses
■ *Circoviridae* □ Other ssDNA viruses

Figure 3

Sign of regression coefficient : ○ Positive (pos) △ Negative (neg)

Taxonomy :
● *Mimiviridae* (*n*=34, 25 pos/9 neg)
● *Phycodnaviridae* (*n*=24, 18 pos/6 neg)
● *Iridoviridae* (*n*=2, 0 pos/2 neg)
● *Picornavirales* (*n*=19, 13 pos/6 neg)
● *Hepeviridae* (*n*=2, 0 pos/2 neg)
● *Partitiviridae* (*n*=1, 1 pos/0 neg)
● *Circoviridae* (*n*=1, 1 pos/0 neg)

Figure 4

1   **Supplemental Information**

2   **Supplemental Figures**

5   **Figure S1:** Distribution of the degree (%) of amino acid identity between environmental
6   sequences and their best BLAST hits to reference sequences for nucleocytoplasmic large
7   DNA viruses (NCLDVs) (**a**), RNA viruses (**b**), and ssDNA viruses (**c**).

8
9 **Figure S2:** Maximum likelihood phylogenetic trees for NCLDV DNA polymerase family B.
10 Environmental sequences are shown in black and references in red. Approximate Shimodaira–
11 Hasegawa (SH)-like local support values greater than 0.8 are shown. Scale bar indicates one
12 change per site.

Tree scale: 1

Bunyavirales

Narnaviridae

Mononegavirales

Other families

Endornaviridae

Tombusviridae

Other families

Bromoviridae

Tymovirales

Flaviviridae

Bunyavirales

Totiviridae

Partitiviridae

Picornavirales

13
14 **Figure S3:** Unrooted maximum likelihood phylogenetic trees for RNA virus RNA-dependent
15 RNA polymerase. Environmental sequences are shown in black and references in red.
16 Approximate SH-like local support values greater than 0.8 are shown. Scale bar indicates one
17 change per site.

Tree scale: 1

Smacoviridae

Nanoviridae

CRESS5

CRESS4

CRESS6

Genomo viridae

Geminiviridae

CRESS1

Bacilladnaviridae

CRESS2

CRESS3

Circoviridae

18
19 **Figure S4:** Unrooted maximum likelihood phylogenetic trees for ssDNA virus replication-
20 associated protein. Environmental sequences are shown in black and references in red.
21 Approximate SH-like local support values greater than 0.8 are shown. Scale bar indicates one
22 change per site.
23

24
Figure S5: Variation in carbon export flux (mg m^{-2} d^{-1}) across sampling depths in the water
column. Dots are average values, and horizontal lines represent standard deviations.
27

Figure S6: The results of PLS regressions using relative abundance profiles of viral marker-genes to explain the variance of carbon export efficiency (CEE) (**a, d, g, j**), carbon export flux at 150 meters (CE$_{150}$) (**b, e, h, k**), and net primary production (NPP) (**c, f, i, l**). **a–c** Bivariate plots between predicted and observed response values in a leave-one-out cross-validation test. The red diagonal line shows the theoretical curve for perfect prediction. **d–f** Variation in root mean squared error of predictions (RMSEP) for the training set (solid black line) and cross-validation set (red dashed line) across the number of components. Blue dashed line shows the number of components selected for the analysis. **g–i** Results of the permutation tests ($n = 10{,}000$) supporting the significance of the association between viruses and the response variable. The histograms show the distribution of Pearson correlation coefficients obtained from PLS models reconstructed based on the permutated response variable and red line show the non-permutated response variable. **j–l** Pearson correlation coefficients between the response variable and abundance profiles of viruses with VIP scores > 2 (VIPs) with the first two components in the PLS regression model using all samples. Viruses with positive regression coefficients are shown with circles, and those with negative coefficients are shown with triangles.

Figure S7: Taxonomic composition of genes predicted in viral genome fragments encoding NCLDV PolBs positively (a and c) or negatively (b and d) associated with CEE (VIP score > 2). **a and b** Metagenome-assembled genomes (MAGs) derived from samples filtered to retain particles of sizes > 0.8 μm. **c and d** Contigs derived from samples filtered to retain particles between 0.2 μm and 3 μm in size. Taxonomic annotations were performed as described in Methods.

Figure S8: Phylogenetic positions of NCLDV PolBs associated with CEE and network-based predicted eukaryotic host groups. The unrooted maximum likelihood phylogenetic tree contains environmental (labeled in red if VIP score > 2 and the regression coefficient is positive, labeled in blue if negative) and reference (labeled in black) sequences of *Prasinovirus* and *Mimiviridae* PolBs. The approximate SH-like local support values are shown in percentages at nodes, and the scale bar indicates one change per site. Host groups predicted at nodes are shown with colored circles. The red arrow points to a clade of viruses predicted to infect Prymnesiales.

61
Figure S9: Phylogenetic position of *Piconavirale*s RdRPs associated with CEE and network-
based predicted eukaryotic host groups. The unrooted maximum likelihood phylogenetic tree
contains environmental (labeled in red if VIP score > 2 and the regression coefficient is
positive, labeled in blue if negative) and reference (labeled in black) sequences of
*Piconavirale*s RdRPs. The approximate SH-like local support values are shown in
percentages at nodes, and the scale bar indicates one change per site. Host groups predicted at
nodes are shown with colored circles.

69
Figure S10: Carbon export efficiency (CEE) is correlated with the change in the slope of
particle size distribution (PSD) that occurred from the surface to deep (below the euphotic
zone). Observed PSDs were fitted in the form $n = ad^b$, where n is the frequency of particles of
a given size, d is the particle diameter, and a and b are parameters (as described by(Guidi et
al., 2008)). b, the PSD slope, is a proxy for particles size. For example $b = -5$ indicates
presence of a large proportion of smaller particles, whereas $b = -3$ indicates a preponderance
of larger particles. A higher b value at deep compared to surface is suggestive of aggregation
or presence of larger organisms at deep compare to surface. The blue line shows the
regression line between CEE and the PSD slope difference between surface and deep. The
shade around the regression line shows the 95% confidence interval.
80

Supplemental Tables

Table S1: Viral lineages associated with CEE

Viruses		VIPs	Positives VIPs	Negative VIPs
NCLDVs	Mimiviridae	34	25	9
	Phycodnaviridae	24	18	6
	Iridoviridae	2	0	2
	Other NCLDVs *	0	0	0
	Total	60	43	17
RNA viruses	Picornavirales (ssRNA+)	19	13	6
	Partitiviridae (dsRNA)	1	1	0
	Narnaviridae (ssRNA+)	0	0	0
	Other families	2*	0	2
	Unclassified	0	0	0
	RNA viruses	0	0	0
	Total	22	14	8
ssDNA viruses	Circoviridae	1	1	0
	Geminiviridae	0	0	0
	Nanoviridae	0	0	0
	Unclassified	0	0	0
	ssDNA viruses	0	0	0
	Total	1	1	0
	All	83	58	25

85 * Two Hepeviridae (ssRNA+).
86

Table S2: Assembly statistics for NCLDV metagenome-assembled genomes and corresponding VIPs

Metagenome-assembled genome	#contigs	N50	L50	Min	Max	Sum	VIPs OTUs (OM-RGC.v1 ID)
TARA_IOS_NCLDV_Bin_127_6	14	21,642	5	8,581	35,822	267,607	PolB 000079111
TARA_IOS_NCLDV_Bin_173_3	12	12,913	3	2,807	34,517	108,412	PolB 000248170
TARA_MED_NCLDV_Bin_284_10	34	10,936	10	2,580	29,722	298,760	PolB 000328966
TARA_MED_NCLDV_Bin_284_14	43	14,837	11	2,756	27,607	439,843	PolB 001175669
TARA_IOS_NCLDV_Bin_127_4	26	5,734	10	2,560	8,505	133,765	PolB 001064263 and 010288541
TARA_AON_NCLDV_Bin_289_4	17	9,468	5	3,044	26,201	153,728	PolB 000200745 and 002503270
TARA_MED_NCLDV_Bin_341_10	5	7,800	2	2,534	7,941	30,478	PolB 002682999
TARA_PON_NCLDV_Bin_65_10	35	13,866	11	3,781	43,080	382,455	PolB 000079078
TARA_PON_NCLDV_Bin_102_1	53	4,608	18	2,606	11,485	239,832	PolB 000495602
TARA_AON_NCLDV_Bin_133_8	8	7,204	3	2,686	10,349	51,009	PolB 000240662

N50: The length of the contigs for which half of the assembly size is contained in contigs with a length greater than N50.
L50: Number of contigs (or scaffolds) with a size greater or equal to N50.

92 **Table S3: Host predictions per viral and host group for 83 VIPs based on**
93 **phylogeny, co-occurrence analysis, and genomic context**

Virus-host relationship	Positive VIPs	Negative VIPs	Total
NCLDV-Mamiellales	10	4	14
NCLDV-Prymnesiales	5	1	6
NCLDV-Pelagophyceae	2	1	3
NCLDV-No prediction	26	11	37
RNA virus-Copepoda	7	2	9
RNA virus-Chaetocerotales	2	0	2
RNA virus-Labyrinthulomycetes	1	0	1
RNA virus-No prediction	4	6	10
ssDNA virus-Copepoda	1	0	1
Total	58	25	83

Table S4: Host prediction per viral OTU for 83 VIPs based on phylogeny, co-occurrence analysis, and genomic context

Virus types	Virus OTUs	Direction of association with CEE	Classification (LCA annotation)	Clade in phylogenic tree used for TIM analysis	TIM-based predicted host	MAGs ID	Genome-based predicted host	Suggested host	Note
NCLDVs	polb_000026723	negative	Mimiviridae	NA	NA	NA	NA	NA	
	polb_000030837	positive	Mimiviridae	Mimiviridae/Mesomimivirinae	Prymnesiales	NA	NA	Prymnesiales	
	polb_000042601	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_000054135	negative	Mimiviridae	Mimiviridae/Mesomimivirinae	Collodaria	NA	NA	Prymnesiales	
	polb_000061559	positive	Mimiviridae	Mimiviridae/Mesomimivirinae	Prymnesiales	NA	NA	Prymnesiales	
	polb_000061999	positive	Mimiviridae	Mimiviridae	NA	NA	NA	NA	
	polb_000073352	negative	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000073545	negative	Mimiviridae	Mimiviridae/CroV relative	NA	NA	NA	NA	
	polb_000079078	negative	Mimiviridae	Mimiviridae/AaV relative	NA	PON_NCLDV_Bin_65_10	Pelagophyceae	Pelagophyceae	
	polb_000079111	positive	Mimiviridae	Mimiviridae/AaV relative	NA	IOS_NCLDV_Bin_127_6	Pelagophyceae	Pelagophyceae	
	polb_000079365	positive	Mimiviridae	Mimiviridae	NA	NA	NA	NA	
	polb_000110630	negative	Mimiviridae	Mimiviridae	NA	NA	NA	NA	
	polb_000129518	positive	Mimiviridae	Mimiviridae/Mesomimivirinae	Prymnesiales	NA	NA	Prymnesiales	
	polb_000159717	positive	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_000161220	positive	Mimiviridae	Mimiviridae/AaV relative	NA	NA	NA	Pelagophyceae	
	polb_000172102	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_000194282	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	Mamiellales	NA	NA	Mamiellales	
	polb_000200745	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	Mamiellales	AON_NCLDV_Bin_289_4	NA	Mamiellales	
	polb_000229407	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000230224	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000232032	negative	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_000236849	negative	Phycodnaviridae	Phycodnaviridae/Prasinovirus	Mamiellales	NA	NA	Mamiellales	
	polb_000239928	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000240662	negative	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000248170	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	Mamiellales	IOS_NCLDV_Bin_173_3	NA	Mamiellales	
	polb_000249074	negative	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000249217	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000251540	negative	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_000328966	positive	Mimiviridae	NA	NA	NCLDV_Bin_284_10	NA	NA	
	polb_000396610	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_000435873	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	NA	NA	NA	Mamiellales	
	polb_000490625	positive	Mimiviridae	Mimiviridae	NA	NA	NA	NA	
	polb_000495602	negative	Iridoviridae	NA	NA	NCLDV_Bin_102_1	NA	NA	
	polb_000503865	positive	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_000673383	negative	Mimiviridae	Mimiviridae	NA	NA	NA	NA	
	polb_000844241	negative	Iridoviridae	NA	NA	NA	NA	NA	
	polb_000912507	positive	Mimiviridae	Mimiviridae/Mesomimivirinae	Collodaria	NA	NA	Prymnesiales	
	polb_001064263	positive	Phycodnaviridae	NA	NA	IOS_NCLDV_Bin_127_4	Mamiellales	Mamiellales	
	polb_001175669	positive	Mimiviridae	NA	NA	MED_NCLDV_Bin_284_14	NA	NA	
	polb_001527691	positive	Mimiviridae	Mimiviridae/Mesomimivirinae	NA	NA	NA	Prymnesiales	
	polb_002035391	positive	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_002503270	positive	Phycodnaviridae	Phycodnaviridae/Prasinovirus	Mamiellales	AON_NCLDV_Bin_289_4	NA	Mamiellales	
	polb_002682999	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_003145223	negative	Mimiviridae	NA	NA	NA	NA	NA	
	polb_003319665	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_003580241	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_004312996	positive	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_004775027	negative	Mimiviridae	NA	NA	NA	NA	NA	
	polb_004804559	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_007102163	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_007423474	negative	Mimiviridae	NA	NA	NA	NA	NA	
	polb_007503502	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_007771300	positive	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_008001141	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_010288541	positive	Phycodnaviridae	NA	NA	IOS_NCLDV_Bin_127_4	Mamiellales	NA	
	polb_013294654	positive	Phycodnaviridae	NA	NA	NA	NA	NA	
	polb_013433452	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_014364115	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_015514497	positive	Mimiviridae	NA	NA	NA	NA	NA	
	polb_015907472	positive	Phycodnaviridae	NA	NA	NA	NA	NA	
RNA viruses	rdrp_105714054	negative	Picornavirales	NA	NA	NA	NA	NA	
	rdrp_107558617	negative	Picornavirales	Picornavirales/Bacillarnavirus	NA	NA	NA	NA	
	rdrp_30787766	positive	Picornavirales	NA	NA	NA	NA	NA	
	rdrp_32150057	positive	Picornavirales	NA	NA	NA	NA	NA	
	rdrp_32150309	positive	Picornavirales	Picornavirales/Labyrnavirus	NA	NA	NA	Labyrinthulomycetes	*1
	rdrp_32202687	positive	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	
	rdrp_33049404	positive	Picornavirales	Picornavirales/Bacillarnavirus	Chaetocerotales	NA	NA	Chaetocerotales	
	rdrp_35179764	positive	Picornavirales	Picornavirales/Bacillarnavirus	Chaetocerotales	NA	NA	Chaetocerotales	
	rdrp_35713768	positive	Partitiviridae	NA	NA	NA	NA	NA	
	rdrp_36496887	positive	Picornavirales	Picornavirales	NA	NA	NA	NA	
	rdrp_36505302	positive	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
	rdrp_42335229	negative	Hepeviridae	NA	NA	NA	NA	NA	
	rdrp_49554577	negative	Picornavirales	Picornavirales	NA	NA	NA	NA	
	rdrp_54294427	positive	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
	rdrp_59731273	negative	Picornavirales	Picornavirales	NA	NA	NA	NA	
	rdrp_77677770	negative	Hepeviridae	NA	NA	NA	NA	NA	
	rdrp_77677810	negative	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
	rdrp_84897402	negative	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
	rdrp_8626697	positive	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
	rdrp_8855752	positive	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
	rdrp_9164160	positive	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
	rdrp_9164163	positive	Picornavirales	Picornavirales/Dicistroviridae	NA	NA	NA	Copepoda	*2
ssDNA viruses	rep_38177659	positive	Circoviridae	NA	NA	NA	NA	Copepoda	*3

Notes

*1 This virus was located in well-separated clade containing Aurantiochytrium single-stranded RNA virus (AsRNAV) which is known to infect Labyrinthulomycetes.

*2 These viruses were grouped within Dicistroviridae (known to infect insects) and may therefore infect marine arthropods such as copepods.

*3 This virus was connected with a copepod, mollusk and Collodaria OTUs in the co-occurrence network reconstructed for the mesoplankton size. Circoviridae-like viruses are known to infect copepod.

105 **Table S5: Statistics for the FlashWeave co-occurrence graphs**

Viral marker gene	Planktonic size fraction*	#Samples	#Viral OTUs	#Eukaryotic OTUs	#Edges in graph	#Virus-to-eukaryote edges	#Viruses connected to a eukaryote (%)
NCLDVs PolB	Piconano	99	2269	4936	20934	3594	1735 (76)
	Nano	51	1775	1872	6704	1027	721 (41)
	Micro	92	2205	2524	12189	2101	1299 (59)
	Meso	95	2238	2250	11624	1796	1126 (50)
RNA viruses RdRP	Piconano	60	125	4484	10754	446	122 (98)
	Nano	36	53	1768	2659	124	46 (87)
	Micro	62	124	2407	5351	367	117 (94)
	Meso	62	48	2100	4329	116	42 (88)
ssDNA viruses Rep	Piconano	60	64	4484	10577	205	63 (98%)
	Nano	36	1	1768	2563	2	1 (100%)
	Micro	62	4	2407	5086	9	4 (100%)
	Meso	62	8	2100	4242	24	8 (100%)

106 * Pico: 0.8 to 5 µm, Nano: 5 to 20 µm, Micro: 20 to 180 µm, Meso: 180 to 2000 µm
107

Table S6: Functional differences between eukaryotes found to be best connected to VIPs and non-VIPs

Functional trait	Positive VIPs ($n = 50$)		Non-VIPs ($n = 983$)		*P*-value (Fisher's exact test, two sided)	Adjusted *P*-value (BH) (*Q*)
	Presence	Absence	Presence	Absence		
Chloroplast	20	30	276	690	0.109	0.164
Silicification	11	39	60	920	0.000	0.001
Calcification	1	49	30	950	1.000	1.000
Functional trait	Negative VIPs ($n = 21$)		Non-VIPs ($n = 983$)		*P*-value (Fisher's exact test, two sided)	Adjusted *P*-value (BH) (*Q*)
	Presence	Absence	Presence	Absence		
Chloroplast	3	17	276	690	0.218	0.655
Silicification	0	21	60	920	0.632	0.947
Calcification	0	21	30	950	1.000	1.000

111 **Table S7: Functional differences between eukaryotes found to be best connected to**
112 **positive and negative VIPs**

Functional trait	Positive VIPs (n = 50)		Negative VIPs (n = 21)		*P*-value (Fisher extact test, two sided)	Adjusted *P*-value (BH) (*Q*)
	Presence	Absence	Presence	Absence		
Chloroplast	20	30	3	17	0.053	0.079
Silicification	11	39	0	21	0.027	0.079
Calcification	1	49	0	21	1.000	1.000

Transparent Methods

Data context

We used publicly available data generated in the framework of the *Tara* Oceans expedition.

Single-copy marker-gene sequences for NCLDVs and RNA viruses were identified from two

gene catalogs: the Ocean Microbial Reference Gene Catalog (OM-RGC) and the Marine Atlas

of *Tara* Oceans Unigenes (MATOU). The viral marker-gene read count profiles used in our

study are as previously reported for prokaryotic-sized metagenomes (size fraction 0.2–3 μm)

(Sunagawa et al., 2015) and eukaryotic-sized metatranscriptomes (Carradec et al., 2018).

Eukaryotic plankton samples (the same samples were used for metatranscriptomes,

metagenomes and 18S rRNA V9-meta-barcodes) were filtered for categorization into the

following size classes: piconano (0.8–5 μm), nano (5–20 μm), micro (20–180 μm), and meso

(180–2,000 μm). For eukaryotic 18S rRNA V9 OTUs (de Vargas et al., 2015), we used an

updated version of the data that included functional trait annotations (chloroplast-bearing,

silicified, and calcified organisms) of V9 OTUs. Occurrence profiles are compositional

matrices in which gene occurrence are expressed as unnormalized (V9 meta-barcode data) or

gene-length normalized (shotgun data) read counts. Indirect measurements of carbon export

($mg\ m^{-2}\ d^{-1}$) in 5-m increments from the surface to a 1,000-m depth were taken from Guidi et

al. (Guidi et al., 2016) The original measurements were derived from the distribution of

particle sizes and abundances collected using an underwater vision profiler. These raw data

are available from PANGEA (Picheral et al., 2014). Net primary production (NPP) data were

extracted and averaged from 8-day composites of the vertically generalized production model

(VGPM) (Behrenfeld and Falkowski, 1997) for the week of sampling. Thus, in this study, the

comparisons between NPP and other parameters were not made at the same time point. This

137   might have affected the results of the regression analysis, especially if there were any short-

138   term massive bloom events, although there was no bloom signal during the sampling period.

Carbon export, carbon export efficiency, and particle size distribution

Carbon flux profiles (mg m^{-2} d^{-1}) were estimated based on particle size distributions and

abundances. The method used for carbon flux estimation was previously calibrated comparing

sediment trap measurement and data from imaging instruments (Guidi et al., 2008). Carbon

flux values from depths of 30 to 970 meters were divided into 20-m bins, each obtained by

averaging the carbon flux values from the designated 20 m in profiles gathered during

biological sampling within a 25-km radius over 24 h when less than 50% of data were missing

(Figure S5). Carbon export (CE) was defined as the carbon flux at 150 m (Guidi et al., 2016).

Carbon export efficiency was calculated as follows: CEE = $CE_{deep}/CE_{surface}$ (Buesseler and

Boyd, 2009). To compare stations with different water column structures, we defined $CE_{surface}$

as the maximum CE (in a 20 m window) within the first 150 m. CE_{deep} is the average CE (also

in a 20 m window) 200 m below this maximum. The 150 m limit serves as a reference point

to automatize the calculation of $CE_{surface}$ and CE_{deep}. The 150m-depth layer was selected

because often used as a reference depth for drifting sediment trap and because most of the

deep chlorophyll maximum (DCM) were shallower except at two (stations 98 (175 m) and

100 (180 m)). The maximum $CE_{surface}$ for these two stations was above 150 m. The sampling

strategy of *Tara* Oceans was designed to study a variety of marine ecosystems and to target

well-defined meso- to large-scale features (based on remote-sensing data). Therefore, this

strategy avoided sampling water with important lateral inputs. Nevertheless, the possibility of

having locations with potential lateral transport cannot be excluded.

We obtained the particle size distribution (PSD) profiles generated by the *Tara* Oceans

expedition and computed the PSD slope at each depth for all profiles. The slope value

(denoted "*b*") is used as the descriptor of the particle size distribution as defined in a previous

work (Guidi et al., 2009). For example, $b = -5$ indicates the presence of a large proportion of

smaller particles, whereas $b = -3$ indicates a preponderance of larger particles. We averaged

the slope values at each sampling site in the same way as for carbon export flux.

Identification of viral marker genes from ocean gene catalogs

Viral genes were collected from two gene catalogs: OM-RGC version 1 and MATOU.

Sequences in these two gene catalogs are representatives of clusters of environmental

sequences (clustered at 95% nucleotide identity). The OM-RGC data were taxonomically re-

annotated, with the NCBI reference tree used to determine the last common ancestor modified

to reflect the current classification of NCLDVs (Carradec et al., 2018). We automatically

classified viral gene sequences as eukaryotic or prokaryotic according to their best BLAST

score against viral sequences in the Virus-Host Database (Mihara et al., 2016). DNA

polymerase B (PolB), RNA-dependent RNA polymerase (RdRP), and replication-associated

protein (Rep) genes were used as markers for NCLDVs, RNA viruses, and ssDNA viruses,

respectively. For PolB, reference proteins from the NCLDV orthologous gene cluster

NCVOG0038 (Yutin et al., 2009) were aligned using MAFFT-*linsi* (Katoh and Standley,

2013). A hidden Markov model (HMM) profile was constructed from the resulting alignment

using *hmmbuild* (Eddy, 2011). This PolB HMM profile was searched against OM-RGC amino

acid sequences and translated MATOU sequences annotated as NCLDVs, and sequences

longer than 200 amino acids that had hits with E-values $< 1 \times 10^{-5}$ were selected as putative

PolBs. RdRP sequences were chosen from the MATOU catalog as follows: sequences

assigned to Pfam profiles PF00680, PF00946, PF00972, PF00978, PF00998, PF02123,

PF04196, PF04197, or PF05919 and annotated as RNA viruses were retained as RdRPs. For

Rep, we reconstructed an HMM profile using a comprehensive set of reference sequences

(Kazlauskas et al., 2018) and searched this profile against the translated MATOU sequences

annotated as ssDNA viruses. We kept sequences that had hits with E-values $< 1 \times 10^{-5}$ and

removed those that contained frameshifts.

The procedure above identified 3,486 PolB sequences in the metagenomic samples and

respectively 975, 388, and 299 RdRP, PolB, and Rep sequences in the metranscriptomes.

Testing for associations between viruses with CEE, CE$_{150}$, and NPP

To test for associations between occurrence of viral marker genes and CEE, CE$_{150}$, and NPP,

we proceeded as follows. Samples with CEE values greater than one and with Z-score greater

than two were considered as outliers and removed (this removed the two samples from station

68). Only marker genes represented by at least two reads in five or more samples were

retained (lowering this minimal number of required samples down to three or four did not

improve the PLS regression model). To cope with the sparsity and composition of the data,

gene-length normalized read count matrices were center log-ratio transformed, separately for

ssDNA viruses, RNA viruses and NCLDVs. We next selected genes with Spearman

correlation coefficients with CEE, CE$_{150}$ or NPP greater than 0.2 or smaller than −0.2 (zero

values were removed). To assess the association between these marker genes and CEE, we

used partial least square (PLS) regression analysis. The number of components selected for

the PLS model was chosen to minimize the root mean square error of prediction (Figure S6).

We assessed the strength of the association between carbon export (the response variable) and

viral marker genes occurrence (the explanatory variable) by correlating leave-one-out cross-

validation predicted values with the measured carbon export values. We tested the

significance of the correlation by comparing the original Pearson coefficients between

explanatory and response variables with the distribution of Pearson coefficients obtained from

PLS models reconstructed based on permutated data (10,000 iterations). We estimated the

contribution of each gene (predictor) according to its variable importance in the projection

(VIP) score derived from the PLS regression model using all samples. The VIP score of a

predictor estimates its contribution in the PLS regression. Predictors with high VIP scores (>

2) were assumed to be important for the PLS prediction of the response variable.

Phylogenetic analysis

Environmental PolB sequences annotated as NCLDVs were searched against reference

NCLDV PolB sequences using BLAST. Environmental sequences with hits to a reference

sequence that had > 40% identity and an alignment length greater than 400 amino acids were

kept and aligned with reference sequences using MAFFT-*linsi*. Environmental RdRP

sequences annotated as were translated into six frames of amino acid sequences, and reference

RNA viruses RdRP sequences collected from the Virus-Host Database were searched against

the Conserved Domain Database (CDD) using rpsBLAST. The resulting alignment was used

to trim reference and environmental RdRP sequences to the conserved part corresponding to

the domain, CDD: 279070, before alignment with MAFFT-*linsi*. Rep sequences annotated as

ssDNA viruses were treated similarly. PolB, RdRP, and Rep multiple sequence alignments

were manually curated to discard poorly aligned sequences. Phylogenetic trees were

reconstructed using the the *build* function of ETE3 (Huerta-Cepas et al., 2016) of the

GenomeNet TREE tool (https://www.genome.jp/tools-bin/ete). Columns were automatically

trimmed using *trimAl* (Capella-Gutiérrez et al., 2009), and trees were constructed using

FastTree with default settings (Price et al., 2009).

A similar procedure was applied for the trees used in the hosts prediction analysis albeit

selecting sequences for the Phycodnaviridae/Mimiviridae (Figure S8) and the Picornavirales

(Figure S9) and removing the ones occurring in fewer than 10 samples, to reduce the size of

the tree.

Virus–eukaryote co-occurrence analysis

We used FlashWeave (Tackmann et al., 2019) with Julia 1.2.0 to predict virus–host interactions based on their co-occurrence patterns. Read count matrices for the 3,486 PolBs, 975 RdRPs, 299 Reps, and 18S rRNA V9 DNA barcodes obtained from samples collected at the same location were fed into FlashWeave. The 18S rRNA V9 data were filtered to retain OTUs with an informative taxonomic annotation. The 18S rRNAV9 OTUs and viral marker sequences were further filtered to conserve only those present in at least five samples. FlashWeave networks were learned for each of the four eukaryotic size fractions with the parameters 'heterogenous' = false and 'sensitive' = true, and edges receiving a weight > 0.2 and a Q-value < 0.01 (the default) were retained. The number of samples per size fraction ranged between 51 and 99 for NCLDVs and between 36 and 62 for RNA and ssDNA viruses. The number of retained OTUs per size fraction varied between 1,775 and 2,269 for NCLDVs and between 48 and 125 for RNA viruses (Table S5).

Mapping of putative hosts onto viral phylogenies

In our association networks, individual viral sequences were often associated with multiple 18S rRNA V9 OTUs belonging to drastically different eukaryotic groups, a situation that can reflect interactions among multiple organisms but also noise associated with this type of analysis (Coenen and Weitz, 2018). To extract meaningful information from these networks, we reasoned as follows. We assumed that evolutionarily related viruses infect evolutionarily related organisms, similar to the case of phycodnaviruses (Clasen and Suttle, 2009). In the interaction networks, the number of connections between viruses in a given clade and the associated eukaryotic host group should accordingly be enriched compared with the number of connections with non-host organisms arising by chance. Following this reasoning, we assigned the most likely eukaryotic host group as follows. The tree constructed from viral marker-gene sequences (PolB, RdRP or Rep) was traversed from root to tips to visit every

node. We counted how many connections existed between leaves of each node and the V9-

OTUs of a given eukaryotic group (order level). We then tested whether the node was

enriched compared with the rest of the tree using Fischer's exact test and applied the

Benjamini–Hochberg procedure to control the false discovery rate among comparisons of

each eukaryotic taxon (order level). To avoid the appearance of significant associations driven

by a few highly connected leaves, we required half of the leaves within a node to be

connected to a given eukaryotic group. Significant enrichment of connections between a virus

clade and a eukaryotic order was considered to be indicative of a possible virus–host

relationship. We refer to the above approach, in which taxon interactions are mapped onto a

phylogenetic tree of a target group using the organism's associations predicted from a species

co-occurrence-based network, as TIM, for Taxon Interaction Mapper. This tool is available at

https://github.com/RomainBlancMathieu/TIM. This approach can be extended to interactions

other than virus–host relationships.

Assembly of NCLDV metagenome-assembled genomes (MAGs)

NCLDV metagenome-assembled genomes (MAGs) were assembled from *Tara* Oceans

metagenomes corresponding to size fractions > 0.8 μm. Metagenomes were first organized

into 11 'metagenomic sets' based upon their geographic coordinates, and each set was co-

assembled using MEGAHIT (Li et al., 2015) v.1.1.1. For each set, scaffolds longer than 2.5

kbp were processed within the bioinformatics platform anvi'o (Eren et al., 2015) v.6.1

following methodology described previously for genome-resolved metagenomics (Delmont et

al., 2018). Briefly, we used the automatic binning algorithm CONCOCT (Alneberg et al.,

2014) to identify large clusters of contigs using both sequence composition and differential

coverage across metagenomes within the set. We then used HMMER (Eddy, 2011) v3.1b2 to

search for a collection of eight NCLDV gene markers (Guglielmini et al., 2019), and

identified NCLDV MAGs by manually binning CONCOCT clusters of interest using the

anvi'o interactive interface. The interface displayed hits for the eight gene markers alongside

coverage values across metagenomes and GC-content. Finally, NCLDV MAGs were

manually curated using the same interface, to minimize contamination as described previously

(Delmont and Eren, 2016).

Taxonomic composition of genes predicted in NCLDV genomes of VIPs

VIP's PolB sequences were searched (using BLAST) against MAGs reconstructed from the

metagenomes of the eukaryotic size fraction (> 0.8 μm) and against contigs used to produce

OM-RGCv1. Genome fragments covering 95% of the length of PolB VIPs with > 95%

nucleotide identity were considered as originating from a same viral OTUs. Genes were

predicted and annotated taxonomically with the same procedure described above

(identification of viral marker genes). Genes contained in viral genome fragments and

annotated as cellular organisms with amino acid identities > 60% were manually inspected

(Supplemental Data 2).

Statistical test

All the statistical significance assessments were performed with two-sided test.

Supplemental References
Alneberg, J., Bjarnason, B.S., Bruijn, I. de, Schirmer, M., Quick, J., Ijaz, U.Z., Lahti, L.,
Loman, N.J., Andersson, A.F., and Quince, C. (2014). Binning metagenomic contigs by
coverage and composition. Nat. Methods *11*, 1144–1146.

Behrenfeld, M.J., and Falkowski, P.G. (1997). Photosynthetic rates derived from satellite-
based chlorophyll concentration. Limnol. Oceanogr. *42*, 1–20.

Buesseler, K.O., and Boyd, P.W. (2009). Shedding light on processes that control particle
export and flux attenuation in the twilight zone of the open ocean. Limnol. Oceanogr. *54*,
1210–1232.

Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009). trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses. Bioinforma. Oxf. Engl.
25, 1972–1973.

Carradec, Q., Pelletier, E., Silva, C.D., Alberti, A., Seeleuthner, Y., Blanc-Mathieu, R., Lima-
Mendez, G., Rocha, F., Tirichine, L., Labadie, K., et al. (2018). A global ocean atlas of
eukaryotic genes. Nat. Commun. *9*, 373.

Clasen, J.L., and Suttle, C.A. (2009). Identification of freshwater Phycodnaviridae and their
potential phytoplankton hosts, using DNA pol sequence fragments and a genetic-distance
analysis. Appl. Environ. Microbiol. *75*, 991–997.

Coenen, A.R., and Weitz, J.S. (2018). Limitations of Correlation-Based Inference in Complex
Virus-Microbe Communities. MSystems *3*, e00084-18.

Delmont, T.O., and Eren, A.M. (2016). Identifying contamination with advanced visualization
and analysis practices: metagenomic approaches for eukaryotic genome assemblies. PeerJ *4*,
e1839.

Delmont, T.O., Quince, C., Shaiber, A., Esen, Ö.C., Lee, S.T., Rappé, M.S., McLellan, S.L.,
Lücker, S., and Eren, A.M. (2018). Nitrogen-fixing populations of Planctomycetes and
Proteobacteria are abundant in surface ocean metagenomes. Nat. Microbiol. *3*, 804–813.

Eddy, S.R. (2011). Accelerated Profile HMM Searches. PLOS Comput. Biol. *7*, e1002195.

Eren, A.M., Esen, Ö.C., Quince, C., Vineis, J.H., Morrison, H.G., Sogin, M.L., and Delmont,
T.O. (2015). Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ *3*,
e1319.

Guglielmini, J., Woo, A.C., Krupovic, M., Forterre, P., and Gaia, M. (2019). Diversification
of giant and large eukaryotic dsDNA viruses predated the origin of modern eukaryotes. Proc.
Natl. Acad. Sci. *116*, 19585–19592.

Guidi, L., Jackson, G.A., Stemmann, L., Miquel, J.C., Picheral, M., and Gorsky, G. (2008).
Relationship between particle size distribution and flux in the mesopelagic zone. Deep Sea
Res. Part Oceanogr. Res. Pap. *55*, 1364–1374.

Guidi, L., Stemmann, L., Jackson, G.A., Ibanez, F., Claustre, H., Legendre, L., Picheral, M.,
and Gorskya, G. (2009). Effects of phytoplankton community on production, size, and export
of large aggregates: A world-ocean analysis. Limnol. Oceanogr. *54*, 1951–1963.

Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlimi, A., Roux, S., Darzi, Y., Audic, S.,
Berline, L., Brum, J.R., et al. (2016). Plankton networks driving carbon export in the
oligotrophic ocean. Nature *532*, 465.

Huerta-Cepas, J., Serra, F., and Bork, P. (2016). ETE 3: Reconstruction, analysis and
visualization of phylogenomic data. Mol. Biol. Evol. msw046.

Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment software version
7: improvements in performance and usability. Mol. Biol. Evol. *30*, 772–780.

Kazlauskas, D., Varsani, A., and Krupovic, M. (2018). Pervasive Chimerism in the
Replication-Associated Proteins of Uncultured Single-Stranded DNA Viruses. Viruses *10*.

Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W. (2015). MEGAHIT: an ultra-fast
single-node solution for large and complex metagenomics assembly via succinct de Bruijn
graph. Bioinforma. Oxf. Engl. *31*, 1674–1676.

Mihara, T., Nishimura, Y., Shimizu, Y., Nishiyama, H., Yoshikawa, G., Uehara, H., Hingamp,
P., Goto, S., and Ogata, H. (2016). Linking Virus Genomes with Host Taxonomy. Viruses *8*,
66.

Picheral, M., Searson, S., Taillandier, V., Bricaud, A., Boss, E., Stemmann, L., Gorsky, G.,
Tara Oceans Consortium, C., and Tara Oceans Expedition, P. (2014). Vertical profiles of
environmental parameters measured from physical, optical and imaging sensors during Tara
Oceans expedition 2009-2013.

Price, M.N., Dehal, P.S., and Arkin, A.P. (2009). FastTree: Computing Large Minimum
Evolution Trees with Profiles instead of a Distance Matrix. Mol. Biol. Evol. *26*, 1641–1650.

Sunagawa, S., Coelho, L.P., Chaffron, S., Kultima, J.R., Labadie, K., Salazar, G.,
Djahanschiri, B., Zeller, G., Mende, D.R., Alberti, A., et al. (2015). Ocean plankton. Structure
and function of the global ocean microbiome. Science *348*, 1261359.

Tackmann, J., Matias Rodrigues, J.F., and von Mering, C. (2019). Rapid Inference of Direct
Interactions in Large-Scale Ecological Networks from Heterogeneous Microbial Sequencing
Data. Cell Syst. *9*, 286-296.e8.

de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., Lara, E., Berney, C.,
Le Bescot, N., Probert, I., et al. (2015). Ocean plankton. Eukaryotic plankton diversity in the
sunlit ocean. Science *348*, 1261605.

Yutin, N., Wolf, Y.I., Raoult, D., and Koonin, E.V. (2009). Eukaryotic large nucleo-
cytoplasmic DNA viruses: Clusters of orthologous genes and reconstruction of viral genome
evolution. Virol. J. *6*, 223.

Supplemental Data 1,2

Click here to access/download
Supplemental File Sets
Supplemental_Data_1_2.xlsx