Quantification of *nosZ* genes and transcripts in activated sludge microbiomes with novel group-

specific qPCR methods validated with metagenomic analyses

DaeHyun D. Kim[a], Doyoung Park[a]* , Hyun Yoon[a]†, Taeho Yun[a] , Min Joon Song[a], and Sukhwan

Yoon[a]#

[a]Department of Civil and Environmental Engineering, Korea Advanced Institute of Science and

Technology (KAIST)

Running Head: Quantification of *nosZ* with group-specific qPCR

#Address correspondence to Sukhwan Yoon, syoon80@kaist.ac.kr

*Present address: Department of Civil and Environmental Engineering, Georgia Institute of

Technology, Atlanta, Georgia, USA

†Present address: Department of Civil and Environmental Engineering, Cornell University, Ithaca,

New York, USA

Abstract

Substantial N_2O emission results from activated sludge nitrogen removal processes. The importance of

N_2O-reducers possessing NosZ-type N_2O reductases have been recognized as the only N_2O sink *in situ* key

to determination of the net N_2O emissions; however, reliable quantification methods for *nosZ* genes and

transcripts have yet to be developed. Here, *nosZ* genes and transcripts in activated sludge tank

microbiomes were analyzed with the group-specific qPCR assays designed *de novo* combining culture-

based and computational approach. A sewage sample was enriched in a batch reactor fed continuous

stream of N_2 containing 20-10,000 ppmv N_2O, where 14 genera of potential N_2O-reducers were identified.

All available amino acid sequences of NosZ affiliated to these taxa were grouped into five subgroups (two

clade I and three clade II groups), and primer/probe sets exclusively and comprehensively targeting the

subgroups were designed and validated with *in silico* PCR. Four distinct activated sludge samples from

three different wastewater treatment plants in Korea were analyzed with the qPCR assays and the results

were validated by comparison with the shotgun metagenome analysis results. With the validated qPCR

assays, the *nosZ* genes and transcripts of six additional activated sludge samples were analyzed and the

results of the analyses clearly indicated the dominance of two clade II *nosZ* subgroups (*Flavobacterium*-

like and *Dechloromonas*-like) among both *nosZ* gene and transcript pools.

Introduction

Nitrous oxide (N_2O) is one of the three major greenhouse gases with the largest contributions to

global warming, along with CO_2 and CH_4.[1] Although the contribution of N_2O is estimated to be only

~6% of the net greenhouse gas emissions in terms of CO_2eq, which is far less than those of CO_2 and

CH_4, eliminating one molecule of N_2O from the atmosphere has the same merit as removing ~300

molecule of CO_2 due to its high global warming potential. Besides, N_2O has also been the most

consequential ozone depletion agent.[2, 3] Thus, global efforts to curb the increase in atmospheric N_2O

concentration are necessitated for sustainable future. A better understanding of the biogeochemical

reactions functioning as N_2O sources and sinks in nitrogen-rich anthropogenic environments, e.g.,

fertilized agricultural soils and wastewater treatment plants (WWTPs), is especially important for N_2O

emission mitigation, as nitrification and denitrification, the biological reactions serving as the major

sources of N_2O, occur at consistent a basis in such environments and thus, they have been estimated

as the hotspots of N_2O emission to the atmosphere.[4, 5]

Particularly, the biological N_2O reduction mediated by Nos-type nitrous oxide reductases (NosZ) has

recently attracted immense scientific attention as the sole sink of N_2O on the Earth's surface at non-

elevated temperature.[6-9] This reaction had been, for long, known merely as one of the stepwise

reactions constituting the denitrification pathway.[4, 10] Only recently has the N_2O-to-N_2 reduction been

recognized as an independent energy-conserving reaction, as diverse organisms possessing either

conventional clade I and newly-discovered clade II *nosZ* were found to be capable of growth with

N_2O as the sole electron acceptor.[9, 11-13] Several independent research groups have reported difference

between clade I and clade II *nosZ*-possessing organisms in terms of their affinities to N_2O, and more

specifically, the organisms with *nosZ* closely affiliated to *Dechloromonas* spp. have been reported

with particularly low whole-cell Michaelis constants ($K_{m,app}$), suggesting that this group of organisms

may be involved in *in situ* consumption of low-concentration N_2O produced via diverse biotic and

abiotic processes [9, 13-16]. The microbial community developed in the laboratory-scale biofilter treating

100 ppmv N_2O included the clade II N_2O reducer *Flavobacterium* spp. as the most abundant *nosZ*-

carrying organisms, also supporting the significance of clade II N_2O as an important N_2O sink.[17, 18]

The NosZ-mediated N_2O reduction has recently been recognized to have an immensely important role

in various engineered wastewater treatment systems as the sole sink of N_2O produced from

denitrification and nitrification.[12, 19-23] In these recent studies, attempts have been made to correlate

N_2O reduction activities or net N_2O emissions to *nosZ* gene/transcript abundances or the *nosZ*-to-

(*nirS*+*nirK*) abundance ratios. Most, if not all, of these *nosZ* gene quantifications were performed

using SYBR Green quantitative polymerase chain reactions (qPCR) targeting the clade I (1840F: 5'-

CGCRACGGCAASAAGGTSMSSGT-3'/ 2090R: 5'-CAKRTGCAKSGCRTGGCAGAA-3') and

clade II *nosZ* (nosZ-II-F: 5'-CTIGGICCIYTKCAYAC-3' / nosZ-II-R: 5'-

SKSACCTTITTRCCITYICG-3').[24, 25] The reliability of these *nosZ* quantification results were

questionable, due to unverified specificity and low amplification efficiency, and even the simplest

questions as to whether clade I or clade II *nosZ* were more abundant in nutrient removal bioreactors

remains controversial.[12, 26, 27] In this study, we have developed TaqMan-based qPCR reactions

targeting four groups (two clade I *nosZ* groups and two clade II *nosZ* groups) of *nosZ* with

amplification efficiency above 90% and verified their coverage and specificity by comparing the

quantification data with the results of shotgun metagenome analyses. Anoxic activated sludge samples

from anoxic tanks of six conventional wastewater treatment plants in Korea with A2O (anaerobic-

anoxic-oxic) configuration were then analyzed with these novel qPCR assays, and without exception,

clade II domination of *nosZ* gene and transcript pools was verified.

Materials and methods

Sample collection

The wastewater inoculum for the N_2O enrichment experiments were grab-sampled from the anoxic

section of the activated sludge tank at Daejeon municipal WWTP (36º23'5" N 127º24'28" E) in

September 2016 (denoted as Daejeon1). Three activated sludge samples for validation of the qPCR

assays by comparison with the metagenomics data were collected at the same WWTP (Daejeon2) in

February 2019 and two other activated sludge WWTPs located in Gwangju and Gapyeong

(35°09'22.4"N 126°49'51.6"E and 37°49'00.1"N 127°31'13.0"E, respectively) in January and

February of 2019, respectively. Activated sludge samples from anoxic tanks of six other A2O

(anaerobic/anoxic/oxic) WWTPs were then collected for group-specific quantification of *nosZ* genes

and transcripts with the developed qPCR assays (Figure S1). Each wastewater sample for analyses of

DNA was collected in a 2-L polyethylene bottle filled up to the brim to minimize oxygen ingress. The

samples for quantification of *nosZ* transcripts were immediately mixed with the same volume of

methanol for RNA fixation.[28] The sample bottles were immediately placed in a cooler and

transported to the laboratory, where they were stored at -80°C until use.

Fed-batch enrichment of activated sludge samples and identification of active N_2O-reducing

groups

A simple fed-batch bioreactor was constructed for enrichment of active N_2O-reducers in the Daejeon1

wastewater sample (Figure S2). A 500-mL glass bottle with a side port (Duran, Mainz, Germany) was

fitted with a GL45 cap with three ports. Two of the ports were used as the inlet for N_2O-carrying gas

and the gas outlet from the bottle and the remaining port was used for aqueous phase sampling. The

side port of the glass bottle was used for sampling of the gaseous phase. The modified MR-1 medium

was prepared by adding per 1 L of deionized water, 0.5 g NaCl, 0.41 g sodium acetate, 0.23 g

KH_2PO_4, 0.46 g K_2HPO_4, 0.026 g NH_4Cl, 1 ml of 1000X trace metal solution, and 1 mL of 1000X

vitamin stock solution.[29] The reactor with 200-mL medium (40% of the total reactor volume) was

continuously supplied with 0 ppmv, 20 ppmv, 200 ppmv, or 10,000 ppmv N_2O prepared in >99.999%

N_2 gas (Samoh Specialty Gas, Daejeon, South Korea) for 30 hours before inoculation. After

inoculating the medium with 2 mL of the Daejeon1 sample, the same gas was bubbled through the

medium at the volumetric flowrate of 20 mL min^{-1} to provide the sole electron acceptor N_2O to the

microbial culture and maintain the reactor at anoxic condition. The reactor operated with >99.999%

N_2 gas served as the control to confirm that the microbial consortia were enriched with N_2O as the

electron acceptor and the contribution of O_2 contamination to microbial growth was kept to minimal.

The N_2O concentration in the headspace of the reactor was monitored with a HP6890 Series gas

chromatography fitted with an HP-PLOT/Q column and an electron capture detector (Agilent, Palo

Alto, CA), with the injector, oven and detector temperatures set to 200, 85, and 250 ℃, respectively.[30]

The O_2 concentration in the fed-batch reactor was monitored with a FireSting-O_2 oxygen meter

(Pyroscience, Aachen, Germany); however, due to the relatively high detection limit of the sensor (the

gas phase concentration of ~0.1% v/v), complete absence of O_2 was not guaranteed.

The growth of bacterial population in the reactor was monitored with qPCR using TaqMan chemistry

targeting the conserved region of eubacterial 16S rRNA genes. At each sampling time point, an 1.5-

mL aliquot was collected from the aqueous phase of the reactor and DNA was extracted from the

pellets using DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) following the protocol provided

by the manufacturer. Quantitative PCR was performed with the 1055f (5'-ATGGCTGTCGTCAGCT-

3') / 1392r (5'-ACGGGCGGTGTGTAC-3') / Bac1115Probe (5'-CAACGAGCGCAACCC-3')

primer and probe set targeting a conserved region of bacterial 16S rRNA genes using a QuantStudio3

platform (Thermo Fisher Scientific, Waltham, MA).[31] Incubation was halted when the population of

the enrichment reached a plateau, as indicated by three consecutive measurements without increased

gene counts. The reactor was dismantled and the aqueous phase was collected for microbial

composition analysis. The hypervariable V6−8 region of the 16S rRNA gene was amplified with 926F:

5′-AAACTYAAAKGAATTGRCGG-3′ / 1392R: 5′-ACGGGCGGTGTGTRC-3′ primer set and

MiSeq sequencing of the amplicons was outsourced to Macrogen Inc. (Seoul, Korea). The raw

sequence reads were deposited in the NCBI short reads archive (SRA) database (accession:

PRJNA552413) and were processed using the QIIME pipeline v 1.9.1 (detailed computational method

provided in the supplementary information).

Design of degenerate primers and probes for group-specific qPCR of *nosZ* genes

The genera assigned to the OTUs with the relative abundances higher than 0.3% in any of the reactor

microbial communities were selected. All *nosZ* gene sequences and corresponding translated NosZ

amino acid sequences belonging to the organisms affiliated to these genera were extracted from the

Uniprot (www.uniprot.org) database (accessed in March 2017) (Table S1). The curated pools of

nosZ/NosZ sequence data included, in total, 174 nucleotide sequences and the corresponding amino

acid sequences from 14 distinct genera. Subsequently, a multiple sequence alignment was performed

with these amino acid sequences, using MUSCLE algorithm with the parameters set to the default

values.[32] The NosZ phylogenetic tree was constructed using the neighbor-joining method in MEGA

7.0 with the bootstrap value set to 500.[33] The *nosZ* gene sequences were clustered into five groups

(NosZG1-5) according to the positions of the corresponding NosZ sequences in the phylogenetic tree,

which consisted of five phylogenetically distinct subbranches.

For each *nosZ* group, a primers and probe set was designed using the PriMux software to

comprehensively and exclusively target the *nosZ* gene sequences within the group.[34] The parameters

were modified from the default values to obtain the optimal candidate degenerate oligonucleotide

sequences for qPCR (length: 18-24 bp, amplicon size: 80-400 bp, T_m: 56-62°C for primers and 68 -

72°C for probes). Several candidate primers and probe sets were generated for each *nosZ* group,

implementing the ***min***, ***max***, and ***combo*** algorithms of the Primux software. The performance of each

candidate primers and probe set was predicted with *in silico* PCR performed with the simulate_PCR

software against all complete genome sequences of the organisms belonging to each of the five *nosZ*

groups (Table S1).[34, 35] Coverage within the target *nosZ* group and mutual exclusivity across the

groups were the two major criteria for assessment of the candidate primers and probe sets. The *in*

silico PCR tests were also performed against the complete genomes of 50 bacterial strains lacking

nosZ to preclude the possibility of unspecific amplification.

Construction of calibration curves for the designed primer and probe sets using model N_2O

reducer strains

A model organism from each group of the selected *nosZ* gene(s) was used for construction of the

calibration curve. The selected model organisms were *Pseudomonas stutzeri* DCP-Ps1 (NosZG1),

Acidovorax soli DSM25157 (NosZG2), *Flavobacterium aquatile* LMG4008 (NosZG3),

Ignavibacterium album JCM16511 (NosZG4), and *Dechloromonas aromatica* RCB (NosZG5).

Acidovorax soli DSM25157 and *F. aquatile* LMG4008 were acquired from Korean Collection for

Type Cultures and *I. album* JCM16511 from Japanese Collection of Microorganisms. The axenic

batch cultures of these organisms were prepared as previously described in the literature or using the

media and incubation conditions specified by the distributors. The cells were harvested at $OD_{600nm} =$

0.1 and the *nosZ* gene(s) in the DNA extracted from the cell pellets were amplified using the designed

primers. The calibration curve for each qPCR reaction was constructed using ten-fold serial dilutions

of PCR®2.1 vectors (Invitrogen, Carlsbad, CA) carrying the *nosZ* amplicons. A uniform thermocycle

was used for the qPCR: 95°C for 10 min and 40 cycles of 95 °C for 30 s, 58 °C for 60 s, and 72 °C for

60 s.

Group specific quantitative PCR targeting *nosZ* gene and transcripts in activated sludge

samples

The group-specific quantification of the *nosZ* genes and transcripts in the activated sludge samples

were performed with the designed primer and probe sets (Table 1) on a QuantStudio 3 real-time PCR

instrument using TaqMan detection chemistry (FAM as the reporter and NFQ-MGB as the quencher).

DNeasy Blood & Tissue Kit (QIAGEN) was used to extract DNA from the activated sludge samples.

From the methanol-treated samples, RNA was extracted with RNeasy Mini Kit (Qiagen), purified

with DNase I (Qiagen) and RNeasy MinElute Cleanup Kit (Qiagen), and reverse-transcribed with

Superscript® III (Thermo Fisher Scientific), as described previously.[36] The luciferase control mRNA

(Promega, Madison, WI, USA) was added as the internal standard to account for RNA loss during the

process. Each 20-µL qPCR reaction mix contained 10 µL of 2X TaqMan master mix (Applied

Biosystems, Foster city, CA, USA), 5 µM each of the forward and reverse primers, 0.5 µM of the

probe, and 2 µL of the DNA or cDNA. Calibration curves prepared with dilution series of the PCR

amplicons were used to calculate the copy numbers of the targeted genes from the C_t values.

Eubacterial 16S rRNA genes in the extracted DNA samples were quantified using the

1055F/1392R/Bac115Probe set for quantification of total bacterial population in the activated sludge

samples (Table S2). The copy numbers of the targeted *nosZ* genes in the wastewater samples were

normalized with the 16S rRNA gene copy numbers to facilitate comparison with the relative

abundances of the *nosZ* groups from the metagenomic analyses.

The PCR amplicons of the Daejeon1 sample amplified with NosZG1-5 primer sets were sequenced

using Illumina Miseq platform (San Diego, CA) at the Center for Health Genomics and Informatics at

University of Calgary. The raw sequence reads have been deposited in the SRA database (accession

number: PRJNA552418). After quality trimming and merging of the paired-end sequences, the

sequences without the probe-binding region were removed, and the remaining reads were clustered

into OTUs with 0.97 cut-off using cd-hit-est v. 4.6.[37] The OTUs were annotated using blastx against

the bacterial Refseq database downloaded in June, 2018, with the e value cut off set to 10^{-3} and word

size to 3 and no seg option selected,

Computational quantification of *nosZ* genes from shotgun metagenomes of the activated sludge

samples

The DNA samples for shotgun metagenome sequencing were extracted with DNeasy PowerSoil Kit

(Qiagen) from 50 mL each of the four activated sludge samples collected for validation of the qPCR

assays. Sequencing of the metagenomic DNA was performed at Macrogen Inc., where Hiseq X Ten

sequencing platform (Illumina, San Diego, CA) was used for generating 5-10 Gb of paired-end reads

data with 150-bp read length. The raw sequence reads have been deposited in the NCBI short reads

archive (SRA) database (accession numbers: PRJNA552406).

The raw reads were then processed using Trimmomatic v0.36 software with the parameters set to the

default values.[38] The trimmed reads were translated *in silico* into amino acid sequences using all six

possible reading frames and screened for clade I and II *nosZ* sequences using hidden Markov models

(HMM). The two HMM algorithms for clade II *nosZ* were downloaded from the Fungene database

(accessed in October 2016). As the HMM for clade I *nosZ* in the database had specificity issues, the

HMM algorithm was constructed *de novo* using *hmmbuild* command of HMMER v3.1b1. The clade I

NosZ sequences used to build the HMM were manually curated from the pool of NosZ sequences

downloaded from the NCBI database (accessed in October 2016), to represent diverse subgroups

within the clade (Table S3).[8, 11] The candidate partial NosZ sequences were extracted from the

translated shotgun metagenome reads using the *hmmsearch* command of HMMER v3.1b1 with the e

value cutoff set to 10^{-5}. The nucleotide sequence reads corresponding to the extracted partial NosZ

sequences (in separate bins for clade I and clade II NosZ) were assembled into contigs using

metaSPAdes v3.12.0 with parameters set to default values.[39] The assembled contigs with lengths

shorter than 200 bp were filtered out. The overlapping contigs appearing in both clade I and clade II

NosZ bins were identified by clustering the two sets of contigs against each other with a nucleotide

identity cutoff of 1.0. These overlapping contigs were manually called to the correct bin according to

the BLASTX results. The trimmed sequence reads were then mapped onto the contigs in the *nosZ*

bins using Bowtie2 v2.2.6, yielding sequence alignments for the contigs and the mapped reads. The

alignment files were further processed using samtools v0.1.19, and the PCR duplicates were removed

using MarkDuplicates function of picard-tools v1.105.[40] The sequence coverages of the contigs were

calculated using bedtools v2.17.0. The number of reads mapped on to the contigs were normalized

with the lengths of the respective contigs. The contigs were assigned taxonomic classification using

blastx, and distributed to the NosZG1-5 bins based on their taxonomic affiliations (Table S4). The

contigs without matching sequence were binned as 'other *nosZ* sequences'. The attempt to use 16S

rRNA sequences extracted with Meta-RNA and assembled with EMIRGE as the template for

mapping was not successful, as the coverage of the extracted 16S rRNA sequences turned out to be an

order of magnitude lower than the *rpoB* coverage (Data not shown).[41, 42] Thus, the sequence coverage

of *rpoB* gene, a single-copy housekeeping gene, was used for normalization of the *nosZ* abundance

data. The HMM algorithm for *rpoB* was downloaded from the Fungene database.

Results

Active N$_2$O reducers enriched in fed-batch incubation with varying N$_2$O concentrations

The microbial communities of the three N$_2$O-reducing enrichments, each prepared with different N$_2$O

concentrations, were analyzed to identify active N$_2$O-reducing groups of microorganisms (Table S5).

Screening for operational taxonomic units (OTUs) with >0.3% abundance in any of the three

enrichments yielded 69 OTUs assigned to 33 genera in total, and 14 of these genera were identified

with phylogenetic subgroups harboring clade I or clade II *nosZ*. The OTUs belonging to these

putatively *nosZ*-harboring genera amounted to 50.8% – 63.2% of the total microbial population in the

enrichments. The abundances of the genera putatively harboring clade II *nosZ* were observed to be

greater than those of the genera harboring clade I of *nosZ* (Table S5). In the enrichment incubated

with 20 ppmv N$_2$O, *Cloacibacterium* (18.9%), *Flavobacterium* (14.2%), and *Acidovorax* (13.5%)

were identified as the dominant genera. *Dechloromonas* (17.3%) and *Flavobacterium* (15.8%) were

the dominant genera in the 200 ppmv enrichment, and *Dechloromonas* was the predominant

population in the enrichment incubated with 10,000 ppmv of N$_2$O, constituting 46.0% of the total

microbial population.

Designing of the degenerate *nosZ* primer/probe sets

With the 174 translated NosZ sequences affiliated to the 14 genera putatively harboring active N_2O-

reducing organisms identified in the activated sludge enrichments, a phylogenetic tree composed of

five distinct branches (NosZG1 – NosZG5) was constructed (Figure S3). NosZG1 and NosZG2 were

identified as clade I NosZ and NosZG3-NosZG5 as clade II NosZ. After the iterative process of

designing candidate primers and probe sets and performing *in silico* PCR tests, the final sets of

degenerate primers/probe sets were designed to comprehensively and exclusively target the five *nosZ*

groups (Table 1). The *in-silico* PCR performed against 174 genomes from which the target *nosZ*

sequences were extracted and 50 genomes without *nosZ* confirmed the high levels of coverage (57.3-

100%) and complete exclusivity for all five degenerate primers/probe sets (Table 1). The qPCR

calibration curves were constructed with the selected model organisms (Figure S4, Table S6), and

despite the high levels of degeneracy (up to 110592), amplification efficiencies above 90% were

attained for all five primer and probe sets after rigorous optimization process.

Cross-checking of the group-specific qPCR quantification of *nosZ* genes with shotgun

metagenome analyses

The *nosZ* gene abundances in four distinct activated sludge samples were quantified using the newly

designed qPCR assays, and the *nosZ* sequences extracted from the shotgun metagenomes of these

same samples were quantitatively analyzed in parallel, for cross-checking of the qPCR results (Figure

1). The qPCR results of the four activated sludge samples invariably showed the dominance of the

clade II *nosZ* genes (NosZG3 and NosZG5) over clade I *nosZ* genes (NosZG1 and NosZG2), with at

least five-fold higher copy numbers (Figure 2). The qPCR targeting the NosZG4 group failed to

amplify the clade II *nosZ* belonging to this group. The NosZG4 primers/probe set was designed from

a single complete genome (*Ignavibacterium album*) and six sequences from metagenome assembled

genomes (MAGs), which may have been insufficient to cover the sequence divergence of this *nosZ*

subgroup.

The distribution of the *nosZ* sequences extracted from the shotgun metagenomes (Table S7) were also

severely biased towards clade II. The compositions of the *nosZ* genes exhibited high level of

similarity across the activated sludge samples. The calculated relative abundances of the two most abundant groups, NosZG3 and NosZG5, were relatively consistent across the samples, varying by less than 1.7-fold. The contributions of NosZG1 and NosZG2 to the total *nosZ* gene abundances were minor in all of the samples and largely variable. The fold differences between the samples with the highest relative abundance of NosZG1 and NosZG2 and those with the lowest abundance were 3.5 and 2.3, respectively. The *nosZ* genes sorted as NosZG4 constituted <1.2% of the total *nosZ* genes grouped as NosZG1-NosZG5, suggesting that this *nosZ* group has less significant role in N_2O reduction in activated sludge tanks than other analyzed groups.

As the qPCR data were normalized with the eubacterial 16S rRNA copy numbers and the metagenome-derived relative abundance data were normalized with the sequence coverages of the single-copy housekeeping gene *rpoB*, direct comparison of the outcomes was not possible. Theoretically, however, the ratios of the *nosZ*/*rpoB* (metagenome) to *nosZ*/16S (qPCR) could be used as indicators of reliability of the qPCR assays, with consistency in the ratios across the *nosZ* groups indicating high reliability. The *nosZ*/*rpoB*-to-*nosZ*/16S ratios were within a narrow range between 1.6 and 2.4 for NosZG5 across the four activated sludge samples. For an unidentified reason, broad gaps between the NosZG1-3 qPCR and the metagenomics data were observed with the Daejeon1 sample. Excluding these results, the *nosZ*/*rpoB*-to-*nosZ*/16S ratio varied from 3.3 to 7.2 for NosZG3, and the *nosZ*/*rpoB*-to-*nosZ*/16S ratios of NosZG1 and NosZG2 ranged from 1.9 to 2.1 and from 1.9 to 9.2, respectively. The *nosZ*/16S-to-*nosZ*/*rpoB* ratios of 12 out of the 16 qPCR assays performed (excluding NosZG4) were within the range between 1.5 and 9.2, indicating that the group-specific qPCR assays enabled reliable quantification of *nosZ* genes.

The *nosZ* amplicons of the Daejeon1 sample, amplified with the newly-designed qPCR primers, were sequenced and analyzed to check whether the qPCR reactions were comprehensive and mutually exclusive (Figure 3). As the NosZG4 primers failed to amplify the targeted genes, NosZG4 was excluded from the downstream analysis. The *nosZ* genes of putative active N_2O reducers captured by the NosZG1 and NosZG2 primer sets accounted for 54.2% of the entire pool of the clade I *nosZ* genes

extracted from the shotgun metagenome. The NosZG3 and NosZG5 primer sets captured 63.4% of the clade II *nosZ* genes recovered from the metagenome (Figure S5). No amplification outside the targeted group occurred for any of the primers, confirming the complete mutual exclusivity of these primers. Each of the qPCR assays were comprehensive within its target groups, as analyzed *nosZ* amplicon sequences covered all genera originally targeted by each primer and probe set.

Overall, the OTUs identified to be abundant from sequence analyses of the PCR amplicons coincided with the dominant *nosZ* OTUs from shotgun metagenome analysis. The OTUs affiliated to *Rhodobacter capsulatus*, Rhodobacteraceae bacterium QY30, *Hyphomicrobium denitrificans*, and *Bradyrhizobium* sp. were the dominant OTUs among the OTUs NosZG1 amplicons. The two most abundant *nosZ* OTUs among the NosZG3 amplicons were affiliated to *Flavobacterium columnare* and *Niastella koreensis*, which were also the most abundant *nosZ* taxa according to the metagenome analyses. Discrepancies between the amplicon sequencing data and the shotgun metagenome data were observed for NosZG2 and NosZG5 to some degree. The OTUs assigned to the genus *Thauera* and *Ruvrivivax gelatinosus* were the dominant among NosZG2, constituting 50.4% of the amplicons, but together constituted only 2.2% of the metagenome-derived *nosZ* sequences belonging to this group. Instead, an OTU affiliated to *Ramlibacter tataouinensis*, closely related to *R. gelatinosus*, was recovered in high relative abundance (15.4%) in the metagenome-derived *nosZ* pool. Likewise, the low relative abundance of the OTUs affiliated to *Dechloromonas aromatica* (0.2%) in the NosZG5 amplicons was coupled to the high relative abundance of the *nosZ* OTU affiliated to the *Azospira oryzae* (54.6%) with >96% translated amino acid identity within the amplified region. Thus, the observed discrepancy may be due to ambiguous OTU assignment among the closely related taxa.

The TaqMan-based qPCR assays were also compared with the most frequently used qPCR assays for clade I and clade II *nosZ* using SYBR green detection chemistry (Figure S6). Despite the low amplification efficiency (79.3%), the abundances of clade I *nosZ*, as determined with the SYBR Green assays, were relatively consistent with the cumulative abundances of NosZG1 and NosZG2 determined with the TaqMan qPCR assays, except for Daejeon1 sample, where the SYBR Green

qPCR yielded 87 times lower copy numbers than the TaqMan qPCR. The SYBR Green qPCR with

nosZII-F–nosZII-R primer set, with a subpar amplification efficiency of 61.9%, underestimated the

clade II *nosZ* copy numbers by several orders of magnitudes for three of the four analyzed samples,

with the largest difference (1.8×10^3-fold) observed with the Daejeon1 sample. The importance of the

clade II *nosZ* among the N_2O-reducing microbial community in these activated sludge samples would

have gone unnoticed due to this underestimation, if the conventional qPCR assays were used as the

sole means of *nosZ* quantification.

Quantitative analyses of *nosZ* genes and transcripts in activated sludge microbiomes

With the new group-specific *nosZ* primers, the activated sludge microbiomes from anoxic tanks of six

A2O WWTPs apart from those used for development and cross-checking processes were analyzed for

the *nosZ* gene and transcript abundances (Figure 4). As observed with the samples from Daejeon1,

Daejeon2, Gwangju, and Gapyeong, the clade II *nosZ* (amplified with NosZG3 and NosZG5), were at

least three-fold more abundant than the clade I *nosZ* (amplified with NosZG1 and NosZG2) in terms

of gene abundance. The *Dechloromonas*-like *nosZ* genes (NosZG5) was the most abundant *nosZ*

group in all samples but from Busan WWTP, where the abundance of *Flavobacterium*-like *nosZ*

genes (NosZG3) was statistically similar to this group. The transcript profiles further highlighted the

significance of clade II *nosZ* in activated sludge microbiomes. Due to the lower gene abundance and

generally low level of transcription observed for clade I *nosZ* (transcript-to-gene ratios of <1.0 for

both NosZG1 and NosZG2 in five out of six samples), transcription of the clade I *nosZ* was at least an

order of magnitude lower than that of the clade II *nosZ* in all samples. The *Pseudomonas*-like *nosZ*

(NosZG1) had remarkably low transcription level (transcript-to-gene ratios of <0.01) in all samples

but from Busan WWTP, suggesting irrelevance of this *nosZ* group as an N_2O sink in the activated

sludge tanks. The relative importance of NosZG3 and NosZG5 is difficult to fathom, as *nosZ*

transcripts targeted by NosZG3 were significantly more abundant in the sample from Busan WWTP

($p<0.05$), while NosZG5 was significantly more abundant in the samples from three other WWTPs

($p<0.05$).

Discussion

Quantification of the functional genes encoding the nitrogen cycle enzymes (e.g., bacterial/archaeal

amoA, comammox *amoA*, *nrfA*, and clade I and II *nosZ*) has been used in increasing number of

studies across diverse disciplines of environmental sciences and engineering, as predictors of nitrogen

cycling activity in diverse soil and aquatic environments.[43-46] Despite the frequent use of qPCR in

determining the potential N_2O-reducing populations, i.e., clade I and clade II *nosZ*-posessing

microorganisms, the predictability of qPCR quantification has rarely been scrutinized in the previous

studies. The new group-specific qPCR assays exhibited definite advantages over these frequently used

primers. Although the targets were limited to the groups of *nosZ*-possessing organisms enriched with

N_2O in *ex situ* cultures, the copy numbers obtained were, in many cases, orders of magnitudes larger

than those measured using previous primer sets. The diversity of the sequences recovered in the

amplicon sequence indicates the exhaustive coverages of the qPCR assays developed in this study

within the targeted groups. Most of the major *nosZ* OTUs (>1% in relative abundance) recovered in

metagenome-based *nosZ* profiling of the examined activated sludge samples were amplified with

exactly one of the four primer sets, thus confirming the absolute mutual exclusiveness of the

primers/probe sets. Any methods for quantification of functional genes are prone to error, and the

nosZ qPCR methods developed here also have certain deficiencies, including inability to amplify *nosZ*

genes affiliated to relatively abundant *Iganvibacterium* spp. In designing the group-specific primers, a

trade-off between coverage and mutual exclusivity was inevitable, which could have been the cause

of modest discrepancy with the metagenome data. Nevertheless, despite these drawbacks, there is

little doubt that the qPCR assays developed in this study are the most reliable tools for real-time

quantification of the environmentally significant *nosZ* genes available to date.

The clade II *nosZ*-possessing organisms of the *Flavobacterium* (amplified with NosZG3),

Dechloromonas, and *Azospira* (amplified with NosZG5) genera have previously been characterized

with high-affinity reduction of N_2O.[13, 14] The whole-cell half-saturation constants as low as 0.324 μM

has been reported for these groups of N_2O reducers. The *nosZ* gene and transcript pools of all six of

the activated sludge samples were dominated by these clade II *nosZ* (targeted by NosZG3 and

NosZG5). This observation, along with the high transcript-to-gene ratios support the previous

hypothesis that the clade II *nosZ*, with higher affinity to N_2O, may have consequential role in reducing

N_2O emissions from soil and aquatic environments, including the activated sludge tanks.[15, 47] The low

transcript-to-gene ratios of the clade I *nosZ* (<0.1 in five out of six samples) targeted by NosZG1 and

NosZG2 were also notable in the *nosZ* transcript profiles. The organisms possessing clade I *nosZ*

genes, being almost exclusively denitrifiers, may favor upstream steps of denitrification reaction in

the anoxic activated sludge tanks, where NO_3^- and/or NO_2^- are constantly available at millimolar

concentrations and N_2O is immediately consumed by their high-affinity cohabitants, which may

prefer N_2O to NO_3^- or NO_2^-. More physiological evidences are warranted in the future research,

however, to verify whether such hypothetical 'division of labor' really exists, as any further

extrapolation from the *nosZ* transcription data alone would be an over-interpretation.

The major *nosZ*-possessing organisms identified in the qPCR and metagenomic analyses differ greatly

from the major *nosZ*-possessing populations of the agricultural soils analyzed previously with shotgun

metagenome sequencing.[48] None of the six most abundant *nosZ* phylogenetic groups in the Havana

and Urbana agricultural research site soils (the *nosZ* genes affiliated to *Anaeromyxobacter, Opitutus*,

Hydrogenobacter, *Ignavibacterium*, *Dyedobacter*, and *Gemmatimonas*) was identified as a major

population in any of the activated sludge samples examined in this study. Neither were they enriched

with N_2O to high relative abundance (e.g., >1%) in the fed-batch reactor. Of the organisms affiliated

to these genera, *Anaeromyxobacter dehalogenans* and *Gemmatimonas aurentiaca* have been

confirmed of N_2O reduction activity; however, the N_2O reduction rates measured *in vitro* were orders

of magnitudes lower for these organisms than other examined N_2O-reducing organisms throughout

the entire range of N_2O concentration, and *G. aurantiaca* lacked the capability to utilize N_2O as the

growth substrate.[11, 13, 30] The differences in the compositions of the *nosZ*-possessing populations may

be attributed to the inherent difference in the rates of the biochemical turnover processes in WWTP

activated sludges and soils. Rapid nitrogen cycling reactions take place on a constant basis in WWTPs,

while the time scale of the biogeochemical processes in agricultural soils is orders of magnitude

longer and often, nitrogen supply provided through fertilization and plant exudation is sparse and

sporadic.[49] Besides, upland agricultural soils are often well-aerated and thus, the capability of N_2O

utilization would not provide specific benefits to the *nosZ*-harboring organisms at ordinary dry

conditions.[50] In fact, incubation of soil microbial consortium with N_2O in the same fed-batch

cultivation resulted in enrichment of the same organismal groups as those observed in the activated

sludge enrichments (e.g., *Pseudomonas*, *Flavobacterium*, *Acidovorax*, and *Chryseobacterium*),

suggesting that under occasions of pulse stimulation of denitrification and N_2O reduction, e.g.,

flooding events immediately following fertilization, these organisms may become the relevant N_2O

sinks in soils, as well (Table S8). Besides, none of the 14 *nosZ*-carrying genera enriched with N_2O in

this study from either the activated sludge and soil belonged to the taxa identified as non-denitrifying

N_2O reducers suggesting that the abundance of non-denitrifier taxa, *per se*, may not be a measure of

the N_2O sink capability, as previously proposed.

Acknowledgements

This work was financially supported by "the R&D Center for reduction of Non-CO2 Greenhouse

Gases (Grant No. 2017002420002)" funded by Korea Ministry of Environment (MOE).

Reference

(1)   Ciais, P.; Sabine, C.; Bala, G.; Bopp, L.; Brovkin, V.; Canadell, J.; Chhabra, A.;

DeFries, R.; Galloway, J.; Heimann, M.; Jones, C.; Le Quere, C.; Myneni, R. B.; Piao,

S.; Thornton, P., Carbon and other biogeochemical cycles. In *Climate change 2013:*

The physical science basis. Contribution of working group I to the fifth assessment

report of the Intergovernmental Panel on Climate Change, Stocker, T. F.; Qin, D.;

Plattner, G.-K.; Tignor, M.; Allen, S. K.; Boschung, J.; Nauels, A.; Xia, Y.; Bex, V.;

Midgley, P. M., Eds. Cambridge University Press: Cambridge, United Kingdom, New

York, NY, USA, 2013; pp 465-570. http://www.ipcc.ch/pdf/assessment-

report/ar5/wg1/WG1AR5_Chapter06_FINAL.pdf

(2)   Ravishankara, A. R.; Daniel, J. S.; Portmann, R. W. Nitrous oxide (N_2O): the dominant

472   ozone-depleting substance emitted in the 21[st] century. *Science* **2009,** *326*, (5949), 123-

473   125.

474   (3)   Portmann, R. W.; Daniel, J. S.; Ravishankara, A. R. Stratospheric ozone depletion due

475   to nitrous oxide: influences of other gases. *Phil. Trans. R. Soc. B* **2012,** *367*, (1593),

476   1256-1264.

477   (4)   Yoon, S.; Song, B.; Phillips, R. L.; Chang, J.; Song, M. J. Ecological and physiological

478   implications of nitrogen oxide reduction pathways on greenhouse gas emissions in

479   agroecosystems. *FEMS Microbiol. Ecol.* **2019**, *95*, (6), fiz066.

480   (5)   Law, Y.; Ye, L.; Pan, Y.; Yuan, Z. Nitrous oxide emissions from wastewater treatment

481   processes. *Phil. Trans. R. Soc. B* **2012,** *367*, (1593), 1265-1277.

482   (6)   Thomson, A. J.; Giannopoulos, G.; Pretty, J.; Baggs, E. M.; Richardson, D. J. *Philos*

483   *Trans Royal Soc B* **2012,** *367*, (1593), 1157-1168.

484   (7)   Frutos, O. D.; Quijano, G.; Aizpuru, A.; Muñoz, R. A state-of-the-art review on nitrous

485   oxide control from waste treatment and industrial sources. *Biotechnol. Adv.* **2018,** *36*,

486   (4), 1025-1037.

487   (8)   Hallin, S.; Philippot, L.; Löffler, F. E.; Sanford, R. A.; Jones, C. M. Genomics and

488   ecology of novel N_2O-reducing microorganisms. *Trends Microbiol.* **2017,** *26*, (1), 43-

489   55.

490   (9)   Suenaga, T.; Hori, T.; Riya, S.; Hosomi, M.; Smets, B. F.; Terada, A. Enrichment,

491   isolation, and characterization of high-affinity N_2O-reducing bacteria in a gas-

492   permeable membrane reactor. *Environ. Sci. Technol.* **2019**, *in press*; DOI:

493   10.1021/acs.est.9b02237

494 (10) Zumft, W. G. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol.*

495   *Rev.* **1997,** *61*, (4), 533-616.

(11) Sanford, R. A.; Wagner, D. D.; Wu, Q.; Chee-Sanford, J. C.; Thomas, S. H.; Cruz-

García, C.; Rodríguez, G.; Massol-Deyá; Krishnani, K. K.; Ritalahti, K. M.; Nissen, S.;

Konstantinidis, K. T.; Löffler, F. E. Unexpected nondenitrifier nitrous oxide reductase

gene diversity and abundance in soils. *Proc. Natl Acad Sci* **2012,** *109*, (48), 19709-

19714.

(12) Conthe, M.; Wittorf, L.; Kuenen, J. G.; Kleerebezem, R.; van Loosdrecht, M. C. M.;

Hallin, S. Life on N_2O: deciphering the ecophysiology of N_2O respiring bacterial

communities in a continuous culture. *ISME J.* **2018,** *12*, (4), 1142-1153.

(13) Yoon, S.; Nissen, S.; Park, D.; Sanford, R. A.; Löffler, F. E. Nitrous oxide reduction

kinetics distinguish bacteria harboring clade I versus clade II NosZ. *Appl. Environ.*

Microbiol. **2016,** *82*, (13),3793-3800.

(14) Suenaga, T.; Riya, S.; Hosomi, M.; Terada, A. Biokinetic characterization and activities

of N_2O-reducing bacteria in response to various oxygen levels. *Front. Microbiol.* **2018,**

9, 697.

(15) Jones, C. M.; Spor, A.; Brennan, F. P.; Breuil, M.-C.; Bru, D.; Lemanceau, P.; Griffiths,

B.; Hallin, S.; Philippot, L. Recently identified microbial guild mediates soil N_2O sink

capacity. *Nat. Clim. Change* **2014,** *4*, 801-805.

(16) Domeignoz-Horta, L. A.; Spor, A.; Bru, D.; Breuil, M.-C.; Bizouard, F.; Leonard, J.;

Philippot, L. The diversity of the N_2O reducers matters for the $N_2O:N_2$ denitrification

end-product ratio across an annual and a perennial cropping system. *Front. Microbiol.*

2015, *6*, 971

(17) Yoon, H.; Song, M. J.; Yoon, S. Design and feasibility analysis of a self-sustaining

biofiltration system for removal of low concentration N_2O emitted from wastewater

treatment plants. *Environ. Sci. Technol.* **2017,** *51*, (18), 10736-10745.

(18) Yoon, H.; Song, M. J.; Kim, D. D.; Sabba, F.; Yoon, S. A serial biofiltration system for

effective removal of low-concentration nitrous oxide in oxic gas streams: mathematical

modeling of reactor performance and experimental validation. *Environ. Sci. Technol.*

2019, *53*, (4), 2063-2074.

(19) Paranychianakis, N. V.; Tsiknia, M.; Kalogerakis, N. Pathways regulating the removal

of nitrogen in planted and unplanted subsurface flow constructed wetlands. *Water Res*.

2016, *102*, 321-329.

(20) Conthe, M.; Kuenen, J. G.; Kleerebezem, R.; van Loosdrecht, M. C. M. Exploring

microbial N_2O reduction: a continuous enrichment in nitrogen free medium. *Environ.*

Microbiol. Rep. **2018**, *10*, (1), 102-107.

(21) Boonnorat, J.; Techkarnjanaruk, S.; Honda, R.; Ghimire, A.; Angthong, S.; Rojviroon,

T.; Phanwilai, S. Enhanced micropollutant biodegradation and assessment of nitrous

oxide concentration reduction in wastewater treated by acclimatized sludge

bioaugmentation. *Sci. Tot. Environ.* **2018,** *637*, 771-779.

(22) Song, K.; Suenaga, T.; Harper, W. F.; Hori, T.; Riya, S.; Hosomi, M.; Terada, A.

Effects of aeration and internal recycle flow on nitrous oxide emissions from a

modified Ludzak–Ettinger process fed with glycerol. *Environ. Sci. Pollut. Res.* **2015,**

22, (24), 19562-19570.

(23) Vieira, A.; Galinha, C. F.; Oehmen, A.; Carvalho, G. The link between nitrous oxide

emissions, microbial community profile and function from three full-scale WWTPs. *Sci.*

Tot. Environ. **2019,** *651*, 2460-2472.

(24) Henry, S.; Bru, D.; Stres, B.; Hallet, S.; Philippot, L. Quantitative detection of the *nosZ*

gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S

rRNA, *narG*, *nirK*, and *nosZ* genes in soils. *Appl. Environ. Microbiol.* **2006,** *72*, (8),

5181-5189.

(25) Jones, C. M.; Graf, D. R.; Bru, D.; Philippot, L.; Hallin, S. The unaccounted yet

abundant nitrous oxide-reducing microbial community: a potential nitrous oxide sink.

ISME J. **2013,** *7*, (2), 417-426.

(26) Di, H. J.; Cameron, K. C.; Podolyan, A.; Robinson, A. Effect of soil moisture status

and a nitrification inhibitor, dicyandiamide, on ammonia oxidizer and denitrifier growth

and nitrous oxide emissions in a grassland soil. *Soil Biol. Biochem.* **2014,** *73*, 59-68.

(27) Stoliker, D. L.; Repert, D. A.; Smith, R. L.; Song, B.; LeBlanc, D. R.; McCobb, T. D.;

Conaway, C. H.; Hyun, S. P.; Koh, D.-C.; Moon, H. S. Hydrologic controls on nitrogen

cycling processes and functional gene abundance in sediments of a groundwater flow-

through lake. *Environ. Sci. Technol.* **2016,** *50*, (7), 3649-3657.

(28) Binder, B. J.; Liu, Y. C., Growth rate regulation of rRNA content of a marine

Synechococcus (Cyanobacterium) strain. *Appl. Environ. Microbiol.* **1998,** *64*, (9), 3346-

3351.

(29) Yoon, S.; Sanford, R. A.; Löffler, F. E., *Shewanella* spp. use acetate as an electron

donor for denitrification but not ferric iron or fumarate reduction. *Appl. Environ.*

Microbiol. **2013,** *79*, (8), 2818-2822.

(30) Park, D.; Kim, H.; Yoon, S. Nitrous oxide reduction by an obligate aerobic bacterium

Gemmatimonas aurantiaca strain T-27. *Appl. Environ. Microbiol.* **2017**, *83*, (12),

e00502-17.

(31) Ritalahti, K. M.; Amos, B. K.; Sung, Y.; Wu, Q.; Koenigsberg, S. S.; Löffler, F. E.

Quantitative PCR targeting 16S rRNA and reductive dehalogenase genes

simultaneously monitors multiple *Dehalococcoides* strains. *Appl. Environ. Microbiol.*

2006, *72*, (4), 2765-2774.

(32) Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res.* **2004,** *32*, (5), 1792-1797.

(33) Kumar, S.; Stecher, G.; Tamura, K. MEGA7: Molecular Evolutionary Genetics

Analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* **2016,** *33*, (7), 1870-1874.

(34) Hysom, D. A.; Naraghi-Arani, P.; Elsheikh, M.; Carrillo, A. C.; Williams, P. L.;

Gardner, S. N. Skip the alignment: degenerate, multiplex primer and probe design using

K-mer matching instead of alignments. *PloS One* **2012,** *7*, (4), e34560.

(35) Gardner, S. N.; Slezak, T. J. B. B. Simulate_PCR for amplicon prediction and

annotation from multiplex, degenerate primers and probes. *BMC Bioinformatics* **2014,**

15, (1), 237.

(36) Yoon, S.; Cruz-Garcia, C.; Sanford, R.; Ritalahti, K. M.; Loffler, F. E. Denitrification

versus respiratory ammonification: environmental controls of two competing

dissimilatory NO_3^-/NO_2^- reduction pathways in *Shewanella loihica* strain PV-4. *ISME*

J. **2015,** *9*, (5), 1093-1104.

(37) Li, W.; Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics* **2006,** *22*, (13), 1658-1659.

(38) Bolger, A. M.; Lohse, M.; Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **2014,** *30*, (15), 2114-2120.

(39) Nurk, S.; Meleshko, D.; Korobeynikov, A.; Pevzner, P. A. metaSPAdes: a new

versatile metagenomic assembler. *Genome Res.* **2017,** *27*, (5), 824-834.

(40) Li, H.; Handsaker, B.; Wysoker, A.; Fennell, T.; Ruan, J.; Homer, N.; Marth, G.;

Abecasis, G.; Durbin, R. The sequence alignment/map format and SAMtools.

Bioinformatics **2009,** *25*, (16), 2078-2079.

(41) Miller, C. S.; Baker, B. J.; Thomas, B. C.; Singer, S. W.; Banfield, J. F. EMIRGE:

reconstruction of full-length ribosomal genes from microbial community short read

sequencing data. *Genome Biol.* **2011,** *12*, (5), R44.

(42) Huang, Y.; Gilna, P.; Li, W. Identification of ribosomal RNA genes in metagenomic

fragments. *Bioinformatics* **2009,** *25*, (10), 1338-1340.

(43) Pjevac, P.; Schauberger, C.; Poghosyan, L.; Herbold, C. W.; van Kessel, M. A. H. J.;

Daebeler, A.; Steinberger, M.; Jetten, M. S. M.; Lücker, S.; Wagner, M.; Daims, H.

amoA-targeted polymerase chain reaction primers for the specific detection and

quantification of comammox *Nitrospira* in the environment. *Front. Microbiol.* **2017,** *8*,

1508.

(44) Meinhardt, K. A.; Bertagnolli, A.; Pannu, M. W.; Strand, S. E.; Brown, S. L.; Stahl, D.

A. Evaluation of revised polymerase chain reaction primers for more inclusive

quantification of ammonia-oxidizing archaea and bacteria. *Environ. Microbiol. Rep.*

2015, *7*, (2), 354-363.

(45) Ma, Y.; Zilles, J. L.; Kent, A. D. An evaluation of primers for detecting denitrifiers via

their functional genes. *Environ Microbiol* **2019,** *21*, (4), 1196-1210.

(46) Chen, X.; Peltier, E.; Sturm, B. S. M.; Young, C. B. Nitrogen removal and nitrifying

and denitrifying bacteria quantification in a stormwater bioretention system. *Water Res.*

2013, *47*, (4), 1691-1700.

(47) Conthe, M.; Wittorf, L.; Kuenen, J. G.; Kleerebezem, R.; Hallin, S.; van Loosdrecht, M.

C. M. Growth yield and selection of *nosZ* clade II types in a continuous enrichment

culture of N_2O respiring bacteria. *Environ. Microbiol. Rep.* **2018,** *10*, (3), 239-244.

(48) Orellana, L. H.; Rodriguez-R, L. M.; Higgins, S.; Chee-Sanford, J. C.; Sanford, R. A.;

Ritalahti, K. M.; Löffler, F. E.; Konstantinidis, K. T. Detecting nitrous oxide reductase

(*nosZ*) genes in soil metagenomes: method development and implications for the

nitrogen cycle. *mBio* **2014,** *5*, (3), e01193-14.

(49) Cassman, K. G.; Dobermann, A.; Walters, D. T. Agroecosystems, nitrogen-use

efficiency, and nitrogen management. *Ambio* **2002,** *31*, (2), 132-140.

(50) Aulakh, M. S.; Khera, T. S.; Doran, J. W. Mineralization and denitrification in upland,

nearly saturated and flooded subtropical soilI. Effect of nitrate and ammoniacal

nitrogen. *Biol. Fert. Soils* **2000,** *31*, (2), 162-167.

Figures

Figure 1. Flow chart for designing of the group-specific *nosZ* primer/probe sets and cross-

validation with *nosZ* sequence data extracted from shotgun metagenome data

Figure 2. The relative abundances of NosZG1-5 *nosZ* genes as quantified using qPCR and

metagenome analysis. The *nosZ* copy numbers from the qPCR assays were normalized with

the copy numbers of eubacterial 16S rRNA genes. The coverages of *nosZ* genes in the

metagenome data were normalized with the coverages of *rpoB* genes. The presented qPCR

quantification data are the averages of triplicate samples processed seperately through

extraction and qPCR procedures, with the error bars representing their standard deviations.

Figure 3. The phylogenetic tree constructed with the reference *nosZ* sequences of the taxa
matching OTUs generated from the partial *nosZ* sequences amplified with NosZG1-5 primers.
The phylogenetic tree was generated using neighbor-joining method with 500 bootstrap
replications. The heatmap shows the relative abundances of the OTUs within each *nosZ*
group, computed from the amplicon sequencing data and the *nosZ* sequence data extracted
from the metagenome

Figure 4. The copy numbers of NosZG1-5 *nosZ* genes and transcripts in activated sludge

samples collected from anoxic tanks of six activated sludge-type WWTPs in Korea, as

quantified using the group-specific *nosZ* qPCR developed in this study. The presented data

are the averages of triplicate samples processed seperately through extraction, purification

and reverse transcription (for analyses of transcripts), and qPCR procedures, with the error

bars representing their standard deviations.

666   Table 1. The primers/probe sets developed in this study for group-specific quantification of
667   *nosZ* genes
668
669

Targeted group	Primers / Probe	Sequence (5' → 3')	Degeneracy	Coverage	Efficiency (%)
NosZG1	NosZG1F	AAG GTN CGB GTN TAC ATG	48	70% (77/110)	91.6
	NosZG1R	CSN NCA TYT CCA TGT GCA	64		
	NosZG1P	FAM-ACT GCM VBT GGT TCT GCC AYG C-MGBNFQ	36		
NosZG2	NosZG2F	GRC ATC WKC MMC GAC AAG	32	84.4% (27/32)	93.4
	NosZG2R	HYC TCG RYG TTG TAC TGG	24		
	NosZG2P	FAM-ACC ACS CGC GTG TTC TGC G-MGBNFQ	2		
NosZG3	NosZG3F	CAY TTT GCW CCD GAY AAT ATT GAA	24	95% (19/22)	90.7
	NosZG3R	BSH WGT TTC ACC BGG CAT	108		
	NosZG3P	FAM-AAY YTR GAA CAA GAY TGG GAT GTA CCK C-MGBNFQ	32		
NosZG4	NosZG4F	ATW GTT GCY GGH GGM AAA	24	57.1% (4/7)	94.9
	NosZG4R	TTC CCA DGT KCC DAW TTT	36		
	NosZG4P	FAM-MGG YGA AGT KSA GAA TCC GGG WT-MGBNFQ	32		
NosZG5	NosZG5F	AAC GAC AAG KCS AAY CCG	8	100% (5/5)	97.7
	NosZG5R	GCG GTC GAA CTT CCA GTA	0		
	NosZG5P	FAM-GCS GTG MTC GAY CTG CGB G-MGBNFQ	24		