# Parallel ecological and evolutionary responses to selection in a natural bacterial community

Elze Hesse[1], Adela M. Luján[1, 2], Siobhan O'Brien[1, 3], Arthur Newbury[1], Terence McAvoy[1], Jesica Soria Pascual[1], Florian Bayer[1], Dave J. Hodgson[4] & Angus Buckling[1]

[1] CEC & ESI, Faculty of Environment, Science and Economy, University of Exeter, Penryn Campus, Cornwall, TR10 9FE, UK.

[2] Centro de Investigación y Desarrollo en Inmunología y Enfermedades Infecciosas (CIDIE), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)/Universidad Católica de Córdoba (UCC), Córdoba, X5016DHK, Argentina. Facultad de Ciencias de la Salud, Universidad Católica de Córdoba (UCC), Córdoba, X5004ASK, Argentina.

[3] Current affiliation: Department of Microbiology, School of Genetics and Microbiology, Trinity College Dublin, College Green, Dublin 2, Ireland.

[4] CEC, Faculty of Environment, Science and Economy, University of Exeter, Penryn Campus, Cornwall, TR10 9FE, UK.

**Corresponding author |** Elze Hesse, CEC & ESI, Faculty of Environment, Science and Economy, University of Exeter, Penryn Campus, Cornwall, TR10 9FE, UK. Email: e.hesse@exeter.ac.uk.

**Abstract |** Evolution can occur over ecological timescales, suggesting a potentially

important role for rapid evolution in shaping community trait distributions. However,

evidence of concordant eco-evolutionary dynamics often comes from *in vitro* studies of

highly simplified communities, and measures of ecological and evolutionary dynamics are

rarely directly comparable. Here, we quantified how ecological species sorting and rapid

evolution simultaneously shape community trait distributions by tracking within and

between-species changes in a key trait in a complex bacterial community. We focused on

the production of siderophores; bacteria use these costly secreted metabolites to scavenge

poorly soluble iron and to detoxify environments polluted with toxic non-ferrous metals.

We found that responses to copper-imposed selection within and between species were

ultimately the same – intermediate siderophore levels were favored – and occurred over

similar timescales. Despite being a social trait, this level of siderophore production was

selected regardless of whether species evolved in isolation or in a community context. Our

study suggests that evolutionary selection can play a pivotal role in shaping community

trait distributions within natural, highly complex, bacterial communities. Furthermore, trait

evolution may not always be qualitatively affected by interactions with other community

members.

**Significance |** Bacterial communities possess remarkable taxonomic and metabolic

diversity and play a key role in nearly every biogeochemical process on Earth. Rapid

evolution (occurring over ecological time scales) can in principle shape these processes,

yet we have little understanding of its importance in natural communities. Here, we

quantified how the production of metal-detoxifying siderophores is driven by species

compositional changes and evolution in a compost community in response to copper stress.

We found that siderophore production converged at intermediate levels, with evolutionary

and ecological changes occurring at similar rates. Understanding how ecological and

evolutionary processes contribute to shaping trait distributions will improve our ability to

predict ecosystem responses to global change, and aid in the engineering of microbial

consortia.

**Main text |** Changes in community trait distributions are typically driven by inter-

specific competition changing the abundance of species carrying certain traits (i.e.

ecological species sorting) (1). In the case of microbial communities, which have massive

population sizes and short generation times, evolution can occur over timescales

commensurate with ecological species sorting (2-4), suggesting a potentially important role

of rapid evolution in shaping community trait distributions (5-7) and responses to

environmental change (8). Understanding the mechanisms underpinning community trait

distributions is not only of fundamental interest, but also has applied relevance (9-11), such

as in the engineering of microbial consortia to perform valuable biological tasks (12),

including bioremediation (13), and the development of microbiome technologies aimed at

improving human health and crop yield (14).

Most work demonstrating a role of rapid evolution in community dynamics has

been carried out in *in vitro* environments (2, 15) and/or has relied on the use of greatly

simplified communities (3, 16, 17). This may overestimate the role of rapid evolution in

community dynamics because, in addition to the typically very strong selection pressures

associated with novel *in vitro* environments (4, 18), ecological complexity itself can

constrain evolution (19-22) (but see (23)). The response to selection – resulting from

ecological species sorting or within-species evolutionary change – typically increases with

trait variation (24). Increasing species diversity can both increase between-species trait

variation and reduce within-species variation – the latter because of reduced population

sizes (25, 26). The relative importance of ecological species sorting versus evolution on

shaping trait distributions is therefore likely to increase with species diversity (1, 19).

While some recent work has demonstrated rapid evolution occurring over ecological

timescales in more complex communities and/or semi-natural environmental contexts (2-

4), the role of ecological species sorting versus evolutionary selection in shaping natural

community trait distributions is unclear.

A useful model system to track simultaneous ecological and evolutionary changes

in functional traits is the ubiquitous production of metal-chelating siderophores in bacterial

communities (27). Although perhaps best known for their function as iron carriers (28, 29),

siderophores can also protect bacteria from toxic metal stress by preventing the diffusion

of toxic metals into bacterial cells (30, 31). Because detoxification takes place in a shared

environment, these costly extracellular compounds not only protect the producer, but

potentially also neighboring conspecifics (32) and other community members (33, 34). By

acting as community-wide public-goods, siderophores play a key role in driving species

interactions (35) and adaptive responses to environmental change (34). We previously

determined the effect of toxic metals on siderophore production in natural bacterial

communities inhabiting mine-degraded soils (33, 36) as well as in experimental compost

communities exposed to copper stress (33). In both cases, the presence of toxic metals

selectively favored bacterial taxa with greater investment in siderophores, thereby

increasing community-wide siderophore levels. However, we have also found that

bacterial taxa can benefit from siderophores produced by conspecifics (32) or individuals

from phylogenetically diverse taxa (34), suggesting that selection for (higher or lower)

siderophore production may also be affected by social interactions.

Here, we investigate ecological changes in siderophore production in response to

copper stress in a compost bacterial community simultaneously with evolutionary change

in siderophore production by a focal bacterium – the common soil-dwelling bacterium

*Pseudomonas fluorescens* (strain SBW25) (37). We hypothesized that ecological changes

in siderophore production resulting from exposure to novel copper stress would be greater

than evolutionary changes that occurred within SBW25. We also expected the community

to alter the evolutionary trajectory of SBW25's siderophore production as a consequence

of both community-imposed evolutionary constraints and greater selection for lower-level

siderophore production resulting from the protective effects afforded by other community

members.

**Results**

**Copper favors siderophore-producing taxa within the compost community**

We set up a fully factorial experiment involving the presence/absence of the natural

compost community, focal strain *Pseudomonas fluorescens* SBW25 and copper pollution

in initially sterile compost (SI Appendix, Fig. S1). These communities were allowed to

evolve for six weeks, after which we quantified changes in siderophore production by

isolating 24 clones of SBW25 and 24 isolates per replicate community where possible (i.e.

$n = 48$ isolates for the SBW25 + community treatments). Isolates were then grown

individually in King's B medium for 48 hrs after which we measured the ability of culture

supernatants to chelate iron using colorific CAS assays, corrected for variation in optical

density (our measure of siderophore production is based on arbitrary unity; see Methods).

We found that copper selected for greater siderophore levels within the compost

community compared to unpolluted control communities (Fig. 1A), irrespective of the

presence of SBW25 (2-way GLM on mean community siderophore production: $F_{1, 20} =$

$0.01$, $P = 0.91$ for copper × SBW25, $F_{1, 21} = 0.04$, $P = 0.85$ for SBW25 and $F_{1, 22} = 8.88$, $P$

$< 0.01$ for copper treatment. GLM estimates of mean siderophore production $\pm$ 95%

confidence intervals [CI] in copper-polluted versus unpolluted communities pooled across

SBW25 levels (i.e. presence/absence): 0.55 [0.53, 0.57] and 0.50 [0.48, 0.52],

respectively).

To determine whether the observed shifts in siderophore production were linked to

any compositional changes in the community, we determined the genus-level identity of all

final-time-point compost isolates assayed for siderophore production ($n$ = 548 isolates in

total). We found that copper-mediated changes in siderophore production were

accompanied by a shift in community composition (2-way PERMANOVA on Bray–Curtis

dissimilarities: copper × social context: $F_{1, 20}$ = 2.47, $P$ = 0.03), with between-community

diversity differing significantly across all treatment combinations (pairwise

PERMANOVAs: $R^2$ > 0.26, all $P_{adj}$ < 0.05; Fig. 1B). In particular, the presence of SBW25

changed the relative abundance of taxa within the compost community by suppressing

*Pseudomonas, Bacillus* and *Arthrobacter* compared to the community-only treatment (all

$P_{adj}$ <0.05; SI Appendix, Fig. S2). We also found that copper tended to select against taxa

that displayed relatively low mean levels siderophore production (i.e. blue taxa in Fig. 1C):

in particular, while intermediate siderophore producers, such as *Cupriavidus*, were

significantly more abundant in copper-polluted communities, *Achromobacter* and

*Lysinibacillus* were negatively affected by copper stress (SI Appendix, Table S1). As a

result of such ecological species sorting, the most abundant community members in

copper-polluted compost were those displaying more intermediate siderophore levels (Fig

1D). Note that evolution may also have contributed to changes in siderophore production,

but this is impossible to quantify from these taxonomic results. We therefore focus on

evolution within a focal strain.

**Copper selects against high siderophore levels within SBW25**

We tracked siderophore changes within replicate populations of SBW25 that had either

evolved in the presence or absence of the compost community. We found the exact

opposite response compared to that observed at the community level – toxic copper

selectively favored SBW25 individuals that, on average, produced lower amounts of

siderophore compared to control populations (LMM: copper effect: $\chi_1^2 = 10.97$, $P < 0.001$.

Mean siderophore production $\pm$ 95% CI in copper-polluted versus control populations:

0.68 [0.67, 0.70] and 0.72 [0.71, 0.73], respectively). Again, the impact of copper stress on

siderophore production did not differ significantly as a function of community

presence/absence (LMM: interaction between copper $\times$ community: $\chi_1^2 = 0.22$, $P = 0.64$;

community effect: $\chi_1^2 = 1.97$, $P = 0.16$; Fig. 2).

**Copper selects against siderophore extremes**

How can we resolve these apparently opposing responses to copper-imposed selection

operating at the species versus community level? To answer this question, we first

compared the shape of the distribution of siderophore production of the compost

community to that of SBW25. We found that copper selected against siderophore

extremes, resulting in trait convergence (Fig. 3), regardless of whether SBW25 and the

community were grown in isolation or together in co-culture.

To quantify this convergence, we focused on the effect that copper has on

siderophore production when the community and SBW25 were grown together, using a

hierarchical Bayesian linear model to account for heteroscedasticity between isolate types

(community versus SBW25; all point estimates are *maximum a posteriori estimates* $\pm$ 95%

credible intervals). The copper-mediated change in siderophore production in SBW25 was

-0.042 [-0.072, -0.014]. In contrast, copper stress increased siderophore production among

compost community isolates by 0.044 [0.009, 0.081]. Thus, the effect was of

approximately equal magnitude, but in the opposite direction. Indeed, combining the

posterior estimates for the effect of copper on siderophore production in SBW25 and in the

rest of the community (positive values indicate that the positive effect on community

isolates is larger than the negative effect in SBW25 and *vice versa*), the estimated

combined effect was 0.001 [-0.044, 0.045]. In the absence of copper, the difference in

siderophore production between SBW25 and the compost community was 0.216 [0.185,

0.251]. Copper pollution significantly reduced this figure by 0.086 [0.041, 0.133] to 0.130

[0.098, 0.161]. Thus, opposing and approximately equal effects of selection lead to a

significant reduction in the difference between (relatively high producing) SBW25 and

(relatively low producing) community isolates under copper stress.

The model described above included separate standard deviation parameters for

each sample type (community/SBW25 × copper/no copper) to estimate their respective

within-microcosm variation in siderophore production. The standard deviation of

siderophore production amongst SBW25 isolates was 0.0294 [0.0267, 0.0340] in the

absence of copper. The copper treatment led to an increase of 0.006 [-0.00031, 0.0131] to

0.0354 [0.0315, 0.0427]. Conversely, variation in siderophore production was reduced

amongst compost community isolates in the presence of copper. Here, the standard

deviation fell from 0.0994 [0.0899, 0.115] to 0.0693 [0.0633, 0.0808] in response to the

copper treatment, a significant difference of 0.0301 [0.0157, 0.0474]. That the overall

variation in siderophore production was much higher in the compost community than

SBW25 is expected, since measurements were taken from a range of unrelated strains. The

decrease in variance due to copper stress suggests ecological species sorting (and some

evolutionary change), whereas the increase in variance within SBW25 isolates is

compatible with evolutionary change.

To further investigate the generality of this result, we focused on isolates of

representative genera (*n* = 8 genera) that occurred in both copper environments and had

been assayed for siderophore production. We found that trait changes were qualitatively

similar: copper selected against siderophore extremes. That is, taxa with high 'baseline'

siderophore levels (i.e. mean siderophore levels > 0.55 in control conditions) produced

fewer siderophores in copper-polluted compost (Cohen's $d < 0$; SI Appendix, Fig. S3). In

contrast, copper stress favored greater siderophore production in bacterial taxa with low

'baseline' siderophore levels (Cohen's $d > 0$). Hence, we found a negative relationship

between effect size and 'baseline' siderophore levels (Spearman rank correlation: $S =$

624.36, $P = 0.005$, $\rho = -0.72$).

**Direct fitness costs of high levels of siderophore production in copper**

Selection for siderophore production in the presence of toxic copper can be explained by

the direct protective effects afforded by siderophores. However, what is less clear is why

there was selection against *high* levels of siderophore production in toxic copper

conditions. There are two general, non-mutually exclusive explanations. First, there may

be a direct cost associated with greater investment in siderophores in copper when

individuals produce more than enough siderophore to detoxify copper, making the excess

an unnecessary metabolic burden. Second, individuals producing high levels of

siderophore may protect competing conspecific and heterospecific individuals, increasing

the relative cost of siderophore production.

We first explored how evolved changes in siderophore production affect individual

performance in toxic copper conditions. Specifically, we determined whether SBW25

clones that had evolved to produce relatively low levels of siderophore displayed lower

copper tolerance ($n = 277$ clones in total, all isolated from the 'no community' treatment;

SI Appendix, Fig S1). To this end, evolved clones were individually grown in King's B

broth supplemented with a toxic dose of copper sulphate (final concentration of 6.17 mM

$CuSO_4$). We recorded changes in optical density ($OD_{600}$) over a 48-hrs period and

estimated clone-specific Malthusian growth parameters ($m$). Our results show that clones

that had evolved in copper-free and copper-polluted compost grew equally well in toxic

copper broth (LMM: effect of evolutionary background on $m$: $\chi^2_1 = 0.76$, $P = 0.39$; SI

Appendix, Fig. S4), suggesting that siderophores are effective in detoxifying copper.

Furthermore, reduced investment in siderophores did not confer lower tolerance,

irrespective of a clones' evolutionary history (LMM on $m$: siderophore production $\times$

copper background: $\chi^2_1 = 0.02$, $P = 0.88$; copper background: $\chi^2_1 = 0.16$, $P = 0.69$;

siderophore production: $\chi^2_1 = 2.95$, $P = 0.09$; SI Appendix, Fig. S4). The observed shift

towards reduced siderophore levels in SWB25 therefore did not alter copper tolerance.

239            We next investigated why high investment in siderophores is costly in the presence

of copper but not in the absence. We speculated that the constrained growth of SBW25 in

the presence of copper (SI Appendix, Fig. S5) may have reduced levels of siderophore

required for the canonical function of siderophores: iron uptake (38). To test this

hypothesis, we manipulated population growth rates (and hence indirectly the need for iron

to sustain growth) by inoculating monocultures of the wild-type pyoverdine-producing

SBW25 strain and an isogenic pyoverdine-deficient mutant (PBR840) (39) in sterile

compost at very low density (allowing for faster growth) and high density (allowing for

slower growth), both in the presence and absence of copper pollution. This mutant does not

produce the main siderophore pyoverdine (which reduces total siderophore production to

21% of the wild- type) and displays levels of siderophore production similar to an average

compost community member.

251            Confirming our observations on copper tolerance in evolved SBW25 populations,

the pyoverdine-deficient mutant performed equally well compared to the pyoverdine

producing strain in copper-polluted compost (GLM on $m$: copper $\times$ strain: $F_{1,43} = 0.52$, $P =$

0.48; Fig. 4A), with copper significantly reducing population growth (main effect copper:

$F_{1,44} = 100.24$, $P < 0.001$. Mean growth $\pm$ 95% CI: $\bar{m}_{control} = 1.50$ [1.45, 1.55] and

$\bar{m}_{copper}$ 1.14 [1.09, 1.19], pairwise contrast: $t$ ratio $= 10.01$, $P < 0.001$), irrespective of

inoculation density (copper × density: $F_{1,40} = 0.22$, $P = 0.65$). We also found that in both

copper environments, the pyoverdine producers benefitted when inoculated at low density

and hence growth was not constrained (strain × density: $F_{1,43} = 16.30$, $P < 0.001$;

$\bar{m}_{producer} = 1.98$ [1.90, 2.05] and $\bar{m}_{deficient}$ 1.88 [1.81, 1.95], pairwise contrast: $t$ ratio =

1.89, $P = 0.07$; Fig. 4A). In contrast, the non-producing mutant grew faster than the

producer under slow (high inoculation density) growth conditions ($\bar{m}_{producer} = 0.61$ [0.54,

0.69] and $\bar{m}_{deficient} = 0.81$ [0.74, 0.88], pairwise contrast: $t$ ratio = -3.82, $P < 0.001$).

Consistent with this monoculture pattern, the relative fitness of the pyoverdine producer

was greater when competing with the mutant under high growth conditions (low

inoculation density), while it was lowest under low growth conditions (GLM on $r_{producer}$:

density × copper interaction: $F_{1,19} = 6.95$, $P = 0.016$) – especially in the presence of copper

(high density: $\bar{r}_{copper} = 0.79$ [0.73, 0.85 ]and $\bar{r}_{control} = 0.91$ [0.85, 0.97], pairwise contrast:

$t$ ratio = 2.92, $P < 0.01$ and low density: $\bar{r}_{copper} = 1.04$ [0.98, 1.10] and $\bar{r}_{control} = 1.00$

[0.94, 1.07], pairwise contrast: $t$ ratio = -0.86, $P = 0.40$. Data on Malthusian growth and

relative fitness are presented in SI Appendix, Fig. S6). These results suggest a cost to high

investment in siderophores when growth is otherwise constrained, while there is no cost

and potentially a benefit to high levels of siderophore production when growth is less

constrained, and these compounds are needed to compete for iron.

**Public-goods exploitation favors reduced investment in siderophores**

While direct costs and benefits could qualitatively explain our results, we next

explored whether public-goods exploitation could have played an additional role in driving

selection against SBW25 individuals producing relatively large amounts of siderophores.

Previous work investigating within-species exploitation of siderophores has demonstrated

that siderophore production is costly to the individual (40, 41), creating incentives for non-

producing 'cheats' to invade and steal these public-goods away from the producer (42).

We have also found that bacterial compost isolates that invest relatively little in

siderophores benefit from other species' detoxifying siderophores under similar levels of

copper stress (34). However, siderophore-mediated facilitation could not fully rescue the

growth of copper-sensitive isolates, nor did it result in exploitation (i.e. no detectable

indirect costs by helping competitors) within the community over ecological timescales

(34). Combined with our current finding of parallel changes in siderophore distribution at

the within-taxon and community level (Fig. 3), we focus on social dynamics *within* species

only.

To determine whether individuals with lower siderophore investment can exploit

relatively high producers we carried out competition assays. In particular, we manipulated

the relative frequencies of the wildtype and pyoverdine-deficient mutant of SBW25 (1% or

99% starting frequency of strains) and assayed their relative performance following 7 days

of growth in copper-polluted and unpolluted compost. Negative frequency-dependent

fitness often characterizes social interactions in structured environments, where rare non-

producing 'cheats' are more likely to readily encounter cooperators rather than other cheats

and grow relatively well, while rare cooperators will mostly encounter cheats, but so will

other cheats; with the latter suffering a greater relative growth cost. We found that the

pyoverdine-deficient mutant experienced a relative selective benefit when rare, in

particular in toxic copper conditions (GLM on selection coefficient ($s$) of pyoverdine-

deficient mutant: frequency $\times$ copper: $F_{1,\,44} = 79.30$, $P <0.001$. Mean $\pm$ [95% CI] for

control: $s_{high}= 0.045$ [-0.08, 0.13] and $s_{low} = 0.33$ [0.25, 0.41]; pairwise contrast; $t$ ratio =

-4.95, $P < 0.001$. Mean $\pm$ [95% CI] for copper: $s_{high}= 0.301$ [0.22, 0.38] and $s_{low} = 1.31$

[1.22, 1.39]; pairwise contrast; $t$ ratio = -17.55, P < 0.001; Fig. 4B. Data on Malthusian

growth are presented in SI Appendix, Fig. S7). This shows that in addition to direct fitness

effects, social exploitation by conspecifics could also have contributed to the evolution of

reduced siderophore production by SBW25 in response to copper stress.

**Discussion**

Using siderophores as a focal trait, we show that evolutionary and ecological processes

operate in parallel and over similar timescales in a natural bacterial community, resulting

in trait convergence towards intermediate siderophore levels in response to copper stress.

In particular, copper-imposed selection changed the relative abundance of bacterial taxa by

selecting against individuals that displayed extremes in siderophore investment. This

pattern was consistent at the community- and within-taxon level. Crucially, changes in

siderophore production not only resulted from ecological species sorting, but also from

evolutionary changes within species: SBW25 adapted to copper stress by lowering its

investment in siderophores to levels more comparable to that of an average compost

community member.

In our system, copper pollution resulted in selection against siderophore extremes.

We propose such stabilizing selection in part resulted from directly incurred costs borne by

taxa with relatively low and high siderophore production. In particular, we have previously

shown that bacterial compost taxa that invest relatively little in siderophores display

reduced copper tolerance and suffer an absolute growth cost (34). Our current findings

indicate that beyond a certain threshold, increased investment in detoxifying siderophores

resulted in diminishing returns, thereby selecting against individuals producing relatively

large amounts of siderophore.

This raises the question as to why siderophore production was maintained at higher

levels in unpolluted compared to copper-polluted environments. It has been well-

established that siderophores play an important role in scavenging insoluble iron from the

environment (43). Ironbound siderophores are typically imported by specific receptors that

constrain their use by other competing strains, thereby limiting siderophore cross-feeding

between strains (but see (44, 45)). As such, siderophores are thought to play a key role in

determining the outcome of interspecific competition for iron (46, 47). Hence,

siderophores were likely positively selected for in copper-free conditions to pillage iron

from competing strains. By contrast, competition for iron was likely reduced under copper

stress because iron no longer limited bacterial growth rates (copper did); the main role of

siderophores in copper-polluted conditions was to detoxify metals.

Social conflict and cooperation are often thought to play a key role in driving the

eco-evolutionary dynamics of microbial communities (48-53). In microbes, social

behaviors are often mediated by secretion of 'public-goods' – metabolically costly

compounds that benefit the producer, but that are open to exploitation by non-producers

that do not contribute their fair share but utilize the benefits afforded by their production

(54). Most of our understanding of public-good dynamics comes from work on within-

species interactions (55), yet many public-goods can simultaneously benefit other species

(56-60), particularly in microbial communities. Key examples include antibiotic-degrading

enzymes (61, 62), resource-scavenging molecules (55, 63), immune-manipulating effectors

(64-67) and potentially siderophore-mediated detoxification. However, we found little

evidence to suggest that inter-specific facilitation and exploitation shaped siderophore

production in this experiment: siderophore production of neither the community nor

SBW25 was qualitatively affected by the other, despite – and in line with previous work –

SBW25 greatly altering community composition over the course of our experiment (68).

We did find some evidence for within-species exploitation: the pyoverdine-deficient

'cheat' accrued a relative fitness benefit when competing with wild-type pyoverdine-

producing SBW25, in particular when rare, both in the presence and absence of copper.

Such frequency dependent dynamics are indicative of intraspecific exploitation of public-

goods in spatially structured environments (55, 69-71), and is predicted to increase

variation within species while decreasing divergence between species, which is borne out

by our results.

In our experiment we quantified the *potential* of bacterial isolates to produce

siderophores rather than their *in-situ* production in compost. The transcription of genes

encoding the proteins that participate in siderophore biosynthesis is tightly regulated by the

presence of iron and/or toxic metals (30). However, the use of a common garden

environment to quantify siderophores was necessary because: (1) for the majority of

environmental isolates we do not exactly know what kind of metal-chelating agents they

produce, meaning a metagenomics approach is currently not viable; (2) it is notoriously

difficult to extract siderophores from soil and link their production to specific taxa and (3)

the presence of soil organic material and other factors in compost interferes with CAS

assays. Notwithstanding, we demonstrate a direct link between relative changes in

siderophore potential and fitness.

Our focus on public-goods traits, such as siderophore production, may have over-

estimated the role of evolution. Previous work has shown that public-goods cheats readily

emerge via loss-of-function mutations, including cheats defective for quorum sensing (72)

or the production of extracellular proteases (73), polysaccharides (74) and siderophores

(70). For example, pyoverdine-deficient cheats typically evolve within days in the lab (75),

often via single SNPs in regulatory genes. However, over-estimation of the role of

evolution may be mitigated to some extent because, compared to many studies on the

evolution of public-goods, we evolved compost communities over an extended period

which also allowed for major shifts in community composition. More broadly, adaptive

evolution of a wide range of non-social traits can result from loss-of-function mutations

through rewiring of regulatory and metabolic networks (76). Finally, gain of functions,

including potential public-goods traits, can also potentially evolve very rapidly via

horizontal gene transfer (HGT; 77, 78, 79) (80). HGT can occur across large phylogenetic

distances (81), which could actually increase the relative importance of evolution *versus*

species sorting in shaping community trait distributions.

To conclude, we show that the importance of rapid evolution in shaping community

trait distributions is not limited to simplified *in vitro* communities but can also play a role

as important as species sorting within highly complex natural microbial communities.

Furthermore, our results suggest that trait evolution may not always be qualitatively

affected by interactions with other community members.

## Methods

394    All supporting data and R code are available at https://zenodo.org/records/11639136.

395    Sequences have been deposited at ENA Project PRJEB29924:

396    https://www.ebi.ac.uk/ena/data/search?query=PRJEB29924.

### Bacterial strains

We used *Pseudomonas fluorescens* strain SBW25 as our focal species (37). This common

soil-dwelling bacterium produces a variety of siderophores, including pyoverdine (45) and

an ornicorrugatin-like siderophore (82), that are able to chelate iron. The strain's primary

siderophore – pyoverdine – has been shown to bind copper (30), which renders the

environment less toxic by preventing diffusion into the bacterial cell (31). To facilitate

isolation of SBW25 from the community, the strain was modified by inserting a LacZ

genetic marker and a gentamicin-resistance cassette, which resulted in blue resistant

colonies when plated out on Lysogeny Broth (LB) agar supplemented with 90μg/mL 5-

Bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-gal) and 25 μg/mL gentamicin (32).

To isolate the compost community, 40 gr of fresh compost (Verve John Innes No.

1) was added to 200 mL of M9 minimal salt solution and incubated shaken at 150 rpm at

28 °C for 24h. The supernatant was plated out on LB agar to verify the presence of a

bacterial community, and also on LB supplemented with 30μg/mL gentamicin to verify the

susceptibility of the community to this antibiotic (32). A 20% glycerol stock solution was

prepared from the aqueous sample suspension and frozen at -80°C for future use.

**Selection experiment**

Compost microcosms were established with 30 gr of twice-autoclaved sterile compost in

round 90 mm petri dishes. To track ecological and evolutionary changes in siderophore

production, we seeded twelve replicate compost microcosms with (1) a soil wash of the

isolated compost community, (2) an overnight culture of SBW25 or (3) both together at a

1:1 ratio ($n = 36$ compost microcosms in total, SI Appendix, Fig. S1), keeping total

inoculation density constant ($\sim 5 \times 10^7$ cells). Note that results of the community-only

treatment were published previously (33). Compost microcosms were incubated at 26°C

and 75% relative humidity for 24h, after which half of the microcosms of the above

treatments ($n = 6$) was supplemented with a toxic dose of copper (2 mL of filter-sterilized

0.25M $CuSO_4$) and the remainder with an equal volume of sterile $dd$H$_2$0 (33). Microcosms

were incubated for a total of 6 weeks. After three weeks, a second dose of copper or

$dd$H$_2$O was added to the relevant microcosms. Samples were taken just before ('ancestral')

and 6 weeks after copper amendment ('evolved') by transferring 1 gr of compost per

microcosm to 6 mL of M9 solution in 30 mL glass vials. Vials were shaken for 2h at 26°C

at 180 rpm, after which supernatants were frozen in glycerol (25% final concentration) at -

80°C for future assays.

**Phenotypic assays on ancestral and evolved isolates**

*Siderophore production*

To quantify siderophore production, serial-diluted freezer stocks (taken just before and 6

weeks after giving the first dose of copper) were plated out on duplicate LB agar plates

supplemented with (i) X-gal to isolate members of the compost community or (ii) X-gal

and gentamicin to isolate SBW25 colonies. Following 48 hours of incubation at 26°C,

individual colonies could be identified, counted and picked. For each unique treatment-

time combination, we picked 24 colonies of the compost community and/or SBW25 per

replicate, with a total of 48 colonies picked from compost microcosms seeded with both

SBW25 and the community. Individual colonies were grown in KB broth for 48h at 26°C,

after which the supernatant was assayed for the extent of iron chelation. Siderophore

production was quantified using the liquid CAS assay described by Schwyn and Neilands

(83). Siderophore production per isolate was estimated using: $[1 - (A_i/A_{ref})] /(OD_i)$, where

$OD_i$ = optical density at 600 nanometer (nm) and $A_i$ = absorbance at 630 nm of the assay

mixture and $A_{ref}$ = absorbance at 630 nm of reference mixture (KB+CAS; $A_{ref}$). For some

samples, absorbance reads were negative following reference correction, so we

standardized siderophore production by setting the minimum observed value to zero. Note

that CAS assays performed in iron-limited KB medium provided qualitatively similar

results (34).

*Copper tolerance*

We determined the direct fitness consequences of copper-imposed reductions in

siderophore production in SBW25. Final-time-point clones ($n = 277$) isolated from

populations that had evolved in the absence of the community were grown individually at

26°C for 24h, after which $\sim 10^4$ cells were inoculated into 96-well plate wells containing

200 µL of King's B broth supplemented with a toxic dose of copper sulphate (final

concentration of 6.17 mM $CuSO_4$). Individual clones were incubated statically at 26°C for

52 hours, and their optical densities ($OD_{600}$) were measured every 8–12 h (Varioskan Flash

plate reader, Thermo Scientific, Waltham, MA, USA). We determined the Malthusian

growth parameter (*m*) for each clone as: ln(final density / start density) / time (hours) (84).

**Sanger sequencing of compost isolates**

The 16S rRNA gene of all final-time-point compost isolates assayed for siderophore

production was sequenced to confirm genus-level identity (i.e. *n* = 24 per replicate). PCRs

were performed in 25µL reactions containing 1x DreamTaq Green PCR Master Mix (2X)

(Thermo Scientific), 200 nM of the 27F and 1492R primers and 3 µL of 1:100 diluted

culture that had undergone 3 freeze-thaw cycles. The thermal cycling parameters were set

to 94ºC for 4 min, followed by 35 cycles of 1 min at 94ºC, 30s at 48ºC and 2 min at 72ºC,

and a final extension of 8 min at 72ºC. Following Exo-AP clean-up, high quality samples

were Sanger sequenced using the 27F primer (Core Genomic Facility, University of

Sheffield). Sequence quality was assessed using the R 'dada2' package (85), and

sequences were trimmed in Genious (86) to achieve an overall quality score >35. Using

Mothur (87), sequences longer then 300bp were then aligned to the Silva.Bacteria.Fasta

database, and taxonomy was classified using the RDP trainset 14 032015 as a reference

database.

**The role of siderophores as iron-scavengers as a function of density**

To determine whether high siderophore production was selected against as a result of iron

being less limited under toxic copper conditions, we conducted competition assays

between the pyoverdine-producing LacZ-marked SBW25 strain (37) and an isogenic

pyoverdine-deficient mutant (PBR840) (39) in 30 mL glass universals containing 5 grams

of twice-autoclaved sterile compost. Using a full-factorial design, strains were grown in

isolation or together at a 1:1 ratio at either low ($10^2$ CFUs) or high inoculation density ($10^6$

CFUs), keeping total inoculation density constant across these social contexts. Compost

microcosms were incubated for 24 hours at 26°C and 75% relative humidity, after which

we added a toxic dose of $CuSO_4$ (1 mL of 0.25M $CuSO_4$) to half of the microcosms and an

equal volume of sterile $dd$H$_2$O to the remainder *(n = 6 replicates per treatment)*. Compost

microcosms were incubated for a total of 7 days, after which soil washes were taken, serial

dilutions of which were plated out on KB agar. For co-cultures, strains could be

distinguished based on their colony pigmentation: wild-type colonies were blue on X-gal

supplemented KB agar, whereas mutant colonies were white. We determined the

Malthusian growth parameter (*m*) for each strain as described above.

**Public-goods exploitation in SBW25**

To determine whether social exploitation could have played an additional role in driving

copper-mediated reductions in siderophore production in SBW25, we conducted

competition assays between the pyoverdine-producing wild-type and pyoverdine-deficient

mutant strain of SBW25 in petri dishes containing 30 gr of twice-autoclaved sterile

compost. Using a full-factorial design, strains were inoculated together at a low (1%, $10^4$

CFUs, $n = 24$) or high (99% $10^6$ CFUs, $n = 24$) frequency. Compost microcosms were

incubated for 24 hours at 26°C and 75% relative humidity, after which we added a toxic

dose of $CuSO_4$ (2 mL of 0.25M $CuSO_4$) to half of the microcosms and an equal volume of

sterile $dd$H$_2$O to the remainder. Following 7 days of incubation, soil washes were taken

and plated out on KB agar supplemented with X-gal to distinguish between strains. We

determined the Malthusian growth parameter (*m*) for each strain as above.

**Data analyses**

For all analyses, we used *R* Version 4.0.3 (R Development Core Team; http://www.r-

project.org). In general, models were compared by sequentially deleting terms and

comparing model fits using F-tests or $\chi^2$-tests (where appropriate), after which pairwise

contrasts were computed using the 'emmeans' packages (88), with $\alpha < 0.05$. In case of

multiple pairwise testing, *P*-values were adjusted using the false discover rate ('fdr')

method unless stated otherwise. We checked residual behaviour using the 'DHARMa'

package (89). All plots were produced using the 'ggplot2' package (90).

*The effect of copper on community structure and mean levels of siderophore production*

To test for the effect of copper on siderophore production in compost communities that had

either evolved with or without SBW25, we initially used a linear mixed effects (LMM)

model (*'lmer'* function in *'lme4'* package) (91) with copper × SBW25 presence as fixed

explanatory variables, as well as their interaction. To account for non-independency of

observations we fitted random intercepts for individual microcosms. However, we

observed strong heteroscedasticity of residuals, which could not be remedied by including

a treatment-specific variance structure. For each replicate community, we therefore

averaged siderophore production across the 24 compost isolates assayed and used a 2-way

generalized linear model (GLM) with a Gaussian error distribution to test for the

interactive effect of copper × SBW25 presence on mean community-wide siderophore

production. This simplified model gave qualitatively very similar results to the LMM

without violating any underlying model assumptions.

We determined the genus-levels identity of all end-point compost isolates assayed

for siderophore production (*n* = 24 per community) and used PCoA ordination plots to

depict pair-wise Bray-Curtis dissimilarities in community composition between

microcosms (excluding observations on SBW25). To test for the interactive effect of

copper × SBW25 presence on community structure we ran a permutational 2-way

ANOVA test on Bray-Curtis dissimilarities using the '*adonis2*' function in the '*vegan*'

package (92) with 9999 permutations. Following this, we ran pairwise permutational

ANOVAs to disentangle which treatments differed significantly from one another using

the '*calc_pairwise_permanovas*' function in the '*mctoolsr*' package

(https://github.com/leffj/mctoolsr/), with *P*-values corrected for multiple testing using the

'fdr' method. We confirmed homogeneity of variance using the '*betadisper*' function in

the '*vegan*' package to estimate group differences in dispersion and treated all copper ×

SBW25 combinations as levels of a single factor in a 1-way AVOVA on dispersion.

To determine which of the compost isolates differed in abundance across

treatments, we fitted a negative binomial GLM to the data using the '*DESeq*' function in

the R package '*DESeq2*' (93). We focused on a subset of ten common compost isolates

that occurred in both unpolluted and copper-polluted communities. Based on the outcome

of pairwise permutational ANOVAs (see above), we initially included the interaction term

between copper × SBW25 presence in the design of our GLM. However, following model

reduction (LRT tests), the minimal adequate model was best described by only including

the additive effect of copper and community context. Using this additive model, we

calculated significant differences in the abundance of taxa using Wald tests and we

corrected *P* –values for multiple testing for each of two contrasts used (presence/absence

of either SBW25 or copper).

*The effect of copper on mean siderophore production in a focal strain*

To test for the interactive effect of copper and community context on siderophore

production in SBW25, we used a LMM model with copper × community presence as fixed

explanatory variables, as well as their interaction. To account for non-independency of

observations, we fitted random intercepts for individual microcosms. Based on the

obtained simulation-based residual plots, we included a dispersion parameter for each of

the copper × community levels, using the *'glmmTMB'* function in the *'glmmTMB'* package

(94).

*The effect of copper on the distribution of siderophore production*

To demonstrate that copper selectively favored intermediate siderophore levels, we

determined the effect size of copper for bacterial taxa that commonly occurred in both

copper treatments and related this to their mean siderophore production under non-polluted

conditions, using a Spearman rank correlation. Effect size was calculated using the

*'cohen.d'* function in the *'effsize'* package (95). Cohen's *d* takes the difference between

two means and expresses it in standard deviation units, with negative values indicating a

reduction in siderophore levels in response to copper, and *vice versa* for positive values.

*Effect-size in compost communities evolved with SBW25*

To determine the impact of copper on the siderophore production of both SBW25 and

community isolates that had evolved together, we used a hierarchical Bayesian linear

model which accounts for the differing variances associated with explanatory variables

(i.e. siderophore measurements from community isolates were taken from diverse bacterial

strains even within a single replicate community, as opposed to the originally isogenic

SBW25 stocks):

$$y_i \sim N(\mu_i, \sigma_{\text{group}[i]})$$

$$\mu_i = \lambda_{\text{group}[i],\text{replicate}[i]}$$

$$\lambda_{j,k} \sim N(\nu_j, \tau_j) \text{ , for } j = 1..4, k = 1..6$$

$$\nu_j = \alpha_{X1[j]} + \beta_{X1[j]} X2_j \text{ , for } j = 1..4$$

$$\alpha_j \sim N^+(0,1) \text{ , for } j = 1..2$$

$$\beta_j \sim N(0,0.25) \text{ , for } j = 1..2$$

$$\sigma_j \sim \text{Exponential}(0.2) \text{ , for } j = 1..4$$

$$\tau_j \sim \text{Exponential}(0.2) \text{, for } j = 1..4$$

where X1 is a vector of indices indicating SBW25 vs community samples; X2 is a vector

indicator indicating the presence/absence of copper; *replicate* and *group* are vectors of

indices denoting the replicate microcosm (from a given group) and the combination of

isolate type (SBW25 vs community) and presence/absence of copper for each data point.

The model above, with separate standard deviation terms ($\sigma$) for each group was chosen

over models with one or two (determined by isolate type or copper presence) standard

deviation terms after model selection by Pareto smoothed importance sampling leave one

out (PSIS-LOO) cross validation (96). Inference was performed by Markov chain Monte

Carlo using Turing.jl (97) in the Julia programming language (98): 4 chains of 1000

samples each using the No U-Turn Sampler (NUTS) algorithm (99), with all R-hat scores

< 1.01. 95% credible intervals were calculated as the 95% highest posterior density interval

(HPDI) using StatisticalRethinking.jl

(https://github.com/StatisticalRethinkingJulia/StatisticalRethinking.jl). Subsequently, point

estimates were obtained by computing the *maximum a posteriori* parameter estimates of

the same model, using Optim.jl (100) in conjunction with Turing.jl. To compare the

positive affect of copper on siderophore production amongst compost community isolates

with the negative affect on SBW25, we computed the posterior distribution of the

combined effect as $\beta_1 + \beta_2$ by element-wise addition of Markov chain samples. The same

approach was used to infer the posterior distribution of the difference between community

and SBW25 siderophore production with copper ($\alpha_2 - \alpha_1$) and without ($\alpha_2 + \beta_2 - (\alpha_1 + \beta_1)$).

*The role of direct and indirect selection in favoring intermediate siderophore levels*

To test for whether copper-adapted clones of SBW25 evolved greater resistance to copper

than those evolved under unpolluted conditions, we used a LMM model on *m* with

evolutionary background as fixed explanatory variable. To account for non-independency

of observations, we fitted random intercepts for individual microcosms. However, the

variation explained by the random intercepts approached zero, causing singularity issues

when fitting the LMM using the '*lme4*' package. We therefore used the default setting of

the nonlinear optimizer in the '*glmmTMB*' function for parameter estimation. We next

tested whether increased investment in siderophores conferred greater copper tolerance by

including siderophore production in the evolutionary LMM described above.

We determined whether siderophores were selectively favored when iron was

needed for growth by propagating a pyoverdine-producing and pyoverdine-deficient

mutant strain of SBW25 in isolation and together at different inoculation densities under

copper-polluted and unpolluted conditions. We first determined whether pyoverdines

provide an *absolute* growth benefit to the wild-type producer under toxic copper conditions

by comparing the Malthusian parameter ($m$) of individually grown strains using a GLM

with density × strain × copper as explanatory variables, as well as all possible 2- and 3-way

interactions. We next determined whether density differentially affected the performance

of the pyoverdine producing and non-producing strains during direct competition, using a

LMM with $m$ as response variable and strain × frequency as explanatory variables and

random intercepts fitted for each microcosm ($n = 24$) to account for non-independency of

paired observations within microcosms. Finally, we were interested in how density

affected the relative fitness of the pyoverdine-producing strain ($r_{producer} =$

$m_{producer}/m_{non-producer}$) (84) during direct competition with the mutant as a function of

copper stress. We expected the pyoverdine-producing strain to be at a selective

disadvantage under low growth conditions (i.e. when iron is not limiting growth). We

tested this using a GLM with $r_{producer}$ as response variable and copper × density as

explanatory variables, as well as their 2-way interaction.

Finally, we determined whether social exploitation could have played an additional

role in selecting against high siderophore levels in response to copper stress. To this end,

we competed the pyoverdine-producing wildtype strain SBW25 and a pyoverdine-deficient

mutant at a low (1%) or high (99%) frequency in copper-polluted and unpolluted compost.

We determined whether copper differentially affected the relative fitness of the

pyoverdine-deficient mutant ($s_{mutant} = m_{mutant} - m_{wild-type}$) (84) as a function of its

initial frequency. Note that we used the selection rate ($s$) as a measure of relative fitness as

some of the Malthusian growth parameters were negative (SI Appendix, Fig. S7). We

expected the pyoverdine-deficient mutant to reap greater fitness benefits when rare and

tested this hypothesis using a GLM with $s_{mutant}$ as response variable and initial frequency

(low or high) × copper as fixed explanatory variables, as well as their interaction.

**Acknowledgments |** We thank the three anonymous reviewers for their valuable

comments and Daniel Padfield for advice on statistical analyses. This work was funded by

AXA Research Fund, Biotechnology and Biological Sciences Research Council (BBSRC:

BB/K00324) and Natural Environment Research Council (NERC: NE/V012347/1) grants

awarded to AB and a UK Research Innovation (UKRI) Future Leader Fellowship (UKRI-

FLF: MR/V022482/1) awarded to EH. SOB was funded by a 'Bridging the Gaps' award

and PhD scholarship from the University of Exeter. AML was supported by FP7 Marie

Skłodowska-Curie International Incoming Fellowship (IIF) (no. 331163).

**Author Contributions |** EH, AML, SOB, DJH, AB conceived and designed the study. EH,

AML, SOB, TM, JSP, FB collected the data. EH, AN and FB carried out data analyses.

EH, AB wrote the first draft of the manuscript, and all authors contributed to revisions.

**Competing Interest Statement |** The authors declare no competing interests.

**Open Access |** For the purpose of open access, the authors have applied a Creative

Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising

from this submission'.

**References**

1. T. G. Barraclough, How do species interactions affect evolutionary dynamics across whole communities? *Annu. Rev. Ecol. Evol. Syst.* **46**, 25-48 (2015).

2. J. Cairns, R. Jokela, L. Becks, V. Mustonen, T. Hiltunen, Repeatable ecological dynamics govern response of experimental community to antibiotic pulse perturbation. *Nat. Ecol. Evol.* **4**, 1385-1394 (2020).

3. R. Evans *et al.*, Eco-evolutionary dynamics set the tempo and trajectory of metabolic evolution in multispecies communities. *Curr. Biol.* **30**, 4984-4988 (2020).

4. A. B. Chase, C. Weihe, J. B. Martiny, Adaptive differentiation and rapid evolution of a soil bacterium along a climate gradient. *PNAS* **118**, e2101254118 (2021).

5. F. Fiegna, A. Moreno-Letelier, T. Bell, T. G. Barraclough, Evolution of species interactions determines microbial community productivity in new environments. *ISME* **9**, 1235-1245 (2015).

6. N. G. Hairston Jr, S. P. Ellner, M. A. Geber, T. Yoshida, J. A. Fox, Rapid evolution and the convergence of ecological and evolutionary time. *Ecol. Lett.* **8**, 1114-1127 (2005).

7. F. A. Gorter, M. Manhart, M. Ackermann, Understanding the evolution of interspecies interactions in microbial communities. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **375**, 20190256 (2020).

8. J. B. Martiny *et al.*, Investigating the eco‐evolutionary response of microbiomes to environmental change. *Ecol. Lett.* **26**, S81-S90 (2023).

9. M. Lässig, V. Mustonen, A. M. Walczak, Predicting evolution. *Nat. Ecol. Evol.* **1**, 1-9 (2017).

692    10.    N. Meroz, N. Tovi, Y. Sorokin, J. Friedman, Community composition of microbial

693           microcosms follows simple assembly rules at evolutionary timescales. *Nat.*

694           *Commun.* **12**, 1-9 (2021).

695    11.    Y. Yang, Emerging patterns of microbial functional traits. *Trends Microbiol.* **29**,

696           874-882 (2021).

697    12.    M. Cavaliere, S. Feng, O. S. Soyer, J. I. Jiménez, Cooperation in microbial

698           communities and their biotechnological applications. *Environ. Microbiol.* **19**, 2949-

699           2963 (2017).

700    13.    S. Shibasaki, S. Mitri, Controlling evolutionary dynamics to optimize microbial

701           bioremediation. *Evol. Appl.* **13**, 2460-2471 (2020).

702    14.    A. N. Yadav *et al.*, Beneficial microbiomes: biodiversity and potential

703           biotechnological applications for sustainable agriculture and human health. *J. appl.*

704           *biol.* **5**, 4-7 (2017).

705    15.    T. Scheuerl *et al.*, Bacterial adaptation is constrained in complex communities. *Nat.*

706           *Commun.* **11**, 1-8 (2020).

707    16.    M. Castledine, D. Padfield, A. Buckling, Experimental (co) evolution in a multi‐

708           species microbial community results in local maladaptation. *Ecol. Lett.* **23**, 1673-

709           1681 (2020).

710    17.    D. Lawrence *et al.*, Species interactions alter evolutionary responses to a novel

711           environment. *PLoS Biol.* **10**, e1001330 (2012).

712    18.    A. Buckling, R. Craig Maclean, M. A. Brockhurst, N. Colegrave, The Beagle in a

713           bottle. *Nature* **457**, 824-829 (2009).

714    19.    C. De Mazancourt, E. Johnson, T. Barraclough, Biodiversity inhibits species'

715           evolutionary responses to changing environments. *Ecol. Lett.* **11**, 380-388 (2008).

716    20.    P. Gómez, A. Buckling, Real-time microbial adaptive diversification in soil. *Ecol.*

717           *Lett.* **16**, 650-655 (2013).

21.    M. A. Brockhurst, N. Colegrave, D. J. Hodgson, A. Buckling, Niche occupation

limits adaptive radiation in experimental microcosms. *PLoS One* **2**, e193 (2007).

22.    R. S. Etienne, B. Haegeman, A conceptual and statistical framework for adaptive

radiations with a key role for diversity dependence. *Am. Nat.* **180**, E75-E89 (2012).

23.    A. Jousset, N. Eisenhauer, M. Merker, N. Mouquet, S. Scheu, High functional

diversity stimulates diversification in experimental microbial communities. *Sci.*

*Adv.* **2**, e1600124 (2016).

24.    D. I. Bolnick *et al.*, Why intraspecific trait variation matters in community ecology.

*TREE* **26**, 183-192 (2011).

25.    R. Lanfear, H. Kokko, A. Eyre-Walker, Population size and the rate of evolution.

*TREE* **29**, 33-41 (2014).

26.    J. Johansson, Evolutionary responses to environmental changes: how does

competition affect adaptation? *Evolution* **62**, 421-435 (2008).

27.    R. Kummerli, K. T. Schiessl, K. McNeill, M. Ackermann, Habitat structure and the

evolution of diffusible siderophores in bacteria. *Ecol. Lett.* **17**, 1536-1655 (2014).

28.    R. C. Hider, X. Kong, Chemistry and biology of siderophores. *Nat. Prod. Rep.* **27**,

637-657 (2010).

29.    C. Ratledge, L. G. Dover, Iron metabolism in pathogenic bacteria. *Annu. Rev.*

*Microbiol.* **54**, 881-941 (2000).

30.    I. J. Schalk, M. Hannauer, A. Braud, New roles for bacterial siderophores in metal

transport and tolerance. *Environ. Microbiol.* **13**, 2844-2854 (2011).

31.    A. Braud, V. Geoffroy, F. Hoegy, G. L. A. Mislin, I. J. Schalk, Presence of the

siderophores pyoverdine and pyochelin in the extracellular medium reduces toxic

metal accumulation in *Pseudomonas auruginosa* and increases bacterial metal

tolerance. *Environ. Microbiol. Rep.* **2**, 419-425 (2010).

32.    S. O'Brien *et al.*, No effect of intraspecific relatedness on public goods cooperation
744           in a complex community. *Evolution* **72**, 1165-1173 (2018).

745    33.    E. Hesse *et al.*, Ecological selection of siderophore-producing microbial taxa in
746           response to heavy metal contamination. *Ecol. Lett.* **21**, 117-127 (2018).

747    34.    E. Hesse *et al.*, Stress causes interspecific facilitation within a compost community.
748           *Ecol. Lett.* **24**, 2169-2177 (2021).

749    35.    J. Kramer, Ö. Özkaya, R. Kümmerli, Bacterial siderophores in community and host
750           interactions. *Nat. Rev. Microbiol.*, 1-12 (2020).

751    36.    E. Hesse *et al.*, Anthropogenic remediation of heavy metals selects against natural
752           microbial remediation. *Proc B* **286**, 20190804 (2019).

753    37.    P. B. Rainey, M. J. Bailey, Physical and genetic map of the Pseudomonas
754           fluorescens SBW25 chromosome. *Molecular Microbiology* **19**, 521-533 (1996).

755    38.    A. M. Lujan, P. Gomez, A. Buckling, Siderophore cooperation of the bacterium
756           *Pseudomonas fluorescens* in soil. *Biol. Lett.* **11**, 20140934 (2015).

757    39.    C. D. Moon *et al.*, Genomic, genetic and structural analysis of pyoverdine-
758           mediated iron acquisition in the plant growth-promoting bacterium *Pseudomonas*
759           *fluorescens* SBW25. *BMC Microbiol.* **8**, 7 (2008).

760    40.    N. Jiricny *et al.*, Fitness correlates with the extent of cheating in a bacterium. *J.*
761           *Evol. Biol.* **23**, 738-747 (2010).

762    41.    A. S. Griffin, S. A. West, A. Buckling, Cooperation and competition in pathogenic
763           bacteria. *Nature* **430**, 1024-1027 (2004).

764    42.    S. Pollak *et al.*, Public good exploitation in natural bacterioplankton communities.
765           *Sci. Adv.* **7**, eabi4717 (2021).

766    43.    J. Kramer, Ö. Özkaya, R. Kümmerli, Bacterial siderophores in community and host
767           interactions. *Nat. Rev. Microbiol.* **18**, 152-163 (2020).

44.   J. Galet *et al.*, *Pseudomonas fluorescens* pirates both ferrioxamine and

ferricoelichelin siderophores from *Streptomyces ambofaciens*. *Appl. Environ.*

*Microbiol.* **81**, 3132-3141 (2015).

45.   P. Cornelis, S. Matthijs, Diversity of siderophore-mediated iron uptake systems in

fluorescent pseudomonads: not only pyoverdines. *Environ. Microbiol.* **4**, 787-798

(2002).

46.   R. Niehus, A. Picot, N. M. Oliveira, S. Mitri, K. R. Foster, The evolution of

siderophore production as a competitive trait. *Evolution* **71**, 1443-1455 (2017).

47.   A. R. Figueiredo, Ö. Özkaya, R. Kümmerli, J. Kramer, Siderophores drive invasion

dynamics in bacterial communities through their dual role as public good versus

public bad. *Ecol. Lett.* **25**, 138-150 (2022).

48.   J. D. Palmer, K. R. Foster, Bacterial species rarely work together. *Science* **376**,

581-582 (2022).

49.   J. J. Morris, R. E. Lenski, E. R. Zinser, The Black Queen hypothesis: evolution of

dependencies through adaptive gene loss. *MBio* **3**, 2 e00036-00012 (2012).

50.   W. D. Hamilton, The genetical evolution of social baviour. I. *J. Theor. Biol.* **7**, 1-16

(1964).

51.   L. J. Belcher, A. E. Dewar, M. Ghoul, S. A. West, Kin selection for cooperation in

natural bacterial populations. *PNAS* **119**, e2119070119 (2022).

52.   M. Garcia-Garcera, E. P. Rocha, Community diversity and habitat structure shape

the repertoire of extracellular proteins in bacteria. *Nat. Commun.* **11**, 1-11 (2020).

53.   O. X. Cordero, L.-A. Ventouras, E. F. DeLong, M. F. Polz, Public good dynamics

drive evolution of iron acquisition strategies in natural bacterioplankton

populations. *PNAS* **109**, 20059-20064 (2012).

54.   S. A. West, A. S. Griffin, A. Gardner, S. P. Diggle, Social evolution theory for

microorganisms. *Nat. Rev. Microbiol.* **4**, 597-608 (2006).

794 55. S. A. West, S. P. Diggle, A. Buckling, A. Gardner, A. S. Griffin, The social lives of

795   microbes. *Annu. Rev. Ecol. Syst.* **38**, 53-77 (2007).

796 56. B. J. Crespi, The evolution of social behavior in microorganisms. *TREE* **16**, 178-

797   183 (2001).

798 57. J. Gore, H. Youk, A. Van Oudenaarden, Snowdrift game dynamics and facultative

799   cheating in yeast. *Nature* **459**, 253-256 (2009).

800 58. G. Hardin, The tragedy of the commons. *Science* **162**, 1243-1248 (1968).

801 59. R. Axelrod, W. D. Hamilton, The evolution of cooperation. *Science* **211**, 1390-

802   1396 (1981).

803 60. J. L. Sachs, U. G. Mueller, T. P. Wilcox, J. J. Bull, The evolution of cooperation.

804   *Q. Rev. Biol.* **79**, 135-160 (2004).

805 61. H. H. Lee, M. N. Molla, C. R. Cantor, J. J. Collins, Bacterial charity work leads to

806   population-wide resistance. *Nature* **467**, 82-85 (2010).

807 62. I. Frost *et al.*, Cooperation, competition and antibiotic resistance in bacterial

808   colonies. *ISME* **12**, 1582-1593 (2018).

809 63. C. D. Nadell, J. B. Xavier, K. R. Foster, The sociobiology of biofilms. *FEMS*

810   *Microbiol. Rev.* **33**, 206-224 (2008).

811 64. K. P. Rumbaugh *et al.*, Quorum sensing and the social evolution of bacterial

812   virulence. *Curr. Biol.* **19**, 341-345 (2009).

813 65. I. Eleftherianos *et al.*, An antibiotic produced by an insect-pathogenic bacterium

814   suppresses host defenses through phenoloxidase inhibition. *PNAS* **104**, 2419-2424

815   (2007).

816 66. B. Raymond, S. A. West, A. S. Griffin, M. B. Bonsall, The dynamics of

817   cooperative bacterial virulence in the field. *Science* **337**, 85-88 (2012).

67.    T. Köhler, G. G. Perron, A. Buckling, C. Van Delden, Quorum sensing inhibition

selects for virulence and cooperation in *Pseudomonas aeruginosa. PLoS Pathog.* **6**,

e1000883 (2010).

68.    P. Gómez *et al.*, Local adaptation of a bacterium is as important as its presence in

structuring a natural microbial community. *Nat. Commun.* **7**, 12453 (2016).

69.    R. Kummerli, A. S. Griffin, S. A. West, A. Buckling, F. Harrison, Viscous medium

promotes cooperation in the pathogenic bacterium *Pseudomonas aeruginosa. Proc*

*B* **276**, 3531-3538 (2009).

70.    E. Butaitė, M. Baumgartner, S. Wyder, R. Kümmerli, Siderophore cheating and

cheating resistance shape competition for iron in soil and freshwater Pseudomonas

communities. *Nat. Commun.* **8**, 414 (2017).

71.    A. Ross-Gillespie, A. Gardner, S. A. West, A. S. Griffin, Frequency dependence

and cooperation: theory and a test with bacteria. *Am. Nat.* **170**, 331-342 (2007).

72.    H. Heuer, K. Smalla, Plasmids foster diversification and adaptation of bacterial

populations in soil. *FEMS Microbiol. Rev.* **36**, 1083-1104 (2012).

73.    T. Robinson, P. Smith, E. R. Alberts, M. Colussi-Pelaez, M. Schuster, Cooperation

and cheating through a secreted aminopeptidase in the *Pseudomonas aeruginosa*

RpoS response. *Mbio* **11**, 10.1128/mbio. 03090-03019 (2020).

74.    A. Dragoš *et al.*, Evolution of exploitative interactions during diversification in

Bacillus subtilis biofilms. *FEMS Microbiol. Ecol.* **94**, fix155 (2018).

75.    S. O'Brien, D. J. Hodgson, A. Buckling, Social evolution of toxic metal

bioremediation in *Pseudomonas aeruginosa. Proc B* **281**, 20140858 (2014).

76.    A. K. Hottes *et al.*, Bacterial adaptation through loss of function. *PLoS genetics* **9**,

e1003617 (2013).

77.    T. Dimitriu *et al.*, Genetic information transfer promotes cooperation in bacteria.

*PNAS* **111**, 11103-11108 (2014).

78.   D. J. Rankin, E. P. Rocha, S. P. Brown, What traits are carried on mobile genetic

elements, and why? *Heredity (Edinb.)* **106**, 1-10 (2011).

79.   A. E. Dewar, L. J. Belcher, T. W. Scott, S. A. West, Genes for cooperation are not

more likely to be carried by plasmids. *Proc B* **291**, 20232549 (2024).

80.   T. Nogueira *et al.*, Horizontal gene transfer of the secretome drives the evolution of

bacterial cooperation and virulence. *Curr. Biol.* **19**, 1683-1691 (2009).

81.   O. X. Cordero, P. Hogeweg, The impact of long-distance horizontal gene transfer

on prokaryotic genome size. *Proceedings of the National Academy of Sciences* **106**,

21748-21753 (2009).

82.   X. Cheng, I. de Bruijn, M. van der Voort, J. E. Loper, J. M. Raaijmakers, The Gac

regulon of *Pseudomonas fluorescen*s SBW25. *Environ. Microbiol. Rep.* **5**, 608-619

(2013).

83.   B. Schwyn, J. B. Neilands, Universal chemical assay for the detection and

determination of siderophores. *Anal. Biochem.* **160**, 47-56 (1987).

84.   R. E. Lenski, M. R. Rose, S. C. Simpson, S. C. Tadler, Long-term experimental

evolution in *Escherichia coli*. I. Adaptation and divergence during 2000

generations. *Am. Nat.* **138**, 1315-1341 (1991).

85.   B. J. Callahan *et al.*, DADA2: high-resolution sample inference from Illumina

amplicon data. *Nat. Methods* **13**, 581-583 (2016).

86.   M. Kearse *et al.*, Geneious Basic: an integrated and extendable desktop software

platform for the organization and analysis of sequence data. *Bioinformatics* **28**,

1647-1649 (2012).

87.   P. D. Schloss *et al.*, Introducing Mothur: open-source, platform-independent,

community-supported software for describing and comparing microbial

communities. *Appl. Environ. Microbiol.* **75**, 7537-7541 (2009).

88.    R. V. Lenth, Least-squares means: the R package lsmeans. *J. Stat. Softw.* **69**, 1-33

870           (2016).

871    89.    F. Hartig (2020) DHARMa: Residual diagnostics for hierachical (multi-level /

872           mixed) regression models.  R package version 0.3.3.0.

873    90.    H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer Verlag, New

874           York, 2016).

875    91.    D. Bates, M. Maechler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models

876           Using lme4. *J. Stat. Softw.* **67**, 1-48 (2015).

877    92.    J. Oksanen *et al.* (2010) vegan: Community Ecology Package. R package version

878           2.5-4.

879    93.    M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and

880           dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

881    94.    M. E. Brooks *et al.*, glmmTMB balances speed and flexibility among packages for

882           zero-inflated generalized linear mixed modelling. *The R Journal* **9**, 378-400 (2017).

883    95.    M. Torchiano (2016) Effsize - A package for efficient effect size computation

884    96.    A. Vehtari, A. Gelman, J. Gabry, Practical Bayesian model evaluation using leave-

885           one-out cross-validation and WAIC. *Stat. Comput.* **27**, 1413-1432 (2017).

886    97.    H. Ge, K. Xu, Z. Ghahramani (2018) Turing: a language for flexible probabilistic

887           inference. in *International conference on artificial intelligence and statistics*

888           (PMLR), pp 1682-1690.

889    98.    J. Bezanson, A. Edelman, S. Karpinski, V. B. Shah, Julia: A fresh approach to

890           numerical computing. *SIAM review* **59**, 65-98 (2017).

891    99.    M. D. Hoffman, A. Gelman, The No-U-Turn sampler: adaptively setting path

892           lengths in Hamiltonian Monte Carlo. *J. Mach. Learn. Res.* **15**, 1593-1623 (2014).

893    100.   P. K. Mogensen, A. N. Riseth, Optim: A mathematical optimization package for

894           Julia. *JOSS* **3** (2018).

**Figure legends**

**Figure 1 | Copper favors siderophore-producing taxa within the compost community.**

Panel **(A)** shows that copper-polluted (copper [+], orange) communities ($n =6$) produced on average larger amounts of siderophore compared to copper-free (copper [-], blue) communities following six weeks of evolution in isolation (light shading) or in co-culture with SBW25 (dark shading) ($n = 6$ communities per treatment). Brown line and shaded area depict mean $\pm$ SE siderophore production of the ancestral communities ($n = 12$) prior to copper amendment. Boxes depict upper and lower quartiles of treatment-specific raw data with the center line showing the median and whiskers providing a measure of 1.5x interquartile range. Points represent the mean value per community based on twenty-four isolates, with within-community variance indicated by different point sizes. Note that for some replicate communities we isolated < 24 isolates, yielding a total sample size of $n = 548$. Panel **(B)** depicts a Principal Coordinate Analysis (PCoA) plot based on Bray-Curtis dissimilarities between compost communities (excluding SWB25). The percentage of variation explained is shown on each axis, calculated from the relevant eigenvalues. Communities belonging to the same treatment are joined with straight colored lines (control$_{no\ SBW25}$ = light blue, control$_{SBW25}$ = dark blue, copper$_{no\ SBW25}$ = orange, copper$_{SBW25}$= red), with large points representing treatment-specific centroids and small points individual microcosms. Dissimilarities were calculated using data on the total abundance of different isolates (identified at genus level) within each community that were assayed for siderophore production ($n = 24$ per community). Panel **(C)** depicts the effect of coper on the abundance of bacterial taxa in the compost community. The log2-fold change in total abundance of the ten most common genera when comparing communities that had evolved in unpolluted versus copper-polluted compost. Of the taxa tested, three significantly differed in terms of total abundance between copper-free versus copper-

polluted treatments: while *Cupviavidus* was positively affected*, Achromobacter and*

*Lysinibacillus* were both negatively affected by copper stress (see SI Appendix, Table S1

table for Wald test and P-values). Points represent estimated log2-fold change in total

abundance of each taxon in response to copper-stress (averaged across culturing

conditions) ± standard error bars. Genera are color-coded based on their mean across-

treatment siderophore production, ranging from relatively low (blue) to intermediate

(orange) and high (red) siderophore production. Panel **(D)** depicts the relative abundance of

the ten most common culturable compost taxa, excluding SWB25, following six weeks of

evolution under different selection regimes. Genera are listed in order of their across-

treatment mean siderophore production, increasing from top to bottom, such that orange-

coral taxa produce relatively intermediate to high levels of siderophores.

**Figure 2 | Toxic copper favors the evolution of reduced siderophore production within**

**SBW25**. Boxplot showing that copper (copper [+], brown) selected for lower mean

siderophore levels in SBW25 populations compared to copper-free conditions (copper [-],

pink) irrespective of whether populations evolved in isolation (light-shading) or in co-

culture with the compost community (dark-shading) ($n = 6$ populations per treatment).

Orange line and shaded area depict mean ± SE siderophore production of ancestral SBW25

populations ($n = 12$) prior to copper amendment. Boxes depict the upper and lower

quartiles of treatment-specific raw data with the center line showing the median and

whiskers providing a measure of 1.5x interquartile range. Points represent the mean value

per population ($n = 6$ per copper treatment) based on twenty-four clones where possible,

with within-population variance given by different point sizes. Note that for some replicate

communities we isolated < 24 clones, yielding a total sample size of $n = 501$.

**Figure 3 | Copper leads to trait convergence by selecting against siderophore extremes.**

Panel **(A)** shows the frequency distributions of siderophore production in ancestral SBW25 populations (blue) and compost communities (brown) prior to copper amendment. Siderophore data were pooled across copper treatments and replicates ($n = 24$ isolates for each of the twelve replicate SBW25 populations / compost communities). Panel **(B)** demonstrates that the distribution of siderophore production in SBW25 (blue) and the community (brown) converges in response to copper stress, independent of whether these evolved in isolation (top panel) or in co-culture (bottom panel). Siderophore data were pooled across replicates ($n = 24$ isolates for replicate population/community). Note that for some replicates we isolated < 24 clones/isolates of SBW25 or the community, yielding a total sample size of $n = 1049$. Brown and blue vertical lines depict mean siderophore production of the ancestral compost communities and SBW25 populations, respectively.

**Figure 4 | Benefit of different siderophore strategies is context dependent.** Boxplot showing **(A)** the Malthusian growth parameter ($m$) of the pyoverdine-deficient mutant (blue) and pyoverdine-producing wild-type SBW25 strain (gold) as a function of inoculation density when individually grown in copper-polluted (copper [+]) and unpolluted (copper [-]) compost **(B)** the pyoverdine-deficient 'cheat' receiving a relative selective benefit ($S_{mutants} > 0$) when rare, in particular when competing with the wild-type producer in copper-polluted compost. Strains were co-inoculated together at different frequencies (1% or 99%) in copper-polluted (copper [+]) and unpolluted (copper [-]) compost. Blue horizontal line depicts equal fitness ($S_{mutants} = 0$) of competing strains. Boxes depict the upper and lower quartiles of treatment-specific raw data with the center line showing the median and whiskers providing a measure of 1.5x interquartile range, and points show the raw data for each replicate population.

**Figure 1**

 **Figure 2**

972 **Figure 3**