Interspecific social interactions shape public goods production in natural

microbial communities

One sentence summary – Interspecific exploitation shapes the evolution and ecology of

public goods production

Elze Hesse[1*], Siobhan O'Brien[1, 2], Adela M. Luján[1, 3], Florian Bayer[1], Eleanor M. van

Veen[4], Dave J. Hodgson[5] & Angus Buckling[1]

[1] CEC & ESI, Biosciences, University of Exeter, Penryn Campus, Cornwall, TR10 9FE,

UK.

[2] Institute of Integrative Biology, University of Liverpool, Biosciences Building, Crown

Street, L69 7ZB, Liverpool, UK.

[3] CIQUIBIC, Departamento de Química Biológica, Facultad de Ciencias Químicas,

CONICET, Universidad Nacional de Córdoba, Córdoba, Argentina.

[4] Camborne School of Mines, CEMPS, University of Exeter, Penryn Campus, Cornwall

TR10 9FE, UK.

[5] CEC, Biosciences, University of Exeter, Penryn Campus, Cornwall, TR10 9FE, UK.

[*] Correspondence: E-mail: elzehesse@gmail.com; Tel: +44 (0) 1326 259470; Fax: +44

(0) 1392 263434

Abstract

Some microbial public goods benefit conspecifics, as well as other species. Here, we use

evolution and competition experiments to determine how exploitation of public goods by

the wider microbial community shapes the production of an interspecific public good:

metal-detoxifying siderophores. By simultaneously studying whole microbial

communities and an embedded focal species, we show that interspecific exploitation

results in both ecological selection against microbial taxa that produce relatively large

amounts of siderophores, and evolution of reduced siderophore production within taxa

over similar time scales. Our findings demonstrate the crucial role of interspecific

interactions in shaping microbial social behaviours.

Keywords – ecology, evolution, microbial communities, public goods, siderophores,

interspecific exploitation, Black Queen Hypothesis.

Main text

Microbes produce a range of metabolically costly public goods that improve their

growth and survival, but that are open to exploitation by non-producing 'cheats' (*1-6*).

Optimal levels of public goods therefore reflect a balance of the benefits received by the

producer and individuals who share the same genes, and costs associated with enhancing

the fitness of exploiting competitors. The evolution of public goods has typically been

studied in the context of within-species interactions (*7*), but many microbial public goods,

including antibiotic-degrading enzymes (*8, 9*), resource-scavenging molecules (*10, 11*)

and immune-manipulating effectors (*12, 13*), potentially benefit not only conspecifics but

also other species. As a consequence, interspecific exploitation may play a key role in

shaping the production of public goods, and may select for the loss of key metabolic

genes (*14, 15*). However, experimental evidence for interspecific exploitation of

community-wide public goods, and the resultant ecological and evolutionary

consequences, is lacking.

Metal-detoxifying siderophores have the potential to act as a community-wide

public good (*16-18*). The canonical function of siderophores is to bind and take up

insoluble iron (*19*), but these extracellular agents can also bind to toxic heavy metals (*20,*

21) to prevent uptake by cells (*22*). Siderophores can therefore provide protection from

toxic metals both to conspecifics (*23*) and to members of different species (*16*). There is

good evidence for intraspecific exploitation of metal-detoxifying siderophores, as low

siderophore-producers of *Pseudomonas aeruginosa* – an opportunistic bacterial pathogen

– grow poorly alone but can invade isogenic wild-type producers (*16*). Here, we explore

whether interspecific exploitation also plays an important role in shaping siderophore

production, both through competition between species and through evolutionary change.

We previously determined how mean siderophore production and community

composition changed as a function of copper pollution in natural soil and experimental

compost microbial communities (*17*). We found that copper favours higher siderophore-

producing microbial taxa, thereby increasing mean levels of siderophore production in

polluted compared to non-polluted communities. This suggests there are direct and/or

kin-selected benefits to producing siderophores (*24, 25*). Re-analysis of this dataset

suggests that high siderophore-producers are also open to exploitation: copper-mediated

increases in mean siderophore levels are accompanied by a reduction in the variation

(dispersion parameter in Gaussian double GLM with copper as explanatory variable: $\chi^2 =$

22.85, df = 1, $P < 0.001$), demonstrating selection against high as well as low

siderophore-producing taxa (Fig. 1A). Moreover, genera producing relatively large

amounts of siderophores in non-polluted compost microcosms displayed lower

production in copper-polluted compost, and *vice versa* for low siderophore-producing

taxa (copper × genus interaction in 2-way ANOVA: $F_{5,\,170} = 2.87$ and $P = 0.02$; Fig. 1B).

7   While selection against high levels of siderophore production could be a

consequence of interspecific exploitation, this may also result solely from intraspecific

exploitation or even not be driven by social interactions at all. To determine if

interspecific exploitation shapes selection for siderophore production, we carried out all

possible pairwise competitions between ten random compost isolates (i.e. isolated from

experimental compost communities) spanning the observed continuum of siderophore

production (Table S1), as well as growing each in isolation. By conducting all possible

pairwise combinations, we tested for the effects of siderophores *per se* over and above

any other competitive differences between isolates. In copper-contaminated compost,

isolates that produced relatively large amounts of siderophore grew better in isolation

than in competition, whereas non- or low-producing isolates benefitted from the presence

of other taxa (linear model on $m_{\text{competition}} - m_{\text{isolation}}$: $F_{1,\,88} = 10.95$, $P = 0.001$ for main

effect of siderophore production; Fig. 2A). This demonstrates that high siderophore-

producing taxa were exploited by low siderophore-producing taxa. The relationship

between siderophore production and relative fitness in the presence *versus* absence of

competitors was not significant in non-polluted compost, where the importance of

siderophores as an interspecific public good is likely to be greatly reduced (linear model:

$F_{1, 88} = 0.05$, $P = 0.82$; Fig. 2B).

The above findings demonstrate that interspecific exploitation imposes selection

on siderophore production, and that this can result in selection against high siderophore-

producing taxa. Given that microbial evolution can occur on time scales concomitant with

changes in community composition, we hypothesised that interspecific exploitation could

result in selection for lower siderophore production in initially high-siderophore

producing taxa in these communities. To test this, we conducted a similar experiment to

our original study (*17*) but this time also followed the evolution of a focal taxon,

Pseudomonas fluorescens SBW25 (*26*). This common soil bacterium produces a range of

siderophores (*27*) known to chelate metals other than iron (*28*).

Changes in community-wide siderophore production in response to copper

addition were qualitatively the same as we found before (*17*), with copper resulting in

greater mean (LME with copper as fixed effect and random intercepts fitted for individual

microcosms: $\chi^2 = 7.16$, df = 1, $P = 0.007$; Fig. 3A), but lower variation in siderophore

production (Gaussian double GLM: $\chi^2 = 14.40$, df = 1, $P < 0.001$, Fig. 3B). Copper-

mediated shifts in siderophore production were also associated with compositional

changes in the microbial assemblage (PERMANOVA: $F_{1, 10} = 6.26$, $P = 0.003$, $R^2 = 0.38$;

multivariate dispersion: $F_{1, 10} = 0.05$, $P = 0.82$; Fig. 3C), such that copper favoured

microbial taxa that on average produced larger amounts of siderophore (Fig. 3D).

P. fluorescens – initially a very high siderophore-producing species – evolved to

produce significantly less siderophore in copper-polluted compost (Fig. 4). However,

evolutionary trajectories of *P. fluorescens* were the same whether the natural community

was present or absent (LME: $\chi^2 = 1.63$, df $= 1$, $P = 0.02$ for main community effect), with

copper consistently reducing mean siderophore levels ($\chi^2 = 10.56$, df $= 1$, $P < 0.001$ for

main copper effect: Fig. 4A–B). There are a number of interpretations of this finding.

First, *P. fluorescens* siderophore production is not driven by exploitation, but may simply

have been higher than needed to detoxify the environment. Second, exploitation occurred,

but it is entirely driven by exploitation within species. These interpretations do not invoke

interspecific exploitation, but the absence of an effect of interspecific exploitation on

siderophore evolution seems surprising given its strong effect on ecological selection

(Fig. 3).

A third interpretation is therefore that intraspecific exploitation compensates for

interspecific exploitation when other species are not present. Given the community-wide

nature of the public good, exploitation by competitors can theoretically have the same

evolutionary consequences on a focal population regardless of whether the exploitation is

by the same or different species (*20*). We hypothesised that in the presence of the

community, *P. fluorescens* siderophore producers were exploited by both other taxa and

evolved *P. fluorescens*, while in the absence of the community the greatly increased

density of *P. fluorescens* (Fig. S1) resulted in comparable levels of exploitation from

conspecifics alone.

To determine whether or not intra- and interspecific exploitation played a role in

shaping *P. fluorescens* evolutionary trajectories, we conducted a series of short-term

(week-long) growth rate assays of wild-type SBW25 and an isogenic pyoverdine knock-

out mutant strain (*29*) in copper-polluted compost. This mutant does not produce the

main siderophore pyoverdine (which reduces total siderophore production to 21% of the

wild-type), and displays levels of siderophore production similar to the average compost

community member. In the absence of the community, the *P. fluorescens* wild-type and

mutant grew equally well as monocultures (Fig. 5A), demonstrating that there is no

abiotic selection against high siderophore production. However, the pyoverdine mutant

outcompeted the wild-type in co-culture in a frequency dependent manner (growth rate:

$F_{2, 56} = 31.67$, $P < 0.001$ for strain × frequency interaction; selection coefficient: $F_{2, 27} =$

179.87, $P < 0.001$ for effect of frequency; Fig. 5B). Such dynamics are indicative of

intraspecific exploitation of high siderophore producers (10, 35). We next determined

how the producer and mutant each independently fared in the presence *versus* absence of

the community. As above, in the absence of the community the siderophore producing

wild-type and pyoverdine mutant grew equally well as monocultures (Fig. 5C). The

presence of the community did not notably affect the growth rate of the mutant,

suggesting little or no net exploitation of detoxifying siderophores by other community

members. By contrast, the growth rate of the wild-type was markedly reduced by the

community, strongly suggesting exploitation of its siderophores (2-way ANOVA: social

background × strain = $F_{1, 39} = 9.42$, $P < 0.01$; Fig. 5C). As a net result, the wild-type had a

significantly lower fitness compared to the mutant when growing together with the

natural compost community (ANOVA: $F_{1, 19} = 12.65$, $P < 0.01$; Fig. 5D). Taken together,

these results suggest both intra- and interspecific exploitation played an important role in

driving the evolution of reduced siderophore production in *P. fluorescens*.

To conclude, we show that interspecific exploitation plays a key role in shaping

community-wide levels of public goods, selecting for and against microbial taxa that

differ in their mean siderophore production. Moreover, interspecific (in addition to

intraspecific) exploitation plays a key role in the driving within-taxon evolutionary

changes. While both ecological and evolutionary processes operate over similar time

frames, we anticipate that evolutionary effects resulting from interspecific interactions

will become increasingly important relative to ecological selection once community

composition has reached equilibrium. Indeed, this is precisely the scenario envisaged by

the Black Queen Hypothesis (*14*), where selection favours intermediate levels of public

goods because of differential benefits and selection to lose traits as a result of

interspecific exploitation. Given that microbial public goods with community-wide

benefits are extremely common (*10, 30*), our results suggest interspecific exploitation

likely plays a key role in shaping microbial community composition and evolution in

nature.

Acknowledgments

We thank Daniel Padfield for providing the R code to plot community composition data

and Uli Klümper for advice on analysing Sanger sequences. We are also grateful to

Michiel Vos for critical comments that improved the manuscript. **Funding:** This work

was funded by the AXA Research Fund, BBSRC and NERC to AB. SOB was funded by

a "Bridging the Gaps" award and PhD scholarship from the University of Exeter. AML

was supported by Marie Curie International Incoming Fellowships within the EU Seventh

Framework Programme. AB acknowledges support from the Royal Society. **Authorship:**

EH, SOB, AB conceived and designed the experiment. EH, SOB, FB, AL, EvV collected

the data. EH carried out data analyses. EH, AB wrote the first draft of the manuscript, and

all authors contributed to revisions. **Competing interests:** none to declare. **Data and**

material availability: Sequences have been deposited as ENA Project PRJEB29924

(https://www.ebi.ac.uk/ena/data/search?query=PRJEB29924). Experimental data will be

uploaded on Dryad.

Supplementary Materials

Material and Methods

Table S1 – Siderophore production of compost isolates used in pairwise competitions.

Table S2 – Pairwise treatment contrasts accompanying Figure 5 in the main text.

Figure S1 – The effects of copper and interspecific competition on *Pseudomonas*

fluorescens population densities following six weeks of evolution.

References

1. B. J. Crespi, The evolution of social behavior in microorganisms. *TREE* **16**, 178-183 (2001).

2. S. A. West, A. S. Griffin, A. Gardner, S. P. Diggle, Social evolution theory for microorganisms. *Nat. Rev. Microbiol.* **4**, 597-608 (2006).

3. J. Gore, H. Youk, A. Van Oudenaarden, Snowdrift game dynamics and facultative cheating in yeast. *Nature* **459**, 253-256 (2009).

4. G. Hardin, The tragedy of the commons. *Science* **162**, 1243-1248 (1968).

5. R. Axelrod, W. D. Hamilton, The evolution of cooperation. *Science* **211**, 1390-1396 (1981).

6. J. L. Sachs, U. G. Mueller, T. P. Wilcox, J. J. Bull, The evolution of cooperation. *Q. Rev. Biol.* **79**, 135-160 (2004).

7. S. Mitri, K. R. Foster, The genotypic view of social interaction in microbial communities. *Annu. Rev. Genet.* **47**, 247-273 (2013).

8. H. H. Lee, M. N. Molla, C. R. Cantor, J. J. Collins, Bacterial charity work leads to population-wide resistance. *Nature* **467**, 82-85 (2010).

9. I. Frost, W. P. Smith, S. Mitri, A. San Millan, Y. Davit, J. M. Osborne, J. M. Pitt-Francis, R. C. MacLean, K. R. Foster, Cooperation, competition and antibiotic resistance in bacterial colonies. *ISME* **12**, 1582-1593 (2018).

10. S. A. West, S. P. Diggle, A. Buckling, A. Gardner, A. S. Griffin, The social lives of microbes. *Annu. Rev. Ecol. Syst.* **38**, 53-77 (2007).

11. C. D. Nadell, J. B. Xavier, K. R. Foster, The sociobiology of biofilms. *FEMS Microbiol. Rev.* **33**, 206-224 (2008).

12.   K. P. Rumbaugh, S. P. Diggle, C. M. Watters, A. Ross-Gillespie, A. S. Griffin, S.

A. West, Quorum sensing and the social evolution of bacterial virulence. *Curr.*

Biol. **19**, 341-345 (2009).

13.   I. Eleftherianos, S. Boundy, S. A. Joyce, S. Aslam, J. W. Marshall, R. J. Cox, T. J.

Simpson, D. J. Clarke, S. E. Reynolds, An antibiotic produced by an insect-

pathogenic bacterium suppresses host defenses through phenoloxidase inhibition.

Proc Natl Acad Sci U S A **104**, 2419-2424 (2007).

14.   J. J. Morris, R. E. Lenski, E. R. Zinser, The Black Queen hypothesis: evolution of

dependencies through adaptive gene loss. *MBio* **3**, 2 e00036-00012 (2012).

15.   O. X. Cordero, L.-A. Ventouras, E. F. DeLong, M. F. Polz, Public good dynamics

drive evolution of iron acquisition strategies in natural bacterioplankton

populations. *Proc Natl Acad Sci U S A* **109**, 20059-20064 (2012).

16.   S. O'Brien, E. Hesse, A. M. Lujan, D. J. Hodgson, A. Gardner, A. Buckling, No

effect of intraspecific relatedness on public goods cooperation in a complex

community. *Evolution* **72**, 1165-1173 (2018).

17.   E. Hesse, S. O'Brien, A. M. Lujan, N. Tromas, F. Bayer, E. van Veen, D. J.

Hodgson, A. Buckling, Ecological selection of siderophore-producing microbial

taxa in response to heavy metal contamination. *Ecol. Lett.* **21**, 117-127 (2018).

18.   S. Estrela, E. Libby, J. Van Cleve, F. Débarre, M. Deforet, W. R. Harcombe, J.

Peña, S. P. Brown, M. E. Hochberg, Environmentally mediated social dilemmas.

TREE **34**, 6-18 (2019).

19.   R. C. Hider, X. Kong, Chemistry and biology of siderophores. *Nat. Prod. Rep.* **27**,

637-657 (2010).

20.   A. Braud, F. Hoegy, K. Jezequel, T. Lebeau, I. J. Schalk, New insights into the

metal specificity of the *Pseudomonas aeruginosa* pyoverdine-iron uptake

pathway. *Environ. Microbiol.* **11**, 1079-1091 (2009).

21.   I. J. Schalk, M. Hannauer, A. Braud, New roles for bacterial siderophores in metal

transport and tolerance. *Environ. Microbiol.* **13**, 2844-2854 (2011).

22.   A. Braud, V. Geoffroy, F. Hoegy, G. L. A. Mislin, I. J. Schalk, Presence of the

siderophores pyoverdine and pyochelin in the extracellular medium reduces toxic

metal accumulation in *Pseudomonas auruginosa* and increases bacterial metal

tolerance. *Environ. Microbiol. Rep.* **2**, 419-425 (2010).

23.   S. O'Brien, D. J. Hodgson, A. Buckling, Social evolution of toxic metal

bioremediation in *Pseudomonas aeruginosa*. *Proc R Soc Lond [Biol]* **281**,

20140858 (2014).

24.   N. M. Oliveira, R. Niehus, K. R. Foster, Evolutionary limits to cooperation in

microbial communities. *Proc Natl Acad Sci U S A* **111**, 17941-17946 (2014).

25.   R. Niehus, A. Picot, N. M. Oliveira, S. Mitri, K. R. Foster, The evolution of

siderophore production as a competitive trait. *Evolution* **71**, 1443-1455 (2017).

26.   P. B. Rainey, M. J. Bailey, Physical and genetic map of the *Pseudomonas*

fluorescens SBW25 chromosome. *Mol. Microbiol.* **19**, 521-533 (1996).

27.   P. Cornelis, S. Matthijs, Diversity of siderophore-mediated iron uptake systems in

fluorescent pseudomonads: not only pyoverdines. *Environ. Microbiol.* **4**, 787-798

(2002).

28. A. Bhattacharya, Siderophore mediated metal uptake by Pseudomonas fluorescens

and its comparison to iron (iii) chelation. *Cey. J. Sci. (Bio. Sci.)* **39**, 147-155

(2011).

29. C. D. Moon, X.-X. Zhang, S. Matthijs, M. Schäfer, H. Budzikiewicz, P. B.

Rainey, Genomic, genetic and structural analysis of pyoverdine-mediated iron

acquisition in the plant growth-promoting bacterium *Pseudomonas fluorescens*

SBW25. *BMC Microbiol.* **8**, 7 (2008).

30. G. D'Souza, S. Shitut, D. Preussger, G. Yousif, S. Waschina, C. Kost, Ecology

and evolution of metabolic cross-feeding interactions in bacteria. *Nat. Prod. Rep.*

35, 455-488 (2018).

31. B. Schwyn, J. B. Neilands, Universal chemical assay for the detection and

determination of siderophores. *Anal. Biochem.* **160**, 47-56 (1987).

32. A. M. Lujan, P. Gomez, A. Buckling, Siderophore cooperation of the bacterium

Pseudomonas fluorescens in soil. *Biol. Lett.* **11**, 20140934 (2015).

33. B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, S. P.

Holmes, DADA2: high-resolution sample inference from Illumina amplicon data.

Nat. Methods **13**, 581-583 (2016).

34. R. E. Lenski, M. R. Rose, S. C. Simpson, S. C. Tadler, Long-term experimental

evolution in *Escherichia coli*. I. Adaptation and divergence during 2000

generations. *Am. Nat.* **138**, 1315-1341 (1991).

35. A. Buckling, P. B. Rainey, Antagonistic coevolution between a bacterium and a

bacteriophage. *Proc R Soc Lond [Biol]* **269**, 931-936 (2002).

36.   D. Bates, M. Maechler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects

Models Using lme4. *J. Stat. Softw.* **67**, 1-48 (2015).

37.   P. K. Dunn, G. K. Smyth, dglm: Double Generalized Linear Models *R package*

version 1.8.3. (2016). (https://CRAN.R-project.org/package=dglm).

38.   G. K. Smyth, Generalized linear models with varying dispersion. *J. R. Stat. Soc.*

B. **51**, 47-60 (1989).

39.   J. Oksanen, F. G. Blanchet, R. Kindt, P. Legendre, R. B. O'Hara, G. L. Simpson,

P. Solymos, M. H. H. Stevens, H. Wagner, vegan: Community Ecology Package.

R package version 2.5-4. (2010). (https://cran.r-project.org/package=vegan).

40.   R. V. Lenth, Least-squares means: the R package lsmeans. *J. Stat. Softw.* **69**, 1-33

(2016).

Figure 1. Variation in siderophore production in compost communities and across individual microbial taxa. (A) Density plot demonstrating that copper (red) selects against very low and very high siderophore-producing taxa following six weeks of evolution in compost. Variation in siderophore production was lower in polluted compared non-polluted (blue) communities. **(B)** Box–whisker plot depicting variation in siderophore production in six common culturable microbial genera as a function of copper pollution (red = polluted and blue = non-polluted communities) following six weeks of evolution. Boxes depict the upper and lower quartiles of the treatment-specific raw data with the centre line showing the median and whiskers providing a measure of the 1.5x interquartile range. Siderophore production was quantified for multiple isolates (n = 13–62) per genus using individual liquid CAS assays (*31*), corrected for variation in cell density as measured by OD_{600}.

Figure 2. Copper pollution changes the relationship between siderophore production and the relative performance of compost isolates during pairwise competition. Species performance ($n = 10$) was calculated as: $d = m_{competition} - m_{isolation}$. In scenarios where $d > 0$, compost isolates benefitted from the presence of other species, whereas $d < 0$ indicates isolates grew better in isolation. Small dots depict d for all possible pairwise competitions ($n = 9$ per isolate) and large dots provide a measure of across-species mean performance in **(A)** copper-polluted and **(B)** non-polluted microcosms. Line and shaded area depict the fitted relationship ± 95% CI for copper ($d = 0.16[-0.07, 0.38] - 31.85[-50.96, -12.73] \times$ siderophores) and control microcosms ($d = -0.32[-0.54, -0.10] - 2.11[-20.58, 16.37] \times$ siderophores). Siderophore production was quantified using individual liquid CAS assays (*31*), corrected for variation in OD_{600}.

Figure 3. The effect of copper pollution on microbial community composition and siderophore production in the presence of *P. fluorescens*. (A) Copper-polluted communities produced larger amounts of siderophore compared to non-polluted communities following six weeks of evolution. Boxes depict upper and lower quartiles of treatment-specific raw data with the centre line showing the median and whiskers providing a measure of 1.5x interquartile range. Points represent mean values per community. Letters denote significant contrasts, with α < 0.05. **(B)** Copper (red) selects against very low and very high siderophore-producing taxa, thereby reducing variation in siderophore production compared to non-polluted (blue) communities. **(C)** Principal Coordinate Analysis (PCoA) plot based on Bray-Curtis dissimilarities between communities. The percentage of variation explained is shown on each axis, calculated

from the relevant eigenvalues. Communities belonging to the same treatment are joined

with straight coloured lines (copper = red and control = blue), with large points

representing treatment-specific centroids and small points individual microcosms. **(D)**

Relative abundance of ten common culturable genera, listed in order of their mean

across-treatment siderophore production, increasing from top to bottom, such that orange-

red taxa are intermediate-high producers. Siderophore production was quantified using

individual CAS assays (*31*), corrected for variation in OD_{600}.

Figure 4. Toxic copper selects against high siderophore-producers in focal

Pseudomonas fluorescens. Box–whisker plots demonstrating that mean siderophore

levels in *P. fluorescens* populations are reduced under copper stress when evolving in the

presence **(A)** and absence **(B)** of the compost microbial community. Boxes depict the

upper and lower quartiles of treatment-specific raw data with the centre line showing the

median and whiskers providing a measure of 1.5x interquartile range, where black, blue

and red depict ancestral (i.e. prior to copper amendment; $n = 12$), non-polluted ($n = 6$)

and copper-polluted evolved populations ($n = 6$), respectively. Points represent the mean

value per population based on twenty-four clones. Siderophore production per clone was

quantified using individual liquid CAS assays (*31*), corrected for variation in OD_{600}.

Letters denote significant contrasts calculated based on the full LME model testing the

interactive effects of copper $\times$ social background on mean siderophore levels by

averaging across social backgrounds (copper – control contrast: *t* ratio = -3.47, *P* =

0.002).

Figure 5. Public goods exploitation decreases the relative fitness of high siderophore producers in copper-polluted compost. Box–whisker plots depicting **(A)** Malthusian growth rates (*m*) of a wild-type pyoverdine producer and isogenic knockout mutant of *Pseudomons fluorescens* when grown as monocultures (black) or together at high (yellow) or low (blue) frequencies. In a separate experiment **(B)**, strains were individually grown in copper-polluted compost in the presence (red) or absence (black) of the compost community. Boxes depict upper and lower quartiles of treatment-specific raw data with the centre line showing the median and whiskers providing a measure of the 1.5x interquartile range. Points represent individual microcosms ($n = 6 – 12$ per treatment). Box–whisker plots depicting the relative fitness of the wild-type producer **(C–D)** under different social conditions (see above for colour coding). A selection coefficient of zero indicates the wild-type and mutant have equal fitness. Different letters denote significant

1 treatment effects based on pairwise Bonferroni-adjusted contrasts, with $\alpha < 0.05$ (Table

2 S2).