The plant mobile domain proteins MAIN and MAIL1 interact with the

phosphatase PP7L to regulate gene expression and silence transposable

elements in *Arabidopsis thaliana*.

Short title: The PMD MAIN/MAIL1 and PP7L complex regulates gene expression and TE silencing.

Melody Nicolau[1,2], Nathalie Picault[1,2], Julie Descombin[1,2], Yasaman Jami-Alahmadi[3], Suhua Feng[4],

Etienne Bucher[5], Steven E. Jacobsen[4,6], Jean-Marc Deragon[1,2], James Wohlschlegel[3] and Guillaume

Moissiard[1,2]*.

1 LGDP-UMR5096, CNRS, Perpignan, France.

2 LGDP-UMR5096, Université de Perpignan Via Domitia, France.

3 Department of Biological Chemistry, University of California at Los Angeles, Los Angeles, CA, USA.

4 Department of Molecular, Cell and Developmental Biology, University of California at Los Angeles,

Los Angeles, CA, USA.

5 Plant Breeding and Genetic Resources, Agroscope, Nyon, Switzerland.

6 Howard Hughes Medical Institute, University of California at Los Angeles, Los Angeles, CA, USA.

* Corresponding author

E-mail: guillaume.moissiard@univ-perp.fr

ABSTRACT

Transposable elements (TEs) are DNA repeats that must remain silenced to ensure cell integrity. Several epigenetic pathways including DNA methylation and histone modifications are involved in the silencing of TEs, and in the regulation of gene expression. In *Arabidopsis thaliana*, the TE-derived plant mobile domain (PMD) proteins have been involved in TE silencing, genome stability, and control of developmental processes. Using a forward genetic screen, we found that the PMD protein MAINTENANCE OF MERISTEMS (MAIN) acts synergistically and redundantly with DNA methylation to silence TEs. We found that MAIN and its close homolog MAIN-LIKE 1 (MAIL1) interact together, as well as with the phosphoprotein phosphatase (PPP) PP7-like (PP7L). Remarkably, *main*, *mail1, pp7l* single and *mail1 pp7l* double mutants display similar developmental phenotypes, and share common subsets of upregulated TEs and misregulated genes. Finally, phylogenetic analyses of PMD and PP7-type PPP domains among the Eudicot lineage suggest neo-association processes between the two protein domains to potentially generate new protein function. We propose that, through this interaction, the PMD and PPP domains may constitute a functional protein module required for the proper expression of a common set of genes, and for silencing of TEs.

AUTHOR SUMMARY

The plant mobile domain (PMD) is a protein domain of unknown function that is widely spread in the angiosperm plants. Although most PMDs are associated with repeated DNA sequences called transposable elements (TEs), plants have domesticated the PMD to produce genic versions that play important roles within the cell. In *Arabidopsis thaliana*, MAINTENANCE OF MERISTEMS (MAIN) and MAIN-LIKE 1 (MAIL1) are genic PMDs that are involved in genome stability, developmental processes, and silencing of TEs. The mechanisms involving MAIN and MAIL1 in these cellular processes remain elusive. Here, we show that MAIN, MAIL1 and the phosphoprotein phosphatase (PPP) named PP7-like (PP7L) interact to form a protein complex that is required for the proper expression of genes, and the

47 silencing of TEs. Phylogenetic analyses revealed that PMD and PP7-type PPP domains are evolutionary

48 connected, and several plant species express proteins carrying both PMD and PPP domains. We

49 propose that interaction of PMD and PPP domains would create a functional protein module involved

50 in mechanisms regulating gene expression and repressing TEs.

52 INTRODUCTION

In eukaryotes, DNA methylation and post-translational modifications of histones are

epigenetic marks involved in chromatin organization, regulation of gene expression and silencing of

DNA repeats such as transposable elements (TEs) [1-3]. Constitutive heterochromatin is highly

condensed and enriched in silenced TEs that are targeted by DNA methylation and histone H3 lysine

9 dimethylation (H3K9me2). Euchromatin is more relaxed and composed of genes that are more

permissive to transcription, depending on the recruitment of transcription factors (TFs), cofactors and

RNA polymerases [1, 4]. In plants, DNA methylation occurs in three different cytosine contexts: CG,

CHG and CHH (where H = A, T or C), involving specialized DNA methyltransferases [5]. In *Arabidopsis*

thaliana, DOMAINS REARRANGED METHYLTRANSFERASE 2 (DRM2) and DRM1 mediate de novo DNA

methylation in all sequence contexts through the RNA-directed DNA methylation (RdDM) pathway,

which involves among other components, RNA-DEPENDENT RNA POLYMERASE 2 (RDR2) and DICER-

LIKE 3 (DCL3) for the production of short interfering (si)RNAs [6, 7]. The maintenance of CG

methylation is specifically performed by METHYLTRANSFERASE 1 (MET1), while CHROMOMETHYLASE

2 (CMT2) and CMT3 are involved in the maintenance at CHG sites [8, 9]. CMT2 can also be involved in

the deposition of CHH methylation at specific genomic location [10, 11]. Finally, DRM2 is mostly

required for the maintenance of CHH methylation through the RdDM pathway [6, 7, 9]. Together with

DNA methylation, additional pathways play important roles in TE silencing. The MICRORCHIDIA 1

(MORC1) and MORC6 ATPases interact together, and are required for heterochromatin condensation

and repression of TEs, acting mostly downstream of DNA methylation and RdDM pathway [12-14].

More recently, the *A. thaliana* plant mobile domain (PMD) proteins MAINTENANCE OF

MERISTEM (MAIN) and MAIN-LIKE 1 (MAIL1) were identified as new factors required for TE silencing

[15]. In addition, these two proteins have been involved in genome stability, and regulation of

developmental processes such as cell division and differentiation [16, 17]. The PMD is a large protein

domain of unknown function that is widely represented among the angiosperms, predominantly

associated with TEs [15, 18]. It has been proposed that genic PMD versions, such as the MAIN and

MAIL1 proteins derived from TEs after gene domestication [15, 18, 19]. Previous studies suggested

that genic PMDs could act as cellular factors related to transcription, possibly acting as transcription

factor (TF)-like, co-factor or repressor proteins regulating this cellular process [16, 18]. Nevertheless,

the role of PMD proteins in the regulation of transcription remains elusive. Most of genic PMD

proteins are standalone versions, however, in some cases, the PMD is fused to another protein

domain, such as protease, kinase or metallo-phosphatase (MPP) domains. For instance in *A. thaliana*,

the MAIL3 protein carries a PMD, which is fused to a putative serine/threonine-specific

phosphoprotein phosphatase (PPP) domain phylogenetically related to the plant-specific protein

phosphatase 7 (PP7) [20]. PP7 is a calmodulin-binding PPP that has been related to cryptochrome

(CRY)-mediated blue-light signaling, and to the control of stomatal aperture [20-22]. PP7 is also

involved in the perception of red/far red light by controlling the phytochrome pathway [23, 24]. In

addition to PP7 and MAIL3 (also known as "long PP7"), the protein PP7-like (PP7L) belongs to the same

phylogenetic clade [20]. PP7L was recently identified as a nuclear protein involved in chloroplast

development and abiotic stress tolerance [25]. The *pp7l* mutant plants showed photosynthetic defects

and strong developmental phenotype associated with misregulation of several genes [25].

In this study, we described a forward genetic screen based on a GFP reporter gene that

allowed us to identify a mutant population in which *MAIN* is mutated, leading to GFP overexpression.

We then deciphered the genetic interaction between the DRM2, CMT3 and MAIN, showing that these

proteins are part of different epigenetic pathways that act redundantly or synergistically to repress

TEs. Biochemical analyses indicated that MAIN and MAIL1 physically interact together. These analyses

also identified PP7L as a robust interactor of the two PMD proteins. In addition, the characterization

of developmental and molecular phenotypes of *pmd* and *pp7l* single and double mutant plants

strongly suggest that these proteins interact together to silence TEs, and regulate the expression of a

common set of genes. Finally, phylogenetic analyses allowed us to determine the distribution of PMD

and PP7/PP7L domains among the Eudicots. Based on these analyses, we have evidences of co-

evolution linked to the neo-association of the PMD and PP7-type PPP domains on single proteins in

several Eudicot species, suggesting a convergent evolution between these two protein domains.

RESULTS

Mutation in *MAIN* is responsible for TE silencing defects.

The *ATCOPIA28* retrotransposon *AT3TE51900* (hereafter called *ATCOPIA28*) is targeted by

several epigenetic pathways such as DNA methylation and the MORC1/6 complex, which altogether

contribute to its repression. We engineered a construct in which the 5' long terminal repeat (LTR)

promoter region of *ATCOPIA28* controls GFP transcription (Fig 1A). While the *ATCOPIA28::GFP*

transgene is fully silenced in wild type (WT) plants, it is weakly expressed in the DNA methylation-

deficient *drm1 drm2 cmt3 (ddc)* triple mutant background (Fig 1B) [26]. We performed an ethyl

methane sulfonate (EMS) mutagenesis using the *ATCOPIA28::GFP ddc* plants as sensitized genetic

material, and screened for mutant populations showing GFP overexpression. Among, the selected

populations, we retrieved two new mutant alleles of *MORC6* carrying missense mutations in either

the GHKL or S5 domains of the protein (S1A-C Fig). We also identified the population *ddc #16* showing

strong overexpression of GFP and misregulation of several endogenous TEs, including *ATCOPIA28* (Fig

1B-D). Mapping experiments based on whole genome resequencing and bulk segregant analysis

indicated that *ddc #16* carries a missense point mutation (C230Y) in the gene *AT1G17930*, previously

named *MAIN* (S1D and S1E Fig). Genetic complementation analyses by crossing the *ddc #16* EMS

mutant with the knock-out (KO) transferred DNA (T-DNA) insertion line *main*-2 generated F1 *ddc #16*

x *main-2* plants that did not express the GFP (S1F Fig). Transcriptional profiling analyses showed,

however, that endogenous TEs, including *ATCOPIA28*, were upregulated in F1 *ddc #16* x *main-2* plants,

but not in F1 control plants generated from the backcross of *ddc #16* with WT Columbia (Col) plants

(S1G Fig). Self-fertilization of F1 *ddc #16* x *main-2* plants allowed us to retrieve several F2 *ddc #16* x

main-2 plants overexpressing the GFP (S1F Fig). Among these GFP positive F2 plants, we identified

individuals that were either homozygote for the EMS mutation in the *MAIN* gene, or plants carrying

both the EMS and T-DNA *main-2* mutant alleles (S1F Fig). Moreover, while all these plants were

homozygote for the *drm2* mutation, half of them segregated the *cmt3* mutation. Thus, altogether,

these analyses suggested that *ATCOPIA28::GFP* silencing is more DRM2- than CMT3-dependent. More

importantly, they confirmed that *MAIN* was the mutated gene causing the upregulation of

ATCOPIA28::GFP and several endogenous TEs. Therefore, *ddc #16* was renamed *ddc main-3*.

MAIN, DRM2 and CMT3 act redundantly or synergistically to repress TEs and DNA-methylated

genes.

To determine the genetic interaction of *ddc* and *main-3* mutations on TE silencing, we carried

out two independent RNA sequencing (RNA-seq) analyses in the hypomorphic *main-3* single, *ddc* triple

and *ddc main-3* quadruple mutant plants (Fig 2A). We observed some variations among the

misregulated loci between the two RNA-seq analyses (S2A-B Fig and S1 Table). Therefore, we decided

to focus our analyses on loci that were reproducibly misregulated in each mutant background (S2C Fig

and S2 Table). As previously described, the *ddc* mutant showed upregulation of several TEs spread

over the five chromosomes (Fig 2B-D and S2D Fig, and S2 Table) [11]. Loss of TE silencing was also

observed to a milder degree in the *main-3* mutant, with the significant enrichment of pericentromeric

TEs among the upregulates TEs (Fig 2B-D and S2D Fig, and S2 Table). The *ddc main-3* mutant showed

an exacerbation of TE silencing defects, with a large number of pericentromeric TEs being specifically

upregulated in this mutant background (Fig 2B-D and S2 Table). Comparative analyses revealed that

upregulated TEs cluster into four distinct classes (Fig 2E and S2E Fig). Class I TEs are upregulated in

both *ddc* and *main-3* mutants, and combining these mutations in *ddc main*-3 lead to synergistic effects

(Fig 2E and S2E-F Fig). Class II and class III TEs are targeted by the MAIN and DRM2/CMT3 pathways,

respectively (Fig 2E and S2E-F Fig). However, the upregulation of class II and class III TEs is further

enhanced in *ddc main-3*, which suggests that the MAIN and DRM2/CMT3 pathways can partially

compensate each other at these genomic locations (S2F Fig). Finally, the most abundant class IV TEs

are only misregulated in *ddc main-3*, which implies that the MAIN and DRM2/CMT3 pathways act

redundantly to silence these TEs (Fig 2E and S2E-F Fig).

Several genes were reproducibly misregulated in the three mutant backgrounds (S2 Table).

Among these genes, a subset was commonly upregulated in *ddc*, *main-3* and *ddc main-3* (S2G Fig).

Remarkably, genes that were upregulated in the three mutants were significantly enriched in

pericentromeric regions of chromosomes, where constitutive heterochromatin resides (S2G Fig). This

is consistent with the fact that, among the upregulated genes, we identified a large proportion of DNA-

methylated genes (S2G Fig). Conversely, we could not identify a subset of genes commonly

downregulated in *ddc, main-3* and *ddc main-3* (S2G Fig). Furthermore, downregulated genes were not

enriched in pericentromeric regions, and most of them were not targeted by DNA methylation (S2G

Fig).

To further dissect the genetic interaction between the DRM2, CMT3 and MAIN pathways, we

generated the *drm1 drm2 main-3 (dd main-3)* and *cmt3 main-3* mutants (S2H Fig). We then analyzed

the expression level of several TEs previously identified as misregulated in *ddc*, *main-3* and/or *ddc*

main-3. The endogenous *ATCOPIA28* was the most expressed in *ddc main-3* and *dd main-3*, and to a

lesser extent, in *cmt3 main-3* (Fig 2F). This is consistent with the fact that all the F2 *ddc #16* x *main-2*

plants overexpressing *ATCOPIA28::GFP* were *drm2* homozygote, although they segregated the *cmt3*

mutation (S1F Fig). Further analyses showed that most of the tested TEs tend to be more expressed

in *cmt3 main-3* than in *dd main-3*, with the exception of *ATIS112A* that was more upregulated in *dd*

main-3 than in *cmt3 main-3* (Fig 2G). In conclusion, these analyses showed complex genetic

interactions between the DRM2, CMT3 and MAIN pathways. Depending on the locus, the three

pathways act redundantly or synergistically to repress TEs and DNA-methylated genes.

MAIN and MAIL1 are required for the proper expression of a common set of genes and TEs.

To study the role of MAIN and MAIL1 in the regulation of gene expression and TE silencing,

we performed two independent RNA-seq experiments in the *main-2* and *mail1-1* null mutants (Exp1

and Exp3, S3 Table), and reanalyzed previously published RNA-seq datasets (Exp2, S3 Table) [15].

Although all the RNA-seq experiments were performed using plants at the same developmental stage

(3-week-old seedlings), we observed important variations among the number of genes and TEs that

were misregulated in the two *pmd* null mutants in each RNA-seq experiment (S3A Fig and S3 Table).

Variations among WT replicates from the three independent RNA-seq experiments were not as

pronounced, and principal component analysis (PCA) showed that all the WT samples tend to cluster

together (S3B Fig). Conversely, the *pmd* mutant samples did not cluster together based on their

genetic background, but rather based on the RNA-seq experiment (S3B Fig). This suggests that *main-*

2 and *mail1-1* perceive their environment in a similar way, leading to the misregulation of similar sets

of loci in each independent RNA-seq experiment. Furthermore, this is consistent with the hypothesis

that MAIN and MAIL1 act in the same pathway [15]. Comparative analyses between the three

independent RNA-seq experiments allowed us to identify genes and TEs that were reproducibly

misregulated in *main-2* and *mail1-1* (S3C Fig). For stringency purpose, we decided to focus on these

lists of reproducibly misregulated loci to perform further analyses (Fig 3A-B and S4 Table).

We then compared the transcriptomes of *main-2* and *mail1-1* null mutants, together with the

hypomorphic *main-3* mutant allele (S2 and S4 Tables). We identified sets of genes and TEs that were

specifically misregulated in each mutant background (Fig 3C and D). These analyses also revealed

some loci that were commonly misregulated in the three mutants (Fig 3C-D, S3D-F Fig and S5 Table).

The biggest overlaps were among the downregulated genes, as most of the downregulated genes in

main-2 were also downregulated in *mail1-1* and, to a smaller proportion, in *main-3* (Fig 3D). As

observed in *main-3* (S2G Fig), the genes that were downregulated in *main-2* and *mail1-1* were mostly

located in the chromosome arms, and they were not DNA-methylated genes (Fig 3E). However, unlike

in *main-3*, the upregulated genes in *main-2* and *mail1-1* were not enriched in pericentromeric regions,

and only one quarter of them were DNA-methylated genes (Fig 3E). This discrepancy suggests that

MAIN and MAIL1 are not only required for the silencing of pericentromeric and DNA-methylated loci,

but also for the repression of genes that are not targeted by DNA methylation. In conclusion, these

comparative analyses allowed us to precisely define the sets of genes and TEs that are commonly

misregulated in *main-2, mail1-1* and *main-3* mutants. Moreover, these results revealed important

overlaps between the misregulated loci in *main-2* and *mail1-1,* which strongly suggests that the two

proteins act in the same pathway to regulate the expression of common sets of loci.

Slight increase in non-CG methylation in the *main-2* mutant does not correlate with changes in gene

expression and TE silencing defect.

Whole genome bisulfite sequencing (BS-seq) analyses showed that, at the chromosome scale,

DNA methylation level is mostly unchanged in *main-2* in comparison to WT, with the exception of a

slight increase in CHG methylation in pericentromeric regions (Fig 4A). Subtle but statistically

significant CHG hypermethylation was further confirmed in pericentromeric TEs and genes, which are

mostly TE genes (Fig 4B and C). Slight CHG and CHH hypermethylation was also detected in TEs located

in chromosome arms (Fig 4D). Conversely, genes located in chromosome arms did not show

significant changes in DNA methylation level in *main-2* (Fig 4E). Identical results were obtained by

analyzing the DNA methylation level at TEs and genes that were specifically misregulated in *main-2*

(Fig 4F-H). We then analyzed the DNA methylation level at genomic locations previously defined as

differentially hypomethylated regions (hypo DMRs) at CHG and CHH sites in *cmt3* and *drm1drm2 (dd)*

mutants, respectively [26]. The *cmt3* and *dd* hypo DMRs are mostly located in TEs. As observed with

pericentromeric genes and all TEs (Fig 4B-D), we found slight increases in CHG and CHH methylation

at *cmt3* and *dd* hypo DMRs, respectively, in *main-2* (S4A and S4B Fig). Finally, DMR calling in *main-2*

using stringent parameters only identified a few DMRs (S4C Fig). Thus, DNA methylation is mostly

unaffected in *main-*2, with the exception of a slight increase in non-CG methylation at pericentromeric

genes and all TEs. Moreover, this subtle non-CG hypermethylation does not correlated with changes

in gene and TE expression observed in *main-2*.

MAIN, MAIL1 and the metallo-phosphatase PP7L physically interact together.

The *main-2* and *mail1-1* null mutants display similar molecular and developmental

phenotypes (Fig 3 and Fig 5A). Thus, we hypothesized that the two PMD proteins may act in the same

pathway, possibly by interacting together. To test this hypothesis, we generated transgenic lines

expressing FLAG- and MYC-tagged genomic PMD versions driven by their endogenous promoters. We

confirmed that epitope-tagged MAIN and MAIL1 proteins were produced at the expected sizes, and

they could complement the respective developmental phenotypes of null mutant plants (Fig 5A and

B). Importantly, they could also efficiently rescue the TE silencing and gene expression defects

observed in the two *pmd* mutants, implying that epitope-tagged MAIN and MAIL1 are functional

proteins (Fig 5C-E). Using FLAG-tagged MAIN and MAIL1 expressing plants, immunoprecipitation

followed by mass spectrometry (IP-MS) analyses were carried out to determine potential protein

interactors. Mass spectrometry (MS) analyses indicated that MAIL1 was strongly immunoprecipitated

with MAIN-FLAG and *vice versa* (Fig 5F). To validate IP-MS results, we crossed the MAIN-FLAG and

MAIL1-MYC lines together. We then performed co-immunoprecipitation (co-IP) experiments using F1

hybrid plants co-expressing the two transgenes, and confirmed that MAIN and MAIL1 interact

together (Fig 5G). MS analyses of MAIN-FLAG and MAIL1-FLAG IP also identified the metallo-

phosphatase PP7L as putative interactor (Fig 5F). MAIN, MAIL1 and PP7L were the only three proteins

reproducibly enriched across multiple replicates (Fig 5F). Co-IP experiments using plants co-expressing

either PP7L-FLAG together with MAIN-MYC or MAIL1-MYC constructs confirmed the interaction

between PP7L and each PMD protein (Fig 5H and I). Thus, the three proteins MAIN, MAIL1 and PP7L

physically interact together.

The *main, mail1* and *pp7l* mutants display similar developmental and molecular phenotypes.

PP7L is a putative metallo-phosphatase that was recently identified as a nuclear protein

required for photosynthesis [20, 25]. The *pp7l-2* null mutant displays abnormal developmental

phenotype reminiscent of the two *pmd* null mutant plants, and 3-week-old *mail1-1 pp7l-2* double

mutant plants do not show exacerbation of this phenotype (Fig 6A). To determine the genetic

interaction between PMD and PP7L, we compared the transcriptomes of *main-2*, *mail1-1, pp7l-2*

single and *mail1-1 pp7l-2* double mutants (RNA-seq Exp3, S3 and S6 Tables). As previously described

for *main-2* and *mail1-1* (S3B Fig), PCA analyses showed that all the mutant samples tend to cluster

together, and away from WT controls (S5A Fig). We identified a large number of misregulated loci in

each mutant background (S5B Fig). Comparative analyses revealed that significant proportions of

these loci were commonly misregulated in all the mutant backgrounds, which is consistent with the

fact that MAIN, MAIL1 and PP7L interact together to possibly regulate gene expression (Fig 6B and C).

These analyses also identified loci that were specifically misregulated in *main-2*, *mail1-1* or *pp7l-2*,

which suggests that each protein is independently required for the proper expression of subsets of

loci (Fig 6C). Besides, these analyses revealed loci that were exclusively misregulated in the *mail1-1*

pp7l-2 double mutant, which implies that PP7L and MAIL1 may act redundantly to ensure the proper

expression of these loci (Fig 6B-C). Further analyses showed that, among the loci that were

misregulated in *mail1-1 pp7l-2*, upregulated genes were significantly more expressed in the double

mutant than in each single mutant, and upregulated TEs were significantly differentially expressed

only between *mail1-1 pp7l-2* and *pp7l-2* mutants (Fig 6D-E). Conversely, there was no significant

difference of expression between the double mutant and single mutants for the downregulated genes

(Fig 6F). Thus, these analyses suggest that combining the *pp7l-2* and *mail1-1* mutations may lead to

synergistic defects mostly at genes that are upregulated in the double mutant. Considering the

variations observed among the misregulated loci in *main-2* and *mail1-1* between independent RNA-

seq experiments, we then decided to focus our analyses on loci previously defined as reproducibly

misregulated in the two *pmd* mutants (Fig 3, S3 Fig and S4 Table). Among these loci, we identified

several genes and TEs that were also misregulated in *pp7l-2* and *mail1-1 pp7l-2*, with the most

significant overlap among downregulated genes (Fig 6G-J and S7 Table). These commonly

misregulated loci did not show any significant difference of expression between the double and single

mutants (Fig6K-M). This suggests that MAIN, MAIL1 and PP7L are equally required for the proper

expression of these subsets of loci (S7 Table). Finally, we performed in silico analyses to identify

enriched DNA motif within a 1kb promoter region upstream of start codon of genes that were up- or

downregulated in the different mutant backgrounds. We could not detect any enrichment of a DNA

motif among any lists of upregulated genes. Likewise, we could not identify a DNA motif significantly

enriched in the lists of downregulated genes from the RNA-seq Exp3. However, we identified a

discrete DNA motif (hereafter called 'DOWN' motif) that was enriched in the promoters of genes that

were reproducibly downregulated in *main-3*, *ddc main-3*, *main-2* and *mail1-1* mutants (S5C Fig). The

'DOWN' motif was further enriched and refined in the promoters of genes commonly downregulated

in all the mutant backgrounds analyzed in this study: eighteen genes out of nineteen, 95% of

enrichment (S5C Fig, S8 and S9 Tables). We analyzed the DNA methylation level of the 'DOWN' motif

in the promoters of these eighteen genes in WT and *main-2*, and found that this DNA motif was not

targeted by DNA methylation. Besides, further analyses showed that only a small fraction of all

Arabidopsis genes carried the 'DOWN' motif in their promoter (11%, S5C Fig), which is in the range of

genomic distribution of several DNA motifs recognized by bZIP TFs (S10 Table). Finally, random test

analyses based on nineteen randomly picked genes strongly suggested that the enrichment of the

'DOWN' motif in the promoter of downregulated genes was substantial (S9 Table). Thus, altogether,

these analyses showed that MAIN, MAIL1 and PP7L are equally required for the proper expression of

a common set of genes that are downregulated in each single mutant as well as in *mail1-1 pp7l-2*

double mutant, and these genes carry the 'DOWN' DNA motif in their promoter.

PP7L is not required for heterochromatin condensation.

WT *Arabidopsis* nuclei at interphase exhibit condensed DNA foci called chromocenters that

are composed of constitutive heterochromatin, and are enriched in H3K9me2 [27]. In several

epigenetic mutants, decondensation of constitutive heterochromatin correlates with disruption of

chromocenters, and loss or diffusion of H3K9me2 in the nucleoplasm [27]. Thus, analyzing H3K9me2

subnuclear distribution by immunofluorescence (IF) experiments has been reproducibly used as a

cytological approach to assay for heterochromatin decondensation [12, 27, 28]. A previous study

showed that subnuclear distributions of chromocenters and H3K9me2 were unchanged in *main-2* and

mail1-1 mutants [15]. However, fluorescent in situ hybridization (FISH) experiments using a DNA

probe for the 106B pericentromeric repeats suggested that heterochromatin was decondensed in the

two *pmd* mutants in comparison to WT plants [15]. We performed IF experiments to analyze the

subnuclear distribution of H3K9me2 in the *pp7l-2* mutant. These analyses did not show any change in

the condensation level of chromocenters in *pp7l-2* nuclei in comparison to WT (Fig 7). Instead, we

observed that *pp7l-2* nuclei were proportionally more condensed than WT nuclei (Fig 7). This is likely

due to the fact that *pp7l-2* mutant displays abnormal phenotype and growth delay in comparison to

WT plants that are entering the floral transition stage, a developmental stage where partial

chromocenter decondensation has been documented [29]. In conclusion, based on the H3K9me2 IF

experiments, we can conclude that *pp7l-2* is not impaired in chromocenter condensation.

The PMD and PP7 domains have co-evolved among the Eudicots.

Among the Angiosperms, most of the genic PMDs, like MAIN and MAIL1, are standalone

versions [18]. However, some genic PMDs can associate with other protein domains, such as for

instance a PPP domain. In *A. thaliana*, the protein MAIL3, which carries a PMD fused to a PPP domain,

is a close homolog of both MAIN/MAIL1 and PP7/PP7L through its PMD and PPP domains, respectively.

Considering that the PMD proteins MAIN and MAIL1 interact with PP7L, and are required for the

expression of similar set of loci, we decided to determine the distribution of related genic PMD and

PPP domains, and to retrace their evolutionary history among plant species. The *A. thaliana* MAIN,

MAIL1 and MAIL3 are all members of the PMD-C family that also includes MAIL2 [15]. Since our

objective is to retrace the evolution of genic (and not TE-containing) PMD-C, we have decided to

restraint our search to Eudicots. Indeed, Eudicot species contain mainly genic PMD-C, while other

angiosperms may contain variable numbers of closely related genic and TE-associated PMD-C motifs

that would be difficult to distinguish in our analysis. To retrace the evolution history of the genic PMD-

C family, we used *A. thaliana* PMD-C genes to search and collect their relatives (paralogues and

orthologues) in 30 genomes representative of the Eudicot diversity (see S11 Table for a list of species

and their corresponding codes used in Fig 8, and S12 Table for motif sequences).

In our phylogenetic analysis, the genic PMD-C family can be clearly separated in two major

clades. The first clade is composed of orthologues of *A. thaliana* MAIL2, MAIL1 and MAIN, while the

second one includes orthologues of *A. thaliana* MAIL3 (Fig 8A). MAIL2 orthologues were found in all

species tested, forming a closely related group, which suggests that they are under strong purifying

selection (see the very short branch lengths linking most MAIL2 genes in Fig 8A). In several species,

additional MAIL2 paralogues were also detected. They were either imbedded in the major MAIL2

group, or forming independent and more divergent subgroups, like in the case of MAIL1 and MAIN

that are Brassicaceae-specific MAIL2 paralogues. By comparison, MAIL3 orthologues were not found

in all Eudicot species tested, and, except in Brassicaceae, *MAIL3* genes appear to be under much

weaker purifying selection compare to *MAIL2* and *MAIL2-like* genes (see the longer branch lengths in

the tree of Fig 8A). Brassicaceae *MAIL3* genes contrast with other *MAIL3*, by forming a closely related

group in the phylogenetic tree. This suggests a clear change in selection pressure, typical of a

neofunctionalization event that could correlate with the acquisition of the PPP motif by these genes

(Fig 8B and see below). Remarkably, another fusion event between PMD-C and PPP motifs occurred

independently in grapevine, but this time involving a MAIL2 paralogue (VvMAIL2.2, Fig 8A).

We then used the PPP motif found in *A. thaliana* MAIL3, to collect orthologous genes and

retrace the evolution history of this motif in the same Eudicot species used above. We confirmed that

these genes can be clearly separated in two distinct clades: PP7 and PP7-like (PP7L) (Fig 8B). All tested

species present one or several closely related PP7 paralogues. Although the Brassicaceae MAIL3 PPP

motif belongs to the PP7 clade, it diverged significantly compared to other standalone PP7 paralogues

(Fig 8B). Same observation was made regarding the PP7 domain of VvMAIL2.2. Thus, as described for

the PMD of Brassicaceae MAIL3 and grapevine VvMAIL2.2, this suggests a fast-evolving period and

neofunctionalization of the PP7 domain in these species, subsequently to the PMD-C/PP7 fusion.

Conversely, PP7L orthologues were not found in all species tested and, accordingly, these genes are

under weaker purifying selection compare to genes belonging to the PP7 subfamily. In conclusion,

phylogenetic analyses showed that, in at least Brassicaceae and grapevine, neo-association of PMD-C

and PP7 domains have potentially create new protein functions that were maintained through

evolution.

DISCUSSION

In *A. thaliana,* MAIN and MAIL1 are standalone PMD proteins that have been involved in

genome integrity, regulation of cell division and differentiation, and silencing of TEs [15-17]. In this

study, we show that TE silencing is widely impaired in the *ddc main-3* higher order mutant, which is

both partially defective in DNA methylation and MAIN activity. We also identify the putative

phosphatase protein PP7L as MAIN and MAIL1 protein interactor, and show that among the loci that

are commonly misregulated in *pmd* and *pp7l* single and double mutants, most of the downregulated

genes carry the 'DOWN' DNA motif in their promoter. Finally, phylogenetic analyses among Eudicots

suggest a mechanism of neofunctionalization between the PMD and PP7-type PPP, to potentially

acquire a functional module that requires the two protein domains.

The PMD MAIN protein acts independently of DRM2- and CMT3 pathways to silence TEs.

Previous analyses showed that some TEs were synergistically upregulated in the *mail1 rdr2*

double mutant plants, suggesting that MAIL1 acts independently of RdDM pathway [15]. In our whole

genome transcriptomic analyses, we show that several TEs are upregulated in both *main-3* and *ddc*

mutants, and their upregulation is dramatically exacerbated in the *ddc main-3* quadruple mutant (Fig

2 and S2 Fig). We also identify TEs that are upregulated in either *ddc* or *main-3* mutants, but display

stronger misregulation in the *ddc main-3* higher order mutant (Fig 2 and S2 Fig). Finally, we identify a

large class of TEs that are only upregulated in *ddc main-3* (Fig 2 and S2 Fig). Altogether, these analyses

reveal complex genetic interaction between the MAIN, DRM2 and CMT3 proteins to silence TE.

Depending on the locus, these proteins act redundantly or synergistically to efficiently silence TE.

Previous work showed that DNA methylation is not impaired in *mail1-1* [15]. We found that DNA

methylation is mostly unaffected in the *main-2* null mutant. However, we detected a mild but

significant hypermethylation at non-CG sites in TEs and pericentromeric genes (Fig 4). One hypothesis

is that CHG and CHH hypermethylation observed in *main-2* is a backup mechanism to compensate for

MAIN loss of function, and to dampen TE silencing defects. Although further studies will be required

to test this hypothesis, it is consistent with the fact that combining the *main-3* and *ddc* mutations

leads to an exacerbation of TE silencing defects. Synergistic effects between different epigenetic

pathways have already been described. For instance, it has been shown that MORPHEUS MOLECULE

1 (MOM1) and MORC1/MORC6 proteins, or MOM1 and the RdDM pathway act synergistically to

efficiently silence TEs [13, 30]. Altogether, these observations contribute to the "mille-feuille" (i.e.

"multiple layers") model, in which different epigenetic pathways converge towards the silencing of

TEs [31].

The putative phosphatase PP7L interacts with the PMD MAIN and MAIL1 protein to regulate a

similar set of genes and TEs.

Recently, the putative phosphoprotein phosphatase PP7L was involved in the biogenesis of

chloroplasts and plant response upon abiotic stress [25]. Here, we show that PP7L interact with MAIN

and MAIL1, and *main-2*, *pp7l-2, mail1-1* single and *mail1-1 pp7l-2* double mutant plants display similar

developmental and molecular phenotypes (Fig 5 and 6). We also show that, as described for *main-2*

and *mail1-1* [15], the subnuclear distribution of chromocenters and H3K9me2 are unaltered in *pp7l-2*

(Fig 7). The 106B pericentromeric repeats appeared decondensed in the two *pmd* mutants [15], future

work will determine if similar phenotype is observed in *pp7l-2*. Although MAIN, MAIL1 and PP7L

interact together, we cannot exclude that an additional protein is required for the interaction. In

addition, PP7L may have additional partners independently of MAIN and MAIL1. Further biochemical

studies such as IP-MS analyses using the FLAG-tagged PP7L line will contribute to addressing these

points.

Transcriptomic analyses revealed complex genetic interaction between MAIN, MAIL1 and PP7L;

the three proteins acting either independently or together to ensure the proper expression of genes,

and to perform TE silencing. Moreover, transcriptome profiling of *mail1-1 pp7l-2* double mutant

revealed that the two mutations may have synergistic effects, specifically at genes that are

upregulated in the mutant. To further study the genetic interaction between the three proteins, it will

be important to analyze the transcriptome of *main-2 mail1-1 pp7l-2* triple mutant. Altogether and

considering that *i)* MAIN, DRM2 and CMT3 pathways genetically interact together, and *ii)* the *main-2*

mutant show a slight increase in DNA methylation at CHG and CHH sites, we cannot rule out that MAIN

is playing a dual role: regulating gene expression through its interaction with MAIL1 and PP7L, and

involved in TE silencing through its genetic interaction with DNA methylation. In the future, it will be

important to analyze DNA methylation in *pp7l-2*, but also in *pmd pp7l-2* higher order mutants. In

parallel, studying the *ddc pp7l-2* mutant will allow to further decipher the genetic interaction between

the PP7L and DNA methylation pathways.

Genes that are commonly downregulated in *main, mail1* and *pp7l* mutants carry the 'DOWN' motif

in their promoters.

Genes that are commonly downregulated in the different *pmd* and *pp7l* mutant backgrounds

show a bigger overlap than other misregulated loci (Fig 6, and S8 Table). Furthermore, eighteen out

of nineteen genes commonly downregulated in the all the mutant backgrounds carry the 'DOWN' DNA

motif in their promoter (S5 Fig). The 'DOWN' motif is also found in the promoter of genes reproducibly

downregulated in *main-2*, *mail1-1*, *main-3* and *ddc main-3*. However, it is not significantly enriched in

the promoter of genes identified as downregulated in the *pmd* and *pp7l* mutants from the RNA-seq

Exp3. One explanation for this discrepancy is that too many loci were identified as downregulated in

each dataset of the RNA-seq Exp3, which created a dilution of the loci carrying the 'DOWN' motif in

their promoter. In the future, further RNA-seq experiments in *pp7l-2* and *mail1-1 pp7l-2* will precise

the lists of reproducibly misregulated loci in these mutant backgrounds*.*

Based on our results, we hypothesize that the 'DOWN' motif may act as a putative cis-

regulatory element (CRE) recognized by an unidentified TF, which would be required for the

transcription of genes identified as downregulated in *pmd* and *pp7l* mutants. This unknown TF could

be recruited or activated by the MAIN/MAIL1/PP7L protein complex. Another hypothesis is that the

'DOWN' motif is directly recognized by the MAIN/MAIL1/PP7L protein complex. Further study will be

required to test if MAIN/MAIL1/PP7L protein complex interact with chromatin, and bind the 'DOWN'

motif. In parallel, further biochemical analyses may allow to identify an uncharacterized putative TF

as MAIN/MAIL1/PP7L protein interactor.

The association of PMD-C and PP7/PP7L domains creates a functional protein module.

In this study, we identified PP7L has a protein partner of the two standalone PMDs MAIN and

MAIL1, and showed that these proteins are required for the proper expression of a common set of

genes, and for TE silencing. Besides, we showed that the Brassicaceae MAIL3 and the grapevine

VvMAIL2.2 proteins carry a PMD fused to a PP7 domain. Based on these results, we hypothesize that

depending on the configuration, the association of PMD-C and PP7/PP7L domains would create a

functional protein module in trans or in cis. It is likely that the cis-association of PMD and PP7 found

in the Brassicaceae MAIL3 proteins occurred in the common ancestors of this Eudicot lineage, possibly

through the process of gene duplication. Since then, the MAIL3 PMD/PP7 fusion was maintained

under strong purifying selection, arguing for a neofunctionalization of the fusion protein. It is likely

that a similar process happened in grapevine, and possibly, in closely related Vitaceae species. To

some extent, the two distinct events that occurred in Brassicaceae and grapevine are reminiscent of

convergent evolution processes leading to the production of a functional PMD/PP7 module.

The occurrence of PMD and PP7/PP7L protein fusion in several Brassicaceae and grapevine is

reminiscent of the concept of Rosetta stone chimera proteins, which describes that two proteins

interacting together in one organism can be found fused together in another species to facilitate

enzymatic activity [32]. There are several examples of Rosetta stone proteins, described for instance

with different subunits of DNA topoisomerase or RNA polymerase [32]. Here, we show that, at least

in *A. thaliana,* the Rosetta stone chimera MAIL3 coexist with its close homologs MAIN/MAIL1 and PP7L

that interact together. The fact that the PMD and PP7 domains are fused together in MAIL3 may be a

strategy to optimize protein activity. Conversely, the enzymatic activity of the MAIN/MAIL1/PP7L

protein complex could be further regulated by allowing, or not, the three proteins to interact together.

Nevertheless, in both scenarios, it is likely that PMD and PP7/PP7L association creates a functional

protein module, which might be specialized in distinct biological processes depending on its

composition. Thus, we hypothesize that the MAIL3 and MAIN/MAIL1/PP7L protein complexes play

different role in the plant. This is consistent with the fact that, unlike *main-2*, *mail1-1* and *pp7l-2*

mutant, the *mail3-2* mutant does not show abnormal developmental phenotype [17]. Further studies

will be required to describe the role of MAIL3 in the plants.

In conclusion, we show here that the two *A. thaliana* PMD MAIN and MAIL1 proteins interact with

PP7L, and are involved in the regulation of a common set of genes and TEs. In addition, we show that

distinct events of PMD-C and PP7 fusions have occurred among the Eudicots (among several

Brassicaceae species and in grapevine), suggesting some convergent evolution processes and a

potential neofunctionalization of PMD/PP7 module in cis. The biological significance of PMD/PP7

fusion proteins will be investigated in the future by studying the role of MAIL3 in *A. thaliana*. In

addition, it will be important to determine whether the PMD proteins play important roles in other

plant species with agronomic value.

MATERIALS AND METHODS

Plant material and growing conditions. Wild-type (WT) and all mutant lines are in the Columbia (Col) ecotype. The *drm1-2* (SALK_031705), *drm2-2* (SALK_150863), *cmt3-11* (SALK_148381), *ddc* triple, *main-2* (GK-728H05), *mail1-1* (GK-840E05) and *pp7l-2* (SALK_003071) null mutant lines were previously described [15-17, 25, 26], and obtained from The Nottingham Arabidopsis Stock Centre. The *mail1-1 pp7l-2* double mutant was obtained by crossing the respective single mutants. T-DNA insertions were confirmed by PCR-based genotyping and RT-qPCR analyses. The *main-3* mutation was genotyped by derived cleaved amplified polymorphic sequences (dCAPS) using the restriction enzyme FokI. Primer sequences are described in S13 Table. All the WT Col and T-DNA mutant plants were grown on soil under a 16h-light/8h-dark cycle. When experiments required to screen for GFP expression under UV light, plants carrying the *ATCOPIA28::GFP* transgene were first grown on Murashige and Skoog (MS) plates under continuous light, 10-day old plants were then screened for GFP expression under UV light, and subsequently transferred onto soil. For *in vitro* plant culture, seeds were surface-sterilized and sowed on solid MS medium containing 0.5% sucrose (w/v).

Cloning of ATCOPIA28::GFP. The pCambia3300-NLS-GFP-T35S vector was previously described [12]. The 5'LTR promoter corresponding to a region of ~1 kb upstream of *ATCOPIA28 (AT3TE51900)* was PCR amplified from WT genomic DNA, and cloned into pCR2.1 TOPO vector (Invitrogen). Quikchange site-directed mutagenesis (Stratagene) was performed according to Manufacturer's instruction to create a polymorphism site (MfeI→NdeI) within the 5'LTR promoter, which was subsequently mobilized into pCambia3300 upstream of NLS-GFP-T35S sequence. *ddc* triple mutant plants were transformed with the AT*COPIA28::GFP* construct using the *Agrobacterium*-mediated floral dip method [33]. Transgenic plants showing GFP fluorescence were backcrossed with a WT plant to promote the silencing of *ATCOPIA28::GFP* in the F1 generation. F1 plants were self-crossed and their F2 progenies were screened for GFP fluorescence, and PCR-based genotyped to obtain *ATCOPIA28::GFP* WT and *ATCOPIA28::GFP ddc* plants. Primer sequences used for *ATCOPIA28::GFP* cloning and PCR genotyping are described in S13 Table.

EMS mutagenesis, GFP screening and mapping analyses. Five thousand seeds of *ATCOPIA28::GFP ddc*

were mutagenized in 0.26% EMS solution for 12 hours with rotation. Seeds were subsequently washed

with water and sown on soil. Fifteen hundred M2 populations were collected, and subsequently

screened for GFP fluorescence under UV light using a SMZ18 Nikon Fluorescence Stereomicroscope

coupled with the C-HGFI intensilight fluorescence filter. Pictures were taken using the DS Qi1MC digital

camera kit. Mapping and identification of the EMS mutation responsible for the phenotype were

performed by bulk segregant analysis coupled with deep genome re-sequencing as previously

described [12], with the following differences. Reads were mapped against the reference genome

(*Arabidopsis* TAIR10) and single nucleotide polymorphisms called in Geneious (Biomatters). Using R,

single nucleotide polymorphisms were filtered for EMS mutations (G:C→A:T) and zygosity called

based on the variant frequency provided by Geneious (≥80% homozygous mutation, ≥45%, and ≤55%

heterozygous mutation). Plots were then created by calculating the ratio of the number of

homozygous and heterozygous and mutations in a 500-kb window as previously described [34].

Cloning of epitope-tagged versions of PMD and PP7L proteins. *MAIN*, *MAIL1* and *PP7L* genomic

regions were PCR amplified and FLAG or Myc epitopes were added to the C-terminus of each protein

as previously described [12]. Each time, the amplified region includes a ~1Kb promoter sequence

upstream of the respective transcriptional start site. For the *MAIN* promoter, a MluI site was modified

to allow LR reaction without changing the sequence integrity of the gene. *main-2*, *mail1-1* and *pp7l-2*

mutant plants were transformed with the *MAIN-FLAG*, *MAIN-MYC*, *MAIL1-MYC* and *PP7L-*

FLAG constructs using the *Agrobacterium*-mediated floral dip method [33]. Primer sequences are

described in S13 Table.

IP and MS analysis. Ten grams of 3-week-old seedling tissue were ground in liquid nitrogen and

resuspended in 50mL ice-cold IP buffer [50mM Tris HCl pH 7.6, 150mM NaCl, 5mM MgCl₂, 0.1%

Nonidet P-40, 10% glycerol (v/v), 0.5mM DTT, 1x Protease Inhibitor Mixture (Roche)] and centrifuged

2 times for 15 min at 4°C at 15 350g. 400µL of M2 magnetic FLAG-beads (Sigma, M8823) were added

to the supernatants, and incubated for 90min rotating at 4°C. M2 magnetic FLAG-beads were washed

seven times in ice-cold IP buffer for 5 min rotating at 4°C, and immunoprecipitated proteins were

eluted 3 times with 150µL 3x-FLAG peptides (Sigma, F4799) for 25 min each at 25°C. The eluted protein

complexes were precipitated by trichloroacetic acid and subjected to MS analyses as previously

described [13]. Peptide and protein-level false discovery rates were calculated by the DTASelect

algorithm using the decoy database approach. Based on a peptide PSM level p-value filter of less than

0.01 and a requirement for at least two peptides per protein, the protein-level false discovery rate

was less than 1% for all proteins detected.

Co-IP and immunoblotting. 0.5 g of 3-week-old seedling tissue were ground in liquid nitrogen,

resuspended in 1.5mL ice-cold IP buffer [50mM Tris pH 7.6, 150mM NaCl, 5mM MgCl2, 0.1% Nonidet

P-40, 10% glycerol, 0.5 mM DTT, 1x Protease Inhibitor Mixture (Roche)], and centrifuged 2 times for

15 min at 4°C, 16 000g. 50µL M2 magnetic FLAG-beads (Sigma, M8823) were added to the

supernatants and incubated for 2 hour rotating at 4°C. Beads were washed 3 times in ice-cold IP buffer

for 10 min rotating at 4°C. Immunoprecipitated proteins were denatured in Laemmli buffer for 5min

at 95°C. 10µL of input and bead elution were run on 10% SDS-PAGE gels, and proteins were detected

by western blotting using either Anti-FLAG M2 monoclonal antibody-peroxidase conjugate (Sigma,

A8592) at a dilution of 1:10000, or c-Myc rat monoclonal antibody (Chromotek, 9E1-100) at a dilution

of 1:1000 followed by goat anti-rat IgG horseradish peroxidase (Abcam, ab205720) used at a dilution

of 1:20000 as secondary antibody. Western blots were developed using Substrat HRP Immobilon

Western (Merck Millipore, WBKLS0500).

RNA extraction. Total RNA was extracted from aerial parts of 3-week-old seedlings grown on soil using

either RNeasy Plant Mini Kit (Qiagen, 74904) or Monarch Total RNA Miniprep Kit (NEB, T2010)

according to the manufacturer's protocols.

RNA sequencing. RNA-seq libraries were generated from 1μg of input RNA using NEBNext Ultra II

Directional RNA Library Prep Kit for Illumina (NEB, E7490) according to the manufacturer's protocols.

Libraries were sequenced on an Illumina HiSeq 4000 or NextSeq 550 machines. Reads were trimmed

using Trimmomatic [35], and mapped to the *A. thaliana* genome (*Arabidopsi*s TAIR10 genome) using

HISAT2 [36]. The sequence alignment files were sorted by name and indexed using SAMtools [37].

Files were converted to BAM files and number of reads mapped onto a gene calculated using HTSeq-

count [38]. Differentially expressed genes were obtained with DESeq2 [39], using a log2 fold-change

≥ 2 (up-regulated genes) or ≤ -2 (down-regulated genes) with an adjusted *p*-value of 0,01. Re-analyses

of previously published RNA-seq datasets from *main-2* and *mail1-1* (PRJEB15202) [15] were

performed as described above.

RT-qPCR. 1 μg of input RNA was converted to cDNA using GoScript Reverse Transcriptase (Promega

A501C) according to the manufacturer's protocol. The final reaction was diluted 6 times with RNase

free water. RT-qPCR experiments were performed with 4μL of cDNA combined to the Takyon No Rox

SYBR MasterMix (Eurogentec, UF-NSMT-B0701), using a LightCycler 480 instrument (Roche).

Amplification conditions were as follows: 95°C 5 min; 45 cycles, 95°C 15s, 60°C 15s, 72°C 30s; melting

curves. RT-qPCR analyses used the $2^{-\Delta\Delta Ct}$ method. For each analysis, ΔCt was first calculated based on

the housekeeping *RHIP1* gene Ct value [40]. ΔΔCt were then obtained by subtracting the wt ΔCt from

the ΔCt of each sample. Values were represented on bar charts relative to WT. Three technical

replicates were performed per biological replicate, and 3 biological replicates were used in all

experiments, unless otherwise stated. Primer sequences are described in S13 Table.

DNA motif detection. The motifs for enhancer sequences (1kb upstream the TSS) were discovered

using MEME (Multiple Em for Motif Elicitation). MEME represents motifs as position-dependent

letter-probability matrices which describe the probability of each possible letter at each position in

the pattern [41].

Bisulfite sequencing. Genomic DNA was extracted from aerial parts of 3-week-old seedlings using

Quick-DNA Plant/Seed Miniprep Kit (Zymo research, D6020) according to the manufacturer's protocol.

Whole genome bisulfite sequencing (WGBS) library was prepared from 50 ng genomic DNA using

NuGen Ovation Ultralow Methyl-Seq kit. Bisulfite treatment was carried out by Qiagen Epitect bisulfite

kit. WGBS libraries were sequenced on an Illumina HiSeq 4000 machine. The raw reads (single end)

were trimmed using Trimmomatic in order to remove adapter sequences [35]. The remaining

sequences were aligned against the *A. thaliana* genome TAIR10 version using Bismark [42]. Duplicated

reads were collapsed into one read. For visualization of the data, we used ViewBS [43].

Sequence selection, multiple sequences alignments and phylogenetic reconstruction.

Blast searches (blastp) were performed starting from known *A. thaliana* PMD-C and PP7/PP7L motifs

on the thirty species representing the diversity of the Eudicot lineages. When necessary tblastn

searches were also used to obtain complete protein sequences. To build the phylogenetic trees, PMD-

C or PP7/PP7L motifs were aligned using the multiple sequence comparison by log-expectation

(MUSCLE v3.7) software [44]. Trees were reconstructed using the fast-maximum likelihood tree

estimation program PHYML [45] using the LG amino acids replacement matrix [46]. Statistical support

for the major clusters were obtained using the approximate likelihood-ratio test (aLRT) [47].

Immunofluorescence and DAPI-staining. Leaves from 3-week-old plants, were fixed for 20 min

rotating at 4°C in 2% formaldehyde in Tris buffer (10 mM Tris-HCl pH 7.5, 10 mM EDTA, 100 mM NaCl),

washed two times for 10 min rotating at 4°C in cold Tris buffer and subsequently chopped in LB01

buffer (15 mM Tris-HCl pH 7.5, 2 mM EDTA, 0.5 mM spermine, 80 mM KCl, 20mM NaCl and 0.1%

Triton- X-100). Nuclei were filtered through a 30 µm cell strainer cap (Sysmex, 04-0042-2316) and 5µl

of the nuclei solution was diluted in 10 µl of sorting buffer (100mM Tris-HCl pH 7.5, 50 mM KCl, 2 mM

MgCl2, 0.05% Tween-20 and 5% sucrose). 20µl of the nuclei dilution were spread onto a polylysine

slide and air-dried for 40 min. Slides were post-fixed in 2% formaldehyde in 1X PBS for 5 min and

washed 2 times with water. Slides were incubated 15 min in 1X PBS, 0.5% Triton X-100 at RT and

washed 3 times with 1X PBS for 5 min. For detection, slides were incubated over night with a mouse

anti-H3K9me2 monoclonal antibody (Abcam, Ab 1220) at 1:500 in 3% BSA, 0.05% Tween in 1X PBS at

4°C in a moist chamber. After 3 washes in 1X PBS for 5 min, slides were incubated 2h with a goat anti-

mouse antibody coupled to Alexa fluor 568 (Invitrogen, A11004) at 1:1000 in 3% BSA, 0.05% Tween in

1X PBS in a moist chamber. Slides were washed 1 time 5 min with 1X PBS, 1 time 10 min with 1X PBS,

1µg/mL DAPI, and 1 time 5 min with 1X PBS. DNA was counterstained with 1µg/mL DAPI in Vectashield

mounting medium (Vector Laboratories). Observation and imaging were performed using a LSM 700

epifluorescence microscope (Zeiss).

Data availability. Nucleotide sequencing data generated in this study have been deposited in

European Nucleotide Archive (ENA) under the accession number PRJEB33240

(http://www.ebi.ac.uk/ena/data/view/PRJEB33240). The proteomics data have been deposited to the

MassIVE data repository (https://massive.ucsd.edu) with the dataset identifier MSV000084089. All

other data and material are available within the manuscript and its supplementary files, or from the

corresponding author upon request.

ACKNOWLEDGMENTS

The authors want to thank Thierry Lagrange, Frederic Pontvianne and other team members for fruitful

discussions, and all the LGDP platform members for their outstanding technical assistance and plant

care. This study is set within the framework of the "Laboratoires d'Excellences (LABEX)" TULIP (ANR-

644   REFERENCES

1.   Grewal SI, Jia S. Heterochromatin revisited. Nat Rev Genet. 2007;8(1):35-46.
2.   Slotkin RK, Martienssen R. Transposable elements and the epigenetic regulation of the
genome. Nat Rev Genet. 2007;8(4):272-85.
3.   Deniz O, Frost JM, Branco MR. Regulation of transposable elements by DNA modifications.
Nat Rev Genet. 2019;20(7):417-31.
4.   Dergai O, Hernandez N. How to Recruit the Correct RNA Polymerase? Lessons from snRNA
Genes. Trends Genet. 2019;35(6):457-69.
5.   Law JA, Jacobsen SE. Establishing, maintaining and modifying DNA methylation patterns in
plants and animals. Nat Rev Genet. 2010;11(3):204-20.
6.   Matzke MA, Mosher RA. RNA-directed DNA methylation: an epigenetic pathway of
increasing complexity. Nat Rev Genet. 2014;15(6):394-408.
7.   Wendte JM, Pikaard CS. The RNAs of RNA-directed DNA methylation. Biochim Biophys Acta.
2016.
8.   Du J, Johnson LM, Jacobsen SE, Patel DJ. DNA methylation pathways and their crosstalk with
histone methylation. Nat Rev Mol Cell Biol. 2015;16(9):519-32.
9.   Zhang H, Lang Z, Zhu JK. Dynamics and function of DNA methylation in plants. Nat Rev Mol
Cell Biol. 2018;19(8):489-506.
10.   Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, et al. The
Arabidopsis nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing
heterochromatin. Cell. 2013;153(1):193-205.
11.   Stroud H, Do T, Du J, Zhong X, Feng S, Johnson L, et al. Non-CG methylation patterns shape
the epigenetic landscape in Arabidopsis. Nat Struct Mol Biol. 2014;21(1):64-72.
12.   Moissiard G, Cokus SJ, Cary J, Feng S, Billi AC, Stroud H, et al. MORC family ATPases required
for heterochromatin condensation and gene silencing. Science. 2012;336(6087):1448-51.
13.   Moissiard G, Bischof S, Husmann D, Pastor WA, Hale CJ, Yen L, et al. Transcriptional gene
silencing by Arabidopsis microrchidia homologues involves the formation of heteromers. Proc Natl
Acad Sci U S A. 2014;111(20):7474-9.
14.   Lorkovic ZJ, Naumann U, Matzke AJ, Matzke M. Involvement of a GHKL ATPase in RNA-
directed DNA methylation in Arabidopsis thaliana. Curr Biol. 2012;22(10):933-8.
15.   Ikeda Y, Pelissier T, Bourguet P, Becker C, Pouch-Pelissier MN, Pogorelcnik R, et al.
Arabidopsis proteins with a transposon-related domain act in gene silencing. Nat Commun.
2017;8:15122.
16.   Wenig U, Meyer S, Stadler R, Fischer S, Werner D, Lauter A, et al. Identification of MAIN, a
factor involved in genome stability in the meristems of Arabidopsis thaliana. Plant J. 2013;75(3):469-
83.
17.   Uhlken C, Horvath B, Stadler R, Sauer N, Weingartner M. MAIN-LIKE1 is a crucial factor for
correct cell division and differentiation in Arabidopsis thaliana. Plant J. 2014;78(1):107-20.
18.   Babu MM, Iyer LM, Balaji S, Aravind L. The natural history of the WRKY-GCM1 zinc fingers
and the relationship between transcription factors and transposons. Nucleic Acids Res.
2006;34(22):6505-20.
19.   Steinbauerova V, Neumann P, Novak P, Macas J. A widespread occurrence of extra open
reading frames in plant Ty3/gypsy retrotransposons. Genetica. 2011;139(11-12):1543-55.
20.   Farkas I, Dombradi V, Miskei M, Szabados L, Koncz C. Arabidopsis PPP family of
serine/threonine phosphatases. Trends Plant Sci. 2007;12(4):169-76.

21.   Sun X, Kang X, Ni M. Hypersensitive to red and blue 1 and its modification by protein
phosphatase 7 are implicated in the control of Arabidopsis stomatal aperture. PLoS Genet.
2012;8(5):e1002674.
22.   Liu HT, Li GL, Chang H, Sun DY, Zhou RG, Li B. Calmodulin-binding protein phosphatase PP7 is
involved in thermotolerance in Arabidopsis. Plant Cell Environ. 2007;30(2):156-64.
23.   Genoud T, Santa Cruz MT, Kulisic T, Sparla F, Fankhauser C, Metraux JP. The protein
phosphatase 7 regulates phytochrome signaling in Arabidopsis. PLoS One. 2008;3(7):e2699.
24.   Uhrig RG, Labandera AM, Moorhead GB. Arabidopsis PPP family of serine/threonine protein
phosphatases: many targets but few engines. Trends Plant Sci. 2013;18(9):505-13.
25.   Xu D, Marino G, Klingl A, Enderle B, Monte E, Kurth J, et al. Extrachloroplastic PP7L Functions
in Chloroplast Development and Abiotic Stress Tolerance. Plant Physiol. 2019.
26.   Stroud H, Greenberg MV, Feng S, Bernatavichute YV, Jacobsen SE. Comprehensive analysis
of silencing mutants reveals complex regulation of the Arabidopsis methylome. Cell. 2013;152(1-
2):352-64.
27.   Fransz P, de Jong H. From nucleosome to chromosome: a dynamic organization of genetic
information. Plant J. 2011;66(1):4-17.
28.   Zhao S, Cheng L, Gao Y, Zhang B, Zheng X, Wang L, et al. Plant HP1 protein ADCP1 links
multivalent H3K9 methylation readout to heterochromatin formation. Cell Res. 2019;29(1):54-66.
29.   Tessadori F, Schulkes RK, van Driel R, Fransz P. Light-regulated large-scale reorganization of
chromatin during the floral transition in Arabidopsis. Plant J. 2007;50(5):848-57.
30.   Yokthongwattana C, Bucher E, Caikovski M, Vaillant I, Nicolet J, Mittelsten Scheid O, et al.
MOM1 and Pol-IV/V interactions regulate the intensity and specificity of transcriptional gene
silencing. Embo J.29(2):340-51.
31.   Rigal M, Mathieu O. A "mille-feuille" of silencing: epigenetic control of transposable
elements. Biochim Biophys Acta. 2011;1809(8):452-8.
32.   Galperin MY, Koonin EV. Who's your neighbor? New computational approaches for
functional genomics. Nat Biotechnol. 2000;18(6):609-13.
33.   Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J. 1998;16(6):735-43.
34.   Hristova E, Fal K, Klemme L, Windels D, Bucher E. HISTONE DEACETYLASE6 Controls Gene
Expression Patterning and DNA Methylation-Independent Euchromatic Silencing. Plant Physiol.
2015;168(4):1298-308.
35.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20.
36.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
requirements. Nat Methods. 2015;12(4):357-60.
37.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
38.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput
sequencing data. Bioinformatics. 2015;31(2):166-9.
39.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15(12):550.
40.   Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible WR. Genome-wide identification and
testing of superior reference genes for transcript normalization in Arabidopsis. Plant Physiol.
2005;139(1):5-17.
41.   Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, et al. MEME SUITE: tools for
motif discovery and searching. Nucleic Acids Res. 2009;37(Web Server issue):W202-8.
42.   Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics. 2011;27(11):1571-2.
43.   Huang X, Zhang S, Li K, Thimmapuram J, Xie S, Wren J. ViewBS: a powerful toolkit for
visualization of high-throughput bisulfite sequencing data. Bioinformatics. 2018;34(4):708-9.

44.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. 2004;32(5):1792-7.
45.   Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies
by maximum likelihood. Syst Biol. 2003;52(5):696-704.
46.   Le SQ, Gascuel O. An improved general amino acid replacement matrix. Mol Biol Evol.
2008;25(7):1307-20.
47.   Anisimova M, Gascuel O. Approximate likelihood-ratio test for branches: A fast, accurate,
and powerful alternative. Syst Biol. 2006;55(4):539-52.
48.   Bernatavichute YV, Zhang X, Cokus S, Pellegrini M, Jacobsen SE. Genome-wide association of
histone H3 lysine nine methylation with CHG DNA methylation in Arabidopsis thaliana. PLoS ONE.
2008;3(9):e3156.

FIGURE CAPTIONS

Fig 1. The *ddc #16* EMS population shows overexpression of *ATCOPIA28::GFP* and upregulation of

endogenous TEs.

(A) Schematic representation of the *ATCOPIA28::GFP* transgene. The 5' long terminal repeat (LTR)

promoter region of an *ATCOPIA28* LTR-retrotransposon (*AT3TE51900*) is used to control the

expression of GFP. The construct carries a Nuclear Localization Signal (NLS) to target the GFP in the

nucleus. (B) WT and *drm1 drm2 cmt3* (*ddc*) triple mutant plants carrying the *ATCOPIA28::GFP*

transgene showed no and weak GFP fluorescence under UV light, respectively. By comparison, the *ddc*

#16 EMS mutant showed strong GFP fluorescence. Insets show plants under white light. (C) Western

blot using anti-GFP antibody confirmed *ATCOPIA28::GFP* overexpression in *ddc #16*. Coomassie

staining of the large Rubisco subunit (rbcL) is used as a loading control. KDa: kilodalton. (D) Relative

expression analyses of *ATCOPIA28::GFP (GFP)* and three endogenous TEs in *ddc* and *ddc #16* assayed

by Real-Time quantitative PCR (RT-qPCR)*.* RT-qPCR analyses were normalized using the housekeeping

RHIP1 gene, and transcript levels in the mutants are represented relative to WT. Error bars indicate

standard deviation based on three independent biological replicates. Screening of EMS mutant

populations was done on MS plates to allow for visualization of GFP-positive individuals under UV

light.

Fig 2. MAIN, DRM2 and CMT3 act synergistically to repress TEs.

(A) Representative pictures showing the developmental phenotype of 3-week-old *ddc*, *main-3* and *ddc main-3* mutants in comparison to WT plant. (B) Number of upregulated TEs in *ddc*, *main-3* and *ddc main-3*, and classified by TE superfamily. (C) Chromosomal distributions of misregulated loci in *ddc*, *main-3* and *ddc main-3* over WT. Chromosome arms are depicted in light grey, pericentromeric regions in dark grey as defined in [48]. Upregulated genes are represented in blue, downregulated genes in green and TEs are represented in red above the chromosomes (upregulated) or below (downregulated). (D) Fraction of upregulated TEs in *ddc*, *main-3* and *ddc main-3* located in chromosome arms or in pericentromeric regions as defined in [48]. Asterisks indicate statistically significant enrichments of TEs in pericentromeric regions in comparison to the genomic distribution of all *A. thaliana* TEs (Chi-Square test, *: p-value≤ 0.05, n.s: not significant). (E) Heatmap showing normalized count reads of upregulated TEs in *ddc*, *main-3* and *ddc main-3* mutants in comparison to WT plants. (F-G) Relative expression analyses of *ATCOPIA28* (F) and several endogenous TEs (G) in *ddc*, *main-3, ddc main-3, cmt3 main-3* and *drm1 drm2 (dd) main-3* assayed by RT-qPCR. RT-qPCR analyses were normalized using the housekeeping *RHIP1* gene, and transcript levels in the mutants are represented relative to WT. Error bars indicate standard deviation based on three independent biological replicates. Analyses described in panels B-E are based on the overlaps of two independent RNA-seq experiments (EMS Exp1 and Exp2, four biological replicates, S1 and S2 Tables). RNA-seq threshold: log2≥2, or log2≤-2 ; p-adj< 0.01.

Fig 3. MAIN and MAIL1 are required for the proper expression of similar genes, and for TE silencing.

(A-B) Number of reproducibly misregulated genes (A) and upregulated TEs (B) in *main-2, mail1-1* and *main-3* mutants in comparison to WT plants. TEs are classified by superfamily. (C) Heatmap showing normalized count reads of misregulated loci in *main-2*, *mail1-1* and *main-3* in comparison to

respective WT controls. Asterisks represents loci that are commonly misregulated in the three mutant

backgrounds. (D) Venn diagrams analyses representing the overlaps between misregulated loci in

main-2, *mail1-1* and *main-3*. Fisher's exact test statistically confirmed the significance of Venn diagram

overlaps (p-value <1,2.10e-14). (E) Fraction of misregulated loci in *main-2* and *mail1-1* located in

chromosome arms or in pericentromeric regions as defined in [48]. Asterisks indicate statistically

significant enrichments of downregulated genes and upregulated TEs in chromosome arms and

pericentromeric regions, respectively, in comparison to the genomic distributions of all *A. thaliana*

genes and TEs (Chi-Square test, *: p-value≤ 0.05, **: p-value≤ 0.01, n.s: not significant). Percentages

of DNA-methylated genes were calculated based on whole genome bisulfite sequencing (BS-seq) of

WT plants described in [26]. n.d.: not determined. Analyses are based on the overlaps of three

independent RNA-seq experiments (Exp1 and Exp2, two biological replicates each, and Exp3, four

biological replicates, S3 and S4 Tables). RNA-seq threshold: log2≥2, or log2≤-2 ; p-adj< 0.01.

Fig 4. The *main-2* mutation has a slight effect on non-CG DNA methylation levels.

(A) Genome-wide DNA methylation levels along the five *Arabidopsis* chromosomes in *main-2* versus

WT plants. Chromosome arms are depicted in light grey, pericentromeric regions in dark grey as

defined in [48]. Mb: megabase. (B-H) Boxplot analyses in two *main-2* and WT biological replicates

showing the DNA methylation levels of all pericentromeric TEs (B) and genes (C), all chromosome arms

TEs (D) and genes (E), TEs that are upregulated in *main-2* (F), and genes that are upregulated (G) and

downregulated (H) in *main-2*. p-values were calculated using a Wilcoxon test. ***: p-value < 2.10e-16.

Fig 5. MAIN, MAIL1 and PP7L physically interact together.

(A) Representative pictures of 3-week-old *main-2* and *mail1-1* mutants, and epitope-tagged

complementing lines in comparison to WT Col plants. (B) Western blots using anti-FLAG and anti-Myc

antibodies showing the accumulation of epitope-tagged PMD proteins at the expected sizes in the

different complementing lines. Coomassie staining of the large Rubisco subunit (rbcL) is used as a

loading control. KDa: kilodalton.(C-E) Relative expression analyses of upregulated TEs (C), upregulated

genes (D) and downregulated genes (E) in the different complementing lines assayed by RT-qPCR. RT-

qPCR analyses were normalized using the housekeeping *RHIP1* gene, and transcript levels in the

complementing lines and mutants are represented relative to WT Col. Error bars indicate standard

deviation based on three independent biological replicates. (F) FLAG-tagged MAIN and MAIL1 proteins

were immunoprecipitated and putative interacting proteins were identified by mass spectrometry.

Numbers of identified spectra, peptides and the normalized spectral abundance factor (NSAFe5) are

shown for two independent experiments, including three *main-2* and two *mail1-1* replicates. WT

replicates are used as a negative control. Only proteins reproducibly enriched in all the FLAG-MAIN

and FLAG-MAIL1 IP, and depleted in WT controls across multiple replicates are described in the table.

(G) MAIL1-MYC was co-immunoprecipitated with MAIN-FLAG in F1 plants obtained by crossing MAIL1-

MYC and MAIN-FLAG lines together. Parental MAIL1-MYC and MAIN-FLAG lines were used as negative

controls. (H) The MAIN-MYC line was supertransformed with the PP7L-FLAG construct, and MAIN-MYC

was co-immunoprecipitated with PP7L-FLAG. Plants expressing only MAIN-MYC or PP7L-FLAG were

used as negative controls. (I) Same as H but using MAIL1-MYC plants supertransformed with the PP7L-

FLAG construct. Epitope-tagged proteins were detected by Western blotting. Arrowheads indicates

expected bands. Asterisks indicates non-specific hybridization. Co-exp: plants co-expressing PP7L-

FLAG and MAIN-MYC (H) or PP7L-FLAG and MAIL1-MYC (I).

Fig 6. *main-2*, *mail1-1*, *pp7l-2* single and *mail1-1 pp7l-2* double mutants display similar

developmental and molecular phenotypes.

(A) Representative pictures of 3-week-old *main-2*, *mail1-1*, *pp7l-2* single and *mail1-1 pp7l-2* double

mutants in comparison to WT Col plant. (B) Heatmap showing normalized count reads of misregulated

loci in *main-2*, *mail1-1*, *pp7l-2* and *mail1-1 pp7l-2* mutants in comparison to WT col plants using the

datasets of RNA-seq Exp3 (4 biological replicates, S3 and S6 Tables). One asterisk defines the loci that

are commonly misregulated in all mutant backgrounds. Two asterisks define the loci that are

misregulated in the *mail1-1 pp7l-2* double mutant. (C) Venn diagrams analyses representing the

overlaps between misregulated loci in *main-2*, *mail1-1*, *pp7l-2* and *mail1-1 pp7l-2*. Fisher's exact test

statistically confirmed the significance of Venn diagram overlaps (p-value <1,2.10e-14). (D-F) Boxplots

analyses showing average RPKM values of upregulated TEs (D), upregulated genes (E) and

downregulated genes (F) in *mail1-1 pp7l-2* in the indicated genotypes of RNA-seq Exp3. These analyses

are based on the misregulated loci datasets defined by ** in panel B. P-values were calculated using

a Wilcoxon test, and only significant p-values are shown. *: p-value< 1.10e-3; ***: p-value< 2.10e-16.

(G) Heatmap showing normalized count reads of reproducibly misregulated loci in *main-2* and *mail1-*

1 in the indicated genotypes of RNA-seq Exp3. Lists of reproducibly misregulated loci in *main-2* and

mail1-1 were as defined in Fig 3 and S4 Table. One asterisk defines the loci that are commonly

misregulated in *main-2* and *mail1-1* mutants (S5 Table). Two asterisks define subsets of loci that are

commonly misregulated in all the indicated genotypes (S7 Table). (H-J) Relative expression analyses

of upregulated TEs (H), upregulated genes (I) and downregulated genes (J) in the different genotypes

assayed by RT-qPCR. RT-qPCR analyses were normalized using the housekeeping *RHIP1* gene, and

transcript levels in the different mutants are represented relative to WT Col. Error bars indicate

standard deviation based on three independent biological replicates. (K-M) Boxplots analyses showing

average RPKM values of commonly upregulated TEs (K), upregulated genes (L) and downregulated

genes (M) in the indicated genotypes of RNA-seq Exp3. These analyses are based on the commonly

misregulated loci datasets defined by ** in panel G. P-value was calculated using a Wilcoxon test. ***:

p-value < 2.10e-16.

Fig 7. Constitutive heterochromatin appears unaltered in *pp7l-2* mutant.

Proportion of nuclei showing condensed, partially decondensed (intermediate), or decondensed

chromocenters in the *pp7l-2* mutant in comparison to WT control (Col) based on H3K9me2

immunostaining of nuclei. Representative pictures of nuclei displaying condensed, partially

decondensed or decondensed chromocenters.   DAPI: DNA stained with 4',6-diamidino-2-

phenylindole.

Fig 8. Evolutionary history of PMD-C and PP7 proteins in plants.

(A) An alignment of the PMD-C motifs from 30 representative Eudicot species was used to construct

a phylogenetic tree. The two major clades (MAIL2/MAIL2-like and MAIL3) are indicated. The species

codes are given in S11 Table, and corresponding protein sequences in S12 Table). In red are genes

presenting a fusion between a PMD-C and a PP7 motif. Statistical supports of key nodes calculated

with the approximate likelihood-ratio test are indicated. Scale bar indicates one substitution/site. The

tree was rooted using the *Amborella trichopoda* PMD-C motif (Atr1PMDC). (B) Phylogenetic tree

constructed using an alignment of the PP7 motif from the same species as in (A). The two major clades

(PP7 and PP7L) are indicated. In red are genes presenting a fusion between a PP7 and a PMD-C motif.

Statistical supports of key nodes calculated with the approximate likelihood-ratio test are indicated.

Scale bar indicates one substitution/site. The tree was rooted using the *A. thaliana* PP5 motif (AtPP5).

885
886

 SUPPORTING INFORMATION CAPTIONS

S1 Fig. *MAIN* is the mutated gene responsible for *ATCOPIA28::GFP* and TE overexpression in the *ddc*

***#16* mutant.**

(A) Representative pictures of *ddc #18* (*ddc morc6-8*) and *ddc #344* (*ddc morc6-9*) mutants in

comparison to *ATCOPIA28::GFP* WT and *ddc* control plants under UV light. Insets show plants under

white light. (B) Enrichment in homozygote/heterozygote ratio of EMS over WT single nucleotide

polymorphisms (SNPs), defining the linkage intervals for the populations *ddc #18* and *ddc #344*. Mb:

megabase. (C) Location of the point mutations corresponding to the *morc6-8* and *morc6-9* alleles

within the *MORC6* genomic sequence. Nucleotide and corresponding amino acid changes are

indicated above the gene. Positions of the mutations are indicated relative to the transcription start

site (+1). Grey boxes represent 5' and 3' UTR, blue boxes and lines represent exons and introns,

respectively. (D) Enrichment in homozygote/heterozygote ratio of EMS over WT single nucleotide

polymorphisms (SNPs), defining the linkage intervals for the population *ddc #16*. (E) Location of the

point mutation corresponding to the *main-3* mutant allele within the *MAIN* genomic sequence. (F)

Genetic complementation analyses using the KO T-DNA insertion line *main-2*. *ddc #16* plants were

crossed with *main-2* plants. F1 plants were self-crossed, and F2 plants were screened under UV light

to select GFP-overexpressing plants. Western blotting using anti-GFP antibodies confirmed GFP

overexpression in selected F2 plants. Coomassie staining of the large Rubisco subunit (rbcL) is used as

a loading control. KDa: kilodalton. Among the selected F2 plants, the presence of *main-3* EMS and

main-2 T-DNA mutant alleles were determined by dCAPS-PCR and PCR analyses, respectively. *DRM2*

and *CMT3* genotyping were determined by PCR analyses. WT: Wild type, Ho: Homozygote mutant. He:

Heterozygote. (G) Relative expression analyses of several TEs in the indicated genotypes assayed by

RT-qPCR. RT-qPCR analyses were normalized using the housekeeping *RHIP1* gene, and transcript levels

in the different genotypes are represented relative to WT. Error bars indicate standard deviation

based on two independent biological replicates. Screening of EMS mutant populations was done on

MS plates to allow for visualization of GFP-positive individuals under UV light.

S2 Fig. Combining the *drm2*, *cmt3* and *main-3* mutations exacerbate TE silencing defects.

(A) Number of misregulated loci in *ddc*, *main-3* and *ddc main-3* in comparison to *ATCOPIA28::GFP* WT

plants in two independent RNA-seq experiments (EMS Exp1 and Exp2, two biological replicates each,

S1 Table). (B) Principal component analysis (PCA) performed on normalized count reads for first two

components of the sixteen samples described in RNA-seq EMS Exp1 and Exp2. (C) Venn diagrams

analyses defining the reproducibly misregulated loci in the different genotypes based on the overlaps

of loci identified as misregulated in RNA-seq EMS Exp1 and Exp2 (S2 Table). (D) Relative expression

analyses of *ATCOPIA28* and *HELITRONY1D* (*AT5TE35950*) in *ddc*, *main-3* and *ddc main-3* assayed by

RT-qPCR. RT-qPCR analyses were normalized using the housekeeping *RHIP1* gene, and transcript levels

in the different genotypes are represented relative to WT. Error bars indicate standard deviation

based on three independent biological replicates. (E) Venn diagrams analysis showing the overlaps

between reproducibly upregulated TEs in *ddc*, *main-3* and *ddc main-3*. The numbers of reproducibly

upregulated TEs for each genotype were defined by the overlaps of upregulated TEs shown in panel

C. Fisher's exact test statistically confirmed the significance of Venn diagram overlaps (p-value

<1,2.10e-14). (F) Same as panel D for TEs defined as class I-IV TEs. Frames of RT-qPCR graphs are using

the same color code as shown in panel E. (G) *top*, Venn diagrams analyses defining the overlaps

between reproducibly up- and downregulated genes in the different genotypes. Reproducibly

misregulated genes were defined based on the overlaps described in panel C. Fisher's exact test

statistically confirmed the significance of Venn diagram overlaps (p-value <1,2.10e-14). *bottom,*

Fraction of misregulated genes in *ddc*, *main-3* and *ddc main-3* located in chromosome arms or in

pericentromeric regions as defined in [48]. Asterisks indicate statistically significant enrichments of

misregulated genes in chromosome arms or pericentromeric regions in comparison to the genomic

distributions of all *A. thaliana* genes (Chi-Square test, *: p-value≤ 0.05, **: p-value≤ 0.01, n.s: not

significant). Percentages of DNA-methylated genes were calculated based on whole genome bisulfite

sequencing (BS-seq) of WT plants described in [26]. n.d.: not determined. (H) Relative expression

analyses of *DRM2* and *CMT3* in *ddc*, *main-3*, *ddc main-3, cmt3 main-3* and *dd main-3* assayed by RT-

qPCR. RT-qPCR analyses were normalized using the housekeeping *RHIP1* gene, and transcript levels in

the different genotypes are represented relative to WT. Error bars indicate standard deviation based

on three independent biological replicates. Screening of EMS mutant populations was done on MS

plates to allow for visualization of GFP-positive individuals under UV light.

S3 Fig. Identification of reproducibly misregulated loci in *main-2*, *mail1-1* and *main-3*.

(A) Number of misregulated loci in *main-2* and *mail1-1* in comparison to WT Col plants in three

independent RNA-seq experiments (Exp1, Exp2 [15], and Exp3; two, two and four biological replicates,

respectively, S3 Table) (B) Principal component analysis (PCA) performed on normalized count reads

for first two components of the twenty-four *main-2*, *mail1-1* and WT Col samples described in RNA-

seq Exp1, Exp2 and Exp3. (C) Venn diagrams analyses defining the reproducibly misregulated loci in

main-2 and *mail1-1* based on the overlaps of loci identified as misregulated in RNA-seq Exp1, Exp2

and Exp3 (S4 Table). (D-F) Relative expression analyses of several upregulated TEs (D), upregulated

genes (E), and downregulated genes (F) in *main-2, mail1-1* and *main-3* assayed by RT-qPCR. RT-qPCR

analyses were normalized using the housekeeping *RHIP1* gene, and transcript levels in the different

genotypes are represented relative to respective WT controls. Error bars indicate standard deviation

based on three independent biological replicates.

S4 Fig. DNA methylation analyses in the *main-2* mutant

(A-B) Boxplot analyses in two *main-2* and WT biological replicates showing the DNA methylation levels

at genomic sites previously defined as hypo CHG differentially methylated regions (DMR) in *cmt3* (A)

and hypo CHH DMR in *drm1 drm2* (B) based on [26]. p-values were calculated using a Wilcoxon test.

*: p-value <5.10e-7, **: p-value <5.10e-10, ***: p-value < 2.10e-16.

S5 Fig. MAIN, MAIL1 and PP7L are required for the proper expression of similar loci, and commonly

downregulated genes carry the 'DOWN' DNA motif in their promoter.

(A) Principal component analysis (PCA) performed on normalized count reads for first two components

of the twenty samples described in RNA-seq Exp3. (B) Number of misregulated loci in the different

genotypes in comparison to WT Col plants from RNA-seq Exp3 (four biological replicates, S3 and S6

Tables). (C) Identification and proportions of the 'DOWN' DNA motif among the promoters of

downregulated genes and all *Arabidopsis* genes using the MEME software. Promoter regions are

defined as 1kb upstream of ATG. The list of all *Arabidopsis* genes used to determine genomic

distributions is based on the TAIR file: TAIR10_upstream_1000_translation_start_20101028.

S6 Fig. Full size images of panels described in Fig 5G-I.

S1 Table. Lists of differentially regulated loci in *ddc*, *main-3* and *ddc main-3* in two independent

RNA-seq experiments.

S2 Table. Lists of reproducibly differentially regulated loci in *ddc*, *main-3* and *ddc main-3*.

S3 Table. Lists of differentially regulated loci in *main-2* and *mail1-1* in three independent RNA-seq

experiments.

S4 Table. Lists of reproducibly differentially regulated loci in *main-2* and *mail1-1*.

S5 Table. Lists of loci reproducibly and commonly misregulated in *main-2*, *mail1-1* and *main-3*.

S6 Table. Lists of differentially regulated loci in *pp7l-2* and *mail1-1pp7l-2* in RNA-seq Exp3.

S7 Table. Lists of loci commonly misregulated in *main-2*, *mail1-1*, *pp7l-2* and *mail1-1pp7l-2*.

S8 Table. Lists of loci commonly misregulated in all mutant backgrounds analyzed in this study.

S9 Table. Lists of commonly downregulated genes displaying the "DOWN" motif in their promoter

and random test analyses.

S10 Table. Distribution of several bZIP DNA motifs among the promoter regions of all *Arabidopsis*

genes.

S11 Table. List of species used to construct the two trees of figure 8, their codes and the

presence/absence of the different PMD-C and PP7 motifs.

S12 Table. (A) PMD-C and (B) PP7/PP7L motifs used to construct the two phylogenetic trees of

Figure 8.

1002

1003   **S13 Table. List of primers used in this study.**

1004

1005   **S14 Table. Next Generation Sequencing (NGS) mapping and coverage statistics.**

1006

1007

1008

1009

Fig1

A

B

C

D

Fig 2

Fig 3

Fig 4

Fig 5

A

Col — MAIN-FLAG MAIN-MYC — MAIL1-FLAG MAIL1-MYC

main-2 *mail1-1*

B

kDa	Col	*main-2*	MAIN-FLAG	*mail1-1*	MAIL1-FLAG	
75–						**FLAG**
Coomassie						**rbcL**
100–75–						**MYC**
Coomassie						**rbcL**

C Relative transcript level
ATCOPIA28 *ATMU1* *ROMANIAT5* *HELITRONY1D*

D *TAF4b* *Cox19L* *SHOC1* *UNK*

E *HAC4* *MKK3* *FAP3*

F

			Spectral count			Peptide count			NSAFe5		
			MAIN	MAIL1	PP7L	MAIN	MAIL1	PP7L	MAIN	MAIL1	PP7L
FLAG IP	**Exp1**	MAIN-FLAG Rep. 1	33	39	35	19	20	22	794.0899	881.3135	670.9659
		MAIN-FLAG Rep. 2	34	42	38	16	18	23	1298.8062	1506.6931	1156.4468
		MAIL1-FLAG Rep. 1	20	25	14	13	14	12	905.1708	1062.5531	504.78354
		WT control Rep. 1	0	0	0	0	0	0	0	0	0
		WT control Rep. 2	0	0	0	0	0	0	0	0	0
	Exp2	MAIN-FLAG Rep. 3	90	75	59	23	33	28	487.18513	381.26144	254.43695
		MAIL1-FLAG Rep. 2	28	57	37	17	27	21	170.688	326.30966	179.68976
		WT control Rep. 3	0	0	0	0	0	0	0	0	0

G INPUT FLAG IP
MAIN-FLAG MAIL1-MYC F1 / MAIN-FLAG MAIL1-MYC F1
FLAG MYC

H INPUT FLAG IP
PP7L-FLAG MAIN-MYC Co-exp. / PP7L-FLAG MAIN-MYC Co-exp.
FLAG MYC

I INPUT FLAG IP
PP7L-FLAG MAIL1-MYC Co-exp. / PP7L-FLAG MAIL1-MYC Co-exp.
FLAG MYC

Fig 6

Fig 7

Fig 8