The plant mobile domain proteins MAIN and MAIL1 interact with the

phosphatase PP7L to regulate gene expression and silence transposable

elements in *Arabidopsis thaliana*.

Short title: The PMD MAIN/MAIL1 and PP7L complex regulates gene expression and TE silencing.

Melody Nicolau[1,2], Nathalie Picault[1,2], Julie Descombin[1,2], Yasaman Jami-Alahmadi[3], Suhua Feng[4],

Etienne Bucher[5], Steven E. Jacobsen[4,6], Jean-Marc Deragon[1,2], James Wohlschlegel[3] and Guillaume

Moissiard[1,2]*.

1 LGDP-UMR5096, CNRS, Perpignan, France.

2 LGDP-UMR5096, Université de Perpignan Via Domitia, France.

3 Department of Biological Chemistry, University of California at Los Angeles, Los Angeles, CA, USA.

4 Department of Molecular, Cell and Developmental Biology, University of California at Los Angeles,

Los Angeles, CA, USA.

5 Plant Breeding and Genetic Resources, Agroscope, Nyon, Switzerland.

6 Howard Hughes Medical Institute, University of California at Los Angeles, Los Angeles, CA, USA.

* Corresponding author

E-mail: guillaume.moissiard@univ-perp.fr

ABSTRACT

Transposable elements (TEs) are DNA repeats that must remain silenced to ensure cell integrity. Several epigenetic pathways including DNA methylation and histone modifications are involved in the silencing of TEs, and in the regulation of gene expression. In *Arabidopsis thaliana*, the TE-derived plant mobile domain (PMD) proteins have been involved in TE silencing, genome stability, and control of developmental processes. Using a forward genetic screen, we found that the PMD protein MAINTENANCE OF MERISTEMS (MAIN) acts synergistically and redundantly with DNA methylation to silence TEs. We found that MAIN and its close homolog MAIN-LIKE 1 (MAIL1) interact together, as well as with the phosphoprotein phosphatase (PPP) PP7-like (PP7L). Remarkably, *main*, *mail1* and *pp7l* mutants display similar developmental phenotypes, and share common subsets of upregulated TEs and misregulated genes. Finally, phylogenetic analyses of PMD and PP7-type PPP domains among the Eudicot lineage suggest neo-association processes between the two protein domains to potentially generate new protein function. We propose that, through this interaction, the PMD and PPP domains may constitute a functional protein module required for the proper expression of genes and silencing of TEs.

AUTHOR SUMMARY

The plant mobile domain (PMD) is a protein domain of unknown function that is widely spread in the angiosperm plants. Although most PMDs are associated with repeated DNA sequences called transposable elements (TEs), plants have domesticated the PMD to produce genic versions that play important roles within the cell. In *Arabidopsis thaliana*, MAINTENANCE OF MERISTEMS (MAIN) and MAIN-LIKE 1 (MAIL1) are genic PMDs that are involved in genome stability, developmental processes, and silencing of TEs. The mechanisms involving MAIN and MAIL1 in these cellular processes remain elusive. Here, we show that MAIN, MAIL1 and the phosphoprotein phosphatase (PPP) named PP7-like (PP7L) interact to form a protein complex that is required for the proper expression of genes, and the

silencing of TEs. Phylogenetic analyses revealed that PMD and PP7-type PPP domains are evolutionary

connected, and several plant species express proteins carrying both PMD and PPP domains. We

propose that interaction of PMD and PPP domains would create a functional protein module involved

in mechanisms regulating gene expression and repressing TEs.

INTRODUCTION

In eukaryotes, DNA methylation and post-translational modifications of histones are

epigenetic marks involved in chromatin organization, regulation of gene expression and silencing of

DNA repeats such as transposable elements (TEs) [1-3]. Constitutive heterochromatin is highly

condensed and enriched in silenced TEs that are heavily DNA methylated, while euchromatin is more

relaxed and composed of genes that are more permissive to transcription, depending on the

recruitment of transcription factors (TFs), cofactors and RNA polymerases [1, 4]. In plants, DNA

methylation occurs in three different cytosine contexts: CG, CHG and CHH (where H = A, T or C),

involving specialized DNA methyltransferases [5]. In *Arabidopsis thaliana,* DOMAINS REARRANGED

METHYLTRANSFERASE 1 (DRM1) and DRM2 mediate de novo DNA methylation in all sequence

contexts through the RNA-directed DNA methylation (RdDM) pathway, which involves among other

components, RNA-DEPENDENT RNA POLYMERASE 2 (RDR2) and DICER-LIKE 3 (DCL3) for the

production of short interfering (si)RNAs [6, 7]. The maintenance of CG methylation is specifically

performed by METHYLTRANSFERASE 1 (MET1), while CHROMOMETHYLASE 2 (CMT2) and CMT3 are

involved in the maintenance at CHG sites [8, 9]. CMT2 can also be involved in the deposition of CHH

methylation at specific genomic location [10, 11]. Finally, DRM1/2 are mostly required for the

maintenance of CHH methylation through the RdDM pathway [6, 7, 9]. Together with DNA

methylation, additional pathways play important roles in TE silencing. MORPHEUS MOLECULE 1

(MOM1) is required for TE silencing [12]. MICRORCHIDIA 1 (MORC1) and MORC6 interact together,

and are required for heterochromatin condensation and repression of TEs, acting mostly downstream

of DNA methylation and RdDM pathway [13-15]. It has been proposed that the MORC1/6 complex

would interact with the RdDM machinery to promote its own recruitment to heterochromatic loci that

must remain condensed and silenced [16].

More recently, the *A. thaliana* plant mobile domain (PMD) proteins MAINTENANCE OF

MERISTEM (MAIN) and MAIN-LIKE 1 (MAIL1) were identified as new factors required for TE silencing

[17]. In addition, these two proteins have been involved in genome stability, and regulation of

developmental processes such as cell division and differentiation [18, 19]. The PMD is a large protein

domain of unknown function that is widely represented among the angiosperms, predominantly

associated with TEs [17, 20]. It has been proposed that genic PMD versions, such as the MAIN and

MAIL1 proteins derived from TEs after gene domestication [17, 20, 21]. Previous studies suggested

that genic PMDs could act as cellular factors related to transcription, possibly acting as transcription

factor (TF)-like, co-factor or repressor proteins regulating this cellular process [18, 20]. Nevertheless,

the role of PMD proteins in the regulation of transcription remains elusive. Most of genic PMD

proteins are standalone versions, however, in some cases, the PMD is fused to another protein

domain, such as protease, kinase or metallo-phosphatase (MPP) domains. For instance in *A. thaliana*,

the MAIL3 protein carries a PMD, which is fused to a putative serine/threonine-specific

phosphoprotein phosphatase (PPP) domain phylogenetically related to the plant-specific protein

phosphatase 7 (PP7) [22]. PP7 is a calmodulin-binding PPP that has been related to cryptochrome

(CRY)-mediated blue-light signaling, and to the control of stomatal aperture [22-24]. PP7 is also

involved in the perception of red/far red light by controlling the phytochrome pathway [25, 26]. In

addition to PP7 and MAIL3 (also known as "long PP7"), the protein PP7-like (PP7L) belongs to the same

phylogenetic clade [22]. PP7L was recently identified as a nuclear protein involved in chloroplast

development and abiotic stress tolerance [27]. The *pp7l* mutant plants showed photosynthetic defects

and strong developmental phenotype associated with misregulation of several genes [27].

In this study, we describe a forward genetic screen based on a GFP reporter gene that allowed

us to identify a mutant population in which *MAIN* is mutated, leading to GFP overexpression. We then

decipher the genetic interaction between the DRM1/2, CMT3 and MAIN, showing that these proteins

act in different epigenetic pathways that act redundantly and synergistically to repress TEs.

Biochemical analyses indicate that MAIN and MAIL1 physically interact together. We have also

identified PP7L as a robust interactor of the two PMD proteins. In addition, the characterization of

developmental and molecular phenotypes of *pmd* and *pp7l* mutant plants strongly suggest that these

proteins interact together to silence TEs, and regulate the expression of a common set of genes.

Finally, phylogenetic analyses allowed us to determine the distribution of PMD, PP7 and PP7L domains

among the Eudicots. Based on these analyses, we have evidences of co-evolution linked to the neo-

association of the PMD and PP7-type PPP domains on single proteins in several Eudicot species,

suggesting a convergent evolution between these two protein domains.

RESULTS

MAIN acts synergistically with DRM1/2 and CMT3 to repress TEs.

The *ATCOPIA28* retrotransposon *AT3TE51900* (hereafter called *ATCOPIA28*) is targeted by

several epigenetic pathways such as DNA methylation and the MORC1/6 complex, which altogether

contribute to its repression. We engineered a construct in which the 5' long terminal repeat (LTR)

promoter region of *ATCOPIA28* controls GFP transcription (S1A Fig). While the *ATCOPIA28::GFP*

transgene is fully silenced in wild type (WT) plants, it is weakly expressed in the DNA methylation-

deficient *drm1 drm2 cmt3 (ddc)* triple mutant background (Fig 1A) [28]. We performed an ethyl

methane sulfonate (EMS) mutagenesis using the *ATCOPIA28::GFP ddc* plants as sensitized genetic

material, and screened for mutant populations showing GFP overexpression. Among, the selected

populations, we retrieved two new mutant alleles of *MORC6* carrying missense mutations in either

the GHKL or S5 domains of the protein (S1B-D Fig). We also identified the population *ddc #16* showing

strong overexpression of GFP and misregulation of several endogenous TEs, including *ATCOPIA28* (Fig

1A-C). Mapping experiments based on whole genome resequencing and bulk segregant analysis

indicated that *ddc #16* carries a missense point mutation ($C_{230} \rightarrow Y$) in the gene *AT1G17930*, previously

named *MAIN* (S1C and S1E Fig). Genetic complementation analyses using the knock-out (KO)

transferred DNA (T-DNA) insertion line *main-2* confirmed that *MAIN* was the mutated gene causing

the overexpression of GFP (S1F Fig). Therefore, *ddc #16* was renamed *ddc main-3.*

To determine the genetic interaction of *ddc* and *main-3* mutations on TE silencing, we carried

out RNA sequencing (RNA-seq) analyses in the hypomorphic *main-3* single, *ddc* triple and *ddc main-3*

quadruple mutant plants (S1G Fig). As previously described, the *ddc* mutant showed upregulation of

several TEs [11], which was also observed to a milder degree in the *main-3* mutant (Figs 1D and S1H,

and S1 Table). The *ddc main-3* mutant showed an exacerbation of TE silencing defects, with a large

number of pericentromeric TEs being specifically upregulated in this mutant background (Figs 1D-E

and S1H-I, and S1 Table). Reverse transcription quantitative PCR (RT-qPCR) analyses confirmed the

RNA-seq results, and showed that *ddc* and *main-3* mutations have synergistic effects on TE

upregulation (Fig 1F). These analyses reveal that DRM1/2, CMT3 and MAIN act redundantly and

synergistically in the silencing of several TEs, strongly suggesting that these proteins act in different

silencing pathways.

MAIN, MAIL1 and the metallo-phosphatase PP7L physically interact together.

The *main-2* and *mail1-1* null mutants show similar developmental phenotypes, and

transcriptomic analyses have revealed that many TEs, including *ATCOPIA28*, are overexpressed in

these two mutants (S2A and B Fig, and S2 Table) [17]. In addition, several genes are misregulated in

the two *pmd* mutants (S2C and D Fig, and S2 Table). It has been suggested that MAIN and MAIL1 act

in the same pathway, possibly by interacting together [17]. To test this hypothesis, we generated

transgenic lines expressing FLAG- and MYC-tagged genomic PMD versions driven by their endogenous

promoters. We showed that epitope-tagged MAIN and MAIL1 proteins were produced at the expected

sizes, and they could complement the respective developmental phenotypes of null mutant plants

(S2A and S2E Fig). Importantly, the different epitope-tagged PMD proteins could also efficiently rescue

the TE silencing defects observed in the two *pmd* mutants (S2F Fig). Using FLAG-tagged MAIN and

MAIL1 expressing plants, immunoprecipitation followed by mass spectrometry (IP-MS) analyses were

carried out to determine potential protein interactors. Mass spectrometry (MS) analyses indicated

that MAIL1 was strongly immunoprecipitated with MAIN-FLAG and *vice versa* (Fig 2A). Co-

immunoprecipitation (co-IP) experiments using F1 transgenic line expressing MAIN-FLAG and MAIL1-

MYC transgenes validated IPMS results (Fig 2B). MS analyses of MAIN-FLAG and MAIL1-FLAG IP also

identified the metallo-phosphatase PP7L as putative interactor (Fig 2A). Co-IP experiments using

plants co-expressing either PP7L-FLAG together with MAIN-MYC, or PP7L-FLAG in combination with

MAIL1-MYC constructs confirmed the interaction between PP7L and the two PMD proteins (Fig 2C and

D). Thus, the three proteins MAIN, MAIL1 and PP7L physically interact together.

The *main, mail1* and *pp7l* mutants display similar developmental and molecular phenotypes.

PP7L is a putative metallo-phosphatase that is related to the phosphatase PP7, and it was

recently identified as a nuclear protein required for photosynthesis [22, 27]. Because PP7L interacts

with MAIN and MAIL1, and the *pp7l-2* null mutant displays abnormal developmental phenotype

reminiscent of the two *pmd* mutant plants (Figs 2 and S3A), we decided to reanalyze available *pp7l*

RNA-seq data [27]. Like in the two *pmd* mutants, several TEs were upregulated and hundreds of genes

were misregulated in *pp7l* (S3B and S3C Fig, and S2 Table). We then compared the transcriptomes of

pmd and *pp7l* mutants, and found that 56% and 60% of TEs upregulated in *main* and *mail1* mutants

were also upregulated in *pp7l* (Fig 3A). Similar analyses with misregulated genes revealed that a

fraction of genes that were down- or upregulated in *pp7l* were similarly misregulated in the two *pmd*

mutants (Fig 3B and C). Moreover, we found that 38% and 47% of genes downregulated in *main* and

mail1 mutants were also repressed in *pp7l*, while only 6 % of genes upregulated in the *pmd* mutant

were similarly overexpressed in *pp7l* (Fig 3B and C). RT-qPCR analyses confirmed the upregulation of

several TEs in the *pp7l-2* null allele (Fig 3D). These analyses also validated that several genes were

similarly misregulated in *pp7l*, *main* and *mail1* mutants (Figs 3E-F and S2D). In addition, whole genome

bisulfite sequencing analyses performed in *main* showed that DNA methylation level is largely

unaffected in the *pmd* mutant in comparison to WT plants (S3D-E Fig). Finally, we performed in silico

analyses to identify any putative DNA cis regulatory elements (CREs) within a 1-kb promoter region

upstream of start codon of genes that were similarly down- or upregulated in the three mutant

backgrounds (Fig 3A-C). We found that a specific DNA decamer motif was significantly enriched in

twenty-one out of the twenty-four downregulated genes (Fig 3G and S3 Table). We also identified a

DNA motif in ten out of the eleven upregulated genes, although the E-value was not significant, most

likely because of the small sample size (S3F Fig and S3 Table). Remarkably, the DNA motif identified

among the dowregulated genes was also strongly enriched in the list of downregulated genes in the

hypomorphic *main-3* mutant (twenty-seven out of twenty-eight genes), and very similar motifs were

retrieved in the lists of downregulated genes in *ddc main-3*, *main-2* and *mail1-1* mutants (S3G Fig and

S3 Table). Thus, the MAIN/MAIL1 protein complex interacts with PP7L, and the three proteins are

required for the proper expression of a common set of loci. Moreover, most of the loci that are

commonly downregulated in the *main*, *mail1* and *pp7l* mutants display a specific DNA motif within

their promoters, suggesting that the transcriptional regulation of these loci could occur through the

recognition of a putative CRE.

The PMD and PP7 domains have co-evolved among the Eudicots.

Among the Angiosperms, most of the genic PMDs, like MAIN and MAIL1, are standalone

versions [20]. However, some genic PMDs can associate with other protein domains, such as for

instance a PPP domain. In *A. thaliana*, the protein MAIL3, which carries a PMD fused to a PPP domain,

is a close homolog of both MAIN/MAIL1 and PP7/PP7L through its PMD and PPP domains, respectively.

Considering that the PMD proteins MAIN and MAIL1 interact with PP7L, and are required for the

expression of similar set of loci, we decided to determine the distribution of related genic PMD and

PPP domains, and to retrace their evolutionary history among plant species. The *A. thaliana* MAIN,

MAIL1 and MAIL3 are all members of the PMD-C family that also includes MAIL2 [17]. Since our

objective is to retrace the evolution of genic (and not TE-containing) PMD-C, we have decided to

restraint our search to Eudicots. Indeed, Eudicot species contain mainly genic PMD-C, while other

angiosperms may contain variable numbers of closely related genic and TE-associated PMD-C motifs

that would be difficult to distinguish in our analysis. To retrace the evolution history of the genic PMD-

C family, we used *A. thaliana* PMD-C genes to search and collect their relatives (paralogues and

orthologues) in 30 genomes representative of the Eudicot diversity (see S4 Table for a list of species

and their corresponding codes used in Fig 4, and S5 Table for motif sequences).

208   In our phylogenetic analysis, the genic PMD-C family can be clearly separated in two major

clades. The first clade is composed of orthologues of *A. thaliana* MAIL2, MAIL1 and MAIN, while the

second one includes orthologues of *A. thaliana* MAIL3 (Fig 4A). Notably, a MAIL2 gene was found in

all species tested. MAIL2 genes form a closely related group suggesting that they are under strong

purifying selection (see the very short branch lengths linking most MAIL2 genes in Fig 4A). Additional

MAIL2 paralogues were also detected in several species. Some MAIL2 paralogues are imbedded in the

major MAIL2 group while others are more divergent and form independent subgroups. This is the case

of MAIL1 and MAIN that are clearly Brassicaceae-specific MAIL2 paralogues. MAIL1 and MAIN diverge

significantly from Brassicaceae MAIL2 following a gene duplication event that likely occurred in the

common ancestor of Brassicaceae, before falling under strong purifying selection, a situation often

encountered in neofunctionalization events. The MAIL3 distribution is irregular compared to MAIL2,

as a MAIL3 orthologue was not found in all Eudicot species tested. Accordingly, except in Brassicaceae,

MAIL3 genes appear to be under much weaker purifying selection compare to *MAIL2* and *MAIL2-like*

genes (see the longer branch lengths in the tree of Fig 4A). Brassicaceae *MAIL3* genes contrast with

other *MAIL3*, by forming a closely related group in the phylogenetic tree. This suggests a clear change

in selection pressure, typical of a neofunctionalization event, this time correlated with the acquisition

of the PPP motif by these genes (Fig 4B and see below). Remarkably, another fusion event between a

PMD-C and a PPP motif occurred independently in grapevine, but this time involving a MAIL2

paralogue (VvMAIL2.2, Fig 4A).

227   We then used the PP7/PP7L-related motif found in *A. thaliana* MAIL3, to collect orthologous

genes and retrace the evolution history of PP7/PP7L in the same Eudicot species used above. We

confirmed by this analysis that these PPP genes can be clearly separated in two distinct clades: PP7

and PP7-like (PP7L) (Fig 4B). All tested species present one or several closely related PP7 paralogues.

Although Brassicaceae MAIL3 PPP motifs belong to the PP7 clade, it diverged significantly compared

to Brassicaceae or any other standalone PP7 paralogues. This suggests a fast-evolving period for

Brassicaceae MAIL3 PP7 motifs subsequently to their fusion with the PMD-C motif. This is in complete

agreement with the apparent change of selection regime that occurred for the MAIL3 PMD motif

following its fusion with PP7 (Fig 4A). It is interesting to note that in grapevine, the PPP motif found

on the same protein as the MAIL2 PMD-C is also of the PP7 type, and is quite divergent compared to

the other grapevine standalone PP7 and MAIL2 domains (VvMAIL2.2 and VvPP7.2 in Fig 4A and B,

respectively). Finally, PP7L orthologues were not found in all species tested and, accordingly, these

genes appear to be under weaker purifying selection compare to genes belonging to the PP7

subfamily. However, this was less obvious among the Brassicaceae clade, where the PP7L domain

appears highly conserved (see the branch lengths in the tree of Fig 4B). In conclusion, phylogenetic

analyses showed that, in at least two distinct events, neo-association of PMD-C and PP7 domains have

potentially create a new protein function that was maintained through evolution.

DISCUSSION

In *A. thaliana,* MAIN and MAIL1 are standalone PMD proteins that have been involved in

genome integrity, regulation of cell division and differentiation, and silencing of TEs [17-19]. In this

study, we showed that TE silencing is widely impaired in the *ddc main* higher order mutant, which is

both partially defective in DNA methylation and MAIN activity. We also identified the putative

phosphatase protein PP7L as a MAIN and MAIL1 protein interactor, and showed that most of the genes

that are commonly misregulated in *main*, *mail1* and *pp7l* mutants share identical CRE motifs. This

suggests that transcription at these loci might be regulated through similar mechanisms involving

specific CRE recognition. Finally, phylogenetic analyses among Eudicots suggest mechanism of

neofunctionalization between the PMD and PP7-type PPP, to potentially acquire a functional module

that requires the two protein domains.

The PMD MAIN protein acts independently of DRM1/2- and CMT3-mediated DNA methylation

pathways to silence TEs.

Previous analyses showed that some TEs were synergistically upregulated in the *mail1 rdr2*

double mutant plants, suggesting that MAIL1 acts independently of RdDM pathway [17]. In our whole

genome transcriptomic analyses, we showed that several TEs are upregulated in both *main-3* and *ddc*

mutants, and their upregulation is dramatically exacerbated in the *ddc main-3* quadruple mutant (Fig

1 E and F, red color code). We also identified TEs that are only upregulated in *ddc* or *main-3* mutants,

but display stronger misregulation in the *ddc main-3* higher order mutant (Fig 1 E and F, blue and

yellow color codes). Finally, we identified a large number of TEs that are only overexpressed in *ddc*

main-3 (Fig 1 E and F, green color code). Thus, the PMD MAIN, DRM1/2 and CMT3 pathways act

synergistically at a subset of TEs (Fig 1 E, blue, red and yellow color codes), but also operate

redundantly to efficiently silence a much larger set of TEs that are predominantly pericentromeric (Fig

1 E, green color code and S1H Fig). Previous work showed that DNA methylation is not impaired in

mail1 [17]. Likewise, we found that DNA methylation is largely unaffected in the *main* null mutant

(S3E Fig). Considering the strong synergistic effect of *main* and *ddc* mutations on TE derepression, we

can exclude the hypothesis that MAIN act downstream of DRM1/2- or CMT3-mediated DNA

methylation (no epistasis). Instead, it is very likely that MAIN, DRM1/2 and CMT3 pathways act

independently to efficiently silence TEs. Notably, synergistic effects between different epigenetic

pathways have already been observed. For instance, it has been shown that MORC1/6 and MOM1

pathways, or RdDM and MOM1 pathways act synergistically and redundantly to efficiently silence TEs

[14, 29]. Altogether, these observations contribute to the "mille-feuille" (i.e. "multiple layers") model,

in which different epigenetic pathways converge towards the silencing of TEs [30].

The putative phosphatase PP7L interacts with the PMD MAIN and MAIL1 protein to regulate a

similar set of genes and TEs.

Recently, PP7L was involved in the biogenesis of chloroplasts and plant response upon abiotic

stress [27]. The *pp7l* mutant shows dramatic developmental phenotype, which correlates with

misregulation of gene expression and upregulation of TEs (S3A-C Fig) [27]. Using IPMS and co-IP

experiments, we first showed that MAIN and MAIL1 interact together, which was already suggested

based on transcriptomic analyses of *main mail1* double mutant plants (Fig 2A and B) [17]. In addition,

we identified PP7L, which is a putative PPP, as a robust interactor of MAIN and MAIL1, suggesting that

the three proteins form heteromers (Fig 2A, C and D).

Comparison of upregulated TEs and misregulated genes in *main*, *mail1* and *pp7l* mutants revealed

that a significant proportion of loci that are misregulated in the two *pmd* mutants are also

misregulated in *pp7l* (Fig 3A-F, S2D and S2F). This suggests that the three proteins act, at least partially,

in the same pathway, which is consistent with the fact that they interact together.

Genes that are commonly downregulated in *main, mail1* and *pp7l* carry an identical DNA motif in

their promoter regions.

In silico analyses identified a specific DNA motif (5'-CATGCCGTTT-3') among the promoter of genes

that are commonly downregulated in *main*, *mail1* and *pp7l* mutants (Fig 3 G). this DNA motif was also

strongly enriched in most of the downregulated genes in the hypomorphic *main-3* mutant*,* and in

approximately half of the downregulated genes in *main-2* and *mail1-1* null mutant (S3G Fig). Based on

these results, we hypothesize that the MAIN/MAIL1/PP7L protein complex would be involved in

cellular processes that positively regulate transcription through the recognition of the putative CRE

motif 5'-CATGCCGTTT-3'. Consistently with this hypothesis, the highly similar DNA pentamer 5'-

TGACGT-3' was previously described as a G-box-like DNA element specifically recognized by

basic/leucine zipper (bZIP) TFs [31]. Therefore, it is likely that either a bZIP-like TF is recruited through

the MAIN/MAIL1/PP7L pathway, or the PMDs/PP7L complex directly recognizes the DNA motif to

promote transcription. In the future, it will be important to test if the MAIN/MAIL1/PP7L protein

complex interact directly with chromatin through the specific recognition of CRE motif. It will also be

essential to determine if this protein complex is required for the activation of specific TFs that bind

the CRE motif to regulate transcription.

The association of PMD-C and PP7/PP7L domains creates a functional protein module.

In this study, we identified PP7L has a protein partner of the two standalone PMDs MAIN and

MAIL1, and showed that *main*, *mail1* and *pp7l* display similar mutant phenotypes (Figs 2, 3 and S3). In

addition, phylogenetic analyses suggest that the association in trans of Brassicaceae PMD

MAIN/MAIL1 with PP7L would have created a selection pressure to optimize the MAIN/MAIL1/PP7L

interaction and function within the plant species. Indeed, both Brassicaceae MAIN/MAIL1 and PP7L

are clearly divergent compared to their closest orthologues and present much shorter branch lengths,

suggesting that a new correlated selection regime resulted from their cooperative functions in this

lineage (Fig 4). Besides, we showed that PMD and PP7 domains are fused in several Brassicaceae

MAIL3 and in the grapevine VvMAIL2.2 proteins (Fig 4). We propose a model in which the MAIL3-

specific PMD/PP7 fusion found in several Brassicaceae would have occurred in the common ancestors

of this Eudicot lineage, leading to a transient fast-evolving period for both motifs (possibly under a

positive selection regime). Since then, the MAIL3 PMD/PP7 fusion was maintained under strong

purifying selection, arguing for a neofunctionalization of the fusion protein. It is likely that similar

process happened in grapevine, and possibly, in closely related Vitaceae species. To some extent, the

two distinct events that occurred in Brassicaceae and grapevine are reminiscent of convergent

evolution processes leading to the production of a functional PMD/PP7 module, in which the two

protein domains would interact in cis. Although the Brassicaceae MAIL3 and grapevine VvMAIL2.2

proteins are probably involved in distinct biological processes, we hypothesize that, like the

MAIN/MAIL1/PP7L complex, they may interact with chromatin at specific genomic locations to

regulate gene expression.

In conclusion, we show here that the two *A. thaliana* PMD MAIN and MAIL1 proteins interact

with PP7L, and are involved in the regulation of a common set of genes and TEs. This association in

trans would have created a selection pressure on PP7L to optimize its interaction with MAIN and

MAIL1. In the future, it will be important to test if the MAIN/MAIL1/PP7L protein complex interact

with chromatin. In addition, we showed that distinct events of PMD-C and PP7 fusions occurred

among the Eudicots (among several Brassicaceae species and in grapevine), suggesting some

convergent evolution processes and a potential neofunctionalization of PMD/PP7 module in cis. In the

future, it will be essential to determine the role of these PMD/PP7 fusion proteins, such as the *A.*

thaliana MAIL3. In addition, it will be important to decipher the function of PMD proteins in plant

species with agronomic value.

MATERIALS AND METHODS

Plant material and growing conditions. Wild-type (WT) and all mutant lines are in the Columbia (Col)

ecotype. The *drm1-2* (SALK_031705), *drm2-2* (SALK_150863) *cmt3-11* (SALK_148381) *ddc* triple

mutant line was previously described [28]. The *main-2* (GK-728H05), *mail1-1* (GK-840E05) and *pp7l-2*

(SALK_003071) were previously described [17-19, 27], and obtained from The Nottingham Arabidopsis

Stock Centre. T-DNA insertions were confirmed by PCR-based genotyping. The *main-3* mutation was

genotyped by derived cleaved amplified polymorphic sequences (dCAPS) using the restriction enzyme

FokI. Primer sequences are described in S6 Table. Plants were grown on soil under a 16h-light/8h-dark

cycle or *in vitro* under continuous light and transferred on soil after 10 days. For *in vitro* plant culture,

seeds were surface-sterilized and sowed on solid Murashige and Skoog medium containing 0.5%

sucrose (w/v).

Cloning of ATCOPIA28::GFP. The pCambia3300-NLS-GFP-T35S vector was previously described [13].

The 5'LTR promoter corresponding to a region of ~1 kb upstream of *ATCOPIA28 (AT3TE51900)* was

PCR amplified from WT genomic DNA, and cloned into pCR2.1 TOPO vector (Invitrogen). Quikchange

site-directed mutagenesis (Stratagene) was performed according to Manufacturer's instruction to

create a polymorphism site (MfeI→NdeI) within the 5'LTR promoter, which was subsequently

mobilized into pCambia3300 upstream of NLS-GFP-T35S sequence. *ddc* triple mutant plants were

transformed with the AT*COPIA28::GFP* construct using the *Agrobacterium*-mediated floral dip

method [32]. Transgenic plants showing GFP fluorescence were backcrossed with a WT plant to

promote the silencing of *ATCOPIA28::GFP* in the F1 generation. F1 plants were self-crossed and their

F2 progenies were screened for GFP fluorescence, and PCR-based genotyped to obtain

ATCOPIA28::GFP WT and *ATCOPIA28::GFP ddc* plants. Primer sequences used for *ATCOPIA28::GFP*

cloning are described in S6 Table.

EMS mutagenesis, GFP screening and mapping analyses. Five thousand seeds of *ATCOPIA28::GFP ddc*

were mutagenized in 0.26% EMS solution for 12 hours with rotation. Seeds were subsequently washed

with water and sown on soil. Fifteen hundred M2 populations were collected, and subsequently

screened for GFP fluorescence under UV light using a SMZ18 Nikon Fluorescence Stereomicroscope

coupled with the C-HGFI intensilight fluorescence filter. Pictures were taken using the DS Qi1MC digital

camera kit. Mapping and identification of the EMS mutation responsible for the phenotype were

performed by bulk segregant analysis coupled with deep genome re-sequencing as previously

described [13], with the following differences. Reads were mapped against the reference genome

(*Arabidopsis* TAIR10) and single nucleotide polymorphisms called in Geneious (Biomatters). Using R,

single nucleotide polymorphisms were filtered for EMS mutations (G:C→A:T) and zygosity called

based on the variant frequency provided by Geneious (≥80% homozygous mutation, ≥45%, and ≤55%

heterozygous mutation). Plots were then created by calculating the ratio of the number of

homozygous and heterozygous and mutations in a 500-kb window as previously described [33].

Cloning of epitope-tagged versions of PMD proteins. *MAIN* and *MAIL1* genomic regions were PCR amplified and FLAG or Myc epitopes were added to the C-terminus of each protein as previously described [13]. In both cases, the amplified region includes a ~1Kb promoter sequence upstream of the respective transcriptional start site. For the *MAIN* promoter, a MluI site was modified to allow LR reaction without changing the sequence integrity of the gene. Primer sequences are described in S6 Table.

IP and MS analysis. Ten grams of 3-week-old seedling tissue were ground in liquid nitrogen and resuspended in 50mL ice-cold IP buffer [50mM Tris HCl pH 7.6, 150mM NaCl, 5mM MgCl$_2$, 0.1% Nonidet P-40, 10% glycerol (v/v), 0.5mM DTT, 1x Protease Inhibitor Mixture (Roche)] and centrifuged 2 times for 15 min at 4°C at 15 350g. 400μL of M2 magnetic FLAG-beads (Sigma, M8823) were added to the supernatants, and incubated for 90min rotating at 4°C. M2 magnetic FLAG-beads were washed seven times in ice-cold IP buffer for 5 min rotating at 4°C, and immunoprecipitated proteins were eluted 3 times with 150μL 3x-FLAG peptides (Sigma, F4799) for 25 min each at 25°C. The eluted protein complexes were precipitated by trichloroacetic acid and subjected to MS analyses as previously described [14].

Co-IP and immunoblotting. 0.5 g of 3-week-old seedling tissue were ground in liquid nitrogen, resuspended in 1.5mL ice-cold IP buffer [50mM Tris pH 7.6, 150mM NaCl, 5mM MgCl2, 0.1% Nonidet P-40, 10% glycerol, 0.5 mM DTT, 1x Protease Inhibitor Mixture (Roche)], and centrifuged 2 times for 15 min at 4°C, 16 000g. 50μL M2 magnetic FLAG-beads (Sigma, M8823) were added to the supernatants and incubated for 2 hour rotating at 4°C. Beads were washed 3 times in ice-cold IP buffer for 10 min rotating at 4°C. Immunoprecipitated proteins were denatured in Laemmli buffer for 5min at 95°C. 10μL of input and bead elution were run on 10% SDS-PAGE gels, and proteins were detected by western blotting using either Anti-FLAG M2 monoclonal antibody-peroxidase conjugate (Sigma,

A8592) at a dilution of 1 :10 000, or c-Myc rat monoclonal antibody (Chromotek, 9E1-100) at a dilution

of 1 : 1000 followed by goat anti-rat IgG horseradish peroxidase (Abcam, ab205720) used at a dilution

of 1 :20000 as secondary antibody. Western blots were developed using Substrat HRP Immobilon

Western (Merck Millipore, WBKLS0500).

RNA extraction. Total RNA was extracted from aerial parts of 3-week-old seedlings using either

RNeasy Plant Mini Kit (Qiagen, 74904) or Monarch Total RNA Miniprep Kit (NEB, T2010) according to

the manufacturer's protocols.

RNA sequencing. RNA-seq libraries were generated from 1µg of input RNA using NEBNext Ultra II

Directional RNA Library Prep Kit for Illumina (NEB, E7490) according to the manufacturer's protocols.

Libraries were sequenced on an Illumina HiSeq 4000 or NextSeq 550 machines. Reads were trimmed

using Trimmomatic [34], and mapped to the *A. thaliana* genome (*Arabidopsi*s TAIR10 genome) using

HISAT2 [35]. The sequence alignment files were sorted by name and indexed using SAMtools [36].

Files were converted to BAM files and number of reads mapped onto a gene calculated using HTSeq-

count [37]. Differentially expressed genes were obtained with DESeq2 [38], using a log2 fold-change

> 2 (up-regulated genes) or < -2 (down-regulated genes) with an adjusted *p*-value of 0,01. Re-analyses

of previously published RNA-seq datasets from *main-2* and *mail1-1* (PRJEB15202) [17], and *pp7l*

mutants (GSE122495) [27] were performed as described above.

RT-qPCR. 1 µg of input RNA was converted to cDNA using GoScript Reverse Transcriptase (Promega

A501C) according to the manufacturer's protocol. The final reaction was diluted 6 times with RNase

free water. RT-qPCR experiments were performed with 4µL of cDNA combined to the Takyon No Rox

SYBR MasterMix (Eurogentec, UF-NSMT-B0701), using a LightCycler 480 instrument (Roche).

Amplification conditions were as follows: 95°C 5 min; 45 cycles, 95°C 15s, 60°C 15s, 72°C 30s; melting

curves. Three technical replicates were performed per biological replicate, and 3 biological replicates

were used in all experiments.

Cis-elements detection. The motifs for enhancer sequences (1kb upstream the TSS) were discovered

using MEME (Multiple Em for Motif Elicitation). MEME represents motifs as position-dependent

letter-probability matrices which describe the probability of each possible letter at each position in

the pattern [39].

Bisulfite sequencing. Genomic DNA was extracted from aerial parts of 3-week-old seedlings using

Quick-DNA Plant/Seed Miniprep Kit (Zymo research, D6020) according to the manufacturer's protocol.

Whole genome bisulfite sequencing (WGBS) library was prepared from 50 ng genomic DNA using

NuGen Ovation Ultralow Methyl-Seq kit. Bisulfite treatment was carried out by Qiagen Epitect bisulfite

kit. WGBS libraries were sequenced on an Illumina HiSeq 4000 machine. The raw reads (single end)

were trimmed using Trimmomatic in order to remove adapter sequences [34]. The remaining

sequences were aligned against the *A. thaliana* genome TAIR10 version using Bismark [40]. Duplicated

reads were collapsed into one read. For visualization of the data, we used ViewBS [41].

Sequence selection, multiple sequences alignments and phylogenetic reconstruction.

Blast searches (blastp) were performed starting from known *A. thaliana* PMD-C and PP7/PP7L motifs

on the thirty species representing the diversity of the Eudicot lineages. When necessary tblastn

searches were also used to obtain complete protein sequences. To build the phylogenetic trees, PMD-

C or PP7/PP7L motifs were aligned using the multiple sequence comparison by log-expectation

(MUSCLE v3.7) software [42]. Trees were reconstructed using the fast-maximum likelihood tree

estimation program PHYML [43] using the LG amino acids replacement matrix [44]. Statistical support

for the major clusters were obtained using the approximate likelihood-ratio test (aLRT) [45].

Data availability. Nucleotide sequencing data generated in this study have been deposited in

European Nucleotide Archive (ENA) under the accession number PRJEB33240

(http://www.ebi.ac.uk/ena/data/view/PRJEB33240). The proteomics data have been deposited to the

MassIVE data repository (https://massive.ucsd.edu) with the dataset identifier MSV000084089. All

other data and material are available within the manuscript and its supplementary files, or from the

corresponding author upon request.

ACKNOWLEDGMENTS

The authors want to thank Thierry Lagrange, Frederic Pontvianne and other team members for fruitful

discussions, and all the LGDP platform members for their outstanding technical assistance and plant

care. This study is set within the framework of the "Laboratoires d'Excellences (LABEX)" TULIP (ANR-

10-LABX-41). S.E.J. is an investigator of the Howard Hughes Medical Institute.

REFERENCES

474
1. Grewal SI, Jia S. Heterochromatin revisited. Nat Rev Genet. 2007;8(1):35-46.
2. Slotkin RK, Martienssen R. Transposable elements and the epigenetic regulation of the
genome. Nat Rev Genet. 2007;8(4):272-85.
3. Deniz O, Frost JM, Branco MR. Regulation of transposable elements by DNA modifications.
Nat Rev Genet. 2019;20(7):417-31.
4. Dergai O, Hernandez N. How to Recruit the Correct RNA Polymerase? Lessons from snRNA
Genes. Trends Genet. 2019;35(6):457-69.
5. Law JA, Jacobsen SE. Establishing, maintaining and modifying DNA methylation patterns in
plants and animals. Nat Rev Genet. 2010;11(3):204-20.
6. Matzke MA, Mosher RA. RNA-directed DNA methylation: an epigenetic pathway of
increasing complexity. Nat Rev Genet. 2014;15(6):394-408.
7. Wendte JM, Pikaard CS. The RNAs of RNA-directed DNA methylation. Biochim Biophys Acta.
2016.
8. Du J, Johnson LM, Jacobsen SE, Patel DJ. DNA methylation pathways and their crosstalk with
histone methylation. Nat Rev Mol Cell Biol. 2015;16(9):519-32.
9. Zhang H, Lang Z, Zhu JK. Dynamics and function of DNA methylation in plants. Nat Rev Mol
Cell Biol. 2018;19(8):489-506.
10. Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, et al. The
Arabidopsis nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing
heterochromatin. Cell. 2013;153(1):193-205.
11. Stroud H, Do T, Du J, Zhong X, Feng S, Johnson L, et al. Non-CG methylation patterns shape
the epigenetic landscape in Arabidopsis. Nat Struct Mol Biol. 2014;21(1):64-72.

12.   Amedeo P, Habu Y, Afsar K, Mittelsten Scheid O, Paszkowski J. Disruption of the plant gene
MOM releases transcriptional silencing of methylated genes. Nature. 2000;405(6783):203-6.
13.   Moissiard G, Cokus SJ, Cary J, Feng S, Billi AC, Stroud H, et al. MORC family ATPases required
for heterochromatin condensation and gene silencing. Science. 2012;336(6087):1448-51.
14.   Moissiard G, Bischof S, Husmann D, Pastor WA, Hale CJ, Yen L, et al. Transcriptional gene
silencing by Arabidopsis microrchidia homologues involves the formation of heteromers. Proc Natl
Acad Sci U S A. 2014;111(20):7474-9.
15.   Lorkovic ZJ, Naumann U, Matzke AJ, Matzke M. Involvement of a GHKL ATPase in RNA-
directed DNA methylation in Arabidopsis thaliana. Curr Biol. 2012;22(10):933-8.
16.   Gallego-Bartolome J, Liu W, Kuo PH, Feng S, Ghoshal B, Gardiner J, et al. Co-targeting RNA
Polymerases IV and V Promotes Efficient De Novo DNA Methylation in Arabidopsis. Cell.
2019;176(5):1068-82 e19.
17.   Ikeda Y, Pelissier T, Bourguet P, Becker C, Pouch-Pelissier MN, Pogorelcnik R, et al.
Arabidopsis proteins with a transposon-related domain act in gene silencing. Nat Commun.
2017;8:15122.
18.   Wenig U, Meyer S, Stadler R, Fischer S, Werner D, Lauter A, et al. Identification of MAIN, a
factor involved in genome stability in the meristems of Arabidopsis thaliana. Plant J. 2013;75(3):469-
83.
19.   Uhlken C, Horvath B, Stadler R, Sauer N, Weingartner M. MAIN-LIKE1 is a crucial factor for
correct cell division and differentiation in Arabidopsis thaliana. Plant J. 2014;78(1):107-20.
20.   Babu MM, Iyer LM, Balaji S, Aravind L. The natural history of the WRKY-GCM1 zinc fingers
and the relationship between transcription factors and transposons. Nucleic Acids Res.
2006;34(22):6505-20.
21.   Steinbauerova V, Neumann P, Novak P, Macas J. A widespread occurrence of extra open
reading frames in plant Ty3/gypsy retrotransposons. Genetica. 2011;139(11-12):1543-55.
22.   Farkas I, Dombradi V, Miskei M, Szabados L, Koncz C. Arabidopsis PPP family of
serine/threonine phosphatases. Trends Plant Sci. 2007;12(4):169-76.
23.   Sun X, Kang X, Ni M. Hypersensitive to red and blue 1 and its modification by protein
phosphatase 7 are implicated in the control of Arabidopsis stomatal aperture. PLoS Genet.
2012;8(5):e1002674.
24.   Liu HT, Li GL, Chang H, Sun DY, Zhou RG, Li B. Calmodulin-binding protein phosphatase PP7 is
involved in thermotolerance in Arabidopsis. Plant Cell Environ. 2007;30(2):156-64.
25.   Genoud T, Santa Cruz MT, Kulisic T, Sparla F, Fankhauser C, Metraux JP. The protein
phosphatase 7 regulates phytochrome signaling in Arabidopsis. PLoS One. 2008;3(7):e2699.
26.   Uhrig RG, Labandera AM, Moorhead GB. Arabidopsis PPP family of serine/threonine protein
phosphatases: many targets but few engines. Trends Plant Sci. 2013;18(9):505-13.
27.   Xu D, Marino G, Klingl A, Enderle B, Monte E, Kurth J, et al. Extrachloroplastic PP7L Functions
in Chloroplast Development and Abiotic Stress Tolerance. Plant Physiol. 2019.
28.   Stroud H, Greenberg MV, Feng S, Bernatavichute YV, Jacobsen SE. Comprehensive analysis
of silencing mutants reveals complex regulation of the Arabidopsis methylome. Cell. 2013;152(1-
2):352-64.
29.   Yokthongwattana C, Bucher E, Caikovski M, Vaillant I, Nicolet J, Mittelsten Scheid O, et al.
MOM1 and Pol-IV/V interactions regulate the intensity and specificity of transcriptional gene
silencing. Embo J.29(2):340-51.
30.   Rigal M, Mathieu O. A "mille-feuille" of silencing: epigenetic control of transposable
elements. Biochim Biophys Acta. 2011;1809(8):452-8.
31.   Schindler U, Beckmann H, Cashmore AR. TGA1 and G-box binding factors: two distinct
classes of Arabidopsis leucine zipper proteins compete for the G-box-like element TGACGTGG. Plant
Cell. 1992;4(10):1309-19.
32.   Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J. 1998;16(6):735-43.

33.   Hristova E, Fal K, Klemme L, Windels D, Bucher E. HISTONE DEACETYLASE6 Controls Gene
Expression Patterning and DNA Methylation-Independent Euchromatic Silencing. Plant Physiol.
2015;168(4):1298-308.
34.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20.
35.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
requirements. Nat Methods. 2015;12(4):357-60.
36.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
37.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput
sequencing data. Bioinformatics. 2015;31(2):166-9.
38.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15(12):550.
39.   Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, et al. MEME SUITE: tools for
motif discovery and searching. Nucleic Acids Res. 2009;37(Web Server issue):W202-8.
40.   Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics. 2011;27(11):1571-2.
41.   Huang X, Zhang S, Li K, Thimmapuram J, Xie S, Wren J. ViewBS: a powerful toolkit for
visualization of high-throughput bisulfite sequencing data. Bioinformatics. 2018;34(4):708-9.
42.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res. 2004;32(5):1792-7.
43.   Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies
by maximum likelihood. Syst Biol. 2003;52(5):696-704.
44.   Le SQ, Gascuel O. An improved general amino acid replacement matrix. Mol Biol Evol.
2008;25(7):1307-20.
45.   Anisimova M, Gascuel O. Approximate likelihood-ratio test for branches: A fast, accurate,
and powerful alternative. Syst Biol. 2006;55(4):539-52.

FIGURE CAPTIONS

Fig 1. MAIN is required for the silencing of TEs, acting redundantly and synergistically with the

DRM1/2 and CMT3 pathways.

(A) WT and *drm1 drm2 cmt3* (*ddc*) triple mutant plants carrying the *ATCOPIA28::GFP* transgene

showed no and weak GFP fluorescence under UV light, respectively. By comparison, the *ddc #16* EMS

mutant showed strong GFP fluorescence. Insets show plants under white light. (B) Western blot using

anti-GFP antibody confirmed *ATCOPIA28::GFP* overexpression in *ddc #16*. Coomassie staining of the

large Rubisco subunit (rbcL) is used as a loading control. (C) Relative expression analyses of

ATCOPIA28::GFP (GFP) and three endogenous TEs in *ddc* and *ddc #16 over WT* assayed by Real-Time

quantitative PCR (RT-qPCR), and normalized to the house keeping gene *RHIP1*. Error bars indicate

standard deviation based on three independent biological replicates. (D) Number of upregulated TEs

in *ddc*, *main-3* and *ddc main-3*, and classified by TE superfamily. Threshold: log2>2, p-adj< 0,01.

Analyses are based on four biological replicates. (E) Venn diagram analyses showing the overlaps

between upregulated TEs in *ddc*, *main-3* and *ddc main-3*. (F) Relative expression analyses of several

TEs in *ddc*, *main-3* and *ddc main-3* over WT assayed by RT-qPCR and normalized to *RHIP1*. Error bars

indicate standard deviation based on three independent biological replicates. The same color code

was used for the Venn diagram (E) and the frames of RT-qPCR graphs (F).

Fig 2. MAIN and MAIL1 physically interacts together and with PP7L.

(A) FLAG-tagged MAIN and MAIL1 proteins were immunoprecipitated and putative interacting

proteins were identified by mass spectrometry. The total number of identified spectra is shown for

three MAIN and two MAIL1 biological replicates. WT replicates are used as a negative control. (B)

MAIL1-MYC was co-immunoprecipitated with MAIN-FLAG in F1 plants co-expressing the two

transgenes. Plants expressing only MAIL1-MYC or MAIN-FLAG were used as negative controls. (C)

Plants expressing the MAIN-MYC transgene were "supertransformed" with the PP7L-FLAG construct,

and MAIN-MYC was co-immunoprecipitated with PP7L-FLAG. Plants expressing only MAIN-MYC or

PP7L-FLAG were used as negative controls. (D) Same as C but using MAIL1-MYC plants

"supertransformed" with the PP7L-FLAG construct. Epitope-tagged proteins were detected by

Western blotting. Arrowheads indicates expected bands. Asterisks indicates non-specific

hybridization.

Fig 3. The PMD proteins and PP7L regulates a same set of loci.

(A-C) Venn diagrams showing the overlaps between upregulated TEs (A), downregulated genes (B) and

upregulated genes (C) in *main-2*, *mail1-1* and *pp7l-1* based on analyses described in S2 and S3 Figs.

(D-E) Relative expression analyses assayed by RT-qPCR of upregulated TEs (D), downregulated genes

and (E) upregulated genes in the *main-2* and *pp7l-2* mutants over WT. Transcript level is normalized

to *RHIP1*. Error bars indicate standard deviation based on three independent biological replicates. (G)

Overrepresentation of a specific DNA motif in the 1-kb promoter regions of genes that are commonly

downregulated in *main*, *mail1* and *pp7l* using the MEME software. Under the logo are represented

the ratio of genes carrying the motif among the list of downregulated loci, and the significant E-value

associated with this motif.

Fig 4. Evolutionary history of PMD-C and PP7 proteins in plants.

(A) An alignment of the PMD-C motifs from 30 representative Eudicot species was used to construct

a phylogenetic tree. The two major clades (MAIL2/MAIL2-like and MAIL3) are indicated. The species

codes are given in S4 Table, and corresponding protein sequences in S5 Table). In red are genes

presenting a fusion between a PMD-C and a PP7 motif. Statistical supports of key nodes calculated

with the approximate likelihood-ratio test are indicated. Scale bar indicates one substitution/site. The

tree was rooted using the *Amborella trichopoda* PMD-C motif (Atr1PMDC). (B) Phylogenetic tree

constructed using an alignment of the PP7 motif from the same species as in (A). The two major clades

(PP7 and PP7L) are indicated. In red are genes presenting a fusion between a PP7 and a PMD-C motif.

Statistical supports of key nodes calculated with the approximate likelihood-ratio test are indicated.

Scale bar indicates one substitution/site. The tree was rooted using the *A. thaliana* PP5 motif (AtPP5).

629
630

 SUPPORTING INFORMATION CAPTIONS

 S1 Fig. *MAIN* is the mutated gene responsible for *ATCOPIA28::GFP* and TE overexpression in the *ddc*

 ***#16* mutant.**

 (A) Schematic representation of the *ATCOPIA28::GFP* construct. The 5' long terminal repeat (LTR)

 promoter region of an ATCOPIA28 LTR-retrotransposon (*AT3TE51900*) is used to control the

 expression of GFP. The construct also carries a Nuclear Localization Signal (NLS) to target the GFP in

 the nucleus. (B) Representative pictures of WT, *ddc* and *ddc #18* (*ddc morc6-8)* and *ddc #344* (*ddc*

 morc6-9) mutant plants under UV light. Insets show plants under white light. (C) Enrichment in

 homozygote/heterozygote ratio of EMS over WT single nucleotide polymorphisms (SNPs), defining

 the linkage intervals for the populations *ddc #18*, *ddc #344* and *ddc #16* (*ddc main-3)*. (D) Location of

 the point mutations corresponding to the *morc6-8 and morc6-9* alleles within the *MORC6* genomic

 sequence. (E) Location of the point mutation corresponding to the *main-3* allele within the *MAIN*

 genomic sequence. Nucleotide and corresponding amino acid changes are indicated above the gene.

 Positions of the mutations are indicated relative to the transcription starting site (+1). Grey boxes

 represent 5' and 3' UTR, blue boxes and lines represent exons and introns, respectively. (F) Genetic

 complementation analysis using the KO T-DNA insertion line *main-2*. *ddc #16* plants were crossed with

 main-2 plants, F1 plants were self-crossed and F2 plants were screened under UV light to analyze GFP

 fluorescence. Western blotting using anti-GFP antibodies confirmed GFP overexpression in selected

 F2 plants. Coomassie staining of the large Rubisco subunit (rbcL) is used as a loading control. The

 DRM2 and *CMT3* genotype of selected F2 plants were subsequently analyzed by PCR using specific

 primers. Ho: Homozygous. He: Heterozygous. All the F2 plants that were selected were homozygote

 for the *drm2* mutation. (G) Representative pictures showing the developmental phenotype of 3-week-

 old WT, *ddc*, *main-3* and *ddc main-3* plants. (H) Chromosomal distributions of misregulated loci in *ddc*,

 main-3 and *ddc main-3* over WT. Chromosome arms are represented in light grey, pericentromeric

regions in dark grey. Upregulated genes are represented in blue, downregulated genes in green and

TEs are represented in red above the chromosomes (upregulated) or below (downregulated). The size

of each bar is proportional to the misregulation of the loci. Threshold: log2>2, or log2<-2; p-adj< 0,01.

(I) Relative expression analyses of *ATCOPIA28* and *HELITRONY1D* (*AT5TE35950*) in *ddc*, *main-3* and *ddc*

main-3 over WT assayed by RT-qPCR. Relative transcript level is normalized to *RHIP1*. Error bars

indicate standard deviation based on three independent biological replicates.

S2 Fig. Epitope-tagged MAIN and MAIL1 constructs complement the developmental and molecular

phenotypes of *main-2* and *mail1-1* null mutants.

(A) Representative pictures of 2-week-old WT, *main-2*, *mail1-1* and epitope-tagged complementing

mutant plants. (B) Number of upregulated TEs in *main-2* and *mail1-1*, classified by TE superfamily. (C)

Number of misregulated genes in *main-2* and *mail1-1* classified by upregulated and downregulated

genes. Threshold: log2>2, or log2<-2; p-adj< 0,01. These analyses are based on four biological

replicates, including the reanalysis of RNA-seq data published in reference [17]. (D) Relative

expression analyses of upregulated genes (above panel) and downregulated genes (below panel) in

main-2 and *mail1-1* over WT assayed by RT-qPCR. Relative transcript level is normalized to *RHIP1*.

Error bars indicate standard deviation based on three independent biological replicates. (E) Western

blots using anti-FLAG and anti-MYC antibodies confirmed the accumulation of epitope-tagged PMD

proteins at the expected sizes in the different complementing lines. (F) Relative expression analyses

of several TEs in the different complementing lines over WT assayed by RT-qPCR and normalized to

RHIP1. The *main-2* and *mail1-1* mutants are used as negative controls. Error bars indicate standard

deviation based on three independent biological replicates.

S3 Fig. Developmental and molecular phenotypes of *pp7l* mutant.

(A) Representative pictures of 5-week-old *main-2*, *mail1-1* and *pp7l-2* mutant plants showing similar

abnormal developmental phenotypes in comparison to WT plant. (B) Number of upregulated TEs in

pp7l-1 and classified by TE superfamily. (C) Number of misregulated genes in *pp7l-1* classified by

upregulated and downregulated genes. Threshold: log2>2, or log2<-2; p-adj< 0,01. These analyses are

based on reanalysis of RNA-seq data published in reference [27]. (D-E) Metaplot analyses showing the

average distribution of DNA methylation levels in upregulated TEs (D) and genes (E) in *main-2* versus

WT plants. Upstream and downstream regions are 2 kb long. (F) Overrepresentation of a specific DNA

motif in the 1-kb promoter regions of genes that are commonly upregulated in *main*, *mail1* and *pp7l*

using the MEME software. Under the logo are represented the ratio of genes carrying the motif among

the list of commonly upregulated loci. Note that the E-value associated with this motif is not

significant. (G) Same as (F) for the hypomorphic and null *pmd* mutants. Only DNA motifs with

significant E-values are shown.

S1 Table. Lists of differentially regulated loci in *ddc*, *main-3* and *ddc main-3*.

S2 Table. Lists of differentially regulated loci in *main-2*, *mail1-1* and *pp7l-1*.

S3 Table. Lists of misregulated genes carrying a specific DNA motif within their promoter.

S4 Table. List of species used to construct the two trees of Fig 4, their codes and the

presence/absence of the different PMD-C and PP7motifs are indicated.

S5 Table. (A) PMD-C and (B) PP7/PP7L motifs used to construct the two phylogenetic trees of Fig 4.

Sequences were collected at the NCBI or JGI databases.

S6 Table. List of primers used in this study.

706   2

Fig 1

Figure 1

Fig 2

Figure 2

Figure 3

Fig 4

Figure 4