Enzyme stoichiometry indicates the variation of microbial nutrient requirements at different soil depths in subtropical forests

Jiebao Liu[1,2], Ji Chen[4], Guangshui Chen[1,2], Jianfen Guo[1,2*], Yiqing Li[1,3*]

[1] School of Geography Science, Fujian Normal University, Fuzhou 350007, China; jpanther@163.com (J. Liu); gshuichen@163.com (G. Chen); jfguo@fjnu.edu.cn (J. Guo)

[2] Key Laboratory for Subtropical Mountain Ecology (Ministry of Science and Technology and Fujian Province Funded), College of Geographical Science, Fujian Normal University, Fuzhou 350007, China

[3] College of Agriculture, Forestry and Natural Resources Management, University of Hawaii, Hilo, HI 96720, USA; yiqing@hawaii.edu (Y. Li)

[4] Dept. of Agroecology, Aarhus University, Blichers Allé 20, Postboks 50, DK-8830 Tjele; ji.chen@agro.au.dk (J. Chen)

Correspondence: jfguo@fjnu.edu.cn(J. Guo); yiqing@hawaii.edu (Y. Li) Tel.: +86 591 83448515

Abstract:

Soil extracellular enzyme activities and associated enzymatic stoichiometry are considered indicators to nutrients availability and microbial substrate limitation. However, many of current studies have been focusing on upper most soil layer with a single enzyme as represent, leading to critical uncertainties in understanding the nutrient availability in deeper soils. In this study, we investigated C-, N- and P-acquiring enzyme activities and microbial C, N and P limitation across a range of soil layers (0 - 10, 10 - 20, 20 - 40 and 40 - 60 cm) in a natural secondary evergreen broad-leaved forest and a Chinese fir (*Cunninghamia lanceolata*) plantation forest in subtropical China. Our results showed that the two typical subtropical forests are commonly co-limited by C and P at all soil depths, but not N. The study found that microbial C and P limitation fluctuated along soil depth, and that higher N was demanded by microbes in soil under 20 cm in the two forests. Our results highlight the asymmetrical patterns of microbial nutrient limitation along the whole soil profile, and provide important information in understanding nutrient limitations in deeper soils. It is feasible to explicitly incorporate this vertical asymmetrical nutrient limitation patterns into future research priority.

Keywords: Deep soil; microbial nutrient limitation; soil enzyme; stoichiometry; subtropics.

1   Introduction

Forests in the tropical and subtropical area have the fastest growth rates, and exerts a more important role on sequestering atmospheric CO_2 when comparing with temperate areas in higher latitudes [1-3]. It is reported that tropical forest ecosystems were generally phosphorus limited, and additionally nitrogen limited in montane locations [4, 5]. Soil micro-organisms play a key role in releasing and maintaining soil nutrients. However, microbial processes are also limited by nutrients availability. Therefore, understanding the nutrient limitations of micro-organisms is essential for understanding the functions of ecosystems and for predicting ecosystem responses to global changes as well.

Soil enzymes, primarily produced by microbes, play a key role in decomposition, turnover and mineralization of soil organic matter, and the relative abundance of enzymes (named enzyme stoichiometry) reflects nutrient demands of microorganisms and nutrients availability in the environment [6-10]. Enzymes involved in the decomposition of cellulose (β-glucosidase (BG), cellobiosidase (CBH)), xylane (β-xylosidase, BX), chitin (β - 1,4-N-acetylglucosaminidase, NAG), polypeptides (leucine aminopeptidase, LAP) and phosphate (acid or alkaline phosphatase, AP), and have been widely recognized [e.g., 11-14]. However, carbon acquiring enzymes in forest soil were usually represented by only β-1,4-glucosidase (BG) and nitrogen acquiring enzymes were also represented by a single enzyme, β - 1,4-N-acetylglucosaminidase (NAG) because they are easier to study [15]. Thus, previous meta-analysis revealed that microbial C:N:P acquisition ratios converged on 1:1:1 when using ratios of ln(BG): ln(LAP + NAG): ln(AP) [15]. Polysaccharide degradation requires the interaction of various enzymes [16], using one single enzyme activity as an indicator of nutrient dynamics in soil may neglect synergistic effects of many other enzymes in forest ecosystem

[17, 18].

Recently, enzymatic stoichiometry has been suggested as an efficient method to indicate the relative

resource limitation of soil micro-organisms [19, 20]. The theory of enzymatic stoichiometry has been

widely used in different ecosystems [4, 15, 21-23], and a new vector analysis of eco-enzyme activities

based on this has been proposed by Moorhead et al. (2016) [24], who plotted the proportion of C:N

versus C:P acquiring enzyme activities to calculate the 'length' and 'angle' of vectors and quantify

relative C vs. nutrient acquisition and the relative P vs. N acquisition. The vector analysis has attracted

increasing attention [19, 23, 25, 26]. However, studies on enzyme activities together with enzymatic

stoichiometry utilizing whole C-acquiring enzymes as a group are rare in forest ecosystems. Paucity

of data on intact cognition of enzyme activities together with enzymatic stoichiometry constrains our

ability to accurately predict how soil nutrient cycle in forests will respond to climate change.

A large-scale study in North-South Transect of Eastern China (NSTEC) with a unique vegetation

belt ranging from boreal forest to tropical rain forest reported that microbial nutrient acquisition was

P-limited in tropical forests [27]. However, other researches indicated that karst and non-karst forests

in southwest China were P-limited, and microbial nutrient acquisition in karst ecosystem was more C-

and P-limited, rather than N-limited [23, 28]. To our best knowledge, however, no public enzymatic

stoichiometry studies have been conducted in the fast-growing forest ecosystem of Chinese fir

plantation (*Cunninghamia lanceolata*) which play a key role in regional and national carbon

sequestration.

Currently, most enzymatic stoichiometry studies focused on top soil (0-20 cm) mainly because

enzyme activity of surface soil is high and easy to change significantly with treatments [e.g., 15, 23,

27, 28]. Soils below 20 cm contribute over 50% of the global soil carbon stocks despite their low

carbon concentration [29]. Limited studies have found no consistent tendency of soil enzymes and

enzymatic stoichiometry below 20 cm [13, 21, 30-32]. Besides, substrate limitation in deep soil is a

major factor limiting microbial activity and controlling C turnover [33, 34]. Hence, exploring

enzymatic stoichiometry patterns together with microbial nutriment limitation in deep soil can provide

insights of biological mechanisms on nutrition cycling.

In this study, paired natural secondary forest (NSF) and fast-growing coniferous plantation forest

(CPF) were selected to examine enzyme activities, enzymatic stoichiometry and microbial CNP

limitation along the whole soil profile. Specifically, we addressed the following questions: (1) Will

microbial C, N, and P acquisition be significantly higher in the Chinese-fir plantation than the

secondary forest in subtropics? (2) How will soil depth affect enzyme activities and enzymatic

stoichiometry in subtropical soils?

2   Materials and methods

2.1   Study sites

The study was conducted in Shunchang county ($117° 29' \sim 118° 14'$ E, $26° 38' \sim 27° 12'$ N),

northwest of Fujian province of China. The climate in the study area is subtropical marine monsoon

with the annual mean temperature of 19.0 °C and the mean annual precipitation of 1629 mm. The

predominant soil type is red soil, which is classified as Oxisol based on the USDA's Soil Taxonomy.

Evergreen broadleaf forests and coniferous plantation forests are the main forest types at the area.

Natural secondary forest (NSF) and Chinese fir plantation forest (CPF) in the study originated from

the same natural evergreen forests. In NSF and adjacent CFP, three blocks of 20×20 m size were

selected as pairing designs with similar slope and aspect (southern). The distance between paired NSF

and CPF in one block is about 50 meters. The distance between the three blocks is within 3 km from

each other.

The dominant tree species in NSF are *Castanopsis fargesii* Franch*, Castanopsis lamontii,*

Castanopsis fargesii, Altingia gracilipes. The main understory plant species in NSF included *Machilus*

chrysotricha, *Woodwardia japonica, Adinandra millettii, Ardisia crispa (*Thunb*.), Sarcandra glabra*

*(*Thunb*.) Nakai, Embelia rudis.* Canopy coverage is approximately 75%, the mean tree height and

stand density were 14.7 m and 1267 stem ha^{-1}, respectively. The CPF is dominated by Chinese fir

(*Cunninghamia lanceolata* (Lamb.) Hook (Taxodiaceae)) and dominant understory species are

Machilus grijsii Hance, *Vaccinium bracteatum* Thunb, *Cleredendrum cwtophyllum* Turcz,

Dicranopteris dichotoma (Thunb.) *Bernh*., *Rhizoma Cibotii.* Canopy coverage, mean tree height and

stand density in CPF were 90%, 18.6 m and 1725 stem ha^{-1}, respectively.

2.2   Soil sampling

Soil sampling was taken in August 2017. After removal of surface debris, soil samples were

collected randomly at four soil depths (0 - 10, 10 - 20, 20 - 40 and 40 - 60 cm) from five random

locations in each plot. In each sampling subplot, five soil cores at each layer were taken and mixed to

represent a composite sample. Samples were immediately placed in fresh-keeping container and

transported to laboratory for chemical analysis. Stones and root fragments were removed, and soils

were sieved with a mesh size of 2 mm before chemical analyses.

2.3   Soil enzyme activities

Extracellular enzyme activities (EEA) were determined following the method of Saiya-Cork [35]

and German [36]. Fluorogenic methylumbelliferone-based (MUB) or methylcoumarin-based (MCA)

artificial substrates were used to estimate the activities of C-cycling [Xylanases (BX),

Cellobiohydrolase (CBH), beta-Glucosidase (BG)], N-cycling [beta -1,4-N-acetylglucosaminidase

(NAG), Leucine aminopeptidase (LAP)], and P-cycling [Acid phosphatase (AP)]. High substrate

concentration of each enzyme was adopted to ensure each enzyme to be assayed under saturation

conditions.

Briefly, suspensions of 1 g soil (dry weight equivalent) with 125 ml of 50 mM acetate buffer (pH

5.0) were dispersed using low-energy sonication (50 J s^{-1} output energy) [37]. The soil slurries and

substrate solutions were incubated for 180 min at 20 ± 0.5 °C. Fluorescence was measured in

microplates at an excitation wavelength of 365 nm and an emission wavelength of 450 nm with a

Synergy H4 Hybrid Microplate Reader (BioTek, USA). To ensure that all enzyme activities were

comparable, consistent parameters of incubation method and quench way were kept [7]. The

incubation plates were shaken at an interval of one every hour to ensure the complete reaction. The

maximum duration of keeping soil slurries in buffer was less than 30 min before adding MUB (MCA)-

linked substrate [38]. Fluorescence was conducted in 1 min following the addition of 10 μl 1.0 M

NaOH solution [36]. The time between the addition of substrate and the fluorescence reading was

standardized to 180 min to eliminate variation of enzyme activity.

Unit for enzyme activity was expressed as nmol activity g^{-1} dry soil h^{-1}. C acquisition was expressed

by the sum of Xylanases (BX), Cellobiohydrolase (CBH), and beta-Glucosidase (BG) activities. N

acquisition was represented by the sum of beta -1,4-N-acetylglucosaminidase (NAG) and Leucine

aminopeptidase (LAP), while P acquisition was showed as acid phosphatase (AP) activity. The microbial C:N acquisition ratio was calculated as sums of hydrolytic enzymes using (BG + BX + CBH): (NAG + LAP), while (BG + BX + CBH): (AP) and (NAG + LAP): (AP) were used for the microbial C: P and N: P acquisition ratios, respectively.

2.4　Statistical analysis

All assayed parameters were reported as means ± standard errors (SE). One-way ANOVA followed by LSD test was used to determine the significance of enzyme activities and soil enzyme stoichiometry among the soil depths in each type of the two forests. Natural-log data transformations were used to meet the assumption of normality and homoscedasticity. All statistical analyses were done by SPSS v 23.0 (IBM SPSS Statistics for Windows, ver. 23.0; IBM Corp., Armonk, NY, USA). Mixed linear model analyses in SPSS v23.0 was selected to determine the effects of forest type, soil depth and their interactions on soil enzyme activities and enzymatic stoichiometry.

The following three approaches were used to examine microbial resource acquisition. The first was based on the method of Hill [22] to draw scatter plot of enzymatic stoichiometry, with Enzyme N/ Enzyme P as x axis and Enzyme C/ Enzyme N as y axis. Different resource constraints (N limitation, P limitation, C&P limitation and N&P limitation) were shown in four district respectively. The second approach was based on enzyme ratios of C:P and N:P. Lower enzyme C:P and N:P ratios are suggestive of a degree of P limitation [4]. The third approach was vector analysis proposed by Moorhead [24].

$$\text{Vector L (unitless)} = \sqrt{X^2 + Y^2} \qquad (1)$$

$$\text{Vector A (degree)} = \text{Degrees (Atan2(X, Y))} \qquad (2)$$

Where

$$X= \frac{Enzyme\ C}{Enzyme\ C\ +\ Enzyme\ P} \quad , \quad Y= \frac{Enzyme\ C}{Enzyme\ C\ +\ Enzyme\ N}$$

A relatively longer vector L indicates greater C- acquisition, and the vector A < 45° and >45°

indicate relative degrees of N- and P-limitation, respectively [19, 28].

3 Results

3.1 Soil enzyme activity

Results from mixed linear model showed that soil depth significantly influenced Enzyme C, N and

P acquisition, while the interactions of forest type × soil depth significantly affected enzyme C (Table

1; $P<0.05$). From the soil depths of 0–10 cm to 40-60 cm, the mean activities of C, N, and P acquisition

enzymes decreased by 64%, 48%, and 76%, respectively (Fig.1). There were no remarkable

differences of C-acquiring (BX+CBH+BG) and P-acquiring (AP) enzyme activities between NSF and

CPF at 0-10 cm, 10-20 cm, 20-40 cm, 40-60 cm depths. N-acquiring (NAG + LAP) enzyme activity

in NSF at 0-10 cm soil depth was significantly higher than that in CPF (Fig.1).

For NSF, C- and N-acquiring enzyme activities had the same distribution trend that surface soil (0-

10 cm) layer had remarkably higher activities than those at the depths of 10-20, 20-40 and 40-60 cm,

among which remarkable difference was not found (Fig.1). There was no significant difference in P-

acquiring enzyme activity between 0-10 cm and 10-20 cm depths, but P-acquiring enzyme activities

at the two soil depths were significantly higher than those at 20-40 and 40-60 cm depths. For CPF, C-

acquiring enzyme activity in 0-10 cm layer was remarkably higher than that in 20-40 and 40-60 cm

soil depths (Fig.1). Similarly, N-acquiring enzyme activity in topsoil (0-10 cm) was significantly

higher than in 10-20 and 40-60 cm soil depths. Enzyme P activity in 0-10 cm was almost 2-fold higher

than in 10-20 cm soil depth, and significantly higher than in 20-40 and 40-60 cm soil depths (Fig.1).

3.2 Soil enzymatic stoichiometry

Throughout the study area, mean soil enzymatic ratios of (BG+BX+CBH):(NAG+LAP), (BG+BX+CBH):AP and (NAG+LAP):AP were 3.88, 0.27 and 0.08, respectively. Enzyme activity ratio of C:P (0.30) in CPF was significantly higher than that of NSF (0.24) (Fig. 2), all of which were significantly affected by soil depth, forest type and their interactions (Table 1, $P<0.05$). While ratio of enzyme N:P was only affected notably by soil depth (Table 1, $P<0.05$). From 0–10 cm to 40-60 cm soil depth, mean soil enzymatic ratios of C:P and N:P increased by 59% and 119%, respectively (Fig. 2).

Enzymatic ratio of C:N in CPF located at 10-20 cm (5.91) was significantly higher than the other three soil layers (Fig. 3). At soil layers of 10-20 cm and 40-60 cm, enzymatic ratios of C:P in CPF were 0.30 and 0.40 respectively, which were significantly higher than those in NSF. The highest enzymatic ratio of C:P was observed at the soil layer of 20-40 cm in NSF (0.34) and 40 -60 cm in CPF (0.40), respectively (Fig. 3). Enzymatic ratios of C:P and N:P in CPF increased with soil depths. Enzymatic ratios of N: P at 0-10 cm and 10-20 cm depths in CPF were remarkably lower than those at 20-40 cm and 40-60 cm depths (Fig. 3)

3.3 Vector analysis

The plots of enzymatic stoichiometry indicated that four soil layers were co-C and P limited and not with N limited in NSF and CPF (Table 2, Fig. S1). Vector analysis showed that vector length and angle were not significantly different between NSF and CPF at 0-60 cm soil depth, indicating no differences of C-limited and P-limited between NSF and CPF at each soil layer (Table 2).

Vector length showed that there were no significant differences among the four soil layers in NSF,

and the third (20-40 cm) layer was notably lower than 0-10 and 10-20 cm soil layers in CPF (Table 2).

Vector angle in CPF decreased significantly along the four soil layers, with angle greater than 45° in

each layer. Although vector A varied significantly with depth in CPF, there was no coincident tendency

of vector A in NSF. In NSF the angles at each soil layer were greater than 45°, and the largest and

smallest angles occurred in the 10-20 cm and 20-40 cm soil depths, respectively. Further, the study did

not find differences in N limitation between NSF and CPF (Table 2).

4   Discussion

4.1   Spatial variations of C-, N- and P- acquiring enzyme activities

Generally, hydrolase activities decreased gradually with the decrease of soil depth [15, 31, 32], we

observed that C and P acquiring enzyme activities exhibited a decreasing trend with soil depth but N-

acquiring enzyme activities had an increasing trend in NSF. The most fluctuated enzyme activity was

at 10-20 cm soil layer, where C- acquiring enzyme activity in NSF and N- acquiring enzyme activity

in NSF and CPF decreased sharply. The results indicated that microbial metabolisms were obviously

higher at the 0-10 cm soil, which can explained by that fine root biomass of natural secondary forest

and Chinese fir plantation were abundantly distributed at the top 0–10 cm soil layer [39] and active

microbial activities due to rich easily-degradable weight organic substances in the topsoil [40, 41].

Despite no remarkable difference of C-acquiring enzyme activities was found between NSF and

CPF at 0-10 cm soil layer such as BG enzyme, BX and CBH enzyme activities were significantly

higher in NSF than in CPF at 0-10 cm (Table S1). Although enzymes are substrate specific and

individual enzyme activity may not reflect total soil nutritional status [42], lots of studies utilized β-

1,4-glucosidase (BG) as whole carbon acquiring enzyme because glucose released from cellulose is

the most abundant monomer in soil [17]. Oligo- and monosaccharides (termed sugars), including

glucose, xylose, arabinose, galactose, mannose and rhamnose, come from decomposition of

polysaccharides by microbial enzymes (cellulases, xylanases, glucosidases and chitinases etc.). These

sugars are soluble and captured rapidly by microbes for metabolism and carbon storage [17].

Complicated belowground biochemical processes indicate that using a single enzyme to describe the

mechanism of microbial nutrient demands may be biased and using more enzymes beyond β-1,4-

glucosidase (BG) to reflect the mechanisms are needed.

In the present study, significant difference in N-acquiring enzyme was found at the surface soil

between the two forests due to higher Leucine aminopeptidase (LAP) activity, which contributed

almost 50% to the total N acquiring enzyme activity (Table S1). The result revealed LAP plays an

important role in forest soil as found by Moorhead et al. (2016) [24]. Nitrogen -acquiring enzyme

activity in CPF did not decrease gradually with soil depth, which was associated with LAP activity

(Fig. 1). The distribution pattern of LAP showed a decreasing trend followed by an increasing trend

with increasing soil depth (Table S1).

It was reported that Chinese fir might uptake phosphorus through soil acidification [43] and produce

less acid phosphates [44], which resulted in high P demand for its rapid growth with relatively lower

enzyme P activity at all soil depths in CPF compared to NSF .

4.2   Enzymatic stoichiometry and microbial nutrient limitations

The lower enzymatic ratios of N: P and C: P (<1) together with the results that vector angles were

higher than 45° in both NSF and CPF at the 0-60 cm soil depth indicated P limitation in NSF and CPF

[4, 45]. Besides, gradual decrease of vector A in CPF indicated that microbial P limitation gradually

reduced along soil depth (Table 2). While microbial P demand fluctuated according to soil depth in

NSF, with highest demand at 10-20 cm and the lowest demand at 20-40 cm soil layer. Former studies

revealed that P restrictions are quite common [5, 46, 47], the study tried to provide evidences of

microbial P limitation in middle-subtropical forests.

In the study, both the scatter plot of enzymatic stoichiometry (Fic. S1) and the vector analysis (Table

2) indicated that the two forests were commonly co-limited by C and P at all soil depths, which were

consistent with the study conducted at karst ecosystems in southwest China [28]. Meanwhile, higher

microbial C acquiring in CPF at 10-20 cm and 40-60 cm soil layers were found due to remarkably

higher enzyme C: P ratio at the two soil layers in CPF than in NSF. The results confirmed that forest

type, soil depth and their interactions have significant effects on the ratio of enzyme C: P (Table 1;

$P<0.05$).

It was also reported that many subtropical ecosystems are N-rich relative to other plant nutrients

[48-50], the study indicates that N was not limited with respect to microbial grow demand from the

results of enzymatic stoichiometry. That is, the C: N ratio may not reach the critical ratio of the two

elements for microbial growth. The sharply increasing trend of enzyme N:P ratio below 20 cm soil

layer indicated the higher N demand in deep soil, which was in line with the result of incubation

experiment of a subtropical mixed forest soil [51]. The higher enzyme C: N ratio in CPF at 0-20 cm

and 40-60 cm soil layers suggested that NSF was more N-limited compared to CPF.

Although the trends of enzyme C:N and C:P ratios remained unchanged at all soil layers between

the two forests when using single BG as whole C acquisition enzyme, enzyme C:P ratio at NSF

displayed a different trend along the soil depths. Enzyme C:P ratio between 20-40 cm and 40-60 cm

soil depths at NSF changed from significance to insignificance (Fig. 3;Fig. S2), which may

underestimate the C demand of NSF at the 20-40 cm soil depth. On the other hand, vector A in NSF

at 40-60 cm soil depth become significantly higher than that in CPF when using single BG as whole

C acquisition enzyme (Table S2), which may overestimate microbial P limitation in CPF at 40-60 cm

soil depth. Thus, utilizing single BG as whole C acquisition enzyme may underestimate or

overestimate microbial C and P demand in deep soils.

East Asian monsoon subtropical forests have one of the highest carbon uptakes of forests worldwide,

which may be caused by young stand ages, and high nitrogen deposition [52, 53]. Nevertheless,

research also indicated that soil total C concentration increased significantly and soil total P

concentration decreased significantly across all soil depths in subtropical China during 1955 to 2016

[49]. To our best knowledge, this study pioneered in examining microbial nutrient demands along a

whole soil profile in middle-subtropical forests, which should contribute to informational references

for forest management in the subtropics.

5   Conclusion

The study utilized enzymatic stoichiometry to evaluate microbial C, N and P limitation in a natural

secondary evergreen broad-leaved forest and a fast-growing plantation forest in the subtropics, where

enzymatic stoichiometry is much less studied. We found that the two subtropical forests were co-

limited by C and P at all soil depths (0-10, 10-20, 20-40 and 40-60 cm) and both forests were not N-

limited. No significant differences of microbial C, N and P limitation between the two forests at all

soil depths revealed the complicated belowground plant-microbe interactions. The results of enzyme

stoichiometry indicate higher N demand by microbes in soils below 20 cm in both plantation forest

and the secondary forest. The finding that C-acquiring (BX+CBH+BG), N-acquiring (NAG + LAP)

and P-acquiring enzyme activities decreased with increasing soil depth but there was no difference in enzyme activities along the whole soil profile indicates an imbalance of microbial nutrient demand at different soil depths. Utilizing single BG as whole C acquisition enzyme could miscalculate microbial C and P demands at different soil layers. This pioneer study in subtropical China in examining microbial nutrient demands in both top and deep soils provides important information for subtropical tree plantation and natural forest management, especially under future climate change.

Acknowledgments: Thanks Moorhead Daryl for his valuable suggestions in preparing the manuscript.

Author Contributions: Y. Li conceived and designed the experiments; J. Liu performed the experiments and wrote the manuscript; G. Chen helped to analyze the data; J. Guo and J. Chen. participated in revising the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Reference

1. Post WM, Kwon KC. Soil carbon sequestration and land-use change: processes and potential. Glob Chang Biol. 2000;6(3):317-27. doi: 10.1046/j.1365-2486.2000.00308.x. PubMed PMID: WOS:000086198100006.

2. Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, et al. A Large and Persistent Carbon Sink in the World's Forests. Science. 2011;333(6045):988-93. doi: 10.1126/science.1201609. PubMed PMID: WOS:000294000400048.

3. Reich PB, Luo Y, Bradford JB, Poorter H, Perry CH, Oleksyn J. Temperature drives global patterns in forest biomass distribution in leaves, stems, and roots. Proc Natl Acad Sci U S A. 2014;111(38):13721-6. Epub 2014/09/17. doi: 10.1073/pnas.1216053111. PubMed PMID: 25225412; PubMed Central PMCID: PMCPMC4183289.

4. Waring BG, Weintraub SR, Sinsabaugh RL. Ecoenzymatic stoichiometry of microbial nutrient acquisition in tropical soils. Biogeochemistry. 2014;117(1):101-13. doi: 10.1007/s10533-013-9849-x. PubMed PMID: WOS:000329606200008.

5. Camenzind T, Hättenschwiler S, Treseder KK, Lehmann A, Rillig MC. Nutrient limitation of soil microbial processes in tropical forests. Ecol Monogr. 2018;88(1):4-21. doi: doi:10.1002/ecm.1279.

6. Allison SD. Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. Ecol Lett. 2005;8(6):626-35. doi: 10.1111/j.1461-0248.2005.00756.x. PubMed PMID: WOS:000229389500007.

7. Burns RG, DeForest JL, Marxsen J, Sinsabaugh RL, Stromberger ME, Wallenstein MD, et al. Soil enzymes in a changing environment: Current knowledge and future directions. Soil Biol Biochem. 2013;58:216-34. doi: 10.1016/j.soilbio.2012.11.009. PubMed PMID: WOS:000317158300027.

8. Moorhead DL, Lashermes G, Sinsabaugh RL. A theoretical model of C- and N-acquiring exoenzyme activities, which balances microbial demands during decomposition. Soil Biol Biochem. 2012;53:133-41. doi: 10.1016/j.soilbio.2012.05.011. PubMed PMID: WOS:000307141400018.

9. Chen J, Luo YQ, Garcia-Palacios P, Cao JJ, Dacal M, Zhou XH, et al. Differential responses of carbon-degrading enzyme activities to warming: Implications for soil respiration. Glob Chang Biol. 2018;24(10):4816-26. doi: 10.1111/gcb.14394. PubMed PMID: WOS:000445728800028.

10. Sinsabaugh RL, Hill BH, Shah JJF. Ecoenzymatic stoichiometry of microbial organic nutrient acquisition in soil and sediment. Nature. 2009;462(7274):795-U117. doi: 10.1038/nature08632. PubMed PMID: WOS:000272559900040.

11. Maly S, Fiala P, Reininger D, Obdrzalkova E. The relationships among microbial parameters and the rate of organic matter mineralization in forest soils, as influenced by forest type. Pedobiologia. 2014;57(4-6):235-44. doi: 10.1016/j.pedobi.2014.09.003. PubMed PMID: WOS:000347578700005.

12. Wittmann C, Kähkönen MA, Ilvesniemi H, Kurola J, Salkinoja-Salonen MS. Areal activities and stratification of hydrolytic enzymes involved in the biochemical cycles of carbon, nitrogen, sulphur and phosphorus in podsolized boreal forest soils. Soil Biol Biochem. 2004;36(3):425-33. doi: 10.1016/j.soilbio.2003.10.019.

13. Jing X, Chen X, Tang M, Ding Z, Jiang L, Li P, et al. Nitrogen deposition has minor effect on soil extracellular enzyme activities in six Chinese forests. Sci Total Environ. 2017;607-608:806-15. Epub 2017/07/18. doi: 10.1016/j.scitotenv.2017.07.060. PubMed PMID: 28711842.

14. Oulehle F, Tahovská K, Chuman T, Evans CD, Hruška J, Růžek M, et al. Comparison of the impacts of acid and nitrogen additions on carbon fluxes in European conifer and broadleaf forests. Environ Pollut. 2018;238:884-93. doi: 10.1016/j.envpol.2018.03.081.

15. Sinsabaugh RL, Lauber CL, Weintraub MN, Ahmed B, Allison SD, Crenshaw C, et al. Stoichiometry of soil enzyme activity at global scale. Ecol Lett. 2008;11(11):1252-64. doi: 10.1111/j.1461-0248.2008.01245.x. PubMed PMID: WOS:000260466400013.

16. Warren RAJ. MICROBIAL HYDROLYSIS OF POLYSACCHARIDES. 1996;50(1):183-212. doi: 10.1146/annurev.micro.50.1.183. PubMed PMID: 8905079.

17. Gunina A, Kuzyakov Y. Sugars in soil and sweets for microorganisms: Review of origin, content, composition and fate. Soil Biol Biochem. 2015;90:87-100. doi: 10.1016/j.soilbio.2015.07.021. PubMed PMID: WOS:000363075500011.

18. Nannipieri P, Giagnoni L, Renella G, Puglisi E, Ceccanti B, Masciandaro G, et al. Soil enzymology: classical and molecular approaches. Biol Fertil Soils. 2012;48(7):743-62. doi: 10.1007/s00374-012-0723-0. PubMed PMID: WOS:000310081800001.

19. Hill BH, Elonen CM, Jicha TM, Kolka RK, Lehto LLP, Sebestyen SD, et al. Ecoenzymatic stoichiometry and microbial processing of organic matter in northern bogs and fens reveals a common P-limitation between peatland types. Biogeochemistry. 2014;120(1-3):203-24. doi: 10.1007/s10533-014-9991-0. PubMed PMID: WOS:000339871700013.

20. Sinsabaugh RL, Shah JJF, Hill BH, Elonen CM. Ecoenzymatic stoichiometry of stream sediments with comparison to terrestrial soils. Biogeochemistry. 2012;111(1-3):455-67. doi: 10.1007/s10533-011-9676-x. PubMed PMID: WOS:000314063200029.

21. Peng XQ, Wang W. Stoichiometry of soil extracellular enzyme activity along a climatic transect in temperate grasslands of northern China. Soil Biol Biochem. 2016;98:74-84. doi: 10.1016/j.soilbio.2016.04.008. PubMed PMID: WOS:000378008900009.

22. Hill BH, Elonen CM, Seifert LR, May AA, Tarquinio E. Microbial enzyme stoichiometry and nutrient limitation in US streams and rivers. Ecol Indicators. 2012;18:540-51. doi: 10.1016/j.ecolind.2012.01.007. PubMed PMID: WOS:000303284500059.

23. Chen H, Li D, Xiao K, Wang K. Soil microbial processes and resource limitation in karst and non-karst forests. Funct Ecol. 2018;32(5):1400-9. doi: 10.1111/1365-2435.13069. PubMed PMID: WOS:000431991700024.

24. Moorhead DL, Sinsabaugh RL, Hill BH, Weintraub MN. Vector analysis of ecoenzyme activities reveal constraints on coupled C, N and P dynamics. Soil Biol Biochem. 2016;93:1-7. doi: 10.1016/j.soilbio.2015.10.019. PubMed PMID: WOS:000369207200001.

25. Fanin N, Moorhead D, Bertrand I. Eco-enzymatic stoichiometry and enzymatic vectors reveal differential C, N, P dynamics in decaying litter along a land-use gradient. Biogeochemistry. 2016;129(1-2):21-36. doi: 10.1007/s10533-016-0217-5. PubMed PMID: WOS:000381287000002.

26. Pei ZQ, Leppert KN, Eichenberg D, Bruelheide H, Niklaus PA, Buscot F, et al. Leaf litter diversity alters microbial activity, microbial abundances, and nutrient cycling in a subtropical forest ecosystem. Biogeochemistry. 2017;134(1-2):163-81. doi: 10.1007/s10533-017-0353-6.

PubMed PMID: WOS:000406191000012.

27. Xu ZW, Yu GR, Zhang XY, He NP, Wang QF, Wang SZ, et al. Soil enzyme activity and stoichiometry in forest ecosystems along the North-South Transect in eastern China (NSTEC). Soil Biol Biochem. 2017;104:152-63. doi: 10.1016/j.soilbio.2016.10.020. PubMed PMID: WOS:000389555900015.

28. Chen H, Li DJ, Mao QG, Xiao KC, Wang KL. Resource limitation of soil microbes in karst ecosystems. Sci Total Environ. 2019;650:241-8. doi: 10.1016/j.scitotenv.2018.09.036. PubMed PMID: WOS:000447092700027.

29. Jobbagy EG, Jackson RB. The Vertical Distribution of Soil Organic Carbon and Its Relation to Climate and Vegetation. Ecol Appl. 2000;10(2):423-36. doi: 10.2307/2641104.

30. Zuo YP, Li JP, Zeng H, Wang W. Vertical pattern and its driving factors in soil extracellular enzyme activity and stoichiometry along mountain grassland belts. Biogeochemistry. 2018;141(1):23-39. doi: 10.1007/s10533-018-0499-x. PubMed PMID: WOS:000447723800002.

31. Stone MM, DeForest JL, Plante AF. Changes in extracellular enzyme activity and microbial community structure with soil depth at the Luquillo Critical Zone Observatory. Soil Biol Biochem. 2014;75:237-47. doi: 10.1016/j.soilbio.2014.04.017. PubMed PMID: WOS:000338619600026.

32. Schnecker J, Wild B, Takriti M, Eloy Alves RJ, Gentsch N, Gittel A, et al. Microbial community composition shapes enzyme patterns in topsoil and subsoil horizons along a latitudinal transect in Western Siberia. Soil Biol Biochem. 2015;83:106-15. doi: http://dx.doi.org/10.1016/j.soilbio.2015.01.016.

33. Tian QX, Yang XL, Wang XG, Liao C, Li QX, Wang M, et al. Microbial community mediated response of organic carbon mineralization to labile carbon and nitrogen addition in topsoil and subsoil. Biogeochemistry. 2016;128(1-2):125-39. doi: 10.1007/s10533-016-0198-4. PubMed PMID: WOS:000374557300008.

34. Heitkotter J, Heinze S, Marschner B. Relevance of substrate quality and nutrients for microbial C-turnover in top-and subsoil of a Dystric Cambisol. Geoderma. 2017;302:89-99. doi: 10.1016/j.geoderma.2017.04.029. PubMed PMID: WOS:000403522800009.

35. Saiya-Cork KR, Sinsabaugh RL, Zak DR. The effects of long term nitrogen deposition on extracellular enzyme activity in an Acer saccharum forest soil. Soil Biol Biochem. 2002;34(9):1309-15. doi: 10.1016/s0038-0717(02)00074-3. PubMed PMID: WOS:000177861500012.

36. German DP, Weintraub MN, Grandy AS, Lauber CL, Rinkes ZL, Allison SD. Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. Soil Biol Biochem. 2011;43(7):1387-97. doi: 10.1016/j.soilbio.2011.03.017. PubMed PMID: WOS:000291576800001.

37. Marx MC, Wood M, Jarvis SC. A microplate fluorimetric assay for the study of enzyme diversity in soils. Soil Biol Biochem. 2001;33(12-13):1633-40. doi: 10.1016/s0038-0717(01)00079-7. PubMed PMID: WOS:000171415500006.

38. DeForest JL. The influence of time, storage temperature, and substrate age on potential soil enzyme activity in acidic forest soils using MUB-linked substrates and L-DOPA. Soil Biol Biochem. 2009;41(6):1180-6. doi: 10.1016/j.soilbio.2009.02.029. PubMed PMID: WOS:000266942900021.

39. Yang Y-S, Chen G-S, Lin P, Xie J-S, Guo J-F. Fine root distribution, seasonal pattern and production in four plantations compared with a natural forest in Subtropical China. Annals of Forest Science. 2004;61(7):617-27. doi: 10.1051/forest:2004062.

40. Phillips RP, Meier IC, Bernhardt ES, Grandy AS, Wickings K, Finzi AC. Roots and fungi accelerate carbon and nitrogen cycling in forests exposed to elevated CO2. Ecol Lett. 2012;15(9):1042-9. doi: 10.1111/j.1461-0248.2012.01827.x. PubMed PMID: 22776588.

41. Hasselquist NJ, Metcalfe DB, Inselsbacher E, Stangl Z, Oren R, Nasholm T, et al. Greater carbon allocation to mycorrhizal fungi reduces tree nitrogen uptake in a boreal forest. Ecology. 2016;97(4):1012-22. doi: 10.1890/15-1222.1. PubMed PMID: WOS:000373923100020.

42. Nannipieri P, Trasar-Cepeda C, Dick RP. Soil enzyme activity: a brief history and biochemistry as a basis for appropriate interpretations and meta-analysis. Biol Fertil Soils. 2018;54(1):11-9. doi: 10.1007/s00374-017-1245-6. PubMed PMID: WOS:000419135400002.

43. Yu Y, Yu J, Shan Q, Fang L, Jiang D. Organic acid exudation from the roots of Cunninghamia lanceolata and Pinus massoniana seedlings under low phosphorus stress. Frontiers of Forestry in China. 2008;3(1):117-20. doi: 10.1007/s11461-008-0014-y.

44. Lang F, Bauhus J, Frossard E, George E, Kaiser K, Kaupenjohann M, et al. Phosphorus in forest ecosystems: New insights from an ecosystem nutrition perspective. J Plant Nutr Soil Sci. 2016;179(2):129-35. doi: 10.1002/jpln.201500541. PubMed PMID: WOS:000374035700001.

45. Luo L, Meng H, Gu J-D. Microbial extracellular enzymes in biogeochemical cycling of ecosystems. J Environ Manage. 2017;197:539-49. doi: 10.1016/j.jenvman.2017.04.023. PubMed PMID: WOS:000401880100052.

46. Zechmeister-Boltenstern S, Keiblinger KM, Mooshammer M, Penuelas J, Richter A, Sardans J, et al. The application of ecological stoichiometry to plant-microbial-soil organic matter transformations. Ecol Monogr. 2015;85(2):133-55. doi: 10.1890/14-0777.1.sm. PubMed PMID: WOS:000353845900001.

47. Schimel JP, Weintraub MN. The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. Soil Biol Biochem. 2003;35(4):549-63. doi: 10.1016/s0038-0717(03)00015-4. PubMed PMID: WOS:000182365300005.

48. Zhang J-b, Cai Z-c, Zhu T-b, Yang W-y, Müller C. Mechanisms for the retention of inorganic N in acidic forest soils of southern China. Sci Rep. 2013;3:2342. doi: 10.1038/srep02342

49. Yu Z, Wang M, Huang Z, Lin TC, Vadeboncoeur MA, Searle EB, et al. Temporal changes in soil C-N-P stoichiometry over the past 60 years across subtropical China. Glob Chang Biol. 2018;24(3):1308-20. Epub 2017/10/14. doi: 10.1111/gcb.13939. PubMed PMID: 29028280.

50. Zhao W, Cai Z-C, Xu Z-H. Does ammonium-based N addition influence nitrification and acidification in humid subtropical soils of China? Plant Soil. 2007;297(1-2):213-21. doi: 10.1007/s11104-007-9334-1. PubMed PMID: WOS:000248856600018.

51. Tian QX, Wang XG, Wang DY, Wang M, Liao C, Yang XL, et al. Decoupled linkage between soil carbon and nitrogen mineralization among soil depths in a subtropical mixed forest. Soil Biol Biochem. 2017;109:135-44. doi: 10.1016/j.soilbio.2017.02.009. PubMed PMID: WOS:000397691700013.

52. Yu G, Chen Z, Piao S, Peng C, Ciais P, Wang Q, et al. High carbon dioxide uptake by subtropical forest ecosystems in the East Asian monsoon region. Proceedings of the National Academy of Sciences. 2014;111(13):4910. doi: 10.1073/pnas.1317065111.

53. Chen J, Luo YQ, van Groenigen KJ, Hungate BA, Cao JJ, Zhou XH, et al. A keystone microbial

enzyme for nitrogen control of soil carbon storage. Science Advances. 2018;4(8). doi: 10.1126/sciadv.aaq1689. PubMed PMID: WOS:000443498100009.

Table 1. Estimated fixed effects of soil depth, forest type and their interactions on soil enzyme, enzyme stoichiometry. C acquisition enzyme was represented by the sum of β-1,4-glucosidase (BG), cellobiohydrolase (CBH), β-xylosidase (BX). The sum of β-1,4-N-acetylglucosaminidase (NAG) and leucine aminopeptidase (LAP) was regarded as N acquisition, while the activity of acid phosphatase (AP) was considered as P acquisition enzyme. Enzymatic stoichiometry of C:N, C:P and N:P was calculated by the C, N and P acquisition enzyme respectively. Significant results ($P<0.05$) are indicated in bold.

Parameters	Intercept			Forest type			Soil depth			Forest type × Soil depth		
	DF	F	*P*	DF	F	*P*	DF	F	*P*	DF	F	*P*
Enzyme C	1	167.9	**< 0.001**	1	0.016	0.904	3	26.40	**< 0.001**	3	4.456	**0.026**
Enzyme N	1	963.7	**< 0.001**	1	6.458	0.060	3	14.31	**< 0.001**	3	1.118	0.388
Enzyme P	1	169.1	**< 0.001**	1	1.495	0.287	3	59.51	**< 0.001**	3	1.916	0.205
Enzyme C:N	1	141.1	**< 0.001**	1	1.098	0.362	3	3.183	0.087	3	1.595	0.268
Enzyme C:P	1	1994.6	**< 0.001**	1	23.29	**< 0.001**	3	21.68	**< 0.001**	3	5.491	**0.013**
Enzyme N:P	1	159.4	**< 0.001**	1	0.271	0.630	3	12.99	**0.001**	3	1.028	0.423

Table 2. Vector analysis between the natural secondary forest (NSF) and Chinese fir plantation forest (CPF) at different soil depth. Values represented mean ± standard error (n=3). Capital letters show the significant difference between the two forests at the same soil depth, and the different lower cases reflect the significant difference between soil depths within one forest.

Depth	Vector L		Vector A	
(cm)	NSF	CPF	NSF	CPF
0-10	0.813±0.02Aa	0.841±0.02Aa	77.64±0.71Aab	78.37±0.62Aa
10-20	0.78±0.06Aa	0.88±0.03Aa	79.14±1.20Aa	74.83±0.78Ab
20-40	0.81±0.04Aa	0.76±0.02Ab	71.75±1.87Ac	72.05±0.50Ac
40-60	0.75±0.02Aa	0.82±0.03Aab	74.54±0.95Abc	69.48±0.79Ad

FIGURE CAPTIONS

Fig. 1. Variations of C, N and P acquiring enzyme activities at different soil depths of the natural secondary forest (NSF) and Chinese fir plantation forest (CPF). Statistically significant differences were assumed at $P < 0.05$. All values are presented as mean ± standard error with 3 replicates. Capital letters show the significant difference between the two forests at the same soil depth, and the different lower cases reflect the significant difference between soil depths within one forest.

Fig. 2. Comparisons of enzymatic stoichiometry at the 0-60 cm soil depth of the natural secondary forest (NSF) and Chinese fir plantation forest (CPF). All values are presented as mean ± standard error. "*" indicates a significance at a $P < 0.05$ level.

Fig. 3. Variations of enzymatic stoichiometry at different soil depths of the natural secondary forest (NSF) and Chinese fir plantation forest (CPF). Capital letters mean the significant difference between the two forests at the same soil depth, and the different lower cases reflect the significant difference between soil depths within one forest. All values are presented as mean ± standard error with 3 replicates.

Fig. 1.

Fig. 2.

Fig. 3.