Regulation of neuronal mRNA splicing and Tau isoform ratio by ATXN3

through deubiquitylation of splicing factors

RUNNING TITLE: Neuronal splicing regulation by ATXN3

Andreia Neves-Carvalho[1,2], Sara Duarte-Silva[1,2], Joana Silva[1,2], Bruno Almeida[1,2],

Sasja Heetveld[3], Ioannis Sotiropoulos[1,2], Peter Heutink[3], Ka Wan Li[4] and Patrícia

Maciel[1,2*]

[1] Life and Health Sciences Research Institute (ICVS), School of Medicine, University

of Minho, Braga, Portugal

[2] ICVS/3B's – PT Government Associate Laboratory, Braga/Guimarães, Portugal

[3] German Center for Neurodegenerative Diseases (DZNE), 72076 Tübingen,

Germany

[4] CNCR, Amsterdam Neuroscience, Vrije Universiteit, Amsterdam, The Netherlands

CORRESPONDING AUTHOR:
Patrícia Maciel, PhD
Associate Professor
School of Medicine
University of Minho
Braga, Portugal
Telephone: 351-253-604824
Fax: 351-253-604820
E-mail: pmaciel@med.uminho.pt
ORCID ID: 0000-0002-4151-6885

FUNDING

This work has been funded by FEDER funds, through the Competitiveness Factors

Operational Programme (COMPETE), and by National funds, through the Foundation

for Science and Technology (FCT), under the scope of the project POCI-01-0145-

FEDER-007038 and through the project POCI-01-0145-FEDER-016818

(PTDC/NEU-NMC/3648/2014) and fellowships [SFRH/BD/51059/2009 to A. N-C,

SFRH/BPD/110728/2015 to B.A, SFRH/BD/78388/2011 to S.D-S,

SFRH/BD/88932/2012 to J.M.S and IF/01799/2013 to I.S.

ABSTRACT

The ubiquitylation/deubiquitylation balance in cells is maintained by

Deubiquitylating enzymes, including ATXN3. The precise role of this protein,

mutated in SCA3, remains elusive, as few substrates for its deubiquitylating

activity were identified. Therefore, we characterized the ubiquitome of

neuronal cells lacking ATXN3, and found altered polyubiquitylation in a large

proportion of proteins involved in RNA metabolism, including splicing factors.

Using transcriptomic analysis and reporter minigenes we confirmed that

splicing was globally altered in these cells. Among the targets with altered

splicing was SRSF7 (9G8), a key regulator of MAPT (Tau) exon 10 splicing.

Loss-of-function of ATXN3 led to a deregulation of MAPT exon 10 splicing

resulting in a decreased 4R/3R-Tau ratio. Similar alterations were found in the

brain of a SCA3 mouse and humans, pointing to a relevant role of this

mechanism in SCA3, and establishing a previously unsuspected link between

two key proteins involved in different neurodegenerative disorders.

Ataxin-3/ Neurodegeneration/ Serine-Arginine rich splicing factor 7 (SRSF7 or

9G8)/ Splicing/ Ubiquitylation

INTRODUCTION

Ubiquitin signaling is now recognized as a fundamental molecular mechanism

tightly regulating a broad range of intracellular events [1-3]. Ubiquitylation is a

highly dynamic biochemical modification in which an ubiquitin (Ub) moiety is

attached to a protein. This process is catalyzed by the sequential actions of

Ub-activating enzyme (E1), Ub-conjugating enzymes (E2) and Ub-protein

ligases (E3) that bind Ub to different lysine (K) residues in the substrate,

resulting in mono or poly Ub chains (Reviewed in [4]). Ubiquitylated substrates

are then recognized by proteins containing ubiquitin binding domains and

directed to different fates. For example, K48 usually targets proteins for

proteasomal degradation [4], while K63-linked polyUb regulates protein

activation, subcellular localization or degradation in the lysosome (autophagy)

and is known to be relevant for DNA repair (Reviewed in [5]). Ubiquitylation and

proteolysis by the Ubiquitin-proteasome pathway (UPP) are now well-known

as important mechanisms in the nervous system as this proteolytic pathway is

known to degrade misfolded or short-lived regulatory proteins (Reviewed in

[1,6]). Impairment of the UPP has been connected to several neurodegenerative

diseases such as Alzheimer's (Reviewed in [7]), Parkinson's (Reviewed in [8])

and Hungtington's (Reviewed in [9]) diseases. Like most post-transcriptional

modifications, ubiquitylation is a reversible signal and is counterbalanced by

deubiquitylating (DUB) enzymes, which remove Ub units and play an

important role in the modulation of proteins by the proteasome as well as the

autophagic pathway [10-12], and in Ub signaling in general. Several pieces of

evidence have suggested that Ub may also regulate splicing: (i) Ub and Ub-

like proteins have been shown to copurify with splicing complexes [13,14], (ii)

ubiquitylated splicing factors have been identified in proteomic screenings [15,16]

and (iii) several functional domains related to Ub pathway have been

identified in important spliceosome proteins, such as Jab1/MPN domain of the

essential U5 SnRNP component Prp8 [17-20]. Thus, the action of DUBs has a

major impact on the ubiquitylated proteome (also known as ubiquitome).

Ataxin-3 (ATXN3) is a protein with DUB activity known to be involved in

Spinocerebellar ataxia type 3 (SCA3)/ Machado Joseph disease (MJD), a

neurodegenerative disorder of adult onset caused by the expansion of a

polyglutamine (polyQ) tract in this protein. Interestingly, in addition to being

involved in global protein quality control responses [21,22], normal ATXN3

appears to participate in cell adhesion and in the organization of cytoskeleton

network [23-26], to be involved in transcriptional regulation and DNA repair [27-29],

and to be required for neuronal differentiation [25]. Nevertheless, the relative

relevance of these functions and the physiological role of this highly

conserved but genetically non-essential protein remains to be clarified and its

substrates are also not well characterized, particularly in neurons. Given the

known relevance of UPP in neuronal function and its link to nervous system

diseases, we sought to clarify the precise role of ATXN3. To identify potential

candidates for the DUB activity of this protein, we characterized changes in

the ubiquitome of neuronal cells lacking ATXN3 (ATXN3[shRNA] cells) by mass-

spectrometry, and identified for the first time a role for ATXN3 in RNA

metabolism. We found a global perturbation of splicing in the absence of this

protein, among which a mis-splicing of tau exon 10 which perturbs the 4R/3R

tau ratio, a mechanism that may contribute for SCA3 pathogenesis, and links

two key proteins involved in different neurodegenerative disorders.

RESULTS

Changes in the ubiquitome of cells lacking ATXN3 suggest a link to RNA

metabolism

With the goal of identifying substrates of the DUB activity in neuronal cells, we

searched for changes in the ubiquitome of neuronally differentiated SH-SY5Y

cells lacking ATXN3 (ATXN3[shRNA] cells - Figure EV1A-B)[25], taking advantage

of a recently developed methodology that combines capture of the whole

spectrum of polyubiquitylated proteins in a cell extract using an enrichment by

Tandem Binding Ubiquitin Entities (TUBEs) followed by mass spectrometry

analysis (TUBEs-LC-MS/MS) [30]. Figure 1A summarizes the steps followed for

the purification and identification of the polyubiquitylated proteins in

ATXN3[shRNA] and scrambled control shRNA (SCR[shRNA]) cells. The integration

of the data and the comparison between the proteins identified in ATXN3[shRNA]

versus SCR[shRNA] cells resulted in a list of proteins with altered

polyubiquitylation in cells lacking ATXN3; among these are potential targets of

the DUB activity of ATXN3, i.e., putative ATXN3 substrates. In each pulldown

experiment, around 1200-1300 proteins were identified. When the results of

all the independent experiments were merged, we observed that many of

these proteins were only sporadically detected across the different

experiments; for the remaining analysis, these proteins were excluded,

reducing the list to 615 consistently identified proteins. From these, 193

proteins showed altered levels in the polyubiquitylated fraction in ATXN3[shRNA]

cells when compared to the SCR[shRNA] controls (p<0.05) (Table EV1). The

majority of these polyubiquitylated proteins were either not detected (44.04%)

or showed decreased levels (24.35%) in ATXN3[shRNA] cells when compared to

the controls. These proteins can be grouped into 10 functional networks, the

most representative being related to: (i) Gene expression, DNA replication,

recombination and repair (17.7%), (ii) RNA post-transcriptional modification

(13.3%), (iii) Molecular transport, RNA trafficking (10.8%), (iv) Cell death and

survival (8.9%) and (v) Organ morphology (8.9%) (Figure 1B). The fact that a

significant proportion of the proteins with altered polyubiquitylation in

ATXN3[shRNA] cells are RNA metabolism-related proteins, around 8% of them

being splicing factors (Table 1), suggested to us that ATXN3 could be playing

a role in mRNA splicing, a previously undescribed role for this DUB.

Cells lacking ATXN3 show global perturbation of splicing

We hypothesized that absence of ATXN3 could lead to a deregulation of the

pre-mRNA splicing process. To address this, we used three hybrid minigene

reporter plasmids: the pyPY minigene, representing splicing events with

alternative competing 3' acceptor sites/splice sites, the AdML minigene,

representing constitutive/strong splicing events, and the α-globin minigene, for

which the alternative splicing (exon skipping) is indicative of the performance

of regulatory splicing factors such as hnRNP and SF proteins [31,32]. As shown

in Figure 2A-D, knockdown of ATXN3 significantly altered the processing of

the three splicing reporters, suggesting a general deregulation of splicing in

the absence of this protein. Interestingly, overexpression of a catalytically inert

version of ATXN3 in SH-SY5Y cells (ATXN3_C14A), previously shown to

have a dominant-negative effect [25] also yielded similar alterations (Figure 2E).

Consistent with a global deregulation of splicing in the absence of ATXN3,

microarray analysis (Figure 2F) using specific arrays containing additional

probes for exon/exon junctions, confirmed that 1993 genes (43%), from the

7450 differentially expressed probes in ATXN3[shRNA] cells, presented

differentially regulated alternative splicing events (Table EV1). These genes

mainly encode components with function in protein degradation systems, cell

adhesion, axon guidance and various signaling pathways (Table EV2), the

majority of splicing events being decreased when compared to control cells

(Table EV3). The most affected alternative splicing event types were exon

cassette usage (34%), complex splicing (20%) and intron retention (15%)

(Figure 2G). Alternative terminal exon usage, which can be mechanistically

linked to alternative splicing, was also seen in 16% of the cases (Figure 2G).

Being a key mechanism in the regulation of gene expression, defects in these

regulatory processes may affect a multiplicity of cellular functions related to

disease.

ATXN3 modulates SRSF7 degradation by the proteasome

Interestingly, among the splicing factors with differential ubiquitylation was the

Arginine/Serine-Rich splicing factor SRSF7 (also known as 9G8), known to be

involved in the regulation of alternative splicing of the neurodegeneration-

related gene *MAPT* [33], encoding the Tau protein. As shown in Figure 3A-B,

we detected decreased levels of ubiquitylated forms of SRSF7 in cells lacking

ATXN3 (p=0.02), validating the data obtained with the proteomic analysis.

Consistent with the hypothesis that the differential ubiquitylation of SRSF7

can have an impact on its degradation, the steady state levels of total (Figure

3A-B) and nuclear (Figure 3C-D) SRSF7 were decreased in ATXN3[shRNA] cells

as compared with the SCR[shRNA] control cells (p=0.025 and p=0.001,

respectively). This suggests that SRSF7 is indeed being more degraded in the

absence of ATXN3, confirming that ATXN3 affects the expression of SRSF7

at the protein and not at the mRNA level (p=0.712) (Figure EV1C).

Based on the fact that SRSF7 was captured using Agarose-TUBEs that have

higher affinity for polyubiquitylated than for monoubiquitylated proteins [30] and

considering that ATXN3 is a DUB, we asked if ATXN3 could modify

ubiquitylation and regulate degradation of SRSF7 through the UPP.

Supporting this possibility, the levels of SRSF7 were significantly increased in

ATXN3[shRNA] cells upon proteasome inhibition with MG132 p=0.0016) (Figure

3E-F). Additionally, we detected lower levels of polyubiquitylated SRSF7

presenting Ub-K48 linkages (Figure 3G), a type of Ub chain that usually

signals proteins for degradation through the UPP. Protein stability analysis

upon inhibition of protein synthesis by cicloheximide treatment showed a

decrease in the half-life of SRSF7 in ATXN3[shRNA] cells (Figure 3H), suggesting

that ATXN3 normally acts to inhibit SRSF7 degradation in neurons. In order

for ATXN3 to modify the ubiquitylation and stability of SRSF7, the two proteins

need to interact. Co-immunoprecipitation using protein extracts from

differentiated neuron-like SH-SY5Y cells confirmed that an interaction

between ATXN3 and SRSF7 occurs in these cells (Figure 3I). This interaction

was further confirmed in live cells by the Protein Ligation Assay (Figure 3J,

Figure EV1E). Globally, our data are compatible with the hypothesis of

SRSF7 being a substrate of the DUB activity of ATXN3 in neurons.

Tau splicing is deregulated in the absence of ATXN3

Given that one of the splicing factors showing differential ubiquitylation was

SRSF7, we next analyzed mRNA levels of Tau isoforms in ATXN3[shRNA] cells

to assess the functional impact of the reduction of this splicing factor in Tau

splicing. We found a significant decrease in the transcript levels of 4R-Tau

isoform ($p=7.4\times10^{-11}$) (Figure 4A), but no alteration of the 3R-Tau isoform

($p=0.270$) in ATXN3[shRNA] cells as compared with SCR[shRNA] control cells

(Figure 4B), leading to an altered 4R/3R-Tau ratio ($p=2,98\times10^{-6}$) and

decreased total tau levels ($p=1.92\times10^{-4}$), both in ATXN3[shRNA] cells (Figure 4D)

as well as in cultured primary cerebellar neurons ($p=3.23\times10^{-9}$) in which

ATXN3 expression was silenced by shRNA (Figure EV1D, Figure 4E-H).

Consistently, catalytically inactive (Dominant negative) ATXN3 expression

also led to alterations in tau expression and in the isoform ratio similar to

those seen in cells with silenced ATXN3: decreased expression of 4R-Tau

($p=1.1\times10^{-5}$) (Figure 4I), no alterations on the 3R-Tau isoform (Figure 4J), and

consequentially a decreased 4R/3R-Tau ratio ($p=0.0088$) (Figure 4K) and

decreased levels of total Tau ($p=1.20\times10^{-6}$) (Figure 4L). To explore the

functional consequences of this reduction in 4R-Tau levels, we performed a

morphometric analysis of these cells. ATXN3[shRNA] cultures presented a

decreased expression of βIII-tubulin and a decreased average length of the

neuritis as compared with the SCR[shRNA] controls, which correlates with a more

immature stage of neuronal differentiation (Figure 4M-N).

Decreased SRSF7 expression and Tau splicing perturbation in SCA3

models and patients

To determine if the presence of an expanded polyglutamine (polyQ) tract within ATXN3 would lead to an alteration in this newly identified physiological function, we analyzed a SH-SY5Y cell line expressing a version of ATXN3 bearing 83 glutamines (ATXN3_83Q), and found a globally decreased efficiency of splicing (Figure 5A). We next asked whether the presence of the polyQ tract could also impact *MAPT* mRNA splicing regulation, altering the 4R/3R-Tau ratio. Interestingly, expression of ATXN3_83Q lead to similar changes in the expression of tau isoforms as observed in cells lacking ATXN3 or expressing a catalytic mutant version of this protein: i) decreased expression of 4R-Tau isoform ($p=6.6\times10^{-5}$) (Figure 5B), ii) no alterations on the 3R-Tau isoform (Figure 5C), iii) a decrease of the 4R/3R-Tau ratio ($p=2.4\times10^{-4}$) (Figure 5D), and (iv) a decrease in the levels of total Tau ($p=7.6\times10^{-6}$) (Figure 5E).. Of notice, the overexpression of the wild type ATXN3 (ATXN3_28Q) also caused some (albeit less important) degree of perturbation (Figure 5 A-E), pointing to the importance of a tight regulation of ATXN3 dosage in neurons.

Tau splicing was also analyzed in SCA3 mouse model (CMVMJD135) that reproduces key features of the disease [34]. A decreased expression of the 4R-Tau isoform was seen in the brainstem (affected brain region) of CMVMJD135 mice (Tg) ($p=7\times10^{-5}$) comparing with the Wild type (Wt) controls (Figure 5F), with no alterations of the 3R-Tau isoform expression ($p=0.84$) (Figure 5G), leading to an altered 4R/3R-Tau ratio ($p=1.2\times10^{-3}$) (Figure 5H) and decreased

levels of total Tau ($p=1.1\times10^{-8}$) (Figure 5I). In addition, SRSF7 protein levels

were also reduced in the brainstem (p=0.0093), spinal cord (p=0.0014) and to

a lesser extent in the cerebellum (p=0.1138) (disease-affected brain areas) of

Tg mice as compared with the Wt littermate controls (Figure 5J-L,

respectively), but no alterations at the mRNA level were detected (Figure

EV1F). This indicates that the impact of ATXN3 on tau splicing through the

control of SRSF7 levels is also occurring in SCA3 and is corroborated by the

decreased levels of SRSF7 protein present in the brains of SCA3 patients as

compared with healthy controls (Figure 5M). To explore the functional

consequences of this reduction in 4R-Tau levels, an isoform of Tau

associated with mature neurons and synaptic function, we performed a

morphometric analysis of neurons in this brain region. A decreased mean

neurite length (p=0.0189) (Figure 5N-O) and a decrease in complexity of the

dendritic trees (p=0.0004) (Figure O, P) were observed in neurons from the

Medial vestibular nuclei in the Brainstem.

Altogether, these results suggest that the functional interaction of ATXN3 with

SRSF7 and its impact on molecular mechanisms regulating tau splicing in

neurons may be relevant in the pathogenesis of SCA3.

DISCUSSION

Ubiquitylation is a post-translational modification that controls several aspects

of neuronal function by regulating protein abundance. Disruption of this

signaling pathway has been described in neurodegenerative diseases.

Indeed, since many of these diseases exhibit ubiquitylated protein

aggregates, loss of ubiquitin homeostasis may be an important contributor for

disease. Considering the importance of DUB enzymes in maintaining the ubiquitylation balance, we focused on the characterization of the ubiquitome of neuronal cells lacking ATXN3 [25]. Using TUBEs, that enable the pulldown of polyubiquitylated proteins without further genetic manipulation or inhibition of the proteasome [30] in combination with LC-MS/MS, we were able to consistently identify 615 proteins per condition, a yield comparable to those described in other studies [35,36]. Among the proteins identified, approximately one third presented altered levels of polyubiquitylation in ATXN3[shRNA] cells. Curiously, the majority of these proteins presented decreased levels in the polyubiquitin-enriched fraction signal, suggesting that normally ATXN3 might be preventing their degradation, for instance by editing the substrate's ubiquitin chains, as we observed before for ITGA5 [24,25]. Therefore, when ATXN3 is silenced, the polyubiquitin signal is not removed, which may result in an increased degradation of the targeted proteins. The fact that a significant proportion of the proteins with altered polyubiquitylation levels in ATXN3[shRNA] cells were splicing factors and proteins involved in RNA processing, is suggestive of ATXN3 playing a role in the pre-mRNA splicing process in neuronal cells. This process is particularly important in generating diversity and specificity in the central nervous system, which requires a wide protein repertoire to generate its highly specialized and adaptable neuronal circuits [37]. Given the importance of RNA processing for neuronal function, it comes as no surprise that altered RNA processing is a contributing factor to the pathogenesis of multiple neurodegenerative diseases [38-42]. Disruption of alternative slipicing (AS) regulation, resulting from mutations or abnormal expression of RNA-binding proteins, can lead to an imbalance or an

inappropriate expression pattern of key protein isoforms, with profound

consequences on cellular and organismal physiology. Indeed, the relative

concentration of splicing factors and hnRNPs, which may be regulated by the

Ub pathway, has been shown to regulate AS [43]. For example, it was shown

that the abundance of several splicing factors is differentially affected by

proteasome inhibition, and abnormal levels of these regulatory proteins were

associated with a different AS pattern [44]. Another study showed that the

splicing factor SRSF5 is targeted to proteasomal degradation as the cells

undergo terminal differentiation [45]. Consistently, assessed general splicing

efficacy was decreased in cells depleted of ATXN3, as shown using artificial

reporter minigenes, further supporting the involvement of ATXN3 in the

regulation of the splicing process. Additionally, genome wide microarray

analysis of endogenous splicing events confirmed that absence of ATXN3

leads to a dramatic alteration in the splicing pattern of a large number of

genes in neuronal cells, including genes related to protein degradation,

adhesion, axon guidance and signaling pathways. This is consistent with our

previous findings showing that absence of ATXN3 leads to an impairment in

neuronal differentiation and adhesion [25], and affects the degradation of

several target proteins [23-25]. Significantly, a portion of the predicted splicing

regulators of these genes with altered splicing in cells lacking ATXN3 were

found to have altered levels of polyubiquitylation in our proteomic analysis, the

majority of them being Serine/Arginine (SR) – rich phosphoproteins. Proteins

of the SR family are key players in the control of AS, regulating the selection

of alternative sites [46,47]. The fact that SRSF7 (9G8), one of those proteins and

a key regulator of MAPT (tau) exon 10 splicing [48,49], was captured using the

Agarose-TUBEs, which have high affinity for polyubiquitin chains [30],

suggested that this splicing factor is polyubiquitylated, and thus might be

degraded through the UPP. Indeed, while it has been described that the

mature human neurons express approximately equal levels of 4R and 3R-Tau

isoforms [50,51], whose expression is developmentally controlled and finely-

tuned, we found that loss of function of ATXN3 disrupts this balance in

neuronal cells leading to a decreased 4R/3R-Tau ratio, that coincides with

(and may be caused by) decreased levels of SRSF7. Very recent works have

shown imbalanced expression in 3R and 4R-Tau, namely a decreased

expression of the 4R isoform as here observed, in the brain of a mouse model

of Down syndrome, but also in human AD and PD patients [52-54], strengthening

the idea that a faulty tau exon 10 splicing might be contributing for the

pathogenesis of several neurodegenerative diseases. In addition, the

decreased levels of the 4R-Tau isoform presented by ATXN3[shRNA] cells, also

correlate with the immature phenotype of these cells, that we previously

described [25]. Indeed Conejero-Godberg and colleagues also assessed the

transcriptional profiles related to 3R and 4R-Tau splice variants, revealing that

a cluster of genes related to neuronal cell morphology and outgrowth of

neurites is upregulated in the presence of 4R isoforms, while genes related to

cellular growth and proliferation were downregulated [55]. This is in agreement

with previous work from Sennvik's team establishing a role for 4R-Tau

isoforms in promoting neuronal differentiation [56]. Taking into account that loss

of function of ATXN3 leads to a deregulation in the amount of SRSF7, at the

protein level but not RNA level, we hypothesized that this splicing factor could

be a substrate of the DUB activity of ATXN3, which could be modulating its

degradation. In line with that, we confirmed that ATXN3 interacts closely with

SRSF7 in neuronal cells, suggesting that these proteins are molecular

partners in such cells, and establishing a functional link between two key

proteins involved in different neurodegenerative diseases. Trying to explore if

this interaction could be indicative that ATXN3 is modulating the degradation

of SRSF7 through the proteasome, we measured the levels of this protein

upon proteasome inhibition; as expected, we saw an accumulation of SRSF7

when the proteasome was inhibited. Interestingly, we also found a decreased

4R/3R-Tau ratio, as well as decreased protein levels of SRSF7, both in cells

expressing the mutant protein, in affected brain regions of our mouse model

for the disease, at an age when the transgenic animals display an overt

phenotype resembling behavioral and pathological characteristics present in

human patients [34]. This suggests that the mis-splicing of tau may be

contributing to SCA3 pathogenesis, as it does to other neurodegenerative

diseases. Considering that we observed a similar phenomenon using cells

lacking ATXN3 and cells expressing a catalytic inactive mutant form of the

protein, it is reasonable to think that the DUB activity of ATXN3 is important in

this cellular process and that the polyglutamine expansion might cause a

partial loss of this normal function of the protein (even in presence of 50% of

the wild type protein), perhaps through a dominant negative mechanism, thus

contributing to SCA3 pathogenesis. Indeed, Winborn and co-workers

demonstrated that while expression of ataxin-3 with a non-pathogenic polyQ

tract leads to a reduction in total ubiquitylated protein levels in cells, this

reduction is not observed upon overexpression of an expanded ataxin-3,

which may reflect inefficient deubiquitylation of cellular proteins, pointing to an

overall loss of function [57]. Overall, our data suggest that ATXN3 is involved in

splicing regulation through the modulation of the ubiquitylation of splicing

factors. This regulatory role may be mediated through the DUB activity of

ATXN3 by modulating activation, degradation and/or subcellular localization of

the splicing factors, or may be an indirect result of the modulation of E3

ligases specific for these targets.

Recently, several pieces of evidence coming from the use of technologies

such as exon arrays and RNA-Seq suggested an association between

perturbation of AS and several neurodegenerative diseases including

Alzheimer's disease, Huntington's disease, Pick's disease, Spinal Muscular

atrophy and Amyotrophic Lateral Sclerosis [58-60]. Importantly, several ataxia-

causing proteins have been described to interact with splicing factors [61]. Also,

expression of mutant ATXN3 bearing an expanded polyglutamine tract was

previously shown to alter the ability of the subnuclear domains known as Cajal

bodies to efficiently participate in small nuclear ribonucleoprotein (snRNP)

biogenesis pathway, and to reduce the efficacy of splicing reporter genes [62].

These findings lead us to propose that ATXN3 plays a role in splicing

regulation in neurons, a novel function for this protein, and that a perturbation

of this physiological function might contribute for SCA3 pathogenesis.

MATERIALS AND METHODS

Cell lines and primary cultures

SH-SY5Y cell cultures: SH-SY5Y human female neuroblastoma cell line

(ATCC number CRL-2266) was cultured in Dulbecco's modified eagle

medium: nutrient mixture (DMEM)/F-12 supplemented with 10% (v/v) fetal

bovine serum (FBS) (Biochrom), 2mM glutaMAX (Invitrogen), 100 U/mL

penicillin (Invitrogen), 100 μg/mL streptomycin (Invitrogen) and 25 ng/mL

puromycin (Sigma Aldrich). Medium was changed every 2 days. Cells were

transfected with a shRNA sequence targeting *ATXN3* or a scrambled shRNA

sequence as described elsewhere [25]. Differentiation of stably infected cell

lines was induced by 0.1 μM all-trans-retinoic acid (RA) (Sigma Aldrich) in

opti-MEM (Invitrogen) supplemented with 0.5% FBS for 7 days. Medium was

replaced every 2 days. *Primary cultures of cerebellum:* cerebellar neuron

cultures were prepared from P7 Wistar rats. Briefly, upon dissection, cerebella

were submitted to a trypsin-based enzymatic digestion followed by

mechanical dissociation. Isolated cells were plated on poly-d-lysine (Sigma

Aldrich) and lysine (Sigma Aldrich) pre-coated 6-well plates at a density of

1.25×10^6 cells/mL using Neurobasal A (Invitrogen) supplemented with 1x B27

(Invitrogen), 0.1 mg/mL Kanamycin (Invitrogen), 1 mM Glutamax I (Invitrogen)

and 10 ng/mL bFGF (Invitrogen) for 7 days. Half of the medium was changed

every 4 days.

Mice

Female CMVMJD135 (background C57BL/6) mice were used in the study.

Transgenic (Tg) and wild-type (Wt) littermates were housed at weaning in

groups of 5 animals in filter-topped polysulfone cages 267x207x140 mm (370

cm^2 floor area) (Tecniplast), with corncob bedding (Scobis Due, Mucedola

SRL) in a conventional animal facility. The mice were genotyped at weaning

by PCR, as we previously described [63]. All animals were maintained under

standard laboratory conditions: an artificial 12 h light/dark cycle (lights on from

8:00 to 20:00 h), with an ambient temperature of 21±1ºC and a relative

humidity of 50-60%; the mice were given a standard diet (4RF25 during the

gestation and postnatal periods, and 4RF21 after weaning (Mucedola SRL)

and water *ad libitum*. Animal facilities and the people directly involved in

animal experiments were certified by the Portuguese regulatory entity –

Direcção Geral de Veterinária. Health monitoring was performed according to

FELASA guidelines[64], confirming the Specified Pathogen Free status of

sentinel animals maintained in the same animal room. Humane endpoints for

the experiment were defined (20% reduction of the body weight, inability to

reach food and water, presence of wounds in the body, dehydration). The

animals were euthanized at 45 weeks of age by decapitation, the brain

regions of interest were macro dissected and store at -80ºC until further

processing.

Pulldown of polyubiquitylated proteins

Cells treated with RA were lysed by sonication on ice in lysis buffer (50 mM

Tris-HCl pH 7.5, 0.15 M NaCl, 1mM EDTA, 1% NP-40, 10% Glycerol,

complete protease inhibitors (Roche) and 50 µM UB/UBl protease inhibitor

PR-619 (LifeSensors). After lysis, 2 mg of total protein extract was incubated

with 100 µL of pre-equilibrated Agarose-TUBEs (LifeSensors), overnight at

4ºC on a rocking platform. Sedimented beads were washed 3 times with

washing buffer (20 mM Tris pH 8.0, 0.15 M NaCl, 0.1% Tween-20) before

being eluted with 1x SDS sample buffer (62.5mM Tris-HCl pH 6.8, 10% glycerol, 2% SDS, Bromophenol Blue). Eluted proteins were immediately boiled at 98ºC for 15min and ran in a 10% SDS-PAGE gel. After incubation with the FK2 anti-ubiquitin primary antibody (1:2000, Millipore) overnight at 4ºC, membranes were incubated with secondary antibody for 1 hour at room temperature (anti-mouse, 1:10.000, Bio-Rad). Antibody binding was detected by chemiluminescence (Clarity kit, Bio-Rad).

Digestion of proteins from preparative 1D-PAGE gel

The 1D PAGE LC-MS/MS approach was used for protein identification as previously described [65]. Eluted proteins were separated using 1.5 mm and 10% SDS-PAGE gels. The quality of purification was controlled by Coomassie Brilliant Blue g-250 (Sigma) staining before MS analysis. Gel image was acquired the Gel Doc™ EZ system (Bio-rad). After Coomassie staining, all the visible blue-stained protein spots were manually excised from the gel. The gel pieces were destained overnight at room temperature using 50% acetonitrile in 25 mM ammonium bicarbonate buffer, pH 8.5, and then dehydrated with 100% acetonitrile. The shrunken pieces were then re-swollen in 50 mM ammonium bicarbonate buffer, dehydrated in 100% acetronitrile and dried in a speedvac® concentrator (Savant) for 30 min. The gel pieces were rehydrated in 60 µL of 20 µg/mL Trypsin (Promega) in 50 mM ammonium bicarbonate solution and incubated for 2h at 55ºC. The gel pieces were then incubated with 0.1% trifluoroacetic acid in 50% acetonitrile for 20 min at room temperature in order to extract the remaining peptides from the gel. The tryptic peptides were dried in a speedvac for 2 h.

Liquid chromatography-tandem mass spectrometry (LC-MS/MS)

After re-dissolution in 17 μL 0.1% acetic acid, samples were separated on a capillary C18 column using a nano LC-ultra 1D plus HPLC system (Eksigent) and analyzed on-line with an electrospray LTQ-Orbitrap Discovery mass spectrometer (Thermo Fisher Scientific). MS/MS spectra were searched against a human database (uniprot_sprot, 2010_01) with the ProteinPilot™ software (version 3.0; AB-sciex) using the Paragon™ algorithm (version 3.0.0.0 [66]) as the search engine. The detected protein threshold (unused protscore (confidence)) in the software was set to 0.10 to achieve 20% confidence and the proteins identified were grouped to minimize redundancy. Peptides with "unused" values < 2 have low confidence and were excluded from analysis. The "unused" value is defined in the handbook of ProteinPilot as a sum of peptide scores from all the non-redundant peptides matched to a protein. Peptides with confidence of ≥ 99% would have a peptide score of 2. Tryptic peptides shared by multiple proteins were assigned to the winner protein.

RNA extraction and Array hybridization

Total RNA was isolated from ATXN3shRNA and SCRshRNA cells using an miRNeasy mini kit (Qiagen) and quality assessment was achieved using RNA 6000 Nano labchip (Bioanalyzer, Agilent) and by a Nanodrop spectrophotometer (Thermo). Total RNAs RIN values were between 8.7 and 9.3 (average: 9.17). Affymetrix Human Transcriptome Array 2.0 ST arrays were hybridized according Affymetrix recommendations using the Ambion WT

protocol (Life technologies, France) and Affymetrix labelling and hybridization

kits. Raw data, transcript data and exon data were controlled with Expression

console (Affymetrix).

Microarray data analysis

Affymetrix Human Transcriptome Array 2.0 ST dataset analysis was

performed using the GenoSplice technology (www.genosplice.com). Data was

normalized using quantile normalization. Background corrections were made

with antigenomic probes and probes were selected according to their %GC,

cross-hybridization status and potential overlap with repeat region as

previously described [PMID:23861464, PMID:23321315, PMID:23284676].

Only probes targeting exons and exon-exon junctions annotated from FAST

DB® transcripts (release fastdb_2013_2) were selected [PMID:16052034,

PMID:17547750]. Only probes with a DABG P value ≤0.05 in at least half of

the arrays were considered for statistical analysis [PMID:23861464,

PMID:23321315, PMID:23284676]. Only genes expressed in at least one

compared condition were analyzed. To be considered to be expressed, the

DABG P-value had to be ≤0.05 for at least half of the gene probes. We

performed an unpaired Student's *t-test* to compare gene intensities between

ATXN3^{shRNA} and SCR^{shRNA} cells. Genes were considered significantly

regulated when fold-change was ≥1.5 and P-value ≤0.05 (unadjusted P-

value). Analysis at the splicing level was first performed taking into account

only exon probes ('EXON analysis) in order to potentially detect new

alternative events that could be differentially regulated (i.e., without taking into

account exon-exon junction probes). Analysis at the splicing level was also

performed by taking into account exon-exon junction probes ('SPLICING PATTERN analysis) using the FAST DB® splicing pattern annotation (i.e., for each gene, all possible splicing patterns were defined and analyzed. All types of alternative events can be analyzed: alternative first exons, alternative terminal exons, cassette exon, mutually exclusive exons, alternative 5' donor splice site, alternative 3' acceptor splice sites and intron retention). EXON and SPLICING PATTERN analyses were performed using unpaired Student's *t-test* on the splicing-index as previously described [PMID:23861464, PMID:23321315, PMID:23284676]. Results were considered statistically significant for unadjusted P-values ≤ 0.05 and fold-changes ≥ 1.5 for SPLICING PATTERN analysis and unadjusted P-values ≤ 0.05 and fold-changes ≥ 2.0 for EXON analysis. Gene Ontology (GO), KEGG and REACTOME analyses of differentially regulated genes were performed using DAVID [PMID:19131956].

Plasmid purification

Hybrid minigene reporter plasmids pyPY, AdML and α-globulin [31,32] were kindly provided by Prof. Juan Valcárcel (Centre de Regulació Genòmica – CRG, Barcelona). Top10 competent cells (Invitrogen) were transformed with 100 ng of plasmid DNA, according to the recommended protocol. Briefly, the cells were incubated with the constructs on ice for 30 min followed by heat shock at 42ºC for 1 min. After incubation on ice for 2 min, 500 μL of LB medium was added to the cell vial and incubated at 150 rpm for 60 min at 37ºC. Cultures were grown overnight at 37ºC in LB/ampicillin plates. The next day, one colony was inoculated in LB/ampicilin (100 mg/mL) at 37ºC

overnight. Plasmid extraction was carried out using the ZR Plasmid

Miniprep™ (Zymo Research) according with the manufacturer's protocol. DNA

concentration was determined using Nanodrop (Alfagene) and integrity

verified by running 200 ng in an agarose gel.

Cell transfection

4×10^5 cells per well were plated in gelatin-coated 6 well plates and incubated

for 24 h. Before transfection, the culture medium was changed to DMEM/F-

12-AA without antibiotics and supplemented with 5% FBS. Cells were

transfected with 200 ng of the reporter plasmids using Lipofectamine® 2000

Transfection Reagent (Invitrogen) according with the manufacturer's

instructions. Briefly, reporter plasmid minigenes and the transfection reagents

were appropriately diluted in Opti-MEM medium separately and incubated for

5 min at room temperature. The mixed reagents were then incubated at room

temperature for 20 min allowing the formation of transfection complexes. The

cells were then incubated for 24 h with the transfection mix.

Semi-quantitative PCR

PCR amplification of pyPY, AdML and α-globin reporter genes was carried out

using Taq DNA Polymerase (Thermo Fisher Scientific) following the

manufacturer's protocol. The cycling conditions were: 95ºC for 5 min followed

by 24 cycles of denaturing at 95ºC for 1 min, annealing at 60ºC for 45 sec,

extension at 72ºC for 1 min and final extension at 72ºC for 5 minutes. The

primers used are listed in Table S1. The PCR product was ran in a 2%

agarose gel. Gel analyses and splicing efficiency calculations were performed

using Image Lab software (Bio-Rad).

Cell fractionation

RA-treated pelleted cells were resuspended in ice RSB buffer (10 mM Tris-

HCl pH 7.4, 10 M NaCl) and incubated on ice 3 min. After centrifugation for 5

min, 4000 rpm at 4ºC, the pellet was resuspended in RSBG40 buffer (10 mM

Tris-HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 10% glycerol, 0.5% NP-40, 0,5

mM DTT) and centrifuged for 3 min, 7000 rpm at 4ºC. The supernatant was

collected as cytoplasmic fraction and stored at -80ºC. The nuclear pellet was

resuspended in 50 µL B1 buffer (20 mM Tris-HCl pH 7.9, 75 mM NaCl, 5%

glycerol, 0.5 mM EDTA, 0.85 mM DTT, 0.125 mM PMSF) and 450 µL B2

buffer (20 mM HEPES pH 7.6, 300 mM NaCl, 0.2 mM EDTA, 1 mM DTT, 7.5

mM $MgCl_2$, 1 M Urea, 1% NP-40). Samples were vortexed for 5 sec,

incubated on ice for 10 min and centrifuged for 5 min, 15000 rpm at 4ºC. The

supernatant was collected as nuclear fraction and stored at -80ºC.

Protein synthesis inhibition and proteasome inhibition

RA-treated SH-SY5Y cells were incubated with 5 µM cicloheximide (Merck)

during 20, 60 and 120 minutes. For proteasome inhibition, RA-treated cells

were incubated with 5 µM MG132 (Calbiochem) for 24h prior to lysis.

High-throughput high-content functional imaging

SH-SY5Y cells were seeded at a density of $4x10^3$ cells/well in flat bottom 96-

well plates previously coated with Matrigel (BD, Biosciences), and 10 µM all-

trans-retinoic acid (Sigma Aldrich) was added the day after plating in DMEM-

F12 with 1% FBS, to induce differentiation. After 5 days, cells were washed in

DMEM/F-12 and incubated with 50 ng/mL BDNF (Prepotech) in DMEM/F-12

without serum for 3 days. Cells were then labeled for β-III tubulin (1:1000,

R&D Systems), scanned at different locations of each well and the

quantitative analysis of neurite length were automatically done using the

automatic imaging system Thermo Scientific Cellomics® ArrayScan® VTI.

Immunoprecipitation

RA-treated cells were washed in ice-cold PBS and lysed by sonication on ice

in NP-40 buffer. Aliquots were taken from protein extracts as 10% inputs. 1

mg of total protein was pre-cleared for 3 h at 4ºC by incubation with

glutathione-coupled sepharose beads (GE Healthcare) previously equilibrated

in Wash buffer 1 (50 mM Tris-HCl pH 7.5, 150 nM NaCl, 1% NP-40, 1x

protease inhibitors (Roche)) for 3 times, 10 min at 4ºC. Beads were then

centrifuged and the supernatant was incubated O/N at 4ºC with 50 µL

equilibrated beads. After centrifugation, the supernatant was discarded and

the beads were washed 2 times with Wash buffer 1 for 10 min at 4ºC. Beads

were then washed twice with Wash buffer 2 (50 mM Tris-HCl pH 7.5, 500 mM

NaCl, 0.1% NP-40) for 10 min at 4ºC, and once with Wash buffer 3 (50 mM

Tris-HCL pH 7.5, 0.1% NP-40) for 20 min at 4ºC. The supernatant was

discarded and the bound proteins were eluted with 1x Laemmli buffer, boiled

for 5 min at 98ºC and run in a 10% SDS-PAGE gel.

Proximity Ligation Assay (PLA)

SH-SY5Y cells were cultured onto polyD-lysine coated cover slips in 24-well

plates using D-MEM media as indicated to above. PLA protocol was

performed as described in [67] and following manufacturer instructions with

adaptations. Cells were fixed with 4% PFA, and permeabilized with 0.5%

Triton in PBS at room temperature. Blocking was performed at 37ºC in a

humidity chamber for 1 hour, using Duolink® PLA Blocking Solution (Sigma).

Primary antibodies were incubated over night at 4ºC followed by two washing

steps of 5 minutes with Wash Buffer A (Sigma) at room temperature. PLUS

and MINUS PLA probes were diluted 1:5 in the Duolink Antibody Diluent

(Sigma), added to the cover slips and incubated for 1h at 37ºC in a humidity

chamber. Washing was performed for 5 minutes two times with Washing

Buffer A at room temperature followed by incubation with Ligase solution

(Sigma) for 30 minutes at 37ºC in a humidity chamber. After washing two

times during 2 minutes with Washing Buffer A at room temperature, cover

slips were incubated with Amplification solution (Sigma) for 100 minutes at

37ºC in a humidity chamber protected from light. Then, cover slips were

washed two times during 10 minutes with Washing Buffer B (Sigma) at room

temperature followed by an additional washing step of 1 minute using 100X

diluted Washing Buffer B. Cover slips were mounted on a glass slide with

mounting media and DAPI and cells analyzed by confocal microscopy.

Immunoblotting

RA-treated cells were pelleted and frozen in liquid nitrogen. For cellular and

brain tissue extracts, 50 µg of total protein isolated in NP-40 buffer (150 mM

NaCl, 50 mM Tris-HCl pH 7.6, 0.5% NP-40, protease inhibitors (Roche)) were

resolved in 10% SDS-PAGE gels and then transferred to a nitrocellulose

membrane. After incubation with the primary antibodies: Tau-5 (1:2000,

Abcam), Tau-4R (1:1000, Millipore), Tau-3R (1:2000, Millipore), Tau (E178)

(1:1000, Abcam), Histone H3 (1:10.000, Abcam), SFRS7(9G8) (1:1000, kindly

provided by Dr. James Stévine), Actin (1:200, DSHB) overnight at 4ºC,

membranes were incubated with secondary antibodies for 1 hour at room

temperature (anti-rabbit or anti-mouse, 1:10.000, Bio-Rad, anti-goat, 1:7500,

Pierce. Antibody binding was detected by chemiluminescence (Clarity kit, Bio-

Rad).

qRT-PCR

1 µg of total RNA purified from differentiated cells or brain tissue samples was

reversed transcribed using the One-step SuperScript kit (Bio-Rad). The qRT-

PCR reaction was performed in a CFX96 Real-time PCR detection system

(Bio-Rad) using the Quantitec SYBR Green kit (Qiagen) and the primers listed

on Table S1. Gene expression was normalized to *HMBS* or *B2M* levels within

each sample. Results are presented as fold change.

Neurostructural analysis

Mice were transcardially perfused with 0.9 % saline (n=9 per group) under

deep anesthesia (mixture of ketamine hydrochloride (150 mg/kg) plus

medetomidine (0.3 mg/kg)). Briefly, brains were immersed in Golgi-Cox

solution[68] for 14 days, processed and cut on a vibratome at 200 µm thick

coronal sections. For each selected neuron, all branches of the dendritic tree

were reconstructed at x1000 magnification, using a motorized microscope

(Zeiss, Thornwood) attached to a camera (Sony) and Neurolucida software

(Microbrightfield). A three-dimensional analysis of the reconstructed neurons

was performed using NeuroExplorer software (Microbrightfield).

Quantification and statistical analysis

Continuous variables with normal distributions (Shapiro-Wilk test p>0.05)

were analyzed with the Student *t-test* or two-way ANOVA. Non-parametric

Mann-Whitney U-test was used when variables were non-continuous or when

a continuous variable did not present a normal distribution (Shapiro-Wilk test

p<0.05). All statistical analysis were performed using SPSS 22.0 (SPSS Inc.,

Chicago, IL). A critical value for significance of p<0.05 was used throughout

the study. For real-time quantitative PCR data, the same approach was used

and results were presented using the ΔΔCt method, as described before [69].

AUTHOR CONTRIBUTIONS

Conceptualization, ANC and PM; Methodology, ANC, PM, IS, SH, PH and

KWL; Formal analysis, ANC and KWL; Investigation, ANC, BA, SDS, JS, SH

and KWL; Writing – original draft, ANC; Writing – review & editing, ANC, PM,

IS, PH and KWL; Visualization, ANC; Funding acquisition, PM; Resources,

PM, KWL, PH; Supervision, PM

ACKNOWLEDGMENTS

This article has been developed under the scope of the project NORTE-01-

0145-FEDER-000013, supported by the Northern Portugal Regional

Operational Programme (NORTE 2020), under the Portugal 2020 Partnership

Agreement, through the European Regional Development Fund (FEDER).

This work has been funded by FEDER funds, through the Competitiveness

Factors Operational Programme (COMPETE), and by National funds, through

the Foundation for Science and Technology (FCT), under the scope of the

project POCI-01-0145-FEDER-007038 and through the project POCI-01-

0145-FEDER-016818 (PTDC/NEU-NMC/3648/2014) and fellowships

[SFRH/BD/51059/2019 to A. N-C, SFRH/BPD/110728/2015 to B.A,

SFRH/BD/78388/2011 to S.D-S, SFRH/BD/88932/2012 to J.M.S and

IF/01799/2013 to I.S.

CONFLIT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Hegde, A.N. & Upadhya, S.C. Role of ubiquitin-proteasome-mediated

proteolysis in nervous system disease. *Biochim Biophys Acta* **1809**, 128-

140 (2011).

2. Yi, J.J. & Ehlers, M.D. Emerging roles for ubiquitin and protein degradation

in neuronal function. *Pharmacol Rev* **59**, 14-39 (2007).

3. Baptista, M.S., Duarte, C.B. & Maciel, P. Role of the ubiquitin-proteasome

system in nervous system function and disease: using C. elegans as a

dissecting tool. *Cell Mol Life Sci* **69**, 2691-2715 (2012).

4. Hershko, A. & Ciechanover, A. The ubiquitin system. *Annu Rev Biochem* **67**,

425-479 (1998).

5.   Komander, D. The emerging complexity of protein ubiquitination.

729   *Biochem Soc Trans* **37**, 937-953 (2009).

730   6.   Tai, H.C. & Schuman, E.M. Ubiquitin, the proteasome and protein

731   degradation in neuronal function and dysfunction. *Nat Rev Neurosci* **9**,

732   826-838 (2008).

733   7.   de Vrij, F.M., Fischer, D.F., van Leeuwen, F.W. & Hol, E.M. Protein quality

734   control in Alzheimer's disease by the ubiquitin proteasome system. *Prog*

735   *Neurobiol* **74**, 249-270 (2004).

736   8.   Upadhya, S.C. & Hegde, A.N. Ubiquitin-proteasome pathway components

737   as therapeutic targets for CNS maladies. *Curr Pharm Des* **11**, 3807-3828

738   (2005).

739   9.   Rubinsztein, D.C. The roles of intracellular protein-degradation pathways

740   in neurodegeneration. *Nature* **443**, 780-786 (2006).

741   10.   Nijman, S.M.*, et al.* A genomic and functional inventory of deubiquitinating

742   enzymes. *Cell* **123**, 773-786 (2005).

743   11.   Komander, D., Clague, M.J. & Urbe, S. Breaking the chains: structure and

744   function of the deubiquitinases. *Nat Rev Mol Cell Biol* **10**, 550-563 (2009).

745   12.   Clague, M.J., Coulson, J.M. & Urbe, S. Cellular functions of the DUBs. *J Cell*

746   *Sci* **125**, 277-286 (2012).

747   13.   Rappsilber, J., Ryder, U., Lamond, A.I. & Mann, M. Large-scale proteomic

748   analysis of the human spliceosome. *Genome Res* **12**, 1231-1245 (2002).

749   14.   Makarov, E.M.*, et al.* Small nuclear ribonucleoprotein remodeling during

750   catalytic activation of the spliceosome. *Science* **298**, 2205-2208 (2002).

751   15.   Wagner, S.A*., et al.* A proteome-wide, quantitative survey of in vivo

752   ubiquitylation sites reveals widespread regulatory roles. *Mol Cell*

753   *Proteomics* **10**, M111 013284 (2011).

754   16.   Peng, J*., et al.* A proteomics approach to understanding protein

755   ubiquitination. *Nat Biotechnol* **21**, 921-926 (2003).

756   17.   Bellare, P., Kutach, A.K., Rines, A.K., Guthrie, C. & Sontheimer, E.J. Ubiquitin

757   binding by a variant Jab1/MPN domain in the essential pre-mRNA

758   splicing factor Prp8p. *RNA* **12**, 292-302 (2006).

759   18.   Kramer, A., Mulhauser, F., Wersig, C., Groning, K. & Bilbe, G. Mammalian

760   splicing factor SF3a120 represents a new member of the SURP family of

761   proteins and is homologous to the essential splicing factor PRP21p of

762   Saccharomyces cerevisiae. *RNA* **1**, 260-272 (1995).

763   19.   Lygerou, Z., Christophides, G. & Seraphin, B. A novel genetic screen for

764   snRNP assembly factors in yeast identifies a conserved protein, Sad1p,

765   also required for pre-mRNA splicing. *Mol Cell Biol* **19**, 2008-2020 (1999).

766   20.   Makarova, O.V., Makarov, E.M. & Luhrmann, R. The 65 and 110 kDa SR-

767   related proteins of the U4/U6.U5 tri-snRNP are essential for the assembly

768   of mature spliceosomes. *EMBO J* **20**, 2553-2563 (2001).

769   21.   Burnett, B., Li, F. & Pittman, R.N. The polyglutamine neurodegenerative

770   protein ataxin-3 binds polyubiquitylated proteins and has ubiquitin

771   protease activity. *Hum Mol Genet* **12**, 3195-3205 (2003).

772   22.   Doss-Pepe, E.W., Stenroos, E.S., Johnson, W.G. & Madura, K. Ataxin-3

773   interactions with rad23 and valosin-containing protein and its

774   associations with ubiquitin chains and the proteasome are consistent

with a role in ubiquitin-mediated proteolysis. *Mol Cell Biol* **23**, 6469-6483

(2003).

23. Rodrigues, A.J.*, et al.* Absence of ataxin-3 leads to cytoskeletal

disorganization and increased cell death. *Biochim Biophys Acta* **1803**,

1154-1163 (2010).

24. do Carmo Costa, M.*, et al.* Ataxin-3 plays a role in mouse myogenic

differentiation through regulation of integrin subunit levels. *PLoS One* **5**,

e11728 (2010).

25. Neves-Carvalho, A.*, et al.* Dominant negative effect of polyglutamine

expansion perturbs normal function of ataxin-3 in neuronal cells. *Hum*

Mol Genet **24**, 100-117 (2015).

26. Zhong, X. & Pittman, R.N. Ataxin-3 binds VCP/p97 and regulates

retrotranslocation of ERAD substrates. *Hum Mol Genet* **15**, 2409-2420

(2006).

27. Wang, G., Sawai, N., Kotliarova, S., Kanazawa, I. & Nukina, N. Ataxin-3, the

MJD1 gene product, interacts with the two human homologs of yeast DNA

repair protein RAD23, HHR23A and HHR23B. *Hum Mol Genet* **9**, 1795-

1803 (2000).

28. Chatterjee, A.*, et al.* The role of the mammalian DNA end-processing

enzyme polynucleotide kinase 3'-phosphatase in spinocerebellar ataxia

type 3 pathogenesis. *PLoS Genet* **11**, e1004749 (2015).

29. Gao, R.*, et al.* Inactivation of PNKP by mutant ATXN3 triggers apoptosis by

activating the DNA damage-response pathway in SCA3. *PLoS Genet* **11**,

e1004834 (2015).

30.   Hjerpe, R*., et al.* Efficient protection and isolation of ubiquitylated

800   proteins using tandem ubiquitin-binding entities. *EMBO Rep* **10**, 1250-

801   1258 (2009).

802   31.   Guth, S., Martinez, C., Gaur, R.K. & Valcarcel, J. Evidence for substrate-

803   specific requirement of the splicing factor U2AF(35) and for its function

804   after polypyrimidine tract recognition by U2AF(65). *Mol Cell Biol* **19**,

805   8263-8271 (1999).

806   32.   Pacheco, T.R., Coelho, M.B., Desterro, J.M., Mollet, I. & Carmo-Fonseca, M.

807   In vivo requirement of the small subunit of U2AF for recognition of a

808   weak 3' splice site. *Mol Cell Biol* **26**, 8183-8190 (2006).

809   33.   Cavaloc, Y., Popielarz, M., Fuchs, J.P., Gattoni, R. & Stevenin, J.

810   Characterization and cloning of the human splicing factor 9G8: a novel 35

811   kDa factor of the serine/arginine protein family. *EMBO J* **13**, 2639-2649

812   (1994).

813   34.   Silva-Fernandes, A*., et al.* Chronic treatment with 17-DMAG improves

814   balance and coordination in a new mouse model of Machado-Joseph

815   disease. *Neurotherapeutics* **11**, 433-449 (2014).

816   35.   Lopitz-Otsoa, F*., et al.* Integrative analysis of the ubiquitin proteome

817   isolated using Tandem Ubiquitin Binding Entities (TUBEs). *J Proteomics*

818   **75**, 2998-3014 (2012).

819   36.   Li, K.W*., et al.* Identifying true protein complex constituents in interaction

820   proteomics: the example of the DMXL2 protein complex. *Proteomics* **12**,

821   2428-2432 (2012).

822   37.   Ule, J. & Darnell, R.B. RNA binding proteins and the regulation of neuronal

823   synaptic plasticity. *Curr Opin Neurobiol* **16**, 102-110 (2006).

38. Anderson, P. & Ivanov, P. tRNA fragments in human health and disease.

FEBS Lett **588**, 4297-4304 (2014).

39. Belzil, V.V., Gendron, T.F. & Petrucelli, L. RNA-mediated toxicity in

neurodegenerative disease. *Mol Cell Neurosci* **56**, 406-419 (2013).

40. Bentmann, E., Haass, C. & Dormann, D. Stress granules in

neurodegeneration--lessons learnt from TAR DNA binding protein of 43

kDa and fused in sarcoma. *FEBS J* **280**, 4348-4370 (2013).

41. Ling, S.C., Polymenidou, M. & Cleveland, D.W. Converging mechanisms in

ALS and FTD: disrupted RNA and protein homeostasis. *Neuron* **79**, 416-

438 (2013).

42. Halliday, G.*, et al.* Mechanisms of disease in frontotemporal lobar

degeneration: gain of function versus loss of function effects. *Acta*

Neuropathol **124**, 373-382 (2012).

43. Busch, A. & Hertel, K.J. Evolution of SR protein and hnRNP splicing

regulatory factors. *Wiley Interdiscip Rev RNA* **3**, 1-12 (2012).

44. Moulton, V.R., Gillooly, A.R. & Tsokos, G.C. Ubiquitination regulates

expression of the serine/arginine-rich splicing factor 1 (SRSF1) in normal

and systemic lupus erythematosus (SLE) T cells. *J Biol Chem* **289**, 4126-

4134 (2014).

45. Breig, O. & Baklouti, F. Proteasome-mediated proteolysis of SRSF5

splicing factor intriguingly co-occurs with SRSF5 mRNA upregulation

during late erythroid differentiation. *PLoS One* **8**, e59137 (2013).

46. Graveley, B.R. Sorting out the complexity of SR protein functions. *RNA* **6**,

1197-1211 (2000).

47. Fu, X.D. The superfamily of arginine/serine-rich splicing factors. *RNA* **1**, 663-680 (1995).

48. Gao, L., Wang, J., Wang, Y. & Andreadis, A. SR protein 9G8 modulates splicing of tau exon 10 via its proximal downstream intron, a clustering region for frontotemporal dementia mutations. *Mol Cell Neurosci* **34**, 48-58 (2007).

49. Ding, S*., et al.* Regulation of alternative splicing of tau exon 10 by 9G8 and Dyrk1A. *Neurobiol Aging* **33**, 1389-1399 (2012).

50. Goedert, M., Spillantini, M.G., Jakes, R., Rutherford, D. & Crowther, R.A. Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron* **3**, 519-526 (1989).

51. Goedert, M. & Jakes, R. Expression of separate isoforms of human tau protein: correlation with the tau pattern in brain and effects on tubulin polymerization. *EMBO J* **9**, 4225-4230 (1990).

52. Yin, X*., et al.* Dyrk1A overexpression leads to increase of 3R-tau expression and cognitive deficits in Ts65Dn Down syndrome mice. *Sci Rep* **7**, 619 (2017).

53. Magdalinou, N.K*., et al.* Identification of candidate cerebrospinal fluid biomarkers in parkinsonism using quantitative proteomics. *Parkinsonism Relat Disord* **37**, 65-71 (2017).

54. Luk, C*., et al.* Development and assessment of sensitive immuno-PCR assays for the quantification of cerebrospinal fluid three- and four-repeat tau isoforms in tauopathies. *J Neurochem* **123**, 396-405 (2012).

55.   Chen, S., Townsend, K., Goldberg, T.E., Davies, P. & Conejero-Goldberg, C.

MAPT isoforms: differential transcriptional profiles related to 3R and 4R

splice variants. *J Alzheimers Dis* **22**, 1313-1329 (2010).

56.   Sennvik, K.*, et al.* Tau-4R suppresses proliferation and promotes neuronal

differentiation in the hippocampus of tau knockin/knockout mice. *FASEB*

J **21**, 2149-2161 (2007).

57.   Winborn, B.J.*, et al.* The deubiquitinating enzyme ataxin-3, a

polyglutamine disease protein, edits Lys63 linkages in mixed linkage

ubiquitin chains. *J Biol Chem* **283**, 26436-26443 (2008).

58.   Singh, N.N. & Singh, R.N. Alternative splicing in spinal muscular atrophy

underscores the role of an intron definition model. *RNA Biol* **8**, 600-606

(2011).

59.   Orozco, D. & Edbauer, D. FUS-mediated alternative splicing in the nervous

system: consequences for ALS and FTLD. *J Mol Med (Berl)* **91**, 1343-1354

(2013).

60.   Mills, J.D. & Janitz, M. Alternative splicing of mRNA in the molecular

pathology of neurodegenerative diseases. *Neurobiol Aging* **33**, 1012

e1011-1024 (2012).

61.   Ranum, L.P. & Cooper, T.A. RNA-mediated neuromuscular disorders. *Annu*

Rev Neurosci **29**, 259-277 (2006).

62.   Sun, J., Xu, H., Negi, S., Subramony, S.H. & Hebert, M.D. Differential effects

of polyglutamine proteins on nuclear organization and artificial reporter

splicing. *J Neurosci Res* **85**, 2306-2317 (2007).

63.   Silva-Fernandes, A.*, et al.* Motor uncoordination and neuropathology in a

transgenic mouse model of Machado-Joseph disease lacking intranuclear

inclusions and ataxin-3 cleavage products. *Neurobiol Dis* **40**, 163-176

(2010).

64.   Nicklas, W.*, et al.* Recommendations for the health monitoring of rodent

and rabbit colonies in breeding and experimental units. *Lab Anim* **36**, 20-

42 (2002).

65.   Colledge, M.*, et al.* Targeting of PKA to glutamate receptors through a

MAGUK-AKAP complex. *Neuron* **27**, 107-119 (2000).

66.   Shilov, I.V.*, et al.* The Paragon Algorithm, a next generation search engine

that uses sequence temperature values and feature probabilities to

identify peptides from tandem mass spectra. *Mol Cell Proteomics* **6**, 1638-

1655 (2007).

67.   Fredriksson, S.*, et al.* Protein detection using proximity-dependent DNA

ligation assays. *Nat Biotechnol* **20**, 473-477 (2002).

68.   Glaser, E.M. & Van der Loos, H. Analysis of thick brain sections by

obverse-reverse computer microscopy: application of a new, high clarity

Golgi-Nissl stain. *J Neurosci Methods* **4**, 117-125 (1981).

69.   Pfaffl, M.W. A new mathematical model for relative quantification in real-

time RT-PCR. *Nucleic Acids Res* **29**, e45 (2001).

FIGURE LEGENDS

Figure 1. Experimental design – purification of polyubiquitylated proteins

using TUBEs. **(A-I)** RA-treated SH-SY5Y cells (with silenced ATXN3 or not)

were lysed and protein extracts were incubated with TUBEs. **(A-II)** TUBEs-

captured proteins were recovered with Laemmli buffer and analyzed by

comassie blue staining of the polyubiquityated proteins purified using TUBEs

run in a 1-D SDS-PAGE gel or western-blot with anti-ubiquitin FK2 antibody.

(A-III) Polyubiquitylated proteins were trypsin-digested and identified by LC-

MS/MS. **(B)** After acquisition, data were processed and integrated in

functional networks.

Table 1. RNA metabolism related proteins with altered ubiquitylation in cells

lacking ATXN3.

Figure 2. Efficacy of RNA processing assessed by splicing reporter

minigenes in ATXN3[shRNA] cells. **(A-E)** Semi-quantitative analysis of minigene'

alternative splicing showed a decreased efficiency of splicing in **(A-D)**

ATXN3[shRNA] cells and **(E)** ATXN3_C14A cells. Schemes for the splicing

products are indicated on the left right. Exons are represented as colored

boxes and introns by red lines. The AdML minigene contains one intron,

giving rise to two bands: the upper band corresponds to the unspliced

transcript, the lower band to the spliced product. The pyPY minigene contains

two alternative splice sites originating three bands: an upper band

corresponding to the unspliced transcript, a middle band corresponding to the

splicing of the weak py tract and a lower band corresponding to the splicing

product of the strong PY tract. The pyPY minigene contains two alternative 3'

splice sites associated with polypirimidine (Py) tracts with different strengths.

The weak Py tract (py) is represented by thin red lines and the strong Py tract

(PY) by a thick red line. The α-globin minigene contains two introns and a set

of G triplets in intron 2 that promote the recognition of the 5' splice site leading

to skipping of exon 2. **(F)** Experimental design – microarray analysis of

alternative splicing using the Affymetrix Human Transcriptome Array 2.0 ST to

assess perturbation of global splicing patterns in ATXN3[shRNA] cells. **(G)**

Distribution of the differentially regulated alternative splicing events in

ATXN3[shRNA] cells. n≥3 independent biological replicates in all experiments.

p<0,05 was taken as the cut-off. T-test was used for comparisons. Asterisks

represent mean ± standard deviation. *p<0,05; **p<0,01; ***p<0,001.

Figure 3. ATXN3[shRNA] cells showed decreased SRSF7 protein levels. **(A-B)**

Western-blot analysis after capture of polyubiquitylated proteins confirmed the

decrease in polyubiquitylated forms of SRSF7 in ATXN3[shRNA] cells. **(A-D)**

Downregulation of both **(A-B)** total and **(C-D)** nuclear SRSF7 protein levels in

ATXN3[shRNA] cells. **(E-F)** Protein levels of SRSF7 are increased in ATXN3[shRNA]

cells upon proteasome inhibition (MG132 treatment). **(G)** Western-blot

analysis after capture of polyubiquitylated proteins with K48-linked

polyubiquitin chains showed decrease levels in K48-polyubiquitylated forms of

SRSF7 in ATXN3[shRNA] cells. Relative band density for each protein was

analyzed. The results were normalized for Histone H3. **(H)** Relative amounts

of SRSF7 in SCR[shRNA] cells and ATXN3[shRNA] cells at various cycloheximide

treatment times. Human ATXN3 co- interacts both **(I)** *in vitro* and **(J)** *in situ*

with SRSF7. Scale bar corresponds to 40 μm. n≥3 independent biological

replicates in all experiments. T-test or ANOVA were used for comparisons.

Asterisks represent mean ± standard deviation. *p<0.05; **p<0.01, ***p<0.01.

Figure 4. Disruption of 4R/3R-Tau ratio in ATXN3[shRNA] cells. **(A)** qRT-PCR

analysis showed that silencing of ATXN3 lead to decreased transcription

levels of 4R-Tau, **(B)** no alterations in the mRNA levels of the 3R-Tau isoform,

(C) a significant decrease of the 4R/3R-Tau ratio and **(D)** decreased levels of

total Tau, both in ATXN3[shRNA] cells and in **(E-H)** cerebellar primary neurons.

(I) ATXN3_C14A cells present decreased mRNA levels of 4R-Tau isoform, **(J)**

no alterations in 3R-Tau expression, **(K)** decreased 4R/3R-Tau ratio and **(L)**

decreased levels of total Tau. **(M)** SH-SY5Y ATXN3[shRNA] cultures presented a

decreased expression of βIII-tubulin as assessed by immunostaining. **(N)** The

average length of the neuritis was reduced in ATXN3[shRNA] cells as compared

with the SCR[shRNA] controls. 4R/3R-Tau ratio was obtained by dividing 4R and

3R-Tau mRNA levels. n≥3 independent biological replicates in all

experiments. T-test was used for comparisons. Asterisks represent mean $\pm$

standard deviation. ***p<0.001.

Figure 5. PolyQ expansion in ATXN3 affects Tau splicing. **(A)** Semi-

quantitative analysis of minigene' alternative splicing showed a decreased

efficiency of splicing in ATXN3_83Q cells. **(B)** Expression of ATXN3_83Q led

to decreased mRNA levels of 4R-Tau isoform, **(C)** no alterations in 3R-Tau

expression, **(D)** decreased 4R/3R-Tau ratio, and **(E)** decreased levels of total

Tau. **(F)** Tg mice presented a decrease in the mRNA levels of 4R-Tau, **(G)** no

alterations in expression of 3R-Tau, **(H)** a decreased 4R/3R-Tau ratio, and **(I)**

a decrease in the levels of total Tau comparing with Wt controls. **(J-L)** Tg

mice showed decreased Sfrs7 protein levels in different disease-affected

brain areas. **(M)** Decreased SRSF7 protein levels in SCA3 patient's brains.

(N-P) Decreased total dendritic length and dendritic tree complexity of

brainstem neurons in the Tg mice.

mRNA levels were normalized to the *HMBS* or *Hprt* genes. Relative band

density for each protein was analyzed. The results were normalized for actin

levels. n≥3 independent biological replicates in all experiments. T-test or

ANOVA were used for comparisons. Asterisks represent mean ± standard

deviation. *p<0.05; **p<0.01; ***p<0.001.

1002

Table S1. Differences in polyubiquitylated proteins identified by TUBEs-MS in

SH-SY5Y cells lacking ATXN3. List of identified proteins with altered levels of

ubiquitylation in RA-treated ATXN3[shRNA] cells as compared with the SCR[shRNA]

controls (p<0.05). These proteins were present in at least 3 independent

experiments. The values are the average of "unused" values given by the

Proteinpilot algorithm. The absent "unused" values indicate the (near)

complete absence of the polyubiquitylated protein. In red and green are

proteins with increased and decreased polyubiquitylated levels, respectively.

$ indicates protein splicing factors.

1012

Figure EV1. No changes in the mRNA levels of SRSF7 in ATXN3[shRNA] cells

and in the CMVMJD135 mice. **(A-B)** Percentage of ATXN3 silencing using

different shRNA clones as compared to the scrambled sequence and empty

vector expression. The different generated cell lines showed a similar

phenotype. **(C)** No significant differences were observed in the mRNA levels

of SRSF7 between the ATXN3[shRNA] cells and the SCR[shRNA] controls. **(D)**

Knockdown of Atxn3 in cerebellar primary neurons. **(E)** PLA negative control

(no antibodies) **(F)** No significant differences were observed in the mRNA

levels of Srsf7 in the Tg mice as compared with the Wt-littermates controls.

mRNA levels were normalized to the *HMBS* or *Hprt* genes. Relative band

density for each protein was analyzed. The results were normalized for actin

levels. n≥3 independent biological replicates in all experiments. T-test was

used for comparisons. Asterisks represent mean $\pm$ standard deviation.

*p<0.05.

1027

1028   **Table EV1.** Differentially regulated alternative splicing events in ATXN3[shRNA]

1029   cells. List of genes with altered splicing in RA-treated ATXN3[shRNA] cells as

1030   compared with the SCR[shRNA] controls.

1031

1032   **Table EV2.** KEGG pathway analysis of genes with altered splicing in

1033   ATXN3[shRNA] cells. The genes identified on the KEGG pathway analysis

1034   presented at least one differentially regulated exon/splicing pattern in

1035   ATXN3[shRNA] cells.

1036

1037   **Table EV3.** Changes in alternative types events in ATXN3[shRNA] cells.

1038

Figure 1

A

I. DIFFERENTIATION AND LYSIS

Retinoic acid (RA) 0.1 µM

SCR^shRNA

ATXN3^shRNA

DUB

II. PULLDOWN (PolyUb proteins)

Elution

SDS-PAGE

**Comassie blue staining
Western-blot (anti-Ub)**

III. LC-MS/MS & DATA ANALYSIS

3-6 Biological replicates

Trypsin digestion

LC_MS/MS

Data acquisition

ProteinPlot™ search engine

**Data integration and
Analysis of the ubiquitome**

B

FUNCTIONAL NETWORKS

- **RNA post-transcriptional modification** 14%
- **Molecular transport, RNA trafficking** 11%
- **Cell death and survival** 9%
- **Organ morphology** 9%
- **DNA replication, recombination and repair, Cell death and survival** 8%
- **Cell cycle, Cell death and survival** 8%
- **Cellular compromise, Cell death and survival, Neurological disease** 8%
- **Cellular growth and proliferation, Gene expression, Cell cycle** 8%
- **Gene expression, DNA replication, recombination and repair** 18%
- **Cellular development, Cell death and survival, Cell cycle** 7%

Table 1

RNA METABOLISM PROTEINS WITH ALTERED UBIQUITYLATION IN ATXN3^{shRNA} CELLS

UPREGULATED	p value
RBM10 (RNA-binding protein 10)	4,87E-19
SNRPB2 (U2 small nuclear ribonucleoprotein B)	3,26E-11
SFRS2 (Splicing factor arginine/serine-rich 2)	5,36E-11
RBM8A (RNA-binding protein 8A)	1,56E-06
PRPF8 (Pre-mRNA-processing-splicing factor 8)	1,00E-03
SFRS5 (Serine/arginine-rich splicing factor 5)	5,00E-03
SNRPN (Small nuclear ribonucleoprotein-associated protein N)	5,00E-03
SRSF9 (Serine/arginine-rich splicing factor 9)	7,00E-03
RBM12B (RNA-binding protein 12B)	2,00E-02
AQR (Intron-binding protein aquarius)	2,00E-02
PRPF40A (Pre-mRNA-processing factor 40 homolog A)	2,00E-02
HNRNPK (Heterogeneous nuclear ribonucleoprotein K)	3,00E-02

DOWNREGULATED	p value
DDX41 (DEAD-box protein abstrakt variant)	6,22E-06
DDX56 (ATP-dependent RNA helicase)	2,38E-05
LUC7L (Putative RNA-binding protein Luc-like 1)	5,00E-04
SFRS7 (Serine/arginine-rich splicing factor)	8,00E-04
DDX10 (Probable ATP-dependent RNA helicase)	1,00E-02
HNRNPA1P10 (Heterogeneous nuclear ribonucleoprotein N)	2,00E-02
FUS (RNA-binding protein)	3,00E-02
RBMX (Heterogeneous nuclear ribonucleoprotein G)	4,00E-02
SRRT (Serrate RNA effector molecule)	4,00E-02

Figure 2

A

bp
pyPY
400
250
150

AdML
200
150

α-globin
600
200

SCR^shRNA   ATXN3^shRNA

B py/PY ratio

Spliced/Unspliced

***

SCR^shRNA
ATXN3^shRNA

C AdML

Spliced/unspliced

***

D α-globin

Spliced/unspliced

*

E DN ATXN3 py/PY ratio

Spliced/Unspliced

**

Empty vector
ATXN3_C14A

F

Retinoic acid (RA) 0.1 μM

SCR^shRNA

ATXN3^shRNA

3 biological replicates
Affymetrix HTA 2.0

Data analysis

7450 differentially
expressed genes
(fold change ≥1.5 and
p-value ≤ 0.5)

1993 genes with
differentially regulated
exon usage events

G

TYPES OF SPLICING AFFECTED

Complex
20%

Altered terminal
exon
16%

Intron retention
15%

Altered donor
splice site
3%

Altered acceptor
splice site
3%

Mutually
exclusive exons
1%

Exon cassette
42%

Figure 3

Figure 4

Figure 5

Figure S1

A

B

C *SRSF7* mRNA

D Cerebellar Primary neurons

E PLA negative control

F Mouse *Srsf7* mRNA