Arl13b regulates Sonic Hedgehog signaling from outside primary cilia

Eduardo D. Gigante[a,b], Megan R. Taylor[c], Anna A. Ivanova[d], Richard A. Kahn[d], Tamara Caspary[b]

[a]Neuroscience Graduate Program, [b]Department of Human Genetics, [c]Emory College of Arts and Sciences, [d]Department of Biochemistry, Emory University School of Medicine, Atlanta, GA 30322, USA

Author e-mails: Eduardo.gigante@emory.edu, meg.taylor@emory.edu, anna.ivanova@emory.edu, rkahn@emory.edu

ORCID ID: Gigante: 0000-0002-1486-5377, Kahn: 0000-0002-0259-0601, Caspary: 0000-0002-6579-7589

⌘**Correspondence**: Tamara Caspary

tcaspar@emory.edu

Abstract

ARL13B is a regulatory GTPase highly enriched in cilia. Complete loss of *Arl13b*

disrupts cilia architecture, protein trafficking and Sonic hedgehog signaling. To

determine whether ARL13B is required within cilia, we knocked in a cilia-excluded

variant of ARL13B (V358A) and showed it retains all known biochemical function. We

found that ARL13B^{V358A} protein was expressed but could not be detected in cilia, even

when retrograde ciliary transport was blocked. We showed *Arl13b$^{V358A/V358A}$* mice are

viable and fertile with normal Shh signal transduction. However, in contrast to wild type

cilia, *Arl13b$^{V358A/V358A}$* cells displayed short cilia and lacked ciliary ARL3 and Inpp5e.

These data indicate that ARL13B's role within cilia can be uncoupled from its function

outside of cilia. Furthermore, these data imply that the cilia defects upon complete

absence of ARL13B do not underlie the alterations in Shh transduction, which is

unexpected given the requirement of cilia for Shh transduction.

Keywords:

ARL13B; Sonic Hedgehog signaling; primary cilia; mouse development

Introduction

The Hedgehog (Hh) signaling pathway is essential for embryogenesis in a wide

variety of organisms. Initially discovered in *Drosophila* where there is a single Hh ligand,

the core components of the Hh pathway are conserved in vertebrates (Nusslein-Volhard

and Wieschaus 1980). These include the vertebrate Hh receptor Patched1 (Ptch1), the

obligate transducer of the pathway Smoothened (Smo), as well as the Gli transcription

factors (Ci in *Drosophila*) that act as both activators (GliA) and repressors (GliR) to

control target gene transcription (Briscoe and Therond 2013). There are three Hh

ligands in vertebrates, Sonic (Shh), Indian (Ihh) and Desert (Dhh), that regulate a

multitude of developmental processes including formation of the limbs and digits, the

bones of the skull and face, and the patterning of the neural tube (Placzek and Briscoe

2018). Diminished Hh signaling during embryogenesis results in birth defects whereas

increased Hh signaling leads to tumors, highlighting the importance of the pathway and

its regulation (Raleigh and Reiter 2019).

Given the deep homology between invertebrate and vertebrate Hh signaling, the

discovery that primary cilia are required for vertebrate, but not invertebrate, Hh signaling

was unexpected (Huangfu et al. 2003; Huangfu and Anderson 2006). Vertebrate Hh

components dynamically traffic within cilia in response to ligand (Corbit et al. 2005;

Haycraft et al. 2005; Rohatgi et al. 2007). In the absence of ligand, Ptch1 is enriched on

the ciliary membrane and Smo is barely detectable (Corbit et al. 2005; Rohatgi et al.

2007). Furthermore, full length Gli proteins traffic to the ciliary tip and back to the

cytoplasm before being cleaved to their repressor form, which actively shuts down Hh

target gene transcription in the nucleus (Kim et al. 2009; Liu et al. 2005; Wen et al.

2010; Humke et al. 2010). In contrast, upon ligand stimulation (Shh, Ihh or Dhh), the

Ptch1 receptor binds the ligand and shuttles out of cilia (Rohatgi et al. 2007).

Subsequently, Smo is enriched in the cilium and is subsequently activated (Kong et al.

2019; Corbit et al. 2005). Activated Smo promotes the processing of full-length Gli

transcription factors into GliA, which turns on target genes (Aza-Blanc et al. 2000; Ruiz i

Altaba 1998). Ablation of cilia results in an absence of GliA and GliR production,

rendering the pathway inert and leading to an absence of transcriptional response

(Huangfu and Anderson 2005). The dynamic ciliary movement of Shh components

appears to be critical to pathway function, as alterations to cilia disrupt pathway output

(Caspary et al. 2007; Huangfu and Anderson 2006; Liem et al. 2012; Liem et al. 2009;

Goetz et al. 2009; Murdoch and Copp 2010; Tuz et al. 2014; Cortellino et al. 2009;

Houde et al. 2006; Liu et al. 2005; Tran et al. 2008; Taylor et al. 2015).

Given that the fundamental logic of the pathway is conserved from flies through

vertebrates, and that flies transduce Hh signals without relying on cilia, both

evolutionary and mechanistic questions are raised as to how vertebrate cells co-opted

the cilium for Hh signal transduction. One distinction lies in the fact that vertebrates use

Hh signaling over a longer distance than flies, leading to the proposal that the cilium is a

critical part of a mechanism underlying long range signaling (Bangs et al. 2015). For

example, in neural patterning Shh is initially expressed in the notochord and is secreted

to specify fates more than 20 cells away (Chiang et al. 1996; Briscoe et al. 2001;

Roelink et al. 1994). At the evolutionary level, comparisons among organisms with cilia

and or Hh have provided some clues. The round worm *C. elegans* have cilia yet do not

possess Hh signaling as they don't have most of the genes encoding the core

components of Hh signal transduction (Consortium 1998; Roy 2012). Curiously, a few

components of Hh signaling such as fused and costal 2 are in the *C. elegans* genome

where they are functionally important for ciliogenesis (Ingham et al. 2011). Additionally,

C. elegans retained a Ptch1 homolog important for development and pattern formation,

but no Hh or Smo (Zugasti et al. 2005; Kuwabara et al. 2000). In contrast, planaria

flatworms possess both cilia and Hh signaling but the cilia are not required to transduce

Hh signaling (Rink et al. 2009). These data suggest that the mechanistic link of cilia and

Hh is limited to vertebrates and originated in the last common ancestor of vertebrates,

the urochordates.

ARL13B is a member of the ARF family of regulatory GTPases and is highly

enriched on the ciliary membrane (Caspary et al. 2007). In mice, a null mutation of

Arl13b leads to short cilia and to alterations in Shh signal transduction (Caspary et al.

2007; Larkins et al. 2011). ARL13 is ancient, predicted to be present in the last common

eukaryotic ancestor. ARL13 appears to have been lost during evolution in organisms

that lack cilia and duplicated to ARL13A and ARL13B in the urochordates, thus ARL13B

is proposed to hold important clues in deciphering the links between cilia and vertebrate

Hh signaling (Schlacht et al. 2013; Li et al. 2004; Kahn et al. 2008; East et al. 2012;

Logsdon and Kahn 2004).

ARF regulatory GTPases, like ARL13B, are best known to play roles in

membrane trafficking (D'Souza-Schorey and Chavrier 2006). As is true for a large

number of regulatory GTPases, ARL13B is functionally diverse (Sztul et al. 2019). It

regulates endocytic traffic (Barral et al. 2012) as well as the phospholipid composition of

the ciliary membrane through recruitment of the lipid phosphatase Inpp5e to the ciliary

membrane (Humbert et al. 2012). ARL13B also has a conserved role as a guanine

nucleotide exchange factor (GEF) for ARL3, another ciliary ARF-like (ARL) protein

(Gotthardt et al. 2015; Zhang et al. 2016; Hanke-Gogokhia et al. 2016; Ivanova et al.

2017). ARL13B regulates intraflagellar transport (IFT), the process that builds and

maintains cilia (Cevik et al. 2010; Li et al. 2010; Nozaki et al. 2017). It is known to

interact with several proteins associated with cilia, including the exocyst, tubulin and

UNC119 (Seixas et al. 2016; Zhang et al. 2016; Larkins et al. 2011; Revenkova et al.

2018). Critical to this work, loss of ARL13B disrupts Shh signal transduction in at least

two distinct ways: Smo enrichment in cilia occurs even in the absence of ligand and Gli

activator production is diminished, although Gli repressor is made normally (Caspary et

al. 2007; Larkins et al. 2011).

Due to the high enrichment of ARL13B on the ciliary membrane, ARL13B is

assumed to function in its diverse roles from within the cilium. However, ARL13B is

present in early endosomes and circular dorsal ruffles on the cell surface (Barral et al.

2012; Casalou et al. 2014). We previously showed that a V358A variant of ARL13B

does not localize to cilia as it disrupts a known VxPx cilia localization sequence

(Higginbotham et al. 2012; Mariani et al. 2016). Exogenous overexpression of a

ARL13B^{V358A} construct in *Arl13b* null cells does not rescue ARL13B-dependent

phenotypes such as cilia length as well as interneuron migration and connectivity,

consistent with ciliary ARL13B mediating these processes (Higginbotham et al. 2012;

Mariani et al. 2016). In contrast, we found that overexpressed ARL13B^{V358A} does rescue

the Shh-dependent ciliary enrichment of Smo in mouse embryonic fibroblasts, arguing

that ARL13B may function outside the cilium to regulate Smo traffic (Mariani et al.

2016). Together, these results raise the question of where ARL13B functions.

To define where ARL13B functions in relation to cilia, we wanted an *in vivo*

model so generated mice carrying the ARL13B^{V358A} point mutation using CRISPR/Cas9.

Here we demonstrate that ARL13B^{V358A} protein was undetectable in *Arl13b$^{V358A/V358A}$*

cilia in both neural tube and mouse embryonic fibroblast cilia, even after blocking

retrograde ciliary traffic. We report that *Arl13b$^{V358A/V358A}$* mice were viable, fertile, and

transduced Shh signal normally. We found ARL3 and Inpp5e did not localize to the

short cilia present in/on *Arl13b$^{V358A/V358A}$* cells. These data indicate that ARL13B's roles

within and outside cilia can be uncoupled; ARL13B's role in regulating cilia length is

from within cilia, whereas its control of Shh signaling appears to be from outside the

cilium. Thus, these data imply that the cilia defects seen in the complete absence of

ARL13B do not underlie the alterations in Shh transduction, which is unexpected given

the requirement of cilia for Shh signal transduction.

Results

ARL13B^{V358A} *displays normal GEF activity*

We previously showed that mouse ARL13B^{V358A} protein retained normal intrinsic

and GAP-stimulated GTP hydrolysis activities by analyzing GST-ARL13B^{V358A} purified

from human embryonic kidney (HEK) cells (Mariani et al. 2016). In order to test

ARL13B^{V358A} GEF activity for ARL3, we used the same GST-ARL13B^{V358A} protein

preparation and measured the rates of spontaneous or GEF-stimulated GDP

dissociation from ARL3 in the presence and absence of ARL13B or ARL13B^{V358A}. ARL3

spontaneously releases pre-bound GDP quite slowly under the conditions in the assay,

though release is linear with time, and by extrapolation requires just under 30 min for

50% of pre-bound [^{3}H]GDP to dissociate (**Figure 1**). In marked contrast, addition of

ARL13B under the same conditions caused the release of 50% of the GDP within ~10

seconds and close to 100% in one minute. We detected no differences between the wild

type and mutant ARL13B^{V358A} proteins in this assay, consistent with this point mutation

having no effect on ARL13B GEF function (**Figure 1**). This result is consistent with GEF

activity being conserved within the protein's GTPase domain while the V358A mutation

is located in the C-terminal domain (Gotthardt et al. 2015). These data indicate that

ARL13B^{V358A} retains all known ARL13B biochemical activities, suggesting that the

V358A mutation only disrupts ARL13B localization.

CRISPR engineered* Arl13b$^{V358A/V358A}$ *mice express cilia-excluded ARL13B protein.

137   To determine the consequences of ARL13B^{V358A} expression *in vivo*, we used

CRISPR/Cas9 editing to change residue 358 from valine to alanine (**Figure 2A**). We

performed sequencing of the region that flanked exon 8, using primers outside the

region of the donor oligonucleotide. We confirmed the heterozygous T to C base pair

change for a valine to alanine change at residue 358 (**Figure 2B**). We backcrossed

heterozygous F1 progeny to FVB/NJ for three generations before analysis to minimize

any off target confounds (see details in Methods).

144   To determine whether ARL13B^{V358A} was detectable in neural tube cilia, we

performed immunofluorescence using antibodies directed against ARL13B and the cilia

marker IFT88. At embryonic day (E)10.5, we saw IFT88 staining, indicating the

presence of cilia. In neural tube cilia of *Arl13b*^{+/+} and *Arl13b*^{V358A/+} embryos, we

observed equivalent ARL13B staining but could not see ARL13B signal in neural tube

cilia of *Arl13b*^{V358A/V358A} embryos, indicating the ARL13B^{V358A} protein is undetectable in

cilia *in vivo* (**Figure 2C**).

One possible explanation for the absence of ciliary ARL13B in the

Arl13b^{V358A/V358A} embryos is that the ARL13B^{V358A} protein is not expressed or is

unstable. To determine whether ARL13B protein expression was affected by the V358A

mutation, we performed Western blots on E10.5 whole embryo lysates (**Figure 2D**). We

found a ~7% (±20%) reduction of ARL13B levels in *Arl13b*^{V358A/+} embryos and a ~37%

(±6%) decrease of ARL13B levels in *Arl13b*^{V358A/V358A} embryos, compared to WT

(**Figure 2E**). This decrease may reflect a change in ARL13B^{V358A} stability compared to

ciliary ARL13B or could signify ARL13B^{V358A} protein having a distinct half-life when

localizing to different cellular compartments. Regardless, these data indicate that

ARL13B^{V358A} is expressed in *Arl13b*^{V358A/V358A} embryos.

***ARL13B*^{V358A} protein is undetectable in cilia in mouse embryonic fibroblasts.**

To more closely investigate whether any ARL13B^{V358A} protein could localize to

cilia, we generated immortalized mouse embryonic fibroblasts (MEFs). We performed

double labeling to first identify cilia using acetylated α-tubulin or IFT88 antibodies and

subsequently determined whether ARL13B is present in cilia. We used five distinct

antibodies against ARL13B: 4 antibodies against amino acids 208-427 of the mouse

ARL13B protein (1 mouse monoclonal (NeuroMab), 3 rabbit polyclonal antibodies

(Caspary et al. 2007)), as well as 1 antibody against the full-length human ARL13B

presence of cilia. In neural tube cilia of *Arl13b*$^{+/+}$ and *Arl13b*$^{V358A/+}$ embryos, we

observed equivalent ARL13B staining but could not see ARL13B signal in neural tube

cilia of *Arl13b*$^{V358A/V358A}$ embryos, indicating the ARL13B^{V358A} protein is undetectable in

cilia *in vivo* (**Figure 2C**).

One possible explanation for the absence of ciliary ARL13B in the

Arl13b$^{V358A/V358A}$ embryos is that the ARL13B^{V358A} protein is not expressed or is

unstable. To determine whether ARL13B protein expression was affected by the V358A

mutation, we performed Western blots on E10.5 whole embryo lysates (**Figure 2D**). We

found a ~7% (±20%) reduction of ARL13B levels in *Arl13b*$^{V358A/+}$ embryos and a ~37%

(±6%) decrease of ARL13B levels in *Arl13b*$^{V358A/V358A}$ embryos, compared to WT

(**Figure 2E**). This decrease may reflect a change in ARL13B^{V358A} stability compared to

ciliary ARL13B or could signify ARL13B^{V358A} protein having a distinct half-life when

localizing to different cellular compartments. Regardless, these data indicate that

ARL13B^{V358A} is expressed in *Arl13b*$^{V358A/V358A}$ embryos.

***ARL13B*^{V358A} protein is undetectable in cilia in mouse embryonic fibroblasts.**

To more closely investigate whether any ARL13B^{V358A} protein could localize to

cilia, we generated immortalized mouse embryonic fibroblasts (MEFs). We performed

double labeling to first identify cilia using acetylated α-tubulin or IFT88 antibodies and

subsequently determined whether ARL13B is present in cilia. We used five distinct

antibodies against ARL13B: 4 antibodies against amino acids 208-427 of the mouse

ARL13B protein (1 mouse monoclonal (NeuroMab), 3 rabbit polyclonal antibodies

(Caspary et al. 2007)), as well as 1 antibody against the full-length human ARL13B

protein (rabbit polyclonal (Protein Tech)). By eye, we observed strong ciliary staining of

ARL13B with each of the antibodies in *Arl13b$^{+/+}$* and *Arl13b$^{V358A/+}$* cells, but we were

unable to identify any evidence of ARL13B staining above background within cilia in

Arl13b$^{V358A/V358A}$ cells with any of these five antibodies (**Figure 3A**). Taken together,

these data support our conclusion that ARL13B^{V358A} is not detectable in cilia using the

currently available, validated ARL13B antibodies.

While we observed no evidence of detectable ciliary ARL13B in *Arl13b$^{V358A/V358A}$*

MEFs, it is possible that a small amount of ARL13B^{V358A} is constantly trafficking in and

out of cilia, at steady-state levels that remain below the limits of detection. To address

this possibility, we blocked retrograde ciliary transport, reasoning that any ARL13B

undergoing trafficking in and out of cilia would accumulate. We treated cells with

ciliobrevin-D, which blocks the retrograde motor protein dynein (Firestone et al. 2012).

As a positive control, we examined IFT88, a ciliary protein known to accumulate at

ciliary tips upon blocking of retrograde transport. We found IFT88 enrichment at ciliary

tips in *Arl13b$^{+/+}$*, *Arl13b$^{V358A/+}$* and *Arl13b$^{V358A/V358A}$* cells upon ciliobrevin-D treatment

relative to the respective DMSO-treated controls with no difference in IFT88 staining

among the 3 genotypes (**Figure 3B**). To determine whether ARL13B accumulated in

cilia of ciliobrevin-D-treated MEFs, we examined cells co-stained for acetylated α-

tubulin and ARL13B. In *Arl13b$^{+/+}$* and *Arl13b$^{V358A/+}$* cells, we saw that about 90% of

acetylated α-tubulin-positive cilia also stained for ARL13B in DMSO-treated control or

ciliobrevin-D treated MEFs (**Figure 3C**). In *Arl13b$^{V358A/V358A}$* MEFs, we saw no ciliary

ARL13B staining in DMSO-treated control or ciliobrevin-D-treated cells (**Figure 3C**).

Thus, even when retrograde traffic out of cilia is blocked, we were unable to detect

ARL13B protein in cilia in $Arl13b^{V358A/V358A}$ MEFs.

We re-examined these data and repeated our analyses using over-exposed

images. After defining the region of interest using the acetylated α-tubulin staining, we

subsequently overexposed the ARL13B channel five-fold relative to the images in

Figure 3 and acquired measurements at the cilium and the cell body (**Supplemental**

Figure 1A). As a control, we used $Arl13b^{hnn/hnn}$ MEFs, which are devoid of any ARL13B,

and obtained a ratio of 1.0, with or without ciliobrevin-D treatment. We found the same

ratio when we analyzed $Arl13b^{V358A/V358A}$ MEFs consistent with ARL13B being absent

from the cilia (**Supplemental Figure 1B, C**). We observed a few instances of ARL13B

appearing to co-localize with acetylated α-tubulin, but these were rare (<5%, 2/49) and

occurred in <u>both</u> $Arl13b^{V358A/V358A}$ and $Arl13b^{hnn/hnn}$ (null) cells indicating this is the

background staining of the primary antibody (**Supplemental Figure 1D**). We extended

our analysis of overexposed images in $Arl13b^{V358A/V358A}$ and $Arl13b^{hnn/hnn}$ neural tube

sections four-fold relative to the images in Figure 2; we identified no overlap of

overexposed ARL13B with cilia marker IFT88 (**Supplemental Figure 1E**). While it is

formally possible that an extremely small amount of ARL13B^{V358A} remains in cilia, we

are not able to find evidence of it; thus, we designate ARL13B^{V358A} protein as cilia-

excluded ARL13B.

Arl13b$^{V358A/V358A}$ mice are viable and fertile.

In order to determine the phenotypic consequence of the $Arl13b^{V358A}$ allele, we

intercrossed $Arl13b^{V358A/+}$ mice. We observed progeny in Mendelian proportions with an

average of 7.3 pups per litter, consistent with the reported FVB/NJ litter size of 7-9

(Murray et al. 2010) (**Table 1**). To test whether homozygous mice breed normally, we

crossed *Arl13b^{V358A/V358A}* males to heterozygous or homozygous females. We found the

Arl13b^{V358A} allele segregated in Mendelian proportions and the litter sizes were normal

indicating that *Arl13b^{V358A}* does not impact viability, fertility or fecundity.

ARL13B^{V358A} permits normal embryonic development and Shh signaling.

Loss of *Arl13b* leads to morphologically abnormal embryos, lethality and neural

patterning defects (Caspary et al. 2007). As ARL13B^{V358A} overcame the embryonic

lethality, we examined overall embryo morphology at E9.5, E10.5 and E12.5. We found

the overall morphology of *Arl13b^{V358A/V358A}* embryos resembled those of *Arl13b^{+/+}* and

Arl13b^{V358A/+} embryos indicating that ARL13B^{V358A} did not lead to gross morphological

defects. (**Figure 4A-I**).

At E9.5, Shh is normally expressed in the notochord as well as the floor plate of

the ventral neural tube. Moving dorsally, the adjacent domains express Nkx2.2 and

subsequently Olig2. In the *Arl13b^{hnn/hnn}* (null) neural tube, the Shh-positive columnar

cells of the floor plate are absent resulting in both ventral and dorsal expansion of

intermediate Shh-dependent cell fates such as Olig2 (Caspary et al. 2007). To

determine whether ARL13B^{V358A} disrupts Shh signaling, we examined neural tube

patterning in *Arl13b^{+/+}*, *Arl13b^{V358A/+}*, and *Arl13b^{V358A/V358A}* embryos at E9.5, E10.5, and

E12.5. At E9.5 we observed no differences in Shh, Nkx2.2 or Olig2 among *Arl13b^{+/+}*,

Arl13b^{V358A/+}, and *Arl13b^{V358A/V358A}* embryos (**Figure 4A-C**). Cell fates in the neural tube

are specified by both the concentration and duration of Shh signaling so we examined

neural patterning at subsequent stages (Dessaud et al. 2007). By E10.5, some Olig2

precursors have differentiated to HB9+ motor neurons and all Shh-responsive cells

express Nkx6.1. We found that Olig2, HB9 and Nkx6.1 positive cells are normally

distributed in all 3 genotypes at E10.5 and E12.5 (**Figure 4D-I**). These data indicate that

ARL13B^{V358A} mediates normal Shh signaling.

ARL13B regulates ciliary enrichment of Shh components from outside cilia.

In *Arl13b$^{hnn/hnn}$* cells, Smo is enriched in cilia in a Shh-independent manner,

which may be due to defective trafficking of Smo, as many ARL family members

regulate protein trafficking (Lim et al. 2011; Larkins et al. 2011). To assess Smo

enrichment in *Arl13b$^{V358A/V358A}$* embryos, we stained for the cilia marker acetylated α-

tubulin and Smo in E10.5 embryos. Smo appeared enriched normally in ventral floor

plate cilia and the dorsal boundary of Smo ciliary enrichment in ventral neural

progenitors was identical in *Arl13b$^{+/+}$*, *Arl13b$^{V358A/+}$*, and *Arl13b$^{V358A/V358A}$* samples

indicating ARL13B^{V358A} mediates normal ciliary Smo enrichment (**Figure 5A**).

To examine Smo traffic in response to Shh stimulation, we treated MEFs with

0.5% FBS or Shh-conditioned media for 24 hours and stained for Smo. As expected in

control *Arl13b$^{hnn/hnn}$* cells, we saw ciliary Smo in unstimulated MEFs which increased

upon stimulation with Shh-conditioned media (Larkins et al. 2011) (**Figure 5B**). In

Arl13b$^{+/+}$, *Arl13b$^{V358A/+}$*, and *Arl13b$^{V358A/V358A}$* MEF cilia, we saw ciliary enrichment of

Smo upon Shh stimulation over their respective unstimulated controls (**Figure 5B**, left).

We found no difference in Smo enrichment among these cell lines (**Figure 5B**, right).

Thus, despite ARL13B being critical for Shh-dependent Smo ciliary enrichment,

ARL13B^{V358A} regulates Smo localization normally, arguing this function of ARL13B can

occur when the protein is not in cilia.

In addition to aberrations in Smo trafficking, loss of *Arl13b* leads to changes in

the cilia localization of other Shh components in MEFs (Larkins et al. 2011). To

determine whether these components are disrupted by ARL13B^{V358A}, we examined Gli2,

Gli3, Sufu and Ptch1 in MEFs. We observed no differences in distribution of Gli2 or

Ptch1 among any of the examined genotypes (**Figure 5C, F**). In contrast, as we

previously reported, we observed more Gli3 at the ciliary tip of *Arl13b$^{hnn/hnn}$* cells and

increased Sufu in *Arl13b$^{hnn/hnn}$* cilia compared to wild type controls (Larkins et al.

2011)(**Figure 5D, E**). Thus, Shh components are normally localized in *Arl13b$^{V358A/V358A}$*

MEFs consistent with the normal Shh signal transduction observed in *Arl13b$^{V358A/V358A}$*

embryos (**Figure 4**).

ARL13B regulates ciliary enrichment of ARL3 and Inpp5e from within cilia.

We next examined the role of ARL13B^{V358A} in the localization of other ciliary

proteins. ARL13B is the GEF for ARL3, and we showed that ARL13B^{V358A} GEF activity

for ARL3 is normal (**Figure 1**). ARL13B is also essential for cilia localization of Inpp5e,

as the proteins are in a common complex (Humbert et al. 2012). To determine whether

the ciliary localization of either ARL3 or Inpp5e was affected by ARL13B^{V358A}, we

performed immunofluorescence on MEFs. Ciliary ARL3 staining appeared the same in

Arl13b$^{+/+}$ and *Arl13b$^{V358A/+}$* MEFs, however ARL3 was not detectable in *Arl13b$^{V358A/V358A}$*

or *Arl13b$^{hnn/hnn}$* cilia (**Figure 5G**). Inpp5e was normally detectable in *Arl13b$^{+/+}$* and

Arl13b$^{V358A/+}$ MEF cilia, but not in *Arl13b$^{V358A/V358A}$* and *Arl13b$^{hnn/hnn}$* MEF cilia (**Figure**

5H). These results indicate that ARL13B is required in cilia for the proper localization of

ARL3 and Inpp5e to the cilium.

Ciliary ARL13B is required for normal cell ciliation and ciliary length

286   Loss of *Arl13b* leads to defects in the percentage of ciliated MEFs and in cilia

length (Caspary et al. 2007; Larkins et al. 2011). To test whether ARL13B^{V358A} impacted

cell ciliation, we examined immortalized MEFs and counted the number of ciliated cells

24 hours after induction of ciliation by growth in low serum (0.5% FBS). We found 73%

(±11%) of *Arl13b$^{+/+}$* and 75%, (±9.1%) of *Arl13b$^{V358A/+}$* MEFs are ciliated, consistent with

published results. In contrast, we found 45% (±8.7%) of *Arl13b$^{V358A/V358A}$* cells and 53%

(±6.4%) of *Arl13b$^{hnn/hnn}$* cells had cilia (**Figure 6A**). Thus, *Arl13b$^{V358A/V358A}$* cells exhibit a

similar deficit in percentage of ciliated MEFs as complete loss of ARL13B function.

294   *Arl13b$^{hnn/hnn}$* cilia are about half as long as wild type in both embryos and MEFs

(Caspary et al. 2007; Larkins et al. 2011). To determine whether cilia length was

affected by ARL13B^{V358A}, we stained MEFs with acetylated α−tubulin and measured the

length of the axoneme. We found the average cilia length in *Arl13b$^{+/+}$* and Arl13b$^{V358A/+}$

MEFs was similar, 2.7 ± 0.8 μm and 2.6 ± 0.9 μm, respectively. However, we found

Arl13b$^{V358A/V358A}$ MEF cilia were shorter, 1.9 ± 0.7 μm, similar to *Arl13b$^{hnn/hnn}$* MEF cilia

which were 2.15 ± 0.8 μm (**Figure 6B**). These results indicate that ARL13B^{V358A}

phenocopies complete loss of ARL13B for ciliation and cilia length, indicating these

ARL13B functions require ARL13B in cilia. Furthermore, these data show that ARL13B

function within cilia is distinct from ARL13B function outside of cilia in a subset of

activities, and that the cilia defects and Shh defects in the complete absence of ARL13B

can be uncoupled.

Discussion

Here we show that ARL13B's role(s) in cell ciliation and cilia length, along with

the ciliary enrichment of a subset of proteins, can be uncoupled from ARL13B's function

in regulating Shh signal transduction (**Figure 7**). Furthermore, we show this functional

distinction correlates with ARL13B spatial localization to the cilium. By generating a

mouse expressing a cilia-excluded variant of ARL13B from the endogenous locus, we

showed ARL13B^{V358A} protein is not detectable in cilia of embryonic neural tube or

MEFs, even upon retrograde ciliary traffic blockade. Furthermore, we detected 30% less

ARL13B protein overall in *Arl13b$^{V358A/V358A}$* embryos compared to control embryos.

While this reduction is statistically significant, it is unclear whether it is biologically

significant given that *Arl13b$^{hnn/hnn}$* null mutations are recessive and no *Arl13b$^{hnn/+}$*

heterozygous phenotype is reported (Caspary et al. 2007; Larkins et al. 2011).

In contrast to the E13.5 lethality of *Arl13b$^{hnn/hnn}$* embryos, we found

Arl13b$^{V358A/V358A}$ mice were viable and fertile with correct patterning of the neural tube,

indicating normal Shh signal transduction. These results are consistent with our data

showing that ARL13B^{V358A} protein retains all known biochemical function including GEF

activity (Mariani et al. 2016; Ivanova et al. 2017). However, we observed several

Arl13b$^{hnn/hnn}$ phenotypes in *Arl13b$^{V358A/V358A}$* cells, including loss of ARL3 and Inpp5e

ciliary enrichment, along with low percentage of ciliated cells and shorter average cilia

length. Taken together, our data show that despite the normal high ciliary enrichment of

wild type ARL13B and the requirement of cilia for Shh signaling, cilia-excluded

ARL13B^{V358A} is sufficient for Shh signal transduction.

We regard ARL13B^{V358A} as cilia-excluded based on several lines of evidence.

We did not detect ciliary ARL13B^{V358A} *in vivo* or *in vitro* using any of five validated

ARL13B antibodies. These antibodies are against two antigens, the full-length protein or

residues 208-427, and were independently generated so are likely to recognize a

number of epitopes. Given that 4 of 5 antibodies are polyclonal and ARL13B^{V358A}

displays normal GTP binding, intrinsic and GAP-stimulated GTP hydrolysis, and ARL3

GEF activity, it is likely that the ARL13B^{V358A} protein retains the wild type structure

enabling antibody recognition. Indeed, we observed little or no loss in sensitivity in

detecting protein in the immunoblot assays. Furthermore, we could not forcibly enrich

ciliary ARL13B^{V358A} through retrograde transport blockade with ciliobrevin-D nor could

we detect any ciliary enrichment of ARL13B^{V358A} upon overexposure of the relevant

fluorescent channel. The fact that we observed ciliary phenotypes, namely short cilia

and abnormal ciliary ARL3 and Inpp5e localization in ARL13B^{V358A} cells, argue that

ARL13B normally performs these functions from within cilia (Caspary et al. 2007;

Larkins et al. 2011; Zhang et al. 2016; Humbert et al. 2012; Nozaki et al. 2017). While

we cannot exclude the possibility that sub-detectable levels of ARL13B^{V358A} are present

and functional in cilia, we note such a level would need to be sufficient for Shh signaling

yet insufficient for ARL3 or Inpp5e ciliary localization along with proper regulation of cilia

length.

Our data support ARL13B regulating different biological processes from its

distinct subcellular localizations. ARL13B^{V358A} disrupts cilia localization of Inpp5e and

ARL3, but not Shh components. It is surprising that ARL13B^{V358A} is sufficient to regulate

Shh signaling because cilia are required for Shh signal transduction and because

ARL13B is highly enriched in cilia (Caspary et al. 2007; Huangfu et al. 2003). These

observations suggested that the ciliary defects observed in *Arl13b$^{hnn/hnn}$* mutants caused

the Shh defects. However, our data with ARL13B^{V358A} indicate that ARL13B regulates

cilia length and Shh signaling through distinct localization and effectors.

Based on the normal cilia trafficking of Shh components in *ARL13B$^{V358A/V358A}$*

mutants and the abnormal cilia trafficking of Shh components in the complete absence

of ARL13B (Larkins et al. 2011), ARL13B likely regulates Shh signaling from the cell

body by controlling Shh component traffic to and/or from the cilium. Both Smo and

ARL13B traffic is linked to endosomes providing one possible non-ciliary organelle

(Barral et al. 2012; Wang et al. 2009; Milenkovic et al. 2009). *Arl13b$^{hnn/hnn}$* cells display

constitutive Smo ciliary enrichment along with little enrichment of Gli proteins at the

ciliary tip (Larkins et al. 2011). Our data do not distinguish between ARL13B playing

direct roles in traffic of multiple Shh components or whether the normal Smo ciliary

enrichment with ARL13B^{V358A} subsequently causes normal cilia traffic of downstream

components. ARL13B regulates a step downstream of activated Smo that controls

transcription factor Gli activation, but not repression (Caspary et al. 2007; Bay et al.

2018). We argue from our results that this step is also intact in the presence of

ARL13B^{V358A}.

The fact that ARL13B^{V358A} can mediate normal ciliary Smo enrichment is

especially interesting given that ARL3 and Inpp5e localization to cilia is compromised by

this mutation. This suggests that ARL13B controls cilia enrichment via multiple

localizations and effectors. This is consistent with our understanding of ARF family

members, as they are known to perform multiple tasks from different sites within a cell

(Sztul et al. 2019). We speculate that ARL3 residence in cilia may require that ARL3 be

in (or at least cycle through) its activated, GTP-bound conformation as ARL13B^{V358A}

retains its ARL3 GEF activity, but not its cilia localization. ARL13B is in a common

protein complex with Inpp5e so absence of ARL13B from cilia may disrupt formation or

maintenance of the complex there (Humbert et al. 2012; Nozaki et al. 2017). Inpp5e

regulates Shh signaling through regulation of the phosphoinositol composition of the

ciliary membrane. Inpp5e loss results in increased ciliary PIP$_2$ and enrichment of Shh

repressors in cilia thus resulting in lowered Shh response (Garcia-Gonzalo et al. 2015;

Chavez et al. 2015; Dyson et al. 2017). It is not clear why the absence of ciliary Inpp5e

in ARL13B^{V358A} cells does not lead to aberrant Shh signal transduction.

In *Arl13b$^{hnn/hnn}$* embryos and MEFs, cilia are shorter than normal and have a

microtubule defect where the B-tubule of the microtubule outer doublet is open

(Caspary et al. 2007; Larkins et al. 2011). Consistent with this, we observed shorter cilia

in *Arl13b$^{V358A/V358A}$* MEFs. Similarly, we observed a comparable reduction in the percent

of cilia in *Arl13b$^{hnn/hnn}$* and *Arl13b$^{V358A/V358A}$* MEFs compared to wild type MEFs. While

loss of ARL13B results in short cilia, increased expression of ARL13B promotes cilia

elongation so it is not yet clear what ARL13B's role is in controlling cilia length (Lu et al.

2015; Hori et al. 2008; Caspary et al. 2007; Larkins et al. 2011). Additionally, it is

unclear whether the mechanism from within cilia through which ARL13B controls length

or percent of ciliated cells is the same, or distinct from, the mechanism through which

ARL13B regulates ARL3 or Inpp5e residence in cilia.

Perhaps the most intriguing implication of our data pertains to understanding the

evolution of cilia and Hh signaling. ARL13 functions in cilia formation and maintenance

in *Chlamydomonas* and *C. elegans*, neither of which have Hh signaling, consistent with

the ancient role of Arl13b controlling ciliation and cilia length (Cevik et al. 2010; Cevik et

al. 2013; Stolc et al. 2005; Miertzschke et al. 2014). Our data support ARL13B retaining

ARL13's ancient role in ciliogenesis. However, our data show that ARL13B does not

work from within the cilium to regulate Shh component ciliary traffic or Shh signal

transduction. As an ARL protein involved in membrane traffic, we speculate that

ARL13B may have linked Shh component trafficking to the cilium, albeit from outside

the cilium. That the ARL13 duplication in the urochordates coincides with when ARL13B

adopted new cellular roles in Hh signaling that linked the pathway to cilia fits well with

this model.

Methods

Protein purification and ARL3 GEF assay

Plasmids directing the expression of mouse GST-ARL13B or GST-ARL13B[V358A]

proteins were transiently transfected into HEK cells and the recombinant proteins were

later purified by affinity chromatography using glutathione-Sepharose, as described

previously (Cavenagh et al. 1994; Ivanova et al. 2017). Human ARL3 (98.4% identical

to mouse ARL3) was expressed in BL21 bacteria and purified as previously described

(Van Valkenburgh et al. 2001). The ARL3 GEF assay was performed as previously

described (Ivanova et al. 2017). Briefly, ARL3 (10μM) was pre-incubated with [^{3}H]GDP

(1 μM; PerkinElmer Life Sciences, specific activity ~3,000 cpm/pmol) for 1 h at 30°C in

25 mM HEPES, pH 7.4, 100 mM NaCl, 10 mM $MgCl_2$. In contrast, pre-loading of

ARL13B (10μM) was achieved by incubation in 100 μM GTPγS for 1 min at room

temperature, due to its much more rapid exchange kinetics. The GEF assay was

initiated upon addition of ARL3 (final = 1μM), ARL13B (final = 0 or 1μM), 10 μM GTPγS,

and 100 μM GDP (to prevent re-binding of any released [^{3}H]GDP), in a final volume of

100 μL. The intrinsic rate of GDP dissociation from ARL3 was determined in parallel as

that released in the absence of any added ARL13B. The reactions were stopped at

different times (0–15 min) by dilution of 10μL of the reaction mixture into 2 ml of ice-cold

buffer (20 mM Tris, pH 7.5, 100 mM NaCl, 10 mM $MgCl_2$, 1 mM dithiothreitol). The

amount of ARL3-bound [^{3}H]GDP was determined by rapid filtration through BA85

nitrocellulose filters (0.45 μm, 25 mm, Whatman), with 3 x 2 mL washes, and quantified

using liquid scintillation counting. Time points are routinely collected in at least duplicate

and each experiment reported was repeated at least twice, yielding very similar results.

Data were analyzed in GraphPad Prism 7 software.

Mouse allele generation and identification

All mice were cared for in accordance with NIH guidelines and Emory's

Institutional Animal Care and Use Committee (IACUC). Lines used were *Arl13b[V358A]*

(*Arl13b[em1Tc]*) [MGI: 6256969], *Arl13b[hnn]* [MGI: 3578151] and FVB/NJ (Jackson

Laboratory). To generate the V358A point mutation in the mouse, a CRISPR gRNA

(CCAGTCAATACAGACGAGTCTA) targeting exon 8 of the *Arl13b* locus along with a

donor oligo (5'-

CCTATATTCTTCTAGAAACAGTAAGAAGAAACCAAGAACTACGAATGAAAAGGA

GTCATCGGG<u>C</u>AGAACCAGT<u>G</u>AATACAGACGAGTCTACTCCAAAGAGTCCCACGCC

TCCCCAAC-3'; underlined bases are engineered) were designed to generate a T-to-C

change creating the V358A point mutation and C-to-G change creating a TspRI

restriction site that could be used for genotyping (Millipore Sigma). The gRNA (50

ng/μl), oligo donor (50 ng/μl) and CRISPR protein (100 ng/μL) were injected by the

Emory Transgenic and Gene Targeting core into one-cell C57Bl/6J zygotes. Zygotes

were cultured to 2-cell before being transferred to pseudopregnant females. Genomic

tail DNA from resulting offspring was amplified using PCR primers (5'-

GAAGCAGGCATGGTGGTAAT-3' & 5'-TGAACCGCTAATGGGAAATC-3') located

upstream and downstream of the donor oligo breakpoints. The products were

sequenced (5'-GAAGCAGGCATGGTGGTAAT-3') and 2 animals were identified

heterozygous for the desired change and with no additional editing. One line (#173) was

bred to FVB/NJ for three generations prior to performing any phenotypic analysis. Males

from at least two distinct meiotic recombination opportunities were used in each

generation that to minimize potential confounds associated with off-target

CRISPR/Cas9 editing.

Genotyping was performed on DNA extracted from ear punch or yolk sac via

PCR using the following primers: Fwd: 5'- AAGAATGAAAAGGAGTCAGCG -3', Rev: 5'-

TGAACCGCTAATGGGAAACT -3'; a SNP was engineered in the forward primer that, in

combination with the V358A mutation, created a Cac8I restriction site. Thus, the PCR

product was digested with Cac8I enzyme, run out on a 4% agarose gel and the relevant

bands were detected: undigested wild type (~192bp) and digested mutant bands

(~171bp).

Phenotypic analysis of embryos

Timed matings of heterozygous intercrosses were performed to generate embryos of

the indicated stage, with somite-matched pairs examined at each stage when

appropriate. Embryos were dissected in cold phosphate-buffered saline and processed

for immunofluorescence staining? as previously described (Horner and Caspary 2011).

Mouse embryonic fibroblasts

Mouse embryonic fibroblasts (MEFs) were isolated and immortalized as previously

described (Mariani et al. 2016). MEFs were maintained in DMEM with 10% fetal bovine

serum (FBS) and 1% penicillin/streptomycin at 37°C in 5% CO_2. For experimental use,

Arl13b$^{+/+}$, *Arl13b$^{V358A/+}$*, and *Arl13b$^{V358A/V358A}$* and *Arl13b$^{hnn/hnn}$* MEFs were grown on

coverslips at a density of 0.5 x 10^6 cells/mL and treated for 24 h with 0.5% FBS Shh

conditioned media or 0.5% FBS media to induce ciliogenesis (Larkins et al. 2011).

Antibodies

Primary antibodies used were: mouse anti-Shh (5E1), mouse anti-Nkx2.2 (74.5A5),

mouse anti-HB9 (81.5C10) (1:5 Developmental Studies Hybridoma Bank); rabbit anti-

Olig2 (1:300 Millipore AB9610); mouse anti-acetylated α-tubulin (1:2500 Millipore

Sigma; T6793), mouse anti-Arl13b (1:1000 NeuroMab N295B/66), rabbit anti-Arl13b

(1:1000, Protein Tech 17711-1-AP); rabbit anti-Arl13b sera from 3 distinct rabbits (503,

504 and 505) (Caspary Lab (Caspary et al. 2007)), rabbit anti-Smoothened (1:1000,

kindly provided by K. Anderson); rabbit anti-IFT88 (1:1000, kindly provided by B.

Yoder); rabbit anti-ARL3 (1:1000, (Cavenagh et al. 1994)); rabbit anti-Inpp5e (1:150,

Protein Tech 17797-1-AP); guinea pig anti-Gli2 (1:200, kindly provided by J.

Eggenschwiler); rabbit anti-Ptch1 (1:150, kindly provided by R. Rohatgi), goat anti-Sufu

(1:100, SC10933, Santa Cruz); goat anti-Gli3 (1:200, R&D AF3690) Alexa Fluor 488

and Alexa Fluor 568 (1:300, ThermoFisher); and Hoechst nuclear stain (1:3000).

Image Quantification

Fluorescent intensities were measured in ImageJ software. Cilia were first identified by

positive acetylated α-tubulin and an outline was hand drawn around the length of the

cilium. The channel was switched to the protein of interest and a measurement of the

average fluorescent intensity was acquired. The same outline was then used to acquire

a background reading of the cell-body that most closely matched the background at the

cilium. In all cases immunofluorescent averages of proteins of interest in cilia were

normalized to cell-body intensities (Supplemental Figure 1A). For samples with

antibodies targeting Gli2, Gli3, and Sufu the ciliary tip was isolated and measured. The

cilia tip was identified by weak acetylated α-tubulin staining or the cilium base was

identified by the presence of the acetylated α-tubulin positive fibrils in the cell body that

cluster at the base of the cilium (Larkins et al. 2011). For samples with antibodies

targeting Smo, Ptch1, ARL3, and Inpp5e the entire cilium was measured. We plotted

the ratio of fluorescence intensity of the protein of interest to the cell body background.

The number of cilia examined per genotype and per condition are listed below their

respective violin plot. Data were analyzed by one-way ANOVA, except for Smo intensity

where a two-way ANOVA was used with Sidak's multiple comparison. In the event of a

significant ANOVA, Tukey's multiple comparisons were employed to determine

significance. Data were analyzed in GraphPad Prism 7 software.

In a similar experiment, retrograde IFT was inhibited by the addition of 30 μM

ciliobrevin-D (Millipore Sigma; 250401) or 0.1% DMSO control in low-serum media for

60 min after ciliation was induced by serum starvation for 24 h prior with 0.5% FBS

media. Cilia were identified by staining with antibodies directed against acetylated α-

tubulin, IFT88, and ARL13B. Fluorescent intensities were measured in ImageJ software.

Again, cilia were first identified by positive acetylated α-tubulin and an outline was hand

drawn. The channel was switched to the protein of interest and a measurement of the

average fluorescent intensity was acquired. The same outline was then used to acquire

a background reading of the cell-body. To measure IFT88, the cilia tip was identified by

weak acetylated α-tubulin staining or the cilium base was identified by the presence of

acetylated α-tubulin fibrils in the cell body that cluster at the base of the cilium (Larkins

et al. 2011). Data were analyzed by two-way ANOVA with Sidak's multiple comparison.

Western Blots

Western blotting was performed as previously described (Mariani et al. 2016), with

antibody against ARL13B (1:1000 Neuromab N295B/66). Lysates were prepared with

RIPA buffer and SigmaFast protease inhibitors (Nachtergaele et al. 2013). Values

presented as volume intensity measured by chemiluminescence detected on ChemiDoc

Touch Imaging System were normalized to total protein as measured on a stain-free

gel. Normalized values were analyzed by one-way ANOVA and Tukey's multiple

comparisons. Data were analyzed in GraphPad Prism 7 software.

Acknowledgements

This work was supported by funding from NIH grants R01NS090029 and R35GM122549 to T.C. and R35GM122568 to R.A.K. E.D.G. was supported by R01NS090029 diversity supplement. Further support came from the Emory Integrated Mouse Transgenic and Gene Targeting Core, which is subsidized by the Emory University School of Medicine and with support by the Georgia Clinical & Translational Science Alliance of the National Institutes of Health under Award Number UL1TR002378 as well as the Emory University Integrated Cellular Imaging Microscopy Core of the Emory Neuroscience NINDS Core Facilities grant, P30NS055077. We are grateful to Alyssa Long and Sarah Suciu for sequencing potential CRISPR founders, and to members of the Caspary lab for discussion and manuscript comments.

Competing Interests

The authors have no competing interests to disclose.

References

Aza-Blanc, P., H. Y. Lin, A. Ruiz i Altaba, and T. B. Kornberg. 2000. 'Expression of the
vertebrate Gli proteins in Drosophila reveals a distribution of activator and repressor
activities', *Development*, 127: 4293-301.
Bangs, F. K., N. Schrode, A. K. Hadjantonakis, and K. V. Anderson. 2015. 'Lineage specificity of
primary cilia in the mouse embryo', *Nat Cell Biol*, 17: 113-22.
Barral, D. C., S. Garg, C. Casalou, G. F. Watts, J. L. Sandoval, J. S. Ramalho, V. W. Hsu, and
M. B. Brenner. 2012. 'Arl13b regulates endocytic recycling traffic', *Proc Natl Acad Sci U*
S A, 109: 21354-9.
Bay, S. N., A. B. Long, and T. Caspary. 2018. 'Disruption of the ciliary GTPase Arl13b
suppresses Sonic hedgehog overactivation and inhibits medulloblastoma formation',
Proc Natl Acad Sci U S A, 115: 1570-75.
Briscoe, J., Y. Chen, T. M. Jessell, and G. Struhl. 2001. 'A hedgehog-insensitive form of
patched provides evidence for direct long-range morphogen activity of sonic hedgehog
in the neural tube', *Mol Cell*, 7: 1279-91.
Briscoe, J., and P. P. Therond. 2013. 'The mechanisms of Hedgehog signalling and its roles in
development and disease', *Nat Rev Mol Cell Biol*, 14: 416-29.
Casalou, C., C. Seixas, A. Portelinha, P. Pintado, M. Barros, J. S. Ramalho, S. S. Lopes, and D.
C. Barral. 2014. 'Arl13b and the non-muscle myosin heavy chain IIA are required for
circular dorsal ruffle formation and cell migration', *J Cell Sci*, 127: 2709-22.
Caspary, T., C. E. Larkins, and K. V. Anderson. 2007. 'The graded response to Sonic Hedgehog
depends on cilia architecture', *Dev Cell*, 12: 767-78.
Cavenagh, M. M., M. Breiner, A. Schurmann, A. G. Rosenwald, T. Terui, C. Zhang, P. A.
Randazzo, M. Adams, H. G. Joost, and R. A. Kahn. 1994. 'ADP-ribosylation factor
(ARF)-like 3, a new member of the ARF family of GTP-binding proteins cloned from
human and rat tissues', *J Biol Chem*, 269: 18937-42.
Cevik, S., Y. Hori, O. I. Kaplan, K. Kida, T. Toivenon, C. Foley-Fisher, D. Cottell, T. Katada, K.
Kontani, and O. E. Blacque. 2010. 'Joubert syndrome Arl13b functions at ciliary
membranes and stabilizes protein transport in Caenorhabditis elegans', *J Cell Biol*, 188:
953-69.
Cevik, S., A. A. Sanders, E. Van Wijk, K. Boldt, L. Clarke, J. van Reeuwijk, Y. Hori, N. Horn, L.
Hetterschijt, A. Wdowicz, A. Mullins, K. Kida, O. I. Kaplan, S. E. van Beersum, K. Man
Wu, S. J. Letteboer, D. A. Mans, T. Katada, K. Kontani, M. Ueffing, R. Roepman, H.
Kremer, and O. E. Blacque. 2013. 'Active transport and diffusion barriers restrict Joubert
Syndrome-associated ARL13B/ARL-13 to an Inv-like ciliary membrane subdomain',
PLoS Genet, 9: e1003977.
Chavez, M., S. Ena, J. Van Sande, A. de Kerchove d'Exaerde, S. Schurmans, and S. N.
Schiffmann. 2015. 'Modulation of Ciliary Phosphoinositide Content Regulates Trafficking
and Sonic Hedgehog Signaling Output', *Dev Cell*, 34: 338-50.
Chiang, C., Y. Litingtung, E. Lee, K. E. Young, J. L. Corden, H. Westphal, and P. A. Beachy.
1996. 'Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene
function', *Nature*, 383: 407-13.
Consortium, C. elegans Sequencing. 1998. 'Genome sequence of the nematode C. elegans: a
platform for investigating biology', *Science*, 282: 2012-8.
Corbit, K. C., P. Aanstad, V. Singla, A. R. Norman, D. Y. Stainier, and J. F. Reiter. 2005.
'Vertebrate Smoothened functions at the primary cilium', *Nature*, 437: 1018-21.
Cortellino, S., C. Wang, B. Wang, M. R. Bassi, E. Caretti, D. Champeval, A. Calmont, M. Jarnik,
J. Burch, K. S. Zaret, L. Larue, and A. Bellacosa. 2009. 'Defective ciliogenesis,
embryonic lethality and severe impairment of the Sonic Hedgehog pathway caused by

590   inactivation of the mouse complex A intraflagellar transport gene Ift122/Wdr10, partially
591   overlapping with the DNA repair gene Med1/Mbd4', *Dev Biol*, 325: 225-37.
592 D'Souza-Schorey, C., and P. Chavrier. 2006. 'ARF proteins: roles in membrane traffic and
593   beyond', *Nat Rev Mol Cell Biol*, 7: 347-58.
594 Dessaud, E., L. L. Yang, K. Hill, B. Cox, F. Ulloa, A. Ribeiro, A. Mynett, B. G. Novitch, and J.
595   Briscoe. 2007. 'Interpretation of the sonic hedgehog morphogen gradient by a temporal
596   adaptation mechanism', *Nature*, 450: 717-20.
597 Dyson, J. M., S. E. Conduit, S. J. Feeney, S. Hakim, T. DiTommaso, A. J. Fulcher, A. Sriratana,
598   G. Ramm, K. A. Horan, R. Gurung, C. Wicking, I. Smyth, and C. A. Mitchell. 2017.
599   'INPP5E regulates phosphoinositide-dependent cilia transition zone function', *J Cell Biol*,
600   216: 247-63.
601 East, M. P., J. B. Bowzard, J. B. Dacks, and R. A. Kahn. 2012. 'ELMO domains, evolutionary
602   and functional characterization of a novel GTPase-activating protein (GAP) domain for
603   Arf protein family GTPases', *J Biol Chem*, 287: 39538-53.
604 Firestone, A. J., J. S. Weinger, M. Maldonado, K. Barlan, L. D. Langston, M. O'Donnell, V. I.
605   Gelfand, T. M. Kapoor, and J. K. Chen. 2012. 'Small-molecule inhibitors of the AAA+
606   ATPase motor cytoplasmic dynein', *Nature*, 484: 125-9.
607 Garcia-Gonzalo, F. R., S. C. Phua, E. C. Roberson, G. Garcia, 3rd, M. Abedin, S. Schurmans,
608   T. Inoue, and J. F. Reiter. 2015. 'Phosphoinositides Regulate Ciliary Protein Trafficking
609   to Modulate Hedgehog Signaling', *Dev Cell*, 34: 400-09.
610 Goetz, S. C., P. J. Ocbina, and K. V. Anderson. 2009. 'The primary cilium as a Hedgehog signal
611   transduction machine', *Methods Cell Biol*, 94: 199-222.
612 Gotthardt, K., M. Lokaj, C. Koerner, N. Falk, A. Giessl, and A. Wittinghofer. 2015. 'A G-protein
613   activation cascade from Arl13B to Arl3 and implications for ciliary targeting of lipidated
614   proteins', *Elife*, 4.
615 Hanke-Gogokhia, C., Z. Wu, C. D. Gerstner, J. M. Frederick, H. Zhang, and W. Baehr. 2016.
616   'Arf-like Protein 3 (ARL3) Regulates Protein Trafficking and Ciliogenesis in Mouse
617   Photoreceptors', *J Biol Chem*, 291: 7142-55.
618 Haycraft, C. J., B. Banizs, Y. Aydin-Son, Q. Zhang, E. J. Michaud, and B. K. Yoder. 2005. 'Gli2
619   and Gli3 localize to cilia and require the intraflagellar transport protein polaris for
620   processing and function', *PLoS Genet*, 1: e53.
621 Higginbotham, H., T. Y. Eom, L. E. Mariani, A. Bachleda, J. Hirt, V. Gukassyan, C. L. Cusack,
622   C. Lai, T. Caspary, and E. S. Anton. 2012. 'Arl13b in primary cilia regulates the migration
623   and placement of interneurons in the developing cerebral cortex', *Dev Cell*, 23: 925-38.
624 Hori, Y., T. Kobayashi, Y. Kikko, K. Kontani, and T. Katada. 2008. 'Domain architecture of the
625   atypical Arf-family GTPase Arl13b involved in cilia formation', *Biochem Biophys Res*
626   *Commun*, 373: 119-24.
627 Horner, V. L., and T. Caspary. 2011. 'Disrupted dorsal neural tube BMP signaling in the cilia
628   mutant Arl13b hnn stems from abnormal Shh signaling', *Dev Biol*, 355: 43-54.
629 Houde, C., R. J. Dickinson, V. M. Houtzager, R. Cullum, R. Montpetit, M. Metzler, E. M.
630   Simpson, S. Roy, M. R. Hayden, P. A. Hoodless, and D. W. Nicholson. 2006. 'Hippi is
631   essential for node cilia assembly and Sonic hedgehog signaling', *Dev Biol*, 300: 523-33.
632 Huangfu, D., and K. V. Anderson. 2005. 'Cilia and Hedgehog responsiveness in the mouse',
633   *Proc Natl Acad Sci U S A*, 102: 11325-30.
634 ———. 2006. 'Signaling from Smo to Ci/Gli: conservation and divergence of Hedgehog
635   pathways from Drosophila to vertebrates', *Development*, 133: 3-14.
636 Huangfu, D., A. Liu, A. S. Rakeman, N. S. Murcia, L. Niswander, and K. V. Anderson. 2003.
637   'Hedgehog signalling in the mouse requires intraflagellar transport proteins', *Nature*, 426:
638   83-7.

Humbert, M. C., K. Weihbrecht, C. C. Searby, Y. Li, R. M. Pope, V. C. Sheffield, and S. Seo.
2012. 'ARL13B, PDE6D, and CEP164 form a functional network for INPP5E ciliary
targeting', *Proc Natl Acad Sci U S A*, 109: 19691-6.
Humke, E. W., K. V. Dorn, L. Milenkovic, M. P. Scott, and R. Rohatgi. 2010. 'The output of
Hedgehog signaling is controlled by the dynamic association between Suppressor of
Fused and the Gli proteins', *Genes Dev*, 24: 670-82.
Ingham, P. W., Y. Nakano, and C. Seger. 2011. 'Mechanisms and functions of Hedgehog
signalling across the metazoa', *Nat Rev Genet*, 12: 393-406.
Ivanova, A. A., T. Caspary, N. T. Seyfried, D. M. Duong, A. B. West, Z. Liu, and R. A. Kahn.
2017. 'Biochemical characterization of purified mammalian ARL13B protein indicates
that it is an atypical GTPase and ARL3 guanine nucleotide exchange factor (GEF)', *J
Biol Chem*, 292: 11091-108.
Kahn, R. A., E. Bruford, H. Inoue, J. M. Logsdon, Jr., Z. Nie, R. T. Premont, P. A. Randazzo, M.
Satake, A. B. Theibert, M. L. Zapp, and D. Cassel. 2008. 'Consensus nomenclature for
the human ArfGAP domain-containing proteins', *J Cell Biol*, 182: 1039-44.
Kim, J., M. Kato, and P. A. Beachy. 2009. 'Gli2 trafficking links Hedgehog-dependent activation
of Smoothened in the primary cilium to transcriptional activation in the nucleus', *Proc
Natl Acad Sci U S A*, 106: 21666-71.
Kong, J. H., C. Siebold, and R. Rohatgi. 2019. 'Biochemical mechanisms of vertebrate
hedgehog signaling', *Development*, 146.
Kuwabara, P. E., M. H. Lee, T. Schedl, and G. S. Jefferis. 2000. 'A C. elegans patched gene,
ptc-1, functions in germ-line cytokinesis', *Genes Dev*, 14: 1933-44.
Larkins, C. E., G. D. Aviles, M. P. East, R. A. Kahn, and T. Caspary. 2011. 'Arl13b regulates
ciliogenesis and the dynamic localization of Shh signaling proteins', *Mol Biol Cell*, 22:
4694-703.
Li, Y., W. G. Kelly, J. M. Logsdon, Jr., A. M. Schurko, B. D. Harfe, K. L. Hill-Harfe, and R. A.
Kahn. 2004. 'Functional genomic analysis of the ADP-ribosylation factor family of
GTPases: phylogeny among diverse eukaryotes and function in C. elegans', *FASEB J*,
18: 1834-50.
Li, Y., Q. Wei, Y. Zhang, K. Ling, and J. Hu. 2010. 'The small GTPases ARL-13 and ARL-3
coordinate intraflagellar transport and ciliogenesis', *J Cell Biol*, 189: 1039-51.
Liem, K. F., Jr., A. Ashe, M. He, P. Satir, J. Moran, D. Beier, C. Wicking, and K. V. Anderson.
2012. 'The IFT-A complex regulates Shh signaling through cilia structure and membrane
protein trafficking', *J Cell Biol*, 197: 789-800.
Liem, K. F., Jr., M. He, P. J. Ocbina, and K. V. Anderson. 2009. 'Mouse Kif7/Costal2 is a cilia-
associated protein that regulates Sonic hedgehog signaling', *Proc Natl Acad Sci U S A*,
106: 13377-82.
Lim, Y. S., C. E. Chua, and B. L. Tang. 2011. 'Rabs and other small GTPases in ciliary
transport', *Biol Cell*, 103: 209-21.
Liu, A., B. Wang, and L. A. Niswander. 2005. 'Mouse intraflagellar transport proteins regulate
both the activator and repressor functions of Gli transcription factors', *Development*, 132:
3103-11.
Logsdon, J. M., Jr., and R. A. Kahn (ed.)^(eds.). 2004. *Arf Family GTPases. In: Kahn RA, editor.*
Lu, H., M. T. Toh, V. Narasimhan, S. K. Thamilselvam, S. P. Choksi, and S. Roy. 2015. 'A
function for the Joubert syndrome protein Arl13b in ciliary membrane extension and
ciliary length regulation', *Dev Biol*, 397: 225-36.
Mariani, L. E., M. F. Bijlsma, A. I. Ivanova, S. K. Suciu, R. A. Kahn, and T. Caspary. 2016.
'Arl13b regulates Shh signaling from both inside and outside the cilium', *Mol Biol Cell*.
Miertzschke, M., C. Koerner, M. Spoerner, and A. Wittinghofer. 2014. 'Structural insights into the
small G-protein Arl13B and implications for Joubert syndrome', *Biochem J*, 457: 301-11.

Milenkovic, L., M. P. Scott, and R. Rohatgi. 2009. 'Lateral transport of Smoothened from the
plasma membrane to the membrane of the cilium', *J Cell Biol*, 187: 365-74.
Murdoch, J. N., and A. J. Copp. 2010. 'The relationship between sonic Hedgehog signaling,
cilia, and neural tube defects', *Birth Defects Res A Clin Mol Teratol*, 88: 633-52.
Murray, S. A., J. L. Morgan, C. Kane, Y. Sharma, C. S. Heffner, J. Lake, and L. R. Donahue.
2010. 'Mouse gestation length is genetically determined', *PLoS One*, 5: e12418.
Nachtergaele, S., D. M. Whalen, L. K. Mydock, Z. Zhao, T. Malinauskas, K. Krishnan, P. W.
Ingham, D. F. Covey, C. Siebold, and R. Rohatgi. 2013. 'Structure and function of the
Smoothened extracellular domain in vertebrate Hedgehog signaling', *Elife*, 2: e01340.
Nozaki, S., Y. Katoh, M. Terada, S. Michisaka, T. Funabashi, S. Takahashi, K. Kontani, and K.
Nakayama. 2017. 'Regulation of ciliary retrograde protein trafficking by the Joubert
syndrome proteins ARL13B and INPP5E', *J Cell Sci*, 130: 563-76.
Nusslein-Volhard, C., and E. Wieschaus. 1980. 'Mutations affecting segment number and
polarity in Drosophila', *Nature*, 287: 795-801.
Placzek, M., and J. Briscoe. 2018. 'Sonic hedgehog in vertebrate neural tube development', *Int
J Dev Biol*, 62: 225-34.
Raleigh, D. R., and J. F. Reiter. 2019. 'Misactivation of Hedgehog signaling causes inherited
and sporadic cancers', *J Clin Invest*, 129: 465-75.
Revenkova, E., Q. Liu, G. L. Gusella, and C. Iomini. 2018. 'The Joubert syndrome protein
ARL13B binds tubulin to maintain uniform distribution of proteins along the ciliary
membrane', *J Cell Sci*, 131.
Rink, J. C., K. A. Gurley, S. A. Elliott, and A. Sanchez Alvarado. 2009. 'Planarian Hh signaling
regulates regeneration polarity and links Hh pathway evolution to cilia', *Science*, 326:
1406-10.
Roelink, H., A. Augsburger, J. Heemskerk, V. Korzh, S. Norlin, A. Ruiz i Altaba, Y. Tanabe, M.
Placzek, T. Edlund, T. M. Jessell, and et al. 1994. 'Floor plate and motor neuron
induction by vhh-1, a vertebrate homolog of hedgehog expressed by the notochord',
Cell, 76: 761-75.
Rohatgi, R., L. Milenkovic, and M. P. Scott. 2007. 'Patched1 regulates hedgehog signaling at
the primary cilium', *Science*, 317: 372-6.
Roy, S. 2012. 'Cilia and Hedgehog: when and how was their marriage solemnized?',
Differentiation, 83: S43-8.
Ruiz i Altaba, A. 1998. 'Combinatorial Gli gene function in floor plate and neuronal inductions by
Sonic hedgehog', *Development*, 125: 2203-12.
Schlacht, A., K. Mowbrey, M. Elias, R. A. Kahn, and J. B. Dacks. 2013. 'Ancient complexity,
opisthokont plasticity, and discovery of the 11th subfamily of Arf GAP proteins', *Traffic*,
14: 636-49.
Seixas, C., S. Y. Choi, N. Polgar, N. L. Umberger, M. P. East, X. Zuo, H. Moreiras, R.
Ghossoub, A. Benmerah, R. A. Kahn, B. Fogelgren, T. Caspary, J. H. Lipschutz, and D.
C. Barral. 2016. 'Arl13b and the exocyst interact synergistically in ciliogenesis', *Mol Biol
Cell*, 27: 308-20.
Stolc, V., M. P. Samanta, W. Tongprasit, and W. F. Marshall. 2005. 'Genome-wide
transcriptional analysis of flagellar regeneration in Chlamydomonas reinhardtii identifies
orthologs of ciliary disease genes', *Proc Natl Acad Sci U S A*, 102: 3703-7.
Sztul, E., P. W. Chen, J. E. Casanova, J. Cherfils, J. B. Dacks, D. G. Lambright, F. S. Lee, P. A.
Randazzo, L. C. Santy, A. Schurmann, I. Wilhelmi, M. E. Yohe, and R. A. Kahn. 2019.
'ARF GTPases and their GEFs and GAPs: concepts and challenges', *Mol Biol Cell*, 30:
1249-71.
Taylor, S. P., T. J. Dantas, I. Duran, S. Wu, R. S. Lachman, Consortium University of
Washington Center for Mendelian Genomics, S. F. Nelson, D. H. Cohn, R. B. Vallee,

and D. Krakow. 2015. 'Mutations in DYNC2LI1 disrupt cilia function and cause short rib
polydactyly syndrome', *Nat Commun*, 6: 7092.
Tran, P. V., C. J. Haycraft, T. Y. Besschetnova, A. Turbe-Doan, R. W. Stottmann, B. J. Herron,
A. L. Chesebro, H. Qiu, P. J. Scherz, J. V. Shah, B. K. Yoder, and D. R. Beier. 2008.
'THM1 negatively modulates mouse sonic hedgehog signal transduction and affects
retrograde intraflagellar transport in cilia', *Nat Genet*, 40: 403-10.
Tuz, K., R. Bachmann-Gagescu, D. R. O'Day, K. Hua, C. R. Isabella, I. G. Phelps, A. E.
Stolarski, B. J. O'Roak, J. C. Dempsey, C. Lourenco, A. Alswaid, C. G. Bonnemann, L.
Medne, S. Nampoothiri, Z. Stark, R. J. Leventer, M. Topcu, A. Cansu, S. Jagadeesh, S.
Done, G. E. Ishak, I. A. Glass, J. Shendure, S. C. Neuhauss, C. R. Haldeman-Englert,
D. Doherty, and R. J. Ferland. 2014. 'Mutations in CSPP1 cause primary cilia
abnormalities and Joubert syndrome with or without Jeune asphyxiating thoracic
dystrophy', *Am J Hum Genet*, 94: 62-72.
Van Valkenburgh, H., J. F. Shern, J. D. Sharer, X. Zhu, and R. A. Kahn. 2001. 'ADP-ribosylation
factors (ARFs) and ARF-like 1 (ARL1) have both specific and shared effectors:
characterizing ARL1-binding proteins', *J Biol Chem*, 276: 22826-37.
Wang, Y., Z. Zhou, C. T. Walsh, and A. P. McMahon. 2009. 'Selective translocation of
intracellular Smoothened to the primary cilium in response to Hedgehog pathway
modulation', *Proc Natl Acad Sci U S A*, 106: 2623-8.
Wen, X., C. K. Lai, M. Evangelista, J. A. Hongo, F. J. de Sauvage, and S. J. Scales. 2010.
'Kinetics of hedgehog-dependent full-length Gli3 accumulation in primary cilia and
subsequent degradation', *Mol Cell Biol*, 30: 1910-22.
Zhang, Q., Y. Li, Y. Zhang, V. E. Torres, P. C. Harris, K. Ling, and J. Hu. 2016. 'GTP-binding of
ARL-3 is activated by ARL-13 as a GEF and stabilized by UNC-119', *Sci Rep*, 6: 24534.
Zugasti, O., J. Rajan, and P. E. Kuwabara. 2005. 'The function and expansion of the Patched-
and Hedgehog-related homologs in C. elegans', *Genome Res*, 15: 1402-10.
765

Figure 1.

767 **ARL3 GEF activity is retained in the ARL13B^{V358A} mutant.** Time course of the

768 release of pre-bound [^{3}H]GDP from purified, recombinant ARL3 in the absence (ARL3)

769 or presence of mouse wild type ARL13B or ARL13B^{V358A} (mARL13B) are shown. See

770 Methods for details. Error bars ± standard deviation.

Figure 2.

Figure 2.

Generation of the *Arl13b*^{V358A/V358A} mouse.

(A) Schematic of *Arl13b* gene and donor oligo (orange bar) at exon 8 used to generate

the V358A causing point mutation. Arrows indicate primers used for allele validation. (B)

Arl13b DNA and relevant amino acid sequence with the RVEP sequence in the orange

box and the T-to-C mutation highlighted in pink. (C) Confocal images of cilia marker

IFT88 (green) and ARL13B (magenta) staining in neural tube of E10.5 somite-matched

embryos. ARL13B-positive cilia are visible in *Arl13b*$^{+/+}$ and *Arl13b*$^{V358A/+}$, but not in

Arl1b$^{V358A/V358A}$ embryos. At least 5 embryos per genotype across five litters were

examined. Scale bars are 50 μm. (D) ARL13B Western blot of E10.5 whole embryo

lysates, wild type (+/+), *Arl13b*$^{V358A/+}$ (*A/+*) and *Arl13b*$^{V358A/V358A}$ (*A/A*) and (E)

quantification presented as average intensity normalized to total protein ± standard

deviation. Representative blot of whole embryo lysates (n = 3 embryos per genotype

with technical duplicate of each). * $p < 0.05$, one-way ANOVA and Tukey's multiple

comparison.

Figure 3.

A

	PT	505	504	503	NM
Arl13b[+/+]	204 (209)	223 (229)	147 (152)	223 (237)	55 (55)
Arl13b[V358A/+]	154 (171)	147 (171)	148 (159)	134 (144)	90 (90)
Arl13b[V358A/V358A]	0 (154)	0 (164)	0 (98)	0 (93)	0 (25)

Arl13b[+/+]

Arl13b[V358A/+]

Arl13b[V358A/V358A]

ARL13B α-Actub   ARL13B IFT88

B

IFT88

Relative Fluorescence (au)

***

19	22	23	18	19	19	n
+/+	A/+	A/A	+/+	A/+	A/A	

DMSO   Ciliobrevin

IFT88   α-Actub

C

	DMSO	30μM C-D
Arl13b[+/+]	122 (129)	103 (112)
Arl13b[V358A/+]	112 (126)	119 (129)
Arl13b[V358A/V358A]	0 (131)	0 (118)

Arl13b[+/+]

Arl13b[V358A/+]

Arl13b[V358A/V358A]

ARL13B α-Actub

Figure 3.

ARL13B^{V358A} is undetectable in cilia and cannot be enriched by inhibition of

retrograde transport.

(A) Antibodies against ciliary markers acetylated α-tubulin or IFT88 (magenta) and

ARL13B (cyan) in *Arl13b$^{V358A/V358A}$* MEFs. Representative images show staining for five

indicated ARL13B antibodies: (PT) polyclonal rabbit antibody against full-length human

ARL13B from ProteinTech, (503, 504, 505) polyclonal rabbit sera from three distinct

rabbits raised against C-terminus of mouse ARL13B (amino acids 208-427) (Caspary et

al. 2007), and (NM) monoclonal mouse antibody against C terminus of mouse ARL13B

from NeuroMab. *Arl13b$^{+/+}$* and *Arl13b$^{V358A/+}$* show ciliary ARL13B staining. Table lists

ARL13B-positive cilia and the total number of cilia identified by acetylated α-tubulin or

IFT88 antibody in parentheses. Cilia appear shorter in *Arl13b$^{V358A/V358A}$* cells (see Fig.

6). (B) IFT88 (green) is enriched in the tips of cilia in *Arl13b$^{+/+}$* (+/+), *Arl13b$^{V358A/+}$* (A/+)

and *Arl13b$^{V358A/V358A}$* (A/A) cells following ciliobrevin-D treatment. Violin plots depict

relative fluorescence of IFT88 at cilia tip to cell body with number of cilia measured (n)

listed beneath each plot. (C) Table lists ARL13B (cyan) positive cilia (rabbit anti-

ARL13B; ProteinTech) and the total number of cilia (acetylated α-tubulin: magenta)

examined in control (DMSO) and ciliobrevin-D treated (30μM CB) cell lines.

Representative images show staining for cilia and ARL13B. Two-way ANOVA and

Sidak's multiple comparisons. (*** $p < 0.001$).

Figure 4.

Arl13b^{V358A} mediates normal Shh signaling and neural tube patterning.

(A-I) Whole embryo and neural tube sections of somite-matched littermates at E9.5,

E10.5, and E12.5 stained with indicated markers of cell fate. Whole embryo scale bars:

2 mm. (A'-I' & A"-F") Shh-dependent neural tube patterning at three separate time

points. Cell fate markers are listed above each image. All neural tube scale bars are 50

μm. Neural tube patterning was examined in five embryos for each embryonic stage. E,

embryonic day.

Figure 5.

ARL13B^{V358A} mediates normal ciliary enrichment of Shh components, but not

ARL3 or Inpp5e.

(A) Smo (green) enrichment in ventral neural tube cilia (acetylated α-tubulin: magenta)

is normal in E10.5 embryos. Images are confocal projections. Scale bar is 25 μm. (B-H,

Top) Quantification of average fluorescence intensity in the tip of the cilium (Gli2, Gli3,

and Sufu) or the entire cilium (Ptch1, Smo, ARL3, and Inpp5e) relative to background

level. Violin plots depict relative fluorescent intensity per cilium with number of cilia

examined below each plot. (B-H, Bottom) Representative images for each condition and

cell type with the cilia marker acetylated α-tubulin (magenta) and indicated protein

(green). Wild type (+/+), heterozygous (*A/+*), homozygous (*A/A*) and *hennin* (*hnn*). Data

analyzed by one-way ANOVA and Tukey's multiple comparisons, except Smo data

analyzed by two-way ANOVA and 16 comparisons, corrected $p < 0.003$ deemed

significant. (** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$)

Figure 6.

ARL13B^{V358A} results in decreased ciliation rates and short cilia.

(A) Quantification of ciliation rates in all cell types; wild type (+/+), heterozygous (A/+),

homozygous (A/A) and *hennin (hnn)*. Fewer *Arl13b$^{V358A/V358A}$* and *Arl13b$^{hnn/hnn}$* MEFs

form cilia compared to *Arl13b$^{+/+}$* or *Arl13b$^{V358A/+}$* cells. (B) Quantification of axoneme

length as labeled by acetylated α-tubulin (magenta) in indicated MEFs. Data are

presented as mean (μm) ± standard deviation with the number of cilia measured per

genotype depicted at the base of each bar. Data analyzed by one-way ANOVA and

Tukey's multiple comparisons. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$)

Figure 7.

Model comparing complete loss of ARL13B function to ciliary loss of ARL13B

function

Wildtype (left), *Arl13b^hnnhnn* (middle), and *Arl13b^V358A/V358A* (right) cilia represented as two

halves. On the left half is the organization of Shh components in the presence of Shh

ligand and on the right half is the organization of ARL13B interactors ARL3 and Inpp5e.

(A) ARL13B associates with the ciliary membrane. In the presence of Shh, Ptch1 is

removed from cilia and Smo is visibly enriched in cilia. Smo is activated which promotes

the processing of full-length Gli transcription factors into their activator forms (GliA), that

are shuttled out of the cilium to promote Shh-dependent gene transcription. In addition,

cilia proteins ARL3 and Inpp5e are localized to the primary cilium. (B) In *Arl13b^hnn/hnn*

cells which are null for ARL13B, cilia are shorter than normal. Ciliary Ptch1 and Smo are

visible, although Smo appears punctate instead of diffuse. In addition, loss of ARL13B

decreases transcription of Shh-dependent genes due to lowered GliA. *Arl13b$^{hnn/hnn}$* cilia

also display a failure of Inpp5e and ARL3 to localize to the cilium. Because ARL13B is

the GEF for ARL3, we speculate in this schematic that ARL3 remains GDP bound in

Arl13b$^{hnn/hnn}$ cells. (C) In *Arl13b$^{V358A/V358A}$* cells, ARL13B^{V358A} is not detectable in cilia

and appears diffuse within the cell body. *Arl13b$^{V358A/V358A}$* cilia, like *Arl13b$^{hnn/hnn}$* cilia, are

shorter than wildtype. We observe normal Shh-dependent ciliary Smo enrichment and

normal Shh transcriptional output. However, ARL3 and Inpp5e are absent from cilia,

indicating that ciliary ARL13B is required for ciliary residence of these proteins.

MICE	Sex	wildtype	*V358A/+*	*V358A/V358A*	
Het x Het	M (46.5%)	3	11	6	
Avg. Litter	F (53.5%)	6	8	9	
7.3	% of Total	20.9	44.2	34.9	$\chi^2 = 0.60$
Het x Hom	M (60%)	-	10	11	
Avg. Litter	F (40%)	-	6	8	
7.5	% of Total	-	45.7	54.3	$\chi^2 = 0.69$
Hom x Hom	M (47.4%)	-	-	9	
Avg. Litter	F (52.6%)	-	-	10	
9.5					

Table 1.

852   Genotype of mice born to heterozygous and/or homozygous carrier parents. Data

853   analyzed by chi-squared test.

Supplemental Figure 1.

Supplemental Figure 1.

Overexposure of *Arl13b$^{V358A/V358A}$* cilia in E10.5 neural tube and MEFs reveals no

clear ARL13B^{V358A} presence in cilia.

(A) We identified acetylated α-tubulin positive cilia and used the marker to outline the

area of the cilium. We quantified ARL13B immunofluorescence detected by Caspary lab

polyclonal anti-ARL13B antibody (Caspary et al. 2007) and used the same trace to

measure background levels in a different area of the cell body. (B) *Arl13b$^{V358A/V358A}$*

(*A/A*) and *Arl13b$^{hnn/hnn}$* (*hnn*) MEFs treated with control or ciliobrevin-D 0.5% FBS

Media. Samples were overexposed five times longer than normal, saturating the

detector in the ARL13B channel. We measured no difference in ciliary ARL13B

immunofluorescence between control and ciliobrevin-D treated *Arl13b$^{V358A/V358A}$* and

Arl13b$^{hnn/hnn}$ MEFs. Data presented as violin plots and analyzed by one-way ANOVA.

Respective number of cilia listed below each plot. Cilia that had ARL13B

immunofluorescent readings above background (*) are shown in panel D. (C)

Representative images of over exposed *Arl13b$^{V358A/V358A}$* and *Arl13b$^{hnn/hnn}$* MEFs in

control and ciliobrevin-D treatment conditions. (D) Representative images of cilia with

ARL13B levels above background, marked by * in panel B. (E) Overexposure of cilia in

E10.5 *Arl13b$^{+/+}$*, *Arl13b$^{V358A/V358A}$*, and *Arl13b$^{hnn/hnn}$* embryonic neural tube, stained for

IFT88 and ARL13B (NeuroMab). The ARL13B channel was overexposed for four

seconds instead of one second.