Major role of iron uptake systems in the intrinsic extra-intestinal virulence of the genus *Escherichia* revealed by a genome-wide association study

Marco Galardini[1,6,*], Olivier Clermont[2], Alexandra Baron[2], Bede Busby[3], Sara Dion[2], Sören Schubert[4], Pedro Beltrao[1], Erick Denamur[2,5,*]

[1]EMBL-EBI, Wellcome Genome Campus, Cambridge, United Kingdom
[2] Université de Paris, IAME, UMR1137, INSERM, Paris, France
[3]Genome Biology Unit, EMBL, Heidelberg, Germany
[4]Max von Pettenkofer Institute of Hygiene and Medical Microbiology, Faculty of Medicine, LMU Munich, Germany
[5]AP-HP, Laboratoire de Génétique Moléculaire, Hôpital Bichat, Paris, France
[6]Current address: Biological Design Center, Boston University, Boston, MA 02215, USA

[*]**Corresponding authors:** mgala@bu.edu and erick.denamur@inserm.fr

Abstract:

The genus *Escherichia* is composed of several species and cryptic clades, including *E. coli*, which behave as a vertebrate gut commensal, but also as an opportunistic pathogen involved in both diarrheic and extra-intestinal diseases. To characterize the genetic determinants of extra-intestinal virulence within the genus, we carried out an unbiased genome-wide association study (GWAS) on 370 commensal, pathogenic and environmental strains representative of the *Escherichia* genus phylogenetic diversity and including *E. albertii* (n=7), *E. fergusonii* (n=5), *Escherichia* clades (n=32) and *E. coli* (n=326), tested in a mouse model of sepsis. We found that the high-pathogenicity island (HPI), a ~35 kbp gene island encoding the yersiniabactin siderophore, is highly associated with death in mice, surpassing other associated genetic factors also related to iron uptake, such as the aerobactin and the *sitABCD* operons. We validated the association *in vivo* by deleting key components of the HPI in *E. coli* strains in two phylogenetic backgrounds, and found that virulence is correlated in *E. coli* with growth in the presence of various stressors including several antimicrobials, which hints at collateral sensitivities associated with intrinsic virulence. This study points to the major role of iron capture systems in the extra-intestinal virulence of the genus *Escherichia* and the collateral effects on cell growth of such systems.

Introduction

Members of the *Escherichia* genus are both commensals of vertebrates[1] and opportunistic pathogens[2] involved in a wide range of intestinal and extra-intestinal infections. Apart from the *E. coli* species, it is composed of the cryptic *Escherichia* clades, and the *E. fergusonii* and *E. albertii* species. The latter taxons are rarely isolated in humans but are more frequently found in the environment and avian species where they can cause intestinal infections[3–5]. In humans, extra-intestinal infections represent a considerable burden[6], with bloodstream infections (bacteraemia) being the most severe with a high attributable mortality of between 10-30%[7–10]. The regular increase over the last 20 years of *E. coli* bloodstream incidence[11] and antibiotic resistance[12] is particularly worrisome. The factors associated with high mortality are mainly linked to host conditions such as age, the presence of underlying diseases and to the portal of entry, with the urinary origin being more protective. These factors outweigh those directly attributable to the bacterial agent[7–9,13].

Nevertheless, the use of animal models has shown a great variability in the intrinsic extra-intestinal virulence potential of natural *Escherichia* isolates. In a mouse model of sepsis where bacteria are inoculated subcutaneously, it has been clearly shown that the intrinsic virulence quantified by the number of animal deaths over the number of inoculated animals for a given strain is dependant on the number of virulence factors such as adhesins, toxins, protectins and iron capture systems[14–18]. One of the most relevant virulence factors is the so-called high-pathogenicity island (HPI), a 36 to 43 kb region encoding the siderophore yersiniabactin, a major bacterial iron uptake system[19], which has also been shown to reduce the efficacy of innate immune cells to cause oxidative stress[20]. The deletion of the HPI results in a decrease in the intrinsic virulence in the mouse model in a strain-dependent manner[16,17,21], indicating complex interactions between the genetic background of each strain and the HPI.

The limitation of these gene KO studies is that they target specific candidate genes. Recently, the development of new approaches in bacterial genome-wide association studies (GWAS)[22–25] allows searching in an unbiased manner for genotypes associated with specific phenotypes such as drug resistance or virulence. In this context, we conducted a GWAS in 370 commensal and extra-intestinal pathogenic strains of *E. coli*, and related *Escherichia* clades, as well as *E. fergusonii* and *E. albertii*, representing the genus phylogenetic diversity, to search for traits associated with virulence in the mouse model of sepsis[26]. Most of the strains were isolated from a human host and are divided between commensals and extra-intestinal pathogenic. Most importantly, many (N=186) of these strains have been recently phenotyped across hundreds of growth conditions, including antibiotics and other chemical and physical stressors[27]. This data could then be used to find phenotype associations with virulence and to generate hypotheses on

70 the function of genetic variants associated with the extra-intestinal virulence phenotype
71 and potential collateral sensitivities associated with them.

Results

GWAS identifies the high-pathogenicity island as the strongest driver of the extra-intestinal virulence phenotype

We studied a 326 strain collection representative of the *E. coli* phylogenetic diversity,
with strains belonging to phylogroups A (N=72), B1 (N=41), B2 (N=111), C (N=36), D
(N=20), E (N=19), F (N=12) and G (N=15). To have a broader phylogenetic
representation, we also included strains from *Escherichia* clades I to V (N=32) and the
species *E. albertii* (N=7) and *E. fergusonii* (N=5)[28]. These strains encompass 170
commensal strains and 187 strains isolated in various extra-intestinal infections, mainly
urinary tract infections and septicaemia[7,14,29–35]. To avoid any bias linked to host
conditions, we assessed the strain virulence as its intrinsic extra-intestinal pathogenic
potential using a well-calibrated mouse model of sepsis[14,26], expressed as the number of
killed mice over the 10 inoculated per strain. In accordance with previous data[14,26,36–38],
phylogroup B2 is the most associated with the virulence phenotype ($2E^{-9}$ Wald test
p-value, Figure 1A, Supplementary Table 1).

We used a bacterial GWAS method to associate unitigs to the virulence phenotype,
allowing us to simultaneously test the contribution of core and accessory genome
variation to pathogenicity[24]. It is generally understood that such methods require large
sample sizes to have sufficient power, partly due to the need to break the long clonal
frames typical of bacterial genomes; the appropriate sample size is also a function of
the penetrance of the causal variants[23,39]. We ran simulations with an unrelated set of
complete *E. coli* genomes and verified that our sample size was appropriate for variants
with high penetrance (i.e. odds ratio above 5, Supplementary Figure 1, Methods). We
reasoned that the genetic determinants of virulence are likely to have a relatively high
penetrance, and that the strains used were genetically diverse, enough to break the
clonal frame.

We uncovered a statistically significant association between 5,214 unitigs and the
virulence phenotype, which were mapped back to 81 genes across the strains'
pangenome (Figure 1B, Supplementary Table 2, Methods). To understand whether the
presence of these genes is directly associated with virulence or if its due to genetic
variants such as SNPs we performed a separate association using genes' presence
absence patterns, showing that most genes have an odds ratio that far exceeds the
required threshold we estimated from simulations, as well as low association p-value
(Figure 1C). Furthermore, 48 out of 81 genes with at least an associated unitig mapped
to them have a frequency across strains that is highly correlated with that of the

associated unitigs (Figure 1D), indicating that it's the presence of those genes to be associated with virulence. Genes belonging to the HPI had the lowest association p-value by far ($<1E^{-28}$); the presence of two additional operons encoding for bacterial siderophores (aerobactin[40] and *sitABCD*[41]) was also found to be associated with virulence (Figure 1E). We found that the HPI structure was highly conserved across the genomes that encode it (Supplementary Figure 2). We also observed that the distribution of known virulence factors didn't match the virulence phenotype as closely as the HPI or the aerobactin and *sitABCD* operons, or had unitigs passing the association threshold (Supplementary Figure 3). The remaining 33 genes have a high frequency in the pangenome, indicating the presence of genetic variants such as SNPs present in core genes; the *mtfA* gene, which is involved in the regulation of carbohydrate metabolism[42], had the lowest association p-value among those genes ($1E^{-14}$).

Figure 1. The HPI is strongly associated with the extra-intestinal virulence phenotype assessed in the mouse sepsis assay. A) Core genome phylogenetic tree of the *Escherichia* strains used in this study rooted on *E. albertii* strains. Outer ring reports virulence as the number of killed mice over the 10 inoculated per strain, inner ring the phylogroup or species each strain belongs to. B) Results of the unitigs association analysis: for each gene the minimum association p-value and average minimum allele frequency (MAF) across all mapped unitigs is reported. The gene length fraction is computed by dividing the total length of mapped unitigs by the length of the gene. C) Results of the gene presence/absence association analysis; each gene tested is represented. Genes from the unitigs association analysis are highlighted. OG: orthologous groups. D) Scatterplot of gene frequency versus frequency of associated unitigs; points on the diagonal indicate hits where the association is most likely due to a gene's presence rather than a SNP. E) The structure of the HPI and of the aerobactin and *sitABCD* operons in strain IAI39; all associated genes are highlighted.

KO gene experiments validate the role of the HPI in the extra-intestinal phenotype
The studies on the role of the HPI in experimental virulence gave contrasting results according to the strains' genetic background[17]. Among B2 phylogroup strains, HPI deletion in the 536 (ST127) strain did not have any effect in the mouse model of sepsis[43] whereas this deletion in the NU14 (ST95) strain dramatically attenuated virulence[17]. Two strains from this study belonging to B2 phylogroup/ST141 (IAI51 and IAI52) deleted in *irp1* have attenuated virulence in the same model[21]. To have a broader view of the role of the HPI in various genetic backgrounds, we constructed *irp2* deletion gene mutants in two strains of phylogroup D (NILS46) and A (NILS9) belonging to STs (Sequence Types) frequently involved in human bacteraemia (ST69 and ST10, respectively)[44]. We first verified that the wild-type strains strongly produced yersiniabactin, whereas the *irp2* mutants did not (Figure 2A). We then tested them in the mouse sepsis model and saw an increase in survival for both mutated strains (log-rank test p-value < 0.0001 and 0.0217 for strain NILS9 and NILS46, respectively, Figure 2B, Supplementary Table 3) with no significant difference between the survival profiles for the two mutants (log-rank test p-value > 0.1).

 We have therefore validated *in vivo* the causal link between the HPI and the virulence phenotype detected by the means of an unbiased association approach, which demonstrates the power and accuracy of bacterial GWAS.

Figure 2. Phenotypic consequences of HPI deletion. A) Deletion of HPI leads to a decrease in production of yersiniabactin. Production of yersiniabactin is measured using a luciferase-based reporter (Methods). Strains marked with a "-" and "+" sign indicate a negative and positive control, respectively. The red dashed line indicates an arbitrary threshold for yersiniabactin production, derived from the average signal recorded from the negative controls plus two standard deviations. RLU, relative light units. B) Deletion of HPI leads to an increase in survival after infection. Survival curves for wild-type strains and the corresponding *irp2* deletion mutant, built after infection of 20 mice for each strain.

High-throughput phenotypic data sheds light on HPI and other iron capture systems function

The main function encoded by the HPI cassette is iron scavenging through the expression of the siderophore yersiniabactin[21], which has been previously validated in *E. coli* through knockout experiments[17]. The aerobactin operon also encodes an iron chelator[40], while the *sitABCD* operon encodes a Mn^{2+}/Fe^{2+} ion transporter[41]. In order to investigate other putative functions of these operons and their relationship with virulence, we leveraged a previously-generated high-throughput phenotypic screening in an *E. coli* strain panel that largely overlaps with the strains used here (186 over 370)[27]. We observed a relatively high correlation between growth profiles in certain conditions and both virulence and presence of the HPI, aerobactin and *sitABCD* operons (Figure 3A-D, Supplementary Table 4).

As expected, we found a positive correlation between growth on the iron-sequestering agent pentetic acid[45] and both virulence and HPI/aerobactin/*sitABCD* presence (Pearson's r: 0.36, 0.48, 0.23 and 0.39, respectively). We also found that growth in the presence of bipyridyl, an iron chelator, was positively correlated with the presence of aerobactin (bipyridyl + tobramycin, Pearson's r: 0.30). We similarly observed a positive correlation between growth with pyocyanin, a redox-active phenazine compound being able to reduce Fe^{3+} to Fe^{2+}[46], and both HPI/aerobactin/*sitABCD* presence (Pearson's r: 0.35, 0.28, 0.26 and 0.27 respectively). All growth conditions agree with the iron scavenging function of the three gene clusters and their importance for virulence.

Interestingly, we also found similarly strong positive correlations with growth on sub-inhibitory concentrations of several antibiotics, such as rifampicin, ciprofloxacin, tetracycline and ß-lactams such as amoxicillin, oxacillin and meropenem, as well as other antimicrobial agents such as cerulenin and colicin. On the other hand we found that growth in presence of indole at 2 mM either in association with sub-inhibitory concentrations of cefsulodin and tobramycin, or alone at 40°C was negatively correlated with both virulence and HPI/aerobactin/*sitABCD* presence. Similar negative correlation was observed with aerobactin presence and the MAC13243 compound that increases outer membrane permeability[47]. These correlations might be due to the presence/absence of acquired resistance alleles and/or genes that are strongly associated with pathogenic strains, or might point to the role of iron homeostasis in intrinsic resistance to antibiotics[48]. To investigate these two hypotheses, we focused on tetracycline resistance, a common occurrence in the genus[33,37,49], and for which resistance genes can be easily found through sequence homology (Methods). We measured the correlation between the presence of tetracycline resistant genes, found in 26.8% of the strains, and virulence (Pearson's r: 0.16), as well with the presence of

either of the three iron capture systems (Pearson's r: 0.21, 0.33 and 0.24 for HPI,
aerobactin and *sitABCD*, respectively), which we found to be all comparable with the
direct correlation between growth on sub-inhibitory concentration of tetracycline and the
presence of resistance genes (Pearson's r: 0.4). As such, both scenarios seem
plausible to explain the relationship between growth in presence of tetracycline and both
virulence and presence of HPI/aerobactin/*sitABCD*.
Given the relatively large number of conditions correlated with both virulence and
presence of iron uptake systems, we tested whether these features could be predicted
from growth profiles. We used the commonly-used random forests machine learning
algorithm with appropriate partitioning of input data to tune hyperparameters and reduce
overfitting, leading to four classifiers for virulence and presence of the HPI, aerobactin
and *sitABCD* operons with high predictive power, with the exception of aerobactin
(Figure 3E and 3F and Methods). We noted that prediction of the gene clusters
presence performs slightly better than virulence, possibly reflecting the complex nature
of the latter phenotype. As expected, we found that conditions with relatively high
correlation with each feature have a higher weight across classifiers (Figure 3G,
Supplementary Table 5), which suggests that a subset of phenotypic tests might be
sufficient to classify pathogenic strains. These results show how phenotypic data can be
used to generate hypotheses for the function of virulence factors.

Figure 3. Growth profiles can predict virulence and presence of virulence factors. A-D)
Volcano plots for the correlation between the strains' growth profiles and: A) virulence levels, B)
presence of the HPI, C) presence of aerobactin, and D) presence of *sitABCD*. E-F) Use of the

strains' growth profiles to build a predictor of virulence levels and presence of the three iron
uptake systems. E) Receiver operating characteristic (ROC) curve and F) Precision-Recall
curve for the four tested predictors. G) Feature importance for the predictors, showing the top
15 conditions contributing to the virulence levels predictor.

Discussion

With the steady decline in the price of genomic sequencing and the increasing
availability of molecular and phenotypic data for bacterial isolates, it has finally become
possible to use statistical genomics approaches such as GWAS to uncover the genetic
determinants of relevant phenotypes. Such approaches have the advantage of being
unbiased, and can then be used to confirm previous targeted findings and potentially
uncover new factors, given sufficient statistical power. The accumulation of other
molecular and phenotypic data can on the other hand uncover variables correlated with
phenotype, which can be used to generate testable hypotheses on the function of
genomic hits and potential collateral sensitivities associated with them. Given the rise of
both *E. coli* extra-intestinal infections and antimicrobial resistance, we reasoned that the
intrinsic virulence assessed in a calibrated mouse model of sepsis[14,26] is a phenotype
worth exploring with such an unbiased approach.

We were able to confirm earlier reports on the importance of the HPI in
extra-intestinal virulence[17–19,21,50,51], which showed the strongest signal in both the unitigs
and accessory genome association analysis, and whose importance was validated *in*
vitro and in an *in vivo*. The distribution of the HPI within the species resulting from
multiple horizontal gene transfers via homologous recombination[52] has probably
facilitated its identification using GWAS. We associated additional genetic factors to
intrinsic virulence, such as the aerobactin and *sitABCD* operons, both related to iron
scavenging together with the HPI. We also found mutations in the *mtfA* core gene to be
associated with virulence, although its role in virulence remains elusive. Additional
factors might have been overlooked by this analysis, due to the relatively small sample
size; we however estimate that those putative additional factors might have a relatively
low penetrance, based on our simulations in an independent dataset. As sequencing of
bacterial isolates is becoming more common in clinical settings[53–55], we expect to be
able to uncover these additional genetic factors in future studies.

The association between both the intrinsic virulence phenotype and the presence of
the virulence factors - such as the HPI - and previously collected growth data allowed us
to generate testable hypotheses on mechanism of pathogenesis and putative additional
functions of these factors. In particular we observed a relatively strong correlation
between growth on various antimicrobial agents and both virulence and
HPI/aerobactin/*sitABCD* presence, which seems to confirm the pressure to acquire
resistance for these isolates, as exemplified for tetracycline resistance genes, but also

raises the question about the potential role of iron homeostasis on antimicrobial resistance[48]. As we observed several classes of antibiotics with different molecular targets, it is possible that this is due to a pleiotropic effect. *E. coli* Fur mutants, a transcriptional regulator that represses iron uptake systems, which accumulate high level of intracellular iron, have been shown to increase resistance to quinolones, aminoglycoside, tetracycline, rifampicin and amoxicillin[56]. Cell envelope permeability can also be modified in response to the presence of iron via two-component systems, rendering the cell more resistant[48]. Sub-lethal doses of antibiotics have also been shown to behave as a stressor for the cells that engage a general core hormetic stress response consisting of increased energy production and translation, allowing growth when nutrients are available despite the constant presence of the stressor[57]. When the resources are instead limited, cell growth is stopped. It can be hypothesised that iron is part of these limiting nutrients and that iron capture systems allow coping with the stressors, i.e. the antibiotics.

The negative correlation between virulence and iron capture systems and growth profiles in the presence of indole associated with stress conditions such as sub-lethal doses of antibiotics or high temperature but not indole alone, points however to the possible deleterious role of iron in such conditions. In lysogeny broth, sub-lethal concentrations of antibiotics increase the endogenous production of indole[57] and, at very high concentration (*e.g.* 5 mM), indole induces the production of reactive oxygen species and is toxic for the cells[58]. A vicious circle is rapidly established as antibiotics increase the production of indole[57], which in turn destabilises the membrane[58], further increasing the penetration of the antibiotics. This toxicity has been shown to be partly iron mediated due to the Fenton reaction, the deletion of TonB, an iron transporter, increasing resistance to the antibiotic[59]. Interestingly, tobramycin is the antibiotic involved in this negative correlation (Fig. 3A to D). Sub-lethal doses of this aminoglycoside lead to an increase of reactive oxygen species in the bacterial cell in relation to intra-cellular iron and Fenton reaction[60]. Thus, cells with increased import of extracellular iron might be more sensitive to sub-lethal doses of antibiotics, suggesting the potential for collateral sensitivities related to both intrinsic virulence and the presence of the iron uptake systems. Altogether, these data bring new light on the "liaisons dangereuses" between iron and antibiotics that could potentially be targeted[48]. More generally, they show that the presence of iron capturing systems can be either advantageous or disadvantageous, depending on the growth conditions.

In conclusion, we showed the power of bacterial GWAS to identify major virulence determinants in bacteria. Within the *Escherichia* genus, iron capture systems are the main predictors of the intrinsic extra-intestinal virulence. Furthermore, this analysis demonstrates how a data-centric approach can increase our knowledge of complex

bacterial phenotypes and guide further empirical work on gene function and its relationship to intrinsic virulence.

Materials and methods

Strains used

The full list of the 370 strains used in the association analysis, together with their main characteristics is reported in Supplementary Table 1. These strains belong to various published collections: ECOR (N=71)[29], IAI (n=81)[14], NILS (N=82)[31], Septicoli (N=39)[10], ROAR (N=30)[32], Guyana (N=12)[30], Coliville (N=8)[33], FN (N=6)[34], COLIRED (N=3)[35], COLIBAFI (N=2)[7], correspond to archetypal strains (N=7) or are miscellaneous strains from our personal collections (N=29). The isolation host is predominantly humans (N=291), followed by animals (N=72) and isolated from the environment (N=6). One hundred and seventy strains were commensal whereas five and 187 were responsible of intestinal and extra-intestinal infections, respectively. The genomes of 295 strains were previously available, while the remaining 75 were sequenced as part of this work by Illumina technology as described previously[35]. The genome sequences of all strains are available through Figshare[61].

The construction of the *irp2* deletion mutants of the NILS9 and NILS46 strains was achieved following a strategy adapted from Datsenko and Wanner[62]. Primers used in the study are listed in Supplementary Table 6. In brief, primers used for gene disruption included 44-46 nucleotide homology extensions to the 5'- and 3' regions of the target gene, respectively, and additional 20 nucleotides of priming sequence for amplification of the resistance cassette on the template plasmids pKD4. The PCR product was then transformed into strains carrying the helper plasmid pKOBEG expressing the lambda red recombinase under control of an arabinose-inducible promoter[63]. Kanamycin resistant transformants were selected and further screened for correct integration of the resistance marker by PCR. For elimination of the antibiotic resistance gene, helper plasmid pCP20 was used according to the published protocol. PCR followed by Sanger sequencing of the mutants were performed to verify the deletion and the presence of the expected scar.

Yersiniabactin detection assay

Production of the siderophore yersiniabactin was detected and quantified using a luciferase reporter assay as described elsewhere[17,64]. Briefly, bacterial strains were cultivated in NBD medium for 24 hours at 37°C. Next, bacteria were pelleted by centrifugation and the supernatant was added to the indicator strain WR 1542 harbouring plasmid pACYC5.3L. All the genes necessary for yersiniabactin uptake are located on the plasmid pACYC5.3L, i.e. *irp6, irp7, irp8, fyuA, ybtA*. Furthermore, this plasmid is equipped with a fusion of the *fyuA* promoter region with the luciferase

reporter gene. The amount of yersiniabactin can be quantified semi-quantitatively, as yersiniabactin-dependant upregulation of *fyuA* expression is determined by luciferase activity of the *fyuA-luc* reporter fusion.

Mouse virulence assay

Ten female mice OF1 of 14-16 g (4 week-old) from Charles River® (L'Arbresle, France) received a subcutaneous injection of 0.2 ml of bacterial suspension in the neck ($2 \cdot 10^8$ colony forming unit). Time to death was recorded during the following 7 days. Mice surviving more than 7 days were considered cured and sacrificed[14]. In each experiment, the *E. coli* CFT073 strain was used as a positive control killing all the inoculated mice whereas the *E. coli* K-12 MG1655 strain was used as a negative control for which all the inoculated mice survive[26]. For the mutant assays, 20 mice per strain were used to obtain statistical relevant data. The data was analysed using the lifeline package v0.21.0[65].

Association analysis

All genome-wide association analysis were carried out using pyseer, version v1.3.4[24]. All input genomes were re-annotated using prokka, version v1.14.5[66], to ensure uniform gene calls and excluding contigs whose size was above 200 base pairs. The core genome phylogenetic tree was generated using ParSNP[67] to generate the core genome alignment and gubbins v2.3.5[68] to generate the phylogenetic tree. The strain's pangenome was estimated using roary v3.13.0[69]. Unitigs distributions from the input genome assemblies were computed using unitig-counter v1.0.5. The association between both unitigs and pangenome and phenotype (expressed as number of mice killed post-infection) was carried out using the FastLMM[70] linear mixed-model implemented in pyseer, using a kinship matrix derived from the phylogenetic tree as population structure. For both association analysis we used the number of unique presence/absence patterns to derive an appropriate p-value threshold for the association likelihood ratio test ($2.16E^{-08}$ and $5.45E^{-06}$ for the unitigs and pangenome analysis, respectively). Unitigs significantly associated with the phenotype were mapped back to each input genome using bwa mem v0.7.17-r1188[71] and betools v.2.29.2[72], using the pangenome analysis to collapse gene hits to individual groups of orthologs. A sample protein sequence for each groups of orthologs where at least one unitig with size 20 or higher was mapped was extracted giving priority to strain IAI39 when available, given it was the only strain with a complete genome available[73]; those sample sequences where used to search for homologs in the uniref50 database from uniprot[74] using blast v2.9.0[75]. Each group of orthologs was then given a gene name using both available literature information and the results of the homology search.

Power simulations

Statistical power was estimated using an unrelated set of 548 complete *E. coli* genomes downloaded from NCBI RefSeq using ncbi-genome-download v0.2.9 on May 24th 2018. Each genome was subject to the same processing as the actual ones used in the real analysis (re-annotation, phylogenetic tree construction, pangenome estimation). The gene presence/absence patterns were used to run the simulations, in a similar way as described in the original SEER implementation[23]. Briefly, for each sample size, a random subset of strains was selected, and the likelihood ratio test p-value threshold was estimated by counting the number of unique gene presence/absence patterns in the reduced roary matrix. For each odds ratio tested, a binary case-control phenotype vector was constructed for the strains subset using the following formulae:

$$P_{case|variant} = \frac{D_e}{MAF}$$

$$P_{case|novariant} = \frac{\frac{S_r}{S_r+1} - D_e}{1 - MAF}$$

Were S_r is the ratio of case/controls (set at 1 in these simulations), MAF as the minimum allele frequency of the target gene in the strains subset, and D_e the number of cases. pyseer's LMM model was then applied to the presence/absence vector of the target gene and the likelihood ratio test p-value was compared with the empirical threshold. The randomization was repeated 100 times and power was defined as the proportion of randomizations for each sample size and odds ratio whose p-value was below the threshold. The *pks2* and *fabG* genes were used as gene targets in the simulations, and both gave very similar results.

Correlations with growth profiles

The previously generated phenotypic data[27] for 186 over 370 strains were used to compute correlations with both the number of mice killed after infection and presence/absence of the associated virulence factors. The data was downloaded from the ecoref website (https://evocellnet.github.io/ecoref/download/) and the pearson correlation with the s-scores was computed together with the correlation p-value. Prediction of tetracycline resistance was carried out using staramr v0.7.1 with the ResFinder database[76]. Four predictors, one for virulence (number of killed mice post-infection) and one for presence of the HPI, aerobactin and the *sitABCD* operon were built using the random forest classifier algorithm implemented in scikit-learn v.022.0[77], using the s-scores as predictors. The input was column imputed, and 33% of the observations were kept as a test dataset, using a "stratified shuffle split" strategy. The remainder was used to train the classifier, using a grid search to select the number

of trees and the maximum number of features used, through 10 rounds of stratified shuffle split with validation set size of 33% the training set and using the F1 measure as score. The performance of the classifiers on the test set were assessed by computing the area under the receiver operating characteristic curve (ROC-curve).

Code and data availability

Ethics statement

All animal experimentations were conducted following European (Directive 2010/63/EU on the protection of animals used for scientific purposes) and national recommendations (French Ministry of Agriculture and French Veterinary Services, accreditation A 75-18-05). The protocol was approved by the Animal Welfare Committee of the Veterinary Faculty in Lugo, University of Santiago de Compostela (AE-LU-002/12/INV MED.02/OUTROS 04).

Acknowledgements

We are grateful to Ivan Matic for discussion on the effect of indole. This work was partially supported by the "Fondation pour la Recherche Médicale" (Equipe FRM 2016, grant number DEQ20161136698).

References

1. Tenaillon, O., Skurnik, D., Picard, B. & Denamur, E. The population genetics of commensal Escherichia coli. *Nat. Rev. Microbiol.* **8**, 207–217 (2010).
2. Croxen, M. A. & Brett Finlay, B. Molecular mechanisms of Escherichia coli pathogenicity. *Nature Reviews Microbiology* vol. 8 26–38 (2010).
3. Oaks, J. L. *et al.* Escherichia albertii in wild and domestic birds. *Emerg. Infect. Dis.* **16**, 638 (2010).
4. Clermont, O., Gordon, D. M., Brisse, S., Walk, S. T. & Denamur, E. Characterization of the cryptic Escherichia lineages: rapid identification and prevalence. *Environ. Microbiol.* **13**, 2468–2477 (2011).
5. Blyton, M. D. J. *et al.* Genetic Structure and Antimicrobial Resistance of Escherichia coli and Cryptic Clades in Birds with Diverse Human Associations. *Appl. Environ. Microbiol.* **81**, 5123–5133 (2015).
6. Russo, T. A. & Johnson, J. R. Medical and economic impact of extraintestinal infections due to Escherichia coli: focus on an increasingly important endemic problem. *Microbes Infect.* **5**, 449–456 (2003).
7. Lefort, A. *et al.* Host Factors and Portal of Entry Outweigh Bacterial Determinants To Predict the Severity of Escherichia coli Bacteremia. *Journal of Clinical Microbiology* vol. 49 777–783 (2011).
8. Burdet, C. *et al.* Escherichia coli bacteremia in children: age and portal of entry are the main

441   predictors of severity. *Pediatr. Infect. Dis. J.* **33**, 872–879 (2014).

442 9. Abernethy, J. K. *et al.* Thirty day all-cause mortality in patients with Escherichia coli bacteraemia in
443   England. *Clin. Microbiol. Infect.* **21**, 251.e1–8 (2015).

444 10. V. de Lastours, C. Laouénan, G. Royer, E. Carbonnelle, R. Lepeule, M. Esposito-Farèse, O.
445   Clermont, X. Duval, B. Fantin, F. Mentré, JW Decousser, E. Denamur, A. Lefort, on behalf of the
446   SEPTICOLI group. Mortality in Escherichia coli bloodstream infections: antibiotic resistance still does
447   not make it. *J. Antimicrob. Chemother.*

448 11. Vihta, K.-D. *et al.* Trends over time in Escherichia coli bloodstream infections, urinary tract infections,
449   and antibiotic susceptibilities in Oxfordshire, UK, 1998–2016: a study of electronic health records.
450   *The Lancet Infectious Diseases* vol. 18 1138–1149 (2018).

451 12. Cassini, A. *et al.* Attributable deaths and disability-adjusted life-years caused by infections with
452   antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a population-level
453   modelling analysis. *Lancet Infect. Dis.* **19**, 56–66 (2019).

454 13. Baudron, C. R. *et al.* Escherichia coli bacteraemia in adults: age-related differences in clinical and
455   bacteriological characteristics, and outcome. *Epidemiology & Infection* **142**, 2672–2683 (2014).

456 14. Picard, B. *et al.* The link between phylogeny and virulence in Escherichia coli extraintestinal infection.
457   *Infect. Immun.* **67**, 546–553 (1999).

458 15. Johnson, J. R. & Kuskowski, M. Clonal origin, virulence factors, and virulence. *Infection and immunity*
459   vol. 68 424–425 (2000).

460 16. Tourret, J., Diard, M., Garry, L., Matic, I. & Denamur, E. Effects of single and multiple pathogenicity
461   island deletions on uropathogenic Escherichia coli strain 536 intrinsic extra-intestinal virulence. *Int. J.*
462   *Med. Microbiol.* **300**, 435–439 (2010).

463 17. Smati, M. *et al.* Strain-specific impact of the high-pathogenicity island on virulence in extra-intestinal
464   pathogenic Escherichia coli. *Int. J. Med. Microbiol.* **307**, 44–56 (2017).

465 18. Johnson, J. R. & Russo, T. A. Molecular Epidemiology of Extraintestinal Pathogenic Escherichia coli.
466   *EcoSal Plus* **8**, (2018).

467 19. Schubert, S., Cuenca, S., Fischer, D. & Heesemann, J. High-pathogenicity island of Yersinia pestis in
468   enterobacteriaceae isolated from blood cultures and urine samples: prevalence and functional
469   expression. *J. Infect. Dis.* **182**, 1268–1271 (2000).

470 20. Paauw, A., Leverstein-van Hall, M. A., van Kessel, K. P. M., Verhoef, J. & Fluit, A. C. Yersiniabactin
471   reduces the respiratory oxidative stress response of innate immune cells. *PLoS One* **4**, e8240
472   (2009).

473 21. Schubert, S., Picard, B., Gouriou, S., Heesemann, J. & Denamur, E. Yersinia high-pathogenicity
474   island contributes to virulence in Escherichia coli causing extraintestinal infections. *Infect. Immun.* **70**,
475   5335–5337 (2002).

476 22. Earle, S. G. *et al.* Identifying lineage effects when controlling for population structure improves power
477   in bacterial association studies. *Nature Microbiology* **1**, 1–8 (2016).

478 23. Lees, J. A. *et al.* Sequence element enrichment analysis to determine the genetic basis of bacterial
479   phenotypes. *Nat. Commun.* **7**, 12797 (2016).

480 24. Lees, J., Galardini, M., Bentley, S. D. & Weiser, J. N. pyseer: a comprehensive tool for microbial
481   pangenome-wide association studies. *bioRxiv* (2018).

482 25. Jaillard, M. *et al.* A fast and agnostic method for bacterial genome-wide association studies: Bridging
483   the gap between k-mers and genetic events. *PLoS Genet.* **14**, e1007758 (2018).

484 26. Johnson, J. R. *et al.* Experimental mouse lethality of Escherichia coli isolates, in relation to accessory
485   traits, phylogenetic group, and ecological source. *J. Infect. Dis.* **194**, 1141–1150 (2006).

486 27. Galardini, M. *et al.* Phenotype inference in an Escherichia coli strain panel. *Elife* **6**, 1–19 (2017).

487 28. Clermont, O., Christenson, J. K., Denamur, E. & Gordon, D. M. The Clermont Escherichia coli
488   phylo-typing method revisited: improvement of specificity and detection of new phylo-groups.
489   *Environ. Microbiol. Rep.* **5**, 58–65 (2013).

490 29. Ochman, H. & Selander, R. K. Standard reference strains of Escherichia coli from natural
491   populations. *J. Bacteriol.* **157**, 690–693 (1984).

492 30. Lescat, M. *et al.* Commensal Escherichia coli strains in Guiana reveal a high genetic diversity with
493   host-dependant population structure. *Environ. Microbiol. Rep.* **5**, 49–57 (2013).

494 31. Bleibtreu, A. *et al.* The rpoS gene is predominantly inactivated during laboratory storage and

undergoes source-sink evolution in Escherichia coli species. *J. Bacteriol.* **196**, 4276–4284 (2014).

32. Skurnik, D. *et al.* Emergence of Antimicrobial-Resistant Escherichia coli of Animal Origin Spreading
in Humans. *Mol. Biol. Evol.* **33**, 898–914 (2016).

33. Massot, M. *et al.* Phylogenetic, virulence and antibiotic resistance characteristics of commensal
strain populations of Escherichia coli from community subjects in the Paris area in 2010 and
evolution over 30 years. *Microbiology* **162**, 642–650 (2016).

34. Nowrouzian, F. L. *et al.* Escherichia coli B2 Phylogenetic Subgroups in the Infant Gut Microbiota:
Predominance of Uropathogenic Lineages in Swedish Infants and Enteropathogenic Lineages in
Pakistani Infants. *Appl. Environ. Microbiol.* **85**, (2019).

35. Bourrel, A. S. *et al.* Colistin resistance in Parisian inpatient faecal Escherichia coli as the result of two
distinct evolutionary pathways. *J. Antimicrob. Chemother.* **74**, 1521–1530 (2019).

36. Moissenet, D. *et al.* Meningitis caused by Escherichia coli producing TEM-52 extended-spectrum
beta-lactamase within an extensive outbreak in a neonatal ward: epidemiological investigation and
characterization of the strain. *J. Clin. Microbiol.* **48**, 2459–2463 (2010).

37. Ingle, D. J. *et al.* Biofilm formation by and thermal niche and virulence characteristics of Escherichia
spp. *Appl. Environ. Microbiol.* **77**, 2695–2700 (2011).

38. Clermont, O. *et al.* Characterization and rapid identification of phylogroup G in Escherichia coli, a
lineage with high virulence and antibiotic resistance potential. *Environ. Microbiol.* **21**, 3107–3117
(2019).

39. Power, R. A., Parkhill, J. & de Oliveira, T. Microbial genome-wide association studies: lessons from
human GWAS. *Nat. Rev. Genet.* **18**, 41–50 (2016).

40. Warner, P. J., Williams, P. H., Bindereif, A. & Neilands, J. B. ColV plasmid-specific aerobactin
synthesis by invasive strains of Escherichia coli. *Infect. Immun.* **33**, 540–545 (1981).

41. Bearden, S. W., Staggs, T. M. & Perry, R. D. An ABC transporter system of Yersinia pestis allows
utilization of chelated iron by Escherichia coli SAB11. *J. Bacteriol.* **180**, 1135–1147 (1998).

42. Becker, A.-K. *et al.* YeeI, a novel protein involved in modulation of the activity of the
glucose-phosphotransferase system in Escherichia coli K-12. *J. Bacteriol.* **188**, 5439–5449 (2006).

43. Diard, M. *et al.* Pathogenicity-associated islands in extraintestinal pathogenic Escherichia coli are
fitness elements involved in intestinal colonization. *J. Bacteriol.* **192**, 4885–4893 (2010).

44. Kallonen, T. *et al.* Systematic longitudinal survey of invasive Escherichia coli in England
demonstrates a stable population structure only transiently disturbed by the emergence of ST131.
Genome Res. (2017) doi:10.1101/gr.216606.116.

45. Pippard, M. J., Jackson, M. J., Hoffman, K., Petrou, M. & Modell, C. B. Iron chelation using
subcutaneous infusions of diethylene triamine penta-acetic acid (DTPA). *Scand. J. Haematol.* **36**,
466–472 (1986).

46. Cornelis, P. & Dingemans, J. Pseudomonas aeruginosa adapts its iron uptake strategies in function
of the type of infections. *Front. Cell. Infect. Microbiol.* **3**, 75 (2013).

47. Muheim, C. *et al.* Increasing the permeability of Escherichia coli using MAC13243. *Scientific Reports*
vol. 7 (2017).

48. Ezraty, B. & Barras, F. The 'liaisons dangereuses' between iron and antibiotics. *FEMS Microbiol. Rev.*
40, 418–435 (2016).

49. Mazel, D., Dychinco, B., Webb, V. A. & Davies, J. Antibiotic resistance in the ECOR collection:
integrons and identification of a novel aad gene. *Antimicrob. Agents Chemother.* **44**, 1568–1574
(2000).

50. Johnson, J. R. *et al.* Contribution of yersiniabactin to the virulence of an Escherichia coli sequence
type 69 ('clonal group A') cystitis isolate in murine models of urinary tract infection and sepsis.
Microb. Pathog. **120**, 128–131 (2018).

51. Johnson, J. R. *et al.* Accessory Traits and Phylogenetic Background Predict Escherichia coli
Extraintestinal Virulence Better Than Does Ecological Source. *J. Infect. Dis.* **219**, 121–132 (2019).

52. Schubert, S. *et al.* Role of Intraspecies Recombination in the Spread of Pathogenicity Islands within
the Escherichia coli Species. *PLoS Pathog.* **5**, e1000257 (2009).

53. Fricke, W. F. & Rasko, D. A. Bacterial genome sequencing in the clinic: bioinformatic challenges and
solutions. *Nat. Rev. Genet.* **15**, 49–55 (2014).

54. Quainoo, S. *et al.* Whole-Genome Sequencing of Bacterial Pathogens: the Future of Nosocomial

Outbreak Analysis. *Clin. Microbiol. Rev.* **30**, 1015–1063 (2017).

55. Tagini, F. & Greub, G. Bacterial genome sequencing in clinical microbiology: a pathogen-oriented
review. *Eur. J. Clin. Microbiol. Infect. Dis.* **36**, 2007–2020 (2017).

56. Nichols, R. J. *et al.* Phenotypic landscape of a bacterial cell. *Cell* **144**, 143–156 (2011).

57. Mathieu, A. *et al.* Discovery and Function of a General Core Hormetic Stress Response in E. coli
Induced by Sublethal Concentrations of Antibiotics. *Cell Rep.* **17**, 46–57 (2016).

58. Garbe, T. R., Kobayashi, M. & Yukawa, H. Indole-inducible proteins in bacteria suggest membrane
and oxidant toxicity. *Arch. Microbiol.* **173**, 78–82 (2000).

59. Giroux, X., Su, W.-L., Bredeche, M.-F. & Matic, I. Maladaptive DNA repair is the ultimate contributor
to the death of trimethoprim-treated cells under aerobic and anaerobic conditions. *Proc. Natl. Acad.*
Sci. U. S. A. **114**, 11512–11517 (2017).

60. Baharoglu, Z., Krin, E. & Mazel, D. RpoS plays a central role in the SOS induction by sub-lethal
aminoglycoside concentrations in Vibrio cholerae. *PLoS Genet.* **9**, e1003421 (2013).

61. Galardini, M. Escherichia coli pathogenicity GWAS: input genome sequences (updated). (2020)
doi:10.6084/m9.figshare.11879340.v1.

62. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in Escherichia coli
K-12 using PCR products. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 6640–6645 (2000).

63. Chaveroche, M. K., Ghigo, J. M. & d'Enfert, C. A rapid method for efficient gene replacement in the
filamentous fungus Aspergillus nidulans. *Nucleic Acids Res.* **28**, E97 (2000).

64. Martin, P. *et al.* Interplay between Siderophores and Colibactin Genotoxin Biosynthetic Pathways in
Escherichia coli. *PLoS Pathogens* vol. 9 e1003437 (2013).

65. Davidson-Pilon, C. *et al. CamDavidsonPilon/lifelines: v0.21.0.* (2019). doi:10.5281/zenodo.2638135.

66. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069 (2014).

67. Treangen, T. J., Ondov, B. D., Koren, S. & Phillippy, A. M. The Harvest suite for rapid core-genome
alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol.* **15**, 524
(2014).

68. Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of recombinant bacterial whole
genome sequences using Gubbins. *Nucleic Acids Res.* **43**, e15 (2015).

69. Page, A. J. *et al.* Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* **31**,
3691–3693 (2015).

70. Lippert, C. *et al.* FaST linear mixed models for genome-wide association studies. *Nature Methods*
vol. 8 833–835 (2011).

71. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*
[q-bio.GN] (2013).

72. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841–842 (2010).

73. Touchon, M. *et al.* Organised genome dynamics in the Escherichia coli species results in highly
diverse adaptive paths. *PLoS Genet.* **5**, e1000344 (2009).

74. UniProt Consortium. UniProt: a hub for protein information. *Nucleic Acids Res.* **43**, D204–12 (2015).

75. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool.
Journal of molecular biology vol. 215 403–410 (1990).

76. Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J. Antimicrob. Chemother.*
67, 2640–2644 (2012).

77. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830
(2011).

78. Van Der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy array: a structure for efficient numerical
computation. *Comput. Sci. Eng.* **13**, 22–30 (2011).

79. Jones, E., Oliphant, T. & Peterson, P. SciPy: Open source scientific tools for Python. *http://www.*
scipy. org/ (2001).

80. Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational molecular biology and
bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

81. Talevich, E., Invergo, B. M., Cock, P. J. & Chapman, B. a. Bio.Phylo: A unified toolkit for processing,
analyzing and visualizing phylogenetic trees in Biopython. *BMC Bioinformatics* vol. 13 209 (2012).

82. McKinney, W. & Others. Data structures for statistical computing in Python. in *Proceedings of the 9th*

603   *Python in Science Conference* vol. 445 51–56 (2010).

603 83. Dale, R. K., Pedersen, B. S. & Quinlan, A. R. Pybedtools: a flexible Python library for manipulating genomic datasets and annotations. *Bioinformatics* **27**, 3423–3424 (2011).

606 84. Sukumaran, J. & Holder, M. T. DendroPy: a Python library for phylogenetic computing. *Bioinformatics* **26**, 1569–1571 (2010).

608 85. Huerta-Cepas, J., Serra, F. & Bork, P. ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic Data. *Mol. Biol. Evol.* **33**, 1635–1638 (2016).

610 86. Seabold, S. & Perktold, J. Statsmodels: Econometric and statistical modeling with python. in *Proceedings of the 9th Python in Science Conference* vol. 57 61 (SciPy society Austin, 2010).

612 87. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science Engineering* **9**, 90–95 (2007).

614 88. Waskom, M. *et al. mwaskom/seaborn: v0.9.0 (July 2018)*. (2018). doi:10.5281/zenodo.1313201.

615 89. Kluyver, T. *et al.* Jupyter Notebooks-a publishing format for reproducible computational workflows. in *ELPUB* 87–90 (2016).

617 90. Köster, J. & Rahmann, S. Snakemake-a scalable bioinformatics workflow engine. *Bioinformatics* **34**, 3600 (2018).

Supplementary Figure 1. Simulations of statistical power on an unrelated set of
complete *E. coli* genomes, using the *pks2* gene as target. The dotted red line indicates
the sample size used in the actual analysis.

Supplementary Figure 2. HPI structure conservation across strains. One strain per
phylogroup or species is shown, using the same color scheme as Figure 1E for each
gene.

Supplementary Figure 3. Presence/absence patterns of known virulence factors. Solid color indicates presence, light grey indicates absence. Phenotypes (number of killed mice) and phylogroup or species of each strain are reported as in Figure 1A. "Other virulence factors" are (from inside the ring towards the outside): *sfaD*, *sfaE*, *ompT*, *traT*, *hra2*, *papC*, *iha*, *ireA*, *neuC*, *hlyC*, *clbQ* and *cnf1*.

Supplementary Information

Supplementary Table 1: Strains' information, including virulence phenotype

Supplementary Table 2: Summary of the 81 genes with at least one mapped unitig

Supplementary Table 3: Survival analysis for NILS9 and NILS46 wild-type and HPI mutants

Supplementary Table 4: Correlation between growth on stress conditions (s-scores) and both virulence and presence of the HPI

Supplementary Table 5: Feature importance for each growth condition in the random forests predictor for virulence and HPI presence

Supplementary Table 6: List of PCR primers used in this study