Diel dynamics of dissolved organic matter and heterotrophic prokaryotes reveal enhanced growth at the ocean's mesopelagic fish layer during daytime

Xosé Anxelu G. Morán[1*], Francisca C. García[1,2], Anders Røstad[1], Luis Silva[1], Najwa Al-Otaibi[1], Xabier Irigoien[3], Maria L. Calleja[1, 4]

[1]King Abdullah University of Science and Technology (KAUST), Red Sea Research Center, Biological and Environmental Science & Engineering Division, 23955-6900 Thuwal, Saudi Arabia

[2]Environment and Sustainability Institute, University of Exeter, TR10 9FE Penryn, United Kingdom

[3]AZTI Tecnalia, 20110 Pasaia, Spain

[4]Max Planck Institute for Chemistry, 55128 Mainz, Germany

*Corresponding Author. Email: xelu.moran@kaust.edu.sa
Present address: Instituto Español de Oceanografía, Centro Oceanográfico de Gijón/Xixón, 33212 Gijón/Xixón, Spain

ABSTRACT

Contrary to epipelagic waters, where biogeochemical processes closely follow the light and dark periods, little is known about diel cycles in the ocean's mesopelagic realm. Here, we monitored the dynamics of dissolved organic matter (DOM) and planktonic heterotrophic prokaryotes every 2 h for one day at 0 and 550 m (a depth occupied by vertically migrating fish during light hours) in oligotrophic waters of the central Red Sea. We additionally performed predator-free seawater incubations of samples collected from the same site both at midnight and at noon. Comparable in situ variability in microbial biomass and dissolved organic carbon concentration suggests a diel supply of fresh DOM in both layers. The presence of fish in the mesopelagic zone during daytime promoted a sustained, longer growth of larger prokaryotic cells. The specific growth rates were consistently higher in the noon experiments from both depths (surface: 0.34 vs. 0.18 d^{-1}, mesopelagic: 0.16 vs. 0.09 d^{-1}). Heterotrophic bacteria and archaea in the mesopelagic fish layer were also more efficient at converting DOM into new biomass. These results suggest that the ocean's twilight zone receives a consistent diurnal supply of labile DOM from diel vertical migrating fishes, enabling an unexpectedly active community of heterotrophic prokaryotes.

Planktonic heterotrophic prokaryotes (HP) pertaining to the domains Bacteria and

Archaea rely on labile dissolved organic matter (DOM) for metabolism and growth [1, 2, 3]. In

surface waters, diel cycles in HP biomass and activity have been related to the photosynthetic

activity of phytoplankton [4], which obviously follows sunlight. Heterotrophic prokaryotes

dependence on DOM derived from planktonic algae [5] was reported to increase offshore, far

from coastal inputs, in temperate and polar ecosystems [6]. Although this relationship, also

known as bacterioplankton-phytoplankton coupling, has been the subject of debate [7,8], in

regions with low DOM advection (e.g. at permanently stratified sites without anthropogenic or

riverine inputs nearby, such as oligotrophic tropical waters), we might expect strong diel signals

in the response of heterotrophic prokaryotes coupled with the activity of primary producers [9].

In this regard, the Red Sea offers a unique opportunity to study biogeochemical processes in

oligotrophic ecosystems. With no permanent rivers, the only allochthonous inputs of DOM

come from urban centers such as Suez, Ghardaqa, Jeddah or Port Sudan, coastal macrophytes

[10, 11] or dust events [12, 13].

While epipelagic processes driven by primary production are well known [14, 15], large gaps

in our understanding of the ecology and biogeochemistry of the mesopelagic zone (i.e. waters

between 200 m and 1000 m) remain [16]. In the mesopelagic realm, trophic interactions between

microbes and metazoa have been long neglected. The available studies have focused mostly on

mesozooplankton[17, 18, 19]. However, recent reports on the large biomass contributed to the

ocean's biota by mesopelagic fishes performing diel vertical migration (DVM) suggest they may

also play an important role as rapid vectors of labile organic mater[20, 21]. DVM can affect only a

fraction of individuals from a given population [20]. In the Red Sea, virtually the entire

populations of mesopelagic fish migrate daily between the surface and the so-called deep

scattering layer (DSL) located at 400-650 m in the mesopelagic zone [22, 23]. DVM fishes have been

recently suggested to generate hotspots for heterotrophic prokaryotes, yielding significantly

higher bacterial growth efficiencies compared with shallower layers [24]. An analysis of a 24 h

intensive sampling at the same location has supported the existence of diel inputs of labile

DOM fueling the HP community at the depths occupied by mesopelagic fishes during daytime

[25]. Both DOC concentrations and high nucleic acid content (HNA) bacteria and archaea, usually

made up of copiotrophic taxa [26, 27] and more active than the low nucleic acid content (LNA)

group[28, 29, 30], fluctuated as widely in waters below 200 m as in the upper layers. However, for

the hypothesis of the mesopelagic labile DOM hotspots to be true, we should be able to

demonstrate that the presence or absence of fishes in the twilight zone does make a difference.

Here, we report on the results of two short-term incubations with water collected from

the epi- and mesopelagic layers (surface and 550 m, respectively) of the central Red Sea at

midnight and at the following midday. After removing protistan grazers and other larger

organisms by filtration, we followed the dynamics of DOM-heterotrophic prokaryotes

interactions for 8 days. In parallel, we conducted a high frequency (every 2 h for a full 24 h

starting at noon) characterization of the same depths focusing on the response of

heterotrophic prokaryotes abundance, cell size and biomass to changes in DOM concentrations

including its fluorescent properties, previously unreported for this basin. The specific objectives

of this study were: i) to assess the diurnal scales of variability in the standing stocks of HP and

DOM in epipelagic and mesopelagic waters of the central Red Sea, and ii) to test for differences

in the specific growth rate, maximum biomass and growth efficiency of HP between nighttime

and daytime in both layers. Our hypothesis is that DOM supplied by DVM fishes in the

mesopelagic zone during the day had a commesurable effect on the above-mentioned

variables.

Fig. 1. Echogram from March 5th to 6th 2016 at the study site showing 2 scattering layers of mesopelagic fish
performing diel vertical migration: up to the surface at night and down to deep waters during daytime. Squares
indicate the depth and time of water collection for the incubation experiments. Surface and mesopelagic (Fish)
depths are represented in green and blue, respectively, for coherence with subsequent figures. Colour scale
indicates backscattering strength (Sv, dB).

Results

Environmental variability of DOC and heterotrophic prokaryotes

The complete diel vertical migration of the mesopelagic fishes present at the study site

can be clearly seen in the echogram of **Fig. 1**, with the deeper, more intense layer (dominated

by *Benthosema pterotum*) occupying the depths between ca. 520 and 630 m during daytime on

March 6th 2016. **Fig. 2** shows the diel variability of DOC concentrations and the biomass of HP at

the station's surface and 550 m depth. Mean DOC values were almost 50% higher at 0 than at

550 m (71.0 ± 1.6 SE and 45.6 ± 1.5 µmol C L^{-1}, respectively). However, both depths showed

similar dynamics, with relative maxima of DOC at noon and also at midnight at the surface (**Fig.**

2A) and a slightly higher CV in the fish layer (8.1% vs. 5.7%). The apparent hourly rates of DOC

production (from 8 am to 12 pm at both depths) and consumption (from 12 am to 8 am at the

surface and from 12 pm to 4 pm at 550 m) were similar within each layer: ca. 1.3 µmol C L^{-1} h^{-1}

at the surface and 2.0 µmol C L^{-1} h^{-1} at the fish layer. Regarding the fluorescent DOM fraction,

the protein (Tyrosine)-like component C4 was on average one order of magnitude higher at the

surface than at 550 m, although it showed more variability at depth (**Table S1**).

Fig. 2. Variability of mean DOC concentration (**A**) and heterotrophic prokaryoplankton biomass (**B**) and cell size (**C**) in two layers of the study site: upper (0-25 m) and mesopelagic occupied by fish during daytime (450-600 m) during the 24 h sampling. Surface and mesopelagic (Fish) depths are represented in green and blue, respectively. Squares indicate initial values at the onset of the experimental incubations. The gray area represents nightime hours at the date of sampling. Error bars represent the standard error of the mean (average of 0 and 25 m in the upper layer and 450, 550 and 600 m in the mesopelagic one).

The abundance of HP at the surface (mean $4.31 \pm 0.17 \times 10^5$ cells mL^{-1}) was also one

order of magnitude higher than at 550 m (mean $9.43 \pm 0.39 \times 10^4$ cells mL^{-1}), but varied similarly

with no clear diel patterns. Although their size was 22% ± 8% larger in the mesopelagic (mean

values of 0.034 and 0.028 μm^3 at the fish and surface layers, respectively, **Fig. 2C**), the

corresponding biomass was driven mostly by changes in abundance, averaging 1.91 ± 0.09 µg C

L^{-1} at the surface and 0.50 ± 0.02 µg C L^{-1} at 550 m. **Fig. 2B** (in µmol C L^{-1} for comparison with

Fig. 2C) shows that HP biomass was equally variable at both layers (CV 16.0 vs. 16.3%). In situ

apparent or net growth rates based on changes in HP cell size were of 0.15 and 0.10 d^{-1} at 0 and

550 m, respectively. These estimated specific growth rates changed cyclically over the 24 h

cycle, especially at the surface. Two maxima, at 20:00 and 8:00, were found at the surface layer

while the maximum at 550 m was observed at 16:00 (**Fig. S2**).

Fig. 3. Dynamics of heterotrophic prokaryoplankton biomass (light colour) and DOC concentration (dark colour) in
the predator-free experimental incubations of samples taken at noon (**A, C**) and at midnight (**B, D**) from the surface
and the mesopelagic (Fish, 550 m depth) layers. Note the different scales for surface and mesopelagic water
experiments. Error bars are standard errors of 3 replicates.

Table 1. Mean ± SE values of specific growth rates (µ), DOC consumption rates or prokaryotic carbon demand
(PCD, see the text), prokaryotic heterotrophic production rates (PHP) and prokaryotic growth efficiency (PGE) in
the surface and fish layer incubation experiments performed at noon (Day) and midnight (Night). Rates were
calculated for each period of exponential growth, also indicated in days. The same period was used for DOC
consumption and biomass production rates. Also indicated are the maximum heterotrophic prokaryotes biomass
reached within the incubation and the corresponding ratio of maximum to initial biomass (Max:t0 biomass ratio).

Layer	Time	Period (d)	μ (d^{-1})	DOC consumption rate (μmol C L^{-1} d^{-1})	PHP rate (μmol C L^{-1} d^{-1})	PGE (%)	Maximum HP biomass (μg C L^{-1})	Max:t0 HP biomass ratio
Surface	Day	0-1.75	0.34 ± 0.07	1.0 ± 0.5	0.040 ± 0.004	4.2	2.69 ± 0.20	2.30
	Night[a]	0-2.25	0.18 ± 0.02	2.7 ± 1.1	0.047 ± 0.006	1.8	2.85 ± 0.16	1.80
Fish	Day[a]	0-2.75	0.16 ± 0.04	0.5 ± 0.3	0.020 ± 0.004	4.2	0.94 ± 0.13	3.27
	Night	0-2.25	0.09 ± 0.01	0.3 ± 0.8	0.010 ± 0.001	3.1	0.68 ± 0.02	2.31

[a]Vertically migrating mesopelagic fish present

Experimental incubations of surface and deep samples

With initial concentrations similar to ambient values (**Fig. 2**), DOC was consumed in the

first 2-3 days in the predator-free experiments (**Fig. 3**), albeit at different daily rates (**Table 1**),

followed by net production after day 4 especially in the Surface incubations. Minimum and

maximum consumption rates were 0.32 and 2.69 μmol C L^{-1} d^{-1}, found in the Fish and Surface

Night experiments, respectively (FN and SN). Values in the other two experiments carried out

with noon samples were below 1 μmol C L^{-1} d^{-1} (0.47 and 0.95 μmol C L^{-1} d^{-1}, respectively in in

FS and SD). The initial fluorescence intensity values of the component C4 in the experiments

was higher but reflected the values measured concurrently in the water column (paired *t*-test,

p>0.05, n=4). C4 showed a very consistent consumption pattern regardless of the layer and

treatment. Heterotrophic prokaryotes in the Surface incubations consumed in 4 days 40 and

50% of the initial values during Night and Day respectively, while bacteria inhabiting deep

waters consumed almost all of it (95-100 %) within the same time frame regardless of the

sampling time (**Fig. 4**). Hereinafter we consider changes in C4 as representative of labile DOM

dynamics. Day and night C4 consumption patterns did not show any significant differences

within the same layer but they displayed significantly higher consumption rates in the Fish layer

(ANOVA, p=0.004, post-hoc Fisher LSD test, **Figure 4** and **Table S1**).

Fig. 4. Dynamics of the concentration of the FDOM protein-like C4 component in the predator-free experimental
incubations of samples taken at noon (**A, C**) and at midnight (**B, D**) from the surface and 550 m depth. Error bars
are standard errors of 3 replicates.
169

Heterotrophic prokaryotes responses in the incubation experiments differed between

depths and sampling times (**Fig. 3**). Consistent differences were found between the specific

growth rates at both depths (**Table 1**), with μ being double in the Surface than in the Fish

experiments. Within each layer, the Day μ values were also higher than the Night ones (t-tests,

p=0.020 and p=0.060 at the Surface and Fish experiment, respectively, n=6). HNA cells always

grew faster than their LNA counterparts resulting in increases in their relative contribution from

43-55% to 55-62%, more noticeable in the FD experiment. The mean size of the cells also

increased substantially in the Fish experiments, from 0.027 to 0.060 μm^3 in the FD incubation

and from 0.028 to 0.047 μm^3 in the FN one, while changes in cell size were much smaller in the

Surface experiments, and virtually the same in both periods, from 0.026 to 0.037 μm^3 (SD) and

from 0.025 to 0.035 μm^3 (SN) (**Fig. S2**). Consequently, cell size played an important role in the

increase in biomass, especially in both Fish experiments (**Fig. 3C, D**). The biomass production

rates of heterotrophic prokaryotes for the same periods of DOC consumption ranged 4-fold,

from 0.010 to 0.047 µmol C L^{-1} d^{-1}, mirroring the changes in the latter variable (**Table 1**). The

rates of heterotrophic prokaryotes biomass production and DOC consumption were used for

estimating prokaryotic growth efficiencies (PGE) in the four experimental incubations. PGE was

uniformly below 5%, ranging from 1.8% (SN) to 4.2% (both SD and FD, **Table 1**). Following the

pattern of in situ values, maximum HP biomass measured in the incubations was higher in the

Surface than in the Fish experiments (**Fig. 3**, **Table 1**), although the increase ratios (i.e. the ratio

of maximum to initial biomass, **Table 1**) were significantly higher in the Fish experiments with

all data pooled (t-test, p=0.048 , n=12).

Discussion

Diel interactions between DOM and heterotrophic prokaryotes

There is consensus that marine biota biomass and activity peak in the upper layers and

decrease exponentially with depth, following the strong vertical gradients in physico-chemical

properties [31]. Heterotrophic prokaryotes inhabiting the Red Sea seem to challenge this view.

Together with diel variations of standing stocks [25], this study), night- and day-initiated

incubations of predator-free ambient assemblages with the DOM pool available at the time of

sampling yielded surprising similarities in epipelagic and mesopelagic waters. A method for

estimating division rates in cyanobacteria based in changes in cell size [32, 33] was adapted to

obtain independent estimates of in situ net growth rates (**Fig. S2**). The diel variability in

heterotrophic prokaryotes cell size showed the same pattern at the surface and the

mesopelagic fish layer (**Fig. 2**), yielding low but comparable net growth estimates at the surface

than at 550 m depth (0.10-0.15 d^{-1}). However, it must be noted here that the already large

prokaryotes inhabiting the mesopelagic fish layer were able to grow much bigger in the absence

of protistan grazers (**Fig. S1**), confirming the results of a previous study conducted at the same

site [24].

Diel cycles in biogeochemical properties and plankton biomass and activity in the upper

ocean layers are reasonably well known, following diurnal changes in photosynthesis and food

web processes [4, 9]. No clear patterns were observed for in situ heterotrophic prokaryotes

abundance or biomass, probably due to strong coupling between growth and mortality due to

protistan grazing [34, 35] and/or viral lysis in the Red Sea (E. Sabbagh et al., in prep.), and in

tropical waters in general [36]. However, likewise early observations in the Caribbean [37] recently

confirmed for this site [25], DOM concentrations displayed a coherent diel pattern suggesting

different timing of production and consumption (**Fig. 1**). Confirming previous experiments [38],

DOC consistently decreased in the first 2-3 days in both Surface and Fish samples, ranging from

0.7 to 6.1 μmol L^{-1}, followed by net production after day 4 especially in the Surface incubations,

coincident with a sharp decay in prokaryotic biomass (**Fig. 3A, B**). Indeed, the strong apparent

uptake of surface DOC in situ during nighttime coincided with the highest DOC consumption

experimental result, poiting to its labile nature, although the buildup of HP biomass was similar

in both SD and SN experiments, resulting in lower specific growth rates and PGE values in the

Night (**Table 1**). Labile DOC incorporation does not automatically inform us of its subsequent

partitioning between metabolism (respiration) and growth (biomass production), as shown by

Condon et al. [39] for DOM originated by jellyfish blooms. Since the alternation between light and

dark periods within the incubator used for Surface seawater was common for SD and SN

experiments, if photoheterotrophy [40, 41] afffected DOM dynamics, it should have been equally

apparent in both Night and Day incubations, which were initiated with only 12 h difference (**Fig.

1**). We therefore ruled out photoheterotrophic processes explaining the differences observed

after 8 days of incubation. Rather, the quality of labile DOM at midday (including recent

photosynthate) was probably higher than at midnight (perhaps with a greater contribution of

DOM coming from sloppy feeding, [42], causing the significantly higher μ values (**Table 1**). That

the quality of labile DOC at noon could have been higher was supported by a faster increase in

bacterial cell size (**Fig. S1B**) and the contribution of HNA cells, which increased by 35% in SD

compared with 15% in SN. The decrease in the protein-like C4 component was also more

sustained in the SD experiment compared to SN after 4 days (13.2% vs. 5.3% was consumed

daily during that period, **Fig. 4**) although the rates were virtually the same in the initial periods

(2.75 and 2.25 days, respectively, **Table S1**). Changes in mesopelagic DOC concentrations and

lability at diel ([25], this study) and seasonal scales [43] in the central Red Sea support the recent

claim that the diverse pool of DOM in the deep ocean fluctuates at timescales much shorter

than previously thought [44]. Since the conditions at the study site were hypoxic in most of the

mesopelagic realm [43], the low in situ oxygen concentrations in the Fish water samples,

consistent from 300 m downwards (0.69 ± 0.03 mg L^{-1}) might have been supplemented by pre-

filtration and sampling from the experimental bottles. Although we did not control for this

potential artefact, the same protocol was followed for the FN and FD experiments. Therefore,

the consistently higher values of DOM consumption, prokaryotic cell size, growth rates and

efficiency in the Day compared with the Night incubation strongly support that the presence of

fish indeed had a major impact on the microbial community.

Surface specific growth rate in the Day experiment was nevertheless notably higher

than the 0.08 d^{-1} measured in a previous study carried out in November 2015 at the same

location [24]. The discrepancy cannot be explained by total DOC or chlorophyll *a* concentrations,

but could instead be related to the availability of labile DOM compounds since C4

concentrations were 61% higher in March than in November (M. L. Calleja, pers. comm.). At a

shallower, nearby site characterized by higher total and labile DOC concentrations, specific

growth rates were still considerably higher, ranging from 0.79 to 1.75 d^{-1} [35].

The daily rates of apparent DOC production and consumption based on changes in in

situ concentration were virtually the same within each of the two layers compared, indicating

no net accumulation. This is the expected result in oligotrophic regions at the short time scale

of one day [37, 45]. However, these rates were still ca. 50% higher in the mesopelagic zone

resulting in apparent turnover of labile DOC of 23.6% d^{-1} in the mesopelagic compared with

14.7% d^{-1} at the surface, if we consider the measured diel variability and maximum

concentrations in each layer. This finding conflicts with the contention that DOM is largely of

refractory nature within the mesopelagic waters of the global ocean [46]. The role of vertically

migrating animals, zooplankton and fishes, as vectors of organic matter to deep layers

complementary to the biological pump [14] has been recently recognized [17, 18, 47]). In this regard,

work at the study site has suggested DVM fishes as a transport mechanism supplying labile

DOM that does not accumulate but fuels heterotrohic bacterial activity in mesopelagic waters

[24]. Here we tested this hypothesis by further examining the fluorescence properties of DOM

and its transformation by heterotrophic prokaryotes in experimental incubations with and

without the fishes present. FDOM are useful tracers for biogeochemical processes in the dark

ocean [48, 49]. Fluorescence intensity of the two aminoacid-like fluorophores C3 and C4 decreased

with depth (data not shown), indicating that these fluorophores were mainly produced

autochthonously in surface waters. Both phytoplankton and bacteria are sources of tryptophan

and tyrosine [50], while Urban-Rich et al. [51] have reported that grazing and excretion by

zooplankton can also release material with amino acid-like fluorescence signals. Our results

strongly suggest that DVM fishes can also provide C4 in the mesopelagic realm.

Contrary to the epipelagic zone, very few studies on the diel variability of DOM-

heterotrophic prokaryotes interactions are available for deep waters [25, 52]. Gasol et al. [53]

suggested that mesopelagic prokaryotes in the subtropical NE Atlantic were as active as in the

epipelagic. We demonstrate here not only that heterotrophic prokaryotes specific growth rates

at 550 m were of the same order of magnitude than in surface waters, clearly challenging the

most accepted view [31, 54], but that those rates were almost double at noon conditions, when

the mesopelagic fishes were present, than at midnight, when the entire population was closer

to the surface [22]. From **Fig. 1** it is clear that the fishes were absent at midnight in the entire

mesopelagic zone but their presence at 550 m had been established for ca. 4 h when the noon

sampling took place. Specific growth rates were nevertheless lower than in the previous study

($0.24\ d^{-1}$, [24]). Although seasonality of C4 in the deep scattering layer was less marked than at

the surface, November 2015 was characterized by 82% higher C4 concentrations than in March

2016 (M. Calleja, pers. comm.). Altogether, these results point out to a major role of protein-

like substances in determining the specific growth rates of heterotrophic prokaryotes

throughout the water column, as recently found for nearby shallow waters in a seasonal study

[35]. C4 fluctuated widely in the 24 h monitoring at 550 m depth (**Table S1**) and was also actively

consumed in all our incubations, thus revealing a clearly labile nature. C4 was consumed faster

in the Fish experiments, at 42.1% and 25.8% d^{-1} in FD and FN, respectively (**Fig. 4, Table S1**),

than in the Surface ones (ca. 13% d^{-1} in both SD and SN). A similar relative consumption of

protein-like FDOM (12% d^{-1}), mostly occurring during the first 5 days, was measured by [55] in

experiments conducted with marine surface waters. Our explanation is that fishes released

DOM directly or it leaked from particles associated to the fish presence (e.g. fecal pellets). That

DOM could have been delivered by sinking particles [56] not related to vertical migration would

not explain the difference between the FD and FN experiments.

Cell size has been used as an indicator of the activity of heterotrophic prokaryotes [57].

While in the Surface experiment growing HP cells were only slightly larger than at time 0 (11%

larger size for both the SD and SN experiments), the cell size increase in the Fish experiment

was dramatic, especially in the Day incubations (118% vs. 68% larger, **Fig. S2**). The contribution

of bigger cells to the observed increase in HP biomass is not at all minor: had we used, as in

many studies, a fixed cellular carbon content of 4 fg C $cell^{-1}$ (corresponding to the initial mean

cell size of 0.027 μm^3 for the two depths and periods, **Fig. 2C**), maximum HP biomass in **Table 1**

would have become 2.10 (SD), 2.32, (SN), 0.43 (FD) and 0.39 (FN) μg C L^{-1}, i.e. between 42 and

54% lower than the actual values for the mesopelagic prokaryotes. We conclude that the

presence of fishes in the mesopelagic zone during daytime resulted in significantly higher

growth rates of markedly larger cells. As a consequence, changes in abundance were

exacerbated when considering biomass units. The maximum biomass of heterotrophic

prokaryotes that could be sustained by extant DOM concentrations was significantly higher

than the initial value in both Fish experiments (**Table 1**). Altogether, these results point out to

substantial inputs of labile DOM during daytime at the mesopelagic fish layer that are rapidly

mobilized by large bacterial taxa. The archaeon *Nitrosopulimus maritimus,* which makes up

much of the heterotrophic prokaryoplankton biomass at these depths [58], were apparently not

the main responders to these DOM hotspots, since their contribution to total numbers at the

end of a similar incubation dropped from 50% to 3% [24]. The typical size of these

Thaumarchaeota is small [59], so it is unlikely that they were the dominant groups growing in our

Fish incubations after 2 days (**Fig. S2C, D**). Besides excreting ammonia that boosts its oxidation

by Thaumarchaeota (formely Chrenarchaeota) [60], mesopelagic fish thus seem capable to fuel

the metabolism of large, copiotrophic bacteria.

Prokaryotic growth efficiencies are typically low in open ocean, oligotrophic

environments [61, 62]. Most of these measurements refer to the epipelagic zone, where

photosynthesis takes place. Here we provide more information from deeper layers, where

export processes are more important. The recently reported low PGE values at this Red Sea site

(1.6-3.4%, [24] are confirmed by this new study, while higher values (2.5-12.8%) were recorded in

a shallow, richer bay located a few km south [35]. Few studies have estimated the vertical

variability in PGE values, but those that have usually depict lower values with depth [62, 63],

related to the increased presence of refractory DOM compounds [46] or to the higher dilution of

the labile ones [64]. Notably, the estimated growth efficiency of heterotrophic prokaryotes in our

Day experiments was exactly the same in Surface and Fish water (4.2%), which can only be

explained by the existence of labile DOC of similar quality within both layers, resulting from

photosynthesis at the surface and fish-mediated export in the mesopelagic layer. PGE in the

mesopelagic fish layer was significantly higher than at shallower depths in the experiment

conducted at noon in November 2015 [24]. However, when averaging our new two estimates, the

mean PGE value at 550 m (3.6%) was still 22% higher than at the surface, strongly supporting

the presence of high quality DOM hotspots [24] in the deep scattering layer.

Although this study is limited to a single 24 hour period, it builds on the previously

demonstrated relatively high growth of mesopelagic heterotrophic prokaryotes [24] by comparing

the outcome of deep samples incubations taken only 12 hours apart, at midnight and noon. We

confirm that the Red Sea mesopelagic zone is not a permanently impoverished environment

but subject to daily inputs of labile DOM compounds similarly to the epipelagic layers. This

novel process driven by mesopelagic fishes, which complements other recently discovered

sources of deep organic carbon [14, 65, 66, 67] seems to have been overlooked due to the tight

coupling between the components of microbial food webs [68]. If vertically migrating fishes are

able to fuel an active and distinct community (T. Huete-Stauffer et al., submitted) of

heterotrophic prokaryotes in the mesopelagic layer of the Red Sea, we might expect this fast

DOM flux to be widespread. The mesopelagic Red Sea has an unusually high temperature,

therefore the effect of colder conditions on fish DOM-microbial interactions remain to be

explored. The implications for global biogeochemical cycling would also vary depending on the

actual biomass of mesopelagic fishes and the fraction performing DVM [20], yet its impact may

increase as deep waters warm up [69]. That these small fishes seem able to sustain the microbial

communities inhabiting the twilight zone also may help reconcile current discrepancies

between carbon pools and fluxes in the global ocean.

357
METHODS

Environmental sampling

We occupied one station located 13.4 km offshore to the north of King Abdullah

Economic City, Saudi Arabia (lat 22.46°N, lon 39.02°E) between midday March 6th and midday

March 7th, 2016 [24, 25]. In situ monitoring and sampling was conducted on board of RV Thuwal.

Continuous acoustic measurements in order to locate the position of the vertically migrating

mesopelagic fishes were recorded with a Simrad EK60 38 kHz echosounder mounted on the

ship's hull. From noon on March 6th until the same time on the following day we conducted CTD

casts every 2 hours. At each cast we sampled discrete depths in the water column with Niskin

bottles mounted on a Rosette sampler, ranging from the surface to 650 m depth. Water filtered

through pre-combusted Whatman GF/F filters was collected for analyzing DOC bulk

concentrations and fluorescent DOM (FDOM) properties (40 mL pre-combusted glass vials).

Unfiltered water was collected for characterizing the community of heterotrophic prokaryotes

(2 mL cryovials).

Hourly in situ apparent DOC production and consumption rates were estimated as the

largest difference between DOC concentration in consecutive sampling times that showed a

consistently increasing and decreasing trend, respectively. The same approach was used for

estimating the apparent biomass production of heterotrophic prokaryotes over the diel cycle.

Experimental incubations

10 L of seawater from the surface and 550 m depth were collected in the midnight and

noon casts on March 7th for conducting the experimental incubations of DOC consumption,

change in FDOM and heterotrophic prokaryotes biomass response. In order of remove

protistan grazers and planktonic organisms larger than bacteria and archaea, water was gently

filtered through pre-combusted Whatman GF/C filters (142 mm, nominal pore size 1.2 μm) and

used to fill 3 x 2 L acid-cleaned polycarbonate bottles, which were subsequently incubated at

the in situ temperature and light regimes (darkness for 550 m samples), so that the possible

role of photoheterotrophs in the processing of DOM was included. Removal of heterotrophic

prokaryotic cells by filtration was minor (83% ± 7% SE of the initial abundance was retrieved in

the water used for the incubations) and mean cell size was virtually unaffected (2.6% ± 1.0%

smaller biovolume than inthe unfiltered water). However, filtration eliminated most

Synechococcus and Prochlorococcus cyanobacteria as well as virtually all the larger protistan

grazers of heterotrophic prokaryotes, since the mean abundance of heterotrophic

nanoflagellates in the GF/C filtrate was 1.5% (E. I. Sabbagh, pers. comm.). We are therefore

confident that no extra sources of DOM were included in our experiments. Subsamples were

taken twice per day on the first 2 days, then daily until day 6 and finally at day 8. DOC and

FDOM subsamples from the incubations were gravity-filtered through pre-rinsed 0.2 Millipore

polycarbonate filters. We will occasionally use the codes S and F to refer to the incubations

made with water from the Surface (0 m) and the Fish layer (550 m), respectively, followed by D

or N to refer to the period of sampling (Day or Night): SD, SN, FD, FN.

DOC analysis

Samples for DOC were acidified with H_3PO_4 and kept in the dark at 4 °C until analysis by

high temperature catalytic oxidation at the laboratory. All glass material used was acid cleaned

and burned (450°C, 4.5 h). Consensus reference material of deep sea carbon (42–45 μmol C L^{-1}

and 31-33 μmol N L^{-1}) and low carbon water (1-2 μmol C L^{-1}), provided by D. A. Hansell and W.

Chen (Univ. of Miami) was used to monitor the accuracy of our DOC concentration

measurements. The analytical error of DOC concentration was 1.4 μmol L^{-1}.

DOM fluorescence measurements and PARAFAC modeling

FDOM samples were stored at 4°C before being analyzed (within 2 days after the

completion of the cruise and incubation sample collection). UV-VIS fluorescence spectroscopy

was measured using a HORIBA Jobin Yvon AquaLog spectrofluorometer with a 1 cm path length

quartz cuvette. Three dimensional fluorescence excitation emission matrices (EEMs) were

recorded by scanning with an excitation wavelength range of 240- 600 nm and emission of 250-

600 nm, both at 3 nm increments and integrating at 8 seconds. To correct and calibrate the

fluorescence spectra post-processing steps were followed according to Murphy et al. [70]. Briefly,

fluorescence spectra were Raman area (RA) normalized by subtracting daily blanks that were

performed using Ultra-Pure Milli-Q sealed water (Certified Reference, Starna Cells). Inner-filter

correction (IFC) was also applied according to McKnight et al. (2001) RA normalization, blank

subtraction, IFC and generation of EEMs were performed using MATLAB (version R2015b).

A total of 165 samples for DOM fluorescence were collected (81 from 7 vertical profiles

and 84 from the experimental incubations). The EEMs obtained were subjected to PARAFAC

modeling using DOMFluor Toolbox [71]. Before the analysis, Rayleigh scatter bands were

trimmed. A four-component model was validated using split-half validation and random

initialization [71]: peak C1 at Ex/Em 240(325)/ 407 nm, peak C2 at Ex/Em 258(390)/492 nm, peak

C3 at Ex/Em 240/337 and peak C4 at Ex/Em 276/312 nm. C1 corresponds to peak M [72] and is

comparable to component 2 identified by Català et al. [49]. C2 represents a combination of peaks

A and C [72] and is comparable to component 1 in [49]. C3 corresponds to peak T [72], attributed to

tryptophane, and is comparable to component 3 in [49]. C4 corresponds to peak B [72], attributed

to tyrosine, and is comparable to component 4 in [49]. The maximum fluorescence (Fmax) is

reported in Raman units (RU).

Heterotrophic prokaryotes abundance and biomass

Triplicate samples (1.8 mL) for estimating the abundance of heterotrophic bacteria and

archaea in situ and in the experimental incubations were fixed with 1% paraformaldehyde and

0.05% glutaraldehyde, deep frozen in liquid nitrogen and stored at -80°C until analysis. Once

thawed, 400 µL aliquots were stained with SYBR-Green run in a BD FACSCanto II flow cytometer

for estimating the abundance of low (LNA) and high (HNA) nucleic acid content cells as detailed

in Gasol and Morán [73]. The few cyanobacteria present were easily distinguished in the surface

samples due to their autofluorescence red signal because of the presence of chlorophyll *a*.

Absolute abundances were estimated based on time and the actual flow rates, which were

calibrated daily using the gravimetric method. The right angle light scatter or side scatter (SSC)

signal relative to the value of 1 μm fluorescent latex beads added to each sample was used to

estimate the cell diameter according to Calvo-Díaz and Morán [74]. LNA and HNA cell numbers

were summed to estimate the total abundance and their specific cell sizes averaged to obtain

the mean cell size of the heterotrophic prokaryote community at both depths and different

times. Assuming spherical shape, the mean cell size (biovolume in μm^3) was converted into

cellular carbon content following Gundersen et al. [75]. Heterotrophic prokaryotes biomass was

then calculated as the product of cell abundance and mean cellular carbon content.

Growth rate estimates

In situ apparent or net growth rates of the heterotrophic prokaryote assemblage at the

surface and the mesopelagic fish layer were estimated from changes in biomass ($\mu g\ C\ L^{-1}$)

resulting from changes in abundance and mean cell size over 24 h. Net growth rates (μ, in units

h^{-1}) were calculated as:

$\mu = \ln (N_1/N_0) / \Delta t$ (1)

where N_1 is the final biomass, N_0 is the initial biomass and Δt is the time interval (2 h).

We modeled the overall daily growth rate from Eq. (1) using the size distribution of the

organisms with the R package ssPopModel, which included a modified version of the size-

structured matrix population model originally developed by Sosik et al. [32]. Matrix population

model assumption is that changes in size distribution are only related to growth and division of

the cells. We adapted and simplified the application of this function as described by Hunter-

Cevera et al. [33] for *Synechococcus* cyanobacteria to be used with heterotrophic prokaryote cells.

Specific growth rates in the incubations were calculated as the slope of the ln-

transformed total abundance vs. time for the linear response period, equivalent to the phase of

exponential growth (usually lasting between 2 and 3 days).

Prokaryotic growth efficiency

Prokaryote heterotrophic production (PHP) in the midnight and midday incubations was
estimated as the rate of increase in bacterial biomass during the exponential phase of growth.
Prokaryotic carbon demand (PCD, i.e. the sum of heterotrophic prokaryotes production and
respiration) was approached by the consumption rate of DOC during the same period.
Prokaryotic growth efficiency (PGE) was therefore calculated as the ratio of PHP to PCD.

Statistical analyses

Model I or ordinary least squares (OLS) linear regressions for estimating specific growth
rates were done separately for each replicate, using a common period for each experiment.
Differences between treatments and/or depths were assessed with one way ANOVAs and
Fisher least significance (LSD) post-hoc tests. General relationships between variables were
represented by Pearson's correlation coefficients. Statistical analyses were done with JMP and
STATISTICA software packages.

ACKNOWLEDGMENTS

We are greatly indebted to the crew of RV Thuwal and the rest of the personnel from
the Coastal and Marine Resources (CMOR) Core Lab at KAUST for their assistance during field
work. Besides participating in the sample collection M. Viegas helped us with the rest of the
work in the Red Sea Research Center (RSRC) lab. We are also grateful to past and current
members of the Microbial Oceanography and Biogeochemistry lab at the RSRC.

X.A.G.M. conceived the research, led the experiment design, data analysis and wrote
the paper. F.C.G. modeled in situ growth rates. A.R. performed the acoustic research. L.S. and
N. A-O. analyzed the heterotrophic prokaryotes. X.I. contributed to the interpretation of results.
M.L.C. was responsible for DOC and FDOM measurements, contributed to experimental design
and data analysis. F.C.G., A.R., X.I. and M.L.C. also contributed to writing.

Competing interests

The authors declare that they have no financial or non-financial competing interests.

REFERENCES

1. Carlson CA, Ducklow HW, Michaels AF. Anual flux of dissolved organic carbon from the euphotic zone in the northwestern Sargasso Sea. *Nature* 1994, **371**: 405-408.

2. Pomeroy LR, Williams PJI, Azam F, Hobbie JE. The Microbial Loop. *Oceanography* 2007, **20**(2): 28-33.

3. Goldman JC, Dennett MR. Growth of marine bacteria in batch and continuous culture under carbon and nitrogen limitation. *Limnol Oceanogr* 2000, **45**(4): 789-800.

4. Gasol JM, Doval MD, Pinhassi J, Calderón-Paz JI, Guixa-Boixareu N, Vaqué D, *et al.* Diel variations in bacterial heterotrophic activity and growth in the northwestern Mediterranean Sea. *Mar Ecol Prog Ser* 1998, **164**: 107-124.

5. Baines SB, Pace ML. The production of dissolved organic matter by phytoplankton and its importance to bacteria: Patterns across marine and freshwater systems. *Limnol Oceanogr* 1991, **36 (6)**: 1078-1090.

6. Morán XAG, Gasol JM, Pedrós-Alió C, Estrada M. Dissolved and particulate primary production and bacterial production in offshore Antarctic waters during austral summer: coupled or uncoupled? *Marine Ecology-Progress Series* 2001, **222**: 25-39.

7. Fouilland E, Mostajir B. Revisited phytoplanktonic carbon dependency of heterotrophic bacteria in freshwaters, transitional, coastal and oceanic waters. *FEMS Microbiology Ecology* 2010, **73**(3): 419-429.

8. Morán XAG, Alonso-Sáez L. Independence of bacteria on phytoplankton? Insufficient support for Fouilland & Mostajir's (2010) suggested new concept. *Fems Microbiology Ecology* 2011, **78**(2): 203-205.

9. Ruiz-González C, Lefort T, Massana R, Simó R, Gasol JM. Diel changes in bulk and single-cell bacterial heterotrophic activity in winter surface waters of the northwestern Mediterranean Sea. *Limnol Oceanogr* 2012, **57**(1): 29-42.

10. Duarte CM, Cebrián J. The fate of marine autotrophic production. *Limnol Oceanogr* 1996, **41**(8): 1758-1766.

11. Alongi DM, Mukhopadhyay SK. Contribution of mangroves to coastal carbon cycling in low latitude seas. *Agr Forest Meteorol* 2015, **213**: 266-272.

12. Lekunberri I, Lefort T, Romero E, Vazquez-Dominguez E, Romera-Castillo C, Marrase C, *et al.* Effects of a dust deposition event on coastal marine microbial abundance and

activity, bacterial community structure and ecosystem function. *J Plankton Res* 2010,
32(4): 381-396.

13.   Bao HY, Niggemann J, Luo L, Dittmar T, Kao SJ. Molecular composition and origin of
water-soluble organic matter in marine aerosols in the Pacific off China. *Atmos Environ*
2018, **191:** 27-35.

14.   Herndl GJ, Reinthaler T. Microbial control of the dark end of the biological pump. *Nat*
Geosci 2013, **6**(9): 718-724.

15.   Henson SA, Sanders R, Madsen E. Global patterns in efficiency of particulate organic
carbon export and transfer to the deep ocean. *Global Biogeochem Cy* 2012, **26**.

16.   Robinson C, Steinberg DK, Anderson TR, Aristegui J, Carlson CA, Frost JR*, et al.*
Mesopelagic zone ecology and biogeochemistry - a synthesis. *Deep-Sea Res Pt Ii* 2010,
57(16): 1504-1518.

17.   Bianchi D, Stock C, Galbraith ED, Sarmiento JL. Diel vertical migration: Ecological controls
and impacts on the biological pump in a one-dimensional ocean model. *Global*
Biogeochem Cy 2013, **27**(2): 478-491.

18.   Isla A, Scharek R, Latasa M. Zooplankton diel vertical migration and contribution to deep
active carbon flux in the NW Mediterranean. *Journal of Marine Systems* 2015, **143:** 86-
97.

19.   Al-Mutairi H, Landry MR. Active export of carbon and nitrogen at Station ALOHA by diel
migrant zooplankton. *Deep-Sea Res Pt Ii* 2001, **48**(8-9): 2083-2103.

20.   Klevjer TA, Irigoien X, Rostad A, Fraile-Nuez E, Benitez-Barrios VM, Kaartvedt S. Large
scale patterns in vertical distribution and behaviour of mesopelagic scattering layers. *Sci*
Rep-Uk 2016, **6**.

21.   Irigoien X, Klevjer TA, Rostad A, Martinez U, Boyra G, Acuña JL*, et al.* Large mesopelagic
fishes biomass and trophic efficiency in the open ocean. *Nat Commun* 2014, **5**.

22.   Klevjer TA, Torres DJ, Kaartvedt S. Distribution and diel vertical movements of
mesopelagic scattering layers in the Red Sea. *Mar Biol* 2012, **159**(8): 1833-1841.

23.   Røstad A, Kaartvedt S, Aksnes DL. Light comfort zones of mesopelagic acoustic
scattering layers in two contrasting optical environments. *Deep-Sea Research Part I-*
Oceanographic Research Papers 2016, **113:** 1-6.

24. Calleja ML, Ansari MI, Røstad A, Silva L, Kaartvedt S, Irigoien X, *et al.* The Mesopelagic
Scattering Layer: A Hotspot for Heterotrophic Prokaryotes in the Red Sea Twilight Zone.
Front Mar Sci 2018, **5**: 259.
587
588 25. García FC, Calleja ML, Al-Otaibi N, Røstad A, Morán XAG. Diel dynamics and coupling of
589 heterotrophic prokaryotes and dissolved organic matter in epipelagic and mesopelagic
590 waters of the central Red Sea. *Environ Microbiol* 2018, **20**(8): 2990-3000.
591
592 26. Vila-Costa M, Gasol JM, Sharma S, Moran MA. Community analysis of high- and low-
593 nucleic acid-containing bacteria in NW Mediterranean coastal waters using 16S rDNA
594 pyrosequencing. *Environ Microbiol* 2012, **14**(6): 1390-1402.
595
596 27. Schattenhofer M, Wulf J, Kostadinov I, Glockner FO, Zubkov MV, Fuchs BM. Phylogenetic
597 characterisation of picoplanktonic populations with high and low nucleic acid content in
598 the North Atlantic Ocean. *Systematic and Applied Microbiology* 2011, **34**(6): 470-475.
599
600 28. Gasol JM, Zweifel UL, Peters F, Fuhrman JA, Hagström Å. Significance of size and nucleic
601 acid content heterogeneity as measured by flow cytometry in natural planktonic
602 bacteria. *Appl Environm Microbiol* 1999, **65**: 4475-4483.
603
604 29. Bouvier T, del Giorgio PA, Gasol JM. A comparative study of the cytometric
605 characteristics of High and Low nucleic-acid bacterioplankton cells from different
606 aquatic ecosystems. *Environmental Microbiology* 2007, **9**(8): 2050-2066.
607
608 30. Morán XAG, Ducklow HW, Erickson M. Single-cell physiological structure and growth
609 rates of heterotrophic bacteria in a temperate estuary (Waquoit Bay, Massachusetts).
610 *Limnol Oceanogr* 2011, **2011**(56): 37-48.
611
612 31. Arístegui J, Gasol JM, Duarte CM, Herndl GJ. Microbial oceanography of the dark ocean's
613 pelagic realm. *Limnol Oceanogr* 2009, **54**(5): 1501-1529.
614
615 32. Sosik HM, Olson RJ, Neubert MG, Shalapyonok A, Solow AR. Growth rates of coastal
616 phytoplankton from time-series measurements with a submersible flow cytometer.
617 *Limnol Oceanogr* 2003, **48**(5): 1756-1765.
618
619 33. Hunter-Cevera KR, Neubert MG, Solow AR, Olson RJ, Shalapyonok A, Sosik HM. Diel size
620 distributions reveal seasonal growth dynamics of a coastal phytoplankter. *P Natl Acad*
621 *Sci USA* 2014, **111**(27): 9852-9857.
622
623 34. Calbet A, Agersted MD, Kaartvedt S, Mohl M, Moller EF, Enghoff-Poulsen S, *et al.*
624 Heterogeneous distribution of plankton within the mixed layer and its implications for
625 bloom formation in tropical seas. *Sci Rep-Uk* 2015, **5**.
626

627   35.   Silva L, Calleja ML, Huete-Stauffer TM, Ivetic S, Ansari MI, Viegas M, *et al.* Low
628   abundances but high growth rates of heterotrophic bacteria in the coastal Red Sea.
629   *Front Microbiol*, **submitted**.
630

631   36.   Morán XAG, Gasol JM, Pernice MC, Mangot JF, Massana R, Lara E, *et al.* Temperature
632   regulation of marine heterotrophic prokaryotes increases latitudinally as a breach
633   between bottom-up and top-down controls. *Global Change Biol* 2017, **23**(9): 3956-3964.
634

635   37.   Johnson KM, Burney CM, Sieburth JM. Enigmatic marine ecosystem metabolism
636   measured by direct diel Sigma-CO2 and O-2 flux in conjunction with DOC release and
637   uptake. *Mar Biol* 1981, **65**(1): 49-60.
638

639   38.   Calleja ML, Ansari MI, Rostad A, Silva L, Kaartvedt S, Irigoien X, *et al.* The Mesopelagic
640   Scattering Layer: A Hotspot for Heterotrophic Prokaryotes in the Red Sea Twilight Zone.
641   *Front Mar Sci* 2018, **5**.
642

643   39.   Condon RH, Steinberg DK, del Giorgio PA, Bouvier TC, Bronk DA, Graham WM, *et al.*
644   Jellyfish blooms result in a major microbial respiratory sink of carbon in marine systems.
645   *P Natl Acad Sci USA* 2011, **108**(25): 10225-10230.
646

647   40.   Béjà O, Aravind L, Koonin EV, Suzuki MT, Hadd A, Nguyen LP, *et al.* Bacterial rhodopsin:
648   evidence for a new type of phototrophy in the sea. *Science* 2000, **289**: 1902-1906.
649

650   41.   Ruiz-González C, Simó R, Sommaruga R, Gasol JM. Away from darkness: a review on the
651   effects of solar radiation on heterotrophic bacterioplankton activity. *Front Microbiol*
652   2013, **4**.
653

654   42.   Nagata T. Production mechanisms of dissolved organic matter. In: Kirchman DL (ed).
655   *Microbial ecology of the oceans*. Wiley-Liss: New York, 2000, pp 121-152.
656

657   43.   Calleja ML, Al-Otaibi N, Morán XAG. Dissolved organic carbon contribution to oxygen
658   respiration in the central Red Sea. *Sci Rep-Uk* 2019, **9**: 4690.
659

660   44.   Follett CL, Repeta DJ, Rothman DH, Xu L, Santinelli C. Hidden cycle of dissolved organic
661   carbon in the deep ocean. *P Natl Acad Sci USA* 2014, **111**(47): 16706-16711.
662

663   45.   Wright RT. Dynamics of pools of dissolved organic carbon. In: Hobbie JE, Williams PJlB
664   (eds). *Heterotrophic activity in the sea*. Plenum Press: New York, 1984, pp 121-155.
665

666   46.   Jiao N, Herndl GJ, Hansell DA, Benner R, Kattner G, Wilhelm SW, *et al.* Microbial
667   production of recalcitrant dissolved organic matter: long-term carbon storage in the
668   global ocean. *Nat Rev Microbiol* 2010, **8**(8): 593-599.
669

670 47. Bianchi D, Galbraith ED, Carozza DA, Mislan KAS, Stock CA. Intensification of open-ocean
671 oxygen depletion by vertically migrating animals. *Nat Geosci* 2013, **6**(7): 545-548.
672

673 48. Nelson NB, Siegel DA. The Global Distribution and Dynamics of Chromophoric Dissolved
674 Organic Matter. *Annual Review of Marine Science, Vol 5* 2013, **5**: 447-476.
675

676 49. Catalá TS, Reche I, Fuentes-Lema A, Romera-Castillo C, Nieto-Cid M, Ortega-Retuerta E,
677 *et al.* Turnover time of fluorescent dissolved organic matter in the dark global ocean.
678 *Nat Commun* 2015, **6**: 5986.
679

680 50. Determann S, Lobbes JM, Reuter R, Rullkotter J. Ultraviolet fluorescence excitation and
681 emission spectroscopy of marine algae and bacteria. *Mar Chem* 1998, **62**(1-2): 137-156.
682

683 51. Urban-Rich J, McCarty JT, Fernandez D, Acuna JL. Larvaceans and copepods excrete
684 fluorescent dissolved organic matter (FDOM). *J Exp Mar Biol Ecol* 2006, **332**(1): 96-105.
685

686 52. Carlucci AF, Craven DB, Robertson KJ, Henrichs SM. Microheterotrophic Utilization of
687 Dissolved Free Amino-Acids in Depth Profiles of Southern-California Borderland Basin
688 Waters. *Oceanol Acta* 1986, **9**(1): 89-96.
689

690 53. Gasol JM, Alonso-Saez L, Vaque D, Baltar F, Calleja ML, Duarte CM, *et al.* Mesopelagic
691 prokaryotic bulk and single-cell heterotrophic activity and community composition in
692 the NW Africa-Canary Islands coastal-transition zone. *Prog Oceanogr* 2009, **83**(1-4): 189-
693 196.
694

695 54. Baltar F, Arístegui J, Gasol JM, Herndl GJ. Prokaryotic carbon utilization in the dark
696 ocean: growth efficiency, leucine-to-carbon conversion factors, and their relation. *Aquat*
697 *Microb Ecol* 2010, **60**(3): 227-232.
698

699 55. Yamashita Y, Tanoue E. In situ production of chromophoric dissolved organic matter in
700 coastal environments. *Geophys Res Lett* 2004, **31**(14).
701

702 56. Smith DC, Simon M, Alldredge AL, Azam F. Intense Hydrolytic Enzyme-Activity on Marine
703 Aggregates and Implications for Rapid Particle Dissolution. *Nature* 1992, **359**(6391): 139-
704 142.
705

706 57. Gasol JM, del Giorgio PA, Massana R, Duarte CM. Active versus inactive bacteria: size-
707 dependence in coastal marine plankton community. *Mar Ecol Prog Ser* 1995, **128**: 91-97.
708

709 58. Ngugi DK, Antunes A, Brune A, Stingl U. Biogeography of pelagic bacterioplankton across
710 an antagonistic temperature-salinity gradient in the Red Sea. *Mol Ecol* 2012, **21**(2): 388-
711 405.
712

713 59. Konneke M, Bernhard AE, de la Torre JR, Walker CB, Waterbury JB, Stahl DA. Isolation of
714   an autotrophic ammonia-oxidizing marine archaeon. *Nature* 2005, **437**(7058): 543-546.
715

716 60. Bianchi D, Babbin AR, Galbraith ED. Enhancement of anammox by the excretion of diel
717   vertical migrators. *P Natl Acad Sci USA* 2014, **111**(44): 15653-15658.
718

719 61. del Giorgio PA, Cole JJ. Bacterial growth efficiency in natural aquatic systems. *Ann Rev
720   Ecol Syst* 1998, **29**: 503-541.
721

722 62. Reinthaler T, van Aken H, Veth C, Arístegui J, Robinson C, Williams PJLR, *et al.*
723   Prokaryotic respiration and production in the meso- and bathypelagic realm of the
724   eastern and western North Atlantic basin. *Limnology and Oceanography* 2006, **51**(3):
725   1262-1273.
726

727 63. Lemée R, Rochelle-Newall E, Van Wambeke F, Pizay MD, Rinaldi P, Gattuso JP. Seasonal
728   variation of bacterial production, respiration and growth efficiency in the open NW
729   Mediterranean Sea. *Aquatic Microbial Ecology* 2002, **29**(3): 227-237.
730

731 64. Arrieta JM, Mayol E, Hansman RL, Herndl GJ, Dittmar T, Duarte CM. Dilution limits
732   dissolved organic carbon utilization in the deep ocean. *Science* 2015, **348**(6232): 331-
733   333.
734

735 65. Dall'Olmo G, Dingle J, Polimene L, Brewin RJW, Claustre H. Substantial energy input to
736   the mesopelagic ecosystem from the seasonal mixed-layer pump. *Nat Geosci* 2016,
737   **9**(11): 820-+.
738

739 66. Giering SLC, Sanders R, Lampitt RS, Anderson TR, Tamburini C, Boutrif M, *et al.*
740   Reconciliation of the carbon budget in the ocean's twilight zone. *Nature* 2014,
741   **507**(7493): 480-+.
742

743 67. Boeuf D, Edwards BR, Eppley JM, Hu SK, Poff KE, Romano AE, *et al.* Biological
744   composition and microbial dynamics of sinking particulate organic matter at abyssal
745   depths in the oligotrophic open ocean. *P Natl Acad Sci USA* 2019, **in press**.
746

747 68. Pernthaler J. Predation on prokaryotes in the water column and its ecological
748   implications. *Nat Rev Microbiol* 2005, **3**(7): 537-546.
749

750 69. Luna GM, Bianchelli S, Decembrini F, De Domenico E, Danovaro R, Dell'Anno A. The dark
751   portion of the Mediterranean Sea is a bioreactor of organic matter cycling. *Global
752   Biogeochem Cy* 2012, **26**.
753

754 70. Murphy KR, Butler KD, Spencer RGM, Stedmon CA, Boehme JR, Aiken GR. Measurement
755   of Dissolved Organic Matter Fluorescence in Aquatic Environments: An Interlaboratory
756   Comparison. *Environ Sci Technol* 2010, **44**(24): 9405-9412.

758 71. Stedmon CA, Bro R. Characterizing dissolved organic matter fluorescence with parallel
759   factor analysis: a tutorial. *Limnol Oceanogr-Meth* 2008, **6:** 572-579.

761 72. Coble PG. Marine optical biogeochemistry: The chemistry of ocean color. *Chem Rev*
762   2007, **107**(2): 402-418.

764 73. Gasol JM, Morán XAG. Flow cytometric determination of microbial abundances and its
765   use to obtain indices of community structure and relative activity. In: McGenity TJ,
766   Timmis KN, Nogales B (eds). *Hydrocarbon and Lipid Microbiology Protocols*. Springer:
767   Berlin, Germany, 2015, pp 159-187.

769 74. Calvo-Díaz A, Morán XAG. Seasonal dynamics of picoplankton in shelf waters of the
770   southern Bay of Biscay. *Aquatic Microbial Ecology* 2006, **42**(2): 159-174.

772 75. Gundersen K, Orcutt KM, Purdie DM, Michaels AF, Knap AH. Particulate organic carbon
773   mass distribution at the Bermuda Atlantic Time-series Study (BATS) site. *Deep-Sea Res II*
774   2001, **48:** 1697-1718.

775
776