Genomic phylogeography of the White-crowned Manakin *Pseudopipra pipra* (Aves: Pipridae) illuminates a continental-scale radiation out of the Andes

Author Affiliations:

Jacob S. Berv[1,2*], Leonardo Campagna[1,2], Teresa J. Feo[3], Ivandy Castro-Astor[4], Camila C. Ribas[5], Richard O. Prum[6], Irby J. Lovette[1,2]

1. Fuller Evolutionary Biology Program, Cornell Lab of Ornithology, 159 Sapsucker Woods Road, Ithaca, NY 14850, USA.

2. Department of Ecology and Evolutionary Biology, Cornell University, 215 Tower Road, Ithaca, NY 14853, USA.

3. Department of Vertebrate Zoology, MRC-116, National Museum of Natural History, Smithsonian Institution, Washington, DC 20013, USA.

4. Department of Biology, City College of New York and CUNY Graduate Center, City University of New York, New York, NY 10031, USA

5. Coordenacão de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Manaus, AM, Brazil

6. Department of Ecology and Evolutionary Biology, and Peabody Museum of Natural History, Yale University, New Haven, Connecticut, 06520, USA.

*Correspondence to: Jacob S. Berv (jsb439@cornell.edu)

Current address: Department of Ecology and Evolutionary Biology, and University of Michigan Museum of Paleontology, 1105 North University Avenue, Biological Sciences Building, Ann Arbor, MI 48109-1085, USA.

A CONTINENTAL-SCALE AVIAN RADIATION OUT OF THE ANDES

Abstract: The complex landscape history of the Neotropics has generated opportunities for

population isolation and subsequent diversification that place this region among the most species-

rich in the world. Detailed phylogeographic studies are required to uncover the biogeographic

histories of Neotropical taxa, to identify evolutionary correlates of diversity, and to reveal patterns

of genetic connectivity, disjunction, and potential differentiation among lineages from different

areas of endemism. The White-crowned Manakin (*Pseudopipra pipra*) is a small suboscine

passerine bird that is broadly distributed through the subtropical rainforests of Central America, the

lower montane cloud forests of the Andes from Colombia to central Peru, the lowlands of

Amazonia and the Guianas, and the Atlantic forest of southeast Brazil. *Pseudopipra* is currently

recognized as a single, polytypic biological species. We studied the effect of the Neotropical

landscape on genetic and phenotypic differentiation within this species using genomic data derived

from double digest restriction site associated DNA sequencing (ddRAD), and mitochondrial DNA.

Most of the genetic breakpoints we identify among populations coincide with physical barriers to

gene flow previously associated with avian areas of endemism. The phylogenetic relationships

among these populations imply a novel pattern of Andean origination for this group, with

subsequent diversification into the Amazonian lowlands. Our analysis of genomic admixture and

gene flow establishes a complex history of introgression between some western Amazonian

populations. These reticulate processes confound our application of standard concatenated and

coalescent phylogenetic methods and raise the question of whether a lineage in the western Napo

area of endemism should be considered a hybrid species. Lastly, analysis of variation in vocal and

plumage phenotypes in the context of our phylogeny supports the hypothesis that *Pseudopipra* is a

species-complex composed of at least 8, and perhaps up to 17 distinct species which have arisen in

the last ~2.5 Ma.

Key words: Neotropics, ddRAD, genomics, phylogeography, manakin, suboscine, speciation,

introgression, admixture, reticulation, multi-species coalescent, coancestry, isolation by distance,

vocalization

1. *Introduction*

Many kinds of geographic and habitat barriers have been proposed to drive population

diversification and speciation in the Neotropics (Cracraft and Prum, 1988; Haffer, 1969, 2008;

Smith et al., 2014; Wallace, 1854). Such barriers partition biodiversity into areas of endemism

(Cracraft, 1985; Cracraft and Prum, 1988; Crother and Murray, 2011; Da Silva et al., 2005; Linder,

2001; Noguera-urbano, 2016) by acting as impediments to gene flow for dispersal-limited

organisms (Brumfield, 2012; Cheviron et al., 2005; Fernandes et al., 2015; Moore et al., 2008;

Ribas et al., 2012). Three of the most prominent features implicated in structuring the biodiversity

of Neotropical forest birds (Figure 1) include the Andes Mountains and other montane regions

(Figure 1, grey relief); the Chaco, Cerrado, and Caatinga biomes, which collectively form a "dry

diagonal" of open habitat separating the Amazon forest from the Atlantic Forest; and the large

rivers of the complex Amazonian drainage system (Brumfield, 2012; Harvey and Brumfield, 2015;

Naka and Brumfield, 2018; Smith et al., 2014).

Montane Andean regions in the Neotropics are known to be exceptionally biodiverse,

encompassing at least 15 areas of avian endemism with biotas shaped by a combination of

vicariance and dispersal events (Hazzi et al., 2018; Morrone, 2015). Hazzi et al. (2018) concluded

that the Marañon, Apurimac, and Yungas Inter-Andean Valleys, as well as the Colombian Massif,

Las Cruces Pass, and the Táchira Depression have all likely played significant roles in structuring

Andean biodiversity. Elevational gradients in the Andes also contribute substantially to Neotropical

diversification metrics and raise fundamental questions about the historical relationships between

lowland rainforest and montane endemics in the Neotropics (Janzen, 1967; Musher et al., 2019;

Quintero and Jetz, 2018; Weir, 2006). From a biogeographic perspective, avian evolution has

proceeded both "into and out of the Andes" (Brumfield and Edwards, 2007; Nylander et al., 2008).

A number of examples illustrate colonization of Andean forest regions by lowland forest ancestors

(Bates and Zink, 1994; Fjeldså, 1992; Ribas et al., 2007), whereas examples of diversification into

the lowlands are so far restricted to open habitat taxa (da Silva, 1995; van Els et al., 2019; Voelker,

1999). For some tanagers (Aves: Thraupidae), the Northern Andes have been a source of lineages

that later dispersed into the Central Andes and Amazonian lowlands (Sedano and Burns, 2010). By

contrast, Southern Andean forests may have been connected to the Atlantic Forest via Pleistocene

expansions of forests into the region occupied today by the Cerrado biome (Cabanne et al., 2019).

Indeed, glacial cycles and climatic fluctuations during the Pleistocene have been implicated as an

important factor in montane diversification, disproportionally affecting high elevation forest

(Hooghiemstra and Van der Hammen, 2004; Weir, 2006) and transiently connecting highland and

lowland habitats (Brumfield and Edwards, 2007; Cabanne et al., 2019; Nylander et al., 2008).

In the lowland Amazon basin, a number of areas of endemism have also been described,

delimited primarily by major tributaries of the Amazon River (Borges and Da Silva, 2012; Cracraft,

1985; Da Silva et al., 2005; Haffer, 1974). Wallace (1854) initially suggested that the Amazon basin

could be divided into four wide bioregions (which he termed "Guyana," "Ecuador," "Peru," and

"Brazil"), based on primate distributions (see also Lynch Alfaro et al., 2015). These areas were

subsequently partitioned by later biogeographers into at least eight major areas of endemism for

terrestrial vertebrates (Figure 1) (Borges and Da Silva, 2012; Cracraft, 1985; Cracraft and Prum,

1988; Da Silva et al., 2002; Da Silva et al., 2005; Haffer, 1978; Haffer, 1985; Haffer, 1992; Naka,

2011). The aggregate of these geographic partitions has been recognized as the "Amazonian areas

of endemism" and is the basis for the riverine barrier hypothesis (Figure 1, Antonelli et al., 2018;

also see Figure 1A in Silva et al., 2019).

Few studies of birds have applied next-generation sequencing datasets to reconstruct the

biogeographic history of a Neotropical radiation distributed across lowland and highland areas of

endemism. Here, we investigate lineage diversification across the Neotropics in the continentally

distributed *Pseudopipra* genus. The monotypic genus *Pseudopipra* (Kirwan et al., 2016), family

Pipridae, currently includes a single biological species, the White-crowned Manakin (*Pseudopipra*

pipra = Dixiphia). *Pseudopipra* is found across a variety of known dispersal barriers, including

elevational gradients of the Andes, major Amazonian rivers, the dry diagonal, tepuis, and the

Isthmus of Panama. Therefore, this group is particularly appropriate for assessing patterns of

phylogenomic differentiation across a nested set of spatial scales in Neotropical forests.

In order to assess the historical relationships among and within Andean and lowland

Amazonian forest populations, and the degree to which population structure may be related to

landscape features, our study design uses fine-scale sampling across a majority of *Pseudopipra's*

range. We use double digest restriction site associated DNA (RAD) sequencing (Peterson et al.,

2012)—a reduced representation genomic sequencing technique— to sample thousands of

independent markers across the genomes of hundreds of individuals spanning many well-known

Amazonian and Andean areas of endemism, from Costa Rica to the Atlantic Forest of Brazil (> 80

localities). We use these data to infer the history of lineages and populations within *Pseudopipra*

using a variety of population genomic, landscape genetic, phylogenetic, and comparative

approaches. Lastly, we investigated congruence between geographic variation in vocal variation,

plumage phenotype, and phylogeographic structure, to evaluate the biological mechanisms that may

be contributing to population differentiation and gene flow, and to propose a new hypothesis of

species delimitation for the genus.

1.1. *Study organism*

The White-crowned Manakin, *Pseudopipra pipra* (Pipridae, hereafter *Pseudopipra*), is a

small (10-12 gram), non-migratory, suboscine passerine bird that is broadly distributed within

Central America, the lower montane cloud forests of the Andes (up to ~2000m) from Colombia to

central Peru the Amazon basin, and the Atlantic Forest (Figure 2) (Kirwan et al., 2016; Kirwan and

Green, 2012). The male's striking white crown and jet-black body makes them among the most

easily identified manakins, though the grey-green females are often confused with other manakin

species in the field. *Pseudopipra* are typically found in dense humid forest and are predominantly

frugivorous. Unlike many manakin species that exhibit concentrated or cooperative lek behavior

(e.g. Prum, 1990; Prum, 1994), *Pseudopipra* males display in dispersed leks of 2-5 males (Castro-

Astor et al., 2007; Snow, 1961). Castro-Astor et al. (2007) described at least 11 components in the

display repertoire of the Atlantic forest *Pseudopipra* population, including rapid turning, jumping,

"to-and-fro" flights, and an about-face.

A recent phylogenetic hypothesis of manakins based on mitochondrial and nuclear DNA

sequences placed *Pseudopipra* as sister group to the five species of *Ceratopipra* (Ohlson et al.,

2013). While the higher level phylogeny and taxonomy of manakins has received significant

attention (Ohlson et al., 2013; Tello et al., 2009), work exploring intraspecific genetic variation

within manakins has been restricted to relatively few species (e.g. Brumfield et al., 2008;

Capurucho et al., 2013; Cheviron et al., 2006; Cheviron et al., 2005; Francisco et al., 2007; Gubili et

al., 2016; Luna et al., 2017; McDonald, 2003).

Based on geographic variation in plumage and vocalizations, many previous authors have

suggested that the biological species *Pseudopipra pipra* likely includes multiple, distinct

phylogenetic species (Freile, 2014; Kirwan and Green, 2012; Ridgely and Greenfield, 2001;

Ridgely and Tudor, 2009; Snow, 2004; Spencer, 2012). Thirteen subspecies of *Pseudopipra pipra*

have been recognized based primarily on subtle variations in plumage coloration, which are

frequently more marked in females than in males (summarized in Dickinson, 2003; Kirwan and

Green, 2012; Snow, 2004). Because suboscine passerines generally do not learn their songs (but see

Saranathan et al., 2007), the presence of substantial vocal variation across *Pseudopipra* populations

further suggests that the genus may contain unrecognized cryptic species (e.g. Campagna et al.,

2012). However, all previous authors awaited new information on genetic differentiation within

Pseudopipra before making taxonomic recommendations.

It also remains unclear how the current subspecific classification reflects evolutionary

history. The genetic variation within *Pseudopipra* has received prior attention by two studies that

used mitochondrial DNA to infer population genetic patterns. Milá et al. (2012) reported

intraspecific divergences of up to 3.5% (n=19) across three Amazonian populations, and the highest

observed nucleotide diversity ($\pi = 0.266$) among 14 widely distributed Amazonian birds species.

Castro-Astor (2014) used a larger sample (n=57), and discovered that at least four *Pseudopipra*

subspecies correspond to well supported mitochondrial clades which were generally congruent with

Amazonian areas of endemism. Castro-Astor (2014) used a molecular clock model to estimate the

age of the *Pseudopipra* complex to coincide with the onset of the Pleistocene ~2.457 Ma (95% HPD

1.45-3.97).

2. *Material and methods*

2.1. *Field and tissue sampling*

Muscle tissue samples were obtained from available vouchered avian material from US and

Brazilian collections and other institutions (see Acknowledgements and Supplemental Table 1). We

sampled from ten of thirteen recognized subspecies (Figure 2) of *Pseudopipra* (Kirwan and Green,

2012). Unfortunately, we were unable to obtain material of adequate quality for ddRAD sequencing

from the Andes of Colombia (*bolivari*, *unica*, *coracina*, *minima*), Ecuador (*minima*) and the

lowlands of Peru (*pygmaea*). However, we were able to obtain mtDNA data from *coracina* and

pygmaea, which allowed us to make a preliminary assessment of their phylogenetic affinities (see

discussion on mtDNA). In sum, after discarding failed samples and samples with high proportions

of missing data, we obtained genetic data from 286 individuals (234 from ddRAD, 168 from

mtDNA), representing ~80 localities and 10 of 13 subspecies. We also obtained comparable ddRAD

and mtDNA data for two specimens of *Ceratopipra rubrocapilla* as outgroups (Supplementary

Table 1).

2.2. *Laboratory methods and data processing*

We extracted DNA from avian tissue specimens (Supplementary Table 1) using the DNeasy

Blood & Tissue Kit (QIAGEN, CA) and generated ddRADtags following the protocol of Peterson

et al. (2012), with modifications described in Thrasher et al. (2018). We used the STACKS 1.44

bioinformatics pipeline (Catchen et al., 2013) to generate assemblies of ddRAD loci, using

Manacus vitellinus as a reference genome (GCA_001715985.1 from www.ncbi.nlm.nih.gov). We

obtained mitochondrial ND2 sequences for a set of individuals using standard Sanger Sequencing

protocols described in Berv and Prum (2014). Additional sequencing, assembly, and alignment

details are reported in the Supplemental Appendix.

For ddRAD loci, we used several filters in the "populations" module from STACKS to

generate different sets of bi-allelic SNPs: a missing data filter, a minimum depth of coverage filter,

a filter that exports only the first SNP per RAD locus, and a minor allele frequency filter (MAF). In

order to explore the sensitivity of genetic cluster inference to these important parameters (Linck and

Battey, 2017; Paris et al., 2017), we produced eight versions of the SNP dataset by combining these

different standard filters (Table 1).

Dataset 1 (D1- labeled arbitrarily) included 1,960 SNPs that were generated by setting the

MAF to >= 5%, requiring minimum coverage of 20x, requiring the presence of a SNP in at least

80% of individuals, and filtering out all but the first SNP in each RAD locus to minimize linkage

across the dataset. To evaluate the sensitivity of cluster analyses to rare variants (Linck and Battey,

2017; Shafer et al., 2017), Dataset 2 (D2) was exported with identical parameters to dataset D1, but

without a minor allele frequency filter; this generated a dataset of 2,581 putatively unlinked SNPs.

We generated datasets D3 and D4 to reflect the same filters as D1 and D2 but to also include all

SNPs from each locus. Lastly, for each dataset D1-D4, we generated a complimentary dataset which

filtered out any sites with missing data. The parameters of these SNP datasets are summarized in

Table 1.

2.3. *Phylogenetic analysis*

Because sparse super-matrices may be more prone to systematic error (e.g. Roure et al.,

2013), we explored the sensitivity of our phylogenetic inferences to increasing amounts of missing

data using three different haplotype datasets (separate from SNP Datasets) with 20%, 50%, and

80% thresholds for missing data at the RAD locus level. For example, for an 80% threshold, if the

RAD locus is present in at least 20% of individuals, then that locus, containing all variable and

invariable sites, was exported. This approach generated datasets comprising 2,584, 4,763, and 7,901

RAD loci, respectively. For each dataset, we generated a concatenated supermatrix using Sequence

Matrix (Vaidya et al., 2011). These combined datasets had up to ~40% missing sites across an entire

matrix, irrespective of columns or rows (i.e., using the 80% threshold above).

We estimated phylogenies using each dataset with a concatenated maximum likelihood and

a coalescent species-tree approach for a total of six phylogenetic analyses. For analysis of

concatenated data, we estimated a maximum likelihood phylogenetic hypothesis using RAxML

8.2.9 (Stamatakis, 2014). We estimated 500 bootstrap replicates followed by a full ML search for

each dataset, using a GTRGAMMAX substitution model. We avoid the need for ascertainment bias

correction because we use whole short-read haplotype alignments including invariant sites (Leaché

et al., 2015). In our case, each locus alignment was 138bp long (supplemental Appendix). We also

tested for an effect of data partitioning on phylogenetic inference by generating optimized locus

partitioning schemes in PartitionFinder 2 (Lanfear et al., 2017), which were then re-analyzed in

RAxML. We found that our topological results were insensitive to data partitioning, and as such we

do not present these results. Lastly, we generated a mitochondrial DNA ND2 gene tree using IQ-

TREE 1.6.10 (Chernomor et al., 2016; Hoang et al., 2017; Kalyaanamoorthy et al., 2017; Schmidt et

al., 2014; Trifinopoulos et al., 2016). We partitioned by codon position and generated a maximum

likelihood tree using the MFP+MERGE model search and partitioning option (Supplementary

Figure 17).

We evaluated coalescent phylogenetic structure using the SVDquartets method implemented

in PAUP* 4.0a159 (Chifman and Kubatko, 2014, 2015; Swofford, 2002). SVDquartets may be a

useful tool for testing hypotheses of species delimitation because it can generate a lineage-tree

under the multispecies coalescent without assigning individuals to species *a priori*. In this approach,

each individual in a dataset is considered to be a separate "species." If a recovered clade in such a

lineage-tree has high bootstrap support, this would mean that the descendant taxa have strong

support as a group under the multispecies coalescent, which may coincide with species boundaries

(*Personal communication*, Laura Kubatko). Our SVDquartets analysis evaluated all possible

quartets with 500 bootstrap replicates (other settings left to defaults). SVDQuartets has the

advantage of being able to work directly on sequence data (above) and has been shown to perform

well in comparison to summary methods which assume fully and correctly resolved gene trees as

input (Chou et al., 2015; Schmidt-Lebuhn et al., 2017). SVDquartets has also been shown to

produce reliable results under very general conditions, and is theoretically robust to variation in

effective population size, molecular clocks, limited gene flow, and incomplete lineage sorting (Long

and Kubatko, 2017).

While a full exploration of our specific usage of SVDquartets is beyond the scope of the

present work, we suggest that this approach may be a useful way to assess support for hypotheses of

species delimitation under the multispecies coalescent. We do not take the perspective that all well

supported clades from this coalescent assessment *necessarily* represent biological species – such an

interpretation often leads to an over-inflation of species estimates (Chambers and Hillis, 2020;

Hillis, 2020; Sukumaran and Knowles, 2017). In general, we prefer a species agnostic approach to

those which require *a priori* delimitation of species boundaries. Our approach instead focuses on

discovering well supported coalescent structure without a prior hypothesis of species delimitation

and then assesses the degree to which metrics of support across coalescent and concatenated tree

inference methods are concordant. We note that there are many approaches for species delimitation

using genetic data (e.g. BP&P, Flouri et al. (2018), BFD, Leaché et al. (2014), PTP, Zhang et al.

(2013), also see Leaché et al. (2018)), and that there is lack of consensus on how best to interpret

these analyses with respect to various species concepts (Carstens et al., 2013). In the present study,

we instead rely on analysis of vocal and plumage phenotypes in combination with genetic patterns,

to make taxonomic recommendations.

2.4. *Descriptive statistics*

We estimated global and population pairwise F_{st} (Weir and Cockerham, 1984), regional

variation in allelic richness (Goudet, 2005), Hardy-Weinberg equilibrium (Paradis, 2010),

inbreeding (Goudet, 2005), linkage disequilibrium (Kamvar et al., 2014), sex-biased dispersal

(Goudet et al., 2002), and an analysis of molecular variance (Kamvar et al., 2014).

Some of these statistics and analytical details are reported as supplementary material, while others

are discussed below when relevant.

2.5. *Cluster inference and patterns of genomic admixture*

We performed cluster analyses intended to quantify the number of similar genetic groups

across the range of *Pseudopipra*. First, we evaluated broad-scale variation across our genomic data

using phenetic K-means clustering implemented in the find.clusters function in the adegenet R

package (Jombart, 2008). This function first transforms the data with a PCA, and then identifies

clusters with a K-means algorithm (MacQueen, 1967). Using each SNP dataset, we evaluated K1:20

successively with the Bayesian Information Criterion (BIC). K-means clustering is based on the

idea of minimizing the variance within clusters and has the potential advantage of making few

assumptions about the underlying processes which generated the data. For each analysis, we

retained all PC axes, set 200 randomly chosen starting centroids, and ran each search for 10^9

iterations.

We also estimated the number of differentiated populations using STRUCTURE 2.3.4

(Pritchard et al., 2000), a widely-used model-based approach which assigns individuals to

populations such that Hardy-Weinberg equilibrium is maximized and linkage among loci is

minimized within groups. By comparing the output of STRUCTURE runs fixed to different

numbers of populations (K), one can assess the degree of fit of the data to various models of K and

assess genomic admixture among individuals. We ran STRUCTURE using SNP dataset D1 (Table

1), from K1:20 with 10 iterations per K value. We ran the program for 700,000 generations,

discarding the first 200,000 as burn-in, implementing the admixture ancestry model with correlated

allele frequencies. We evaluated K by examining $Pr(X|K)$ or $L(K)$ after summarizing our results

using Structure Harvester (Earl and vonHoldt, 2012) and CLUMPP (Jakobsson and Rosenberg,

2007). We also used the Evanno method to estimate the rate of change in the log probability of data

between values of K, which is a useful metric to detect the uppermost hierarchical level of structure

in the data (Evanno et al., 2005). Lastly, we generated 95% Bayesian Credible Intervals around

admixture coefficients (presented as averages across individuals for population assignment).

To investigate finer scale population genetic differentiation and the genomic composition of

populations defined at different hierarchical levels, we used the programs fineRADstructure and

RADpainter (Malinsky et al., 2018). RADpainter takes advantage of the information in haplotype

data and considers each individual in a dataset as a "recipient" whose genomes are reconstructed

using chunks of "donor" DNA from all other available individuals. This approach generates a

"coancestry matrix" which combines the information that can be derived from both PCA and

model-based clustering approaches and thus can be more sensitive to subtle population structure

(Lawson et al., 2012). The fineRADstructure software uses an MCMC approach to explore the

space of population assignments based on the coancestry matrix, using an algorithm which merges

and splits populations, or moves individuals among populations. We ran fineRADstructure with

default priors, but increased the burn in to 200000 iterations, followed by 1000000 iterations,

sampling every 1000. We then assessed convergence by 1) considering the assignment of

population membership across multiple independent runs, 2) visualizing the MCMC traces of

estimated parameters to ensure convergence on the same posterior distributions, and 3) running

each chain long enough to achieve effective parameter sample sizes > 100.

Both RADpainter and fineRADstructure are based on the recently developed chromoPainter

and finestructure software (Lawson et al., 2012), but are optimized to take advantage of the linkage

properties of RAD-like datasets. As our data are assembled relative to a *Manacus* reference

genome, the order in which loci appear in our data files are related to positions on assembly

scaffolds, even though the ddRAD loci in our data are generally unlinked when considered at the

population level (see Supplementary Appendix).

We compared the inferred population assignments from fineRADstructure and

STRUCTURE with K-means clustering (as above) of the coancestry matrix as well as SNP datasets.

To determine how much of the variation in our dataset was captured by the RADpainter analysis

relative to a standard PCoA of SNP data, we used the normalized PCA approach of (Lawson et al.,

2012), using the mypca function provided in the "FinestructureLibrary.R" library in the

fineRADstructure package.

2.6. *Phylogenetic reticulation in the western Napo lineage*

Model-based, phenetic, and population genetic analyses revealed that a lineage of

individuals presently restricted to the western Amazonian Napo area of endemism has a complex

history of introgression between southern and northern Amazonian clades (see Results). To further

investigate this subset of individuals and their history, we generated two additional versions of the

dataset optimized for different analyses. First- we generated a haplotype dataset of 52 individuals,

which included all putatively introgressed western Napo individuals (n=10, clade E), as well as

samples from proximate neighbor populations. These populations consist of 1) a geographically

adjacent population inferred to have mostly unadmixed ancestry (Southwestern Amazon: Inambari,

n=17, clade C1), and 2) the implied source of introgressed genotypes (north-central Amazonian Jaú

+ eastern Napo, n=25, clade D2).

We used a 5% minor allele frequency cutoff, required 20x minimum coverage, and allowed

up to 20% missing data per locus—generating a dataset comprising 2,370 loci, including 4,979

SNPs (Dataset H1). Dataset H1 is analogous to that generated for coancestry (fineRADstructure)

analysis of the entire dataset (above). We also generated a similar dataset (Dataset H2), without a

minor allele frequency cutoff—comprising 2,947 loci including variable and invariable sites—as

required for inference of demographic parameters (Gronau et al., 2011).

Hybrid populations are expected to have higher mean coancestry and lower F_{st} with each of

their progenitor lineages than their progenitor lineages will have with each other (Barrera-Guzmán

et al., 2018). To test this prediction, we first estimated coancestry values using Dataset H1 and

RADpainter (as above). We then compared inferred coancestry of population pairs with a standard

ANOVA and the glht function in the multcomp R package (Hothorn et al., 2013). We estimated F_{st}

with Weir and Cockerham's estimator (Weir and Cockerham, 1984), and tested the prediction using

1000 bootstrapped datasets to estimate 95% confidence intervals.

To test whether or not the relationships among these focal populations is best represented by

a phylogenetic network, we used PhyloNet 3.6.4 (Wen et al., 2018; Zhu et al., 2018). PhyloNet

takes SNP data and uses a reversible-jump MCMC technique to explore the posterior distribution of

phylogenetic networks and bi-furcating topologies, while accommodating both reticulation and

incomplete lineage sorting. PhyloNet searches the set of all possible reticulation models without

requiring *a priori* model specification.

For computational tractability, we used SNP dataset D1 (1960 unlinked SNPs), and pruned it

to the 52 individuals comprising the groups of interest. We pruned out all sites with missing data,

leaving 572 unlinked biallelic SNPs, as required by PhyloNet. For our final runs of the PhyloNet

program MCMC_BiMarkers, we randomly subsampled five diploid individuals from each group, as

preliminary program runs indicated a severe computational bottleneck with more than five diploid

individuals or with larger datasets. We assigned each group of individuals to a lineage, allowed for

a maximum of one reticulation, set the chain length for 800000 iterations (sampling every 100), set

a burn in of 2.5%, and left all priors as default (i.e. assuming population mutation rates are

constant).

Using the final network topology estimated from PhyloNet, we estimated demographic

parameters using G-PhoCS version 1.3 (Gronau et al., 2011). We ran the program for 750,000

iterations with a 10% burn-in, estimating a total of 13 parameters: the effective population sizes of

the three focal lineages, two ancestral population sizes, two splitting times, and six directional

migration parameters. Parameter MCMC traces were inspected in Tracer 1.7.1 (Rambaut et al.,

2018). We converted the median and 95% Bayesian credible intervals from mutation scale to

generations and individuals as described in Campagna et al. (2015), assuming an approximate

mutation rate of 10^{-9} per base pair per generation (Kumar and Subramanian, 2002; Smeds et al.,

2016). We calculated the number of migrants per generation as the product of the per generation

migration rate multiplied by ¼θ for the receiving population (m_{a-b} * ($\theta_b/4$)) (Gronau et al., 2011),

and considered migration rates to be significant if the 95% lower bound was > 1 (the level of gene

flow expected to counteract the effects of drift; Wright, 1931).

2.7. *Spatial distribution of genetic variation*

Dispersal barriers often create sharp genetic discontinuities between adjacent populations

that would otherwise be continuous (Petkova et al., 2015). To assess the degree to which spatial

variation in genetic diversity within *Pseudopipra* may be attributed to landscape features, we

investigated how discrete and continuous patterns of isolation-by-distance (IBD) vary across the

landscape. First, we used Mantel and partial Mantel tests to investigate IBD effects within sampling

regions delimited by prominent physical barriers, and then separately tested the roles of specific

dispersal barriers in structuring genetic variation. Mantel tests evaluate the correlation among two

or more matrices, in our case representing pairwise genetic and geographic distance. Significance is

assessed by permuting the rows and columns of one of these matrices (Mantel, 1967). Partial

Mantel tests incorporate a third "barrier matrix," which contains information about environmental

or ecological distance. By evaluating the correlation between genetic and ecological distance while

controlling for geographic distance, a partial Mantel test can be used to investigate the effect of a

particular geographic barrier on genetic differentiation. Because mantel tests may have an inflated

type I error rate (e.g. Bradburd et al., 2018; Guillot and Rousset, 2013; Legendre and Fortin, 2010),

we follow the recommendations of Diniz-Filho et al. (2013), and only reject the null hypothesis of

no correlation if $p < 0.001$ (~10x more stringent than a standard correction for multiple tests would

require).

For these tests and others, we generated a pairwise geographic distance matrix by

calculating great circle distances using the distm function in the geosphere R package (Hijmans et

al., 2015). Individuals sampled at the same GPS coordinates were set to have an arbitrarily small

geographic distance of 0.0001 meters. In order to assess spatial variation *within and between*

populations, we use estimates of pairwise genetic distance at the individual level, rather than at the

population level. Our analyses compare patristic phylogenetic distance between all pairs of

individuals (estimated from RAxML) as well as coancestry (from fineRADstructure), as proxies of

genetic distance among individuals. These procedures, including the generation of an appropriate

barrier matrix, are developed in a set of R functions and scripts that operate on adegenet GENIND

objects (see supplementary R code) and internally use the Mantel implementation from the ecodist

R package (Goslee and Urban, 2007).

We also analyzed continuous patterns of spatial variation in our data with a population

genetic model that relates effective migration rates across geographic space to expected genetic

dissimilarities among individuals (Petkova et al., 2015). This recently proposed method, termed

Estimated Effective Migration Surfaces (EEMS), produces a visualization that emphasizes

deviations from isolation by distance to discover migratory corridors and barriers to gene flow. This

method is complimentary to more explicit hypothesis-testing approaches (such as Mantel tests),

because it seeks to identify where the assumption of constant IBD is violated in geographic space,

and thereby can highlight areas where strong barriers to gene flow may exist. In brief, a region is

covered with a dense regular grid connecting subpopulations (demes), among which individuals can

migrate with rates varying by location. EEMS uses an MCMC approach to estimate expected

genetic dissimilarity between two individuals, integrating over all possible migration histories, and

adjusting migration rates among graph edges to match the genetic differences in the data. A

migration surface is then interpolated across a region to indicate where genetic similarities decay

faster than what a constant isolation by distance model predicts, visually highlighting barriers to

gene flow and migration corridors. An effective diversity parameter is also estimated for every

deme, reflecting local deviations in heterozygosity. We ran several analyses using default priors,

iteratively increasing the grid density and chain length, with up to 2000 estimated demes. For each

test, we ran three chains, and after examining chain convergence, we assessed model fit by

comparing the observed and fitted dissimilarity between and within demes.

2.8. Vocal variation

We obtained sound recordings of vocalizations of *Pseudopipra* from the Macaulay Library

at the Cornell Lab of Ornithology (https://www.macaulaylibrary.org) and the Xeno-canto

(www.xeno-canto.org) collections (Supplementary Table 4). Vocalizations were analyzed in Raven

Pro 1.4 (Bioacoustics_Research_Program, 2011), and converted into spectrograms using a 512-

sample Hann window with 50% overlap. Audio recordings ranged from a cut of a single

vocalization to longer recordings of a lek with multiple individuals vocalizing. For our initial

assessment, we identified qualitatively distinct vocalization types based on our own survey of the

available recordings, without regard to the geographic location of the recordings or subspecies

identity.

Pseudopipra vocalizations have 1-3 buzzy and/or tonal notes. We measured: 1) starting

frequency, 2) ending frequency, 3) minimum frequency, 4) maximum frequency, 5) number of

notes, and 6) duration of the entire vocalization (Supplementary Figure 13). We performed principal

components analysis (PCA) and logistic regression on these vocal measurements to test for

significant differences between vocal types and to reduce the dimensionality of the data for

comparison to results from analysis of genetic data. Because no sound records were directly

associated with genetic samples in this study, we used geographic proximity to vocalization

recordings and localization to areas of endemism or areas bounded by clear physical barriers to

associate vocal types to genetic samples and taxonomic ranks. This approach assumes that

genetically and geographically proximate individuals are likely to share the same vocal type, and

enabled us to perform a preliminary assessment of how variation in vocalization type maps onto

existing genetic variation. Testing the fine-scale association of genetic and vocalization boundaries

will require extensive field sampling from individual manakins. Additional analytical details and

analyses are reported as supplementary material.

3. Results

3.1. Genetic *data collection*

Our total genomic data set included an average of 1 ± 0.4 million sequences of 138 bp from

232 individuals of *Pseudopipra* and two *Ceratopipra rubrocapilla,* which were used as an outgroup.

Our reference-based assembly produced a catalogue with 47,046 RAD loci. Filtered SNP datasets

requiring a SNP to be present in at least 80% of individuals ranged from 1,960 unlinked SNPs

(dataset D1: 5% MAF with 1st SNP on each locus) to 41,024 SNPs (dataset D4: all SNPs on all loci,

no MAF) (Table 1). Datasets requiring a SNP to be present in 100% of individuals were

considerably smaller, as expected, and ranged from 175 SNPs in D1mod, to 3,832 SNPs in D4mod

(Table 1). We also obtained 168 mtDNA ND2 sequences using PCR methods which broadly

overlapped with our ddRAD dataset. Our mtDNA dataset included two subspecies, *pygmaea* and

coracina, from which we were not able to obtain enough high-quality material for ddRAD

sequencing.

3.2. *Phylogenetic analyses*

RAxML and SVDquartets analyses generated a consistently well-supported phylogenetic

hypothesis (Figure 3). The monophyletic lineages of *Pseudopipra* have distributions that are largely

congruent with previously recognized avian areas of endemism, and further reveal much finer scale

patterns of local biogeographic differentiation. Using the two *C. rubrocapilla* individuals to root the

phylogeny, we detected support for at least five major clades across all phylogenetic analyses of

nuclear genomic data, with consistently high bootstrap support in both concatenated and coalescent

analysis (clades A-E in Figures 3-5). We first describe the structure of these major clades, followed

by descriptions of well supported substructure within each of these regional groupings. We indicate

six bootstrap values for each clade with at least some high support, from each of the three datasets

analyzed in RAxML and SVDquartets (see heatmaps, Figure 3). Phylogenetic analysis of mtDNA

(Supplemental Figure 17) generated a congruent result, except where noted below.

3.2.1 *Phylogenetic structure*

The sister group to all the other *Pseudopipra* is a clade including individuals from

subtropical Central America and the northern Andes. The northern Andes are represented in our

ddRAD dataset by a single individual from subtropical forests (~1,000 m) of the eastern slopes of

the Andes in San Martín, Peru. This subtropical San Martín specimen (identified as *P. pipra*

occulta) was the only viable sample in our genomic dataset from the entire northern Andes region

(A1 in Figure 3). Mitochondrial data from three additional specimens from the Andes of southern

Ecuador (subspecies *coracina*) form a monophyletic group with the San Martín specimen; thus, this

northern Andean sister lineage to the Central American lineage likely includes all montane Andean

populations of *Pseudopipra* in Ecuador, Colombia, and Venezuela. A well supported Central

American clade (A2 in Figure 3) was identified in all analyses, with Costa Rican (A3 in Figure 3)

and Panamanian subclades (A4 in Figure 3) supported in RAxML.

The second successively nested clade in *Pseudopipra* includes individuals from subtropical

forests of the Andes of Southern Peru, which are sister to all remaining lineages (Clade B in Figure

3). The southern Andean Peru clade is further subdivided into two very well resolved clades: a

northern clade (B2 in Figure 3), sampled from the southern Cordillera Azul in southwestern Loreto,

and a southern clade from Pasco, Junín, and Cusco (labeled B1 in Figure 3). This southern Andean

Peru clade is the sister group to a monophyletic, lowland clade (F in Figure 3) that is further

subdivided into a southern Amazon clade + Atlantic Forest clade (Clade C in Figure 3), and a

Northern Amazonian + Guiana Shield clade (Clade D in Figure 3). Our analysis of mtDNA with

sampling of subspecies *pygmaea*, from the tropical forest of lower Rio Huallaga Valley, Peru,

places this population as sister group to the entire lowland radiation (Clade F; Supplemental

Appendix).

Within the broad lowland Northern Amazonian + Guiana Shield clade (Clade D), virtually

no phylogenetic substructure is associated with strong bootstrap support. The maximum likelihood

tree however does have substantial biogeographic coherence, which we use as additional

justification for population subdivision when monophyletic groups are identified to be coincident

with geographic features. For example, all but one individual from coastal Suriname and Amapá

form a monophyletic assemblage (D1 in Figure 3), east of the Essequibo river. Likewise, all but one

individual from the far eastern Napo area of endemism, east of the Rio Putumayo, form a clade (D2

in Figure 3). This coherence with landscape features is unlikely to be coincidental, though support

for these groups is weak. We treat clade D1 and D2 as well as two other Guianan regions, as

separate populations for descriptive population genetic statistics.

A widely distributed set of individuals encompassing the western portion of the Napo area

of endemism (from the northern bank of the confluence of the Rio Solimões and the Rio Napo north

of Iquitos, then west and south) was recovered as the sister group to the Guiana Shield + Northern

Amazon clade (Clade E in Figure 3). Analysis of mtDNA (Supplementary Figure 17) further

resolved (BS > 99) two subgroups delimited by the Rio Napo. Importantly, and in contrast to

nuDNA, we infer that these mtDNA haplotypes are derived from Inambari lineages (BS = 98,

Clades C1 + C3 in Figure 3), and are not sister to Clade E. These conflicting patterns are explained

by subsequent population genetic analyses, which indicate that this Napo population is a distinct

lineage of the southern Amazonian clade (Clade C in Figure 3) that has experienced substantial

hybrid introgression from the northern Amazonian clade D (see Population Genetic analyses and

Discussion).

Within the large lowland Southern Amazonia clade (Clade C in Figure 3), we recovered

eight moderately supported and hierarchically nested clades, which are successive sister groups to

each other, and are congruent in distribution with recognized areas of endemism. First is a

moderately supported clade (C1 in Figure 3) of individuals from the western Inambari area of

endemism, from west of the Rio Ucayali to the Rio Purus. Although the first lineages to diverge

within clade C1 have low bootstrap support, a monophyletic subgroup of three montane (>1000m)

individuals from the highlands between the Rio Huallaga and Rio Ucayali cluster together with high

bootstrap support (C2 in Figure 3).

The next successively nested clade includes individuals from eastern Inambari, between the

Rio Purus and the Rio Madeira (C3 in Figure 3). The next successive clade includes two individuals

sampled from the Rondônia area of endemism, between the Rio Madeira and Rio Tapajós (C4 in

Figure 3). The next successive clade includes a group of individuals from the Tapajós area of

endemism between the Rio Tapajós and the Rio Xingu (C5 in Figure 3). Next, a clade comprised of

individuals from the northern Xingu area of endemism, east of the Rio Xingu (C6 in Figure 3), is

sister to a well-supported Brazilian Atlantic Forest clade (C7 in Figure 3), which is further

subdivided into three successive clades (north to south) in Bahia, Espírito Santo, and Rio de Janeiro,

respectively (C8, C9, and C10 in Figure 3). In summary, the subdivisions of the Southern

Amazonian clade reflect a stepwise west-east progression, with a hierarchically nested structure:

(West Inambari (East Inambari (Rondônia (Tapajós (Xingu (Bahia (Espírito Santo (Rio de Janeiro))))))).

3.3. *Cluster inference and patterns of genomic admixture*

3.3.1. *Broad-scale population structure and admixture*

Weir and Cockerham's F_{st} was moderate when comparing populations within *Pseudopipra*;

for Dataset 1, overall F_{st} was 0.195 [95% CI: 0.187-0.203], and never dipped below 0.171 [0.168-

0.175] (Dataset 4). Notably, in SNP datasets filtered to remove sites with missing data, overall F_{st}

estimates were on average ~20% higher and ranged from 0.188 [0.175-0.201] (D4) to 0.261 [0.191-

0.336] (D2). Pairwise population F_{st} estimates from Dataset 1 (other datasets showed similar

patterns, see Supplementary Figure 10a-h), ranged from essentially undifferentiated, to almost

entirely distinct. At the extremes: comparing the geographically proximate Jaú and Guiana Shield

populations indicated a particularly low F_{st}: 0.005 [0.003 - 0.007]. By contrast, comparing the

Atlantic Forest Espírito Santo population to the Panama population recovered an F_{st} of 0.81 [0.79 -

0.83], or almost entirely differentiated. For reference, Hartl and Clark (1997) indicate $F_{st} > 0.15$

may be considered "great" genetic differentiation.

526   K-means clustering of SNP datasets on the basis of BIC score identified groups of

527 individuals that differed somewhat with respect to dataset filtering strategies (Summarized in Figure

528 5, right). For D1, K-means clustering identified five major groups: Central America (Clade A2),

529 South Andean Peru (Clade B), Atlantic Forest (Clade C7), Southern Amazon including the western

530 Napo population (the rest of Clade C + Clade E), all Guiana Shield + Northern Amazon (Clade D)

531 (Supplementary Figure 2a-h). For dataset D1, the montane sample from San Martín (North Andean

532 Peru, 5444.PE.MAR) clustered with Central American populations (Clade A), as it does in all

533 phylogenetic analyses. These clusters are essentially similar to those detected by STRUCTURE at

534 K=5 (Figure 4, 5, below). All other SNP datasets grouped the San Martín specimen with the

535 Southern Amazon populations. D1m further identified the Tapajós + Xingu individuals as a distinct

536 cluster, while D2, D3, D3m, D4, and D4m generated sets of clusters identical to D1, with the

537 exception of the clustering of the San Martín specimen, which clustered with the Southern Amazon

538 in those datasets. Additional field sampling of the North Andean population is required to

539 understand why this lineage may share signals of ancestry with northern and southern groups. K-

540 means clustering of D2m failed to discriminate any lowland Amazonian populations

541 (Supplementary Figure 2d). Thus, only 75 low frequency SNPs which are removed in D1m but are

542 present in D2m obfuscate the signal of population differentiation detected by K-means clustering of

543 D1m. Lastly, across datasets, standard PCoA explained 13-26% of the variance in the SNP data on

544 the first two axes (Supplementary Figure 2).

545   Visual inspection of $Pr(X|K)$ from summarized STRUCTURE runs on dataset D1 indicated

546 that the likelihood of each successive K from 1:20 plateaued at K = 5, with the standard deviation

547 across runs increasing rapidly after this point (Supplementary Figure 3). We describe patterns of

548 inferred admixture as they appear in Figure 5, reading from top to bottom. For K5, individuals from

549 Central America are unambiguously assigned to their own cluster (Clade A2 in Figure 3, 5). The

north Andean montane specimen from San Martín, (North Andean Peru, 5444.PE.MAR) is

reconstructed as being highly admixed, with a genomic composition including 4 of the 5 inferred

clusters. However, this inference is tenuous because we have only one individual to represent this

likely distinct and highly diverse north Andean lineage (though see RADpainter below, which is

consistent with these results). Samples from the southern Amazon and Atlantic Forest (Clade C),

have no detected admixture with northern populations, except for some individuals from the

western (Clade C1), and to a more limited degree, the eastern (Clade C3) Inambari populations,

which are inferred to have limited (~<5%) admixture with individuals assigned to the Northern

Amazon + Guiana Shield (Clade D, see Figure 5, 6, and discussion below).

Going from West to East across the Southern Amazon, very limited admixture with Atlantic

Forest alleles (green, Clade C7) is first inferred in the Tapajos (Clade C5) area of Endemism. This

admixture component increases in the Xingu area of endemism (Clade C8) and increases further in

a geographic and genetic cline to 100% assignment probability to a distinct cluster in the Rio de

Janeiro clade (Clade C10). Given the geographic context of this cline in admixture, we suggest this

pattern may a product of isolation by distance, hierarchical structure, and geographic barriers

(Bradburd et al., 2018), and not isolated introgression events. We note that the conStruct method

(Bradburd et al., 2018) may be useful for explicitly testing this hypothesis (Supplemental

Appendix), however our evaluation of this method suggested that it is presently intractable for a

dataset of this size.

Next, all individuals from the western portion of the Napo area of endemism are inferred to

be heavily admixed (Clade E; Figure 5). On average, at K=5, STRUCTURE assigns

40.2% of the these individuals' (n=10) genetic composition to the Guiana Shield population (SD

4.7%, average Bayesian credible interval (aBCI): [0.33-0.47]), 50.4% to the Southern Amazon

population (SD 3.4%, aBCI: [0.43-0.58]), as well as limited 4.15% probability to Central Peru (SD

3.1%, aBCI: [0.01-0.08]), and 5.2% to Central America (SD 4.8%, aBCI: [0.02-0.1]). The three

localities included in this clade– (1) north of Iquitos, on the east bank of the confluence of the Rio

Solimões and the Rio Napo, (2) southwest of Iquitos in Allpahuayo Mishana Reserve (represented

by three mtDNA samples), and (3) near San Jacinto, Ecuador– indicate that this heavily admixed

population (corresponding to subspecies *discolor*) may occur across a large area of the western

Napo region of endemism, and therefore may not represent a narrow hybrid zone (Figures 2-7).

Lastly, at K=5, all members of the Guiana Shield + Northern Amazon clade (Clade D) are

inferred to be virtually unadmixed. Individuals in the far eastern Napo area of endemism on the east

bank of the Rio Putumayo (brown dots in Figure 3, 4) cluster with other Guiana Shield groups in

phylogenetic analyses but are detected to have limited (~5%) admixture with pure individuals of

Southern Amazon provenance. At higher values of K, the broad-scale population assignments

inferred at K=5 are mostly unchanged, however additional admixture components are inferred for

most groups. The introgressed western Napo clade is eventually placed into its own cluster at K=9-

10 (see Supplementary Figure 4, Supplementary Figure 8).

3.3.2. *Fine-scale population structure and admixture*

Admixture analysis using the RAD locus optimized chromosome painting approach in

RADpainter and then population assignment with fineRADstructure (Lawson et al., 2012; Malinsky

et al., 2018), was congruent with other analyses, and also detected finer partitions of population

structure. The fineRADstructure program detected 15 groups of individuals which were each

supported with a posterior probability of 100% and which also overlapped with sampling regions

which we delimited for population genetic analyses based on geographic barriers (Black dots in

Figure 3, Supplementary Figure 5). The only sampling regions which fineRADstructure failed to

confidently partition into separate groups consisted of a group of individuals in the Jaú area of

endemism, and a group of individuals in the eastern Napo area of endemism (both groups are poorly

resolved in phylogenetic analyses, but unambiguously members of the broader Northern Amazon +

Guiana Shield clade D). The eastern Napo individuals (D2 in figure 3) are identified as a low

support group ($pp = 0.49$), while the Jaú individuals are lumped in with other western Guiana Shield

individuals (D in figure 3). See Supplementary Figure 5 (fineRADstructure dendrogram), and the

full dataset coancestry matrix plots Supplementary Figures 6-7.

Patterns of coancestry show a complex mosaic of shared ancestry among populations

(Supplementary Figures 6-7). In general, coancestry patterns appear consistent with admixture

patterns detected by STRUCTURE, including the case of the highly introgressed western Napo

lineage. Therefore, we interpret patterns indicated by STRUCTURE to be reliable. Notably, the

high level of admixture initially inferred for our San Martín sample is recapitulated in RADpainter

coancestry analysis – this sample has relatively high coancestry across most populations (as does

the pattern of coancestry we report for the western Napo lineage, below).

Lastly, the Lawson et al. (2012) "normalized PCA" approach provided with

fineRADstructure captured 89% of the variance in the genetic data on the first four component axes

(51.4%, 24.9%, 7.77%, 4.94%) indicating that the coancestry matrix captures much more

information about genetic variance than standard PCoA/PCA of SNP data (Supplementary Figure

8a-c). It is therefore not surprising that K-means clustering of coancestry principle component axes

better identified groups of individuals which match previously described subspecies designations

and phylogenetic structure (Figure 5, CoA).

3.4. *Phylogenetic reticulation in the western Napo lineage*

Separate RADpainter and PhyloNet analysis of the populations in the western Napo area of

endemism and their putative progenitor lineages support the hypothesis that this region has a

complex history of introgression. One-tailed tests of mean coancestry indicated the null hypothesis

that the difference in means among groups is less than or equal to zero can be strongly rejected (ghlt

p < 2e-16), and that individuals from the western Napo population have greater median/mean

coancestry with each progenitor lineage than their progenitor lineages share with each other

(PHvGS/progenitors = ~1.078, PHvInambari/progenitors = ~1.16). These analyses establish the

asymmetry in admixture patterns detected by STRUCTURE. On average, western Napo individuals

share ~ 7.5% greater coancestry with Inambari individuals (southern Amazon) than they do with

other Northern Amazon individuals (glht p < 2e-16), indicating these sampled populations do not

include F1 hybrids. We did not observe any patterns of clinal variation within our samples of this

lineage (spanning ~20,500 km^2).

fineRADstructure analysis of the coancestry matrix, as well as visualization of the

coancestry matrix indicated that the western Napo population may be differentiated, as well as

intermediate when compared to its progenitors (Figure 6a-d). Principle components analysis (Figure

6d) of the coancestry matrix indicate that the western Napo lineage is intermediate on PC1

(compared to progenitor lineages), but differentiated on PC2, which would be expected if the Napo

lineage has had sufficient time in reproductive isolation for sorting of ancestral alleles or for the

evolution of new alleles (Barrera-Guzmán et al., 2018). This hypothesis is supported by the

observation that this lineage has distinct mitochondrial haplotypes derived from Inambari

populations (~separated from Inambari haplotypes by ~6.5 substitutions or 0.63%), as well as

analyses of IBD (below).

Bootstrapped estimates of F_{st} indicated the same general patterns: 95%CI; western Napo vs

Eastern Napo (Northern progenitor, 0.07 - 0.090), western Napo v Inambari (Southern progenitor,

0.058 - 0.077): Eastern Napo vs Inambari: (between progenitors, 0.122 - 0.144). These patterns of

F_{st} estimates were robust to alternative assignments of geographically proximate individuals to

progenitor linages (for example, including an additional adjacent population for both of the putative

progenitor lineages). Thus, the introgressed western Napo lineage is less differentiated from *each* of

its progenitor lineages than its progenitor lineages are from each other and it is *also* biased in its

genetic composition toward Inambari populations.

PhyloNet detected that an evolutionary reticulation is the best hypothesis to explain the

origin of the western Napo lineage (Figure 6a), which provides a plausible evolutionary scenario

upon which we can interpret other statistics. All parameters in the RJMCMC search achieved an

effective sample size > 150 (most >1000), and PhyloNet recovered two network topologies

representing 98.6% of the posterior distribution. The top ranked network (*pp* = 57%, Figure 6a)

implies the MRCA of the northern Amazonian (Eastern Napo + Jaú) and southern (Inambari +

Western Napo) lineages diverged first (clade F). This initial split was followed by a split within the

southern lineage that led to the Inambari (S1) and progenitor western Napo (S2a) populations

(MRCA of S1 and S2a: Figure 6a). At some point, genetically distinct nuclear genomes from the

Northern lineage (clade D/D2) introgressed into S2a (ancestral western Napo), which gave rise to

the contemporary western Napo population (S2).

The second-best network (*pp* = 41%, Supplemental Data) depicts a scenario of classic

hybrid speciation. First, northern and southern lineages of Clade F first diverged, as seen for the top

ranked network. Then, members of S1 and the NW Amazonian lineages merged to form a hybrid

lineage. We base our subsequent discussion on the top ranked network, as this hypothesis is more

consistent with patterns we detect in other analyses. Support for these two reticulation scenarios is

similar, however, and they present interesting alternatives which both require a complex history of

reticulation.

Taken in the context of STRUCTURE (Figure 5) and fineRADstructure (Figure 6) results,

the gene flow implied by the top ranked network seems to have been asymmetrical, leaving very

little evidence of southern admixture among northern lineages (see Figure 5, "weakly resolved

eastern Napo," which are detected to have very limited (< ~5%) southern admixture). This

observation, taken alone, could have been produced by sampling biases. However, demographic

parameters estimated with G-PhoCS also support this hypothesis: the only significant migration rate

(lower 95% CI > 1 migrant per generation) was inferred from the eastern Napo (member of Guiana

Shield lineage) lineage into S2/S2a (Figure 6, Supplementary Table 5).

G-PhoCS also allows us to place a preliminary maximum age constraint on the timing of

introgression into S2a. Assuming 2.47 years per generation in *Pseudopipra* (estimated in Bird et al.,

2020) and a constant mutation rate, the absolute splitting time between S2a and S1 may have been

~1.2 Ma (Figure 6, Supplementary Table 5). Though we emphasize that this is a rough

approximation, this result is loosely consistent with prior divergence time estimates for several

groups in the genus based on molecular clock analysis of mtDNA (Castro-Astor, 2014); that study

estimated ssp *pipra* (Clade D; Northern Amazon + Guiana Shield lineage) to have branched off

from its closest relatives ~ 0.67 Ma [95% HPD: 0.3506-1.1287]. Therefore, introgression into S2a

must have occurred more recently.

Lastly, patterns of admixture (Figure 5, Supplemental Figure 6, 7) suggest some evidence of

limited introgression of Northern Amazon alleles into the southwestern Amazonian Inambari

population (Figure 5). This could be a consequence of limited gene flow into the Inambari

population concurrently with or after the primary introgression event(s) which generated the

introgressed western Napo lineage. While our interpretations are conditional on whether or not our

genetic sample is indicative of a broader pattern, the product of this process appears to be a distinct,

geographically isolated and introgressed lineage within the western Napo area of endemism.

3.5. *Spatial distribution of genetic variation*

Under isolation by distance (IBD), the empirical relationship between geographic distance

and genetic distance is related to offspring dispersal distance and population size (Rousset, 1997).

Isolation by distance effects are thus predicted when individual dispersal distances are smaller than

a species' range (Teske et al., 2018). Fifteen out of eighteen considered areas (see Population

Genetics, Figure 4, and Supplemental Appendix for a description of how these areas were defined)

contained > 2 individuals, and thus could be evaluated for isolation by distance effects within them.

Five of these regions were identified to have a significant signal of IBD within them at the 0.001

alpha level (Table 2). This signal, however, was not consistent across different proxies of genetic

and geographic distance, suggesting that different processes may be contributing to spatial

correlations within these areas.

Partial Mantel tests of barriers to gene flow indicated that all but one evaluated physical

barrier likely play significant roles in structuring genetic variation between areas (Table 2). Taken

together, Mantel and partial Mantel tests suggest individuals are generally able to disperse within

localized regions (with some signal of spatial autocorrelation within regions) but are unable to cross

prominent geographic barriers delimiting these areas (Table 2). Notably, the Rio Negro was the

only evaluated barrier which is not detected to exhibit a significant effect— (Table 2 and

Supplementary Table 2).

Model based analysis of migration rates with EEMS largely corroborated the specific

hypotheses implied by Mantel tests. For dissimilarities between pairs of demes, the correlation

between observed and fitted values was high – $R^2 = 0.871$ – within demes somewhat less – $R^2 =$

0.579 (Supplementary Figures 10, 11). Thus, the EEMS model appears to do a good job of

describing spatially structured variation in this dataset, and a non-stationary migration surface is a

better fit to the data than simple isolation by distance. The EEMS visualization detects many strong

signals (posterior probability > 0.9) of barriers to gene flow (Figure 7). As expected, the Andes and

the Amazon river were unambiguously detected as the most significant barriers to gene flow and are

clearly recovered as landscape features, even though the model does not know these features exist.

Remarkably, EEMS almost perfectly infers the spatial extents of the lowland Amazonian range

(range map outlined in black), despite very limited genetic sampling toward the edge of the range.

722   EEMS detects signals of deviation from constant IBD that likely correspond to several

Amazonian tributaries in the Amazon basin which delimit areas of endemism (Figure 7). An

additional strong barrier to gene flow was inferred in the south-central amazon (south-central brown

patch in Figure 7), corresponding to the Caatinga - Cerrado – Chaco "dry-diagonal" biomes. Strong

signals of genetic connectivity are inferred in many localities (blue), notably west of the Andes,

connecting north Andean and Central American lineages (despite a lack of sampling in that region).

This link perhaps reflects the common ancestry among montane lineages in these regions.

Consistent with other analyses, EEMS detects the large Guianan + Northern Amazonian region (E.

Napo + Jaú + Guiana) to have relatively weak population structure across a variety of barriers

(including the Rio Negro). EEMS detects several discrete areas of continuous gene flow within the

Atlantic Forest, likely corresponding to the Coastal Bahia and Serra do Mar areas identified by da

Silva et al. (2004) (Figure 7). The introgressed hybrid lineage from the western Napo area is

inferred to be genetically isolated from neighboring populations, with apparent reductions in gene

flow on all sides, perhaps corresponding to the effects of the Rio Marañón in the south and Rio

Putumayo/Içá to the north.

737   Lastly, EEMS detected two broad clusters of relatively high genetic diversity (relative

heterozygosity: Supplementary Figure 12). One is centered along the Amazon river and was

relatively uniform within the northern Amazonian basin, reflecting relatively high genetic diversity

in Guiana Shield, but also some southern Amazonian populations. Notably, the other prominent

high diversity area overlaps with our sampling of the introgressed western Napo lineage. Areas of

relatively low genetic diversity included the Atlantic Forest, the Peruvian Andes, and Central

American lineages. These patterns are consistent with direct estimates of regional variation in allelic

richness (Supplementary Appendix).

3.6. *Vocal variation*

We examined 198 recordings of *Pseudopipra* and identified 14 distinct vocalizations that

can be diagnosed by ear or visual inspection of sonograms (arbitrarily labeled numerically in Figure

8). This set of vocalization types includes both advertisement vocalizations from males at leks and

calls by males and females in other contexts. After examining recordist notes, we generated a

preliminary hypothesis which distinguishes eleven highly variable and distinct, lekking

advertisements (Figure 8a) from three broad ranging call types (Figure 8b). Our delimitation of

vocalizations into advertisement and call categories partly reflects expert opinion, as the behavioral

context of vocal records was not always clear (see Supplemental Appendix). Work by Castro-Astor

et al. (2007) provides important insights of the vocal behavior of Atlantic forest populations, but

most *Pseudopipra* populations lack similarly detailed studies. Vocalization types within each

category can be statistically discriminated based on a combination of the frequency, duration, and

number of notes. PCA and logistic regression on lekking vocalizations (n=114, Supplementary

Table 3), and calls (n=33) found significant differences among types ($p < 0.001$). The first three

axes of a PCA explained ~90% of the variation in lekking vocalization characters, with PC1 (~64%)

primarily explaining variation in note number and frequency (Supplementary Figure 15).

Our assignments of vocal types to well supported genetic groups are preliminary in the

absence of recordings from the same individuals from which we have genetic samples.

Nevertheless, we found that most of the 11 lek advertisement vocalizations are phylogenetically

diagnostic of the populations in which they were recorded (except for two cases of apparent

plesiomorphy, Supplementary Figure 16b). In contrast, the three call vocalization types we

identified were diagnostic of one lineage (type 5, clade D), or shared plesiomorphically across two

ancient clades (type 10 and 13, Supplementary Figure 16a). See Taxonomic Summary, below.

4. Discussion

In comparison to the great diversity of Neotropical organisms available for study, there have

been few studies focused on understanding both fine and broad scale diversity patterns within

broadly distributed Neotropical taxa, and this has slowed the biogeographic interpretation of the

effects that many landscape features have on the evolution of Neotropical organisms (Fernandes et

al., 2013; Harvey et al., 2017; Harvey and Brumfield, 2015; Marks et al., 2002; Milá et al., 2012;

Nyári, 2007). We investigated the phylogenetic and population genetic history of a continentally

distributed taxon currently recognized as a single biological species– *Pseudopipra pipra*– using a

genomic dataset which samples thousands of loci across the genomes of 232 individuals

representing eight of thirteen *Pseudopipra* subspecies. Two additional subspecies, *pygmaea* and

coracina, were represented by mtDNA only. Our genetic sampling encompasses a majority of the

broad range of *P. pipra*, spanning Neotropical forests from Central America to the Atlantic Forest

of Brazil, including highland and lowland areas of endemism. Our sampling of vocal variation fills

in many important gaps in our genetic sampling—therefore, our intersecting datasets allow us to

assess at least some data from all thirteen previously identified subspecies.

The predominant biogeographic signal we detect in our dataset is of two successively nested

montane clades as the sister groups to several major lowland Amazonian clades (Figure 5, see

Historical Biogeography, below). This pattern implies that lowland Amazonian and montane

Central American populations of *Pseudopipra* expanded historically "out of the Andes." Ancient

lineages endemic to the Andes may have originated around the onset of the Pleistocene ~2.5 Ma

(molecular clock estimates from Castro-Astor, 2014). The lowland clade within *Pseudopipra*

originated (~1.085 [0.57-1.79] Ma) in southwestern Amazonia (Figure 4) and expanded, occupying

Amazonia on both margins of the Amazon river, and eventually reached the Atlantic forest (see

below). Therefore, our study provides an unusual example of a widespread suboscine passerine

which originated in the Andean highlands.

4.1 *Population genetic variation*

We examined our dataset using a variety of population genetic approaches intended to assess

the number of natural genetic groupings that may be present in *Pseudopipra*. Phenetic K-means

clustering of SNP data and analysis with STRUCTURE estimated a minimum of 5 genetic clusters

with broad geographic ranges (Figure 5, Supplementary Figure 2, Supplementary Figure 4). K-

means clustering of the coancestry matrix (which captured substantially more variation than

standard PCoA) more finely partitioned the dataset into 8 genetic clusters (Figure 5, Supplementary

Figure 8), which closely recapitulates previously identified subspecies boundaries (Figure 2).

Phylogenetic (RAxML, SVDquartets) and model-based population genetic (fineRADstructure)

analyses of nuclear genomic data detected fifteen identical groups of individuals which coincided

with sampling regions delimited by clear geographic barriers (results, Figure 3, 4, Supplementary

Appendix). Two additional subspecies represented by mtDNA (results) bring the total number of

distinct genetic populations delimited by geographic barriers to at least seventeen. The close

concordance observed between population structure and landscape features implies that the

evolutionary history of *Pseudopipra* within the Neotropics is deeply connected to the South

American landscape, adding support to a rich body of literature endorsing this hypothesis

(Brumfield, 2012; Cracraft and Prum, 1988; Silva et al., 2019).

Our sampling reveals a "horseshoe" pattern of genetic differentiation across the Amazon

basin (Figure 4), with no evidence of gene flow across the Amazon river, except in the western

headwaters where the Amazon is narrowest and floodplains are dynamic and young (Haffer, 1992;

Pupim et al., 2019; Räsänen et al., 1987). Given that the genus originated in the west, we

hypothesize that northern and southern clades diverged across the Amazon river in this region. For

virtually all evaluated cases, we find significant effects of geographic barriers on subdividing

genetic variation within this species complex, including across the Amazon River and most of its

larger tributaries, with the notable exception of the Rio Negro (Figure 7 and Supplementary Table

2a, 2b). As expected, the "dry-diagonal" Caatinga, Cerrado, and Chaco belt is a strong barrier to

gene flow, isolating Atlantic Forest lineages from their southeastern Amazonian Xingu relatives—

as are the Andes, which exhibit a disproportionate effect on divergence between Peruvian foothills

populations and Central American lineages (see Supplementary Appendix for additional

discussion).

Multiple analyses indicate that the western Napo population has a complex history of initial

and early differentiation from the broader Southern Amazonian clade, followed by subsequent,

asymmetric genetic introgression from Jaú and eastern Napo populations of the Guianan Shield +

Northern Amazon clade (Clade D; see discussion below). When the genetic data are analyzed under

either concatenated or coalescent phylogenetic frameworks which do not allow for phylogenetic

reticulation, the western Napo population is reconstructed as most closely related to Guiana Shield

+ Northern Amazon populations (Figure 3, Figure 5, Clade D). In contrast, phylogenetic network

analysis indicates that this population is derived from individuals of Inambari (southern Amazon)

provenance (Figure 6, also summarized in Figure 4). This incongruence between methods is likely

due to the complexity of the constituent gene trees in this population-level differentiation induced

by asymmetric introgression from the Jaú and eastern Napo populations (Thom et al., 2018). We

hypothesize that introgression has introduced more recently derived variation from the Guiana

Shield + Northern Amazon clade D into an older, differentiated, population of the Southern

Amazonian clade, pulling the phylogenetic affinity toward the northern clade.

4.2 *History of an introgressed hybrid lineage*

The substantial level of observed introgression (>40%) in the western Napo individuals, as

well as a suite of genomic and phenotypic evidence, suggests that this genetic population may be

best recognized as a hybrid lineage, though it is uncertain if it can be considered a product of hybrid

speciation. To date, five prior examples of hybrid species formation in birds have been evaluated

with genomic data: (1) the Golden-crowned manakin *Lepidothrix vilasboasi* (Barrera-Guzmán et al.,

2018), (2), the Audubon's warbler *Setophaga auduboni* (Brelsford et al., 2011), (3) the Italian

sparrow *Passer italiae* (Hermansen et al., 2011), (4) the Hawaiian duck *Anas wyvilliana* (Lavretsky

et al., 2015), and (5) a putative species of Darwin's finches *Geospiza* (Lamichhaney et al., 2018).

These examples of mosaic genome hybrid speciation, in which the fusion of two or more progenitor

lineages leads to a reproductively isolated hybrid species (Counterman, 2016; Jiggins et al., 2008),

repeatedly demonstrate that modern patterns of avian diversity may have been significantly

influenced by historical phylogenetic reticulation (Suh, 2016). Our analyses of genomic data across

Pseudopipra imply that the western Napo lineage is likely a differentiated, allopatric phylogenetic

species which has a genomic composition derived from heavy historical admixture with a non-sister

lineage. This historical scenario is not the same as those examples cited above, but it has produced a

population genetic signature which is extremely similar (compare to Barrera-Guzmán et al., 2018).

These observations force us to ask whether historical process, or contemporary genetic pattern

should be paramount in delineating a hybrid species.

Some authors have proposed strict population genetic criteria for identifying homoploid

hybrid speciation, including demonstrating (1) evidence of reproductive isolation from parental

species, (2) evidence of past hybridization, and (3) evidence that intrinsic reproductive isolation was

derived from the hybridization event (Schumer et al., 2014). The first two conditions are satisfied

by the western Napo *Pseudopipra* lineage. First, phylogenetic and population genetic analyses are

suggestive of contemporary reproductive isolation. Phylogenetic analyses of nuclear genome data,

including concatenated and coalescent analyses (Figure 3, 4, 6), recover this lineage as a well-

supported reciprocally monophyletic group. Differentiated mtDNA haplotypes from a broader

geographic sample (Figure 6) also form subgroups derived from southern Inambari haplotypes

(results, Supplementary Figure 17), implying reproductive isolation from female northern

progenitors at a minimum.

PCA-based and model-based population genetic cluster analyses (Figure 6b, 6c,

Supplementary Figures 4-9), help establish this lineage as intermediate *and* differentiated when

compared to the living descendants of its inferred progenitors. The level of differentiation on PC2

corresponds to about 10% of the total variation explained by PC1 and PC2 (~85%). At the same

time, Mantel tests, as well as model-based analysis of isolation by distance, support the hypothesis

that contemporary gene flow into and out of this population is at least restricted and probably not

occurring (Figure 7, Supplementary Figure 12). Thus, we interpret our G-PhoCS results of a

significant migration parameter (results) as indicative of a window of historical introgression, which

could be further explored with simulation studies. Estimates of inbreeding coefficients support this

hypothesis and suggest similar levels of inbreeding among *Pseudopipra* populations in the western

Amazon ($F_{IS} > 0.1$, Supplementary Figure 9, and Supplementary Appendix). The western Napo

population is thus not detected to be more outbred than most populations (i.e. recently introgressed).

In sum, these patterns observed in our genetic data are consistent with the hypothesis of

contemporary reproductive isolation for the introgressed Western Napo populations.

The second criterion from Schumer et al. (2014), evidence of past hybridization, is directly

supported by multiple model-based analysis of genomic admixture and phylogenetic reticulation

(Figure 5, 6), and indirectly supported by model based and phenetic cluster analysis (Supplementary

Figures 4-9). We hypothesize that hybridization initially occurred between progenitor lineage S2a

and western lineages of the Guiana Shield + Northern Amazon clade (Figure 6a). Lineage S2a was

then replaced by the novel introgressed lineage, which exchanged genes with neighboring

populations within a finite time window.

The third criterion from Schumer et al. (2014), evidence that intrinsic reproductive isolation

was derived from the hybridization event, is recognized as the most difficult to demonstrate, and

probably the most tenuous. As discussed by Nieto Feliner et al. (2017) in critique of Schumer et al.

(2014), this criterion may be useful for framing the strength of evidence for or against the existence

of a homoploid hybrid species, but only if one adheres strictly to the biological species concept

(Mayr, 1942). If one applies a more evolutionary definition of species, like the metapopulation

lineage concept, in which the only necessary requirement is demonstration of separately evolving

lineages (de Queiroz, 2005), the importance of demonstrating intrinsic reproductive isolation to

claim species status for a putative hybrid lineage (or any lineage at all) is less clear to us. Speciation

in birds in particular is thought to proceed relatively fast, and may be more affected by external

factors than internal ones (Grant and Grant, 1997). Further, requiring hybridization to be the *source*

of intrinsic isolation excludes the possibility that extrinsic or pre-zygotic isolation may evolve from

changes that enable hybrids access to new niches, which may also be geographically isolated

relative to parental forms (Nieto Feliner et al., 2017).

Intriguingly, the evolutionary scenario implied by phylogenetic reticulation analysis for the

introgressed western Napo hybrids provides a mechanism to explain a confusing distribution of

vocalization types (Figure 8, Supplementary Appendix). The males from the Napo area of

endemism share vocalization type 2 with birds in the southeastern Amazon (Figure 8,

Supplementary Figure 16), and with the restricted Huallaga valley *pygmaea* subspecies (the sister

group to all lowland Amazonian forms, according to mtDNA). Manakins are suboscine passerines

and have innate vocalizations (though see Saranathan et al., 2007). Therefore, the introgressed Napo

lineage appears to have inherited its vocalizations from the MRCA of groups S1 and S2 (Figure 6a)

and retained it through subsequent introgression with northern lineages that have a different vocal

type (Type 3 and/or 5). In this scenario, vocal type 7 in the western Inambari represents more

recently derived variation in the southern Amazonian clade (Figure 8, Supplementary Figure 16).

As noted earlier, the western Napo lineage also has mtDNA haplotypes derived from

Inambari populations (Supplementary Figure 17). In addition to implying reproductive isolation

from the Guiana Shield clade, this mitochondrial pattern implies that the introgressed Napo lineage

may be descended from the union of progenitor female S2a and progenitor western males of the

Guiana Shield + Northern Amazon clade, perhaps through male biased dispersal or mate choice (a

preliminary test of male biased dispersal did not detect a signal, Supplemental Appendix).

Regardless, this pattern raises fascinating questions about why and how the western Napo lineage

may have retained its southern song type.

Taking all of the available genetic and phenotypic evidence into account, we suggest that the

western Napo lineage may be an additional example of the formation of a hybrid lineage in birds,

and one that was produced via an underappreciated historical mechanism. Assuming that our

limited sampling is representative of the *discolor* subspecies, the patterns we observe appear to be

most similar to "hybrid trait speciation," in which a hybrid species is formed after introgression

from one species into a genomic background of a close relative (Counterman, 2016; Jiggins et al.,

2008; Marques et al., 2019). This type of combinatorial hybrid speciation was originally proposed

from studies of *Heliconius* butterflies (Jiggins et al., 2008; Salazar et al., 2010).

4.3 *Comparative and Historical Biogeography*

The spatial pattern of population structure and evolutionary relationships we identify here

are broadly congruent with other recent studies of Neotropical biogeography in upland forest birds,

contributing another example of the substantial phylogeographic pseudocongruence across taxa at

this wide scale (e.g. Cunningham and Collins, 1994; Harvey et al., 2017). As expected, the two

primary barriers identified for *Pseudopipra* in the South American lowlands are the Amazon River

and the dry diagonal. Amazonian tributaries seem to delimit recently isolated populations in

southern Amazonia, while northern *Pseudopipra* populations show weaker evidence of being

isolated by the Negro and Japurá rivers. By contrast, southern populations seem to be strongly

isolated by rivers with smaller discharge, such as the Xingu, Tapajós and Purus.

A survey of the suboscine ovenbirds in the *Xenops* complex (Harvey and Brumfield, 2015)

identified similar patterns of genetic structuring coincident with major Amazonian tributaries, as did

Ribas et al. (2012) in the *Psophia* trumpeters, though the inferred relationships among areas of

endemism are not the same. A recent comparative analysis shows that large rivers delimit genetic

clusters in as many as 23 groups of upland Amazonian forest birds, strongly arguing in favor of

these rivers acting as barriers to gene flow (Silva et al., 2019). The somewhat distinct pattern found

for *Pseudopipra* may be related to its recent expansion throughout the eastern Andean lowlands and

its habitat tolerance.

Within the manakins, Ohlson et al. (2013) placed *Pseudopipra* as sister to the genus

Ceratopipra, which includes five well-recognized species that are extensively codistribtuted with

Pseudopipra. The montane origin of *Pseudopipra* is congruent with the observation that

Pseudopipra is the sister group to *Ceratopipra,* which is also broadly distributed in the Neotropical

lowlands from Central America to the Atlantic Forest, with two secondary expansions into the

subtropical forests in the tepuis (*C. cornuta*) and the southern Andes (*C. chloromeros*) (Ohlson et

al., 2013). The breakpoints among these *Ceratopipra* species are highly concordant with the

breakpoints among the genetic clusters within the *Pseudopipra* complex that we have presented

here, implying that these closely related taxa have components of their phylogeographic history in

common. See the Supplemental Appendix for additional notes, as well as Castro-Astor (2014).

A key feature of the phylogeny of *Pseudopipra* is that montane Andean lineages are sister

groups to both the montane Central American and the lowland Amazonian-southeast Brazilian

lineages (Figure 5). Ancestral character reconstruction under Bayesian stochastic mapping and

maximum parsimony therefore unambiguously reconstructs the most recent common ancestor of

Pseudopipra as Andean (Figure 3-5, Supplemental Appendix). Thus, it appears that the most recent

common ancestor of *Pseudopipra* was an Andean lineage restricted to subtropical, lower montane

forest. The earliest diversification event in the genus was likely the differentiation of the ancestral

Andean populations north and south of the headwaters of the Rio Huallaga, Peru. The northern

lineage gave rise to the subtropical montane lineages of the northern Andes and Central America

(Clade A). The southern lineage gave rise to a subtropical montane Southern Peruvian lineage

(Clade B) and a southwestern lowland Amazonian lineage (*pygmaea*, represented in our dataset by

mtDNA). Subsequently, the Amazonian lineage expanded to the eastern lowlands and differentiated

into the northern lowland Amazonian + Guiana Shield (Clade D in Figure 3) and southern lowland

Amazonian/Atlantic Forest (Clade C in Figure 3) clades. Within the Southern Amazonian clade,

differentiation proceeded from west to east: (Figure 3, 4):

(West Inambari (East Inambari (Rondônia (Tapajós (Xingu (Bahia (Espírito Santo (Rio de Janerio)))))))).

Contrastingly, the northern Amazonian lineage also expanded all the way to the Atlantic Ocean, but

without producing strong phylogenetic structure (Figures 3, 4). In the Napo area of endemism,

substantial introgression occurred from east Napo and Jaú populations into the western Napo

population, with limited backcrossing into Inambari and east Napo populations.

Intriguingly, a well-supported distinct montane (>1000m) (BS: 100) (Clade C2 in Figure 3;

Cushabatay area, northern Cordillera Azul, Loreto Prov., Peru) is recovered as nested within the

well supported lowland western Inambari subclade (Clade C1 in Figure 3). Thus, there may have

been at least one secondary invasion of montane Andean habitats from a lowland ancestor. Patterns

of admixture support this hypothesis – montane individuals from the northern Cordillera Azul and

their proximate lowland relatives all show limited genomic admixture from derived lineages of the

Guiana Shield + Northern Amazon clade (yellow in Figure 5). This result is consistent with a

scenario of secondary invasion of the highlands from lowland Inambari lineages after weak

introgression from the western Napo (see below).

4.4 *North Andean diversity*

Our final genomic data set does not include any samples from Andean Ecuador or

Colombia, a region that includes four previously recognized subspecies and at least four distinct

vocalization types (see Taxonomic Summary). Unfortunately, the three Andean Ecuadorian tissue

(representing subspecies *coracina*) specimens in our original sample were not of high enough

preservation quality to be viable for ddRAD sequencing (Supplementary Table 1). We were able to

obtain mtDNA from these samples however, which unambiguously formed a monophyletic cluster

with our single San Martín (North Andean Peru) specimen in mtDNA gene tree analyses

(Supplemental Appendix, Supplementary Figure 17). A prior study by Castro-Astor (2014) based

on mtDNA also included one Andean Ecuadorian sample and one Central American sample, which

unambiguously clustered together (*pp*=100). Thus, it is likely that our unsampled Andean lineages

of Ecuador and Colombia (including the subspecies *minimus, bolivari, unica,* and *coracina*) are

likely members of the northern Andean clade represented in our genomic data set by our single San

Martín specimen. As noted earlier, our San Martín specimen had approximately equal probability of

assignment to multiple populations, which could indicate heavy admixture, or that this is a single

sample from a highly distinct population which has its own distinct history. These patterns may also

be consistent with the hypothesis that the North Andean clade is the source of montane and lowland

Pseudopipra diversity.

1010

4.5 *Phenotypic and Vocal Evolution*

Thirteen subspecies have been previously described based on variations in plumage color

and, to a lesser extent, size (See Taxonomic Summary in Supplementary Appendix). Male plumage

coloration varies among the subspecies of *Pseudopipra* in the glossiness of the black body feathers,

the length of the white crown feathers, and the color– white, gray, or black– of the bases of these

crown feathers. Female plumage coloration often shows more striking differentiation among

subspecies than does male plumage, including variation in the shade of olive green on the body, the

shade and extent of gray on the crown and face, and olive, yellow, or gray coloration on the belly

(Chapman et al., 1914; de Schauensee, 1945, 1950; Ridgway, 1906; Wetmore, 1972; Zimmer,

1936).

Although we did not conduct a detailed analysis of plumage variation among populations of

Pseudopipra, we did observe study skin specimens of all but one (*bolivari*) of the previously

described subspecies in museum collections. All of the ten subspecies that were included in our

genetic samples were identified as distinct, diagnosable, monophyletic groups. In other words, in all

cases that we were able to test, traditional taxonomic practices conducted between 1758 through

1936 successfully identified and named distinct evolutionary lineages within *Pseudopipra*. We did

not have genetic samples of the northern Andean taxa *minima, unica,* and *bolivari*, but we note that

they all share entirely white crown feather bases (forecrown feathers only in *bolivari*)(de

Schauensee, 1945, 1950), implying that this unique plumage character state is derived within

Pseudopipra, and that *minima*, *unica*, and *bolivari* form a clade. One lineage containing two

subspecies– *separabilis* and *cephaleucos*- has evolved another unique, shared, derived plumage

character– a distinctive, second, predefinitive male plumage which has also further differentiated

between the two subspecies. In *cephaleucos,* predefinitive males have an olive green back, a white

crown, and slate gray face and belly. In *separabilis,* predefinitive males are similar with lighter gray

belly, and a medium gray, instead of white, crown. In conclusion, plumage coloration appears to

provide highly informative evidence of evolutionary lineage status in this genus.

Our analysis of vocal behavior indicates that vocalizations are also highly informative of

lineage identity within *Pseudopipra*. The complex phylogenetic history we have documented with

genetic data has clearly had a strong impact on the evolution of vocalizations in *Pseudopipra.*

Because lek vocalizations are a focus of female choice in manakins and because suboscine

passerines have largely innate vocalizations, the extensive vocal and genetic differentiation among

populations of *Pseudopipra* strongly indicates the existence of many species within *Pseudopipra*.

We identified fourteen distinct vocalization types (Figure 8), and nearly all were restricted to

and diagnostic of a single previously recognized subspecies or a broader monophyletic group (See

Expanded Taxonomic Summary in Supplementary Appendix for details). One of the two exceptions

was vocalization type 2, which may be shared plesiomorphically across the Amazon basin between

populations of *pygmaea* in the Huallaga Valley, *discolor* in the western Napo region of Ecuador and

Peru, and *separabilis* in Para, Brazil, but not in the intervening southwestern Amazonian

populations of *microlopha* (Supplementary Figure 16). Unfortunately, available vocal sampling is

very limited in the intervening regions along the south side of the Amazon Basin, which are

populated by a nested series of genetically distinct lineages with successively closer relationships to

southern Amazonian *separabilis* and *cephaleucos* from the Atlantic forest of Brazil.

1053   The other exception is the sharing of vocal type 1 ('*trill-dink*') between populations of

occulta from the eastern slopes of the Andes from San Martin, Peru to southeastern Ecuador (A1)

and populations of *comata* from the southern Cordillera Azul in Loreto (B1). The song of the

southern clade (B2) from southern Huánuco (AMNH 820866, 820952), Pasco, Junín, and Cusco is

currently unknown. Surprisingly, these geographically adjacent lineages sit on either side of the

most basal phylogenetic separation within *Pseudopipra*. The *comata* populations are

phylogenetically closer to northern Andean and Central American lineages that exhibit substantial

vocal diversity. Likewise, the southern Cordillera Azul populations are more closely related to other

south Peruvian and Amazonian populations which also exhibit great vocal diversity. This result

implies that the song type 1 was shared by the ancestral lineage and has been retained in these

lineages (Supplemental Figure 16). We identified the same pattern for call vocalization type 10.

1064   Our phylogenetic analysis also identified a distinct montane clade from the northern

Cordillera Azul (Clade C2; Cushabatay area; Alverson et al. 2001) which is closely related to

populations from lowland forests of eastern Peru south of the Rio Marañón and east to Rio Purus,

Brazil, currently recognized as *P. p. microlopha*. The hypothesis that this lineage is a secondary

expansion into montane forests from the nearby Amazonian populations is strongly supported by

the evidence that this montane population shares same song type 7 ('*jeer*') with the lowland

populations of *microlopha* (ML-190390, ML-190388, and ML-190262).

1071

4.6 *Taxonomy and Revised Classification*

The present set of analyses demonstrate a number of genetically well-differentiated and

phenotypically diagnosable lineages and provides compelling new evidence for evaluating the

species status of *Pseudopipra*. Although there are gaps in our sampling, we find that there are at

least eight genetically well-differentiated, and phenotypically diagnosable lineages of *Pseudopipra*,

and this is reflected in our proposed taxonomic revision below. Nonetheless, it should be noted that

there is more genetic variation than observed vocal variation implies, and more vocal variation

beyond our geographic coverage of genetic samples (especially in northern Andes). Thus, our

taxonomic recommendations for *Pseudopipra* almost certainly represent an underestimate of extant

diversity, underscoring the need for continued reassessment of species limits and of diversity

patterns in the Neotropics.

Our analysis provides the first comprehensive opportunity to reevaluate species limits within

Pseudopipra based on phylogenetic, population genetic, vocal, and plumage differentiation among

populations and named subspecies (See the Expanded Taxonomic Summary in the Supplementary

Appendix). This proposed classification is a conservative treatment that recognizes the limitations

on our genetic sampling of populations in the northern Andean clade from Ecuador and Colombia.

Four of these unsampled (and under-sampled) northern Andean subspecies – *coracina*, *minima*,

bolivari, and *unica* – have unique and highly differentiated vocalization types, and diagnosable

plumage differences. Our phylogenetic results strongly suggest that most of the *Pseudopipra*

subspecies are distinct evolutionary lineages deserving species status under either the phylogenetic

or biological species concepts. In sum, *Pseudopipra* may include 15-17 distinct species that have

rapidly arisen in the last ~2.5 Ma, but further reclassification will require more vocal data and

detailed analyses of plumage variation.

1095

1096   **5. Taxonomic Summary**

1097

Pseudopipra coracina (Sclater 1856)   **Andean White-crowned Manakin**

Distribution: Subtropical Andes from Venezuela south to Esmeraldas, Ecuador and San Martín,

Peru.

Phylogenetic Position: Clade A1

P. c. coracina (Sclater 1856)

Type Locality: Villavicencio, Meta, Colombia.

Distribution: Subtropical forests of the eastern slope of the Andes from western Venezuela to

Morona-Santiago, Ecuador.

Phylogenetic Position: (mtdna) Member of Clade A1

Lek Vocal Type: 8 (*errrwer*).

Call Vocal Type: Unknown

1109

P. c. minima (Chapman 1914)

Type Locality: West of Popayan, Cauca, Colombia

Distribution: Subtropical forests of western Cauca, Colombia south to Esmeraldas, Ecuador

Phylogenetic Position: (unsampled) Member of Clade A/A1

Lek Vocal Type: 9 (*reeee*)

Call Vocal Type: unknown

1116

P. c. bolivari (de Schauensee 1950)

Type Locality: Murucucu, Bolivar, Colombia

Distribution: Subtropical forests of southern Córdoba, Colombia. (Not Sampled)

Phylogenetic Position: (unsampled) Member of Clade A1

Lek Vocal Type: Unknown

Call Vocal Type: Unknown

1123

P. c. unica (de Schauensee 1945)

Type Locality: Lomas de Isnos, Huila, Colombia

Distribution: Subtropical forests of Magdalena Valley, Antioquia to Huila, Colombia.

Phylogenetic Position: (unsampled) Member of Clade A1

Lek Vocal Type: 11a (*weer-dink*) and 11b (*shureeep*)

Call Vocal Type: Unknown

1130

P. c. occulta (Zimmer 1936)

Type Locality: Ucho, east of Chachapoyas, Amazonas/San Martin, Peru

Distribution: Eastern slope of the Andes from Zamora-Chinchipe, Ecuador (Freile, 2014)

south to San Martín, and Huánuco, Peru, west of the Rio Huallaga

Phylogenetic Position: Member of Clade A/A1

Lek Vocal Type: 1 (*trill-dink*)

Call Vocal Type: 10 (*bree*)

1138

Pseudopipra anthracina (Ridgway 1906)   **Western White-crowned Manakin**

Type Locality: Moravia, Costa Rica

Distribution: Subtropical Costa Rica to Western Panama

Phylogenetic Position: Clade A2

Lek Vocal Type: 4 (*jureeee*)

Call Vocal Type: Unknown

1145

Pseudopipra comata (Berlepsch and Stolzmann 1894)   **Junín White-crowned Manakin**

Type Locality: Garita del Sol, Junin, Peru.

Distribution: Subtropical Andes of Peru from Cordillera Azul, Loreto (east and south of the Rio

Huallaga) to southern Huánuco Pasco, Junín, and northern Cusco.

Phylogenetic Position: Clade B

Lek Vocal Type: 1 (*trill-dink*)

Call Vocal Type: 10 (*bree*)

1153

Pseudopipra pygmaea (Zimmer 1936)   **Huallaga White-crowned Manakin**

Type Locality: Chamicuros, Loreto, Peru

Distribution: Tropical forest of lower Rio Huallaga Valley, Peru

Phylogeneic Position: (mtDNA) Sister to Clade F

Lek Vocal Type: 2 (*deeeer*)

Call Vocal Type: 13 (*weeu*)

1160

Pseudopipra discolor (Zimmer 1936)   **Napo White-crowned Manakin**

Type Locality: Puerto Indiana, Loreto, Peru

Distribution: Tropical forest in Napo, Ecuador and Loreto, Peru south to the Rio Marañón.

Phylogenetic Position: Clade E

Lek Vocal Type: 2 (*deeeer*)

Call Vocal Type: 13 (*weeu*)

1167

Pseudopipra pipra (Linneaus 1758)   **Northern White-crowned Manakin**

Type Locality: Suriname

Distribution: Tropical forest of eastern Colombia, southern Venezuela, the Guianas, and Brazil

north of the Amazon and west to the right (west) bank of the Rio Putumayo, Colombia.

Phylogenetic Position: Clade D

Lek Vocal Type: 3 (*buzzzz*)

Call Vocal Type: 5 (*zeee*)

1175

1176 *Pseudopipra microlopha* (Zimmer 1929)　　　　**Southern White-crowned Manakin**

1177 **Distribution**: Tropical forest of eastern Peru south of the Rio Huallaga, Rio Marañón, and the

1178 Amazon River east to Pará, Brazil, and subtropical forests between the Rio Huallaga and Rio

1179 Ucayali.

1180 **Phylogenetic Position:** Paraphyletic, including Clade C without Clade C7

1181

P. m. **undescribed subspecies**

Distribution: Subtropical forest from the Cushabatay area, northern Cordillera Azul, between

Rio Huallaga and Rio Ucayali, Loreto Province, Peru.

Phylogenetic Position: Clade C2

Lek Vocal Type: 7 (*jeer*)

Call Vocal Type: 13 (*weeu*)

1188

P. m. microlopha (Zimmer 1929)

Distribution: Eastern Peru south of the Rio Marañón and Rio Huallaga, west to Rio Juruá and

Rio Purus, Brazil.

Phylogenetic Position: Clade C1 excluding C2

Lek Vocal Type: 7 (*jeer*)

Call Vocal Type: 13 (*weeu*)

1195

P. m. **undescribed subspecies**

Distribution: Right (east) bank of the Rio Purus to the left (west) bank Rio Madeira

Phylogenetic Position: Clade C3

Plumage: Not examined

Lek Vocal Type: Unknown

Call Vocal Type: 13 (*weeu*)

1202

P. m. **undescribed subspecies**

Distribution: Right (east) bank of the Rio Madeira to the left (west) bank the Rio Tapajos.

Phylogenetic Position: Clade C4

Plumage: Not examined

Lek Vocal Type: Unknown

Call Vocal Type: 13 (*weeu*)

1209

P. m. **undescribed subspecies**

Distribution: Right (east) bank of the Rio Tapajos to the left (west) bank of the Rio Xingu

Phylogenetic Position: Clade C5

Plumage: Not examined

Lek Vocal Type: 6b (*zeeeee-tonk*)

Call Vocal Type: 13 (*weeu*)

1216

P. m. separabilis (Zimmer 1936)

Type Locality: Tapara, Rio Xingu, Brazil

Distribution: Tropical forest of Rio Xingu east to central and southern Pará.

Phylogenetic Position: Clade C6

Lek Vocal Type: 2 (*deeeeer*)

Call Vocal Type: 13 (*weeu*)

Pseudopipra cephaleucos (Thunberg 1822) **Atlantic White-crowned Manakin**

Type Locality: Bahia, Brazil

Distribution: Tropical forest from Bahia south to northern Rio de Janeiro, Brazil. **Phylogenetic**

Position: Clade C7

Lek Vocal Type: 6a (*zeeeee-tonk*)

Call Vocal Type: 13 (*weeu*)

1229

1230   **Supplementary Material:** Supplementary Data, including data files and R code can be found in the

1231   Mendeley Data Repository [temporary link for review only]:

1232   https://data.mendeley.com/datasets/...

1233   and GenBank XXX-XXX.

1234

Funding: J. S. B. was supported by a National Science Foundation Graduate Research Fellowship,

NSF Doctoral Dissertation Improvement Grant [DGE-1650441, DEB-1700786], and University of

Michigan Life Sciences Fellowship. L. C. was supported by NSF grant DEB-1555754 to I.J.L. T. J.

F. was supported by an NSF Post-doctoral Fellowship in Biology [DBI-1523857]. C. C. R. was

funded by CNPq (308927/2016-8), FAPEAM, FAPESP/NSF (NSF DEB-1241066, FAPESP, grant

#2012/50260-6) and USAID (PEER AID-OAA-A-11-00012, cycle 5). J. S. B. and R. O. P. were

funded by the W. R. Coe Fund of Yale University. Data collection was paid for by Athena Funds

from the Cornell Laboratory of Ornithology, and W. R. Coe Fund from Yale University. This work

was also supported in part by the facilities and staff of the Yale University Faculty of Arts and

Sciences High Performance Computing Center, and with the resources of the Cornell University

BRC Bioinformatics Facility, which is partially funded by Microsoft Corporation.

1246

1247 Acknowledgements:

We thank the numerous researchers whose field collection efforts made this research possible

including: Angelo P. Capparella, Camila Ribas, Ivandy Castro-Astor, Cecilia Fox, David E.

Willard, Diego Ocampo, Donna Dittmann, Jacob S. Berv, John P, O'Neill, Karla Conejo, Katya

Balta, Kenneth V Rosenberg, Louise M. Augustine, Peter E. Scott, Richard O. Prum, Steven W.

Cardiff, Susan Allen-Stotz, Thomas Valqui, Tristan J. Davis, Ivan Prates and others. Tissue loans or

skins were kindly received from the Divisão de Aves do Museu de Zoologia da Universidade

Estadual de Feira de Santana (MZFS), the Field Museum of Natural History (FMNH), Instituto

Nacional de Pesquisas da Amazônia (INPA), Kansas University Natural History Museum

(KUNHM), Louisiana Museum of Natural History (LSUMZ), Museu de Zoologia da Universidade

de São Paulo (MZUSP), Museu Paraense Emilio Goeldi (MPEG), the Academy of Natural Sciences

of Drexel University (ANSP), Universidade do Estado do Rio de Janeiro (UERJ), the University of

Alaska Museum (UAM), the University of Nevada, Las Vegas (UNLV), the Yale Peabody Museum

of Natural History (YPM), the Cornell Museum of Vertebrates (CUMV), the American Museum of

Natural History (AMNH), the Museum of Comparative Zoology (MCZ), and the United States

Museum of Natural History (NMNH). We thank the curators and collections managers of these

institutions for facilitating these loans: Camila Ribas, John Bates, Mark Robbins, Donna Dittman,

Alexandre Aleixo, Caio Graco Machado, Joel Cracraft, Luís Fábio Silveira, Kevin Winker, Maria

Alice Santos Alves, Kristof Zyskowski, Nate Rice, Charles Dardia, John Klicka, and others. We

thank Eliot Miller, Emma Greig, Matthew Medler, Greg Budney, Eduardo Iñigo and Andrew

Spencer at the Macaulay Library for assistance with audio recordings. We thank members of the

Lovette Lab, Daniel Lane and **XXX** anonymous reviewers, for comments on the research and the

manuscript. The illustration depicted in Fig. 1 is reproduced with permission from the Handbook of

Birds of the World, Lynx Editions, Barcelona. J. S. B. and R. O. P. conceived of the research. J. S.

B. and L. C. performed laboratory and bioinformatic processing, with input and support from I. J. L.

We thank Bronwyn Butcher for help with laboratory protocols. J. S. B. and L. C. analyzed the

genomic data and interpretation of the results was supported by input from all authors. T. J. F.

downloaded, processed, and analyzed lekking vocalization data. C. C. R. and I. C. R. facilitated

critical sample acquisitions from Amazonia and the Atlantic Forest. I. J. L. and R. O. P. provided

detailed guidance and feedback on the manuscript. R. O. P. examined plumage phenotypes and

proposed a revised taxonomy. The manuscript was written by J. S. B. with input from all authors.

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289 **Literature Cited**
1290

Antonelli, A., Ariza, M., Albert, J., Andermann, T., Azevedo, J., Bacon, C., Faurby, S., Guedes, T.,

Hoorn, C., Lohmann, L.G., Matos-Maraví, P., Ritter, C.D., Sanmartín, I., Silvestro, D., Tejedor, M.,

ter Steege, H., Tuomisto, H., Werneck, F.P., Zizka, A., Edwards, S.V., 2018. Conceptual and

empirical advances in Neotropical biodiversity research. PeerJ 6, e5644.

Barrera-Guzmán, A.O., Aleixo, A., Shawkey, M.D., Weir, J.T., 2018. Hybrid speciation leads to

novel male secondary sexual ornamentation of an Amazonian bird. Proceedings of the National

Academy of Sciences 115, E218-E225.

Bates, J.M., Zink, R.M., 1994. Evolution into the Andes: Molecular Evidence for Species

Relationships in the Genus Leptopogon. The Auk 111, 507-515.

Berv, J.S., Prum, R.O., 2014. A comprehensive multilocus phylogeny of the Neotropical cotingas

(Cotingidae, Aves) with a comparative evolutionary analysis of breeding system and plumage

dimorphism and a revised phylogenetic classification. Molecular Phylogenetics and Evolution 81,

120-136.

Bioacoustics_Research_Program, 2011. Raven Pro: Interactive Sound Analysis Software (Version

1.4) [Computer software]. The Cornell Laboratory of Ornithology. Available from

http://www.birds.cornell.edu/raven., Ithaca, NY.

Bird, J.P., Martin, R., Akçakaya, H.R., Gilroy, J., Burfield, I.J., Garnett, S., Symes, A., Taylor, J.,

Şekercioğlu, Ç.H., Butchart, S.H.M., 2020. Generation lengths of the world's birds and their

implications for extinction risk. Conservation Biology n/a.

Borges, S.H., Da Silva, J.M.C., 2012. A New Area of Endemism for Amazonian Birds in the Rio

Negro Basin. The Wilson Journal of Ornithology 124, 15-23.

Bradburd, G.S., Coop, G.M., Ralph, P.L., 2018. Inferring Continuous and Discrete Population

Genetic Structure Across Space. Genetics 210, 33-52.

Brelsford, A., Milá, B., Irwin, D.E., 2011. Hybrid origin of Audubon's warbler. Molecular Ecology

20, 2380-2389.

Brumfield, R.T., 2012. Inferring the origins of lowland Neotropical birds. The Auk 129, 367-376.

Brumfield, R.T., Edwards, S.V., 2007. EVOLUTION INTO AND OUT OF THE ANDES: A

BAYESIAN ANALYSIS OF HISTORICAL DIVERSIFICATION IN THAMNOPHILUS

ANTSHRIKES. Evolution 61, 346-367.

Brumfield, R.T., Liu, L., Lum, D.E., Edwards, S.V., 2008. Comparison of Species Tree Methods for

Reconstructing the Phylogeny of Bearded Manakins (Aves: Pipridae, Manacus) from Multilocus

Sequence Data. Systematic Biology 57, 719-731.

Cabanne, G.S., Campagna, L., Trujillo-Arias, N., Naoki, K., Gómez, I., Miyaki, C.Y., Santos, F.R.,

Dantas, G.P.M., Aleixo, A., Claramunt, S., Rocha, A., Caparroz, R., Lovette, I.J., Tubaro, P.L.,

2019. Phylogeographic variation within the Buff-browed Foliage-gleaner (Aves: Furnariidae:

Syndactyla rufosuperciliata) supports an Andean-Atlantic forests connection via the Cerrado.

Molecular Phylogenetics and Evolution 133, 198-213.

Campagna, L., Gronau, I., Silveira, L.F., Siepel, A., Lovette, I.J., 2015. Distinguishing noise from

signal in patterns of genomic divergence in a highly polymorphic avian radiation. Molecular

Ecology 24, 4238-4251.

Campagna, L., St. Clair, J.J.H., Lougheed, S.C., Woods, R.W., Imberti, S., Tubaro, P.L., 2012.

Divergence between passerine populations from the Malvinas – Falkland Islands and their

continental counterparts: a comparative phylogeographical study. Biological Journal of the Linnean

Society 106, 865-879.

Capurucho, J.M.G., Cornelius, C., Borges, S.H., Cohn-Haft, M., Aleixo, A., Metzger, J.P., Ribas,

C.C., 2013. Combining phylogeography and landscape genetics of Xenopipo atronitens (Aves:

Pipridae), a white sand campina specialist, to understand Pleistocene landscape evolution in

Amazonia. Biological Journal of the Linnean Society 110, 60-76.

Carstens, B.C., Pelletier, T.A., Reid, N.M., Satler, J.D., 2013. How to fail at species delimitation.

Molecular Ecology 22, 4369-4383.

Castro-Astor, I.N., 2014. A Phylogeographic study of two neotropical bird species, the Red-Headed

Manakin (Pipna rubrocapilla) and the White-crowned Manakin (Dixiphia pipra). Biology. CUNY,

CUNY Academic Works.

Castro-Astor, I.N., Alves, M.A.S., Cavalcanti, R.B., 2007. DISPLAY BEHAVIOR AND SPATIAL

DISTRIBUTION OF THE WHITE-CROWNED MANAKIN IN THE ATLANTIC FOREST OF

BRAZIL. The Condor 109, 155-166.

Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A., Cresko, W.A., 2013. Stacks: an analysis

tool set for population genomics. Molecular Ecology 22, 3124-3140.

Chambers, E.A., Hillis, D.M., 2020. The Multispecies Coalescent Over-Splits Species in the Case

of Geographically Widespread Taxa. Systematic Biology 69, 184-193.

Chapman, F.M., Miller, L.E., Allen, A.A., Richardson, W.B., Cherrie, G.K., 1914. Diagnoses of

apparently new Colombian birds. 3. Bulletin of the AMNH; v. 33, article 40.

Chernomor, O., Minh, B.Q., von Haeseler, A., 2016. Terrace Aware Data Structure for

Phylogenomic Inference from Supermatrices. Systematic Biology 65, 997-1008.

Cheviron, Z.A., Hackett, S.J., Brumfield, R.T., 2006. Sequence variation in the coding region of the

melanocortin-1 receptor gene (MC1R) is not associated with plumage variation in the blue-crowned

manakin (Lepidothrix coronata). Proceedings of the Royal Society B: Biological Sciences 273,

1613-1618.

Cheviron, Z.A., Hackett, S.J., Capparella, A.P., 2005. Complex evolutionary history of a

Neotropical lowland forest bird (Lepidothrix coronata) and its implications for historical hypotheses

of the origin of Neotropical avian diversity. Molecular Phylogenetics and Evolution 36, 338-357.

Chifman, J., Kubatko, L., 2014. Quartet Inference from SNP Data Under the Coalescent Model.

Bioinformatics 30, 3317-3324.

Chifman, J., Kubatko, L., 2015. Identifiability of the unrooted species tree topology under the

coalescent model with time-reversible substitution processes, site-specific rate variation, and

invariable sites. Journal of Theoretical Biology 374, 35-47.

Chou, J., Gupta, A., Yaduvanshi, S., Davidson, R., Nute, M., Mirarab, S., Warnow, T., 2015. A

comparative study of SVDquartets and other coalescent-based species tree estimation methods.

BMC Genomics 16, S2.

Counterman, B.A., 2016. Hybrid Speciation. In: Kliman, R.M. (Ed.), Encyclopedia of Evolutionary

Biology. Academic Press, Oxford, pp. 242-248.

Cracraft, J., 1985. Historical Biogeography and Patterns of Differentiation within the South

American Avifauna: Areas of Endemism. Ornithological Monographs, 49-84.

Cracraft, J., Prum, R.O., 1988. PATTERNS AND PROCESSES OF DIVERSIFICATION:

SPECIATION AND HISTORICAL CONGRUENCE IN SOME NEOTROPICAL BIRDS.

Evolution 42, 603-620.

Crother, B.I., Murray, C.M., 2011. Ontology of areas of endemism. Journal of Biogeography 38,

1009-1015.

Cunningham, C.W., Collins, T.M., 1994. Developing model systems for molecular biogeography:

Vicariance and interchange in marine invertebrates. In: Schierwater, B., Streit, B., Wagner, G.P.,

DeSalle, R. (Eds.), Molecular Ecology and Evolution: Approaches and Applications. Birkhäuser

Basel, Basel, pp. 405-433.

Da Silva, J., Novaes, F., Oren, D., 2002. Differentiation of Xiphocolaptes (Dendrocolaptidae)

across the river Xingu, Brazilian Amazonia: recognition of a new phylogenetic species and

biogeographic implications. BULLETIN-BRITISH ORNITHOLOGISTS CLUB 122, 185-193.

da Silva, J.M.C., 1995. Biogeographic analysis of the South American Cerrado avifauna.

Steenstrupia 21, 49-67.

da Silva, J.M.C., de Sousa, M.C., H. M. Castelletti, C., 2004. Areas of Endemism for Passerine

Birds in the Atlantic Forest, South America. Global Ecology and Biogeography 13, 85-92.

Da Silva, J.M.C., Rylands , A.B., Da Fonseca, G.A.B., 2005. The Fate of the Amazonian Areas of

Endemism. Conservation Biology 19, 689-694.

de Queiroz, K., 2005. Ernst Mayr and the modern concept of species. Proceedings of the National

Academy of Sciences 102, 6600-6607.

de Schauensee, R.M., 1945. Notes on Colombian flycatchers, manakins and cotingas. Proceedings

of the Academy of Natural Sciences of Philadelphia 97, 41-57.

de Schauensee, R.M., 1950. Colombian zoological survey. Part V: New birds from Colombia.

Notulae Naturae (221), 1-13.

Dickinson, E., 2003. The Howard and Moore Complete Checklist of the Birds of the World

Christopher Helm. London.

Diniz-Filho, J.A.F., Soares, T.N., Lima, J.S., Dobrovolski, R., Landeiro, V.L., de Campos Telles,

M.P., Rangel, T.F., Bini, L.M., 2013. Mantel test in population genetics. Genetics and Molecular

Biology 36, 475-485.

Earl, D.A., vonHoldt, B.M., 2012. STRUCTURE HARVESTER: a website and program for

visualizing STRUCTURE output and implementing the Evanno method. Conservation Genet

Resour 4, 359-361.

Evanno, G., Regnaut, S., Goudet, J., 2005. Detecting the number of clusters of individuals using the

software structure: a simulation study. Molecular Ecology 14, 2611-2620.

Fernandes, A., Cohn-Haft, M., Hrbek, T., Farias, I., 2015. Rivers acting as barriers for bird dispersal

in the Amazon. Revista Brasileira de Ornitologia-Brazilian Journal of Ornithology 22, 363-373.

Fernandes, A.M., Gonzalez, J., Wink, M., Aleixo, A., 2013. Multilocus phylogeography of the

Wedge-billed Woodcreeper Glyphorynchus spirurus (Aves, Furnariidae) in lowland Amazonia:

Widespread cryptic diversity and paraphyly reveal a complex diversification pattern. Molecular

Phylogenetics and Evolution 66, 270-282.

Fjeldså, J., 1992. Biogeographic patterns and evolution of the avifauna of relict high-attitude

woodlands of the Andes. Steenstrupia, 9-62.

Flouri, T., Jiao, X., Rannala, B., Yang, Z., 2018. Species Tree Inference with BPP Using Genomic

Sequences and the Multispecies Coalescent. Molecular Biology and Evolution 35, 2585-2593.

Francisco, M.R., Gibbs, H.L., Galetti, M., Lunardi, V.O., Junior, P.M.G., 2007. Genetic structure in

a tropical lek-breeding bird, the blue manakin (Chiroxiphia caudata) in the Brazilian Atlantic Forest.

Molecular Ecology 16, 4908-4918.

Freile, J.F., 2014. First record of White-crowned Manakin Dixiphia pipra in western Ecuador. Bull.

Brit. Orn. Club 134, 238-240.

Goslee, S.C., Urban, D.L., 2007. The ecodist package for dissimilarity-based analysis of ecological

data. Journal of Statistical Software 22, 1-19.

Goudet, J., 2005. hierfstat, a package for r to compute and test hierarchical F-statistics. Molecular

Ecology Notes 5, 184-186.

Goudet, J., Perrin, N., Waser, P., 2002. Tests for sex-biased dispersal using bi-parentally inherited

genetic markers. Molecular Ecology 11, 1103-1114.

Grant, P.R., Grant, B.R., 1997. Genetics and the origin of bird species. Proceedings of the National

Academy of Sciences 94, 7768-7775.

Gronau, I., Hubisz, M.J., Gulko, B., Danko, C.G., Siepel, A., 2011. Bayesian inference of ancient

human demography from individual genome sequences. Nature Genetics 43, 1031.

Gubili, C., Ritter, C.D., Motta, J., Farias, I.P., Bates, J., Canton, R., Capurucho, J.M.G., Cornelius,

C., Feldheim, K.A., Ribas, C.C., 2016. Isolation and characterization of polymorphic microsatellite

DNA Markers from an Amazonian white-sand vegetation specialist bird, Xenopipo atronitens

(Aves: Pipridae). The Wilson Journal of Ornithology 128, 668-672.

Guillot, G., Rousset, F., 2013. Dismantling the Mantel tests. Methods in Ecology and Evolution 4,

336-344.

Haffer, J., 1969. Speciation in Amazonian Forest Birds. Science 165, 131-137.

Haffer, J., 1974. Avian speciation in tropical South America, with a systematic survey of the

toucans (Ramphastidae) and jacamars (Galbulidae) (Publications of the Nuttall Ornithological Club

No. 14, Cambridge, MA).

Haffer, J., 1978. Distribution of Amazon forest birds. Bonner Zoologische Beiträge 29, 38-78.

Haffer, J., 1985. Avian zoogeography of the Neotropical lowlands. Ornithological Monographs,

113-146.

Haffer, J., 1992. On the river effect in some forest birds of southern Amazonia. Boletim do Museu

Paraense Emílio Goeldi. Nova série. Zoologia 8, 217-245.

Haffer, J., 2008. Hypotheses to explain the origin of species in Amazonia. Brazilian Journal of

Biology 68, 917-947.

Hartl, D.L., Clark, A.G., 1997. Principles of population genetics. Sinauer associates Sunderland,

MA.

Harvey, M.G., Aleixo, A., Ribas, C.C., Brumfield, R.T., 2017. Habitat Association Predicts Genetic

Diversity and Population Divergence in Amazonian Birds. The American Naturalist 190, 631-648.

Harvey, M.G., Brumfield, R.T., 2015. Genomic variation in a widespread Neotropical bird (Xenops

minutus) reveals divergence, population expansion, and gene flow. Molecular Phylogenetics and

Evolution 83, 305-316.

Hazzi, N.A., Moreno, J.S., Ortiz-Movliav, C., Palacio, R.D., 2018. Biogeographic regions and

events of isolation and diversification of the endemic biota of the tropical Andes. Proceedings of the

National Academy of Sciences.

Hermansen, J.S., Sæther, S.A., Elgvin, T.O., Borge, T., Hjelle, E., Sætre, G.-P., 2011. Hybrid

speciation in sparrows I: phenotypic intermediacy, genetic admixture and barriers to gene flow.

Molecular Ecology 20, 3812-3822.

Hijmans, R.J., Williams, E., Vennes, C., 2015. Geosphere: spherical trigonometry. R package

version 1.3-11. URL: https://CRAN. R-project. org/package= geosphere.

Hillis, D.M., 2020. The Detection and Naming of Geographic Variation Within Species.

Herpetological Review, 51:52-56.

Hoang, D.T., Vinh, L.S., Chernomor, O., Minh, B.Q., von Haeseler, A., 2017. UFBoot2: Improving

the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution 35, 518-522.

Hooghiemstra, H., Van der Hammen, T., 2004. Quaternary Ice-Age dynamics in the Colombian

Andes: developing an understanding of our legacy. Philosophical Transactions of the Royal Society

of London. Series B: Biological Sciences 359, 173-181.

Hothorn, T., Bretz, F., Westfall, P., Heiberger, R.M., Schuetzenmeister, A., Scheibe, S., 2013.

multcomp: simultaneous inference in general parametric models. R package version 1.2-18.

Foundation for Statistical Computing, Vienna, Austria.

Jakobsson, M., Rosenberg, N.A., 2007. CLUMPP: a cluster matching and permutation program for

dealing with label switching and multimodality in analysis of population structure. Bioinformatics

23, 1801-1806.

Janzen, D.H., 1967. Why Mountain Passes are Higher in the Tropics. The American Naturalist 101,

233-249.

Jiggins, C.D., Salazar, C., Linares, M., Mavarez, J., 2008. Hybrid trait speciation and Heliconius

butterflies. Philosophical Transactions of the Royal Society B: Biological Sciences 363, 3047-3054.

Jombart, T., 2008. adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics 24, 1403-1405.

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermiin, L.S., 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods 14, 587.

Kamvar, Z.N., Tabima, J.F., Grünwald, N.J., 2014. Poppr: an R package for genetic analysis of

populations with clonal, partially clonal, and/or sexual reproduction. PeerJ 2, e281.

Kirwan, G.M., David, N., Gregory, S.M., Jobling, J.A., Steinheimer, F.D., Brito, G.R.R., 2016. The

mistaken manakin: a new genus-group name for Parus pipra Linnaeus, 1758 (Aves: Passeriformes:

Pipridae). Zootaxa 4121, 89-94.

Kirwan, G.M., Green, G., 2012. Cotingas and Manakins. Princeton University Press, Princeton, NJ.

Kumar, S., Subramanian, S., 2002. Mutation rates in mammalian genomes. Proceedings of the

National Academy of Sciences 99, 803-808.

Lamichhaney, S., Han, F., Webster, M.T., Andersson, L., Grant, B.R., Grant, P.R., 2018. Rapid

hybrid speciation in Darwin's finches. Science 359, 224-228.

Lanfear, R., Frandsen, P.B., Wright, A.M., Senfeld, T., Calcott, B., 2017. PartitionFinder 2: New

Methods for Selecting Partitioned Models of Evolution for Molecular and Morphological

Phylogenetic Analyses. Molecular Biology and Evolution 34, 772-773.

Lavretsky, P., Engilis, A., Eadie, J.M., Peters, J.L., 2015. Genetic admixture supports an ancient

hybrid origin of the endangered Hawaiian duck. Journal of Evolutionary Biology 28, 1005-1015.

Lawson, D.J., Hellenthal, G., Myers, S., Falush, D., 2012. Inference of Population Structure using

Dense Haplotype Data. PLOS Genetics 8, e1002453.

Leaché, A.D., Banbury, B.L., Felsenstein, J., de Oca, A.n.-M., Stamatakis, A., 2015. Short Tree,

Long Tree, Right Tree, Wrong Tree: New Acquisition Bias Corrections for Inferring SNP

Phylogenies. Systematic Biology 64, 1032-1047.

Leaché, A.D., Fujita, M.K., Minin, V.N., Bouckaert, R.R., 2014. Species Delimitation using

Genome-Wide SNP Data. Systematic Biology 63, 534-542.

Leaché, A.D., Zhu, T., Rannala, B., Yang, Z., 2018. The Spectre of Too Many Species. Systematic

Biology, syy051-syy051.

Legendre, P., Fortin, M.-J., 2010. Comparison of the Mantel test and alternative approaches for

detecting complex multivariate relationships in the spatial analysis of genetic data. Molecular

Ecology Resources 10, 831-844.

Linck, E.B., Battey, C.J., 2017. Minor allele frequency thresholds strongly affect population

structure inference with genomic datasets. bioRxiv.

Linder, H.P., 2001. On Areas of Endemism, with an Example from the African Restionaceae.

Systematic Biology 50, 892-912.

Long, C., Kubatko, L., 2017. The effect of gene flow on coalescent-based species-tree inference.

arXiv:1710.03806.

Luna, L.W., Souza, T.O., e Silva, W.A.d.G., Schneider, H., Sampaio, I., 2017. Genetic variation of

the endangered Araripe Manakin (Antilophia bokermanni) indicates a history of demographic

decline. Revista Brasileira de Ornitologia 25, 60-66.

Lynch Alfaro, J.W., Boubli, J.P., Paim, F.P., Ribas, C.C., Silva, M.N.F.d., Messias, M.R., Röhe, F.,

Mercês, M.P., Silva Júnior, J.S., Silva, C.R., Pinho, G.M., Koshkarian, G., Nguyen, M.T.T.,

Harada, M.L., Rabelo, R.M., Queiroz, H.L., Alfaro, M.E., Farias, I.P., 2015. Biogeography of

squirrel monkeys (genus Saimiri): South-central Amazon origin and rapid pan-Amazonian

diversification of a lowland primate. Molecular Phylogenetics and Evolution 82, 436-454.

MacQueen, J., 1967. Some methods for classification and analysis of multivariate observations.

Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability, Volume

1: Statistics. University of California Press, Berkeley, Calif., pp. 281-297.

Malinsky, M., Trucchi, E., Lawson, D.J., Falush, D., 2018. RADpainter and fineRADstructure:

Population Inference from RADseq Data. Molecular Biology and Evolution 35, 1284-1290.

Mantel, N., 1967. The detection of disease clustering and a generalized regression approach. Cancer

research 27, 209-220.

Marks, B.D., Hackett, S.J., Capparella, A.P., 2002. Historical relationships among Neotropical

lowland forest areas of endemism as determined by mitochondrial DNA sequence variation within

the Wedge-billed Woodcreeper (Aves: Dendrocolaptidae: Glyphorynchus spirurus). Molecular

Phylogenetics and Evolution 24, 153-167.

Marques, D.A., Meier, J.I., Seehausen, O., 2019. A Combinatorial View on Speciation and

Adaptive Radiation. Trends in Ecology & Evolution.

Mayr, E., 1942. Systematics and the origin of species from the viewpoint of a zoologist. Columbia

University Press.

McDonald, D.B., 2003. MICROSATELLITE DNA EVIDENCE FOR GENE FLOW IN

NEOTROPICAL LEK-MATING LONG-TAILED MANAKINS. The Condor 105, 580-586.

Milá, B., Tavares, E.S., Muñoz Saldaña, A., Karubian, J., Smith, T.B., Baker, A.J., 2012. A Trans-

Amazonian Screening of mtDNA Reveals Deep Intraspecific Divergence in Forest Birds and

Suggests a Vast Underestimation of Species Diversity. PLOS ONE 7, e40541.

Moore, R.P., Robinson, W.D., Lovette, I.J., Robinson, T.R., 2008. Experimental evidence for

extreme dispersal limitation in tropical forest birds. Ecology Letters 11, 960-968.

Morrone, J.J., 2015. Biogeographical regionalisation of the Andean region. Zootaxa 3936, 207-236.

Musher, L.J., Ferreira, M., Auerbach, A.L., McKay, J., Cracraft, J., 2019. Why is Amazonia a

source of biodiversity? Climate-mediated dispersal and synchronous speciation across the Andes in

an avian group (Tityrinae). Proceedings of the Royal Society B: Biological Sciences 286,

20182343.

Naka, L.N., 2011. Avian distribution patterns in the Guiana Shield: implications for the delimitation

of Amazonian areas of endemism. Journal of Biogeography 38, 681-696.

Naka, L.N., Brumfield, R.T., 2018. The dual role of Amazonian rivers in the generation and

maintenance of avian diversity. Science Advances 4.

Nieto Feliner, G., Álvarez, I., Fuertes-Aguilar, J., Heuertz, M., Marques, I., Moharrek, F., Piñeiro,

R., Riina, R., Rosselló, J.A., Soltis, P.S., Villa-Machío, I., 2017. Is homoploid hybrid speciation that

rare? An empiricist's view. Heredity 118, 513.

Noguera-urbano, E.A., 2016. Areas of endemism: travelling through space and the unexplored

dimension. Systematics and Biodiversity 14, 131-139.

Nyári, Á.S., 2007. Phylogeographic patterns, molecular and vocal differentiation, and species limits

in Schiffornis turdina (Aves). Molecular Phylogenetics and Evolution 44, 154-164.

Nylander, J.A.A., Olsson, U., Alström, P., Sanmartín, I., 2008. Accounting for Phylogenetic

Uncertainty in Biogeography: A Bayesian Approach to Dispersal-Vicariance Analysis of the

Thrushes (Aves: Turdus). Systematic Biology 57, 257-268.

Ohlson, J.I., Fjeldså, J., Ericson, P.G.P., 2013. Molecular phylogeny of the manakins (Aves:

Passeriformes: Pipridae), with a new classification and the description of a new genus. Molecular

Phylogenetics and Evolution 69, 796-804.

Paradis, E., 2010. pegas: an R package for population genetics with an integrated–modular

approach. Bioinformatics 26, 419-420.

Paris, J.R., Stevens, J.R., Catchen, J.M., Johnston, S., 2017. Lost in parameter space: a road map for

stacks. Methods in Ecology and Evolution 8, 1360-1373.

Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S., Hoekstra, H.E., 2012. Double Digest RADseq:

An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-Model

Species. PLOS ONE 7, e37135.

Petkova, D., Novembre, J., Stephens, M., 2015. Visualizing spatial population structure with

estimated effective migration surfaces. Nature Genetics 48, 94.

Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of Population Structure Using

Multilocus Genotype Data. Genetics 155, 945-959.

Prum, R.O., 1990. Phylogenetic Analysis of the Evolution of Display Behavior in the Neotropical

Manakins (Aves: Pipridae). Ethology 84, 202-231.

Prum, R.O., 1994. PHYLOGENETIC ANALYSIS OF THE EVOLUTION OF ALTERNATIVE

SOCIAL BEHAVIOR IN THE MANAKINS (AVES: PIPRIDAE). Evolution 48, 1657-1675.

Pupim, F.N., Sawakuchi, A.O., Almeida, R.P., Ribas, C.C., Kern, A.K., Hartmann, G.A., Chiessi,

C.M., Tamura, L.N., Mineli, T.D., Savian, J.F., Grohmann, C.H., Bertassoli, D.J., Stern, A.G., Cruz,

F.W., Cracraft, J., 2019. Chronology of Terra Firme formation in Amazonian lowlands reveals a

dynamic Quaternary landscape. Quaternary Science Reviews 210, 154-163.

Quintero, I., Jetz, W., 2018. Global elevational diversity and diversification of birds. Nature 555,

246.

Rambaut, A., Drummond, A.J., Xie, D., Baele, G., Suchard, M.A., 2018. Posterior Summarization

in Bayesian Phylogenetics Using Tracer 1.7. Systematic Biology 67, 901-904.

Räsänen, M.E., Salo, J.S., Kalliola, R.J., 1987. Fluvial perturbance in the western Amazon basin:

regulation by long-term sub-Andean tectonics. Science 238, 1398-1401.

Ribas, C.C., Aleixo, A., Nogueira, A.C.R., Miyaki, C.Y., Cracraft, J., 2012. A palaeobiogeographic

model for biotic diversification within Amazonia over the past three million years. Proceedings of

the Royal Society B: Biological Sciences 279, 681-689.

Ribas, C.C., Moyle, R.G., Miyaki, C.Y., Cracraft, J., 2007. The assembly of montane biotas: linking

Andean tectonics and climatic oscillations to independent regimes of diversification in Pionus

parrots. Proceedings of the Royal Society B: Biological Sciences 274, 2399-2408.

Ridgely, R.S., Greenfield, P.J., 2001. The birds of Ecuador: status, distribution, and taxonomy.

Cornell University Press.

Ridgely, R.S., Tudor, G., 2009. Field guide to the songbirds of South America: the passerines.

University of Texas Press.

Ridgway, R., 1906. Descriptions of some new forms of oligomyodian birds. Proceedings of the

Biological Society of Washington, 115-120.

Roure, B., Baurain, D., Philippe, H., 2013. Impact of Missing Data on Phylogenies Inferred from

Empirical Phylogenomic Data Sets. Molecular Biology and Evolution 30, 197-214.

Rousset, F., 1997. Genetic Differentiation and Estimation of Gene Flow from F Statistics Under

Isolation by Distance. Genetics 145, 1219-1228.

Salazar, C., Baxter, S.W., Pardo-Diaz, C., Wu, G., Surridge, A., Linares, M., Bermingham, E.,

Jiggins, C.D., 2010. Genetic Evidence for Hybrid Trait Speciation in Heliconius Butterflies. PLOS

Genetics 6, e1000930.

Saranathan, V., Hamilton, D., Powell, G.V.N., Kroodsma, D.E., Prum, R.O., 2007. Genetic

evidence supports song learning in the three-wattled bellbird Procnias tricarunculata (Cotingidae).

Molecular Ecology 16, 3689-3702.

Schmidt, H.A., Minh, B.Q., von Haeseler, A., Nguyen, L.-T., 2014. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Molecular Biology and

Evolution 32, 268-274.

Schmidt-Lebuhn, A.N., Aitken, N.C., Chuah, A., 2017. Species trees from consensus single

nucleotide polymorphism (SNP) data: Testing phylogenetic approaches with simulated and

empirical data. Molecular Phylogenetics and Evolution 116, 192-201.

Schumer, M., Rosenthal, G.G., Andolfatto, P., 2014. HOW COMMON IS HOMOPLOID HYBRID

SPECIATION? Evolution 68, 1553-1560.

Sedano, R.E., Burns, K.J., 2010. Are the Northern Andes a species pump for Neotropical birds?

Phylogenetics and biogeography of a clade of Neotropical tanagers (Aves: Thraupini). Journal of

Biogeography 37, 325-343.

Shafer, A.B.A., Peart, C.R., Tusso, S., Maayan, I., Brelsford, A., Wheat, C.W., Wolf, J.B.W., 2017.

Bioinformatic processing of RAD-seq data dramatically impacts downstream population genetic

inference. Methods in Ecology and Evolution 8, 907-917.

Silva, S.M., Peterson, A.T., Carneiro, L., Burlamaqui, T.C.T., Ribas, C.C., Sousa-Neves, T.,

Miranda, L.S., Fernandes, A.M., d'Horta, F.M., Araújo-Silva, L.E., Batista, R., Bandeira,

C.H.M.M., Dantas, S.M., Ferreira, M., Martins, D.M., Oliveira, J., Rocha, T.C., Sardelli, C.H.,

Thom, G., Rêgo, P.S., Santos, M.P., Sequeira, F., Vallinoto, M., Aleixo, A., 2019. A dynamic

continental moisture gradient drove Amazonian bird diversification. Science Advances 5, eaat5752.

Smeds, L., Qvarnström, A., Ellegren, H., 2016. Direct estimate of the rate of germline mutation in a

bird. Genome Research 26, 1211-1218.

Smith, B.T., McCormack, J.E., Cuervo, A.M., Hickerson, M.J., Aleixo, A., Cadena, C.D., Perez-

Eman, J., Burney, C.W., Xie, X., Harvey, M.G., Faircloth, B.C., Glenn, T.C., Derryberry, E.P.,

Prejean, J., Fields, S., Brumfield, R.T., 2014. The drivers of tropical speciation. Nature 515, 406-

409.

Snow, D., 1961. THE DISPLAYS OF THE MANAKINS PIPRA PIPRA AND TYRANNEUTES

VIRESCENS*. Ibis 103A, 110-113.

Snow, D.W., 2004. Family Pipridae (Manakins). In: del Hoyo, J., Elliot, A., Christie, D.A. (Eds.),

Handbook of the Birds of the World. Lynx Edicions, Barcelona, pp. 110-169.

Spencer, A., 2012. White-crowned Manakin vocal variation. Xeno-Cango.org.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics 30, 1312-1313.

Suh, A., 2016. The phylogenomic forest of bird trees contains a hard polytomy at the root of

Neoaves. Zoologica Scripta 45, 50-62.

Sukumaran, J., Knowles, L.L., 2017. Multispecies coalescent delimits structure, not species.

Proceedings of the National Academy of Sciences 114, 1607-1612.

Swofford, D.L., 2002. Phylogenetic analysis using parsimony (*and Other Methods). Version 4.

Sinauer Associates, Sunderland, Massachusetts.

Tello, J.G., Moyle, R.G., Marchese, D.J., Cracraft, J., 2009. Phylogeny and phylogenetic

classification of the tyrant flycatchers, cotingas, manakins, and their allies (Aves: Tyrannides).

Cladistics 25, 429-467.

Teske, P.R., Golla, T.R., Sandoval-Castillo, J., Emami-Khoyi, A., van der Lingen, C.D., von der

Heyden, S., Chiazzari, B., Jansen van Vuuren, B., Beheregaray, L.B., 2018. Mitochondrial DNA is

unsuitable to test for isolation by distance. Scientific Reports 8, 8448.

Thom, G., Amaral, F.R.D., Hickerson, M.J., Aleixo, A., Araujo-Silva, L.E., Ribas, C.C., Choueri,

E., Miyaki, C.Y., 2018. Phenotypic and Genetic Structure Support Gene Flow Generating Gene

Tree Discordances in an Amazonian Floodplain Endemic Species. Systematic Biology 67, 700-718.

Thrasher, D.J., Butcher, B.G., Campagna, L., Webster, M.S., Lovette, I.J., 2018. Double-digest

RAD sequencing outperforms microsatellite loci at assigning paternity and estimating relatedness:

A proof of concept in a highly promiscuous bird. Molecular Ecology Resources 0.

Trifinopoulos, J., Nguyen, L.-T., Minh, B.Q., von Haeseler, A., 2016. W-IQ-TREE: a fast online

phylogenetic tool for maximum likelihood analysis. Nucleic Acids Research 44, W232-W235.

Vaidya, G., Lohman, D.J., Meier, R., 2011. SequenceMatrix: concatenation software for the fast

assembly of multi-gene datasets with character set and codon information. Cladistics 27, 171-180.

van Els, P., Norambuena, H.V., Etienne, R.S., 2019. From pampa to puna: Biogeography and

diversification of a group of Neotropical obligate grassland birds (Anthus: Motacillidae). Journal of

Zoological Systematics and Evolutionary Research 00: 1– 12.

Voelker, G., 1999. DISPERSAL, VICARIANCE, AND CLOCKS: HISTORICAL

BIOGEOGRAPHY AND SPECIATION IN A COSMOPOLITAN PASSERINE GENUS

(ANTHUS: MOTACILLIDAE). Evolution 53, 1536-1552.

Wallace, A.R., 1854. On the monkeys of the Amazon. Annals and Magazine of Natural History 14,

451-454.

Weir, B.S., Cockerham, C.C., 1984. ESTIMATING F-STATISTICS FOR THE ANALYSIS OF

POPULATION STRUCTURE. Evolution 38, 1358-1370.

Weir, J.T., 2006. DIVERGENT TIMING AND PATTERNS OF SPECIES ACCUMULATION IN

LOWLAND AND HIGHLAND NEOTROPICAL BIRDS. Evolution 60, 842-855.

Wen, D., Yu, Y., Zhu, J., Nakhleh, L., 2018. Inferring Phylogenetic Networks Using PhyloNet.

Systematic Biology, syy015-syy015.

Wetmore, A., 1972. The birds of the Republic of Panama. Part 3—Passeriformes: Dendrocolaptidae

(woodcreepers) to Oxyruncidae (sharpbills). Smithsonian Miscellaneous Collection 150.

Smithsonian Institution Press, Washington, DC.

Wright, S., 1931. Evolution in Mendelian populations. Genetics 16, 97.

Zhang, J., Kapli, P., Pavlidis, P., Stamatakis, A., 2013. A general species delimitation method with

applications to phylogenetic placements. Bioinformatics 29, 2869-2876.

Zhu, J., Wen, D., Yu, Y., Meudt, H.M., Nakhleh, L., 2018. Bayesian inference of phylogenetic

networks from bi-allelic genetic markers. PLOS Computational Biology 14, e1005932.

Zimmer, J.T., 1936. Studies of Peruvian birds. XXII. Notes on the Pipridae. American Museum

novitates; no. 889. 1-29.

1696

1697

1698

1699

1700

1701

1702

1703

1704

1705

1706

1707

1708

1709

1710

1711

Figure 1. Neotropical areas of endemism and sampling localities. Amazonian lowland areas of

endemism as portrayed in Da Silva et al. (2005) are highlighted in shades of green, as well as the

Jaú area of endemism (Borges and Da Silva, 2012). Montane Andean, Central American, Guianan,

and dry diagonal regions (>1000m) are emphasized in grey. As summarized in Da Silva et al.

(2005), a sequence of authors identified seven areas of endemism for lowland birds that retained

Wallace's "Guyana" (Wallace, 1854), split "Ecuador" into "Imeri" and "Napo," renamed "Peru" to

"Inambari," and split "Brazil" into "Rondônia," "Pará," and "Belém" (Cracraft, 1985; Cracraft and

Prum, 1988; Haffer, 1978; Haffer, 1985; Haffer, 1992). "Pará" was subsequently further partitioned

into two regions, "Tapajós" and "Xingu," (Da Silva et al., 2002). Recently, additional sub-partitions

have been proposed for the "Napo" (Jaú - Borges and Da Silva, 2012), and Guianan areas of

endemism (Naka, 2011). Plotted points indicate field sampling localities for 286 *Pseudopipra*

individuals sequenced for genetic analysis in the present study. Inset top right is an illustration of

the *pipra* subspecies, reproduced from the Handbook of the Birds of the World, Lynx Edicions.

1725

1726

1727

1728

1729

1730

1731

1732

1733

1734

CENTRAL
AMERICA

-80

-70

-60

-50

-40

GUIANA

Essequibo river

IMERI

Negro river

NAPO

Napo river

Japurá river

Negro river

Amazonas river

Putumayo river

JAÚ

Solimões river

BELEM

Amazonas river

Marañón river

Purus river

Tapajós river

Xingu river

INAMBARI

Madeira river

XINGU

Caatinga

Huallaga river

Ucayali river

RONDÔNIA

TAPAJÓS

-10

Cerrado

ATLANTIC
FOREST

ANDEAN
MOUNTAINS

Chaco

-20

Figure 2. Subspecies sampling. Our genetic sampling of *Pseudopipra pipra* intersects with the

ranges of ten named subspecies. In this figure, we indicate sampling localities as they overlap with

previously identified subspecies ranges. Areas greater than 1000m in elevation are emphasized in

light grey; major Amazonian rivers are indicated in light blue. Overall distribution reflects the

BirdLife International approximation, updated to reflect current knowledge (BirdLife International

Species factsheet: *Pseudopipra pipra*, 2018, also see Supplemental Figure 1). The ranges of Andean

taxa were adjusted based on unpublished data provided Andrés Cuervo (*personal communication*).

Hypothetical subspecies ranges are based on descriptions in Kirwan and Green (2012) and imply

maximal ranges highlighting key geographic barriers delimiting subspecies. Black dots indicate

sampling for ddRADseq, while crosses indicate sampling for mtDNA ND2. Triangles indicate

sampling localities for male advertisement vocalizations (upward) and call vocalizations

(downward). Our sampling encompasses the majority of the lowland extents of the recognized

range of *Pseudopipra*, as well as key montane populations in Peru, Ecuador, and Central America.

Ssp *anthracina, bolivari, minima, unica, coracina, occulta, and comata* are exclusively montane.

1749

1750

1751

1752

1753

1754

1755

1756

1757

Figure 3. Phylogenetic hypothesis for *Pseudopipra*. Phylogenetic analysis of concatenated ddRAD

sequences generated a well-resolved phylogenetic hypothesis which was largely congruent across

datasets and analytical approaches. Montane clades, including a Northern Andean (A1) + Central

American (A2) clade (A), and a central Peruvian clade (B), are recovered as nested sister groups to

two wide ranging lowland clades (C and D), indicating that the lowland clades are descended from

Andean lineages (Figure 5). Shown is the RAxML topology generated using the 50% haplotype

dataset. Branch lengths are set to equal for display (see Supplementary Data for original tree files).

Colored circles at tips indicate group membership to one of eighteen population-areas (matching

Figure 4 and 5). Heatmaps at key nodes indicate bootstrap scores from each of six phylogenetic

analyses for 20/50/80% datasets, as indicated in the legend. Black dots on branches indicate

populations which are identified with 100% posterior probability in fineRADstructure analysis.

Deep phylogenetic structure is recovered by both RAxML and SVDquartets with high support,

though RAxML recovers additional low-support substructure in the Northern Amazonian + Guiana

Shield clade (D) which is coincident with several geographic features (clades D1 and D2). A clade

of introgressed western Napo individuals (E) is recovered by all standard phylogenetic analyses as

sister to the northern Amazonian + Guiana shield clade (D). As the history of this clade is

characterized by a complex introgression scenario between northern and southern lineages, a purely

bifurcating tree model is inadequate to describe its relations with other groups (see text and Figure

6). Also shown are the well supported phylogenetic positions of two distinct subspecies lineages

(*pygmaea* and *coracaina*) from which we were able to obtain mtDNA (Supplemental Appendix).

1778

1779

1780

Figure 4. Mapping of phylogenetic relationships into geographic space. Here, a simplified topology

is plotted with tips anchored to sampling coordinates. The colors of convex hulls (i.e., hypotheses of

minimum clade ranges) match tip colors in Figure 3 and 5. These colored hulls represent distinct

non-overlapping geographic areas delimited by physical barriers (though this is somewhat visually

confounded in the Northern Amazon + Guiana Shield clade (D), with two weakly resolved lineages

(e.g. pale yellow Jaú and green Guiana Shield population groups appear to have some overlap in

this figure, but they are delimited by the Rio Negro). Clade D bipartitions are shown with dashed

branches (top) to indicate low support for phylogenetic structure (Figure 3, results). The ancestral

habit of the genus is inferred to be Andean (Figure 5, Supplementary Appendix), with a single

origin of lowland lineages. Also shown at the tips of the simplified tree are median admixture

proportions across sampling localities as inferred with STRUCTURE (Figure 5).

1792

1793

1794

1795

1796

1797

1798

1799

1800

1801

1802

1803

D

D2

D1

A3

A4

coracina

A1

Andean
MRCA

B2

B1

pygmaea

E   C1/2   C3   C4   C5   C6   C10   C9   C8

Lowland MRCA

Figure 5. Cluster and admixture analysis of SNP data. Left: RAxML topology using the 50%

haplotype dataset, with tip labels and colors indicating group membership to one of eighteen

population-areas (matching those in Figures 3, 4). Branch colors indicate ancestral area, according

to the results from SIMMAP and maximum parsimony analysis (Supplemental Appendix), and

unambiguously reconstruct the *Pseudopipra* MRCA to have been Andean. Center: STRUCTURE

output for Dataset 1, depicting population assignment and admixture estimates for K=5. The

likelihood of each evaluated number of K clusters from 1:20 plateaued at K = 5, with the standard

deviation across runs increasing rapidly after this point (Supplemental Figure 2). The five optimized

clusters broadly correspond to wide biogeographic Neotropical regions which coincide with

lowland areas of endemism, Central American, and Peruvian montane regions (Figure 4). Given

these five clusters, each labeled focal region is inferred to have a unique combination of admixture

proportions. The admixture cline inferred from the Xingu to the southern Atlantic Forest appears to

be a product of isolation by distance (Supplementary Appendix), while the signature of

introgression inferred for western Napo lineages appears to be a product of substantial historical

introgression from a northern lineage, into a distinct Southern Amazonian lineage (see Figure 6 and

discussion). See Supplementary Figure 3 for K2-10 results. Right: clustering solutions for SNP

dataset D1-D4m (described in Table 1, also see Supplementary Figure 2), as well as for the

coancestry matrix (CoA). Black bars indicate distinct clusters which are also detected as

monophyletic groups in the phylogenetic analysis. Red bars indicate one distinct cluster in a given

dataset, but which is paraphyletic with respect to the phylogenetic analysis. The grey bars for D2m

indicate several groups of individuals that were not resolvable in this dataset (Supplementary Figure

2).

1826

RAXML   **STRUCTURE k=5**   D1 D1m D2 D2m D3 D3m D4 D4m CoA   k-means

North Andean

Montane Central America

South Andean

Inambari (W)

Montane

Inambari (E)

Rondônia

Tapajós

Xingu

Atlantic Forest

Bahia to Rio

Lowland Southern Amazon

Introgressed Western Napo Lineage

weakly resolved eastern Napo

Lowland Northern Amazon/Guiana Shield

weakly resolved Suriname, Amapá

Andean

Central American

Amazonian/Lowland

Figure 6. Examination of the introgressed lineage in the western Napo area of endemism. All of the

available genomic evidence is consistent with the hypothesis that individuals in the western Napo

area of endemism are derived from a distinct southern Amazonian lineage (S2a in Panel A) which is

not represented in contemporary samples. This lineage (S2a) experienced substantial introgression

from the ancestral populations of what are today eastern Napo and Jaú restricted linages. Panel A

shows the phylogenetic network (in black) inferred by PhyloNet, with arrows indicating the

direction of introgression estimated in G-PhoCS. The red arrows along the PhyloNet reticulation

indicate inferred introgression from the northern Amazonian lineages (clade C1 in Figure 3) into

this southern Amazonian lineage (S2a). The dashed grey line connecting S2a/S2 to S1 indicates

possible introgression into Inambari lineages, which may have occurred after or concurrently with

the primary introgression event and brought northern (Clade D) alleles into SW Amazonian

populations (see text and Figure 5). Panel B displays the individual level coancestry matrix for

introgressed Napo individuals and their source/progenitor lineages; darker colors indicate greater

coancestry. Introgressed (PH: putative hybrid) hybrids have greater coancestry with each of their

progenitor lineages (S2a and Northern Amazon+Guiana shield (abbreviated GS)) than their

progenitor lineages share with each other (PHvGS/progenitors = 1.078, PHvInambari/progenitors =

1.17, see text). In the proposed scenario, the southern progenitor lineage (S2a) is either unsampled

or extinct and has not persisted sympatrically with S2. Thus, the western Inambari southern lineage

(S1) comprises the closest living relatives to S2a (see discussion). Panel C summarizes coancestry

variance as projected into component space after normalizing. Panel D summarizes the minimum

implied geographic extents of the groups involved (as sampled in this study). For the convex hull

representing the range of the western Napo lineage, dark shading indicates localities for ddRAD

sampling, while light shading indicates indicates additional localities of the Napo lineage as implied

by mtDNA samples. G-PhoCS demographic parameter estimates are also depicted in panel D (m:

migrants per generation, when lower 95% CI > 1, τ: splitting time in generations ($\times10^6$), θ: effective

population size (10^6), Supplemental Table 5). In A-D, two subpopulations (East and West) are

indicated for each progenitor group, as outlined with outlined minimum convex hulls in D.

1854

1855

1856

1857

1858

1859

1860

1861

1862

1863

1864

1865

1866

1867

1868

1869

1870

1871

1872

1873

A)

SW Amazon
Inambari
S1 (Clade C1)

S2a

S2 (Clade E)

F

NW Amazon
Jaú + eastern Napo

B)

Jaú   Eastern Napo   Introgressed lineage   West Inambari   East Inambari

Northwest Amazon   Southern Amazon clade

C) coancestry PCA of introgressed + progenitors

+ Eastern Napo + Jaú
○ Western Napo (introgressed)
△ West + East Inambari

Component 1 explains 76.6% of the point variability

Component 2 explains 8.2% of the point variability

D)

Napo

major introgression
m: 1.75 [1.05 - 2.59]

Eastern Napo   Jaú

possible
minor introgression

θ~2.81

S2a/S2 θ~4.95   τ~1.03

F

θ~1.43

Inambari   τ~0.485   θ~5.36

S1 θ~2.78

Rondônia

Lat

Long

Figure 7. Estimated Effective migration surface (EEMS). The estimated migration surface links the

topography and drainage system of the Neotropics to spatial patterns of genetic variation. Shown

are results estimated from a model with 2000 demes (vertices). Blue colors indicate areas where

gene flow is more likely to be occurring (i.e., more connectivity than isolation by distance), while

brown colors indicate areas where barriers to gene flow are highly probable (i.e. less connectivity

than isolation by distance). In this analysis, the Andes and the Amazon River are the most clearly

inferred barriers to gene flow. Major rivers, as well as a >1000m contour (light grey) are overlaid to

aid interpretation. See text for additional discussion.

1882

1883

1884

1885

1886

1887

1888

1889

1890

1891

1892

1893

1894

1895

1896

Posterior mean migration rates

log10(m)

Andes Mountains

Isthmus of Panama

Essequibo River

Wilhelmina Mountains

Rio Japurá

Roraima grasslands

Rio Amazonas

Rio Marañón

Caatinga

Rio Tapajós

Cerrado

Rio Xingu

Rio Purus

Coastal Bahia

Chaco

Serra do Mar

Figure 8. Panel A: Summary of lek vocalization phenotypes. Our analysis identified 11

qualitatively distinct lek vocalization types that can be easily diagnosed by ear, or by visual

inspection of sonograms (arbitrarily labeled numerically). See Supplementary Figure 13 for

vocalization measures. The inset map indicates the minimum implied ranges for each identified

vocalization type, with colors and plotting symbols matching those in PCA plots (Supplemental

Figures 14, 15). The thicker pale blue dashed line indicates where the 'southern Amazon'

vocalization type 2 may be shared, apparently plesiomorphically (Supplementary Figure 16).

Vocalization type 2 may therefore occur in other unsampled areas of the southern Amazon. Type 1

also appears to be shared pleisiomorphically between populations on the eastern slopes of the Andes

(Clade A1) and in populations from the southern Cordillera Azul in Loreto (Clade B1). Panel B:

Summary of call vocalization phenotypes. Our analysis identified three distinct but similar call

vocalization types that can be diagnosed by ear, or by visual inspection of sonograms. Vocalization

type 5 was primarily recorded in the northern Amazon basin, coinciding with the broad ranging

Northern Amazon + Guiana Shield lineage (Clade D). From our initial classification, one recording

from the Atlantic Forest (XC427315) was assigned to type 5, suggesting this type may be shared

between the Guiana Shield clade and the Atlantic Forest clade (not shown, supplemental

discussion). As with lek vocalization type 1, call vocalization type 10 was shared pleisiomorphically

across Andean clades (A1 + B1). Call vocalization type 13 was entirely restricted to but shared

across diverse lowland southern Amazonian forms (Clade C). Letters to the top right of sonograms

indicate explicit [i.e., *A*] or implicit clade associations [i.e., (*A*)]. See Supplemental Appendix for

additional details and discussion.

1918

1919

1920   **Table 1.** Summary of dataset filtering parameters. We generated eight SNP datasets under different

1921   filtering schemes to evaluate the sensitivity of cluster analyses. For eight SNP datasets, dataset D1 –

1922   D4mod, we note the parameters for each filter: the required sequencing coverage, the missing data

1923   threshold, whether or not a single SNP (single nucleotide polymorphism) or all SNPs per locus were

1924   retained, the minor allele frequency (MAF), the number of loci, the total number of SNPS

1925   represented in a given dataset, and the average number of SNPs per locus.

1926

1927

1928

1929

1930

1931

1932

1933

1934

1935

1936

1937

1938

1939

1940

1941

1942

Table 1: Summary of dataset filtering parameters

Dataset	coverage	SNP present in	linkage	MAF	loci	SNPs	SNPs/locus
D1	>20x	>80%	1st SNP	>= 5%	1960	1960	1
D1mod	>20x	100%	1st SNP	>= 5%	175	175	1
D2	>20x	>80%	1st SNP	NA	2581	2581	1
D2mod	>20x	100%	1st SNP	NA	250	250	1
D3	>20x	>80%	all SNPs	>= 5%	1960	5099	2.6
D3mod	>20x	100%	all SNPs	>= 5%	177	391	2.2
D4	>20x	>80%	all SNPs	NA	2581	41024	15.9
D4mod	>20x	100%	all SNPs	NA	251	3832	15.3

1943   **Table 2.** Summary of Mantel Tests. For each geographic barrier and sampling region we were able

1944   to evaluate, we report results for analyses using phylogenetic patristic distance and coancestry,

1945   under both linear and log scales. Asterisks indicate *p* values less than 0.001, which we interpret as

1946   statistical significance (see discussion in text).

Table 2: Summary of Mantel Tests

	Patristic distance		Coancestry	
Approximate Barrier	p < 0.001	p < 0.001	p < 0.001	p < 0.001
Cordillera de Talamanca	-	*	-	-
Andes (1)	-	*	-	*
Andes (2)	-	*	-	*
Andes (3)	*	*	-	*
Rio Ucayali	*	*	*	*
Eastern Marañón + Hauallaga Rivers	*	*	*	*
Rio Putumayo	*	*	*	*
Rio Purus	*	*	-	*
Rio Madeira	*	*	*	-
Rio Tapajós	-	*	-	-
Rio Xingu	*	*	-	-
Cerrado (1)	*	*	-	*
Cerrado (2)	*	*	*	*
Rio Japurá	-	*	*	*
Rio Negro	-	-	-	-
Rio Essequibo	*	*	*	*
Rio Amazonas	*	*	*	*
Regions				
Full dataset	*	*	*	*
Central America - Costa Rica (Clade A3)	-	-	-	-
Central America - Panama (Clade A4)	-	-	-	-
North Andean – San Martín (Clade A1)	?	?	?	?
South Andean Peru (North) (Clade B2)	-	-	-	-
South Andean Peru (South) (Clade B1)	-	-	-	-
weakly resolved Guiana Shield (Clade D)	-	-	-	*
unresolved Jaú	-	*	-	*
weakly resolved eastern Napo (Clade D2)	-	-	-	-
weakly resolved Suriname + Amapá (D1)	-	-	-	*
Western Napo introgressed lineage (Clade E)	-	-	-	-
Western Inambari endemic (Clade C1)	*	*	-	*
Eastern Inambari endemic (Clade C3)	-	-	-	-
Rondônia endemic (Clade C4)	?	?	?	?
Tapajós endemic (Clade C5)	-	-	-	*
Xingu endemic (Clade C6)	-	-	-	-
Atlantic Forest – Bahia (Clade C8)	-	-	-	-
Atlantic Forest – Espírito Santo (Clade C9)	?	?	?	?
Atlantic Forest – Rio (Clade C10)	*	-	*	*
	linear	**log**	**linear**	**log**