Studying the interplay of sleep and aging:
Methodological challenges

Beate E. Muehlroth[a] and Markus Werkle-Bergner[a,b]

[a]Center for Lifespan Psychology, Max Planck Institute for Human Development, Berlin, Germany
[b]Institute for Psychology, Universität der Bundeswehr München, Neubiberg, Germany

Correspondence concerning this article should be addressed to: Beate E. Muehlroth (muehlroth@mpib-berlin.mpg.de; +49 30 82406-255) or Markus Werkle-Bergner (werkle@mpib-berlin.mpg.de; +49 30 82406-447); Max Planck Institute for Human Development, Lentzeallee 94, 14195 Berlin, Germany

Abstract

In quest of new avenues to explain, predict, and treat pathophysiological conditions during aging, research on sleep and aging has flourished. Despite the great scientific potential to pinpoint mechanistic pathways between sleep, aging, and pathology, only little attention has been paid to the suitability of analytic procedures applied to study these interrelations. On the basis of electrophysiological sleep and structural brain data of healthy younger and older adults, we identify, illustrate, and resolve methodological core challenges in the study of sleep and aging. We demonstrate potential biases in common analytic approaches when applied to older populations. We argue that uncovering age-dependent alterations in the physiology of sleep requires the development of age-fair and individualized analytic procedures that filter out age-independent inter-individual differences. Age-adapted methodological approaches are thus required to foster the development of valid and reliable biomarkers of age-associated cognitive pathologies.

Introduction

In search of novel biomarkers and treatment targets for age-related pathologies, sleep has recently attracted much attention. Disrupted sleep represents one of the earliest symptoms of Alzheimer's disease (Lim et al., 2013; Lucey et al., 2019), but might also potentiate, accelerate, or even cause cognitive pathology in old age (Shokri-Kojori et al., 2018; Vaou et al., 2018). Because of the assumed bidirectional relationship between sleep and Alzheimer's pathology (Ju et al., 2014; Mander et al., 2016; Noble and Spires-Jones, 2019; Vaou et al., 2018), studying the interrelations between sleep and neurocognitive aging has become increasingly popular.

Even in the absence of any diagnostically identifiable pathology, healthy aging entails fundamental yet very characteristic changes in sleep (Mander, 2017; Ohayon et al., 2004; Vitiello, 2006). These alterations are evident in the global distribution and succession of sleep stages (i.e., the *macro-structure* of sleep; Feinberg, 1974; Ohayon et al., 2004), but also involve a changed signature of electrical oscillations that define and characterize different sleep states (i.e., the *micro-structure* of sleep; cf. Box 1; Carrier et al., 2011; Crowley et al., 2002). In particular, older adults' sleep is characterized by a drastic reduction of deep non-rapid eye movement (NREM) sleep, so-called slow-wave sleep (SWS), that is accompanied by an increase of light NREM sleep (Carrier et al., 2011; Danker-Hopfe et al., 2005; Ohayon et al., 2004; Redline et al., 2004). With advancing age, high-amplitude slow oscillations (< 1 Hz), slow delta waves (1–4 Hz), and discrete sleep spindles (11–16 Hz) defining NREM sleep appear less often and with reduced amplitudes and altered topography (Crowley et al., 2002; Dubé et al., 2015; Fogel et al., 2012; Landolt and Borbély, 2001; Landolt et al., 1996; Martin et al., 2013). Diminished homeostatic sleep pressure, circadian shifts, dysregulation of neurotransmitters, and structural brain alterations may contribute to these pronounced changes in sleep during aging (Mander et al., 2017; Zhong et al., 2019).

Standard criteria to describe these age-related alterations in sleep physiology are only rarely tested and validated within heterogeneous samples that include older adults (but see, e.g., Ujma et al., 2015, 2019; Warby et al., 2014). Therefore, in the following, we aim to demonstrate the possible benefits and limitations of applying established indicators of sleep physiology in the context of aging research. We aim to raise awareness for the complexity of aging that entails widespread structural and functional brain changes that affect sleep in multiple ways (Dubé et al., 2015; Fogel et al., 2012, 2017; Muehlroth et al., 2019b). To identify meaningful biomarkers that can explain or even predict cognitive decline in old age, we need to promote the use of sensitive and age-fair methodology that captures within-person

age-dependent changes in the physiology of sleep, rather than age-inflated inter-individual differences (Lindenberger et al., 2011).

Preamble: How do we define 'old age' and 'aging'?

As an inevitable prerequisite, the identification of changes in sleep physiology during aging requires an appropriate and transparent rationale on how research should delineate and investigate 'old age' and 'aging.' The literature in (cognitive) neuroscience is dominated by cross-sectional group comparisons of 'younger' (~20–30 years) and 'older' adults (~60–80 years) (Browning and Spilich, 1981; Hedden and Gabrieli, 2004). This is grounded in the notion that the brain's structure and function has already undergone profound alterations when humans reach their 60's (e.g., Li et al., 2004; Ohayon et al., 2004; Raz et al., 2005; Rönnlund et al., 2005; Ziegler et al., 2012). Beyond stressing the importance of a general agreement on the definition of 'old age' to ensure the comparability of research on aging, we want to highlight two challenges that accompany such a definition: First, the *chronological* age of individuals can deviate from their *physiological* age. Second, age *differences* derived from such group comparisons cannot capture aging in terms of age-related *change*.

First and foremost, one should bear in mind that *chronological* age, i.e., the time that has passed since birth, is a variable that does not induce or explain any developmental change in itself (Wohlwill, 1970). At best, it can serve as a proxy for an individual's expected functional capacity that relies on true mechanistic alterations taking place as humans age (Li and Schmiedek, 2002; MacDonald et al., 2011). The conceptual and methodological pitfalls of using chronological age as an index of development or aging have been discussed elsewhere in detail (e.g., Cole et al., 2017; Lindenberger and Pötter, 1998; MacDonald et al., 2011; Wohlwill, 1970). Due to great inter-individual variation in the manifestation of aging effects (Cabeza et al., 2018; Habib and Nyberg, 2007; Lindenberger, 2014), particularly in old age, chronological age can deviate greatly from an individual's physiological age (Cole et al., 2017; MacDonald et al., 2011; Steffener et al., 2016). Importantly, physiological brain age, as for instance derived from estimates of structural brain integrity, could represent a better and more informative predictor of differences in aging, individual (brain) health, functional capacity, and mortality (Burzynska et al., 2013; Cole et al., 2018, 2017; Franke et al., 2010). Notably, also brain age derived from the sleep electroencephalogram (EEG) has lately shown its potential to serve as a sensitive index of aging and pathology (Sun et al., 2019). In the course of this paper, we will suggest that the sensitivity of sleep

physiology as a potential biomarker of brain age, aging, and age-related disease is closely tied to an age-fair investigation and evaluation of sleep in aged individuals.

Strictly speaking, the term aging describes a *process of change*, taking place as organisms grow older. Cross-sectional age comparisons are not appropriate to inform us about these changes (Hertzog and Nesselroade, 2003; Hofer and Sliwinski, 2001; Li and Schmiedek, 2002; Lindenberger and Pötter, 1998; Lindenberger et al., 2011; Overton, 2010; Raz and Lindenberger, 2011). When sampling age-heterogeneous groups cross-sectionally, essential confounds such as cohort effects or sampling bias may covary with age and thus prevent or bias conclusions on intra-individual dynamics of change (Hertzog and Nesselroade, 2003; Hofer and Sliwinski, 2001; Li and Schmiedek, 2002; Lindenberger et al., 2011; Rönnlund et al., 2005; Wohlwill, 1970). For convenience, however, we will use the phrase aging independently of the respective study's sampling design in this article. The lack of longitudinal studies on sleep in adulthood does not allow for the identification of 'real' age-related changes of sleep physiology. Hence, we emphasize that longitudinal study designs are essential to study aging and to identify the lead-lag relations between age-related alterations in brain structure and function (Hofer and Sliwinski, 2001; Li and Schmiedek, 2002; Lindenberger, 2014; Raz and Lindenberger, 2011).

Box 1. Sleep electrophysiology at a glance

In most animal species, three different and discrete states of vigilance can be differentiated: wakefulness, REM (rapid eye movement), and NREM (non-rapid eye movement) sleep (Vassalli and Dijk, 2009). REM and NREM sleep can be clearly distinguished based on electrophysiological characteristics (Aserinsky and Kleitman, 1953; Dement and Wolpert, 1958; Feinberg and Evarts, 1969; Loomis et al., 1935, 1962). REM sleep is marked by the occurrence of phasic irregular and rapid eye movements, muscle atony, and desynchronized wake-like electroencephalographic activity. NREM sleep, by contrast, is characterized by synchronous, low-frequency, high-amplitude EEG oscillations (Iber et al., 2007). NREM sleep can further be divided into three to four sleep stages (Iber et al., 2007; Rechtschaffen and Kales, 1968). These NREM sub-stages form a continuum of increasing arousal threshold and sleep depth (Carskadon and Dement, 2011). At the same time, they serve as a proxy for the dominance of different sleep-specific oscillatory brain signals: Lighter NREM sleep (stage 2 or N2 sleep) is defined by the occurrence of discrete sleep spindles (i.e., rhythmically waxing and waning oscillatory events with a typical frequency of 11–16 Hz), and K-complexes (i.e., phasic negative high-amplitude EEG deflections that are followed by a positive EEG component). With deeper sleep, the number and power of EEG slow waves increases. This predominance of slow EEG waves defines the presence of SWS (stage 3 or N3 sleep).

Challenge 1: Ambiguous sleep stage definitions across age groups

*Standardized visual scoring of sleep stages is regarded as the "gold-standard" of sleep research. Defined sleep stages are thereby considered sensitive indicators of specific physiological events. However, current sleep stage definitions may not capture the same physiological processes across age groups. We suggest that genuine sleep analysis – in line with similar views from animal research – should consider the continuous nature of NREM sleep. A focus on the presence of predefined electrophysiological events, like the occurrence or coupling of slow oscillations and sleep spindles, during **both** stage 2 sleep and SWS is required.*

Pitfalls of visually scoring SWS

In human research, the classification of sleep stages (cf. Box 1) is based on widely accepted standardized rules that are used to score sleep by means of EEG activity, muscle tone, and eye movements (Iber et al., 2007; Rechtschaffen and Kales, 1968). SWS, as defined by the *American Academy for Sleep Medicine* (AASM; Iber et al., 2007), requires the presence of 0.5–2 Hz waves with a minimum amplitude of 75 μV covering more than 20 % of a sleep time segment. Already in 1982, Webb noted that the frequently reported striking age-related reduction in SWS likely results from the use of a fixed amplitude threshold to define slow waves (Webb, 1982; Webb and Dreblow, 1982; cf. Figure 1A, Table 1). As also acknowledged by the AASM (Silber et al., 2007), reliably measuring SWS using an amplitude criterion below 75 μV is possible. Still, the current version of the manual for sleep scoring that defines the prevailing gold-standard, relies on a minimum amplitude criterion of 75 μV. Crucially, the age-related reduction in slow wave (< 4 Hz) and slow oscillation (< 1 Hz) amplitudes is one of the most consistent findings in age-comparative sleep studies (Carrier et al., 2011; Dubé et al., 2015; Mander et al., 2017; Muehlroth et al., 2019b; Ujma et al., 2019). Typically, SWS is easily identifiable in younger adults because slow waves exceed the required amplitude threshold. By contrast, the identification of true SWS in older adults is challenging: Due to the reduction of slow wave amplitudes below 75 μV (cf. Figure 1B and C, Table 2), time segments can often not be scored as SWS but are rather assigned to stage 2 sleep – although all other characteristics of the sleep epoch suggest the presence of SWS (i.e., the predominance of low-frequency EEG waves but absence of eye movements along with reduced muscle tone; cf. Figure 2).

Re-analyzing our own previously published data from 29 healthy younger (range: 19–28 years, M_{age} = 23.55 years, SD_{age} = 2.58; 16 females) and 34 healthy older adults (range: 63–74 years, M_{age} = 68.85 years, SD_{age} = 3.11; 15 females), we demonstrate the effects of fixed amplitude thresholds for visual sleep scoring in age-comparative studies. Participants' nighttime sleep was monitored using ambulatory polysomnography (PSG) before and after completing an associative learning task (cf. Supplementary Methods; Muehlroth et al., 2019a,b). We show that, overall, in our dataset the extent of age-related reductions in visually scored SWS matched the increase in stage 2 sleep in older adults (Cohen's $d_{stage\ 2}$ = –2.03 [–2.49; –1.58]; Cohen's d_{SWS} = 2.02 [1.57; 2.48]; Table 1). Visually scoring the data from older adults, a reduction – and in some participants even "absence" – of SWS became evident in comparison to younger adults (Table 1, Figure 1A). Closely inspecting the sleep architecture of older adults in our sample, it became evident that the older age group was divided into subjects not exhibiting any or only few SWS, and subjects showing a more "youth-like" proportion of SWS (Figure 1A). In part, this bisection of older subjects might be due to inter-individual differences in the amplitude of slow oscillations, with half of the older adults showing slow oscillation amplitudes below the proposed 75 μV threshold on average (Md_{OA} = 68.75 μV; Figure 1B). For older adults, falling above or below this threshold divided the sample into individuals with and without SWS according to the standard scoring system (Figure 1C). None of the younger adults undercut an average slow oscillation amplitude threshold of 75 μV (Md_{YA} = 121.94 μV) and, accordingly, SWS could be effectively scored in this age group (Figure 1B and C).

Taken together, it appears that due to age-related reductions in slow wave amplitudes, the fixed amplitude criterion of 75 μV for scoring SWS results in a biased estimate of SWS presence in older populations. Descriptively providing standardized estimates of sleep architecture is important to ensure the comparability and validity of studies. In studies with heterogeneous samples, though, researchers should acknowledge that such derived sleep stage estimates may be severely biased. As Rechtschaffen and Kales themselves put it in 1968 when establishing the standardized scheme for visual sleep scoring: *"Even among human subjects, however, there are some individuals or groups whose polygraph recordings may require further description or elaboration than that provided by the stages proposed here"* (Rechtschaffen and Kales, 1968).

Figure 1. Visual sleep scoring in younger and older adults. (A) Distribution of visually scored sleep stages in younger (orange) and older (blue) adults. In older adults, the proportion of SWS is significantly reduced, whereas lighter stage 2 sleep is increased. Note the bimodal distribution of SWS in older adults with some participants displaying levels of SWS comparable to younger adults, and others clustering around 0 %. (B) Distribution of the average slow oscillation amplitude for all younger (orange) and older adults (blue). Age group medians are inserted as dashed lines. The median slow oscillation amplitude in older adults lies below the amplitude criterion of 75 µV (green line). (C) The proportion of visually scored SWS scales with the average slow oscillation amplitude (respective age group medians marked by dashed lines). Older adults falling below the 75 µV criterion (inserted in green) show little or no SWS. The smoothed local regression line (in black, fitted using LOESS) and the corresponding standard error (grey shading) further illustrate the association. SO: slow oscillation.

Table 1. Age differences in sleep architecture

	Younger Adults	Older adults			Cohen's *d*
	Md [1st qrt; 3rd qrt]	*Md* [1st qrt; 3rd qrt]	Z	p	[95 % CI]
TST (min)	462.00 [428.00; 494.88]	417.50 [371.50; 459.50]	2.60	.001	0.6 [0.22; 0.96]
WASO (min)	6.25 [2.50; 12.00]	20.50 [9.75; 33.75]	3.47	<.001	–0.75 [–1.13; –0.37]
Stage 1 (%)	3.73 [2.63; 6.05]	7.10 [4.78; 9.96]	–5.20	<.001	–0.90 [–1.29; –0.52]
Stage 2 (%)	51.83 [46.40; 56.86]	65.51 [61.52; 69.82]	–4.80	<.001	–2.03 [–2.49; –1.58]
SWS (%)	20.06 [15.90; 27.33]	1.45 [0.00; 11.27]	–8.08	<.001	2.02 [1.57; 2.48]
REM (%)	23.32 [19.35; 26.62]	21.65 [16.73; 24.58]	1.86	.063	0.35 [–0.03; 0.72]

Note. Z and p-values were derived from non-parametric Mann-Whitney *U* tests comparing sleep parameters between younger and older adults. Sleep stage percentages are calculated as proportions of TST. CI: confidence interval; qrt: quartile; TST: total sleep time; WASO: wake after sleep onset (i.e., time awake between sleep onset and final morning awakening); SWS: slow-wave sleep; REM: rapid eye movement sleep.

The need to acknowledge the continuous nature of NREM sleep

The potential bias of current sleep stage definitions across age groups has direct theoretical and methodological implications. These affect the way we should conceive and treat NREM sleep and its sub-stages in our theories and analyses. The transition between stage 2 and SWS happens continuously with slow waves increasing in amplitude and abundance as NREM sleep gets deeper. On the contrary, sleep spindles show a reversed pattern with predominance during lighter NREM sleep. Hence, these two sleep stages, which reflect different poles of the NREM sleep continuum, are often treated as proxies for the dominance of different sleep-specific oscillatory brain signals (Genzel et al., 2014; Hobson, 1968). Although generally accepted in human sleep research, the distinction between sleep stage 2 and SWS based on a certain amplitude and time criterion (Iber et al., 2007) is not without problems (Genzel et al., 2014; Lacroix et al., 2018; Webb, 1982; Silber et al., 2007). Challenges include the handling of potentially biased sleep stage estimates in different age groups and the comparability of estimates between human and animal research (e.g., Genzel et al., 2014; Lacroix et al., 2018).

As discussed above, the current criteria to define the presence of SWS along with great inter-individual variation in slow wave amplitudes, may severely bias the assignment of a specific sleep segment to stage 2 sleep or SWS. Between the ages of 5 and 90, the proportion of NREM sleep decreases only little (Ohayon et al., 2004). The composition of visually scored NREM sleep itself, however, changes drastically across the lifespan. Typically, visually scored SWS is reduced or even absent in old age (Danker-Hopfe et al., 2005; Ohayon et al., 2004) (Figure 1A, Table 1). This might at least partly stem from the fact that time segments "equivalent" to SWS may be scored as stage 2 sleep because the amplitude criterion of 75 µV is not met. This results in two major problems: First, the assignment of SWS-like time segments to stage 2 sleep in older adults blurs the boundary between the two stages. As a consequence, sleep stages can no longer be regarded as a direct reflection or proxy of underlying neural and physiological processes (Hobson, 1968). Second, due to skewed distributions and reduced variance, any correlational analysis relying on measures of SWS is biased, or even prevented, in samples of older adults – especially in samples of older adults with a large percentage of "no-SWS" participants. As a result, a clear-cut association between a given physiological state – identified via a corresponding sleep stage – and an outcome measure, like the ability to maintain memories across sleep (e.g., Diekelmann and Born, 2010; Rasch and Born, 2013), is hardly tangible in age-comparative studies. If the definitions used to identify a given sleep stage do not capture the same phenomenon (e.g., the presence of slow oscillations) to the same degree across age groups, age differences in associations with outcome measures

can hardly capture the intended mechanism. For example, the observation that memory performance is more consistently associated with SWS in younger than in older adults, could be taken as evidence that slow oscillations forfeit their functionality for memory processing during aging (e.g., Baran et al., 2016; Scullin, 2013; Sonni and Spencer, 2015). However, if slow oscillations are present in older adults but just attributed to a different sleep stage, the statistics might be correct, but the functional interpretation would be different. In this case, slow oscillations found in a different sleep stage may continue to contribute to the consolidation of memories even in old age – a matter of fact only revealed when focusing on the oscillatory phenomena themselves (e.g., Helfrich et al., 2018; Mander et al., 2013; Muehlroth et al., 2019b; Varga et al., 2016; Westerberg et al., 2012).

Given the described challenges a NREM sleep subdivision bears, it may be hardly surprising that research in other species refrains from applying such a subclassification (but see Lacroix et al., 2018, for a recent attempt for 'human-like' subclassification of NREM sleep in rodents). The term "SWS" is typically used to describe the whole continuum of NREM sleep, which has evoked pivotal misunderstandings when comparing human and animal research (Genzel et al., 2014). This conception of NREM sleep may arise from the prevailing sleep scoring procedure in rodents that, can easily distinguish between REM and NREM sleep on the basis of movements and spectral parameters of electrical neuronal activity during sleep, but less so between NREM sub-stages (Bagur et al., 2018; Lacroix et al., 2018). In humans, contemplating whole-night spectral EEG data irrespective of participants' age, the distinct natures of REM and NREM sleep become evident, whereas differences between SWS and stage 2 sleep are more subtle (cf. Figure 2). Crucially, this does not signify that NREM sleep should be regarded as a uniform state, rather that the continuous nature of NREM sub-stages must be acknowledged, which makes a clear-cut distinction and definition of stage 2 and SWS problematic, if not impossible.

To conclude, an explicit distinction between stage 2 and SWS bears not only methodological but also theoretical difficulties. A coherent methodology to study the complex and dynamic nature of sleep not only within, but also between species is required to enable informative study comparisons. Research on the causes and consequences of inter-individual and age-related differences in sleep physiology should thus transcend stage-based analyses and rather consider the actual *exploranda*, ie., the underlying neurophysiological processes (for instance, the occurrence of rhythmic thalamocortical sleep spindles) present during **both** stage 2 sleep and SWS.

Figure 2. Whole-night spectral EEG characteristics. Hypnogram (upper panels) from one exemplary younger (A; female, 24.63 years) and older (B; female, 71.55 years) adult with the corresponding log-transformed whole-night spectral EEG data at electrode Cz (lower panels). Both younger and older adults show a similarly increased low-frequency power during periods of NREM sleep along with a distinct increase in power in the fast spindle frequency range (approx. 12.5–16 Hz) that disappears during REM sleep. Periods of wake are characterized by an additional strong enhancement of alpha (8–12 Hz) and high frequency power. Although the older adult only has a small amount of visually scored SWS, the spectrogram clearly shows periods of strongly enhanced low frequency and spindle power – in line with the canonical SWS characteristics found in younger adults. No clear distinction between stage 2 and SWS is observed on the basis of spectral EEG characteristics. Note the differences in absolute sleep duration (x-axis). YA: younger adult; REM: rapid eye movement sleep; SWS: slow-wave sleep; OA: older adult.

Challenge 2: Multiple ways to describe sleep physiology

The multitude of indicators and available algorithms used to describe sleep-specific neural activity hampers the comparability of results between studies. We argue that the validity and value of age-comparative sleep research depends greatly on the precise definition of indicators that reflect a neural process of interest as directly as possible. In our view, indicators of sleep micro-structure should always be preferred as they more closely mirror the dynamics and characteristics of sleep-specific neural activity.

Everyone engaging in sleep research will soon realize the multitude and inconsistency of indicators available to describe neurophysiological activity during sleep (Supplementary Figure 1). Comparing different indicators of slow oscillatory activity during NREM sleep in our own data set (namely % SWS, relative SWA, and the number, density, frequency, and amplitude of detected slow oscillations), we noticed a generally poor agreement between the different variables both across and within age groups ($ICC = 0.180$ [–.07; .39], $F(116,580) = 1.22$, $p = .0734$; $ICC_{YA} = 0.11$ [–.31; .44], $F_{YA}(53,265) = 1.13$, $p_{YA} = .266$; $ICC_{OA} = 0.08$ [–.32; .39], $F_{OA}(62,310) = 1.09$, $p_{OA} = .314$; cf. Supplementary Table 1). In contrast, the test-retest reliability of the same variable between two consecutively recorded nights was good to excellent (all $ICC \geq .78$, all $F \geq 4.61$, all $p < .001$; cf. Supplementary Table 1). A moderate test-retest reliability of relative SWA in the group of older adults was the only exception ($ICC = .68$ [.33; .85], $F(33,33) = 3.69$, $p < .001$). Critically, all participants engaged in an extensive associative learning task between the two recorded nights (cf. Muehlroth et al., 2019a,b). As demanding learning is known to influence indicators of spindle and slow oscillatory activity (e.g., Gais et al., 2002; Mölle et al., 2011, 2009), the reported estimates should even be considered as the lower bounder of reliability. Taken together, even though sleep physiology can be very reliably measured by applying common analytic approaches, the great disparity of available indicators calls for a thorough consideration of the way we describe sleep physiology in our analyses.

In general, the validity of indicators to quantify neural activity during sleep scales with their resolution and precision: Age differences in sleep are best captured by describing the dynamics and features of neural activity in as much detail and as directly as possible. Using machine learning on 3500 sleep recordings covering the ages 18 to 80, Sun and colleagues (2019) recently demonstrated that oscillatory dynamics during sleep – like fluctuations in spectral power – (i.e., the *micro-structure* of sleep) can reliably predict participants' age (mean absolute deviation 7.8 years; see Sun et al., 2019).

Global sleep stage composition (i.e., the *macro-structure* of sleep), in contrast, turned out to be a less appropriate predictor of chronological age (mean absolute deviation 23.3 years; Sun et al., 2019). In line with these results, we believe that the analysis of sleep EEG data should transcend beyond the report of sleep stages and global spectral features and focus on the micro-structure of sleep.

To avoid using the proportion of SWS as a proxy for the presence of slow waves, analyses increasingly report on *slow-wave activity* (SWA). SWA measures the spectral delta power (i.e., 0.5–4.5 Hz) via a transformation of the data in the frequency domain (e.g., a Fast Fourier Transformation, FFT). In conjunction with a normalization procedure – e.g., dividing the respective frequency band by the total power of the whole spectrum – the influence of possible confounds such as skull thickness can be minimized (see, e.g., Dannhauer et al., 2011; Frodl et al., 2001; Leissner et al., 1970; Segalowitz and Davies, 2004; Werkle-Bergner et al., 2009, for arguments on how inter-individual differences in anatomical properties of the brain and skull influence the EEG signal). A normalized measure of SWA increases the comparability across diverse study populations. However, as spectral power conflates both the duration and the amplitude of rhythmic neural activity (e.g., Kosciessa et al., 2018), an interpretation of SWA in terms of abundance or strength of slow waves remains difficult (Amzica and Steriade, 1998; Mensen et al., 2016).

Technological developments over the past decades are increasingly facilitating analysis of specific features and characteristics of neural events during sleep (Mensen et al., 2016; Mölle et al., 2002). Such analyses can incorporate the amplitude, oscillatory frequency, the density of occurrence (i.e., number of events per time interval), and the slope of specific oscillatory events. These different indicators are often used interchangeably although they signify differential physiological properties and, accordingly, show differential age dependency (cf. Box 2, Figure 4; cf. Supplementary Figure 1). If applied correctly, though, they hold great explanatory power and can provide more informed insights into the aging brain (Fogel et al., 2017; Sun et al., 2019; Ujma et al., 2019). Moreover, the possibility to identify individual oscillatory events also entails the opportunity to investigate how different oscillations and neural processes interact (Helfrich et al., 2018; Klinzing et al., 2016; Muehlroth et al., 2019b; Staresina et al., 2015). At the moment, due to the variety of algorithms available to study oscillatory brain signals and their interaction during sleep, these analyses are still time-consuming and between-study comparability of indicators can be challenging (Figure 3; cf. Warby et al., 2014). Identifying and resolving of these discrepancies will lay the grounds for a mechanistic understanding of the causes and consequences of age-related changes in sleep (Helfrich et al., 2018; Muehlroth et al., 2019b).

Box 2. Guidance for the interpretation of frequently reported oscillatory characteristics

Spectral power. The spectral power in predefined frequency bands is believed to mirror the presence of rhythmic neural activity, i.e., periodic alternations in the excitability of neuronal populations (Buzsáki, 2006; Buzsáki et al., 2016; Cohen, 2014). Conventional estimates of spectral power index a combination of the duration and the amplitude of oscillations within a specific frequency range (Cohen, 2014; Kosciessa et al., 2018).

Amplitude. The amplitude of an oscillation reflects the degree of synchronized neuronal firing: Small amplitudes relate to more local synchronous firing, whereas large amplitudes reflect more global synchronous changes in membrane potentials (Buzsáki, 2006; Nir et al., 2011). The basis for this cortical synchronization is the integrity of cortical areas involved in the generation and propagation of a certain oscillation (Dubé et al., 2015; Saletin et al., 2013). In aging, cortical thinning and atrophy may diminish the scalp-detectable range of amplitudes. Age differences in oscillatory amplitudes may, on the one hand, reflect a minimized extent of synchronous neuronal firing as direct consequence of structural brain changes in the course of aging (Dubé et al., 2015). On the other hand, anatomical alterations of the brain and skull may influence the conduction of the EEG signal and thus attenuate the amplitude of an oscillation measured on the scalp (Leissner et al., 1970; Segalowitz and Davies, 2004). Besides, reduced slow wave amplitudes in old age may arise from alterations in homeostatic sleep pressure, which is typically reduced as humans age (Dijk et al., 2000; Esser et al., 2007; Vyazovskiy et al., 2009).

Frequency. In general, the inherent frequency of an oscillation may index the efficiency and speed of neural information transfer (Dubois et al., 2008; Grandy et al., 2013; Klimesch, 1999; Posthuma et al., 2001). The typical developmental acceleration of various oscillatory components including sleep spindles (Campbell and Feinberg, 2016; Purcell et al., 2017) is believed to depend on the differentiation of thalamocortical feedback loops due to maturation, optimization of synaptic connections, and neuronal myelination (Campbell and Feinberg, 2016; Clawson et al., 2016; Dubois et al., 2008; Klimesch, 1999; Piantoni et al., 2013). Maintained or even increased frequency of fast spindles in old age (Crowley et al., 2002; Fogel et al., 2017; Nicolas et al., 2001; but see Martin et al., 2013) may indicate a largely preserved functionality of thalamocortical cells during aging. In contrast, the consistently reported slowing and flattening of slow oscillations in old age (Carrier et al., 2011; Dubé et al., 2015; Mander et al., 2017) may be the consequence of reduced synaptic load and homeostatic sleep pressure in old age and indicative of decelerated and less synchronous switches between phases of neuronal excitability and inhibition (Bersagliere and Achermann, 2010; Carrier et al., 2011; Dijk et al., 2000; Dubé et al., 2015; Fogel et al., 2012; Vyazovskiy et al., 2009).

Challenge 3: Amplitude reductions – towards algorithms with adjusted thresholds

The benefits of a direct investigation of slow oscillatory events are constrained by a lacking consensus on the detection thresholds to be applied. We suggest that an age-fair assessment of true slow oscillatory activity requires the use of amplitude thresholds that are adjusted individually within participants.

Studying rhythmic neural activity during sleep requires a reliable detection of oscillatory events themselves. Here, automated slow oscillation detection methods offer an efficient way to identify the presence and characteristics of individual slow oscillations (eg., Massimini et al., 2004; Mensen et al., 2016; Mölle et al., 2002; Nir et al., 2011). Typically, oscillation detection algorithms rely on a bandpass-filtered EEG signal within a theoretically implied frequency range (eg., 0.5–1 Hz for slow oscillations). The presence of true oscillatory components is then inferred by applying certain amplitude and peak detection criteria (cf. Supplementary Methods). Amplitude criteria can either be based on fixed thresholds or on variable thresholds adjusted within individuals (e.g., Massimini et al., 2004; Mölle et al., 2009, 2002; Staresina et al., 2015). In analogy to the visual scoring of SWS, many algorithms rely on a criterion of 75 μV (Carrier et al., 2011; Dubé et al., 2015; Piantoni et al., 2013) or even higher amplitudes (Heib et al., 2013; Massimini et al., 2004) for the detection of slow oscillations during NREM sleep. But an increasing number of studies opt for thresholds adjusted individually for each participant (e.g., Klinzing et al., 2016; Mölle et al., 2011; Ngo et al., 2015). As yet, there is neither a gold-standard or consensus on the most suitable method nor on the threshold settings to be applied when identifying the presence of oscillatory events (Mensen et al., 2016; Ujma et al., 2019). However, the choice of detection thresholds has profound effects on the analysis of age differences.

Irrespective of the applied detection threshold (i.e., a fixed threshold of 75 μV vs. an individually adjusted threshold of 1.25 times the mean amplitude of all putative slow oscillations; cf. Supplementary Methods), in the older adults in our data, the number, density (i.e., the number of slow oscillations per minute of NREM sleep), amplitude, and frequency of slow oscillations were all reduced (Table 2). Nevertheless, compared to an individually adjusted amplitude criterion, the extent of age effects was augmented when using a fixed detection threshold of 75 μV (non-overlapping confidence intervals for Cohen's *d* of slow oscillation number and density, cf. Table 2). Using individually adjusted thresholds, as displayed in Figure 1B, in older adults, the average slow oscillation amplitude fell below the 75 μV criterion in 58.06 % of the cases (compared to 1.96 % in younger subjects). On a median level,

69.52 % of the detected slow oscillations in older adults did not reach the amplitude criterion of 75 μV, compared to only 3.11 % in younger adults (Figure 3A) – in other words, all these slow oscillations are missed in case of a fixed amplitude criterion of 75 μV as the detection is biased towards higher-amplitude oscillations (Figure 3C). As a consequence, the congruency between slow oscillation density estimates that are derived using individually adjusted vs. fixed detection thresholds is higher in younger than in older adults. Accordingly, estimates are biased in an age-dependent manner (Figure 3B).

Critically, inter-individual differences in slow oscillation amplitudes might not only carry functional information but may be confounded by differences in homeostatic sleep regulation (Dijk et al., 2000), brain integrity (Dubé et al., 2015; Saletin et al., 2013), and skull thickness (Leissner et al., 1970; Frodl et al., 2001). Hence, an individually adjusted amplitude threshold that tags slow oscillations with the largest amplitudes within individuals may have advantages. Only when slow oscillations are selected independently of inter-individual differences in EEG amplitudes, do (age-)fair comparisons become possible. Persisting age differences in the amplitude, density, and morphology of slow oscillations can then be interpreted as valid age group differences. Compared to fixed amplitude detection criteria, especially in age-comparative settings, the use of individually adjusted amplitude criteria for detecting slow oscillations during NREM sleep provides a more realistic and at the same time age-fair measure of true slow oscillatory activity.

Table 2. Individually adjusted vs. fixed threshold settings for slow oscillation detection

	Younger Adults	Older adults			Cohen's *d*
	Md [1st qrt; 3rd qrt]	*Md* [1st qrt; 3rd qrt]	*Z*	*p*	[95 % CI]
Individually adjusted threshold					
number	1171 [1086.75; 1358.25]	915.25 [786.75; 1148.13]	5.06	<.001	1 [0.59; 1.4]
density (events/min)	3.76 [3.32; 4.16]	3.38 [2.93; 3.86]	3.02	.003	0.62 [0.23; 1.00]
frequency (Hz)	0.79 [0.78; 0.81]	0.77 [0.75; 0.79]	4.15	<.001	0.86 [0.46; 1.25]
amplitude (μV)	121.94 [104.93; 137.06]	68.75 [58.78; 86.66]	8.16	<.001	2.2 [1.72; 2.68]
Fixed threshold					
number	1353 [1060.50; 1687.25]	377 [128.00; 583.25]	7.86	<.001	1.93 [1.48; 2.39]
density (events/min)	4.23 [3.06; 5.39]	1.14 [0.46; 2.36]	7.8	<.001	1.83 [1.38; 2.28]
frequency (Hz)	0.79 [0.78; 0.81]	0.76 [0.73; 0.79]	5.04	<.001	1.12 [0.71; 1.52]
amplitude (μV)	120.48 [113.92; 135.07]	96.09 [90.31; 101.91]	8.09	<.001	2.04 [1.58; 2.51]

Note. *Z* and *p*-values were derived from non-parametric Mann-Whitney *U* tests comparing slow oscillation characteristics between younger and older adults. CI: confidence interval; YA: younger adults; OA: older adults; qrt: quartile.

Figure 3. Comparison of fixed (75 µV) and individually adjusted slow oscillation detection thresholds. (A) Amount of detected slow oscillations with an amplitude below 75 µV when using individually adjusted amplitude thresholds. Whereas in younger adults (orange) only few slow oscillations fall below the 75 µV amplitude criterion, in older adults (blue) a large proportion of the detected slow oscillations falls below this threshold. (B) Relation between slow oscillation density estimates derived by the use of individually adjusted vs. fixed detection thresholds. In younger adults (orange), the two detection thresholds produce comparable results. In older adults (blue), especially for individuals with small slow oscillation amplitudes (smaller square size), a fixed detection threshold results in an underestimation of slow oscillation density. (C) Individual average slow oscillations and age group averages of slow oscillations detected with individually adjusted and fixed amplitude thresholds. Left panel: In younger adults, both algorithms result in comparable slow oscillation detection. Right panel: In older adults, individually adjusted thresholds (blue) result in the detection of slow oscillations with smaller amplitudes compared to higher-amplitude fixed threshold detection (green). OA: older adults; SO: slow oscillation; YA: younger adults.

Challenge 4: Differential frequency shifts in sleep oscillations – towards individualized frequency ranges

The exact frequency band in which an oscillatory event can be observed varies between individuals and is influenced by age. To prevent missing true oscillatory events during detection and mixing up functionally distinct oscillatory components, cut-off frequencies have to be determined carefully. We argue that individually determined frequency ranges are the prerequisite to derive meaningful age- and person-specific characteristics of oscillatory components.

Contemplating frequency cut-offs for low-frequency oscillatory components

Age-related changes in the appearance of sleep oscillations do not only include alterations in amplitudes, but also entail shifts in the intrinsically expressed frequencies of the respective neural events. With respect to low-frequency components, a global slowing has consistently been reported (Carrier et al., 2011; Dubé et al., 2015; Mander et al., 2017) (Figure 4I, Box 2). These effects comprise both slow oscillations (< 1 Hz) as well as slow waves with higher frequencies (1–4 Hz). Although slow oscillations and slow waves constitute different entities on a theoretical level (Achermann and Borbély, 1997; Amzica and Steriade, 1998; Steriade et al., 1993), the applied cut-off frequencies defining low-frequency EEG components are often not much attended to.

Empirically, the problem of identifying the exact frequency ranges for specific oscillatory components in individual participants can be solved by considering the NREM sleep power spectrum. Peaks in the power spectrum signify the presence of rhythmic neural activity at a given frequency (Aru et al., 2015; Kosciessa et al., 2018; Whitten et al., 2011). When visually inspecting the age-specific power spectra during NREM sleep in our data, it became evident that low-frequency power was not only globally reduced in older compared to younger adults, but that the peak in low-frequency power was shifted to even lower frequencies in older adults (Supplementary Figure 2). This slowing in low frequencies was statistically reliable (average frequency peak: $t(115) = 6.18$, $p < .001$, $M_{YA} = 0.73$ Hz, $M_{OA} = 0.53$ Hz) and was also evident for discrete slow oscillation events ($Z = 4.15$, $p < .001$, $Md_{YA} = 0.79$ Hz, $Md_{OA} = 0.77$ Hz; see Table 2, Figure 3C). Although statistically detectable, in contrast to the dramatic reduction in slow oscillation amplitudes, inter-individual and age differences in slow oscillation frequencies were relatively small ($d_{frequency} = 0.86$ [0.46; 1.25]; $d_{amplitude} = 2.2$ [1.72; 2.68], non-overlapping confidence interval; cf. Table 2).

Taken together, it appears that the magnitude of the slowing of slow oscillations in old age does not reach an extent that would bias detection results when established frequency ranges (e.g., 0.5–1 Hz for slow oscillations) are utilized. Nonetheless, we emphasize the need to carefully consider, justify, and report the selected frequency range when publishing sleep research findings on older adults. Slow oscillations and "faster" slow waves have been associated with differential homeostatic regulation (Bersagliere and Achermann, 2010; Werth et al., 1997) as well as with divergent roles in the context of cognitive pathology in aging (Lucey et al., 2019; Mander et al., 2015, 2016). Yet, detection algorithms often deploy frequency bands that comprise both oscillatory components (e.g., Carrier et al., 2011; Kurth et al., 2010). We thus underscore the need for clear and coherent terminology along with appropriate detection criteria when reporting on age differences in low-frequency oscillatory activity.

Inter-individual (age) variation in fast and slow spindle frequency bands

In contrast to slow oscillations whose detection seems mostly dependent on appropriate amplitude settings, the detection of sleep spindles is heavily influenced by the selection of frequency ranges. In particular, the detection of spindle events is challenged by the presence of two spindle types. Slow (ca. 9–12.5 Hz) and fast spindles (ca. 12.5–16 Hz) differ in frequency, topography (Figure 4) (De Gennaro and Ferrara, 2003; Doran, 2003; Klinzing et al., 2016; Schabus et al., 2007), generating mechanisms (Ayoub et al., 2013; Schabus et al., 2007; Timofeev and Chauvette, 2013), and not least in their functionality (Barakat et al., 2011; Fogel and Smith, 2011; Mölle et al., 2011). Although the existence of two spindle types is now widely acknowledged, there is no general consensus on the exact definition of the frequency range in which to expect slow and fast spindles should be expected (Cox et al., 2017). Critically, spindles undergo a complex developmental shift and differentiation in their inherent frequency, power and topography (Campbell and Feinberg, 2016; Crowley et al., 2002; Nicolas et al., 2001; Purcell et al., 2017). This not only challenges research on younger adults (Cox et al., 2017), but also research that focuses on the entire lifespan.

In line with these observations, slow and fast spindle frequencies were differentially affected by age in our data. Average fast spindle frequency was increased in older adults ($F(1,90) = 21.01$, $p < .001$, M_{YA} = 13.4 Hz, M_{OA} = 13.69 Hz) (Figure 4C, Box 2). In contrast, the average frequency of slow spindles was reduced in older adults (Figure 4F; cf. Fogel et al., 2017). This effect was only pronounced at electrode Cz, however (age by channel interaction: $F(3,270) = 16.16$, $p < .001$, $Z_{Cz} = 4.64$, $p < .001$, $Md_{YA} = 11.25$ Hz, $Md_{OA} = 10.54$ Hz; all other $Z \leq 2.37$, all other p $\geq$ .018, α-level adjusted to .0125). The complexity

of frequency shifts during aging is thus even enhanced by the topographical intricacy of the observed effects.

The multiple facets of age differences in sleep spindle appearance require the careful consideration of an age-fair definition of spindle frequency ranges. Based on individually identified peaks in the NREM sleep power spectrum at topographical locations where slow and fast spindles are expected (cf. Supplementary Figure 2), individually adjusted spindle frequency ranges can be defined – even if they might not match conventionally prescribed bands exactly (Adamczyk et al., 2015; Bódizs et al., 2009; Cox et al., 2017; Doppelmayr et al., 1998; Grandy et al., 2013; Klimesch, 1999; Ujma et al., 2015). Comparisons between spindle detection algorithms that use commonly predefined frequency ranges and those that adjust their frequency range definitions within individuals, provide a strikingly poor agreement with conventional (fixed frequency) algorithms. The latter have been shown to miss up to 25 % of fast and slow spindles (Adamczyk et al., 2015; Cox et al., 2017; Ujma et al., 2015). To allow for the investigation of differences in spindle characteristics, age-comparative research should ensure that the frequency bands applied to define slow and fast spindles are congruent with age- and person-specific spindle properties. Moreover, frequency definitions need to be broad enough to capture sleep spindles in their complexity and entirety. Independently of an individual's age, person-specific definitions of spindle frequencies form the basis for these improvements (Adamczyk et al., 2015; Cox et al., 2017; Ujma et al., 2015).

Figure 4. Topographical distribution of sleep measures. (A) Centroparietal prevalence of fast spindles in younger adults (orange), compared to a parietal-occipital prevalence in older adults (blue). (B) The amplitude of parietal-occipital fast spindles is preserved in older adults. (C) Overall increased average fast spindle frequency in older adults, with reduced effects at frontal channels. (D) Frontal prevalence in slow spindle density in both younger and older adults, as well as anterior to posterior attenuation of age differences. (E) Reduced slow spindle amplitude over centroparietal regions in younger adults, but more uniform amplitude modulation in older adults. (F) Age-comparable average slow spindle frequency at Oz, but, apart from that, decreases in slow spindle frequency in older adults. (G) Overall reduced slow oscillation density in older adults but preserved frontal dominance. (H) Overall reduced slow oscillation amplitude in older adults but anterior to posterior attenuation of age differences (I) Uniform decrease in slow oscillation frequency in old age at all channels. (J) Anterior dominance of relative SWA in both younger and older adults. Note. All slow oscillation and spindle characteristics were derived using individually adjusted amplitude thresholds. OA: older adults; SO: slow oscillation; SWA: slow-wave activity; YA: younger adults.

Figure 5. Age differences in fast spindle density mirror cortical integrity. (A) Cohen's *d* of age differences in fast spindle density at different electrode sites. Confidence intervals are displayed as error bars. Age differences are most pronounced at Cz but are attenuated over more posterior sites. (B) Sagittal, coronal, and axial depiction of the association between gray matter volume and age (yellow), frontal (violet), and Cz (cyan) fast spindle density (separate voxel-vise multiple regression models for each predictor, controlled for total intracranial volume). In contrast to a global age-related decrease in gray matter volume (see Supplementary Figure 3, Supplementary Table 2), associations with spindle density are more specific. Whereas inter-individual differences in fast spindle density at Pz and Oz are not related to gray matter volume, fast spindle density at Cz shows a strong association with structural integrity in frontal and temporal brain regions (see Supplementary Figure 4, Supplementary Table 3). Clusters are displayed and considered significant at a FWE-corrected voxel threshold of $p < .05$, and a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$. All clusters are corrected for nonstationary smoothness. FWE: family-wise error; YA: younger adults; OA: older adults.

Challenge 5: Topographical disparities in age-related sleep changes

Age-related alterations in sleep physiology do not show a spatially coherent pattern. We highlight that alterations in sleep oscillations that are measurable on the scalp do not necessarily reflect an altered initiation of underlying physiological sleep processes per se, but could rather represent age-related alterations in structural properties of the brain. Taking into account that intrinsic characteristics of sleep-specific oscillatory components may be well preserved during aging can help our understanding of the functionality of sleep in old age.

Across the lifespan, the human brain undergoes complex structural alterations (Buckner, 2004; Cabeza et al., 2005; Fjell et al., 2013; Fjell and Walhovd, 2010; Raz et al., 2005), which are also reflected in the topography of the sleep EEG (Buchmann et al., 2011; Carrier et al., 2011; Kurth et al., 2010; Landolt and Borbély, 2001; Martin et al., 2013; Sprecher et al., 2016). From childhood to adolescence, along with cortical maturation in frontal brain regions, maximal SWA shifts from posterior to frontal brain regions (Kurth et al., 2010; Whitford et al., 2007). In old age, frontal brain regions are particularly prone to age-related structural white and gray matter loss (Giorgio et al., 2010; Raz and Rodrigue, 2006). As a consequence, age-related decline in spectral EEG power and amplitude is particularly pronounced over frontal areas (Carrier et al., 2011; Landolt and Borbély, 2001; Mander et al., 2017; Martin et al., 2013; Sprecher et al., 2016).

Topographical shifts in sleep parameters are diverse. For example, in our own data, the centroparietal prevalence of fast spindle density and amplitude in younger adults was shifted posterior in older adults, resulting in reduced or even absent age differences over parietal and occipital regions (Figure 4A and B; Supplementary Figure 2). In contrast, slow spindles, slow oscillations, and relative SWA (cf. Supplementary Methods) maintained their frontal dominance in old age. Whereas relative SWA and the amplitude and density of slow oscillations were globally and uniformly reduced in old age (Figure 4G–J), the density and amplitude of slow spindles showed most pronounced age-related reductions over frontal areas (Figure 4D and E). Thus, the extracted oscillatory event or indicator (i.e., slow spindle, fast spindle, slow oscillation, and SWA), their derived characteristics (i.e., density, amplitude, and frequency), and not least an individual's age interact and influence where, and to what extent, age differences in sleep can be observed.

Crucially, the topographical heterogeneity of age effects detectable on the scalp level may be meaningful and indicative of alterations in underlying brain structure and cellular properties. To illustrate this effect, we tested for the association between gray matter volume and fast spindle density measured at different electrode sites (separate voxel-vise multiple regression models for each predictor; see Figure 5; cf. Supplementary Methods; Supplementary Figure 4, Supplementary Table 3). In line with the observed posterior preservation of fast spindles in old age (Figure 4A; Supplementary Figure 2), fast spindle density at Pz and Oz did not relate to gray matter volume. In contrast, fast spindle density measured frontally and at Cz, that was most reduced on older adults, was associated with structural integrity in frontal and temporal brain regions (Figure 5; Supplementary Figure 4, Supplementary Table 3). In comparison, gray matter clusters for the association with slow spindle and slow oscillation density were small and less consistent (Supplementary Figures 5 and 6, Supplementary Tables 4 and 5). Based

on these exemplary analyses, it becomes clear that structural brain integrity does not globally shape the generation and propagation of all oscillatory components. However, one should note that chronological age, in general, was associated with global reductions in gray matter volume that were strongest in frontal brain regions (Figure 5; Supplementary Figure 3, Supplementary Table 2). The extent of structural brain alterations in aging can thus overshadow more focalized and potentially meaningful structure–function associations. Moreover,

To derive a meaningful interpretation of the observed structure–function associations, one needs to consider that the extent to which we can detect oscillatory events on the scalp depends on both their generation and propagation. Thus, we can encounter the situation that oscillations are properly generated, but their propagation to certain brain areas is impaired by altered structural properties of the involved brain circuits (Landolt and Borbély, 2001; Martin et al., 2013; Nir et al., 2011; Sprecher et al., 2016). In line with this notion and the gray matter associations reported above, the maximum of fast spindle density over the posterior cortex has been interpreted in terms of aggravated fast spindle propagation to frontal and central brain regions that is severely affected by brain atrophy during aging. In contrast, the generation of sleep spindles is often regarded as less affected by aging (Crowley et al., 2002; Landolt and Borbély, 2001; Martin et al., 2013; Nir et al., 2011; Sprecher et al., 2016; see also Box 2). Still, we must acknowledge that the available cross-sectional correlative data neither allows us to disentangle whether differences in oscillatory components stem from disparities in their generation or their propagation, nor whether structural brain alterations cause, accompany, or result from alterations in sleep physiology. To solve these questions, longitudinal study designs that identify and track the sources of oscillatory components and link them to aging changes in structural brain integrity are required.

To conclude, the topographic distinctiveness of age-related sleep alterations is an important but often overlooked factor in age-comparative sleep studies. With the choice *where* to study age differences, age effects can either be maximized or reduced. Hence, minimal electrode setups may bias effect sizes in age-comparative studies. Crucially, the neglected topography of sleep parameters can cause pivotal misinterpretations. The fact that sleep parameters are selectively altered over specific brain regions, bit not over other areas, casts doubt on the reported extent of age-related impairments in sleep processes. Brain atrophy likely results in a diminished and altered generation of certain sleep parameters (Dubé et al., 2015; Mander et al., 2013), but may also affect the propagation of electrical events that, per se, may be less prone to age-related changes. Future research should interpret the reported effects with care and provide solutions to disentangle age-related impairments in slow wave and spindle generation and propagation. Rather than placing a continued focus on the deficiency of sleep in old age, age-fair

sleep studies require a shift towards maintained and intact sleep characteristics to determine altered or preserved functionality of sleep in old age.

Conclusion

Fundamental changes in human sleep during aging are consistently reported and easy to observe – but, as discussed in this paper, it is nontrivial to quantify and measure them. Intact sleep and the functional succession of different sleep states is enabled by the balance of different neurotransmitter systems (Aston-Jones et al., 2007; Cirelli et al., 2005; Ramm, 1979; Steriade et al., 1990), and synergistic activity in different cortical and sub-cortical brain regions (Hobson, 2005; Saletin et al., 2013; Steriade et al., 1990). Furthermore, it is embedded in a system of regulatory and homeostatic processes (Achermann and Borbély, 1999; Dijk et al., 2000). As complex as sleep is in its regulation and organization, as manifold and diverse are the according age-related changes in sleep (Dubé et al., 2015; Fogel et al., 2012, 2017; Mander et al., 2017; Muehlroth et al., 2019b; Zhong et al., 2019).

Standard criteria applied to measure these alterations are commonly optimized to capture inter-individual variance in healthy younger adults but can be severely biased when applied in other populations. As a result, not all age differences documented via the use of conventional analysis pipelines capture real alterations in underlying neural processing. We thus stress the need to develop age-fair and individually adjusted analytic approaches in order to disentangle true age-dependent effects from age-independent inter-individual differences. But how could such an age-fair research process work?

We suggest that the basis for research on age-dependent differences in a given (neural) process is rooted in thorough theoretical considerations that incorporate the complex and dynamic nature of aging (e.g., Cabeza et al., 2018, 2005; Lindenberger et al., 2006) and the individual as the primary unit of analysis (Molenaar and Campbell, 2009; Nesselroade et al., 2007; Nesselroade and Molenaar, 2016; Rose et al., 2013). Ideally, such a research process starts out with an exact definition of a neural or physiological phenomenon that we intend to investigate. For instances we could ask whether the *intensity* of sleep (in terms of sleep depth) is reduced in old age. In a next step, potential indicators that mirror the phenomenon of interest along with its crucial features have to be considered and the most appropriate indicator(s) need(s) to be selected. In our example, we could decide to use SWA and slow oscillation density as markers of low-frequency oscillatory activity during sleep that index the depth of sleep (e.g., Achermann and Borbély, 1999; Bersagliere and Achermann, 2010; Blake and Gerard, 1937). Next, we need to determine how to best measure the respective indicator. Here, the individual has to be at the center

of the analysis. Inter-individual differences that are irrelevant for our research question and thus introduce "noise" (e.g., differences in anatomical skull properties) should be filtered out by adjusting analytic detection criteria at the level of the individual person (Karch et al., 2015; Molenaar and Campbell, 2009; Nesselroade et al., 2007; Nesselroade and Molenaar, 2016; Rose et al., 2013). In this way, we can create the conditions that facilitate an accurate interpretation of remaining inter-individual differences (Nesselroade et al., 2007). To be as (age-)fair as possible, we thus believe that research on sleep and aging should strive to identify the best possible analysis strategy to study a certain phenomenon of interest – even if this means moving away from conventional analytic approaches.

Complementarily, we stress the inevitable need to shift research foci from cross-sectional towards longitudinal studies. The currently available evidence cannot inform about 'real' age changes in sleep (Hertzog and Nesselroade, 2003; Hofer and Sliwinski, 2001; Li and Schmiedek, 2002; Lindenberger and Pötter, 1998; Lindenberger et al., 2011; Overton, 2010; Raz and Lindenberger, 2011). To disentangle adaptive from maladaptive aging processes, and to identify the lead-lag relations between age-related alterations in brain structure and sleep, study designs that follow the same individual across multiple measurement occasions are required (Hofer and Sliwinski, 2001; Li and Schmiedek, 2002; Lindenberger, 2014; Raz and Lindenberger, 2011).

Finally, we want to emphasize that age-dependent alterations in sleep physiology should not necessarily be interpreted in terms of "shortage" or "deficiency." The brain is an *adaptive* organ that changes its structure and function in reaction to mismatches between environmental demands and organismic supplies (Lövden et al., 2010; Park and Reuter-Lorenz, 2009). First, this means that not all observed age differences in sleep necessarily indicate a loss (Baltes, 1987; Baltes et al., 1998). Reduced SWA, for instance, might merely be the consequence of reduced sleep pressure and an altered need for sleep in old age (Dijk et al., 2000). Second, due to the adaptive nature of the human brain, persons with very different developmental trajectories and physiological conditions can display similar sleep patterns (Muehlroth et al., 2019a). By identifying the similarities among individuals, and specifically among younger and older adults (Nesselroade et al., 2007), we can determine whether and how sleep is preserved in old age. Moreover, we can determine whether maintained sleep physiology is also reflected on the level of behavioral outcomes (Cabeza et al., 2018; Muehlroth et al., 2019a; Nyberg et al., 2012; Nyberg and Pudas, 2019; Park and Reuter-Lorenz, 2009). Age-fair research that shifts its focus towards the successful retention of sleep characteristics represents the foundation to gain a mechanistic understanding of the consequences entailed in altered sleep physiology. This is the basis for identifying the diagnostic value of sleep changes (Ferrarelli et al., 2019; Grandy et al., 2013; Prinz, 1977).

Recent attempts to establish sleep as a novel biomarker and treatment target for Alzheimer's disease have instantly evoked a massive rise in research on sleep and aging (Ju et al., 2014; Mander et al., 2016; Noble and Spires-Jones, 2019; Vaou et al., 2018). Methodological considerations, though, call for a clarification of *what* we want to measure and *how* we can best assess the respective phenomena. Improved methodological approaches will clearly speed up the search for a mechanistic understanding of the association between sleep and cognition in aged individuals.

Box 3. Methodological advice for age-comparative sleep studies

1. Closely inspect your sleep EEG data visually. Knowing what the data looks like will ease your understanding of what to consider and what to adjust in your analyses.

2. Analyses should focus on micro-structural rather than macro-structural sleep features.

3. Define the neural process of interest as well as possible. Avoid proxies wherever possible.

4. Consider that seemingly similar measures (e.g., frequency, amplitude, density, power of the very same spindle event) can reflect very different physiological and neuronal properties. Be precise on which features you extract and interpret them carefully.

5. NREM sleep forms a continuum. Acknowlegde that neurophysiological processes that are captured by sleep spindles and slow oscillations are present during both stage 2 sleep and SWS.

6. Use individually adjusted algorithms (with regard to amplitude and frequency) for sleep oscillation detection.

7. Use electrode setups that enable topographical comparisons. Try to incorporate the mechanisms that are driving topographical heterogeneity in age effects into your theories and methods (e.g., by carrying out structural brain imaging or source localization techniques in addition).

8. Instead of focusing on deficits only, consider what is intact or maintained in old age. Do your very best to apply age-fair methodology and comparisons.

Acknowledgements

The authors thank Ann-Kathrin Joechner for helpful comments on an earlier version of the manuscript and Julia Delius for editorial assistance. We are grateful to everyone who contributed to the collection of data utilized in this article.

References

Achermann, P. and Borbély, A. A. (1997). Low-frequency (< 1H z) oscillations in the human sleep electroencephalogram. *Neuroscience*, 81(1):213–222.

Achermann, P. and Borbély, A. a. (1999). Sleep homeostasis and models of sleep regulation. *Journal of Biological Rhythms*, 14(6):559–568.

Adamczyk, M., Genzel, L., Dresler, M., Steiger, A., and Friess, E. (2015). Automatic sleep spindle detection and genetic influence: Estimation using continuous wavelet transform. *Frontiers in Human Neuroscience*, 9(624).

Amzica, F. and Steriade, M. (1998). Electrophysiological correlates of sleep delta waves. *Electroencephalography and Clinical Neurophysiology*, 107:69–83.

Aru, J., Aru, J., Priesemann, V., Wibral, M., Lana, L., Pipa, G., Singer, W., and Vicente, R. (2015). Untangling cross-frequency coupling in neuroscience. *Current Opinion in Neurobiology*, 31:51–61.

Aserinsky, E. and Kleitman, N. (1953). Regularly occurring periods of eye motility, and concomitant phenomena, during sleep. *Science*, 118(3062):4–5.

Ashburner, J. and Friston, K. J. (2000). Voxel-based morphometry: The methods. *NeuroImage*, 11(6):805–821.

Aston-Jones, G., Gonzalez, M., and Doran, S. (2007). Role of the locus coeruleus-norepinephrine system in arousal and circadian regulation of the sleep–wake cycle. In Ordway, G. A., Schwartz, M. A., and Frazer, A., editors, *Brain Norepinephrine: Neurobiology and Therapeutics*, pages 157–195. University Press, Cambridge.

Ayoub, A., Aumann, D., Hörschelmann, A., Kouchekmanesch, A., Paul, P., Born, J., and Marshall, L. (2013). Differential effects on fast and slow spindle activity, and the sleep slow oscillation in humans with carbamazepine and flunarizine to antagonize voltage-dependent Na+ and Ca2+ channel activity. *Sleep*, 36(6):905–911.

Bagur, S., Lacroix, M. M., de Lavilléon, G., Lefort, J. M., Geoffroy, H., and Benchenane, K. (2018). Harnessing olfactory bulb oscillations to perform fully brain-based sleep-scoring and real-time monitoring of anaesthesia depth. *PLoS Biology*, 16(11):e2005458.

Baltes, P. B. (1987). Theoretical propositions of life-span developmental psychology: On the dynamics between growth and decline. *Developmental Psychology*, 23(5):611–626.

Baltes, P. B., Lindenberger, U., and Staudinger, U. M. (1998). Life-span theory in developmental psychology. In Damon, W. and Lerner, R. M., editors, *Handbook of child psychology: Vol. 1. Theoretical models of human development*, pages 1029–1143. Wiley, New York, NY.

Barakat, M., Doyon, J., Debas, K., Vandewalle, G., Morin, A., Poirier, G., Martin, N., Lafortune, M., Karni, A., Ungerleider, L. G., Benali, H., and Carrier, J. (2011). Fast and slow spindle involvement in the consolidation of a new motor sequence. *Behavioural Brain Research*, 217:117–121.

Baran, B., Mantua, J., and Spencer, R. M. C. (2016). Age-related changes in the sleep-dependent reorganization of declarative memories. *Journal of Cognitive Neuroscience*, 28(6):792–802.

Bersagliere, A. and Achermann, P. (2010). Slow oscillations in human non-rapid eye movement sleep electroencephalogram: Effects of increased sleep pressure. *Journal of Sleep Research*, 19(1):228–237.

Blake, H. and Gerard, R. (1937). Brain potentials during sleep. *American Journal of Physiology*, 119:692–703.

Bódizs, R., Körmendi, J., Rigó, P., and Lázár, A. S. (2009). The individual adjustment method of sleep spindle analysis: Methodological improvements and roots in the fingerprint paradigm. *Journal of Neuroscience Methods*, 178(1):205–213.

Browning, G. B. and Spilich, G. J. (1981). Some important methodological issues in the study of aging and cognition. *Experimental Aging Research*, 7(2):175–187.

Buchmann, A., Ringli, M., Kurth, S., Schaerer, M., Geiger, A., Jenni, O. G., and Huber, R. (2011). EEG sleep slow-wave activity as a mirror of cortical maturation. *Cerebral Cortex*, 21(3):607–615.

Buckner, R. L. (2004). Memory and executive function in aging and AD: Multiple factors that cause decline and reserve factors that compensate. *Neuron*, 44:195–208.

Burzynska, A. Z., Garrett, D. D., Preuschhof, C., Nagel, I. E., and Li, S.-C. (2013). A scaffold for efficiency in the human brain. *Journal of Neuroscience*, 33(43):17150–17159.

Buysse, D. J., Reynolds III, C. F., Monk, T. H., Berman, S. R., and Kupfer, D. J. (1989). The Pittsburg Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Research*, 28(2):193–213.

Buzsáki, G. (2006). *Rhythms of the brain*. Oxford University Press, Oxford.

Buzsáki, G., Anastassiou, C. A., and Koch, C. (2016). The origin of extracellular fields and currents — EEG, ECoG, LFP and spikes. *Nature Reviews Neuroscience*, 13(6):407–420.

Cabeza, R., Albert, M., Belleville, S., M Craik, F. I., Duarte, A., Grady, C. L., Lindenberger, U., Nyberg, L., Park, D. C., Reuter-Lorenz, P. A., Rugg, M. D., Steffener, J., and Rajah, N. M. (2018). Maintenance, reserve and compensation: The cognitive neuroscience of healthy ageing. *Nature Reviews Neuroscience*, 19:701–710.

Cabeza, R., Nyberg, L., and Park, D., editors (2005). *Cognitive neuroscience of aging: Linking cognitive and cerebral aging*. Oxford University Press, New York.

Campbell, I. G. and Feinberg, I. (2016). Maturational patterns of sigma frequency power across childhood and adolescence: A longitudinal study. *Sleep*, 39(1):193–201.

Carrier, J., Viens, I., Poirier, G., Robillard, R., Lafortune, M., Vandewalle, G., Martin, N., Barakat, M., Paquet, J., and Filipini, D. (2011). Sleep slow wave changes during the middle years of life. *European Journal of Neuroscience*, 33:758–766.

Carskadon, M. A. and Dement, W. C. (2011). Monitoring and staging human sleep. In Kryger, M. H., Roth, T., and Dement, W. C., editors, *Principles and Practice of sSeep Medicine*, pages 16–26. Elsevier Saunders, St. Louis, 5th edition.

Cirelli, C., Huber, R., Gopalakrishnan, A., Southard, T. L., and Tononi, G. (2005). Locus ceruleus control of slow-wave homeostasis. *Journal of Neuroscience*, 25(18):4503–4511.

Clawson, B. C., Durkin, J., and Aton, S. J. (2016). Form and function of sleep spindles across the lifespan. *Neural Plasticity*, 2016:6936381.

Cohen, M. X. (2014). *Analyzing neural time series data: Theory and practice*. The MIT Press, London, England.

Cole, J. H., Poudel, R. P., Tsagkrasoulis, D., Caan, M. W., Steves, C., Spector, T. D., and Montana, G. (2017). Predicting brain age with deep learning from raw imaging data results in a reliable and heritable biomarker. *NeuroImage*, 163:115–124.

Cole, J. H., Ritchie, S. J., Bastin, M. E., Valdés Hernández, M. C., Muñoz Maniega, S., Royle, N., Corley, J., Pattie, A., Harris, S. E., Zhang, Q., Wray, N. R., Redmond, P., Marioni, R. E., Starr, J. M., Cox,

S. R., Wardlaw, J. M., Sharp, D. J., and Deary, I. J. (2018). Brain age predicts mortality. *Molecular Psychiatry*, 23(5):1385–1392.

Coppieters 't Wallant, D., Maquet, P., and Phillips, C. (2016). Sleep spindles as an electrographic element: Description and automatic detection methods. *Neural Plasticity*, 2016:6783812.

Cox, R., Schapiro, A. C., Manoach, D. S., and Stickgold, R. (2017). Individual differences in frequency and topography of slow and fast sleep spindles. *Frontiers in Human Neuroscience*, 11:433.

Crowley, K., Trinder, J., Kim, Y., Carrington, M., and Colrain, I. M. (2002). The effects of normal aging on sleep spindle and K-complex production. *Clinical Neurophysiology*, 113:1615–1622.

Danker-Hopfe, H., Scha, M., Dorn, H., Anderer, P., Saletu, B., Zeitlhofer, J., Kunz, D., Barbanoj, M.-J., Himanen, S. L., Penzel, T., and Ro, J. (2005). Percentile reference charts for selected sleep parameters for 20- to 80-year-old healthy subjects from the SIESTA database. *Somnologie*, 9:3–14.

Dannhauer, M., Lanfer, B., Wolters, C. H., and Knösche, T. R. (2011). Modeling of the human skull in EEG source analysis. *Human Brain Mapping*, 32(9):1383–1399.

De Gennaro, L. and Ferrara, M. (2003). Sleep spindles: An overview. *Sleep Medicine Reviews*, 7(5):423–440.

Dement, W. C. and Wolpert, E. A. (1958). The relation of eye movements, body motility, and external stimuli to dream content. *Journal of Experimental Psychology*, 55(6):543–553.

Diekelmann, S. and Born, J. (2010). The memory function of sleep. *Neuroscience*, 11(2):114–126.

Dijk, D.-J., Duffy, J. F., and Czeisler, C. A. (2000). Contribution of circadian physiology and sleep homeostasis to age-related changes in human sleep. *Chronobiology International*, 17(3):285–311.

Doppelmayr, M., Klimesch, W., Pachinger, T., and Ripper, B. (1998). Individual differences in brain dynamics: Important implications for the calculation of event-related band power. *Biological Cybernetics*, 57:49–57.

Doran, S. M. (2003). The dynamic topography of individual sleep spindles. *Sleep Research Online*, 5(4):133–139.

Dubé, J., Lafortune, M., Bedetti, C., Bouchard, M., Franc, J., Doyon, J., Evans, A. C., Lina, J.-m., and Carrier, X. J. (2015). Cortical thinning explains changes in sleep slow waves during adulthood. *Journal of Neuroscience*, 35(20):7795–7807.

Dubois, J., Dehaene-Lambertz, G., Soares, C., Cointepas, Y., Le Bihan, D., and Hertz-Pannier, L. (2008). Microstructural correlates of infant functional development: Example of the visual pathways. *Journal of Neuroscience*, 28(8):1943–1948.

Esser, S. K., Hill, S. L., and Tononi, G. (2007). Sleep homeostasis and cortical synchronization: I. Modeling the effects of synaptic strength on sleep slow waves. *Sleep*, 30(12):1617–1630.

Fandakova, Y., Sander, M. C., Grandy, T. H., Cabeza, R., Werkle-Bergner, M., and Shing, Y. L. (2018). Age differences in false memory: The importance of retrieval monitoring processes and their modulation by memory quality. *Psychology and Aging*, 33(1):119–135.

Feinberg, I. (1974). Changes in sleep cycle patterns with age. *Journal of Psychiatric Research*, 10:283–306.

Feinberg, I. and Evarts, E. V. (1969). Changing concepts of the function of sleep: Discovery of intense brain activity during sleep calls for revision of hypotheses as to its function. *Biological Psychiatry*, 1(4):331–348.

Ferrarelli, F., Kaskie, R., Laxminarayan, S., Ramakrishnan, S., Reifman, J., and Germain, A. (2019). An increase in sleep slow waves predicts better working memory performance in healthy individuals. *NeuroImage*, 191(January):1–9.

Fjell, A. M. and Walhovd, K. B. (2010). Structural brain changes in aging: Courses, causes and cognitive consequences. *Reviews in the Neurosciences*, 21(3):187–221.

Fjell, A. M., Westlye, L. T., Grydeland, H., Amlien, I., Reinvang, I., Raz, N., Holland, D., Dale, A. M., and Walhovd, K. B. (2013). Critical ages in the life-course of the adult brain: Nonlinear subcortical aging. *Neurobiology of Aging*, 34(10):2239–2247.

Fogel, S. M., Martin, N., Lafortune, M., Barakat, M., Debas, K., Laventure, S., Latreille, V., Gagnon, J.-F., Doyon, J., and Carrier, J. (2012). NREM sleep oscillations and brain plasticity in aging. *Frontiers in Neurology*, 3:176.

Fogel, S. M. and Smith, C. T. (2011). The function of the sleep spindle: A physiological index of intelligence and a mechanism for sleep-dependent memory consolidation. *Neuroscience and Biobehavioral Reviews*, 35(5):1154–1165.

Fogel, S. M., Vien, C., Karni, A., Benali, H., Carrier, J., and Doyon, J. (2017). Sleep spindles: A physiological marker of age-related changes in gray matter in brain regions supporting motor skill memory consolidation. *Neurobiology of Aging*, 49:154–164.

Folstein, M. F., Folstein, S. E., and McHugh, P. R. (1975). "Mini-Mental State": A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12:189–198.

Franke, K., Ziegler, G., Klöppel, S., and Gaser, C. (2010). Estimating the age of healthy subjects from T1-weighted MRI scans using kernel methods: Exploring the influence of various parameters. *NeuroImage*, 50(3):883–892.

Frodl, T., Meisenzahl, E. M., Müller, D., Leinsinger, G., Juckel, G., Hahn, K., Möller, H. J., and Hegerl, U. (2001). The effect of the skull on event-related P300. *Clinical Neurophysiology*, 112(9):1773–1776.

Gais, S., Mölle, M., Helms, K., and Born, J. (2002). Learning-dependent increases in sleep spindle density. *Journal of Neuroscience*, 22(15):6830–6834.

Genzel, L., Kroes, M. C. W., Dresler, M., and Battaglia, F. P. (2014). Light sleep versus slow wave sleep in memory consolidation: A question of global versus local processes? *Trends in Neurosciences*, 37(1):10–19.

Giorgio, A., Santelli, L., Tomassini, V., Bosnell, R., Smith, S., De Stefano, N., and Johansen-Berg, H. (2010). Age-related changes in grey and white matter structure throughout adulthood. *NeuroImage*, 51:943–951.

Grandy, T. H., Werkle-Bergner, M., Chicherio, C., Schmiedek, F., Lövdén, M., and Lindenberger, U. (2013). Peak individual alpha frequency qualifies as a stable neurophysiological trait marker in healthy younger and older adults. *Psychophysiology*, 50(6):570–582.

Habib, R. and Nyberg, L. (2007). Neural correlates of availability and accessibility in memory. *Cerebral Cortex*, 18(7):1720–1726.

Hedden, T. and Gabrieli, J. D. E. (2004). Insights into the ageing mind: A view from cognitive neuroscience. *Nature Reviews Neuroscience*, 5:87–96.

Heib, D. P. J., Hoedlmoser, K., Anderer, P., Zeitlhofer, J., Gruber, G., Klimesch, W., and Schabus, M. (2013). Slow oscillation amplitudes and up-state lengths relate to memory improvement. *PLoS ONE*, 8(12):e82049.

Helfrich, R. F., Mander, B. A., Jagust, W. J., Knight, R. T., and Walker, M. P. (2018). Old brains come uncoupled in sleep: Slow wave-spindle synchrony, brain atrophy, and forgetting. *Neuron*, 97:221–230.e7.

Hertzog, C. and Nesselroade, J. R. (2003). Assessing psychological change in adulthood: An overview of methodological issues. *Psychology and Aging*, 18(4):639–657.

Hobson, J. A. (1968). Book review: A manual of standardized terminology, techniques and scoring system for sleep stages of human subject. *Electroencephalography and Clinical Neurophysiology*, 26(6):644.

Hobson, J. A. (2005). Sleep is of the brain, by the brain and for the brain. *Nature*, 437:1254–1256.

Hofer, S. M. and Sliwinski, M. J. (2001). Understanding ageing. *Gerontology*, 47:341–352.

Iber, C., Ancoli-Israel, S., Chesson, A. L., and Quan, S. F. (2007). *The AASM manual for the scoring of sleep and associted events: Rules, terminology and technical specifications*. American Academy of Sleep Medicine, Westchester, IL, 1st edition.

Jasper, H. H. (1958). The ten-twenty electrode system of the International Federation. *Electroencephalography and Clinical Neurophysiology*, 10:371–375.

Ju, Y.-E. S., Lucey, B. P., and Holtzman, D. M. (2014). Sleep and Alzheimer disease pathology—a bidirectional relationship. *Nature Reviews Neuroscience*, 10(2):115–119.

Karch, J. D., Sander, M. C., von Oertzen, T., Brandmaier, A. M., and Werkle-Bergner, M. (2015). Using within-subject pattern classification to understand lifespan age differences in oscillatory mechanisms of working memory selection and maintenance. *NeuroImage*, 118:538–552.

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis. *Brain Research Reviews*, 29(2-3):169–195.

Klinzing, J. G., Kugler, S., Soekadar, S. R., Rasch, B., Born, J., and Diekelmann, S. (2018). Odor cueing during slow-wave sleep benefits memory independently of low cholinergic tone. *Psychopharmacology*, 235:291–299.

Klinzing, J. G., Mölle, M., Weber, F., Supp, G., Hipp, J. F., Engel, A. K., and Born, J. (2016). Spindle activity phase-locked to sleep slow oscillations. *NeuroImage*, 134:607–616.

Koo, T. K. and Li, M. Y. (2016). A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *Journal of Chiropractic Medicine*, 15(2):155–163.

Kosciessa, J. Q., Grandy, T. H., Garrett, D. D., and Werkle-Bergner, M. (2018). Single-trial characterization of neural rhythms: Potential and challenges. *bioRxiv*, page 356089.

Kurth, S., Jenni, O. G., Riedner, B. A., Tononi, G., Carskadon, M. A., and Huber, R. (2010). Characteristics of sleep slow waves in children and adolescents. *Sleep*, 33(4):475–80.

Lacroix, M. M., Lefort, J., Kanbi, K. E., Laventure, S., Dauvilliers, Y., Peyron, C., Benchenane, K., and Unit, B. P. (2018). Improved sleep scoring in mice reveals human-like stages. *bioRxiv*, page 489005.

Landolt, H.-P. and Borbély, A. A. (2001). Age-dependent changes in sleep EEG topography. *Clinical Neurophysiology*, 112:369–377.

Landolt, H.-P., Dijk, D. J., Achermann, P., and Borbély, A. A. (1996). Effect of age on the sleep EEG: Slow-wave activity and spindle frequency activity in young and middle-aged men. *Brain Research*, 738(2):205–212.

Leissner, P., Lindholm, L.-E., and Petersen, I. (1970). Alpha amplitude dependence on skull thickness as measured by ultrasound technique. *Electroencephalography and Clinical Neurophysiology*, 29:392–399.

Li, S.-C., Lindenberger, U., Hommel, B., Aschersleben, G., Prinz, W., and Baltes, P. B. (2004). Transformations in the couplings among intellectual abilities and constituent cognitive processes across the life span. *Psychological Science*, 15:155–163.

Li, S.-C. and Schmiedek, F. (2002). Age is not necessarily aging: Another step towards understanding the clocks that time aging. *Gerontology*, 48:5–12.

Lim, A. S. P., Kowgier, M., Yu, L., Buchman, A. S., and Bennett, D. A. (2013). Sleep fragmentation and the risk of incident Alzheimer's disease and cognitive decline in older persons. *Sleep*, 36(7):1027–1032.

Lindenberger, U. (2014). Human cognitive aging: Corriger la fortune? *Science*, 346(6209):572–578.

Lindenberger, U., Li, S. C., and Bäckman, L. (2006). Delineating brain-behavior mappings across the lifespan: Substantive and methodological advances in developmental neuroscience. *Neuroscience and Biobehavioral Reviews*, 30(6):713–717.

Lindenberger, U. and Pötter, U. (1998). The complex nature of unique and shared effects in hierarchical linear regression: Implications for developmental psychology. *Psychological Methods*, 3(2):218–230.

Lindenberger, U., von Oertzen, T., Ghisletta, P., and Hertzog, C. (2011). Cross-sectional age variance extraction: What's change got to do with it? *Psychology and Aging*, 26(1):34–47.

Loomis, A. L., Harvey, E. N., and Hobart, G. (1935). Further observations on the potential rhythms of the cerebral cortex during sleep. *Science*, 82(2122):198–200.

Loomis, A. L., Harvey, E. N., and Hobart, G. A. I. (1962). Cerebral states during sleep. as studied by human brain potentials. *Journal of Experimental Psychology*, 64(1):127–144.

Lövden, M., Bäckmann, L., Lindenberger, U., Schaefer, S., and Schmiedek, F. (2010). A theoretical framework for the study of adult cognitive plasticity. *Psychological Bulletin*, 136(4):659–676.

Lucey, B. P., McCullough, A., Landsness, E. C., Toedebusch, C. D., McLeland, J. S., Zaza, A. M., Fagan, A. M., McCue, L., Xiong, C., Morris, J. C., Benzinger, T. L. S., and Holtzman, D. M. (2019). Reduced non–rapid eye movement sleep is associated with tau pathology in early Alzheimer's disease. *Science Translational Medicine*, 11(474):eaau6550.

MacDonald, S. W. S., DeCarlo, C. A., and Dixon, R. A. (2011). Linking biological and cognitive aging: Toward improving characterizations of developmental time. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 66B:i59–i70.

Mander, B. A. (2017). A restless night makes for a rising tide of amyloid. *Brain*, 140:2066–2069.

Mander, B. A., Marks, S. M., Vogel, J. W., Rao, V., Lu, B., Saletin, J. M., Ancoli-Israel, S., Jagust, W. J., and Walker, M. P. (2015). Beta-amyloid disrupts human NREM slow waves and related hippocampus-dependent memory consolidation. *Nature Neuroscience*, 18(7):1051–1057.

Mander, B. A., Rao, V., Lu, B., Saletin, J. M., Lindquist, J. R., Ancoli-Israel, S., Jagust, W., and Walker, M. P. (2013). Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. *Nature Neuroscience*, 16(3):357–364.

Mander, B. A., Winer, J., Jagust, W. J., and Walker, M. P. (2016). Sleep: A novel mechanistic pathway, biomarker, and treatment target in the pathology of Alzheimer's Disease? *Trends in Neurosciences*, 39(8):552–566.

Mander, B. A., Zhu, A. H., Lindquist, J. R., Villeneuve, S., Rao, V., Lu, B., Saletin, J. M., Ancoli-Israel, S., Jagust, W., and Walker, M. P. (2017). White matter structure in older adults moderates the benefit of sleep spindles on motor memory consolidation. *The Journal of Neuroscience*, pages 3033–16.

Martin, N., Lafortune, M., Godbout, J., Barakat, M., Robillard, R., Poirier, G., Bastien, C., and Carrier, J. (2013). Topography of age-related changes in sleep spindles. *Neurobiology of Aging*, 34:468–476.

Massimini, M., Huber, R., Ferrarelli, F., Hill, S. L., and Tononi, G. (2004). The sleep slow oscillation as a traveling wave. *Journal of Neuroscience*, 24(31):6862–6870.

Mensen, A., Zhang, Z., Qi, M., and Khatami, R. (2016). The occurrence of individual slow waves in sleep is predicted by heart rate. *Scientific Reports*, 6(July):29671.

Molenaar, P. C. M. and Campbell, C. G. (2009). The new person-specifc paradigm in psychology. *Psychological Science*, 18(2):112–117.

Mölle, M., Bergmann, T. O., Marshall, L., and Born, J. (2011). Fast and slow spindles during the sleep slow oscillation: Disparate coalescence and engagement in memory processing. *Sleep*, 34(10):1411–1421.

Mölle, M., Eschenko, O., Gais, S., Sara, S. J., and Born, J. (2009). The influence of learning on sleep slow oscillations and associated spindles and ripples in humans and rats. *European Journal of Neuroscience*, 29:1071–1081.

Mölle, M., Marshall, L., Gais, S., and Born, J. (2002). Grouping of spindle activity during slow oscillations in human non-rapid eye movement sleep. *Journal of Neuroscience*, 22(24):10941–10947.

Muehlroth, B. E., Sander, M. C., Fandakova, Y., Grandy, T. H., Rasch, B., Shing, Y. L., and Werkle-Bergner, M. (2019a). Memory quality modulates the effect of aging on memory consolidation during sleep: Reduced maintenance but intact gain. *bioRxiv*, 547448.

Muehlroth, B. E., Sander, M. C., Fandakova, Y., Grandy, T. H., Rasch, B., Shing, Y. L., and Werkle-Bergner, M. (2019b). Precise slow oscillation–spindle coupling promotes memory consolidation in younger and older adults. *Scientific Reports*, 9(1):1940.

Nesselroade, J. R., Gerstorf, D., Hardy, S. A., and Ram, N. (2007). Focus article: Idiographic filters for psychological constructs. *Measurement: Interdisciplinary Research & Perspective*, 5(4):217–235.

Nesselroade, J. R. and Molenaar, P. C. (2016). Some behavioral science measurement concerns and proposals. *Multivariate Behavioral Research*, 51(2-3):396–412.

Ngo, H.-V. V., Martinetz, T., Born, J., and Mölle, M. (2013). Auditory closed-loop stimulation of the sleep slow oscillation enhances memory. *Neuron*, 78:545–553.

Ngo, H.-V. V., Miedema, A., Faude, I., Martinetz, T., Mölle, M., and Born, J. (2015). Driving sleep slow oscillations by auditory closed-loop stimulation: A self-limiting process. *Journal of Neuroscience*, 35(17):6630–6638.

Nicolas, A., Petit, D., Rompré, S., and Montplaisir, J. (2001). Sleep spindle characteristics in healthy subjects of different age groups. *Clinical Neurophysiology*, 112(3):521–527.

Nir, Y., Staba, R. J., Andrillon, T., Vyazovskiy, V. V., Cirelli, C., Fried, I., and Tononi, G. (2011). Regional slow waves and spindles in human sleep. *Neuron*, 70(1):153–169.

Noble, W. and Spires-Jones, T. L. (2019). Sleep well to slow Alzheimer's progression? *Science*, 363(6429):813–814.

Nyberg, L., Lövdén, M., Riklund, K., Lindenberger, U., and Bäckman, L. (2012). Memory aging and brain maintenance. *Trends in Cognitive Sciences*, 16(5):292–305.

Nyberg, L. and Pudas, S. (2019). Successful memory aging. *Annual Review of Psychology*, 70:219–243.

Ohayon, M. M., Carskadon, M. A., Guilleminault, C., and Vitiello, M. V. (2004). Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: Developing normative sleep values across the human lifespan. *Sleep*, 27(7):1255–1273.

Oostenveld, R., Fries, P., Maris, E., and Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, page 156869.

Oostenveld, R. and Praamstra, P. (2001). The five percent electrode system for high-resolution EEG and ERP measurement. *Clinical Neurophysiology*, 112:713–719.

Overton, W. (2010). Life-span development – Concepts and issues. In Lerner, R. M., Lamb, M. E., and Freund, A. M., editors, *The Handbook of Life-Span Development: Vol. 1.*, chapter 1. John Wiley & Sons.

Park, D. C. and Reuter-Lorenz, P. (2009). The adaptive brain: Aging and neurocognitive scaffolding. *Annual Review of Psychology*, 60:173–96.

Piantoni, G., Astill, R. G., Raymann, R. J. E. M., Vis, J. C., Coppens, J. E., and Van Someren, E. J. W. (2013). Modulation of gamma and spindle-range power by slow oscillations in scalp sleep EEG of children. *International Journal of Psychophysiology*, 89:252–258.

Posthuma, D., Neale, M. C., Boomsma, D. I., and Geus, E. J. C. D. (2001). Are smarter brains running faster? Heritability of alpha peak frequency, IQ, and their interrelation. *Behavior Genetics*, 31(6):567–579.

Prinz, P. N. (1977). Sleep patterns in the healthy aged: Relationship with intellectual function. *Journal of Gerontology*, 32(2):179–186.

Purcell, S. M., Manoach, D. S., Demanuele, C., Cade, B. E., Mariani, S., Cox, R., Panagiotaropoulou, G., Saxena, R., Pan, J. Q., Smoller, J. W., Redline, S., and Stickgold, R. (2017). Characterizing sleep spindles in 11,630 individuals from the National Sleep Research Resource. *Nature Communications*, 8:15930.

Ramm, P. (1979). The locus coeruleus, catecholamines, and REM sleep: A critical review. *Behavioral and Neural Biology*, 25(4):415–448.

Rasch, B. and Born, J. (2013). About sleep's role in memory. *Physiological Reviews*, 93:681–766.

Raz, N. and Lindenberger, U. (2011). Only time will tell: Cross-sectional studies offer no solution to the age-brain-cognition triangle: Comment on Salthouse (2011). *Psychological Bulletin*, 137(5):790–795.

Raz, N., Lindenberger, U., Rodrigue, K. M., Kennedy, K. M., Head, D., Williamson, A., Dahle, C., Gerstorf, D., and Acker, J. D. (2005). Regional brain changes in aging healthy adults: General trends, individual differences and modifiers. *Cerebral Cortex*, 15:1676–1689.

Raz, N. and Rodrigue, K. M. (2006). Differential aging of the brain: Patterns, cognitive correlates and modifiers. *Neuroscience and Biobehavioral Reviews*, 30:730–748.

Rechtschaffen, A. and Kales, A. (1968). *A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects*. Public Health Service, US Government Printing Office, Washington DC.

Redline, S., Kirchner, H. L., Quan, S. F., Gottlieb, D. J., Kapur, V., and Newmann, A. (2004). The effects of age, sex, ethnicity, and sleep-disordered breathing on sleep architecture. *Archives of Internal Medicine*, 164:406–418.

Rönnlund, M., Nyberg, L., Bäckman, L., and Nilsson, L.-G. (2005). Stability, growth, and decline in adult life span development of declarative memory: Cross-sectional and longitudinal data from a population-based study. *Psychology and Aging*, 20(1):3–18.

Rose, L. T., Rouhani, P., and Fischer, K. W. (2013). The science of the individual. *Mind, Brain, and Education*, 7(3):152–158.

Saletin, J. M., van der Helm, E., and Walker, M. P. (2013). Structural brain correlates of human sleep oscillations. *NeuroImage*, 83:658–668.

Sander, M., Fandakova, Y., Grandy, T., Shing, Y., and Werkle-Bergner, M. (2019). Oscillatory mechanisms of successful memory formation in younger and older adults are related to structural integrity. *bioRxiv*, 530121.

Schabus, M., Dang-Vu, T. T., Albouy, G., Balteau, E., Boly, M., Carrier, J., Darsaud, A., Degueldre, C., Desseilles, M., Gais, S., Phillips, C., Rauchs, G., Schnakers, C., Sterpenich, V., Vandewalle, G., Luxen, A., and Maquet, P. (2007). Hemodynamic cerebral correlates of sleep spindles during human non-rapid eye movement sleep. *Proceedings of the National Academy of Sciences of the United States of America*, 104(32):13164–13169.

Scullin, M. K. (2013). Sleep, memory, and aging: The link between slow-wave sleep and episodic memory changes from younger to older adults. *Psychology and Aging*, 28(1):105–114.

Segalowitz, S. J. and Davies, P. L. (2004). Charting the maturation of the frontal lobe: An electrophysiological strategy. *Brain and Cognition*, 55:116–133.

Shokri-Kojori, E., Wang, G.-J., Wiers, C. E., Demiral, S. B., Guo, M., Kim, S. W., Lindgren, E., Ramirez, V., Zehra, A., Freeman, C., Miller, G., Manza, P., Srivastava, T., De Santi, S., Tomasi, D., Benveniste, H., and Volkow, N. D. (2018). β-Amyloid accumulation in the human brain after one night of sleep deprivation. *Proceedings of the National Academy of Sciences*, 115(17):4483–4488.

Silber, M. H., Ancoli-Israel, S., Bonnet, M. H., Chokroverty, S., Grigg-Damberger, M. M., Hirshkowitz, M., Kapen, S., Keenan, S. A., Kryger, M. H., Penzel, T., Pressman, M. R., and Iber, C. (2007). The visual scoring of sleep in adults. *Journal of Clinical Sleep Medicine*, 3(2):121–131.

Sommer, V., Fandakova, Y., Grandy, T., Shing, Y., Werkle-Bergner, M., and Sander, M. (2019). Neural pattern similarity differentially affects memory performance of younger and older adults. *bioRxiv*, 528620.

Sonni, A. and Spencer, R. M. C. (2015). Sleep protects memories from interference in older adults. *Neurobiology of Aging*, 36(7):2272–2281.

Sprecher, K. E., Riedner, B. A., Smith, R. F., Tononi, G., Davidson, R. J., and Benca, R. M. (2016). High resolution topography of age-related changes in non-rapid eye movement sleep electroencephalography. *PLoS ONE*, 11(2):e0149770.

Staresina, B. P., Bergmann, T. O., Bonnefond, M., Van Der Meij, R., Jensen, O., Deuker, L., Elger, C. E., Axmacher, N., and Fell, J. (2015). Supplements: Hierarchical nesting of slow oscillations, spindles and ripples in the human hippocampus during sleep. *Nature Neuroscience*, 18(11):1679–1686.

Steffener, J., Habeck, C., O'Shea, D., Razlighi, Q., Bherer, L., and Stern, Y. (2016). Differences between chronological and brain age are related to education and self-reported physical activity. *Neurobiology of Aging*, 25(3):289–313.

Steriade, M., Datta, S., Paré, D., Oakson, G., and Curró Dossi, R. C. (1990). Neuronal activities in brain-stem cholinergic nuclei related to tonic activation processes in thalamocortical systems. *The Journal of Neuroscience*, 10(8):2541–2559.

Steriade, M., Nunez, A., and Amzica, F. (1993). A novel slow (<1 Hz) oscillation of neocortical neurons in vivo: Depolarizing and hyperpolarizing components. *The Journal of Neuroscience*, 13(8):3252–3265.

Sun, H., Paixao, L., Oliva, J. T., Goparaju, B., Carvalho, D. Z., van Leeuwen, K. G., Akeju, O., Thomas, R. J., Cash, S. S., Bianchi, M. T., and Westover, M. B. (2019). Brain age from the electroencephalogram of sleep. *Neurobiology of Aging*, 74:112–120.

Timofeev, I. and Chauvette, S. (2013). The spindles: Are they still thalamic? *Sleep*, 36(6):825–826.

Ujma, P. P., Gombos, F., Genzel, L., Konrad, B. N., Simor, P., Steiger, A., Dresler, M., and Bódizs, R. (2015). A comparison of two sleep spindle detection methods based on all night averages: individually adjusted vs. fixed frequencies. *Frontiers in Human Neuroscience*, 9(February):52.

Ujma, P. P., Simor, P., Steiger, A., Dresler, M., and Bódizs, R. (2019). Individual slow wave morphology is a marker of ageing. *Neurobiology of Aging*, 80:71–82.

Vaou, O. E., Lin, S. H., Branson, C., and Auerbach, S. (2018). Sleep and dementia. *Current Sleep Medicine Reports*, pages 134–142.

Varga, A. W., Ducca, E. L., Kishi, A., Fischer, E., Parekh, A., Koushyk, V., Yau, P. L., Gumb, T., Leibert, D. P., Wohlleber, M. E., Burschtin, O. E., Convit, A., Rapoport, D. M., Osorio, R. S., and Ayappa, I. (2016). Effects of aging on slow wave sleep dynamics and human spatial navigational memory consolidation. *Neurobiology of Aging*, 42:142–149.

Vassalli, A. and Dijk, D.-J. (2009). Sleep function: Current questions and new approaches. *The European journal of neuroscience*, 29:1830–1841.

Vitiello, M. V. (2006). Sleep in normal aging. *Sleep Medicine Clinics*, 1:171–176.

Vyazovskiy, V. V., Olcese, U., Lazimy, Y. M., Faraguna, U., Esser, S. K., Williams, J. C., Cirelli, C., and Tononi, G. (2009). Cortical firing and sleep homeostasis. *Neuron*, 63(6):865–878.

Warby, S. C., Wendt, S. L., Welinder, P., Munk, E. G. S., Carrillo, O., Sorensen, H. B. D., Jennum, P., Peppard, P. E., Perona, P., and Mignot, E. (2014). Sleep-spindle detection: Crowdsourcing and evaluating performance of experts, non-experts and automated methods. *Nature Methods*, 11(4):385–392.

Webb, W. B. (1982). The measurement and characteristics of sleep in older persons. *Neurobiology of Aging*, 3:311–319.

Webb, W. B. and Dreblow, L. M. (1982). A modified method for scoring slow wave sleep of older subjects. *Sleep*, 5:195–199.

Werkle-Bergner, M., Shing, Y. L., Müller, V., Li, S.-C., and Lindenberger, U. (2009). Clinical Neurophysiology EEG gamma-band synchronization in visual coding from childhood to old age: Evidence from evoked power and inter-trial phase locking. *Clinical Neurophysiology*, 120:1291–1302.

Werth, E., Achermann, P., Dijk, D.-j., and Borbe, A. A. (1997). Spindle frequency activity in the sleep EEG: Individual differences and topographic distribution. *Electroencephalography and Clinical Neurophysiology*, 103:535–542.

Westerberg, C. E., Mander, B. A., Florczak, S. M., Weintraub, S., Mesulam, M., Zee, P. C., and Paller, K. A. (2012). Concurrent impairments in sleep and memory in amnestic mild cognitive impairment. *Journal of the International Neuropsychological Society*, 18:490–500.

Whitford, T. J., Rennie, C. J., Grieve, S. M., Clark, C. R., Gordon, E., and Williams, L. M. (2007). Brain maturation in adolescence: Concurrent changes in neuroanatomy and neurophysiology. *Human Brain Mapping*, 28(3):228–237.

Whitten, T. A., Hughes, A. M., Dickson, C. T., and Caplan, J. B. (2011). A better oscillation detection method robustly extracts EEG rhythms across brain state changes: The human alpha rhythm as a test case. *NeuroImage*, 54(2):860–874.

Wohlwill, J. F. (1970). The age variable in psychological research. *Psychological Review*, 1:49–64.

Zhong, H.-H., Yu, B., Luo, D., Yang, L.-Y., Zhang, J., Jiang, S.-S., Hu, S.-J., Luo, Y.-Y., Yang, M.-w., Hong, F.-f., and Yang, S.-L. (2019). Roles of aging in sleep. *Neuroscience & Biobehavioral Reviews*, 98:177–187.

Ziegler, G., Dahnke, R., Jäncke, L., Yotter, R. A., May, A., and Gaser, C. (2012). Brain structural trajectories over the adult lifespan. *Human Brain Mapping*, 33(10):2377–2389.

Supplementary Methods

Overview of data

In the present article, we report on data from 29 healthy younger (range: 19–28 years, M_{age} = 23.55 years, SD_{age} = 2.58; 16 females) and 34 healthy older adults (range: 63–74 years, M_{age} = 68.85 years, SD_{age} = 3.11; 15 females) whose sleep was monitored in two consecutive nights at their homes using ambulatory polysomnography (PSG). This resulted in a dataset consisting of 54 complete nights of sleep recordings from younger adults (M_{age} = 23.52 years, SD_{age} = 2.53; 29 female) and 63 from older adults (M_{age} = 68.77 years, SD_{age} = 3.07; 28 female). Analyses involving structural magnet resonance imaging (MRI) data, were based on data from 29 younger adults (M_{age} = 23.66 years, SD_{age} = 2.6; 17 female) and 32 of the older adults (M_{age} = 68.78 years, SD_{age} = 2.94; 14 female). All participants were right-handed native German speakers with no reported history of psychiatric or neurological disease, or any use of psychoactive medication. Older participants were screened for cognitive deficits using the Mini-Mental State Examination (MMSE; M = 29.24, SD = 1.12, range: 26–30; Folstein et al., 1975). General subjective sleep quality was assessed using the Pittsburgh Sleep Quality Index (PSQI; Buysse et al., 1989).

All displayed data were collected as part of a research project investigating age differences in the encoding, consolidation, and retrieval of associative memories (for more details on the experimental procedure, see Muehlroth et al., 2019a,b; Fandakova et al., 2018; Sander et al., 2019; Sommer et al., 2019). Participants' sleep was monitored in two successive nights before and after engaging in an extensive associative learning task. An adaptation night prior to the first experimental night familiarized participants with the PSG procedure. MRI data were collected after the second experimental night. The research protocol was approved by the Ethics Committee of the *Deutsche Gesellschaft für Psychologie* (DGPs) and conducted at the *Max Planck Institute for Human Development* in Berlin. Informed written consent was obtained from all participants before study participation.

Sleep data acquisition

Sleep was recorded in two consecutive nights using an ambulatory PSG device (SOMNOscreen plus; SOMNOmedics, Germany). Eight scalp electrodes were attached according to the international 10–20 system for electrode positioning (Fz, F3, F4, C3, C4, Cz, Pz, Oz; Jasper, 1958; Oostenveld and Praamstra, 2001) along with two electrodes on the mastoids A1 and A2, which later served as the offline reference.

All impedances were kept below 6 kΩ. Data were recorded using Cz as the online reference for all EEG derivations and AFz as ground. Additionally, a bilateral electrooculogram (EOG) was assessed. Two submental electromyogram channels (EMG) were positioned left and right, below the labial angle, and referenced against one chin electrode. Electrical activity of the heart was recorded using two electrocardiogram channels (ECG). EEG channels were recorded between 0.2–75 Hz with a sampling rate of 128 Hz.

EEG pre-processing

Data was first preprocessed using *BrainVision Analyzer 2.1* (Brain Products, Germany). Here, all EEG channels were re-referenced against the average of A1 and A2. To visually score the sleep data according to the standard criteria suggested by the *American Academy for Sleep Medicine* (AASM; Iber et al., 2007), the data were filtered using their recommended settings (EEG: 0.3–35 Hz; EOG: 0.3–35 Hz; EMG: 10–100 Hz). A Notch (50 Hz) filter was applied to the EMG and EOG derivations to remove external electrical interference. Afterwards, sleep stages 1 and 2, SWS, REM sleep, awakenings, and body movements were visually scored in 30-second epochs using *SchlafAus 1.4* (Gais & Werner, Lübeck, Germany). Total sleep time (TST) was calculated as the time spent in stage 1, stage 2, SWS, and REM sleep. Wake after sleep onset (WASO) was defined as the time participants were awake between sleep onset and final morning awakening.

Further analyses of the sleep EEG data were conducted using *Matlab R2014b* (Mathworks Inc., Sherbom, MA), the open-source toolbox *Fieldtrip* (Oostenveld et al., 2011), and custom-made code. Bad EEG channels were visually rejected. For the remaining channels, artefact detection was implemented on 1-second segments. Segments that were visually identified as body movement and those that exhibited amplitude differences of more than 500 µV were excluded. To further exclude segments that strongly deviated from the observed overall amplitude distribution, mean amplitude differences for each segment were z-standardized within each channel. Segments with a z-score of more than 5 in any of the channels were excluded.

Slow oscillation detection

Slow oscillation detection at frontal electrodes was implemented based on previously established algorithms (Mölle et al., 2002; Ngo et al., 2013; cf. Muehlroth et al., 2019b). For NREM epochs (i.e., stage 2 sleep, and SWS), EEG data was band-pass filtered between 0.2 and 4 Hz using a 6th-order Butterworth filter (in forward and backward directions to prevent phase distortions). The whole signal

was then divided into negative and positive half-waves that were separated by zero-crossings of the filtered signal. The combination of a negative half wave with the succeeding positive half wave was considered a putative slow oscillation when its frequency was between 0.5 and 1 Hz. The amplitude of each potential slow oscillation was calculated as the distance between the slow oscillation trough and positive peak, defined as its maximal negative and positive potential, respectively. Selection of real slow oscillations was then performed in two ways:

1. Adaptive amplitude thresholds were defined separately for each participant: Putative slow oscillations exceeding a trough of 1.25 times the mean trough of all putative slow oscillations as well as an amplitude of 1.25 times the average amplitude of all potential slow oscillations, were considered real slow oscillations (Ngo et al., 2013).

2. As an alternative detection criterion, fixed amplitude thresholds were defined: Putative slow oscillations exceeding a trough of –40 µV as well as an amplitude of 75 µV, were selected (Carrier et al., 2011; Dubé et al., 2015). Only artefact-free slow oscillations were considered in further analyses.

Spindle detection

Spindles were detected using an established automated algorithm (Klinzing et al., 2016, 2018; Mölle et al., 2011). For NREM epochs, EEG data were band-pass filtered using a 6th-order Butterworth filter (in forward and backward directions) between 9 and 12.5 Hz for the detection of slow spindles, and between 12.5 and 16 Hz for fast spindles, respectively (Cox et al., 2017; Ujma et al., 2015). The root-mean-square (RMS) representation of the signal was calculated using a sliding window of 200 ms at each sample point. Afterwards, the RMS signal was smoothed by applying a moving average of 200 ms. To increase sensitivity and specificity of our algorithm (Coppieters 't Wallant et al., 2016), we accounted for individual differences in EEG amplitude by anchoring the spindle identification to individually determined amplitude thresholds. A potential spindle was tagged if the amplitude of the smoothed RMS signal exceeded its mean by 1.5 SD of the filtered signal for 0.5 to 3 seconds. Spindles with boundaries closer than 0.25 seconds were eventually merged in the following way: In each processing run, starting with the smallest boundary difference, two spindles were merged if the resulting spindle event remained within the time limit of 3 seconds. As soon as a spindle was combined with another one, it could not be merged with another spindle in the same run again. Only if all boundary differences were revised and no further merging of putative spindles was possible, the new resulting spindle events were processed

in a next run. Starting again with the smallest boundary difference, two putative spindles could be merged if the merged event remained within the required time limit. These runs were repeated until no further merging was possible. Finally, only spindles that did not overlap with previously marked artefact segments were considered real spindles.

Power spectral analysis

To derive whole-night characteristics of spectral EEG power (cf. Figure 2), Fast Fourier Transformation (FFT) with frequency limits between 0.2 and 30 Hz was applied at electrode Cz on 5-second intervals using a Hanning function with no overlap. The resulting whole-night spectral data was log-transformed. Power spectral analyses of the sleep EEG data were conducted, to obtain finer-grained indicators of neural processes during deep NREM sleep. FFT with frequency limits between 0.5 and 30 Hz was applied on 5-second intervals using a Hanning function with no overlap and averaged across all NREM sleep. Log-transformed power spectra were derived for all channels. Slow-wave activity (SWA), defined as spectral power in the delta frequency range (0.5–4.5 Hz), was calculated for NREM epochs and normalized at the total power of the whole spectrum to minimize confounding effects like skull thickness (e.g., Dannhauer et al., 2011; Frodl et al., 2001; Leissner et al., 1970).

Statistical analysis

Due to non-normal data distributions, we calculated non-parametric Mann-Whitney U Tests for independent samples and reported median and quartile values of the variables o identify age differences in sleep architecture and the expressed sleep oscillations between younger and older adults. Effect sizes were quantified by calculating Cohen's d along with its 95 % confidence interval. Mixed factorial ANOVAs with the between factor AGE GROUP and the within factor CHANNEL were calculated to investigate the topographical differences in age-related sleep alterations. Post-hoc testing, again, was conducted using non-parametric Mann-Whitney U Tests for independent samples as well as Wilcoxon signed-rank tests for matched pairs. Associations between variables were estimated using Spearman's rank-order correlation coefficient. The agreement between different sleep variables was quantified by calculating the *intraclass correlation coefficient* (ICC; Koo and Li, 2016) based on a two-way mixed effects model and consistency (= ICC(3,k)). To derive the test-retest reliability of the same sleep variable between the two consecutive nights, ICC calculation was based on a two-way mixed effects model and absolute agreement (= ICC(2,k)). In general, significance levels were set to $\alpha = .05$ and tested two-sided.

We controlled statistical significance for multiple comparisons by Bonferroni-correcting the α-value and dividing it by the number of performed comparisons.

Structural MRI data acquisition and analyses

Whole-brain MRI data was acquired with a Siemens Magnetom 3T TimTrio machine. A high-resolution T1-weighted MPRAGE sequence (TR = 2500 ms, TE = 4.77 ms, FOV = 256 mm, voxel size = 1 × 1 × 1 mm^3) was collected from each participant. To quantify gray matter volume voxel-wise, we applied voxel-based morphometry (VBM; Ashburner and Friston, 2000) using *Statistical Parametric Mapping Software* (SPM12, http://www.fil.ion.ucl.ac.uk/spm) and the *Computational Anatomy Toolbox* (CAT 12, http://www.neuro.uni-jena.de/cat). Images were normalized to Montreal Neurological Institute (MNI) space and segmented into gray matter, white matter, and cerebrospinal fluid. Data were then modulated, i.e., corrected by the volume changes due to spatial normalization. For this, each voxel value was multiplied with the Jacobian determinant derived from the spatial normalization step. Afterwards images were smoothed with a 8 mm full width at half maximum (FWHM) kernel. Total intracranial volume (TIV) was estimated by summing volume of gray matter, white matter, and cerebrospinal fluid.

To derive the association between gray matter volume, age, and specific sleep variables, voxel-vise multiple regression models were implemented. For each subject, sleep estimates were averaged between the two experimental nights. All regression models were controlled for inter-individual differences in TIV by entering TIV as a covariate into the model. For masking, an absolute threshold of .02 was used. A family-wise error (FWE) correction was applied to all statistical maps, and α was set to .05. Further, we deployed a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$ as well as a correction for nonstationary smoothness. Finally, we used the *xjView toolbox* (http://www.alivelearn.net/xjview/) to plot the resulting statistical t-maps onto the Ch2bet MNI template and to derive cluster labels.

Supplementary Material

Supplementary Figure 1. Correspondence between different indicators of slow oscillatory activity. Spearman's rank-order correlation coefficients within and across age groups were calculated between the different indicators (corresponding *p*-values in brackets). All slow oscillation characteristics were derived using individually adjusted amplitude thresholds. The distribution of the respective variable within younger (orange) and older (blue) is displayed on the diagonal. Most variables show strong differences between age groups. Correlation coefficients and corresponding scatter plots are displayed in the upper right and lower left triangle respectively. Only few variables are closely correlated. OA: older adults; SO: slow oscillation; SWA: slow-wave activity; SWS: slow-wave sleep; YA: younger adults.

Supplementary Figure 2. Log-transformed EEG power during NREM sleep. (A) In younger adults (YA) a clear peak in the fast spindle frequency range (12.5–16 Hz; shaded in dark gray) is observed in centroparietal electrodes. Frontal channels locally peak in lower frequencies at the transition from the slow (9–12.5 Hz, shaded in light gray) to fast spindle band. (B) In older adults (OA) we observe the same fast spindle power peak, but in Pz only. (C) For each age group the average power spectrum is depicted with its 95 % bootstrap confidence interval. Overall, the power spectra are flatter in OA (blue) than in YA (orange). YA show a stronger fast spindle power peak (shaded in dark grey) and higher power values in low frequencies (0–9 Hz) compared to OA.

Supplementary Figure 3. Significant negative cluster of the association between gray matter volume and age (multiple regression, controlled for total intracranial volume). The cluster's peak intensity voxel is highlighted (glass brain (A): orange arrow; Ch2bet template brain (B): blue hair cross). The cluster is displayed and considered significant at a FWE-corrected voxel threshold of $p < .05$, and a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$. It is corrected for nonstationary smoothness.

Supplementary Figure 4. Significant negative clusters of the association between gray matter volume and fast spindle density at different electrode sites (multiple regression, controlled for total intracranial volume). The clusters' peak intensity voxel is highlighted (glass brain: orange arrow; Ch2bet template brain: blue hair cross). Clusters are displayed and considered significant at a FWE-corrected voxel threshold of $p < .05$, and a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$. All clusters are corrected for nonstationary smoothness.

Supplementary Figure 5. Significant negative clusters of the association between gray matter volume and slow spindle density at different electrode sites (multiple regression, controlled for total intracranial volume). The clusters' global peak intensity voxel is highlighted (glass brain: orange arrow; Ch2bet template brain: blue hair cross). Clusters are displayed and considered significant at a FWE-corrected voxel threshold of $p < .05$, and a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$. All clusters are corrected for nonstationary smoothness.

Supplementary Figure 6. Significant negative cluster of the association between gray matter volume and slow oscillation density measured at Oz (multiple regression, controlled for total intracranial volume). The cluster's global peak intensity voxel is highlighted (glass brain: orange arrow; Ch2bet template brain: blue hair cross). The cluster is displayed and considered significant at a FWE-corrected voxel threshold of $p < .05$, and a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$. All clusters are corrected for nonstationary smoothness.

Supplementary Figure 7. Significant negative clusters of the association between gray matter volume SWA at different electrode sites (multiple regression, controlled for total intracranial volume). The clusters' global peak intensity voxel is highlighted (glass brain: orange arrow; Ch2bet template brain: blue hair cross). Clusters are displayed and considered significant at a FWE-corrected voxel threshold of $p < .05$, and a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$. All clusters are corrected for nonstationary smoothness.

Supplementary Table 1. Test-retest reliability for different markers of slow oscillatory activity

		ICC [95 % CI]	$F(df_1, df_2)$	p
% SWS	Whole sample	.96 [.93; .98]	$F(62,62) = 25.17$	$< .001$
	YA	.9 [.78; .95]	$F(28,28) = 9.52$	$< .001$
	OA	.94 [.75; .98]	$F(33,33) = 23.83$	$< .001$
Relative SWA	Whole sample	.82 [.69; .90]	$F(62,62) = 6.19$	$< .001$
	YA	.83 [.64; .92]	$F(28,28) = 5.95$	$< .001$
	OA	.68 [.33; .85]	$F(33,33) = 3.69$	$< .001$
SO amplitude	Whole sample	.96 [.94; .98]	$F(62,62) = 25.96$	$< .001$
	YA	.94 [.87; .97]	$F(28,28) = 16.42$	$< .001$
	OA	.86 [.73; .93]	$F(33,33) = 7.35$	$< .001$
SO frequency	Whole sample	.89 [.82; .93]	$F(62,62) = 8.99$	$< .001$
	YA	.88 [.73; .94]	$F(28,28) = 7.98$	$< .001$
	OA	.85 [.74; .93]	$F(33,33) = 7.58$	$< .001$
SO number	Whole sample	.84 [.74; .91]	$F(62,62) = 6.35$	$< .001$
	YA	.83 [.64; .92]	$F(28,28) = 5.98$	$< .001$
	OA	.78 [.57; .89]	$F(33,33) = 4.61$	$< .001$
SO density	Whole sample	.88 [.79; .92]	$F(62,62) = 7.97$	$< .001$
	YA	.92 [.84; .96]	$F(28,28) = 13.08$	$< .001$
	OA	.81 [.62; .91]	$F(33,33) = 5.33$	$< .001$

Note. Test-retest reliability was calculated between the two recorded experimental nights based on a two-way mixed effects model and absolute agreement. CI: confidence interval; ICC: intraclass correlation coefficient; OA: older adults; SO: slow oscillation; YA: younger adults.

Supplementary Table 2. Brain regions associated with age in a voxel-vise multiple regression, controlled for total intracranial volume.

Anatomical peak region	Hemisphere	Peak intensity (*t*)	Peak MNI coordinates			Number of voxels
			x	y	z	
Middle frontal gyrus, orbital part	R	−15.6536	39	36	−19.5	167967

Note. The table lists cluster size along with the cluster's peak intensity (maximal *t*-value, FWE-corrected) and corresponding MNI coordinates. Anatomical classification is based on the respective peak voxel. Reported clusters are significant with a FWE-corrected voxel threshold of $p < .05$, a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$ and corrected for nonstationary smoothness.

Supplementary Table 3. Brain regions associated with fast spindle density at different electrode sites (voxel-vise multiple regressions, controlled for total intracranial volume).

Anatomical peak region	Hemisphere	Peak intensity (t)	Peak MNI coordinates			Number of voxels
			x	y	z	
Frontal fast spindle density						
Middle frontal gyrus	R	8.9326	33	52.5	25.5	2223
Brodmann area 25	R	8.5816	6	19.5	−3	780
Middle frontal gyrus	L	8.3087	−28.5	39	−10.5	782
Inferior frontal gyrus, triangular part	L	8.2467	−40.5	27	24	410
Anterior cingulate	R	8.1811	12	48	0	2328
Lingual gyrus	L	8.0199	−6	−81	−1.5	884
Inferior occipital gyrus	L	7.9767	−42	−81	−6	267
Superior temporal gyrus	L	7.8768	−57	-46.5	13.5	1037
Angular gyrus	R	7.7104	48	−61.5	31.5	266
Inferior frontal gyrus, triangular part	R	7.1719	40.5	13.5	22.5	253
Brodmann area 7	R	7.1272	7.5	−69	31.5	288
Brodmann area 18	L	7.1108	−30	−93	0	239
Brodmann area 21	R	6.8967	58.5	−28.5	−4.5	898
Brodmann area 22	L	6.0636	−49.5	−4.5	4.5	138
Superior frontal gyrus	L	6.0436	−22.5	52.5	10.5	241
Cz fast spindle density						
brodmann area 11	L	12.1947	−30	40.5	−12	20652
Brodmann area 21	R	9.8022	61.5	−27	−4.5	4135
Brodmann area 22	L	8.5415	−54	3	0	2316
Caudate head	L	8.4566	−3	9	−4.5	2404
Brodmann area 7	R	8.2929	9	−64.5	30	811
Brodmann area 22	L	8.2923	−58.5	−49.5	10.5	2274
Angular Gyrus	R	8.1438	49.5	−63	31.5	496
Postcentral gyrus	L	7.4516	−43.5	−36	51	597
Declive	L	7.3935	−37.5	−58.5	−22.5	449
Brodmann area 9	R	6.8413	46.5	12	30	223
Postcentral gyrus	R	6.7044	54	−25.5	37.5	130
Brodmann area 22	R	6.6719	49.5	3	−1.5	242
Brodmann area 18	L	6.5612	−6	−88.5	6	207
Brodmann area 34	L	6.1602	−28.5	1.5	−12	105
brodmann area 31	L	5.9249	−9	−49.5	37.5	109
Cingulate gyrus	R	7.883	7.5	−34.5	42	967
Middle occipital gyrus	L	7.759	−42	−78	−4.5	272
Parahippocampal gyrus	R	6.998	37.5	−28.5	−15	114
Parahippocampal gyrus	R	5.974	22.5	−37.5	−6	197
Pz fast spindle density						
No significant cluster						
Oz fast spindle density						
No significant cluster						

Note. The table lists cluster size along with the cluster's peak intensity (maximal *t*-value, FWE-corrected) and corresponding MNI coordinates. Anatomical classification is based on the respective peak voxel. Reported clusters are significant with a FWE-corrected voxel threshold of $p < .05$, a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$, and corrected for nonstationary smoothness.

Supplementary Table 4. Brain regions associated with slow spindle density at different electrode sites (voxel-vise multiple regressions, controlled for total intracranial volume).

Anatomical peak region	Hemisphere	Peak intensity (*t*)	Peak MNI coordinates			Number of voxels
			x	y	z	
Frontal slow spindle density						
Inferior frontal gyrus	R	6.788	46.5	9	1.5	215
Superior frontal gyrus	L	6.7206	−19.5	15	54	136
Cz slow spindle density						
Brodmann area 9	R	7.2931	52.5	4.5	28.5	100
Pz slow spindle density						
No significant cluster						
Oz slow spindle density						
No significant cluster						

Note. The table lists cluster size along with the cluster's peak intensity (maximal *t*-value, FWE-corrected) and corresponding MNI coordinates. Anatomical classification is based on the respective peak voxel. Reported clusters are significant with a FWE-corrected voxel threshold of $p < .05$, a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$, and corrected for nonstationary smoothness.

Supplementary Table 5. Brain regions associated with slow oscillation density at different electrode sites (voxel-vise multiple regressions, controlled for total intracranial volume).

Anatomical peak region	Hemisphere	Peak intensity (*t*)	Peak MNI coordinates			Number of voxels
			x	y	z	
Frontal slow oscillation density						
No significant cluster						
Cz slow oscillation density						
No significant cluster						
Pz slow oscillation density						
No significant cluster						
Oz slow oscillation density						
Middle temporal gyrus	L	6.4661	−51	−63	−1.5	155

Note. The table lists cluster size along with the cluster's peak intensity (maximal *t*-value, FWE-corrected) and corresponding MNI coordinates. Anatomical classification is based on the respective peak voxel. Reported clusters are significant with a FWE-corrected voxel threshold of $p < .05$, a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$, and corrected for nonstationary smoothness.

Supplementary Table 6. Brain regions associated with SWA at different electrode sites (voxel-vise multiple regressions, controlled for total intracranial volume).

Anatomical peak region	Hemisphere	Peak intensity (t)	Peak MNI coordinates x	y	z	Number of voxels
Frontal SWA						
Insula	R	9.1294	45	−4.5	−1.5	6160
Gyrus rectus	R	9.01080	0	57	−19.5	2676
Insula	L	8.7764	−43.5	−9	−6	4975
Brodmann area 31	L	7.7713	−1.5	−30	48	1114
Middle frontal gyrus	L	7.6467	−25.5	31.5	45	376
Brodmann area 46	R	7.5113	46.5	43.5	10.5	335
Middle temporal gyrus	L	7.1541	−52.5	−70.5	24	507
Cingulate gyrus	L	7.1391	-6	−10.5	49.5	401
Parahippocampa gyrus	L	7.0887	−22.5	−21	−15	265
Middle frontal gyrus	R	6.9864	37.5	37.5	39	220
Declive	R	6.8567	9	−64.5	−19.5	1056
Supramarginal gyrus	L	6.5642	52.5	−55.5	27	328
Precuneus	R	6.1072	−3	−70.5	27	255
Brodmann area 46	L	9.222	−49.5	39	7.5	1423
Middle frontal gyrus	R	6.941	52.5	25.5	27	121
Cz SWA						
Superior temporal gyrus	R	7.4072	36	3	−19.5	1381
Insula	L	6.3966	−43.5	−9	−6	218
Inferior frontal gyrus	L	6.3871	−46.5	24	−3	154
Brodmann area 46	L	7.3826	−48	39	15	432
Middle frontal gyrus	R	6.3796	31.5	52.5	7.5	145
Brodmann area 9	L	7.3294	−15	58.5	31.5	960
Transverse temporal gyrus	R	6.2499	49.5	−24	10.5	159
Superior temporal gyrus	L	6.1291	−40.5	9	−15	130
Brodmann area 46	R	7.0373	48	42	9	224
Declive	R	6.4692	19.5	−72	−24	786
Superior temporal gyrus	L	5.8842	−40.5	−21	6	108
Superior temporal gyrus	R	5.9135	54	−57	27	167
Middle temporal gyrus	R	6.141	63	−21	−6	123
Pz SWA						
Insula	R	8.0873	45	−7.5	1.5	3170
Inferior parietal lobule	L	7.8406	−54	−28.5	24	4601
Brodmann area 10	L	7.8054	−3	66	7.5	2513
Middle frontal gyrus	L	7.6457	−25.5	31.5	45	186
Inferior frontal gyrus	R	7.5282	51	37.5	16.5	289
Brodmann area 6	L	7.3516	−46.5	−3	52.5	349
Medial frontal gyrus	R	7.2276	1.5	−19.5	60	746
Inferior occipital gyrus	R	7.1093	45	−63	−16.5	178
Middle temporal gyrus	L	6.9624	−48	−78	18	205
Medial frontal gyrus	L	6.8258	−9	27	−13.5	285
Supramarginal gyrus	R	6.6716	52.5	−55.5	27	407
Inferior frontal gyrus	R	6.5307	43.5	24	−4.5	470
Cuneus	R	6.1539	0	−69	30	174
Brodmann area 32	L	6.0906	−1.5	16.5	34.5	168
Declive	R	6.68	7.5	−66	−21	588
Oz SWA						
No significant cluster						

Note. The table lists cluster size along with the cluster's peak intensity (maximal t-value, FWE-corrected) and corresponding MNI coordinates. Anatomical classification is based on the respective peak voxel. Reported clusters are significant with a FWE-corrected voxel threshold of $p < .05$, a FWE-corrected cluster extent threshold of $p < .05$ and $k > 100$, and corrected for nonstationary smoothness.