Primate hippocampus size and organization are predicted by sociality but not diet

Orlin S. Todorov[1], Vera Weisbecker[1], Emmanuel Gilissen[2], Karl Zilles[3], and Alexandra A. de

Sousa[4]

[1] School of Biological Sciences, The University of Queensland, St. Lucia, QLD,

4072, Australia

[2] Department of African Zoology, Royal Museum for Central Africa, Leuvensesteenweg,

3080 Tervuren, Belgium and Laboratory of Histology and Neuropathology, Université Libre

de Bruxelles, Brussels, Belgium

[3] Research Centre Ju☐lich, Institute of Neuroscience and Medicine (INM-1), Ju☐lich,

Germany,

[4] Psychology, Culture and Environment, Bath Spa University, Bath, BA2 9BN, UK

26   hippocampus, social group, spatial cognition, primates

28   **Abstract**

The hippocampus is well known for its roles in spatial navigation and memory, but it is

organized into regions that have different connections and functional specializations. Notably,

the region CA2 has a role in social and not spatial cognition, as is the case for the regions

CA1 and CA3 that surround it. Here we investigated the evolution of the hippocampus in

terms of its size and organization in relation to the evolution of social and ecological variables

in primates, namely home range, diet and different measures of group size. We found that the

volumes within the whole cornu ammonis coevolve with group size, while only the volume of

CA1 and subiculum can also be predicted by home range. On the other hand, diet, expressed

as a shift from folivory toward frugivory, was shown to not be related to hippocampal

volume. Interestingly, CA2 was shown to exhibit phylogenetic signal only against certain

measures of group size but not with ecological factors. We also found that sex differences in

the hippocampus are related to body size sex dimorphism. This is in line with reports of sex

differences in hippocampal volume in non-primates that are related to social structure and sex

differences in behaviour. Our findings support the notion that in primates, the hippocampus

might be more important in social, rather than in spatial processing.

1. Introduction

The relationship between vertebrate behaviour and brain size and proportions has been the topic of intensive research for decades, with works on mammals focusing mainly on the question of how the exceedingly large brains of primates, and particularly humans, could evolve. However, while there is an emerging consensus on the energetic constraints on the evolution of brain enlargement [1, 2], the search for behavioural correlates of relative brain size has a long history of producing a frustrating diversity of results [3]. In primates, a long-standing debate is whether brain size is correlated with social or ecological intelligence. Models of primate social intelligence and brain size emphasize social skills, including managing social complexity, theory of mind, social learning, and culture [4, 5]. On the other hand, models of ecological intelligence demonstrate an important impact of home range size and/or diet [6-9] on relative brain size. Evidence about which of these (home range or diet) is the main determinant of brain size is ambiguous [6] [7], but both possibly relate to the memory demands of locating and identifying unpredictable food sources or mates [10-13]. An important caveat to studies of "intelligence" and brain morphology is the fact that most analyses of brain morphology addressing social and ecological factors across primate evolution only consider how they relate to overall brain size. However, there is increasing awareness that overall brain size may only be a rough proxy of brain capacity [14], consistent with long-standing evidence that the brain is a mosaic of different regions, which may respond differentially to selection for specific behaviours [15-18]

The mammalian hippocampus is of particular interest in terms of dissecting the morphological correlates of ecological and social behaviour. This is particularly the regarding "ecological intelligence" which is often framed in terms of spatial awareness of an animal in its environment. The hippocampus contains a population of neurons ('place cells') that respond whenever an animal is in a specific location [19] and these produce a dynamic 'cognitive

map' of the environment by firing in a concerted fashion [20]. Similarly, the entorhinal

cortex, a structure neighbouring the hippocampus in the larger "hippocampal complex", has a

population of "grid cells", which fire when an animal enters an environment with

geometrically patterned locations [21]. Another component of hippocampus-related

"ecological intelligence" is its function in memory consolidation [22], possibly through a

spatial-based mechanism [23]. The hippocampus also has a role in behavioural inhibition [24]

and olfactory memory [25].

Relatively little is known about the correspondence between hippocampal organization and

ecological variables. In desert rodents, the bannertail kangaroo rat has relatively low spatial

memory requirements and has a small hippocampus, whereas Merriam's kangaroo rat uses

spatial memory to relocate its caches in scattered locations, and larger hippocampus [26]. The

"avian hippocampus" in the medial pallial zone is homologous to that in mammals and also

functions in spatial memory [27]. This is consistent with the fact that food-storing birds have

relatively larger hippocampi [28, 29].

The size and internal organization of the hippocampus is also subject to within-species

variation and individual plasticity. Volumetric reorganization of the hippocampus has been

related to the occupational specialization in humans [30]. In birds, hippocampal size and

structure is plastic, being affected by experience [31], and seasonality [32]. In arboreal

primates, a relationship was found between hippocampus size and home range size [33], but

overall, this relationship remains unclear [33] [34]. The possibility for a predictive function of

the hippocampus is particularly evident from studies of sexual dimorphism in hippocampal

size and spatial ability. Whereas male and female meadow voles are sexually dimorphic in

their performance on spatial tasks, hippocampus volume, and home range size, pine voles are

not [35]. Further, in two other polygamous rodent species the relative size of the hippocampus

is greater in males than in females [26], while males and females of the monogamous desert

kangaroo rat do not differ in home range nor in spatial ability [36]. Similarly, during breeding

season, deer mice are polygynous and males have larger home ranges, and outperform

females on spatial tasks [37]. Sex differences in spatial ability and home range size are also

related in two species of carnivores - males exhibit larger home ranges and superior spatial

ability compared to females in the promiscuous giant pandas, but not in the monogamous

Asian small clawed otter [38]. Consistent with the hypothesis that function drives anatomy,

the sex differences are reversed in wider ranging females. In a brood parasite bird species, the

brown-headed cowbirds, females which travel further than males have larger hippocampi [39]

and exhibit superior spatial memory [40].

As of recently, some light has been shed on the role of the hippocampus in social behaviour

and cognition. Hippocampal place cells are involved in processing the presence of

conspecifics in bats [41] and hippocampal volume has been related to social phobia as part of

adjacent circuits in humans [42]. Although the representation system of the hippocampal

complex is itself spatial, this coordinate system is capable of processing other spatially

representable information – such is the case of its role as a "memory map" for encoding

declarative memories [23], or social information [43]. In rats, support for the mechanism

comes from studies finding the hippocampus (specifically a substructure described below,

CA2) uses place fields to encode information about conspecifics [44]. Given these novel

insights into hippocampus function, in species where social behaviour plays an important role,

the involvement of the hippocampus in social information processing might be greater. This

also has implications for linking social and spatial cognition more generally, as they can be

represented in the same cognitive systems [45].

Hippocampal regions

All fields of the hippocampus formation (retrohippocampus, RH) receive inputs from the

entorhinal cortex (EC) along the perforant pathway [46]. Part of it, hippocampus proper,

refers to the cornu ammonis (CA) and the fascia dentata (FD); more commonly these same

regions are divided up into CA1-3 and the dentate gyrus (DG) (Table 1). DG has traditionally

been considered the gateway of the hippocampus because it blocks or filters excitatory

afferents from the EC [47]. Sensory and associative projections from the EC synapse in the

DG [48]. DG arranges sensory inputs to create a metric spatial representation and is involved

in episodic memory and spontaneous exploration of novel environments [49]. DG can be

further subdivided into the fascia dentata (FD) and the hilus (part of the CA). Adjacent to the

FD, the CA is comprised of four fields arranged in a loop, beginning with the hilus (i.e., CA4)

[50]. The hilus is situated along the mossy fiber pathway from the granular stratum of FD to

CA3 and is involved in spatial learning and memory retrieval [51]. It has a role in sequence

learning [52], and local lesions affect pattern separation, particularly for highly similar inputs

[53]. Next are the sequential CA regions in descending order - CA3, CA2, CA1. CA3 receives

connections from the mossy fibers of FD, which it projects to CA1 and back, bypassing CA2.

There are associational bilateral (ipsilateral and contralateral) connections to CA3 [54]. CA3

can be further divided into subregions: CA3a and CA3b encode spatial information into short-

term memory, while CA3c processes environmental geometry along with DG [55]. CA1

receives projections from CA3 and is involved in spatial memory [56]. The spatial properties

of CA1 and CA3 are due to these regions being the primary locations of 'place cells',

responding differentially according to the spatial location of the animal [57]. Adjacent to

CA3, the subiculum has inputs from EC and bilateral connections with perirhinal cortex and

CA1 [58]. It is is a major output of the hippocampus with pronounced dorso-ventral

segregation of function: the dorsal component is involved in processing of spatial information

and information related to movement and memory, while the ventral is a type of interface

between the hippocampus and the hypothalamic–pituitary–adrenal axis, a feedback system

that regulates homeostasis and stress [58]. The subiculum receives projections mostly from

CA1 and these are organized in a simple pattern - all sections of CA1 project to the subiculum

and all parts of the subiculum receive input from CA1 [59]. Moreover, subicular neurons

exhibit spatially-selective firing [58] with a robust location signal [60].

<<Figure 1 about here>>

CA2 has been suggested to act as an interface between emotion and cognition [61]. CA2

receives strong inhibitory inputs from EC, CA3, and DG, and has outputs to CA1 [62]. It is

also influenced by many neuromodulators, receiving unique input from hypothalamic nuclei

associated with social context, reward, and novelty [61] – supramammillary, paraventricular,

median raphe, septal, and the vertical and horizontal limbs of the nucleus of diagonal band of

Broca [62]. CA2 has outputs to septum and the supramammillary nucleus. Unlike CA1 and

CA3, lesions to CA2 do not affect spatial memory in Morris water maze test, nor impact

locomotor ability, anxiety or fear memory in rodents [63]. Rather, CA2 is involved in social

memory and recognition of conspecifics [63]. There are some indications its size may be

particularly adaptive to social and emotional experiences - decrease in CA2 neuron numbers

is associated with schizophrenia and bipolar disorder [64] and stress-related increases in the

density of brain-derived neurotrophic factor neurons are greater in CA2 than CA3 [65].

Here we investigate the evolution of hippocampal size and organization in primates, in

relation to social and ecological pressures. Given the importance of the hippocampus in

spatial cognition, and the subiculum, CA1, CA3, and FD in particular, we predict that these

will be related to variation in ecological variables: home range size and/or dietary complexity.

Additionally, we predict CA2 volume to be related to social memory, measured through

group size. We also expect that amongst brain areas, dimorphism in hippocampal size will be

the best predictor of dimorphism in body size.

175   **2. Methods**

(a) Anatomical data

The morphometric structure of the hippocampus was determined from previously published

data, on bilateral volumes [66]. For measurements, the retrohippocampus (RH) has been

divided into: dentate gyrus (reported in [66] as *fascia dentata*, FD), *hilus* (HIL), *CA3*, *CA2*,

CA1, and *subiculum* (SUB). Volume measurements include the white matter comprising the

rest of the hippocampus [66, 67] measured together as *HP+HS+fibers*, that is the

hippocampus praecommissuralis (HP) plus the hippocampus supracommissuralis (HS) plus

the fibers of the fimbria/fornix complex. Volumes for whole *brain* were taken from the same

source [66]. Volumes for *neocortex* (white and grey matter; NEO) were obtained from the

same research group [67].

Unpublished data on brain component volumes of males and females were used to determine

averages for each sex in a subsample of primates, and correspond to anatomical definitions in

[67]. The brain components include 7 telencephalic components: *bulbus olfactorius + bulbus*

olfactorius accessorius (bulbus olfactorius accessorius is absent in higher primates; BOL),

lobus piriformis (palaeocortex and amygdala; PAL), *septum* (septum pellucidum, septum

verum, Broca's diagonal band, bed nuclei of the anterior commissure and stria terminalis;

SEP), *striatum* (caudate nucleus, putamen, nucleus accumbens, and the parts of the capsula

interna running through the striatum; STR), *schizocortex* (ento- and perirhinal, pre- and

parasubicular cortices and the underlying white matter; SCH), *hippocampus* (including all

regions; HIP), *neocortex* (white and grey matter; NEO). Iincluded were *diencephalon* (plus

globus pallidus without hypophysis; DIE), *mesencephalon* (without substantia reticularis;

MES), *cerebellum* (brachium and nuclei pontis, CER), and *medulla oblongata* (plus

substantia reticularis; MED). Body weight (BoW) data was available for the same individuals,

except for *Miopithecus talapoin* female body weight, which was taken from [68]. Sexual size

dimorphism was determined from BoW and calculated as the ratio of male BoW divided by

female BoW. Sexual dimorphism in each of the brain structures was calculated as the ratio of

the volume in males vs females.

(b) Social and ecological data

Data were collated from three different sources. Home range area in hectares "HR size

average" (HR) were from Powell et al. [7], frugivory "% fruit" were from DeCasien et al. [6],

"group size combined" were from [7]. Further, "social group size" data are from [6] and

"mean group size" and "mean number of females per group" are from Dunbar et al. [69].

These different studies use different methods for collating the datasets, where it is not always

clear whether group size indicates social or foraging group, or whether diet information has

been calculated uniformly and reliably.

(c) Phylogeny

The consensus phylogenetic tree of 43 species of apes and monkeys was obtained from 10k

Trees [70] and information about phylogenetic non-independence was incorporated in all

analysis. Changes in taxonomic nomenclature were considered for matching species names

from the brain dataset to the tree.

(d) Statistical analysis

All continuous variables were natural log transformed and min-max standardized, except for % fruit, which was standardized only. Bonferroni correction was applied on the α level ("significance cut-off" of 0.05) on models tested multiple times by dividing it by the number of comparisons with the same dependent variable (6 models resulting in corrected α of 0.0083).

Analyses were run on R version 3.6.1 [71] using the packages phytools [72], caper [73] and MuMIn [74]. Using the fastanc function in phytools we estimated the ancestral states and painted them on the tree using Fancytree. We used caper for all PGLS analyses. Phylogenetic signal (Pagel's λ) was estimated using Maximum Likelihood and kappa (k) and delta (δ) were fixed to 1. We tested three specific models that might help explain the variation in hippocampus and hippocampal region volumes: two ecological and social. Additionally, we explored the relationship between neocortex and brain volume with hippocampus volume. F statistics were obtained via sequential sum of squares ANOVA.

The volume of the region of interest was always used as the dependent variable in our models, and brain volume was included as a covariate within each model. All variables were shown to be normally distributed, and variance inflation factors of each models were shown to be <3.5 meaning that there was no problem with collinearity. Interactions between predictors were not included as to avoid high cross-collinearity.

We did not test a 'full model' including home range, diet and group sizes together, because of the way PGLS deals with missing data (see Supplementary for summary of the missing data). This would have reduced our sample size substantially (down to 16 degrees of freedom) causing a loss of 66% of the data available (but see 'full model' in Supplement).

In order to verify our findings, we conducted additional model dredging with the MuMIn package, where we used hippocampus and every region volume as dependent variable, against

245   all possible ecological and social variables. This approach tests all possible model

246   combinations of predictor combinations and ranks them using AICc (Akaike Information

247   Criterion, with penalising correction for sample size). [74].

249   **3. Results**

(a) Ancestral state estimation

An exploratory ancestral state estimation revealed that in species where relative hippocampal

volume has decreased (calculated as the residuals from the phylogenetic regression with total

brain volume) have nonetheless undergone an increase in absolute hippocampus volume (Fig.

2). We further tested this observation using PGLS and found that hippocampus volume

increased with a shallower slope compared to both brain and neocortex volumes i.e. species

that evolve towards greater neocorticalization have smaller relative hippocampi. (See

Neocortex section). An exception is the pygmy marmoset (*Callithrix pygmaea)*, for which

both absolute and relative volume have decreased. This finding is unsurprising due to the

expected effects of dwarfism in this species and the limitation this exerts on brain size [75]. In

the case of the lar gibbon (*Hylobates lar)* the analysis revealed an increase in both volumes

from the ancestral state, possibly reflecting the complexity of its habitats and the subsequent

expansion of both hippocampus volume and brain volume.

<<Figure 2 about here>>

(b) PGLS

(i) Ecological model: Home range

We tested the prediction that variation in hippocampal and regional volumes can be predicted

by variation in home range size. Only subiculum and CA1 were related to home range, after

controlling for brain volume (subiculum: $\lambda=0.57$, $F = 8.12$, p=0.007 and CA1: $\lambda=0.80$, $F =$

9.02, p=0.005 on 2 and 36 degrees of freedom). See Table 2 and Supplementary for detailed

results of all analysis.

(ii) Ecological model: Diet

We tested whether diet, that is, increase in frugivory, indicated as fraction fruit, can predict

variation in hippocampal or regional volumes. We found no support for such hypothesis.

(iii) Social model

We tested whether variation in social group size can be predictive of variation in hippocampal

or regional volumes and found support for such relation using 4 different measures of group

size (see Methods - Social and ecological data). All measures of group size could explain

volumetric variation in hippocampus and most subregions, except hilus (0 out of 4 measures),

HP+HS+fibers (1 out of 4), and subiculum (3 out of 4). Interestingly, CA2 volume showed no

phylogenetic signal in any ecological or social model, except for when tested against social

group size and female group size (see Table 3 for phylogenetic signal of the residuals in every

model).

<<Tables 2 and 3 about here>>

<<Figure 3 here>>

(c) Model dredging

As we couldn't test the 'full model' against hippocampus volume and its subcomponents, we

dredged the 'full model' against the anatomical measurements and identified the best model

ranked by AICc. Subsequently, all models within 5 AICc points of the best models were

tested with PGLS and the respective best model (diet, home range or social) was chosen based

on weight. In case where multiple social variables were included in the best model, only the

most significant one in a PGLS regression was chosen. (See Table 4)

<<Table 4 here>>

(d) Neocortex

Following up on the observation that 1) both hippocampus and all its subcomponents were

positively related to brain volume, 2) many interactions between predictors and brain volume

were yielding negative slopes and 3) with increase in absolute hippocampal volume in some

species there was nonetheless a decrease in the relative hippocampal volume, we investigated

whether that relationship is driven by variation in neocortex volume as it comprises

significant proportion of the total brain volume. We found that hippocampal volume is

strongly negatively related to neocortex volume ($\lambda = 0$, slope = -3.81, t=-8.11, p<0.0001, 3,

40 df) even after accounting for brain volume (see Supplementary results).

(e) Sexual size dimorphism

We further explored the relationship between somatic and brain structure sexual dimorphism

in a separate dataset of 12 primate species. Somatic sexual dimorphism was best predicted by

hippocampus volume dimorphism, ($\lambda = 0$, slope = 1.87, Std. error = 0.35, t = 5.19, p=0.0004

on 1 and 10 df). Even though dimorphism in mesencephalon ($\lambda = 0$, slope = 1.35, Std. error =

0.45, t = 2.96, p=0.014 on 1 and 10 df) and lobus piriformis ($\lambda = 0.78$, slope = 0.67, Std. error

= 0.23, t = 2.83, p=0.018 on 1 and 10 df) were also significant predictors of somatic sexual

dimorphism, these relationships didn't stand after correction for multiple comparisons. The

new level of α for this batch of analysis was fixed to 0.0045 (dividing 0.05 by 11 structures)

and was sufficed by hippocampus volume alone. None of the other structure volumes (OBL,

CER, DIE, BOL, SCH, SEP, STR, NEO) showed a relationship with somatic sexual

dimorphism.

3. Discussion

We find that in primates, hippocampal volume and most of its subcomponents can be reliably

predicted by different measures of group size and home range to a certain extent, but not diet.

Moreover, we suggest that as brains get larger, the neocortex may take on functions shared

with the hippocampus and thus hippocampus size relative to the rest of the brain gets smaller.

Alternatively, the size of the hippocampus might be under strong developmental constraint.

Hippocampal structures crucial to spatial memory, CA1 and subiculum, evolve in line with

ecological (spatial) and social demands. CA2, CA3 and fascia dentata were shown to evolve

in line only with social demands, unlike the hilus, for which volume could not be predicted by

any of our models. No relationship between hippocampal volume and any of its

subcomponents was detected with increased fruit consumption in the primate's diet.

First, neocorticalization outpaces the enlargement of the hippocampus, as indicated in the

ancestral state estimation and the subsequent follow-up analysis. This is likely due to a

reallocation of functions such as memory, spatial cognition, and inhibition from the

hippocampus to the neocortex. With neocorticalization, parallel systems are thought to have

emerged, leading to an increased neocortex ratio [76] and allocation of functions to the

neocortex [77]. Whereas in smaller brained species the hippocampus is of utmost importance

in many cognitive abilities, as the neocortex expands there may be a greater proportion of

these functions allocated to it, or the neocortex might be taking up on an array of new social

functions that do not exist in smaller brained species. The neocortex, like the hippocampus,

provides mappings used in information acquisition, retention and use. Compared to rodents,

in highly neocorticalized humans, the hippocampus may not have as prominent a role in

spatial cognition [78].

On the flip side, in primates, the neocortex may also have an increased role in spatial

processing. Parietal association areas of the neocortex are also crucial to spatial perception

and may provide navigational information and are the focus of spatial cognition studies [79].

The interplay between the parietal and hippocampal neural networks remains poorly

understood [80] although it has been suggested that both are involved in spatial navigation.

Parietal representations provide an egocentric frame of reference and may map movements

along a route according to route-centred positional information [81].

Second, of the hippocampal regions, both CA1 and CA3 residuals show phylogenetic signal

and coevolve with home range (CA1) and group size (CA1 and CA3). This is consistent with

the notion that the hippocampus is involved in both social and ecological behaviour [41, 44].

Compared to other brain component volumes, hippocampus volume was found to be the best

predictor for cognitive tasks measuring executive function in primates [82]. This is the first

study linking theses specific hippocampal substructures to both social and ecological factors

across primates. This is in line with work in other taxa linking species-specific requirements

for spatial memory and hippocampus volume [26], but the implication - which would benefit

from future study - is that in primates the role of the hippocampus may be even more

influenced by social factors.

We found no relationship between the percentage fruit in diet and the size of the hippocampus

or any of its subcomponents. While fruit acquisition may play an important role in

intelligence [10, 83] and brain size [6], our findings suggest that the primary contribution of

diet to these features may be the generalised support of the brain's high metabolic costs [9]

rather than specifically influencing neural systems specialised for spatial ability. On the other

hand, non-dietary social-spatial memory factors, such as the ability to code for the locations

of conspecifics, may be linked to hippocampus size.

Third, CA2 volume residuals showed no phylogenetic signal, except for in a regression with

social and female group sizes. Thus, CA2 seems not to be under phylogenetic constraint

related to home range or diet but is only shaped by social pressures. This finding can be

interpreted as an indicator of the relative functional decoupling of this zone to the rest of the

hippocampus. CA2 may show species-specific adaptations related to behavioural niche which

deviate from trends within a clade. Recent work on the function of CA2 in mice found that it

has a special role in social memory [63] and it has a different gene expression profile from

CA1 and CA3 [44]. On the other hand, the adaptability of CA2 might come at a cost in terms

of maintaining elementary functions shared across species - unlike CA1 and CA3 it is a

smaller region and is not involved in spatial tasks [63]. Additionally, hilus was one of the

structures that showed no relationship to social group size. It is important in spatial and

memory functions and may be less adaptable to changes in social structure.

We further investigated how hippocampus size is related to sexual dimorphism in primates

since sex differences in hippocampal anatomy, spatial cognition, and home range size seem to

be linked in some taxa [84]. We found that, of all brain structures examined, sexual

dimorphism in the hippocampus is most closely related to somatic sexual dimorphism. It

should be considered that spatial functions, like other brain functions, have become more

corticalised in taxonomic groups with larger palliums such as primates [77]. However, the

nature of the link is debated, for example, male superiority in spatial cognition may be a by-

product of sex hormones rather than driven by ecological demands [85]. This provides a

preliminary attempt to understand sex differences in the primate hippocampus.

Overall, we show that group size can predict the size of most hippocampus regions, while diet

seems to be unrelated to hippocampal size at all. Moreover, group size was the only predictor

that was related to total hippocampal size. Social group size is thought to be related to an

increase in neocortex size, but this is mainly because of its role in higher cognitive social

processes that are more demanding than simply remembering other individuals [86] [76].

Social memories seem to be structured within the spatial framework of the hippocampus too

[23]. In fact, social memory might in part be an exaptation that "reuses" neural circuitry of the

hippocampus for spatial maps in an ancestral mammal [87, 88]. In line with this, the role of

hippocampus in spatial cognition is pronounced in rodents, but less well understood in

primates; in humans, it is argued that the hippocampus appears to function in memory rather

than spatial cognition [78]. Given the importance of social skills in primates, it is possible that

in this order, social memory (overlain onto spatial maps originally for navigation) has

increased in dominance over spatial mapping. The importance of the increasing evidence that

social and spatial cognition rely on the same underlying representations in humans, such that

spatial maps provide a means for mapping social relations, is developing into applications

ranging from design considerations in the built environment to clinical implications [45].

We thank Anna van Oosterzee for stimulating discussions, Heiko Frahm for providing data

and Simone Blomberg for statistical advice.

References

1.   van Woerden J.T., Willems E.P., van Schaik C.P., Isler K. 2012 Large brains buffer
energetic effects of seasonal habitats in catarrhine primates. *Evolution* **66**, 191-199.
(doi:10.1111/j.1558-5646.2011.01434.x).
2.   Weisbecker V., Blomberg S.P., Goldizen A.W., Brown M., Fisher D.O. 2015 The
evolution of relative brain size in marsupials is energetically constrained but not driven
by behavioural complexity. *Brain, behavior and evolution.* (doi:10.1159/000377666).
3.   Healy S.D., Rowe C. 2007 A critique of comparative studies of brain size.
Proceedings of the Royal Society, London Series B **274**, 453-464.
(doi:10.1098/rspb.2006.3748).
4.   Whiten A., van de Waal E. 2017 Social learning, culture and the 'socio-cultural
brain' of human and non-human primates. *Neurosci Biobehav Rev* **82**, 58-75.
(doi:10.1016/j.neubiorev.2016.12.018).
5.   Dunbar R.I.M., Shultz S. 2017 Why are there so many explanations for primate
brain evolution? *Philos Trans R Soc Lond B Biol Sci* **372**(1727).
(doi:10.1098/rstb.2016.0244).
6.   DeCasien A.R., Williams S.A., Higham J.P. 2017 Primate brain size is predicted by
diet but not sociality. *Nat Ecol Evol* **1**(5), 112. (doi:10.1038/s41559-017-0112).
7.   Powell L.E., Isler K., Barton R.A. 2017 Re-evaluating the link between brain size
and behavioural ecology in primates. *Proc Biol Sci* **284**(1865).
(doi:10.1098/rspb.2017.1765).
8.   Herculano-Houzel S. 2011 Scaling of brain metabolism with a fixed energy budget
per neuron: implications for neuronal activity, plasticity and evolution. *PLoS One* **6**(3),
e17514. (doi:10.1371/journal.pone.0017514).
9.   Fonseca-Azevedo K., Herculano-Houzel S. 2012 Metabolic constraint imposes
tradeoff between body size and number of brain neurons in human evolution. *Proc Natl
Acad Sci U S A* **109**(45), 18571-18576. (doi:10.1073/pnas.1206390109).
10.   Milton K. 1981 Distribution Patterns of Tropical Plant Foods as an Evolutionary
Stimulus to Primate Mental Development. *Am Anthropol* **83**(3), 534-548.
(doi:10.1525/Aa.1981.83.3.02a00020
10.2307/676751).
11.   Milton K. 1988 Foraging behaviour and the evolution of primate intelligence. In
*Machiavellian intelligence: Social expertise and the evolution of intellect in monkeys, apes,
and humans* (eds. Byrne R.W., Whiten A.), pp. 285-305. Oxford, Clarendon Press.
12.   Clutton-Brock T.H., Harvey P.H. 1980 Primates, Brains and Ecology. *Journal of
Zoology* **190**(MAR), 309-323.
13.   Mace G.M., Harvey P.H., Cluttonbrock T.H. 1981 Brain size and ecology in small
mammals. *Journal of Zoology* **193**(Mar), 333-354. (doi:10.1111/j.1469-
7998.1981.tb03449.x).
14.   Weisbecker V., Goswami A. 2011 Neonatal maturity as the key to understanding
brain size evolution in homeothermic vertebrates. *BioEssays* **33**(3), 155-158.
15.   De Winter W., Oxnard C.E. 2001 Evolutionary radiations and convergences in the
structural organization of mammalian brains. *Nature* **409**(6821), 710-714.
16.   Gomez-Robles A., Hopkins W.D., Sherwood C.C. 2014 Modular structure facilitates
mosaic evolution of the brain in chimpanzees and humans. *Nature communications* **5**,
4469. (doi:10.1038/ncomms5469).
17.   Barton R.A., Harvey P.H. 2000 Mosaic evolution of brain structure in mammals.
Nature **405**(6790), 1055-1058. (doi:10.1038/35016580).

18.   Carlisle A., Selwood L., Hinds L.A., Saunders N., Habgood M., Mardon K.,
Weisbecker V. 2017 Testing hypotheses of developmental constraints on mammalian
brain partition evolution, using marsupials. *Sci Rep* **7**(1), 4241. (doi:10.1038/s41598-
017-02726-9).
19.   O'Keefe J. 1976 Place units in the hippocampus of the freely moving rat. *Exp*
Neurol **51**(1), 78-109. (doi:10.1016/0014-4886(76)90055-8).
20.   O'Keefe J., Dostrovsky J. 1971 The hippocampus as a spatial map. Preliminary
evidence from unit activity in the freely-moving rat. *Brain Res* **34**(1), 171-175.
(doi:10.1016/0006-8993(71)90358-1).
21.   Hafting T., Fyhn M., Molden S., Moser M.B., Moser E.I. 2005 Microstructure of a
spatial map in the entorhinal cortex. *Nature* **436**(7052), 801-806.
(doi:10.1038/nature03721).
22.   Scoville W.B., Milner B. 1957 Loss of recent memory after bilateral hippocampal
lesions. *J Neurol Neurosurg Psychiatry* **20**(1), 11-21. (doi:10.1136/jnnp.20.1.11).
23.   Jeffery K.J. 2018 The Hippocampus: From Memory, to Map, to Memory Map.
Trends Neurosci **41**(2), 64-66. (doi:10.1016/j.tins.2017.12.004).
24.   Taylor A.M., Bus T., Sprengel R., Seeburg P.H., Rawlins J.N., Bannerman D.M. 2014
Hippocampal NMDA receptors are important for behavioural inhibition but not for
encoding associative spatial memories. *Philos Trans R Soc Lond B Biol Sci* **369**(1633),
20130149. (doi:10.1098/rstb.2013.0149).
25.   Kaut K.P., Bunsey M.D., Riccio D.C. 2003 Olfactory learning and memory
impairments following lesions to the hippocampus and perirhinal-entorhinal cortex.
Behav Neurosci **117**(2), 304-319. (doi:10.1037/0735-7044.117.2.304).
26.   Jacobs L.F., Spencer W.D. 1994 Natural space-use patterns and hippocampal size
in kangaroo rats. *Brain Behav Evol* **44**(3), 125-132. (doi:10.1159/000113584).
27.   Rodriguez F., Lopez J.C., Vargas J.P., Broglio C., Gomez Y., Salas C. 2002 Spatial
memory and hippocampal pallium through vertebrate evolution: insights from reptiles
and teleost fish. *Brain Res Bull* **57**(3-4), 499-503.
28.   Garamszegi L.Z., Eens M. 2004 The evolution of hippocampus volume and brain
size in relation to food hoarding in birds. *Ecology letters* **7**(12), 1216-1224.
(doi:10.1111/j.1461-0248.2004.00685.x).
29.   Lucas J.R., Brodin A., de Kort S.R., Clayton N.S. 2004 Does hippocampal size
correlate with the degree of caching specialization? *Proc Biol Sci* **271**(1556), 2423-2429.
(doi:10.1098/rspb.2004.2912).
30.   Maguire E.A., Gadian D.G., Johnsrude I.S., Good C.D., Ashburner J., Frackowiak R.S.,
Frith C.D. 2000 Navigation-related structural change in the hippocampi of taxi drivers.
Proc Natl Acad Sci U S A **97**(8), 4398-4403. (doi:10.1073/pnas.070039597
070039597 [pii]).
31.   Clayton N.S., Krebs J.R. 1994 Hippocampal growth and attrition in birds affected
by experience. *Proc Natl Acad Sci U S A* **91**(16), 7410-7414.
32.   Smulders T.V., Shiflett M.W., Sperling A.J., DeVoogd T.J. 2000 Seasonal changes in
neuron numbers in the hippocampal formation of a food-hoarding bird: the black-
capped chickadee. *J Neurobiol* **44**(4), 414-422.
33.   Barton R.A., Purvis A. 1994 Primate brains and ecology: looking beneath the
surface. In *Current Primatology* (eds. Thierry B., Roeder A., Herrenschmid J.), pp. 1-9.
Strasbourg, Université Louis Pasteu.
34.   Edler M. 2007 A comparative analysis of hippocampus size and ecological factors
in primates, Kent State University.

35.   Jacobs L.F., Gaulin S.J., Sherry D.F., Hoffman G.E. 1990 Evolution of spatial
cognition: sex-specific patterns of spatial behavior predict hippocampal size. *Proc Natl*
Acad Sci U S A **87**(16), 6349-6352. (doi:10.1073/pnas.87.16.6349).
36.   Langley C.M. 1994 Spatial memory in the desert kangaroo rat (Dipodomys
deserti). *J Comp Psychol* **108**(1), 3-14.
37.   Galea L.A., Kavaliers M., Ossenkopp K.P. 1996 Sexually dimorphic spatial learning
in meadow voles Microtus pennsylvanicus and deer mice Peromyscus maniculatus. *J Exp*
Biol **199**(Pt 1), 195-200.
38.   Perdue B.M., Snyder R.J., Zhihe Z., Marr M.J., Maple T.L. 2011 Sex differences in
spatial ability: a test of the range size hypothesis in the order Carnivora. *Biology Letters*
7(3), 380-383. (doi:10.1098/rsbl.2010.1116).
39.   Sherry D.F. 2006 Neuroecology. *Annu Rev Psychol* **57**(1), 167-197.
(doi:10.1146/annurev.psych.56.091103.070324).
40.   Guigueno M.F., Snow D.A., Macdougall-Shackleton S.A., Sherry D.F. 2014 Female
cowbirds have more accurate spatial memory than males. *Biol Lett* **10**(2), 20140026.
(doi:10.1098/rsbl.2014.0026).
41.   Omer D.B., Maimon S.R., Las L., Ulanovsky N. 2018 Social place-cells in the bat
hippocampus. *Science* **359**(6372), 218-224. (doi:10.1126/science.aao3474).
42.   Irle E., Ruhleder M., Lange C., Seidler-Brandler U., Salzer S., Dechent P., Weniger
G., Leibing E., Leichsenring F. 2010 Reduced amygdalar and hippocampal size in adults
with generalized social phobia. *J Psychiatry Neurosci* **35**(2), 126-131.
43.   Rubin R.D., Watson P.D., Duff M.C., Cohen N.J. 2014 The role of the hippocampus
in flexible cognition and social behavior. *Front Hum Neurosci* **8**.
44.   Alexander G.M., Farris S., Pirone J.R., Zheng C., Colgin L.L., Dudek S.M. 2016 Social
and novel contexts modify hippocampal CA2 representations of space. *Nature*
communications **7**, 10300. (doi:10.1038/ncomms10300).
45.   Proulx M.J., Todorov O.S., Taylor Aiken A., de Sousa A.A. 2016 Where am I? Who
am I? The relation between spatial cognition, social cognition and individual differences
in the built environment. *Front Psychol* **7**, 64. (doi:10.3389/fpsyg.2016.00064).
46.   Witter M.P., Naber P.A., van Haeften T., Machielsen W.C.M., Rombouts S.A.R.B.,
Barkhof F., Scheltens P., Lopes da Silva F.H. 2000 Cortico-hippocampal communication
by way of parallel parahippocampal-subicular pathways. *Hippocampus* **10**(4), 398-410.
(doi:10.1002/1098-1063(2000)10:4<398::Aid-hipo6>3.0.Co;2-k).
47.   Hsu D. 2007 The dentate gyrus as a filter or gate: a look back and a look ahead.
Prog Brain Res **163**, 601-613. (doi:10.1016/s0079-6123(07)63032-5).
48.   de la Rosa-Prieto C., Ubeda-Banon I., Mohedano-Moriano A., Pro-Sistiaga P., Saiz-
Sanchez D., Insausti R., Martinez-Marcos A. 2009 Subicular and CA1 hippocampal
projections to the accessory olfactory bulb. *Hippocampus* **19**(2), 124-129.
(doi:10.1002/hipo.20495).
49.   Kesner R.P., Lee I., Gilbert P. 2004 A behavioral assessment of hippocampal
function based on a subregional analysis. *Rev Neurosci* **15**(5), 333-351.
50.   Amaral D.G. 1978 A golgi study of cell types in the hilar region of the
hippocampus in the rat. *The Journal of Comparative Neurology* **182**(5), 851-914.
(doi:10.1002/cne.901820508).
51.   Andrews-Zwilling Y., Gillespie A.K., Kravitz A.V., Nelson A.B., Devidze N., Lo I.,
Yoon S.Y., Bien-Ly N., Ring K., Zwilling D., et al. 2012 Hilar GABAergic interneuron
activity controls spatial learning and memory retrieval. *PLoS One* **7**(7), e40555.
(doi:10.1371/journal.pone.0040555).

52.   Lorincz A., Buzsaki G. 2000 Two-phase computational model training long-term
memories in the entorhinal-hippocampal region. *Ann N Y Acad Sci* **911**, 83-111.
53.   Myers C.E., Scharfman H.E. 2009 A role for hilar cells in pattern separation in the
dentate gyrus: a computational approach. *Hippocampus* **19**(4), 321-337.
(doi:10.1002/hipo.20516).
54.   Witter M.P. 2007 Intrinsic and extrinsic wiring of CA3: indications for
connectional heterogeneity. *Learn Mem* **14**(11), 705-713. (doi:10.1101/lm.725207).
55.   Kesner R.P. 2007 Behavioral functions of the CA3 subregion of the hippocampus.
Learn Mem **14**(11), 771-781. (doi:10.1101/lm.688207).
56.   Tsien J.Z., Huerta P.T., Tonegawa S. 1996 The essential role of hippocampal CA1
NMDA receptor-dependent synaptic plasticity in spatial memory. *Cell* **87**(7), 1327-1338.
(doi:10.1016/S0092-8674(00)81827-9).
57.   Moser E.I., Moser M.B. 2009 Hippocampus and Neural Representations. In
Encyclopedia of Neuroscience (pp. 1129-1136.
58.   O'Mara S. 2005 The subiculum: what it does, what it might do, and what
neuroanatomy has yet to tell us. *J Anat* **207**(3), 271-282. (doi:10.1111/j.1469-
7580.2005.00446.x).
59.   Amaral D.G., Dolorfo C., Alvarez-Royo P. 1991 Organization of CA1 projections to
the subiculum: a PHA-L analysis in the rat. *Hippocampus* **1**(4), 415-435.
(doi:10.1002/hipo.450010410).
60.   Sharp P.E., Green C. 1994 Spatial correlates of firing patterns of single cells in the
subiculum of the freely moving rat. *J Neurosci* **14**(4), 2339-2356.
61.   Chevaleyre V., Piskorowski R.A. 2016 Hippocampal Area CA2: An Overlooked but
Promising Therapeutic Target. *Trends Mol Med* **22**(8), 645-655.
(doi:10.1016/j.molmed.2016.06.007).
62.   Cui Z., Gerfen C.R., Young W.S., 3rd. 2013 Hypothalamic and other connections
with dorsal CA2 area of the mouse hippocampus. *J Comp Neurol* **521**(8), 1844-1866.
(doi:10.1002/cne.23263).
63.   Hitti F.L., Siegelbaum S.A. 2014 The hippocampal CA2 region is essential for
social memory. *Nature*. (doi:10.1038/nature13028).
64.   Benes F.M., Kwok E.W., Vincent S.L., Todtenkopf M.S. 1998 A reduction of
nonpyramidal cells in sector CA2 of schizophrenics and manic depressives. *Biol*
Psychiatry **44**(2), 88-97. (doi:10.1016/s0006-3223(98)00138-3).
65.   Badowska Szalewska E., Ludkiewicz B., Krawczyk R., Melka N., Moryś J. 2017
Comparison of the influence of two models of mild stress on hippocampal brain derived
neurotrophin factor (BDNF) immunoreactivity in old age rats. *Acta Neurobiol Exp (Wars)*
77(1), 68-76. (doi:10.21307/ane-2017-037).
66.   Frahm H.D., Zilles K. 1994 Volumetric comparison of hippocampal regions in 44
primate species. *J Hirnforsch* **35**(3), 343-354.
67.   Stephan H., Frahm H., Baron G. 1981 New and revised data on volumes of brain
structures in insectivores and primates. *Folia Primatol (Basel)* **35**(1), 1-29.
(doi:10.1159/000155963).
68.   Gautier-Hion A., Gautier J.P. 1976 [Growth, sexual maturity, social maturity and
reproduction in African Forest Cercopithecines (author's transl)]. *Folia Primatol (Basel)*
26(3), 165-184. (doi:10.1159/000155749).
69.   Dunbar R.I.M., Mac Carron P., Shultz S. 2018 Primate social group sizes exhibit a
regular scaling pattern with natural attractors. *Biol Lett* **14**(1).
(doi:10.1098/rsbl.2017.0490).

70.   Arnold C., Matthews L.J., Nunn C.L. 2010 The 10ktrees website: A new online
resource for primate phylogeny. *Evolutionary Anthropology: Issues, News, and Reviews*
19(3), 114-118. (doi:10.1002/evan.20251).
71.   R Core Team. 2018 R: A language and environment for statistical computing. . In
R Foundation for Statistical Computing (Vienna, Austria.
72.   Revell L.J. 2012 phytools: an R package for phylogenetic comparative biology
(and other things). *Methods in Ecology and Evolution* **3**(2), 217-223.
(doi:10.1111/j.2041-210X.2011.00169.x).
73.   Orme D., Freckleton R.P., Thomas G.H., Petzoldt T., Fritz S.A., Isaac N. 2013 *CAPER:*
*comparative analyses of phylogenetics and evolution in R*145-151 p.
74.   Burnham K., Anderson D. 2004 Model selection and multi-model inference.
Second NY: Springer-Verlag **63**.
75.   Montgomery S.H., Mundy N.I. 2013 Parallel episodes of phyletic dwarfism in
callitrichid and cheirogaleid primates. *J Evol Biol*, n/a-n/a. (doi:10.1111/jeb.12097).
76.   Dunbar R.I.M. 1992 Neocortex size as a constraint on group-size in primates. *J*
Hum Evol **22**(6), 469-493. (doi:10.1016/0047-2484(92)90081-J).
77.   Boraud T., Leblois A., Rougier N.P. 2018 A natural history of skills. *Prog Neurobiol*.
(doi:10.1016/j.pneurobio.2018.08.003).
78.   Kim S., Dede A.J., Hopkins R.O., Squire L.R. 2015 Memory, scene construction, and
the human hippocampus. *Proc Natl Acad Sci U S A* **112**(15), 4767-4772.
(doi:10.1073/pnas.1503863112).
79.   Arbib M.A. 1997 From visual affordances in monkey parietal cortex to
hippocampo-parietal interactions underlying rat navigation. *Philos Trans R Soc Lond B*
Biol Sci **352**(1360), 1429-1436. (doi:10.1098/rstb.1997.0129).
80.   Whitlock J.R., Sutherland R.J., Witter M.P., Moser M.B., Moser E.I. 2008 Navigating
from hippocampus to parietal cortex. *Proc Natl Acad Sci U S A* **105**(39), 14755-14762.
(doi:10.1073/pnas.0804216105).
81.   Nitz D. 2009 Parietal cortex, navigation, and the construction of arbitrary
reference frames for spatial information. *Neurobiol Learn Mem* **91**(2), 179-185.
(doi:10.1016/j.nlm.2008.08.007).
82.   Shultz S., Dunbar R.I.M. 2010 Species differences in executive function correlate
with hippocampus volume and neocortex ratio across nonhuman primates. *J Comp Psych*
124(3), 252-260. (doi:10.1037/a0018894).
83.   Milton K., May M.L. 1976 Body weight, diet and home range area in primates.
Nature **259**(5543), 459-462. (doi:10.1038/259459a0).
84.   Jones C.M., Braithwaite V.A., Healy S.D. 2003 The evolution of sex differences in
spatial ability. *Behav Neurosci* **117**(3), 403-411. (doi:10.1037/0735-7044.117.3.403).
85.   Clint E.K., Sober E., Garland T., Jr., Rhodes J.S. 2012 Male superiority in spatial
navigation: adaptation or side effect? *Q Rev Biol* **87**(4), 289-313. (doi:10.1086/668168).
86.   Lewis P.A., Birch A., Hall A., Dunbar R.I.M. 2017 Higher order intentionality tasks
are cognitively more demanding. *Soc Cogn Affect Neurosci* **12**(7), 1063-1071.
(doi:10.1093/scan/nsx034).
87.   Anderson M.L. 2010 Neural reuse: a fundamental organizational principle of the
brain. *Behav Brain Sci* **33**(4), 245-266; discussion 266-313.
(doi:10.1017/s0140525x10000853).
88.   Dehaene S., Cohen L. 2007 Cultural recycling of cortical maps. *Neuron* **56**(2), 384-
398. (doi:10.1016/j.neuron.2007.10.004).
650

652 **Appendices**

655 **Table 1. Hippocampal regions investigated**

Hippocampus total (=Hippocampal formation; HIP)		
Retrocommissural hippocampus (= Retrohippocampus; HR)		
Hippocampus proper		
Cornu ammonis		
CA1		
CA2		
CA3		
CA4	"Hilus region" or "Hilus"	Dentate
"deep stratum multiforme of the dentate gyrus"		gyrus
Fascia dentata (FD)		
Prosubiculum	"Subiculum" (SUB)	
Subiculum		
"HP+HS+fibers" or "fibers"		
Hippocampus praecommissuralis (HP)		
Hippocampus supracommissuralis (HS)		
fimbria/fornix complex		

Note. The subicular complex includes pro-, pre-, parasubiculum and subiculum. The region called "subiculum (SUB)" in Frahm and Zilles (1994) includes both the prosubiculum and subiculum. The pre- and parasubiculum were not included in Frahm and Zilles (1994) as part of the hippocampus volumes and were not examined here. In Stephan et al 1981, the pre- and parasubiculum were included along with entorhinum and perirhinum as the Schizocortex.

656 **Table 2.** Sequential SS for PGLS results of home range, fraction fruit, group size (Powell) (S1), social group size (S2),
657 group size (Dunbar) (S3), and female group size (S4) controlling for total brain volume. DF - degrees of freedom for
658 each model. For list of abbreviations, see Methods – Anatomical data and Table 1.
659 Only predictors with p<0.0083 are filled in grey. See supplement for full test results, including F, t statistics and exact
660 p values.

	HIP	HP+HS+ fibers	RH	Subiculum	Hilus	CA1	CA2	CA3	FD	DF
Total brain	■	■	■	■	■	■	■	■	■	
Home Range				■		■				36
Fraction fruit										33
S1	■		■			■	■	■	■	37
S2	■		■	■		■	■	■	■	36
S3	■		■	■		■	■	■	■	39
S4	■		■	■		■	■	■	■	26

Table 3. Phylogenetic signal (λ) of the residuals in the regression models for each hippocampal component tested from Table 2. All models include total brain size.

	HIP	HP+HS+ fibers	RH	Subiculum	Hilus	CA1	CA2	CA3	FD
λ Home Range	.67	.23	.77	.57	.27	.80	0	.74	.79
λ Fraction Fruit	.80	.06	.99	.80	.64	.98	0	.86	.81
λ S1	.60	.21	.64	.53	.49	.62	0	.72	.50
λ S2	.64	.23	.66	.42	.52	.62	.84	.78	.55
λ S3	.68	.34	.73	.59	.63	.71	.28	.85	.51
λ S4	.67	.12	.8	.68	.73	.75	0	.86	.59

Table 4. Model dredging

Dredging the full model against hippocampal volume and its subcomponents. Shown are the predictors, weight of each best model against all other possible combinations (to the sum of 1), and best model type. Asterisk marks a

672 group size measure that was chosen from a model including more than one group size variable, based on significance

673 in PGLS.

Region	Predictors in model	Weight	Best model type
Hippocampus	Total Brain + Group size (Dunbar)	0.99	Social
HP+HS+ fibers	Total Brain + Group size (Dunbar)*	0.32	Social
HR	Total Brain + Group size (Dunbar)	0.195	Social
Subiculum	Total Brain + Female group size + Home Range	0.158	Social + Home Range
Hilus	Total Brain + Female group size + Home Range	0.552	Social + Home Range
CA1	Total Brain + Group size (Dunbar) + Home Range	0.325	Social + Home Range
CA2	Total Brain + Social group size	0.26	Social
CA3	Total Brain + Social group size	0.147	Social
FD	Total Brain + Female group size	0.35	Social

677   **Figures Captions**

Figure 1. Hippocampus of *Miopithecus talapoin*. Hil: Hilus, FD: Fascia dentata, S:

Subiculum, 1: CA1, 2: CA2, 3: CA3. From Frahm & Zilles, 1994.

Figure 2. Ancestral trait estimations of absolute hippocampal volume (left), and the residuals

from the phylogenetic regression with total brain size (right). We observe that most species

that had increase in absolute hippocampal volume had a reciprocal decrease in hippocampal

volume relative to the whole brain.

Figure 3. Plot of regression of total brain volume (black solid line and black circles), fraction

fruit (red small-dashed line and red squares) and home range (blue long-dashed line and blue

triangles), and group size (green dotted line and green pluses) against hippocampal volume.

Total hippocampus volume

Relative hippocampus volume

Mya

trait value

decrease   increase