Social networks with strong spatial embedding generate non-standard epidemic dynamics driven by higher-order clustering

David J. Haw[1a], Rachael Pung[1a], Jonathan M. Read[2], Steven Riley[1*]

1. MRC Centre for Outbreak Analysis and Modelling, Department of Infectious Disease Epidemiology, School of Public Health, Imperial College London, London, UK

2. Centre for Health Informatics Computing and Statistics, Lancaster Medical School, Lancaster University, Lancaster, UK

[a] [*]Corresponding author. Email address s.riley@imperial.ac.uk

Abstract

Some directly transmitted human pathogens such as influenza and measles generate sustained exponential growth in incidence, and have a peak incidence that is consistent with age-group-specific depletion of susceptible individuals. Many do not. While a prolonged exponential phase typically arises in traditional disease-dynamic models, quantitative descriptions of non-exponential growth are either highly abstract, phenomenological or ignore well-described social structuring. Here, we create large socio-spatial human contact networks using population density data, a previously developed fitting algorithm, and gravity-like mobility kernels. We define the basic reproductive number for this system to be analogous to the traditional compartmental model definition. We then explore networks with a household-workplace structure in which between-household contacts can be formed with varying degrees of spatial correlation, determined by a single parameter from the gravity-like kernel. By varying this single parameter and simulating epidemic spread, we are able to identify the strength of spatial correlation required to induce sub-exponential outbreak dynamics with lower, later peaks. We investigate the topological properties of our networks via a generalized clustering coefficient that extends beyond immediate neighbourhoods, identifying topological properties of these networks that correlate with strictly sub-exponential spread. Our results motivate the joint observation of incidence and socio-spatial human behaviour during epidemics that exhibit non-standard incidence patterns.

Author Summary

Epidemics are typically described using a standard set of mathematical models that do not capture social interactions or the way those interactions are determined by geography. Here we propose a model that can reflect social networks influenced strongly by the way people travel and we show that they lead to very different epidemic profiles. This type of model will likely be useful for forecasting.

Introduction

Our understanding of key features of epidemic incidence curves is underpinned by standard epidemic dynamics [1, 18]. Numerical solutions to multi-type compartmental models can explain: the initial growth phase [27], the timing and amplitude of the peak [40], epidemic duration [20] and the total number of cases [16]. These models can capture many different types of complexity, such as age-specific susceptibility and transmissibility [15], behavioural risk groups [4] and, with increasing frequency, geographical location [31]. By definition, standard compartmental models have well defined periods of exponential growth [17] and a total cumulative incidence driven by the mix of infectious types [8].

Infectious disease transmission can also be represented by stochastic processes and networks. Directly transmitted pathogens propagate through susceptible human populations and create drected infection trees with an offspring-like process [14]. Each node may be a different type (e.g. children may be more infectious than adults [39]) and individuals with many contacts may be more infectious than those with fewer contacts [23]. Although difficult to observe, infection trees describe a real biological process: these pathogens cannot survive outside of a human host for more than a few hours, so the founding pathogen population for an infectee comes directly from their infector. Further, we can conceptualise that infection trees occur when a true offspring process is constrained to pass through a social network [3, 37], with infection occurring according to a specified probability when an edge exists between a susceptible and an infectious individual. For a limited set of network models, we can calculate key properties of their resulting epidemics, such as the probability of epidemic establishment and cumulative incidence [25]. Higher order network moments, beyond the degree distribution, likely have a role in shaping the outcome of epidemics [21].

The basic reproductive number has been defined for both compartmental models and for network models. For compartmental models, the reproduction number is defined as the average number of new infections generated by a typically infectious individual in an otherwise infectious population [17]. The word "typically" is somewhat overloaded in this definition: during the exponential phase, the system will find the optimal balance of infectious types to maximize the growth of incidence. More infectious individuals will be over-represented in the mix during the exponential phase [36]. The word "typically" reflects this eigenstate of the renewal process. Conversely, for network models, the basic reproductive can be defined as expected number of infections in the second generation, i.e. the product of the average degree and the probability of transmission per link per generation [31]. Here, we suggest that network models also undergo exponential phases of incidence growth and we therefore propose a definition of the basic reproductive number that is consistent with the definition for compartmental models. We use R^* to denote the expected number of infections in the second generation and R_0 for the definition of the basic reproductive number that is consistent with that used for compartmental models.

Both compartmental and network models can be embedded in space [31]. Each node can have a location in space while each compartment can refer to a single unit of space. Node density can be assigned according to known population densities and compartments can be assigned equal spatially areas but different numbers of hosts. In general, the risk of infection passing between two people decreases as the distance between their home location increases. We measure the propensity of nodes to form links across space or for infection to spread between compartments using mobility models borrowed from geography [11], such as the gravity and radiation models. Here, we are specifically interested in how the overall topology of a spatially-embedded network model can be driven by different movement assumptions and thus drive the gross features of the

epidemics that occur on the network.

Results

We used an existing variant of the Metropolis-Hastings algorithm [32] to create a spatially-embedded bipartite network of homes and workplaces consistent with the population density of Monrovia, Liberia, and with three illustrative movement scenarios (SI Appendix, Fig S1). An individual's propensity to choose a given workplace was determined by the distance between their home and workplace and parameters of a gravity-like kernel. The kernel was inversely proportional to distance raised to the power α, with movement scenarios generated solely by changing the value of α: a control value $\alpha = 0$ that removed the embedding and produced a *non-spatial* model; a *wide* kernel with $\alpha = 3$ typical of developed populations [32, 34]; and a *highly local* kernel with $\alpha = 6$ representing less developed populations (SI Appendix, Fig S1 part C compared with rural Huangshan in Ref [13]). The resulting distributions of distances from home to work were driven strongly by our choice of α, with 95% of journeys: less than 24.12km for $\alpha = 0$; less than 12.91km for $\alpha = 3$; and less than 6.68km for $\alpha = 6$. Workplace links were dissolved into links between individuals in different households resulting in a network of cliques (households) that were linked according to α.

The choice of movement kernel used to create the household-workplace networks affected gross features of simulated epidemics, even when controlling for other aspects of the network topology (Fig 1). Unipartite contact networks between households were obtained from the bipartite network of households and workplaces and were dependent on three parameters: mean household size h, mean number of workplace links v, and probability of forming a link in the workplace p_w. The mean workplace size w and mean degree of the network were determined by these parameters: $w = v/p_w + 1$, $\langle k \rangle = h - 1 + v$. Across a broad range of plausible values for h, v and p_w, very local movement ($\alpha = 6$) produced later epidemics than did typical developed-population movement ($\alpha = 3$) or spatially random mixing ($\alpha = 0$, Fig 1A). Similarly, time to extinction was later for very local movement ($\alpha = 6$) compared with more frequent longer-distance movement ($\alpha = 3$) or the absence of spatial embedding ($\alpha = 0$).

Each simulation is assigned a value of R^*, the average number of cases in the first generation per seed infection. For moderate-to-high values of the first generation ratio R^*, there was very little difference in the final size of the outbreak for the different movement assumptions. However, for low values of $R^* < 1.8$, the average final size of the outbreak was substantially smaller for more local kernels. This was driven by a higher probability of extinction when more local movement was assumed. The difference in final size driven by α was no longer present when we controlled for extinction (SI Appendix, Fig S2).

The choice of movement scenario had a substantial impact on peak incidence, even when R^* was high and there was little difference in the final sizes (Fig 1B, Fig 2 rows 1 and 2). For example, for parameters with first generation ratios in the range $[1.8, 2.2]$, average peak daily incidence as a fraction of the total population was 6.5×10^{-3} for random spatial movement, 5.4×10^{-3} for movement assumptions typical of developed populations and 3.0×10^{-3} when highly local movement was assumed. The relationship between peak height and first generation ratio appeared to be strongly linear, with correlation coefficients 0.9778, 0.9826 and 0.9806 for $\alpha = 0, 3$ and 6 respectively.

The relationship between peak incidence and final size for the three movement scenarios illustrates further how the topology of an embedded network could directly affect gross features of an epidemic. Peak incidence is observed prior to final size during an epidemic. For the same

peak height, local movement gave substantially larger final sizes. For peak daily incidences in the range $[3 \times 10^{-3}, 6.5 \times 10^{-3}]$ the final size of the outbreak was 68% when random spatial movement was assumed, 74% when movement was assumed to be typical of developed populations and 84% when highly-local movement was assumed.

For all movement scenarios, the basic reproductive number R_0 was smaller than the first generation ratio R^* and different from the expected number of secondary cases generated by a single seed in an otherwise susceptible population. The duration of the exponential phase can be seen when incidence is plotted on a log scale: a constant gradient of log incidence is evidence exponential growth (figure 2, third row). However, in a network model with clearly defined generations, the generation ratio can also be used to define exponential growth: if the ratio of incidence between generation $n + 1$ and n is the same as the ratio between generations n and $n - 1$, then we can claim to have identified a period of exponential growth (Methods, Fig 2). The value of that constant observed ratio is the basic reproductive number R_0 [17].

Incidence grew exponentially for a much shorter time for highly-local movement than it did for a wider movement kernel, or for non-spatial networks, even when we controlled for R_0 to be within a narrow range (e.g. $(2, 2.2]$, Fig 2). Despite this being a relatively large population, there was no obvious period of exponential growth when we assumed highly local movement. Therefore, given that the basic reproductive number is defined for a genuine renewal process – and its implied exponential growth [17] – it could be argued that R_0 does not exist for some of these networks for our model parameters. However, we did assign a value of R_0 for all simulations based on the most similar subset of consecutive early generations (see Methods). The amplitude of the difference was not driven in any obvious way by the underlying assumptions used to create the networks. These patterns were not specific to the range of values for R_0 (Figs. S3, S4, S5).

Analysis of the higher-order structure of the networks suggests that movement scenarios were driving the observed characteristics of epidemics such as peak timing and attack rate via increased fourth order clustering. We use the term first order clustering for the quantity typically described as the local clustering coefficient [37]: the link density of the immediate neighbourhood of a given node. By extension, we defined order-m clustering coefficient to be the expected proportion of neighbours within m steps on the network who were also neighbours of each other within m steps (Fig 3). We found no relationship between our assumed pattern of movement (α) and first or second order clustering coefficients. There was a weak relationship between α and third order clustering and then a very strong relationship between α and forth order clustering. Patterns between epidemic properties and fourth order clustering for individuals were similar to those between epidemic properties and second order clustering of households, as would be expected, given the bipartite algorithm used to create individual-level networks.

Final size increased with spatial correlation, despite peak size displaying the opposite trend for controlled R^* or R_0. We therefore suggest peak size/final size as a crude measure of adherence to standard epidemic models (with fixed recovery rate $\gamma = 10$ days^{-1}. There was a strong linear relationship between order-m clustering and peak size/final size, that could be explained by α, the strength of spatial embedding, when we control for R_0 (Fig 4B). The gradient of the relationship decreased with order of clustering. Second order household clustering showed the same relationship with peak size as did fourth order individual clustering (Fig 4C). These strong linear relationships only exist when we effectively control for R_0, rather than R^*, and become even less noisy when we reduce the interval of R_0 (not shown).

We conducted a number of sensitivity analyses for these network simulation results. Analytic

approximations for degree distribution $P(K = k)$ and expected first order clustering $\langle CC^1 \rangle$ in our networks are given in Protocol S1, and are independent of α. We confirmed these relationships in Figure S6 by computing these quantities on a set of networks that differ in α. Figure S7 shows the relationship between α and clustering order 1 to 4 on networks generated using a uniform population density. Figure S8 shows the relationship between order-m clustering CC^m and peak size for different values of R_0. Figure S9 provides an illustration of the relationship between higher-order clustering and rewiring probability on a commonly used network model with spatial embedding: the Watts-Strogatz Small World Network [37].

Finally, we map our network model onto a deterministic metapopulation framework so as to relate our simulations of incidence to prior analytical approximations of travelling spatial waves (Protocol S1 for analytic construction). Figure 5 shows the results of simulating on a grid of evenly spaced households of size $h = 4$, where a single continuous variable describes prevalence in each household, and spatial coupling between households used in the force of infection is exactly the kernel used in the construction of our spatially embedded networks. We simulate with randomly spaced seeds (as above), and with a central seed (the center most 4 households), tracking global incidence and local time of peak incidence. The former case yields global incidence curves similar to those generated in our network model (which was seeded similarly). The latter case allows us to identify 4 distinct stages in the propagation of spatial waves that contribute to observed sub-exponential outbreak dynamics in more complex, network-based systems. Figure S11 shows simulation results in 1 spatial dimension with $\alpha = 6$ and $\alpha = 12$, alongside statistical properties of prevalence, which further clarify these growth phases (c.f. figure captions for details and protocol S1 for mathematical analysis).

Discussion

We have shown that non-standard epidemic dynamics can arise from strongly spatially embedded social networks. Using a flexible algorithm of assigning individuals to households and then creating a social networks with widely varying topologies, we can explain the absence of exponential growth and increased attack rate for a given peak height in terms of higher order social structure, while maintaining a standard low-variance offspring distribution. We observe consistent patterns when we control for the basic reproductive number, as measured as directly as possible from a constant ratio of incidence between generations.

The algorithm we used [32] captures the key social contexts of home and workplace while using few parameters, which has allowed us to isolate specific relationships within the epidemic dynamics, across a broad range of network topologies. However, its simplicity is a potential limitation. Specifically, an individual only belongs to a single workplace (which may represent a school or social club). In reality, people will gather non-household contacts from a variety of sources. Also, our networks are not dynamic, which may limit the generalisability of the results to short generation time pathogens.

Accurate empirical data about higher order social contacts would allow us to address some of these issue. There are a number of different approaches to gathering social contact data, including contact diaries, mobile phone apps and tag-based location tracking [28]. Diary methods and current analytical approaches can provide accurate estimates of 1st order moments (degree distribution [29]) and valuable insights into second order moments (clustering [41]). However, these data and current analytical approaches are limited for the estimation of higher order moments. It seems likely that either high resolution mobile phone location data [6] or very high coverage tag-based studies will be needed to reveal these patterns [5]. In addition, further work is needed on the use of algorithms similar to that used here to explicitly fit fully enumerated

social networks to egocentric sample data from a subset of the population (or low coverage non-egocentric data) [22].

Our results can be compared with other disease-dynamic models that produce non-standard incidence profiles. Different functional forms have been suggested for the force-of-infection term in compartmental models that give polynomial growth in the early stages of an epidemic [7, 17]. However, the key features of these model structures may be captured by a more straightforward underlying process [19]. Faster than exponential growth can be achieved with very high variance offspring distributions, which have been inferred by diary studies of social contacts [24]. There is also an extensive literature of much more abstract grid-based models of infectious disease that produce non-standard epidemic dynamic because of very local spatial processes (cellular automata [38]). We note that short periods of super-exponential growth were observed in our results for the simplified 2 dimensional metapopulation example (Fig 5B), arising from from accelerating spatial waves of incidence, not driven by the variance of the offspring distribution.

Prospective forecasting of infectious disease incidence during outbreaks [26] and seasonal epidemics [2] is an active area of public health research. Although non-mechanistic [12] and simple compartmental models [30, 35] have proven most reliable up to now, modern computing capacity enables studies to explore the possibility that incidence forecasts can be improved by the incorporation of realistic social network topology [33, 9]. For example, incidence of Ebola in west Africa in 2013-2016 and currently in central Africa exhibits strong spatial clustering and highly non-standard incidence dynamic, with short periods of exponential growth followed by low sustained peaks in incidence [10]. Future forecasting studies should explore the possibility that that sparse population density and short distances between contacts result in higher-order clustering in the social networks and the resulting non-standard incidence profiles.

Methods

The Model

We simulate 10 independent epidemics for each of 200 parameter sets (h, v, p_w, R^*) drawn from a Latin hypercube, each seeded in 10 randomly selected individuals, and for each $\alpha = 0, 3, 6$. The ranges of values used in the Latin hypercube are given in table S1, and complete parameter sets for all networks are given in supplementary table S1. Our simulations allowed us to track disease incidence and disease generation of each infection.

Assigning R_0 to each simulation

For each simulation output, we calculate the mean reproductive ratio for each generation. For generations 1 to 9 and for each possible consecutive string of 3, 4 or 5 values, we perform a linear regression fit. We define R_0 as mean reproductive ratio over the set of values for which the gradient of this fit is closest to 0 (and all values the remain larger than 1). This allows us to assign a value R_0 to every simulation output.

Higher order clustering

We compute our higher-order clustering coefficients on a subset of 1000 nodes in each network, chosen at random. The algorithm involves storing the network structure as lists of neighbours for each node, and performing an effective contact-tracing procedure. Though it is possible to compute these metrics for all nodes via successive multiplication of adjacency matrices, this procedure becomes computationally expensive in higher orders at networks become large.

Figure 1: For each set of parameters drawn from the Latin hypercube, and for $\alpha = 0, 3, 6$, we show relationships between R^*, R_0 time of peak incidence, and epidemic final size: **(A)** R^*/peak time, **(B)** R^*/peak size, **(C)** R^*/extinction time, **(D)** R^*/final size, **(E)** peak size/final size, **(F)** R^*/R_0 (with the line $R_0 = R^*$ shown in black).

Figure 2: Columns correspond to network structures with $\alpha = 0, 3$ and 6 and simulations with $R_0 \in (2, 2.2]$. Exponential growth in real time is indicated by **(A)** straight lines and **(B)** horizontal lines; **(C)** horizontal lines indicate exponential growth by generation. Figures S3 to S5 show results for a wider range of R_0 values for $\alpha = 0, 3, 6$.

Figure 3: A schematic showing the generalization of clustering coefficient C^1 to higher orders CC^m: CC_i^m measures the density of paths of length $d \leq m$ between the up-to-m neighbours of node i (where node i is shown in gray).

Figure 4: **(A)** $25-$, $50-$ and 75-percentiles of order-m clustering CC^m on networks constructed with different values of α and $h = 5, w = 50, p_w = 0.14, \langle k \rangle = 11$ and $R_0 \in [2, 2.2)$. Plot shows mean values over 3 different networks for each parameter set; **(B)** Using peak size as a crude metric for sub-exponential growth (given a fixed range for R_0), we see linear trends emerging with higher orders of clustering. Plot shows one point per network, with 3 networks generated for each parameter set, and the mean peak size over 10 independently simulated epidemics, All points are numbered with the corresponding value of α; **(C)** Similarly for the household-only networks. Solid lines show linear fits to data, and dotted lines show 95% confidence intervals. Values of linear correlation coefficient and gradient of fits are given in table S2.

Figure 5: Mean-field approximation with $R_0 = 2.2$, $\langle k \rangle = 10$, $h = 4$, using a 100×100 grid of uniformly spaced households: **(A)** seeding in 10 randomly selected households (the same households are used in each simulation), and **(B)** seeding in the centre only. Incidence is given as a proportion of the total population for α ranging from 2 to 6. Supplementary figure S10 shows time of peak incidence in the case $\alpha = 6$ seeded as above.

References

[1] Roy M Anderson, Robert M May, and B Anderson. *Infectious diseases of humans: dynamics and control*, volume 28. Wiley Online Library, 1992.

[2] Matthew Biggerstaff, Michael Johansson, David Alper, Logan C Brooks, Prithwish Chakraborty, David C Farrow, Sangwon Hyun, Sasikiran Kandula, Craig McGowan, Naren Ramakrishnan, Roni Rosenfeld, Jeffrey Shaman, Rob Tibshirani, Ryan J Tibshirani, Alessandro Vespignani, Wan Yang, Qian Zhang, and Carrie Reed. Results from the second year of a collaborative effort to forecast influenza seasons in the united states. *Epidemics*, February 2018.

[3] S P Blythe, C Castillo-Chavez, J S Palmer, and M Cheng. Toward a unified theory of sexual mixing and pair formation. *Math. Biosci.*, 107(2):379–405, December 1991.

[4] M-C Boily, C Lowndes, and M Alary. The impact of hiv epidemic phases on the effectiveness of core group interventions: insights from mathematical models. *Sexually Transmitted Infections*, 78(suppl 1):i78–i90, 2002.

[5] Ciro Cattuto, Wouter Van den Broeck, Alain Barrat, Vittoria Colizza, Jean-François Pinton, and Alessandro Vespignani. Dynamics of person-to-person interactions from distributed RFID sensor networks. *PLoS One*, 5(7):e11596, July 2010.

[6] Eunjoon Cho, Seth A Myers, and Jure Leskovec. Friendship and mobility: User movement in location-based social networks. In *Proceedings of the 17th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '11, pages 1082–1090, New York, NY, USA, 2011. ACM.

[7] Gerardo Chowell, Lisa Sattenspiel, Shweta Bansal, and Cécile Viboud. Mathematical models to characterize early epidemic growth: A review. *Physics of Life Reviews*, 18:66–97, 2016.

[8] Paul C. Cross, Philip L.F. Johnson, James O. Lloyd-Smith, and Wayne M. Getz. Utility of R0 as a predictor of disease invasion in structured populations. *Journal of the Royal Society Interface*, 4(13):315–324, 2007.

[9] John M Drake, Rajreni B Kaul, Laura W Alexander, Suzanne M O'Regan, Andrew M Kramer, J Tomlin Pulliam, Matthew J Ferrari, and Andrew W Park. Ebola cases and health system demand in liberia. *PLoS Biol.*, 13(1):e1002056, January 2015.

[10] Ebola Outbreak Epidemiology Team. Outbreak of ebola virus disease in the democratic republic of the congo, April-May, 2018: an epidemiological study. *Lancet*, 392(10143):213–221, July 2018.

[11] Paul Expert, Tim S. Evans, Vincent D. Blondel, and Renaud Lambiotte. Uncovering space-independent communities in spatial networks. *Proceedings of the National Academy of Sciences*, 108(19):7663–7668, 2011.

[12] David C Farrow, Logan C Brooks, Sangwon Hyun, Ryan J Tibshirani, Donald S Burke, and Roni Rosenfeld. A human judgment approach to epidemiological forecasting. *PLoS Comput. Biol.*, 13(3):e1005248, March 2017.

[13] Tini Garske, Hongjie Yu, Zhibin Peng, Min Ye, Hang Zhou, Xiaowen Cheng, Jiabing Wu, and Neil Ferguson. Travel patterns in china. *PLoS One*, 6(2):e16364, February 2011.

[14] Nicholas C Grassly and Christophe Fraser. Mathematical models of infectious disease transmission. *Nat. Rev. Microbiol.*, 6(6):477–487, May 2008.

[15] Jeff Griffiths, Dawn Lowrie, and Janet Williams. Age-structured model for the AIDS epidemic. *European Journal of Operational Research*, 124(1):1–14, 2000.

[16] David J Haw, Derek A T Cummings, Justin Lessler, Henrik Salje, Jonathan M Read, and Steven Riley. Differential mobility and local variation in infection attack rate, 2019.

[17] J A P Heesterbeek. A brief history of R0 and a recipe for its calculation. *Acta Biotheor.*, 50(3):189–204, December 2001.

[18] H W Hethcote. The mathematics of infectious diseases. *SIAM Rev.*, 42(4):599–653, October 2000.

[19] Thomas House. A general theory of early growth?: Comment on: ???Mathematical models to characterize early epidemic growth: A review??? by Gerardo Chowell et al. *Physics of Life Reviews*, 18:109–111, 2016.

[20] M. Jesse, P. Ezanno, S. Davis, and J.A.P. Heesterbeek. A fully coupled, mechanistic model for infectious disease dynamics in a metapopulation: Movement and epidemic duration. *Journal of Theoretical Biology*, 254(2):331 – 338, 2008.

[21] Matt J Keeling and Ken T D Eames. Networks and epidemic models. *J. R. Soc. Interface*, 2(4):295–307, September 2005.

[22] Pavel N Krivitsky and Martina Morris. Inference for social network models from ego-centrically sampled data, with application to understanding persistent racial disparities in HIV prevalence in the US. *Ann. Appl. Stat.*, 11(1):427–455, March 2017.

[23] Kin O Kwok, Benjamin J Cowling, Vivian W I Wei, Kendra M Wu, Jonathan M Read, Justin Lessler, Derek A Cummings, J S Malik Peiris, and Steven Riley. Social contacts and the locations in which they occur as risk factors for influenza infection. *Proc. Biol. Sci.*, 281(1789):20140709–20140709, August 2014.

[24] Quan-Hui Liu, Marco Ajelli, Alberto Aleta, Stefano Merler, Yamir Moreno, and Alessandro Vespignani. Measurability of the epidemic reproduction number in data-driven contact networks. *Proceedings of the National Academy of Sciences*, 115(50):12680–12685, 2018.

[25] Joel C Miller. Bounding the size and probability of epidemics on networks, 2008.

[26] Martha I Nelson, James O Lloyd-Smith, Lone Simonsen, Andrew Rambaut, Edward C Holmes, Gerardo Chowell, Mark A Miller, David J Spiro, Bryan Grenfell, and Cécile Viboud. Fogarty international center collaborative networks in infectious disease modeling: Lessons learnt in research and capacity building. *Epidemics*, October 2018.

[27] Hiroshi Nishiura, Gerardo Chowell, Muntaser Safan, and Carlos Castillo-Chavez. Pros and cons of estimating the reproduction number from early epidemic growth rate of influenza a (H1N1) 2009. *Theor. Biol. Med. Model.*, 7:1, January 2010.

[28] Jonathan M Read, Ken T D Eames, and W John Edmunds. Dynamic social networks and the implications for the spread of infectious disease. *J. R. Soc. Interface*, 5(26):1001–1007, January 2008.

[29] Jonathan M Read, Justin Lessler, Steven Riley, Shuying Wang, Li Jiu Tan, Kin On Kwok, Yi Guan, Chao Qiang Jiang, and Derek A T Cummings. Social mixing patterns in rural and

urban areas of southern china. *Proceedings Of The Royal Society B-Biological Sciences*, 281(1785):20140268–20140268, June 2014.

[30] Nicholas G Reich, Logan C Brooks, Spencer J Fox, Sasikiran Kandula, Craig J McGowan, Evan Moore, Dave Osthus, Evan L Ray, Abhinav Tushar, Teresa K Yamana, Matthew Biggerstaff, Michael A Johansson, Roni Rosenfeld, and Jeffrey Shaman. A collaborative multiyear, multimodel assessment of seasonal influenza forecasting in the united states. *Proc. Natl. Acad. Sci. U. S. A.*, January 2019.

[31] Steven Riley, Ken Eames, Valerie Isham, Denis Mollison, and Pieter Trapman. Five challenges for spatial epidemic models. *Epidemics*, 10:68 – 71, 2015. Challenges in Modelling Infectious DIsease Dynamics.

[32] Steven Riley and Neil M Ferguson. Smallpox transmission and control: spatial dynamics in great britain. *Proc. Natl. Acad. Sci. U. S. A.*, 103(33):12637–12642, August 2006.

[33] Michele Tizzoni, Paolo Bajardi, Chiara Poletto, José J Ramasco, Duygu Balcan, Bruno Gonçalves, Nicola Perra, Vittoria Colizza, and Alessandro Vespignani. Real-time numerical forecast of global epidemic spreading: case study of 2009 A/H1N1pdm. *BMC Med.*, 10(1):165, December 2012.

[34] James Truscott and Neil M Ferguson. Evaluating the adequacy of gravity models as a description of human mobility for epidemic modelling. *PLoS Comput. Biol.*, 8(10):e1002699, October 2012.

[35] Cécile Viboud, Kaiyuan Sun, Robert Gaffey, Marco Ajelli, Laura Fumanelli, Stefano Merler, Qian Zhang, Gerardo Chowell, Lone Simonsen, Alessandro Vespignani, and RAPIDD Ebola Forecasting Challenge group. The RAPIDD ebola forecasting challenge: Synthesis and lessons learnt. *Epidemics*, August 2017.

[36] Jacco Wallinga, Michiel van Boven, and Marc Lipsitch. Optimizing infectious disease interventions during an emerging epidemic. *Proceedings of the National Academy of Sciences*, 107(2):923–928, 2010.

[37] D J Watts and S H Strogatz. Collective dynamics of 'small-world' networks. *Nature News*, 393(6684):440–442, June 1998.

[38] S. Hoya White, A. Martín del Rey, and G. Rodríguez Sánchez. Modeling epidemics using cellular automata. *Applied Mathematics and Computation*, 186(1):193 – 202, 2007.

[39] WHO Ebola Response Team, Junerlyn Agua-Agum, Archchun Ariyarajah, Isobel M Blake, Anne Cori, Christl A Donnelly, Ilaria Dorigatti, Christopher Dye, Tim Eckmanns, Neil M Ferguson, Robert A Fowler, Christophe Fraser, Tini Garske, Wes Hinsley, Thibaut Jombart, Harriet L Mills, Srinivas Murthy, Gemma Nedjati Gilani, Pierre Nouvellet, Louise Pelletier, Steven Riley, Dirk Schumacher, Anita Shah, and Maria D Van Kerkhove. Ebola virus disease among children in west africa. *N. Engl. J. Med.*, 372(13):1274–1277, March 2015.

[40] Joseph T Wu, Steven Riley, Christophe Fraser, and Gabriel M Leung. Reducing the impact of the next influenza pandemic using household-based public health interventions. *PLOS Medicine*, 3(9):1–9, 08 2006.

[41] Xiong Xiao, Albert Jan van Hoek, Michael G Kenward, Alessia Melegaro, and Mark Jit. Clustering of contacts relevant to the spread of infectious disease. *Epidemics*, 17:1–9, December 2016.