Title: Taxon appearance from extraction and amplification steps demonstrates the value of multiple

controls in tick microbiome analysis

Emilie Lejal [1], Agustín Estrada-Peña[2], Maud Marsot[3], Jean-François Cosson[1], Olivier Rué[4], Mahendra

Mariadassou[4], Cédric Midoux[4,5], Muriel Vayssier-Taussat[6] and Thomas Pollet[1].

[1] UMR BIPAR, Animal Health Laboratory, INRA, ANSES, Ecole Nationale Vétérinaire d'Alfort, Université

Paris-Est, Maisons-Alfort, France

[2] Faculty of Veterinary Medicine, University of Zaragoza, Zaragoza, Spain

[3] ANSES, University Paris-Est, Laboratory for Animal Health, Epidemiology Unit, Maisons-Alfort, France

[4] MaIAGE, INRA, Université Paris-Saclay, 78350, Jouy-en-Josas, France

[5] Irstea, UR PROSE, 92761 Antony, France

[6] INRA, Animal health department, Nouzilly, France

Contacts:

emilie.lejal@vet-alfort.fr

thomas.pollet@inra.fr

muriel.vayssier@inra.fr

Abstract

- **Background**

The development of high throughput sequencing (HTS) technologies has substantially

improved analysis of bacterial community diversity, composition, and functions. Over the last decade,

HTS has been used extensively to identify the diversity and composition of tick microbial communities.

However, a growing number of studies are warning about the impact of contamination brought along

the different steps of the analytical process, from DNA extraction to amplification. In low biomass

samples, e.g. individual tick samples, these contaminants may represent a large part of the obtained

sequences, and thus generate considerable errors in downstream analyses and in the interpretation

of results. Most studies of tick microbiota either do not mention the inclusion of controls during the

DNA extraction or amplification steps, or consider the lack of an electrophoresis signal as an absence

of contamination. In this context, we aimed to assess the proportion of contaminant sequences

resulting from these steps. We analyzed the microbiota of individual *Ixodes ricinus* ticks by including

several categories of controls throughout the analytical process: crushing, DNA extraction, and DNA

amplification.

- **Results**

Controls yielded a significant number of sequences (1,126 to 13,198 mean sequences,

depending on the control category). Some operational taxonomic units (OTUs) detected in these

controls belong to genera reported in previous tick microbiota studies. In this study, these OTUs

accounted for 50.9% of the total number of sequences in our samples, and were considered

contaminants. Contamination levels (i.e. the percentage of sequences belonging to OTUs identified as

contaminants) varied with tick stage and gender: 76.3% of nymphs and 75% of males demonstrated

contamination over 50%, while most females (65.7%) had rates lower than 20%. Contamination mainly

corresponded to OTUs detected in crushing and DNA extraction controls, highlighting the importance

of carefully controlling these steps.

- **Conclusion**

Here, we showed that contaminant OTUs from extraction and amplification steps can

represent more than half the total sequence yield in sequencing runs, and lead to unreliable results

when characterizing tick microbial communities. We thus strongly advise the routine use of blanks and

negative controls in tick microbiota studies, and more generally in studies involving low biomass

samples.

Keywords: Contaminant, Low biomass sample, Tick, Microbiota, High-throughput sequencing

Background

Analysis of microbial community composition by 16S tag sequencing represents a powerful tool commonly used to evaluate the prokaryotic diversity in many ecosystems. Thanks to high-throughput sequencing (HTS) approaches, it is now feasible to identify microbiota with high coverage and reveal the exceptional and complex microbial diversity of environments ranging from the wide ocean (1,2) to small arthropods (3,4). With the constant improvement of sensitivity associated with its decreasing cost, 16S tag sequencing has spread to many areas, including those dealing with low biomass communities such as tick microbiota.

Despite the limitation induced by low biomass, a high number of tick microbiota studies are focusing on individual ticks (4–12) or even on individual or pooled tick organs (9,13–19). The results obtained from these studies have improved our knowledge of tick microbiota, while demonstrating links between its composition and several factors, such as tick stages and gender (5,8,12,18), organ of origin (18), living environment (4,18,20), host blood meal (12), or tick engorgement (15,21,22). The presence of tick-borne pathogens has also been demonstrated to be related to differences in tick microbiota composition and diversity (9,15,19). The continuous improvements in our knowledge, and in particular the assessment of potential links between tick-borne pathogens and particular microbiota members, require us to continue performing analyses on individual ticks, or even better, on individual tick organs.

Several potential biases, inherent to the PCR amplification (chimeric sequences) or sequencing steps (random sequencing errors), are now well known (23), and pipelines have been developed to detect and remove these errors during the sequence analysis. However, several studies have begun to warn scientists about the potential distortion of results due to contamination from the extraction/amplification steps or from the environment during sample processing (23–28). This potential contamination is able to significantly modify results concerning microbe diversity and composition, particularly in the case of low biomass samples, and can have serious consequences on the interpretation of results. For example, while investigating 16s rRNA gene profiles from

nasopharyngeal swabs regularly sampled from a cohort of children from 1 to 24 months, Salter *et al*.

(24) demonstrated that the apparent clustering of samples by age, observed by principal coordinate

analysis, in fact corresponded to the commercial kit used to perform the DNA extraction of these

samples. While studying the influence of laboratory processing on diluted samples, these authors also

observed that the lower the initial biomass of samples, the higher the proportion of contaminants

associated with the kits or the environment. In the context of tick microbiota studies, performing using

blanks or negative controls would therefore be essential to detect potential contamination, and

efficiently distinguish this contamination from genuine tick microbial communities. On the basis of an

exhaustive review of the scientific literature, it is very clear that routine multiple control analyses

(extraction and PCR negative controls) are not considered at their true value, and are only rarely

implemented in tick microbial community analyses (4,6,7). In this context, we performed a 16s rRNA

gene sequencing analysis on individual ticks (females, males and nymphs) considering concomitantly

several negative controls at each step of the process, prior to the sequencing run (tick crushing, DNA

extraction and amplification).

Methods

Tick collection and sample selection

Questing *Ixodes ricinus* nymphs and adults were collected for three years by dragging (from April

2014 to May 2017) in the Sénart forest in the south of Paris. More details on the sampling location and

design, and tick collection, are available in Lejal *et al*. (29). After morphological identification, ticks were

stored at -80°C until further analysis.

Tick washing, crushing, and DNA extraction

Ticks were first washed once in ethanol 70% for 5 minutes and rinsed twice in sterile MilliQ water

for 5 minutes each time. Ticks were then individually crushed in 375 µL of Dulbecco's Modified Eagle

Medium with decomplemented fetal calf serum (10%) and six steel beads using the homogenizer

Precellys®24 Dual (Bertin, France) at 5,500 rpm for 20 seconds. DNA extraction was performed on 100

µL of tick crushing, using the NucleoSpin® Tissue DNA extraction kit (Macherey-Nagel, Germany). For

these previous steps, sterile tubes and filter tips were used on a dedicated area for DNA extraction.

Bench and materials were decontaminated with DNA remover (Molecular BioProducts, San Diego, CA,

USA) before and after each use. Each time a tick crushing or DNA extraction was performed, a crushing

or extraction control was added for the corresponding step.

DNA amplification and multiplexing

In total, 32 males, 35 females, and 557 nymphs were subjected to sequencing analysis. DNA

amplifications were performed on the V4 region of the 16s rRNA gene using the primer pair used by

Galan *et al*. (2016) (16S-V4F: 5'-GTGCCAGCMGCCGCGGTAA-3' and 16S-V4R: 5'-

GGACTACHVGGGTWTCTAATCC-3'), producing a 251 bp amplicon. This primer pair was chosen for its

ability to specifically amplify prokaryotic DNA (86% or 95% of referenced bacterial DNA in the Silva

database, and 52.3% or 93% of Archaeal DNA, when considering 0 or 1 mismatch, respectively). This

characteristic is crucial in the context of tick microbiota analysis, due to the high amount of tick DNA

compared to prokaryotic DNA. An 8 bp-index was added to the forward and reverse primers. The

individual amplification of each sample was performed using a combination of 22 and 32 index-tagged

forward and reverse primers, allowing for amplification and multiplexing of a maximum of 704 samples

that can be loaded together on an Illumina MiSeq flow cell. Amplification and multiplexing steps were

performed under a sterile hood, decontaminated before and after use with DNA remover and UVs,

using tube strips with individual caps to prevent cross-contamination. All the PCR amplifications were

carried out using the Phusion® High-Fidelity DNA Polymerase amplification kit (Thermo Scientific,

Lithuania). For each sample, 5 µL of DNA extract were amplified in a 50 µL final reaction volume,

containing 1X Phusion HF buffer, 0.2 µM of dNTPs, 0.2 U/mL of Phusion DNA polymerase, and 0.35 µM

of forward and reverse primer. The following thermal cycling procedure was used: initial denaturation

at 98°C for 30 s, 35 cycles of denaturation at 98°C for 10 s, annealing at 55°C for 30 s, followed by

extension at 72°C for 30 s. The final extension was carried out at 72°C for 10 minutes. Amplification

step was performed on a peqSTAR 2x thermocycler (PEQLAB, Germany). For each PCR run, negative

controls were performed by using the reaction mixture without template, but using ultrapure water

(Invitrogen, United Kingdom). PCR products were checked on 1.5% agarose gels.

DNA purification, quantification, and pooling

Amplicons were cleaned and quantified using a SequalPrep normalization plate kit (Invitrogen,

USA), and a Qubit dsDNA High Sensitivity Assay kit (Invitrogen, USA), respectively. Amplicons and

controls were pooled at equimolar concentrations and sent to the sequencing platform (GenoScreen,

France).

Controls

In total, 45 negative controls were included and analyzed using the same process as for tick

samples (Figure 1).

12 Crushing controls (CCs): Tube containing crushing solution, remaining open in the room while

putting ticks inside other tubes between the cleaning and crushing steps. DNA extraction and

amplification were then performed on the crushing control in the same conditions as for any other

sample.

12 DNA extraction controls (ECs): For these controls, the different DNA extraction steps were

performed under the same conditions as for any other samples, but without adding sample. DNA

amplification was then performed on the extraction control in the same conditions as for any other

sample.

21 DNA amplification controls (ACs): The DNA amplification step was performed on ultrapure

water (Invitrogen, United Kingdom), in the same conditions as for any other sample.

166   **Figure 1. Process followed by samples and added control steps.**

Sequencing and data processing

The equimolar mix was concentrated and sequenced by GenoScreen (Lille, France) using MiSeq Illumina 2 x 250 bp chemistry. Quality control checks on raw data were performed using FastQC v0.11.3 (30), and no problems were detected. First, pairs of amplicons were merged using Pear v0.9.11 (31) with a minimum overlap of 20 bp, a minimum assembly length of 100 bp, a maximum assembly length of 480 bp, and an e-value of 0.01. Then, the sequencing orientation of extended sequences was checked by using cutadapt v1.12 (32), with primers in forward and reverse strands (without removing them). Sequences with a reversed orientation were reverse-complemented to obtain a set of sequences with the same orientation (5' to 3'). The sequence analysis was performed using FROGS (33). Primers were removed using cutadapt and resulting sequences were dereplicated. Operational taxonomic units (OTUs) were built using Swarm v2.1.12 (34) with parameter d = 3, and chimeras were filtered using vsearch v1.4 (35) in a *de novo* mode. Finally, OTUs with low abundance (< 0.005% of the total sequence count) were filtered out, resulting in 608 OTUs (corresponding to 2,835,506 sequences). OTUs were affiliated by blasting cluster seed sequences against the Silva database v132 (36). OTUs corresponding to mitochondrial/chloroplast DNA or not affiliated to any domain were removed, leading to a dataset composed of 513 OTUs and 2,726,287 sequences.

Data processing to distinguish contaminant versus tick microbiota OTUs

The control composition was used to distinguish tick microbial OTUs from contaminants. All the OTUs corresponding to more than 1% of the total number of sequences obtained in each category of control were identified as main contaminants. However, due to possible errors in sequence assignment occurring during sequencing due to the formation of mixed clusters on the flow cell (37), negative controls can also contain a few sequences of tick microbial OTUs, complicating contaminant identification under this 1% threshold. Therefore, for each OTU, the number of sequences expected to spread to other samples due to false assignment (i.e. expected number) was determined by calculating

0.02% of the total number of each OTU in the dataset, corresponding to a false assignment rate

estimated previously by Galan *et al*. (23). This expected number was compared to the maximum

number of obtained sequences in controls (i.e. maximum number). As this rate was not estimated from

our dataset, we decided to allow variability around this threshold, while calculating a 99% confidence

interval around the expected number of 0.02%. OTUs whose maximum sequence number was above

the upper threshold were considered contaminants. Importantly, cross-contamination from samples to

controls was considered negligible, as no negative controls presented a higher maximum sequence

number than the expected number for an OTU belonging to the most abundant and characteristic

genus of tick microbiota in our dataset (Midichloria), which is also the 3[rd] most abundant OTU in the

dataset (266,516 sequences).

Statistical analyses

The percentage of samples presenting a contamination rate higher than 50% was compared

between samples categories (nymphs, males and females) according to a Chi^2 statistical test.

Alpha diversity was estimated before and after sample cleaning. Samples containing fewer than

500 sequences after cleaning were removed from the analysis, even if they contained more than 500

sequences before cleaning. Several measures of alpha diversity were calculated. Basically, the observed

number of OTUs (species richness) was used as well as the Faith Phylogenetic Diversity (PD) index,

corresponding to the total branch length of the subtree spanned by the community, and also two

indexes taking into account the number of sequences: the Shannon index (more influenced by rare

OTUs) and Simpson index (more influenced by dominant OTUs). Phylogenetic diversity was estimated

using the Picante package (38), while other measures were assessed thanks to Easy 16S

(http://genome.jouy.inra.fr/shiny/easy16S/), based on the phyloseq package (39). Alpha diversity

measure comparison was performed between cleaned and not cleaned groups, as well as between

stages, using non-parametric statistical tests (Wilcoxon and Kruskal–Wallis, respectively). Statistical

computations were performed in R 3.6.0 (40).

Results

Nymphs, males and females presented a mean number of sequences per sample of 3,721,

2,689 and 5,316, respectively. Concerning controls, a mean number of 13,198, 11,235 and 1,126

sequences per sample was obtained for crushing, extraction and amplification controls (CCs, ECs and

ACs), respectively.

The most abundant OTUs, with relative abundance higher than 1% in at least one category of

samples or controls, are represented in Figure 2. All other OTUs, i.e. those with relative abundance

lower than 1% in all categories were grouped together in the "Others" category.

The examination of the at-least-1% OTU composition of the control (CCs, ECs, ACs) allowed to

identify OTUs belonging to the genera: *Pseudomonas*, *Acinetobacter*, *Halomonas*, *Shewanella*,

Tepidimonas, *Esherichia-Shigella*, *Photobacterium*, *Polaromonas*, *Deinococcus* and *Reyranella*, as well

as two OTUs belonging to the Enterobacteriaceae family but multi-affiliated at the genus level, and

one OTU only affiliated to Gammaproteobacteria at the Class level (Figure 2). OTU composition was

contrasted across the control types. While the same OTUs were identified in CCs and ECs and belonged

to *Pseudomonas* [70.7% and 69.3%, respectively], multi-affiliated Gammaproteobacteria [12.2% and

16.7%, respectively], *Acinetobacter* [3.1% and 2.4%, respectively], and multi-affiliated

Enterobacteriaceae [2.7% and 3.6%, respectively]), the microbial community identified in AC controls

was dominated by OTUs belonging to *Halomonas* (58.8%), *Shewanella* (13.6%), *Tepidimonas* (5.1%),

Escherichia-Shigella (2.3%), *Photobacterium* (1.5%), *Polaromonas* (1.5%), *Deinococcus* (1.5%),

Reyranella (1.2%) and multi-affiliated Enterobacteriaceae (1.2%). These OTUs represented 38.9% of

the total number of sequences coming from the tick samples. Focusing on the sample category, these

OTUs correspond to 11.9%, 40.4% and 41.3% of sequences detected in females, males and nymphs,

respectively (Figure 2). These OTUs were therefore considered to be the main contaminants of the

dataset. It is interesting to note that most of these OTUs correspond to those detected in both EC and

CC controls. In fact, only 2.8% of sequences detected in the different categories of samples (nymphs,

females and males) correspond to OTUs detected in the AC controls and identified as main

contaminants.

Figure 2. 16S rRNA gene sequencing taxonomic assignment comparison between negative controls and samples. Percentages are represented for the OTUs corresponding to at least 1% of the total number of sequences assigned to a category of samples or controls, those corresponding to less than 1% are grouped under the term "Others". OTUs belonging to the same genus or family (if multi-affiliated at the scale of genus) were grouped under the same name. One OTU was only identified until the class level (multi-affiliated Gammaproteobacteria).

A more in-depth identification of contaminants, including those corresponding to less than 1% of the total number of sequences was performed, allowing us to identify 160 OTUs as contaminants among the 513 initially identified OTUs in the dataset. This identification increased the proportions of identified contaminants in the samples, which reach 50.9% of the total sequence count from the tick samples, and 20.1%, 51.2% and 53.7% of sequences detected in females, males and nymphs, respectively. Furthermore, after these cleaning steps, 201 samples contained less than 500 sequences compared to 8 before cleaning. For 76.6% of nymph and 75% of male samples, contaminant OTUs represented more than 50% of the total sequence count (Figure 3). These sample contamination rates

271   are significantly higher than those obtained for females ($p > 0.05$ according to a Chi^2 test), for which

272   most of the samples (65.7%) exhibited a contamination rate below 20%.

287   **Figure 3. Impact of contamination on the final number of sequence per sample according to tick stage/gender.**

289   The impact of this contamination on diversity, according to the sample category was estimated

(Figure 4). As expected, for all the studied diversity measures, cleaning led to a decrease in diversity.

Concerning nymphs and females, this decrease was significant for all the tested diversity indexes, while

for males, it was only significant for the observed and phylogenetic diversity measures, suggesting that

for males, contamination concerned mostly low-abundance taxa. The impact of this cleaning on alpha

diversity comparison between tick stages was been studied. Without cleaning, significant differences

in alpha diversity were found between tick stages for observed, Shannon and Simpson measures.

Concerning cleaned data, significant differences were still observed between stages when using

Shannon and Simpson indexes, but disappeared when using the observed diversity measure.

Figure 4. Impact of contamination on alpha diversity estimations according to tick stages. Alpha diversity has

been estimated according to the observed number of OTUs, the Shannon and Simpson indexes and the Faith's

phylogenetic diversity (PD), for each stages on cleaned (turquoise) and not cleaned (orange-red) datasets. Results

of the statistical test comparing alpha diversity measurement between stages, as well as between the cleaned

and uncleaned groups, are represented on coloured bands at the top and at the bottom of the frame respectively:

NS= Not Significant, ***= p < 0.001, **= p < 0.01, *= p < 0.05.

Discussion

In this study, we sequenced the microbiota of 624 individual *I. ricinus* ticks including several

categories of negative controls to identify and quantify the contaminating sequences, in particular

those produced during the extraction or amplification steps. The performance of multiple negative

controls allowed us to perform in-depth identification of contaminant OTUs. These OTUs represented

50.9% of the total sequence yield from tick samples, and had a significant impact on microbial diversity

estimation. This shows the importance of contamination potentially arising from processing steps, and

demonstrates that its consequences on data analyses can be significant and should not be neglected.

A high number of sequences in each category of controls was detected, particularly in CCs and

ECs, in which the mean number of sequences was even higher than the mean sequence number

detected in samples. Importantly, this higher number might be partly overestimated, first due to the

presence of tick DNA during the PCR that may have inhibited the reaction on the 16s rRNA gene in

samples while not in controls, and second during the measure of DNA concentrations that may have

led to a higher estimation of concentrations in samples compared to controls, before the pooling step.

Despite this, the high number of sequences in controls raises the issue of contaminant presence, while

performing each step of the sample processing.

Each processing step leads to a different contamination pattern. Most of the main

contaminants do not seem to originate from the DNA amplification step. Furthermore, as the

composition of crushing and extraction controls was the same, while the reagents used were different

and crushing control also underwent the extraction step, the contaminant sequences most probably

came from the DNA extraction step, rather than from the crushing step. The key role of the extraction

step in the contamination of low biomass samples has previously been demonstrated by Salter *et al*.

(24). Furthermore, three of the four main contaminants identified in our analysis, corresponding to

Pseudomonas, *Acinetobacter*, and multi-affiliated Enterobacteriaceae, belong to families and genera

identified by these authors as contaminants coming from DNA extraction kits. Our observations

differed from those obtain by Galan *et al*. (23), who mainly detected contaminants coming from the

DNA amplification step, while this step represented only 2% of the main contaminants detected in our

samples. In their study of kit contaminants, Salter *et al.* (24) observed that the extent of contamination

arising from DNA extraction was different depending on the kit brand used to perform this step. While

using the less contaminating kit, they extracted DNA from several 10-fold cascade dilutions of a pure

suspension of *Salmonella bongori*, and subsequently assessed 16s rRNA gene concentrations via q-

PCR. They demonstrated a linear relationship between the 16s rRNA gene copy number and the

dilution fold only for the first dilution points, after which levels started to stabilize and to reach a

threshold of 10^5 mean copy number/mL. Consequently, the higher rate of extraction, rather than

amplification contaminants in samples, could be due either to variations of contaminating load

between our extraction and amplification kits, that are not sourced from the same company, or to the

presence of a basal level of extraction contaminant bacterial DNA, potentially limiting the

contamination arising from the next step (amplification).

Interestingly, we showed that in terms of contamination rates, the influence of contaminant

presence was significantly higher in nymph and male samples compared to female ones. Because low

biomass samples were shown to be particularly sensitive to contamination (24), we might suppose that

the amount of bacterial DNA would be lower in males and nymphs, explaining the higher presence of

contaminants in these samples. It would be useful to develop epifluorescence or flow cytometry

protocols adapted to tick samples, to assess whether the amount of bacteria composing the female

microbiota is effectively higher than in the male and nymph. Additionally, previous studies dealing with

tick microbes have identified higher microbiota diversity in males and nymphs compared to females,

while investigating richness and PD (8,18) as well as Shannon and Simpson indexes (18). However, no

negative controls and data cleaning were reported in these studies, and they could therefore result

from contamination. In the present study, we determined that the presence of contaminants can lead

to an overestimation of microbial diversity in all the tick stages, and that this is the case for several

diversity indexes, especially richness and PD estimations, which are the measures most affected by the

cleaning process. Furthermore, the impact of removing contaminants on diversity comparisons

between stages was estimated and showed that this cleaning step can influence the final results, and

particularly in our case, concerning the observed diversity estimation. Our results therefore call for

caution in attempts to draw conclusion about tick stage diversity without considering negative

controls.

As already mentioned by Narasimhan and Fikrig (26), the understanding of arthropod microbial

communities in the context of arthropod survival and pathogen transmission may open new routes in

arthropod-borne pathogen control strategies. In this context, many studies have assessed tick

microbiota composition in individual ticks or even in tick organs (4–19). However, very few reported

the use of negative controls or mentioned the identification and removal of subsequently identified

contaminants (4,6,7), thus obscuring the risk that identified micro-organisms as members of the

microbiota could correspond to contaminants. Nonetheless, the bias generated by the presence of

contaminants can have a considerable influence on the characterization of tick microbiota composition

and structure. Here, we showed that contaminants represented almost 38.9% of the total sequence

yield of tick samples. Furthermore, several contaminants we identified belong to genera (especially

Pseudomonas and *Acinetobacter*) previously reported in several tick microbiota studies (reviewed by

26). This does not necessarily mean that these OTUs are contaminants, as several OTUs belonging to

the same genera may correspond either to tick microbiota members or to contaminants. However,

this points out the lack of reliability concerning this crucial point in the understanding of tick microbiota

composition, and the need to perform negative controls to clean the data set and to rule out the

hypothesis of potential contamination.

Conclusions

Because of their ease of use and the fact that they employ safer chemicals than those in

conventional methods (i.e. phenol chloroform extraction), the use of commercial DNA extraction kits

has become standard, particularly in the context of studies dealing with the identification of tick

microbial communities. However, the issue of contaminants coming from DNA extraction kits has

recently been raised for microbiota studies dealing with low biomass samples, such as blood, intestinal

tissue, mucosal tissue, or nasopharyngeal samples. Our study adds further evidence supporting these

important concerns in the field of tick microbiota. Here, our results suggest that many negative

controls should be performed at each step of sample processing prior to performing microbiota

analyses, allowing us to efficiently identify a potentially serious bias in the study of the actual tick

microbial community. While more and more studies try to identify tick microbiota to in fine potentially

propose hypothesis for the development of new tick-borne pathogen control strategies, we strongly

advise the routine use of negative controls in microbiota studies, particularly in the context of low

biomass samples like ticks or other small arthropods. These procedures should be used routinely in

order to obtain the most reliable results possible.

Declarations

Competing interests

No conflicts of interest are declared.

Acknowledgements

The authors would like to thanks Sabine Delannoy and the IdentyPath genomic platform that allowed us to perform part of the experiments in their laboratory. Thank you also to Maxime Galan for simulating discussions and for providing index sequences.

Funding

This research work was funded by the French National Institute of Agronomic Research (INRA) through grants obtained from the Métaprogramme: "Adaptation of Agriculture and Forests to Climate Change" and "Meta-omics and Microbial Ecosystems". The salary of Emilie Lejal, the PhD student working on this project was funded by the Ile-de-France region.

Authors' contributions

Conceived and designed the experiments: TP, MVT, JFC, EL. Performed the experiments: EL. Analysed the data: EL, AE, TP, MM, MM, CM, OR. Wrote the paper: EL, AE, MM, JFC, OR, MM, CM, MVT, TP.

References

1. Sogin ML, Morrison HG, Huber JA, Welch DM, Huse SM, Neal PR, et al. Microbial diversity in the deep sea and the underexplored "rare biosphere". Proc Natl Acad Sci. 2006 Aug 8;103(32):12115–20.

2. Hamdan LJ, Coffin RB, Sikaroodi M, Greinert J, Treude T, Gillevet PM. Ocean currents shape the microbiome of Arctic marine sediments. ISME J. 2013 Apr;7(4):685–96.

3. Degli Esposti M, Martinez Romero E. The functional microbiome of arthropods. PLOS ONE. 2017
May 5;12(5):e0176573.

4. Estrada-Peña A, Cabezas-Cruz A, Pollet T, Vayssier-Taussat M, Cosson J-F. High Throughput
Sequencing and Network Analysis Disentangle the Microbial Communities of Ticks and Hosts
Within and Between Ecosystems. Front Cell Infect Microbiol [Internet]. 2018 [cited 2019 Mar
15];8. Available from: https://www.frontiersin.org/articles/10.3389/fcimb.2018.00236/full

5. Lalzar I, Harrus S, Mumcuoglu KY, Gottlieb Y. Composition and Seasonal Variation of
Rhipicephalus turanicus and Rhipicephalus sanguineus Bacterial Communities. Appl Env
Microbiol. 2012 Jun 15;78(12):4110–6.

6. Hawlena H, Rynkiewicz E, Toh E, Alfred A, Durden LA, Hastriter MW, et al. The arthropod, but not
the vertebrate host or its environment, dictates bacterial community composition of fleas and
ticks. ISME J. 2013 Jan;7(1):221–3.

7. Rynkiewicz EC, Hemmerich C, Rusch DB, Fuqua C, Clay K. Concordance of bacterial communities
of two tick species and blood of their shared rodent host. Mol Ecol. 2015;24(10):2566–79.

8. Van Treuren W, Ponnusamy L, Brinkerhoff RJ, Gonzalez A, Parobek CM, Juliano JJ, et al. Variation
in the Microbiota of Ixodes Ticks with Regard to Geography, Species, and Sex. Appl Env Microbiol.
2015 Sep 15;81(18):6200–9.

9. Gall CA, Reif KE, Scoles GA, Mason KL, Mousel M, Noh SM, et al. The bacterial microbiome of
Dermacentor andersoni ticks influences pathogen susceptibility. ISME J. 2016 Aug;10(8):1846–
55.

10. Khoo J-J, Chen F, Kho KL, Ahmad Shanizza AI, Lim F-S, Tan K-K, et al. Bacterial community in
Haemaphysalis ticks of domesticated animals from the Orang Asli communities in Malaysia. Ticks
Tick-Borne Dis. 2016 Jul 1;7(5):929–37.

11. Kwan JY, Griggs R, Chicana B, Miller C, Swei A. Vertical vs. horizontal transmission of the
microbiome in a key disease vector, Ixodes pacificus. Mol Ecol. 2017;26(23):6578–89.

12. Swei A, Kwan JY. Tick microbiome and pathogen acquisition altered by host blood meal. ISME J.
2017 Mar;11(3):813–6.

13. Andreotti R, Pérez de León AA, Dowd SE, Guerrero FD, Bendele KG, Scoles GA. Assessment of
bacterial diversity in the cattle tick Rhipicephalus (Boophilus) microplusthrough tag-encoded
pyrosequencing. BMC Microbiol. 2011 Jan 6;11(1):6.

14. Budachetri K, Browning RE, Adamson SW, Dowd SE, Chao C-C, Ching W-M, et al. An Insight Into
the Microbiome of the Amblyomma maculatum (Acari: Ixodidae). J Med Entomol. 2014 Jan
1;51(1):119–29.

15. Narasimhan S, Rajeevan N, Liu L, Zhao YO, Heisig J, Pan J, et al. Gut Microbiota of the Tick Vector
Ixodes scapularis Modulate Colonization of the Lyme Disease Spirochete. Cell Host Microbe. 2014
Jan 15;15(1):58–71.

16. Qiu Y, Nakao R, Ohnuma A, Kawamori F, Sugimoto C. Microbial Population Analysis of the Salivary
Glands of Ticks; A Possible Strategy for the Surveillance of Bacterial Pathogens. PLOS ONE. 2014
Aug 4;9(8):e103961.

17. Clayton KA, Gall CA, Mason KL, Scoles GA, Brayton KA. The characterization and manipulation of
the bacterial microbiome of the Rocky Mountain wood tick, Dermacentor andersoni. Parasit
Vectors. 2015 Dec 10;8(1):632.

18. Zolnik CP, Prill RJ, Falco RC, Daniels TJ, Kolokotronis S-O. Microbiome changes through ontogeny
of a tick pathogen vector. Mol Ecol. 2016;25(19):4963–77.

19. Abraham NM, Liu L, Jutras BL, Yadav AK, Narasimhan S, Gopalakrishnan V, et al. Pathogen-
mediated manipulation of arthropod microbiota to promote infection. Proc Natl Acad Sci. 2017
Jan 31;114(5):E781–90.

20. Carpi G, Cagnacci F, Wittekindt NE, Zhao F, Qi J, Tomsho LP, et al. Metagenomic Profile of the
Bacterial Communities Associated with Ixodes ricinus Ticks. PLOS ONE. 2011 Oct
13;6(10):e25604.

21. Moreno CX, Moy F, Daniels TJ, Godfrey HP, Cabello FC. Molecular analysis of microbial
communities identified in different developmental stages of Ixodes scapularis ticks from
Westchester and Dutchess Counties, New York. Environ Microbiol. 2006 May 1;8(5):761–72.

22. Heise SR, Elshahed MS, Little SE. Bacterial Diversity in Amblyomma americanum (Acari: Ixodidae)
With a Focus on Members of the Genus Rickettsia. J Med Entomol. 2010 Mar 1;47(2):258–68.

23. Galan M, Razzauti M, Bard E, Bernard M, Brouat C, Charbonnel N, et al. 16S rRNA Amplicon
Sequencing for Epidemiological Surveys of Bacteria in Wildlife. mSystems. 2016 Aug
30;1(4):e00032-16.

24. Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, et al. Reagent and laboratory
contamination can critically impact sequence-based microbiome analyses. BMC Biol. 2014 Nov
12;12(1):87.

25. Weiss S, Amir A, Hyde ER, Metcalf JL, Song SJ, Knight R. Tracking down the sources of
experimental contamination in microbiome studies. Genome Biol. 2014 Dec 17;15(12):564.

26. Narasimhan S, Fikrig E. Tick microbiome: the force within. Trends Parasitol. 2015 Jul 1;31(7):315–
23.

27. Glassing A, Dowd SE, Galandiuk S, Davis B, Chiodini RJ. Inherent bacterial DNA contamination of
extraction and sequencing reagents may affect interpretation of microbiota in low bacterial
biomass samples. Gut Pathog. 2016 May 26;8(1):24.

28. Knight R, Vrbanac A, Taylor BC, Aksenov A, Callewaert C, Debelius J, et al. Best practices for
analysing microbiomes. Nat Rev Microbiol. 2018 Jul;16(7):410.

29. Lejal E, Marsot M, Chalvet-Monfray K, Cosson J-F, Moutailler S, Vayssier-Taussat M, et al. A three-
years assessment of Ixodes ricinus-borne pathogens in a French peri-urban forest. bioRxiv. 2019
Apr 5;597013.

30. Andrews S. FastQC: a quality control tool for high throughput sequence data. [Internet]. 2010.
Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

31. Zhang J, Kobert K, Flouri T, Stamatakis A. PEAR: a fast and accurate Illumina Paired-End reAd
mergeR. Bioinformatics. 2014 Mar 1;30(5):614–20.

32. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 2011 May 2;17(1):10–2.

33. Escudié F, Auer L, Bernard M, Mariadassou M, Cauquil L, Vidal K, et al. FROGS: Find, Rapidly, OTUs
with Galaxy Solution. Bioinformatics. 2018 Apr 15;34(8):1287–94.

34. Mahé F, Rognes T, Quince C, Vargas C de, Dunthorn M. Swarm v2: highly-scalable and high-
resolution amplicon clustering. PeerJ. 2015 Dec 10;3:e1420.

35. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool for
metagenomics. PeerJ. 2016 Oct 18;4:e2584.

36. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene
database project: improved data processing and web-based tools. Nucleic Acids Res. 2013 Jan
1;41(D1):D590–6.

37. Kircher M, Sawyer S, Meyer M. Double indexing overcomes inaccuracies in multiplex sequencing
on the Illumina platform. Nucleic Acids Res. 2012 Jan 1;40(1):e3–e3.

38. Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, et al. Picante: R tools
for integrating phylogenies and ecology. Bioinformatics. 2010 Jun 1;26(11):1463–4.

39. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLOS ONE. 2013 Apr 22;8(4):e61217.

40. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2018. Available from: https://www.R-
project.org/

531