Microglia depletion disrupts normal functional development of adult-born neurons in the olfactory bulb

Jenelle Wallace[1,2,3,4], Julia Lord[3], Lasse Dissing-Olesen[4,5], Beth Stevens[4,5,6,7,]*, Venkatesh Murthy[1,2,3,]*

[1]Molecules, Cells, and Organisms training program

[2]Center for Brain Science, Harvard University, Cambridge, MA 02138, USA

[3]Department of Molecular & Cellular Biology, Harvard University, Cambridge, MA 02138, USA

[4]Boston Children's Hospital, F.M. Kirby Neurobiology Center, Boston, MA, USA

[5]Harvard Medical School, Boston, MA, USA

[6]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[7]Howard Hughes Medical Institute, Boston Children's Hospital, Boston, MA, USA

*Correspondence: beth.stevens@childrens.harvard.edu (B.S.), vnmurthy@fas.harvard.edu (V.M.)

Abstract

Microglia play key roles in regulating synapse development and refinement in the developing brain, but

it is unknown whether they are similarly involved during adult neurogenesis. By transiently ablating

microglia from the healthy adult mouse brain, we show that microglia are necessary for the normal

functional development of adult-born granule cells (abGCs) in the olfactory bulb. Microglia ablation

reduces the odor responses of developing, but not preexisting GCs *in vivo* in both awake and

anesthetized mice. Microglia preferentially target their motile processes to interact with mushroom

spines on abGCs, and when microglia are absent, abGCs develop smaller spines and receive weaker

excitatory synaptic inputs. These results suggest that microglia promote the development of excitatory

synapses onto developing abGCs, which may impact the function of these cells in the olfactory circuit.

Introduction

Microglia are critically important for normal brain development in the embryonic and early postnatal

stages (Hammond et al., 2018). Originally thought to be primarily involved in injury and disease, many

recent studies have implicated microglia in diverse neurodevelopmental functions (Tremblay et al.,

2011)(Salter and Beggs, 2014)(Wu et al., 2015)(Hong et al., 2016). However, much less is known about

what role microglia might play in the healthy adult brain, even during the process of adult neurogenesis,

which can be thought of as an extension of developmental processes throughout the lifespan.

During early postnatal development, microglia have been implicated in the regulation of

synaptic development, including activity-dependent synaptic pruning (Stevens et al., 2007)(Schafer et

al., 2012)(Tremblay et al., 2010)(Paolicelli et al., 2011)(Gunner et al., 2019) on one hand and promotion

of synaptic development and maturation on the other (Hoshiko et al., 2012)(Zhan et al.,

2014)(Miyamoto et al., 2016)(Nakayama et al., 2018). Although microglia seem well-positioned to

perform similar roles to facilitate the integration of adult-born neurons into circuits in the adult brain in

the dentate gyrus (DG) and olfactory bulb (OB) (Ekdahl, 2012)(Rodríguez-Iglesias et al., 2019), most

studies on microglial regulation of adult neurogenesis to date have focused on early stages of the

process occurring in the neurogenic niches. For example, hippocampal adult neurogenesis is impaired in

models of neuroinflammation (Monje et al., 2003)(Ekdahl et al., 2003) and in immune-deficient mice (Ziv

et al., 2006). Microglia regulate adult neurogenesis in the subgranular zone of the DG through ongoing

phagocytosis of apoptotic neuroblasts (Sierra et al., 2010), although they do not seem to be similarly

involved in the subventricular zone (SVZ) or rostral migratory stream (RMS) (Kyle et al., 2019) (but see

(Ribeiro Xavier et al., 2015) for counterevidence).

Most of what is known about microglial involvement in later stages of neurogenesis is based on

injury and disease models. Microglial activation via sensory deafferentation in the OB decreases the

number of adult-born neurons (Lazarini et al., 2012) and their spine density (Denizet et al., 2016). In

addition, lipopolysaccharide injection or CX3CR1 knockout activates microglia in the hippocampus and

alters both inhibitory (Jakubs et al., 2008) and excitatory (Bolós et al., 2018) synapses onto adult-born

neurons in the DG. These studies suggest that microglia can modulate the synaptic integration of adult-

born neurons under inflammatory conditions, but raise the question of whether they are similarly

involved in the healthy adult brain.

A recent study documented increased activity in principal neurons in the OB after microglia

ablation (Reshef et al., 2017). Here we investigate the cellular and circuit mechanisms behind this effect

by analyzing the functional development of adult-born granule cells (abGCs) in the absence of microglia.

We demonstrate that microglia ablation during, but not after abGC development reduces activity in

abGCs that make inhibitory connections with principal cells in the OB. We show that microglia normally

interact with spines in developing abGCs, and the volume of these spines is reduced when microglia are

ablated. This is accompanied by a reduction in the amplitude of excitatory but not inhibitory inputs to

abGCs, suggesting that microglia are essential for proper integration of abGCs in adult circuits.

Results

Microglia preferentially interact with spines on abGCs

We labeled cohorts of abGCs born using lentiviral injection into the RMS (Consiglio et al.,

2004)(Livneh and Mizrahi, 2012) of adult 8-12 week old mice. To visualize interactions between

microglia and abGCs, we performed time-lapse *in vivo* two-photon imaging of the dendrites of dTomato-

labeled abGCs in the external plexiform layer (EPL) of the OB over the first four weeks after injection in

CX3CR1-GFP +/- mice, in which microglia are labeled with GFP (Figure 1A). Consistent with previous

observations (Nimmerjahn et al., 2005)(Tremblay et al., 2010), we found that microglial processes were

highly motile and occasionally appeared in close proximity to labeled dendritic spines (Figure 1B). To

quantify whether microglia preferentially interact with dendritic spines (defined as colocalization of a

microglial process with at least 5% of the area of a spine head, see Methods, *Analysis of microglia-spine*

interactions) on abGCs compared to encountering them by chance during the course of continuous

motility, we compared the frequency of interactions between microglial processes and spine heads in

the actual imaging data with the frequency of interactions in a series of images in which the microglia

channel was arbitrarily shifted with respect to the dendritic imaging channel ("Offsets").

Microglia exhibited an impressive degree of motility, interacting with 38.5% of abGC dendritic

spines classified as "mushroom" spines (Figure 1C) and 27.2% of spines classified as "filopodial" spines

(Figure 1H) during the course of our 30-90 minute imaging sessions, which was not significantly different

from the offset data (p = 0.13 and p = 0.39, respectively) (Figure 1D,I). However, we found that microglia

interacted significantly more often with individual mushroom spines than predicted by chance (Data:

mean 0.15 ± 0.00039 interactions/10 min vs. Offsets: mean 0.12 ± 0.016 interactions/10 min, p =

0.048)(Figure 1E) though the length of individual interactions was not significantly higher (p =

0.96)(Figure 1F). In contrast, microglia did not interact with filopodial spines at levels above chance

(Figure 1I-L). Microglia did not cover significantly more of the spine than predicted by chance during

interactions with either spine type (Mushroom: p = 0.72, Filopodial: p = 0.84)(Figure 1G,L).

These results suggest that microglia specifically interact with spines that likely contain functional

synapses (Whitman and Greer, 2007), positioning them to influence synaptic stabilization and

maturation during the early development of abGCs.

Odor responses are reduced in abGCs that mature in the absence of microglia

To assess whether microglial functions are essential for the development of abGCs, we ablated

microglia during the entire timecourse of abGC development, beginning three weeks before lentiviral

labeling (Figure 2A). Microglia ablation using the CSF1R inhibitor PLX5622 formulated in chow as

previously described (Elmore et al., 2014) efficiently ablated microglia from the OB (85% ablation as

assessed by immunostaining, 96% ablation as assessed by flow cytometry) within one week and ablation

could be maintained at similar levels for up to nine weeks with ongoing delivery (Figure 2B, Figure 2—

figure supplements 1 and 2). We found no evidence of any largescale inflammatory response to

microglia ablation, as assessed by immunostaining of glial fibrillary acidic protein (GFAP) (Figure 2–figure

supplement 3), consistent with previous reports of the effects of PLX5622 on the whole brain (Elmore et

al., 2014) and the olfactory bulb (Reshef et al., 2017).

At five to six weeks post injection, when abGCs have reached a functionally mature state

(Wallace et al., 2017), we used two-photon imaging to visualize abGC dendrites *in vivo*. Microglia

ablation did not affect the overall number of adult-born neurons in the OB (Figure 2—figure supplement

4), consistent with other reports, (Reshef et al., 2017)(Kyle et al., 2019), and we could readily identify

dTomato-labeled abGC dendrites in control and PLX-treated mice. Since GCs in the OB are axonless and

their release sites are located at dendodendritic synapses on spines in the EPL (Rall et al., 1966), we

chose to record calcium responses in these dendrites. We first recorded responses in anesthetized mice

to a panel of 15 monomolecular odors (Methods, *Odor stimulation*, Table 1), while simultaneously

imaging morphology in the dTomato channel to aid in region of interest identification and image

alignment (Figure 2C). AbGC responses to odors were sparse as previously described (Figure 2D)

(Wallace et al., 2017), but across the population we could identify dendrites responding to most of the

odors in our panel (Figure 2E). We characterized responses by taking the mean $\Delta F/F_\sigma$ value over a five-

second period following the onset of a two-second odor stimulus and plotted the cumulative

distribution of dendritic responses across all odors. The distribution was shifted left towards lower

responsiveness in PLX-treated mice (p = 2.56e-08) while the noise distributions constructed from blank

trials were not different (p = 0.96)(Figure 2F). Dendrites in PLX-treated mice also responded to fewer

odors (for threshold response criteria, see Methods, *In vivo imaging analysis, Thresholds*) (median (IQR):

Control: 3 (1-6), PLX: 1 (0-4), p = 1.16e-04) (Figure 2G). We also found that lifetime sparseness (Willmore

and Tolhurst, 2001), (bounded between 0 and 1, a low score indicates a sparser representation) was

lower in dendrites in PLX-treated mice (median (IQR): Control: 0.18 (0.067-0.32), PLX: 0.067 (0-0.25), p =

4.18e-04)(Figure 2H). These effects were also significant when we performed hierarchical bootstrapping

(Saravanan et al., 2019) to take into account the fact that we imaged dozens of dendrites from each

mouse, with dendrites in PLX-treated mice responding to fewer odors (mean Control: 3.6 ± 0.38, PLX:

2.8 ± 0.26, p = 0.050)(Figure 2—figure supplement 5B) and having lower median response amplitudes

(median Control: 0.14 ± 0.039, PLX: 0.070 ± 0.016, p = 0.0011) (Figure 2—figure supplement 5D).

However, there was no difference in the median amplitude of responses above threshold (p = 0.77)

(Figure 2—figure supplement 5F), suggesting that the difference in overall responses was mostly

mediated by an increase in the proportion of dendrites that did not respond to any of the odors in our

panel, which was significantly higher in PLX-treated mice (Control: 0.14 ± 0.039, PLX: 0.32 ± 0.039, p =

0.0013) (Figure 2—figure supplement 5E).

While imaging in anesthetized mice allows better control of breathing rate, brain motion, and

possible motivational influences on brain state, granule cell odor representation is significantly different

in awake mice (Kato et al., 2012)(Wienisch and Murthy, 2016)(Wallace et al., 2017). Therefore, we also

imaged abGC dendrites in awake mice and found similar effects as in anesthetized mice (Figure 3A,

Figure 3—figure supplement 1). Dendrites in PLX-treated mice had lower responsiveness (p = 0.037)

(Figure 3B), responding to a lower median number of odors (median (IQR) Control: 1 (0-6), PLX: 0 (0-3), p

= 0.052)(Figure 3C) and having lower lifetime sparseness (median (IQR) Control: 0.067 (0-0.28), PLX: 0 (0-

0.16), p = 0.056) (Figure 3D). Interestingly, while response time courses were similar between control

and PLX-treated mice in the anesthetized state (p = 0.30), allowing us to characterize responses with a

simple mean across the odor analysis period, principal components analysis of the $\Delta F/F_\sigma$ traces for all

cells' responses to all odors revealed different response time courses in awake mice (p = 0.004) (Figure

3—figure supplement 1). These differences were likely not due to changes in active sampling of odors

since sniffing rates were not different during baseline or odor presentation periods (mean Baseline:

Control = 3.49 Hz, PLX = 3.52 Hz, p = 0.91; Odor: Control = 4.00 Hz, PLX = 3.94 Hz, p = 0.81) (Figure 3—

figure supplement 3). To ensure that our analysis of response amplitudes was not complicated by this

possible change in response timing, we also applied an event detection analysis method (using a sliding

window across the response period to detect increases in fluorescence above a noise threshold, see

Methods, *In vivo Imaging analysis, Event detection*) to the awake data and found similar results with

dendrites in PLX mice still characterized by responses to a lower median number of odors (p = 0.037)

(Figure 3—figure supplement 1).

Microglia ablation after development has no effect on odor responses

We next wondered whether the effect of microglia ablation was specific to abGCs developing in

the absence of microglia or whether it might affect abGCs more generally. To address this question, we

modified our experimental timeline to label abGCs and wait three months for them to mature fully

(Figure 4A) before imaging their responses to the same set of odors (Figure 4B) in the same mice before

and after three weeks of PLX5622. In this case we found no significant differences in the distribution of

responses (p = 0.45)(Figure 4C), median number of odors evoking a significant response (median (IQR)

Control: 1.5 (0-6), PLX: 2 (0-5), p = 1.00) (Figure 4D), or lifetime sparseness (median (IQR) Control: 0.096

(0-0.31), PLX: 0.099 (0-0.27), p = 0.86). We verified that our imaging paradigm was stable since there

was also no change in responses when we imaged the same mice for two sessions three weeks apart

without any PLX treatment (Figure 4—figure supplement 1). Even after 9 weeks of PLX treatment, the

level of responsiveness remained stable in mature abGCs (Figure 4—figure supplement 2).

Synapse development in abGCs that mature in the absence of microglia

Since we found that microglia ablation reduces the functional responses of abGCs, we wondered

if there were accompanying changes in excitatory synapses made on abGCs. We studied spines on the

apical dendrites of abGCs in the EPL since our *in vivo* imaging showed lower calcium responses to odors

in these dendrites, which could reflect fewer or weaker synaptic inputs. Four weeks after lentiviral

labeling, we examined spines on apical dendrites in abGCs in tissue sections from control and PLX-

treated mice (Figure 5A,B). We found higher spine density (median (IQR): Control: 0.30 (0.21-0.36)

spines/μm, PLX: 0.40 (0.31-0.48) spines/μm, p = 0.039) (Figure 5C), but smaller spine head volumes in

PLX-treated mice (mean Control: 0.37 ± 0.03 μm^3, PLX: 0.31 ± 0.02 μm^3, p = 0.057) (Figure 5D).

Furthermore, most of the increase in spine density in PLX-treated mice was due to an increase in the

density of filopodial spines that had similar spine head volumes as in control mice (Figure 5—figure

supplement 1). In contrast, mushroom spines had significantly smaller head volumes in PLX-treated mice

(mean Control: 0.45 ± 0.04 μm^3, PLX: 0.36 ± 0.03 μm^3, p = 0.036)(Figure 5—figure supplement 1).

We next investigated the electrophysiological correlates of the observed differences in spine

head size by recording spontaneous excitatory postsynaptic currents (sEPSCs) in abGCs. We used the

same timeline as our *in vivo* imaging experiments, namely microglia ablation beginning three weeks

before lentiviral labeling, continuing until electrophysiological recordings from labeled cells in brain

slices at five to six weeks post injection (Figure 6A,B). We found a similar frequency of sEPSCs in cells

from control and PLX-treated mice (mean Control: 30.8 ± 2.2 Hz, PLX: 30.6 ± 2.4 Hz, p = 0.48) (Figure 6C),

but their amplitude was reduced (mean Control: 10.8 ± 0.6 pA, PLX: 9.0 ± 0.4 pA, p = 0.007) (Figure 6D).

Passive membrane properties including membrane resistance (median Control: 597 MΩ, PLX: 532 MΩ, p

= 0.31) and capacitance (median Control: 14.2 pF, PLX: 13.6 pF, p = 0.61) were unchanged, signifying no

differences in cell surface area or resting membrane properties (Figure 6—figure supplement 1). We also

confirmed that our recording conditions were consistent by verifying that series resistance and the

distance of the recorded cells from the mitral cell layer were not different between groups (Figure 6—

figure supplement 1).

To check whether there might be accompanying changes in inhibition that could offset or

augment the observed changes in excitation, we also recorded spontaneous inhibitory postsynaptic

currents (sIPSCs) in the same cells (Figure 6E). We found no difference in the frequency (p = 0.77)(Figure

6F) or amplitude of sIPSCs (p = 0.79)(Figure 6G), suggesting that abGCs that mature in the absence of

microglia receive weaker excitatory inputs without noticeable accompanying changes in inhibition.

Microglia ablation after development has no effect on synaptic inputs

Since there was no significant change in functional responses in abGCs that matured before

microglia ablation, we checked whether synaptic inputs were also unchanged in this condition using the

same experimental timeline as before and recording sEPSCs in abGCs that experienced three weeks of

microglia ablation after three months of maturation (Figure 7A). There was no significant change in the

frequency (p = 0.76)(Figure 7B) or amplitude of sEPSCs (p = 0.09)(Figure 7C). Inhibitory inputs were also

unchanged (Figure 7—figure supplement 1). These results suggest that microglia ablation only affects

synaptic inputs to abGCs when it occurs during the first five to six weeks of the cells' development

rather than after maturation.

Discussion

We delineate an important role for microglia in the regulation of adult-born neuron integration.

Our study adds to the growing literature documenting physiological and/or behavioral effects following

microglia ablation from the healthy adult brain (Parkhurst et al., 2013)(Torres et al., 2016)(Reshef et al.,

2017). Importantly, our study is the first to link microglia ablation directly to changes in the *in vivo*

activity and functional inputs of a specific affected cell population. We show that eliminating microglia

has functional consequences for abGCs incorporating into the circuitry of the OB, reducing their

responses to stimuli. Furthermore, we go on to investigate the physiological mechanisms and show that

the dampened responses we observe are consistent with reduced spine volume and weaker excitatory

inputs in neurons that develop in the absence of microglia.

Methodological considerations

There are several technical caveats that must be acknowledged to allow proper interpretation of

our study. While ablation of microglia with PLX5622 is highly efficient, it necessarily involves a period of

microglial cell death, and it is unknown how this debris may be cleared or what effects it may have on

other cell types. We have attempted to minimize these effects by beginning all experiments at least

three weeks after initiation of PLX5622 treatment, when the number of microglia should have reached

steady state levels (Elmore et al., 2014), but we cannot exclude the possibility of long-term effects

resulting from microglial cell death. Encouragingly, there seems to be no upregulation of cytokines or

GFAP with PLX treatment generally (Elmore et al., 2014) or in the OB, specifically (Figure 2—figure

supplement 3; Reshef et al., 2017) in contrast to other methods for microglia ablation, which may cause

a significant increase in GFAP+ astrocytes and a cytokine storm (Bruttger et al., 2015). However, we

acknowledge that the array of possible inflammatory responses to microglial disruption is still poorly

understood (Liddelow and Barres, 2017) and may involve both responses to microglial cell death as well

as reactions to the absence of normal homeostatic cues. Future work will be needed to characterize

changes in crosstalk among different cell populations after microglia ablation and implications for the

inflammatory environment. However, that inflammation in the OB has been shown to generally reduce

adult-born neuron numbers (Lazarini et al., 2012) and spine density (Denizet et al., 2016) as opposed to

increasing spine density and reducing excitatory synapse strength as we show here, so it is unlikely that

undetected inflammation relating to microglia ablation can fully account for the results we observe.

We also note that microglia ablation with PLX5622 is brain-wide. AbGCs receive local inputs

from within the OB as well as feedback from other cortical areas and neuromodulatory inputs (Lepousez

et al., 2013), so we cannot unambiguously attribute the effects on abGCs to changes within the OB.

Future work will be necessary to investigate whether there is a change in the balance of distal

(predominantly feedforward) and proximal (mostly feedback) inputs or instead a more general effect on

excitatory synapse maturation and/or maintenance.

AbGCs in the olfactory circuit

AbGCs become responsive to odor stimuli soon after they arrive in the OB and then undergo a

period of functional refinement during which their initially broadly tuned responses become more

selective (Wallace et al., 2017) (although there may also be a subpopulation of abGCs that shows

broadening of responses with maturation (Quast et al., 2017)(Wallace et al., 2017)). Although the

mechanism behind this increase in stimulus selectivity is not well-understood, it may involve both a

decrease in some aspects of excitability (Carleton et al., 2003)(Nissant et al., 2009), especially dendritic

excitability (Wallace et al., 2017), as well as selective reorganization of synaptic inputs such that mature

abGCs become more responsive to a particular glomerular module at the expense of other inputs. We

propose that abGCs in PLX-treated mice experience a normal drop in excitability (likely regulated by cell-

intrinsic mechanisms) without a concomitant selective strengthening of specific synaptic inputs, leading

to sparser odor responses (Figure 8). This is consistent with the higher density of more filopodial-like

spines in PLX-treated mice (characteristic of abGCs at an earlier stage of maturation (Breton-Provencher

et al., 2014)) along with smaller mushroom spines (Figure 5) and lower amplitude sEPSCs (Figure 6). It is

also possible that abGCs in PLX-treated mice could have generally lower excitability, but we regard this

as unlikely since excitability has been mainly shown to affect survival rather than synaptic inputs in

abGCs (Lin et al., 2010), in contrast to what we observe here. Future work will be needed to disentangle

the different aspects of abGC maturation and more fully elucidate which are affected by microglia.

Though the effects on synaptic development that we observe in individual cells are modest

(~15% reduction in median sEPSC amplitude in PLX-treated mice, Figure 6C), these effects combined

nonlinearly across many synapses likely become important near response threshold *in vivo*. Indeed, we

observed larger effects on calcium responses recorded *in vivo* (median response amplitudes halved

across all abGCs imaged in each PLX-treated animal, Figure 2—figure supplement 5D), which likely

represent dendritic calcium spikes or global calcium events accompanying somatic action potentials that

result from simultaneous activation of many feedforward and feedback synapses (Egger et al.,

2005)(Egger, 2008).

Microglia-neuron interactions in the adult brain

Though several studies have examined microglial motility (Nimmerjahn et al., 2005) and

interactions between microglia and neuronal elements (Wake et al., 2009)(Tremblay et al., 2010)(Sipe et

al., 2016), it has been unclear whether microglia interact with synapses more often than would be

expected by chance, given the dense synaptic milieu of the adult brain and the high degree of microglial

process motility. Using automated methods to segment microglial processes from *in vivo* two photon

time lapse imaging experiments and shuffle the resulting images for comparison, we demonstrated

targeted motility towards mushroom spines on abGCs, which are likely to contain established excitatory

synapses (Whitman and Greer, 2007). In the absence of microglia, we suggest that fewer spines attain

mushroom morphology, leading to overall reduced spine volume and correspondingly weaker excitatory

synapses. Concomitantly, we observed an increase in spine density in microglia-ablated animals, which

is likely accounted for by filopodial spines which did not have functional or stable synapses (since we did

not see an accompanying increase in the frequency of synaptic currents in PLX-treated mice). One

potential unifying hypothesis is that microglia may prune weaker synapses, allowing stronger synapses

to strengthen further (Stevens et al., 2007)(Schafer et al., 2012)(Tremblay et al., 2010)(Paolicelli et al.,

2011). However, unlike other sensory systems that experience largescale pruning events during

development, abGCs develop an increasing number of synapses over time (Carleton et al., 2003)(Breton-

Provencher et al., 2014). This does not preclude a role for microglia in pruning inputs to these cells (in a

situation in which synaptic formation occurs at a higher rate than elimination) but suggests that if this

occurs, it is likely to be more subtle and difficult to detect with conventional methods. Although we did

not directly observe pruning of dendritic spines in abGCs in our time lapse imaging experiments, it

remains possible that microglia could prune presynaptic elements (Schafer et al., 2012)(Gunner et al.,

2019)(which we did not image) or that our frame rate was too slow to observe such events (Weinhard et

al., 2018).

278   In addition to synaptic pruning, microglia may be involved in promoting synaptic strengthening.

Deficits in microglial CX3CR1 have been linked to weaker synapses and impaired connectivity (Zhan et

al., 2014)(Reshef et al., 2017) as well as delays in maturation of the AMPA/NMDA ratio (Hoshiko et al.,

2012). It is still unclear whether these effects, similar to those that we observe in the PLX ablation

model, are related to differences in spine surveillance and direct microglial-neuron signaling or due to

altered microglial behavior, including changes in release of moderating factors or interactions with other

cell types, such as astrocytes. We expect that in the future, a new generation of tools for more

specifically modifying microglial gene expression and function during defined time windows will help

clarify the mechanisms by which microglia affect synaptic development.

287   In addition, although most previous work has focused on microglial regulation of synapses,

microglia may also regulate neuronal excitability through other mechanisms (Li et al., 2012)(Peng et al.,

2019). Accordingly, although we found no differences in passive membrane properties between abGCs

in control versus PLX-treated animals, we cannot exclude additional possible effects of microglia on

nonsynaptic aspects of abGC development, including active dendritic properties, that could also

contribute to the reduced responsiveness we observe when microglia are ablated.

Timing of microglia ablation

295   While our results suggest that the effect of microglia ablation is specific to developing abGCs

(since our results with microglia ablation after abGC development did not reach statistical significance in

most cases), we saw similar trends in reduced responses and excitatory inputs regardless of the timing

of microglia ablation. This could be because lentiviral labeling is an imperfect method for isolating a

single cohort of abGCs (so some abGCs in the "after development group" may still be at an earlier stage

of development – see discussion of newcomer cells in (Wallace et al., 2017)). Another possibility is that

microglia ablation also affects mature abGCs, although to a lesser extent than developing abGCs. This

could be because microglia may be involved in the general maturation or strengthening of newly formed

excitatory synapses in GCs, and even mature GCs have high rates of synapse formation (and elimination)

compared to cells in other brain regions (Mizrahi, 2007)(Sailor et al., 2016). In this scheme, developing

abGCs would demonstrate the most significant phenotype due to their higher rates of spine dynamics,

but even mature GCs might accrue smaller effects over time.

The larger effect of microglia ablation on developing rather than mature adult-born neurons

highlights the role of microglia during developmental stages and may explain why some studies have not

found significant effects of microglia ablation in other areas of the healthy adult brain (Elmore et al.,

2014)(Torres et al., 2016).

Consequences for olfactory processing

The reduced activity that we observe in mature abGCs after microglia ablation is consistent with

increased activity in the principal cells they inhibit (Reshef et al., 2017). Since reduced inhibition from

GCs to OB principal neurons has been directly linked to an increase in the time needed to discriminate

odors in challenging olfactory tasks (Abraham et al., 2010), this alteration in the OB circuitry could have

functional consequences (Egger and Urban, 2006). Furthermore, given that abGCs may have an outsized

role in the plasticity that underlies complex olfactory behaviors (Breton-Provencher et al., 2009)(Li et al.,

2018) (Mandairon et al., 2018), microglial regulation of their development may contribute to ongoing

plasticity in the olfactory system.

Acknowledgements
We thank Martin Wienisch, Joseph Zak, Vikrant Kapoor, and Julien Grimaud from the Murthy lab for
assistance with equipment setup and training in relevant techniques. Thanks to Samuel Marsh and the
members of the Murthy and Stevens labs for productive discussions. Thanks to Arnaud Frouin and
Alanna Carey from the Stevens lab for assistance with cloning and production of lentiviral plasmids. We
thank Chen Wang and the Viral Core at Boston Children's Hospital for production of lentivirus.
328

329 **Materials and methods**

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
genetic reagent (*M. musculus*)	CX3CR1-GFP heterozygote	Jung et al. 2000	Cat# Jax: 008451	
genetic reagent (lentivirus)	pLenti-TRE-t2A-dTomato-GCaMP5	Wallace et al. 2017		Produced in house (see Methods, "Viral Vectors)
genetic reagent (lentivirus)	pLenti-TRE-t2A-dTomato-GCaMP6s	Wallace et al. 2017		Produced in house (see Methods, "Viral Vectors)
genetic reagent (lentivirus)	pLenti-TRE-t2A-dTomato	Wallace et al. 2017		Boston Children's Hospital Viral Core
genetic reagent (lentivirus)	pLenti-hSynapsin -tTad	Hioki et al. 2009		Produced in house or by Boston Children's Hospital Viral Core
antibody	Anti-Iba1 (Rabbit polyclonal)	Wako	Cat # 019-19741 RRID:AB_839504	1:500
antibody	Anti-GFAP (Rabbit polyclonal)	Dako	Cat # Z0334 RRID:AB_10013382	1:1000
antibody	Anti-BrdU (Rat monoclonal)	Abcam	Cat # ab6326	1:200

antibody	Anti-NeuN (Mouse monoclonal)	Millipore	MAB377	1:200
antibody	Anti-CD45 (Rat monoclonal)	Biolegend	103116	1:100 (final concentration of 2 µg/mL)
antibody	Anti-CD11b (Rat monoclonal)	Biolegend	101217	1:200 (final concentration of 2.5 µg/mL)
antibody	Anti CD16/CD32 (Rat)	BD Bioscience	553141	1:50 (final concentration of 10 µg/mL)
chemical compound, drug	PLX5622	Plexxikon	PLX5622	1200 mg/kg
chemical compound, drug	BrdU	Sigma Aldrich	Cat # 19-160	100 mg/kg
software, algorithm	FlowJo	FlowJo	Version 10.2	https://www.flowjo.com/solutions/flowjo/downloads/previous-versions
software, algorithm	Imaris	Oxford Instruments	Versions 8 and 9	https://imaris.oxinst.com/
software, algorithm	pClamp	Molecular Devices	Version 10.3	
software, algorithm	Matlab	Mathworks	Version 2019b	

software, algorithm	MiniAnalysis	Synaptosoft	Version 6.0.3	http://www.synaptosoft.com/MiniAnalysis/
other	Blue Dead Cell Stain	ThermoFisher	L34961	
other	Counting Beads (CountBright)	ThermoFisher	C36950	

Mice
Mice were C57BL/6J males (Jackson Laboratories) or CX3CR1-GFP heterozygotes (Jackson Laboratories
stock #008451) (Jung et al., 2000) that were 8 to 12-weeks-old at the beginning of the experiment. Mice
were singly housed after chronic cranial window implantation or housed with littermates for
experiments that did not require an implant. In both cases, they were housed on a 12-hour reversed
light/dark cycle after window implantation or viral injection. Littermates were randomly assigned to
experimental groups (control vs. PLX). All procedures were performed using approved protocols in
accordance with institutional (Harvard University Institutional Animal Care and Use Committee) and
national guidelines.
339

Viral vectors
For all experiments involving lentiviral labeling of abGCs, we used a Tet-Off lentiviral system (Hioki et al.,
2009) with one construct expressing the transactivator tTAad under control of a synapsin promotor
(lenti-STB) and a second construct expressing structural and/or activity markers. For spine quantification
in fixed tissue (Figure 5) and electrophysiology (Figures 6 and 7), we used lenti-STB and lenti-dTomato
produced by the Boston Children's hospital. For imaging microglia-spine interactions (Figure 1), we used
lenti-STB and lenti-dTomato-t2A-GCaMP5 and for all experiments measuring odor responses (Figures 2,
3 and 4) we used lenti-STB and lenti-dTomato-t2A-GCaMP6s produced in house. VSV-G pseudotyped
lentiviral vectors were produced by transfection of human embryonic kidney cells (HEK293FT) with
third-generation lentivirus plasmids using lipofection (Mirus TransIT®-293). Supernatant was collected
48 h after transfection and concentrated using ultrafiltration (Centricon Plus-20 PLGC centrifuge filter
units).
352

Lentiviral injections and cranial window surgeries
Reproduced from (Wallace et al., 2017):
"Mice were anesthetized with an intraperitoneal injection of ketamine (100 mg/kg) and xylazine (10
mg/kg) and body temperature was maintained at 37°C by a heating pad. A small craniotomy was made
bilaterally over the RMS injection sites (coordinates from bregma: A +3.3, L +0.82, from the brain
surface: V-2.9 and -2.7) and 250 nL of lentivirus (1:1 mixture of both constructs or 1:50 dilution of the
tTA-containing construct to achieve sparser labeling for spine quantification in Figure 5) was injected at
each of the two depths using a pulled glass micropipette (tip diameter approximately 10-20 um)." For
cranial windows, the surface of the brain was kept moist with artificial cerebrospinal fluid (125mMNaCl,
5mMKCl, 10mMGlucose, 10mMHEPES, 2mM CaCl2 and 2mM MgSO4 [pH 7.4]) and Gelfoam (Patterson
Veterinary) and a glass window consisting of two 3 mm No. 1 coverslips (Warner) glued together with
optical glue (Norland Optical Adhesive 61) was implanted as previously described (Adam and Mizrahi,

2011). For mice used for awake imaging, Kwik Sil (World Precision Instruments) was placed between the
coverslip and the brain surface to reduce movement. In this case, the coverslip consisted of two 3 mm
and one 4 mm No. 0 coverslips forming a plug (Dombeck and Tank, 2014) with the 4 mm coverslip cut
with a diamond knife to fit between the mouse's eyes. In both cases, the edges around the coverslip
were sealed with Vetbond (3M) and then C&B-Metabond dental cement (Parkell, Inc.). A custom-made
titanium headplate (eMachineShop) was cemented to the skull. After surgery, mice were treated with
carprofen (6 mg/kg) every 24 hours and buprenorphine (0.1 mg/kg) every 12 hours for 5 days."
372

Two photon imaging of microglia-spine interactions (Figure 1)
A custom-built two-photon microscope (Wienisch et al., 2012) was used for in vivo imaging.
Fluorophores were excited and imaged with a water immersion objective (20x, 0.95 NA, Olympus) at
950 nm using a Ti:Sapphire laser (Mai Tai HP, Spectra-Physics). The point spread function of the
microscope was measured to be 0.66 x 0.66 x 2.26 μm. Image acquisition and scanning were controlled
by custom-written software in Labview. Emitted light was routed through two dichroic mirrors (680dcxr,
Chroma and FF555- Di02, Semrock) and collected by two photomultiplier tubes (R3896, Hamamatsu)
using filters in the 500-550 nm range (green channel, FF01-525/50, Semrock) and 572-642 nm range (red
channel, FF01-607/70, Semrock). Fields of view were 75x75 μm square spanning 800x800 pixels. Z-stacks
of approximately 30 μm depth with a 1 μm z step for both channels (16 bit) were taken every 3 minutes
(0.5 Hz frame rate with 3x averaging during acquisition) for periods of 30-90 minutes. Two or three fields
of view were imaged in each mouse.
385

Analysis of microglia-spine interactions (Figure 1)
Since both channels exhibited bleed-through with our imaging parameters, the ImageJ spectral unmixing
plugin (Author: J. Walter) was used to calculate and apply unmixing matrices for each image stack prior
to further analysis. In Fiji, spine heads were delineated manually for each time point in the frame where
they appeared brightest using the oval or polygon tools and ROI Manager and classified as either
mushroom (spines whose spine head was wider than the spine neck at all timepoints) or filopodial
(without a well-defined head). The Weka segmentation plugin (Arganda-Carreras et al., 2017) was used
to perform binary segmentation of microglial processes from background after training on 5 frames
(that were fully segmented manually) selected to represent a variety of microglial morphologies and
brightness variation across the three mice. To optimize our resolution, we segmented very
conservatively by using z stacks to mark microglial processes only in the plane where they appeared
brightest. This strategy combined with only delineating spine heads in the brightest frame means that
we only detected the closest interactions between microglial processes and spine heads. The features
chosen for segmentation training in Weka were Gaussian blur, Sobel filter, Hessian, and Difference of
Gaussians. This approach allowed us to segment complex microglia morphology from background
automatically in every image frame and obviated the need for corrections for bleaching or variations in
brightness across different imaging fields. Imaging frames that were too dim to segment (usually due to
loss of immersion water) were excluded. Each segmented image stack was checked manually to ensure
that any residual bleed-through from the red channel did not appear in the segmentation. After
segmentation, ROIs representing spine heads were exported to Matlab using a custom-written macro
with the command getSelectionCoordinates. In Matlab, the ROIs were loaded onto the segmented
image and the mean value of the binary microglia channel within each ROI at each timepoint (0 if there
was no colocalization or up to 1 if all pixels were colocalized with a segmented microglial process) was

measured. The frame was quantified as containing a microglia-spine interaction if the value of
colocalization was greater than 0.05 (at least 5% of the spine head area overlapped by a microglial
process). To compare the overlap between microglial processes and spine heads in the real data
compared to what might be expected by chance, we iteratively translated the microglia channel relative
to the marked spine head ROIs to calculate distributions for all the measured interaction parameters
(Dunn et al., 2011). To do this, we took the original segmented image stack and translated it
horizontally, vertically, or horizontally and vertically first by the maximum spine width across the whole
data set (~32 pixels) to ensure none of the offsets would overlap the real data and then iteratively by
the mean spine width (~10 pixels) to ensure each offset would be as uncorrelated as possible for a total
of 231 offsets. Note that this method likely underestimates the significance of interactions in the real
dataset because microglia cell bodies were never colocalized with spines in the real dataset (dendrites
overlapping microglia cell bodies were not chosen for imaging) but were likely colocalized and quantified
as interacting with spines in some of the offset datasets. To produce Video 2, frames were first
registered with the MultiStackReg plugin (Author: Brad Busse) based on the magenta channel and
bleaching was corrected with histogram matching in ImageJ (these steps were not necessary for analysis
because we segmented each image frame separately as described above).
425
Two photon imaging of odor-evoked responses (Figures 2 and 4)
Animals were matched in littermate pairs before cranial window surgery. All animals with a clear region
of the cranial window and visible lentiviral expression were used for imaging (2/4 control mice and 2/4
PLX-treated mice in the first experiment, 1/2 control mice and 1/2 PLX mice in a second experiment, and
2/4 control mice and 4/4 PLX-treated mice in a third experiment). Imaging was performed at 930 nm
with the same two photon microscope described above.
Reproduced from our previous work (Wallace et al., 2017):
"Animals were anesthetized with an intraperitoneal injection of ketamine and xylazine (90% of dose
used for surgery) and body temperature was maintained at 37°C by a heating pad. Frame rates were 4
Hz, the pixel size was 0.5 μm, and fields of view measured 150 x 150 μm. To locate regions for imaging, a
low magnification z stack (~300-500 μm square) at slow scanning speed (usually 0.5 Hz) with a 1-2 μm z
step was taken from the surface of the dura to the granule cell layer. Planes with many dendrites
perpendicular to the imaging axis were chosen for imaging during odor stimulation."
439
Two photon imaging in awake animals (Figure 3)
A subset of the animals that were imaged under anesthesia were chosen for awake imaging (all animals
that had sufficiently stable cranial windows were chosen from each of the two experiments). Animals
were water-restricted beginning 1-2 days before being handled and accustomed to head-fixation in a
restraining tube (Guo et al., 2014) for 1-2 days with manual delivery of water rewards (approximately 30
minute sessions each). They were then acclimated to the sound of the scan mirrors and odor delivery
(using the full odor set) on the day before imaging with manual delivery of water rewards before
imaging and periodically between sets of repetitions. The same protocol was repeated for 1 or 2 days of
imaging for each mouse.
449
Odor stimulation
Odor lists are found in Table 1. Odors (Sigma) were delivered with a custom-built 16 channel
olfactometer at a nominal volumetric concentration of 16 % (v/v) in mineral oil and further diluted by 16

times in air to a final concentration of approximately 1% (except for isoeugenol which was not diluted in

the olfactometer and therefore had a final concentration of approximately 6.25%). Odors were

presented for 2 s with an interstimulus interval of 40 s with 3-5 times repetitions. The order of odor

delivery was not randomized. A "no odor" trial with the same parameters but in which no odor valve

opened was included with each set of repetitions. Odors were delivered through a mask with balanced

input and output air flow that also allowed us to record respiration (Grimaud and Murthy, 2018). The

positioning of the mask was adjusted daily to ensure optimal signal to noise. A photoionization detector

(miniPID, Aurora Scientific) was used to confirm that odor concentrations were consistent between trials

with these parameters. Odors were replaced before each set of experiments.

Table 1

Ethyl tiglate		Valeraldehyde	
Ethyl valerate		Isoamyl acetate	
Valeric acid		Anisole	
Allyl butyrate		Ethyl propionate	
Carvone		Propyl acetate	
2-methoxypyrazine		2-heptanone	
Isoeugenol		Acetophenone	
Allyl tiglate			

464
465

In vivo imaging analysis (Figures 2, 3, 4)
Data were analyzed offline using custom-written scripts in MATLAB (Mathworks). Experimenters were
blind to fluorescence changes during data analysis but not to experimental group.

Regions of interest (ROIs)
Dendritic ROIs from abGCs were chosen based on average intensity projections in the dTomato channel
as previously described (Wallace et al., 2017). Fields of view were non-overlapping and separated by at
least 100 μm to minimize the chance of the same dendrites appearing in multiple fields of view. Z stacks
of each imaging region were taken with a 2 μm z step from the surface of the dura down to the
convergence of GC dendrites into a single apical dendrite. The density of labeling precluded tracing of all
dendrites back to their parent cell, but we used these z stacks to ensure that multiple ROIs were not
chosen from the same dendrite. Therefore, these data should be considered a sample from a population
of dendrites rather than cells since some dendrites may have originated from the same cell. We
previously estimated with our labeling and imaging strategy that about 1.4 times more dendrites than
cells are represented in the dataset (and this is almost certainly an overestimate since this was based on
the first week of imaging, where labeling is significantly sparser than at later timepoints). For ablation in
abGCs after development (Figure 4), the same fields of view were imaged before and after PLX
treatment. For Figure 4—Supplement 2, we chose matching ROIs for the two sessions (any ROIs without
similar morphology between the two sessions were excluded as described previously (Wallace et al.,
2017)) to quantify how much our imaging conditions changed between sessions without PLX treatment.
However, we opted to choose ROIs independently for Figure 4 since matching ROIs always results in the
exclusion of many ROIs, reducing sample size.

Motion correction
To correct for fast lateral motion and image drift, all image frames for a given field of view were aligned
to the average of the first trial using cross-correlation based on rigid body translation (ImageJ plugin
Moco)(Dubbs et al., 2016). Frames with out of frame motion were removed based on the cosine
similarity between each frame and the average intensity projection of the first trial (or the user-
determined trial with the least motion for awake imaging). Image frames with cosine similarity that
differed by more than 25% (30% for awake imaging) from the mean value for the user-determined best
trial for that field of view or more than 20% (15% for awake imaging) from the mean of the baseline
period for that trial were discarded. In some trials, immersion water dried up or laser power fluctuated,
so trials were removed if their average brightness was less than half of the brightness of the average of
the first three trials or if difference in brightness between the baseline and odor periods was greater
than three times the standard deviation of brightness in the first trial. The entire trial was removed if,
after these corrections, it contained less than 75% of the original frames during either the baseline or
odor analysis period.

Fluorescence changes
The average intensity in the GCaMP channel was calculated for each ROI, for each frame and for each
odor. A response value for each cell-odor pair was calculated as the average $\Delta F/F_\sigma$ value over the odor
analysis period (5 s following odor onset) where F_σ represents the standard deviation of fluorescence
during the baseline period. We used F_σ because we found that in many cases the baseline GCaMP6s
fluorescence was so low in abGC dendrites that we could not reliably subtract the background as
described previously (Wallace et al., 2017). Bleaching was corrected by fitting a single exponential to the
florescence during the baseline period and taking the value at the end of the baseline period as the
baseline mean, only for ROIs where the fluorescence during the last 2.5 s of the baseline period was
greater than 1.1 times the fluorescence during the first 2.5 s. If, after correction for bleaching, baseline
F_σ was greater than 30% of the mean baseline fluorescence, that ROI was considered too noisy and was
removed from the analysis.

Event detection (Figure 3—Supplement 1)
Events were detected separately in each trace using the calculated ROC threshold (see next section) and
any frames that were included in an event in any repeat were averaged across all repeats to obtain a
mean event trace and the mean value in a 1 second period around the peak in this mean trace was then
calculated as the average response to that odor. For latency, the mean latency across repeats was
calculated for all repeats that had detected events.

Thresholds
For all figures where a threshold was applied to the data, thresholds were calculated based on the
distribution of "no odor" trials. An area under the receiver operating curve analysis was performed and
the lowest threshold yielding a 10% false positive rate was chosen. Thresholds were calculated for each
figure by combining responses from both control and PLX-treated groups and performing ROC analysis
on the combined data. For event detection, we used ROC analysis to find the optimal combined
threshold for the number of frames and standard deviation above baseline and chose the threshold that
gave closest to a 10% false positive rate.

Lifetime sparseness
After applying a threshold to the data, we used the following equation to calculate lifetime sparseness:
(Willmore and Tolhurst, 2001)

$$LS = \frac{\left(\Sigma_{j=1}^{m}\frac{r_j}{m}\right)^2}{\left(\Sigma_{j=1}^{m}\frac{r_j^2}{m}\right)}$$

where m = number of odors, r_j = response of the neuron to odor j.

If all stimuli activate a cell nearly uniformly, LS will be close to 1, and if only a small fraction of the stimuli

activate a cell significantly, LS will be close to 0. For any cells with all responses below threshold, we set

LS = 0, interpreting this as the sparsest possible representation.

Temporal dynamics

Principal component analysis of the time course of responses was performed in Matlab using centered

data and singular value decomposition as described previously (Wienisch and Murthy, 2016). To

compare timecourses for the control and PLX-treated groups, principal components were calculated on

each dataset (all traces from all cell-odor pairs) separately and the angle between the two spaces

spanned by the coefficient vectors for the first three principal components from the beginning to the

end of the odor analysis period was calculated. Then a permutation test was performed in which the

group to which each trace belonged was shuffled 1,000 times, and the angles between new coefficient

vectors were calculated based on random division into two groups of the same size as the original

datasets. The actual angle was then compared to this distribution to obtain a p value.

Respiration measurements

Peaks were extracted from respiration traces using the findPeaks function in Matlab with a minimum

peak distance of 10 Hz.

Raincloud plots

Raincloud plots of the type in Figure 2G were created using the Matlab version of the RainCloudPlots

package (Allen et al., 2018).

Microglia ablation with CSF1R inhibitor PLX5622

CSF1R inhibitor PLX5622 was generously provided by Plexxikon (Berkeley, CA) and mixed into

standard rodent diet at 1200 mg per kilogram of chow (Research Diets: AIN-76A diet). Control

diet was formulated identically, but without the inhibitor.

Flow Cytometry (Figure 2--Supplement 1)

Microglia ablation was confirmed via flow cytometry. A single cell suspension enriched for microglia was

generated as previously described (Hammond et al., Immunity 2019). Briefly, mice were deeply

anesthetized with a mixture of ketamine (100 mg/kg) and xylazine (10 mg/kg) and transcardially

perfused with 20 mL of cold Hank's balanced salt solution (HBSS, GIBCO, 14175-079). Bulbs and brains

were minced using a razor blade (Electron Microscopy Science, 71960) and homogenized using a dounce

tissue grinder (Wheaton, 357542). Microglia were enriched via centrifugation in 40% Percoll (Sigma-

Aldrich, 17-0891-01) at 500g for 1h at 4°C. Samples were incubated for 20 min with Blue Dead Cell Stain

(Thermo Fisher, L34961) and Fc blocking antibody (Rat Anti-Mouse CD16/CD32, BD Bioscience, 553141)

in HBSS + 2 mM EDTA. Cells were additionally stained for 20 min with antibodies against CD45

(Biolegend, 103116) and CD11b (Biolegend, 101217) in buffer (HBSS + 2 mM EDTA + 0.5% BSA).

Counting Beads (CountBright, Thermo Fisher Scientific, C36950) were added and samples analyzed using

a FACS Aria II 'SORP'. All events were collected until a total of 8000 counting beads had been acquired

for each sample. The data were analyzed in FlowJo 10.2.

BrdU injections (Figure 2—Supplement 3)
Mice received two intraperitoneal injections of BrdU (Sigma, 100 mg/kg in 0.9% saline) 12 hours apart.
574

Fixed tissue preparation
Mice were deeply anesthetized with a ketamine/xylazine mixture and perfused transcardially with 20 mL
of PBS (pH 7.4) first, followed by 30-50 mL of 4% paraformaldehyde (diluted from 16% stock,
Electron Microscopy Sciences) in 0.1 M phosphate buffered saline (pH 7.4). Brains were removed from
the skull and placed in 5 ml 4% paraformaldehyde for 2 hours. They were then rinsed with PBS and one
hemisphere for each mouse was sliced coronally at 100 μm with a vibratome (Leica) for imaging of
dTomato-labeled abGC spines (Figure #), while the other hemisphere was sliced sagitally at 35-40 μm for
immunostaining (Iba, Figure # or BrdU, Figure #).
583

Immunohistochemistry
For Iba-1 staining, 3-4 slices per mouse spanning the olfactory bulb were permeabilized and blocked
with a solution containing 0.1% Triton X-100 (Fisher), and 5% goat serum in PBS for 1 hour at room
temperature or blocked with Starting Block (ThermoFisher) with 0.3% TritonX-100 for 1 hour at room
temperature and then incubated overnight at 4°C with the primary antibodies rabbit anti-Iba-1 (Wako:
019-19741, RRID:AB_839504) at 1:500 or rabbit anti-GFAP (Dako: Z0334, RRID: AB_10013382) at 1:1000
and then secondary antibodies (Alexa goat-647 anti-Rabbit) for 2 hours at room temperature. For
BrdU/NeuN staining, one of every eight slices per mouse was chosen. Slices were washed in PBS with
0.1% Triton X three times for five minutes each before being incubated for in 2N HCl for 10 minutes at
room temperature and then 20 minutes at 37°C. They were then placed in 0.1M Boric Acid buffer for 15
min and washed again three times with PBS. All slices were then incubated in starting block
(ThermoFisher) with 0.3% Triton X for one hour before being staining in in PBS with 0.3% Triton X with
rat anti-BrdU (Abcam: 6326 at 1:200), and mouse anti-NeuN (Millipore: MAB377 at 1:200) primary
antibodies for 36-48 hours at 4°C and then secondary antibodies (Alexa Fluor 488 and 594 at 1:200) for 2
hours at room temperature. Slices were treated with 0.2% w/v Sudan Black in 70% EtOH for 5 min
before mounting.
600

Confocal imaging and quantification
Slices were mounted with DAPI mounting media (Vectashield DAPI) and imaged with a confocal
microscope (LSM 880, Zeiss). Reported cell densities were calculated based on distances in fixed tissue,
uncorrected for volume changes due to fixation and mounting. All imaging and quantification were
performed blind to the experimental group of the animal (PLX-treated or control).
Iba1 and GFAP Quantification (Figure 2—figure supplements 2 and 3)
For the 1- and 4-week timepoints, one z-stack per animal was imaged at 10X with pixel size 0.42 x 0.42 x
1 μm spanning the thickness of the slice. For the 9-week timepoint, two z-stacks per animal were
imaged at 20X and the counts from both were averaged. Stacks were imaged with pixel size 0.59 x 0.59 x
1 μm spanning 10 μm and converted to maximum intensity projections. The polygon tool was used to
outline the granule cell layer in each image and the area was measured. Iba-1 or GFAP positive cells
were counted in this area manually using the Cell Counter plugin on maximum intensity projection
images in ImageJ. Cells were counted only if the cell body was fully included in the image stack.

BrdU Quantification (Figure 2—figure supplement 4)
For BrdU/NeuN, two z-stacks per OB (one centered dorsally and one centered ventrally) were taken at
20X with pixel size 0.52 x 0.52 x 0.89 μm spanning 9.8 μm. BrdU counts were performed using the
automatic spots function in Imaris (Bitplane) with the same quality settings for spot detection
for all images (quality threshold 2370, number of voxels threshold 524). Cells were counted as positive if
they were located in the granule cell layer and were also positive for NeuN. The area of the granule cell
layer in each image was measured in ImageJ, and the density of BrdU/NeuN positive cells was calculated
for each image and averaged for all images for each mouse.

Spine quantification (Figure 5)
All images were 20-40 μm z-stacks with 0.42 z-step taken with a 63x oil immersion objective, and four
fields of view were imaged per mouse. Care was taken to ensure minimal saturation (less than 5% of
pixels). Secondary and tertiary apical dendrites that were judged to be sufficiently bright, well-separated
from adjacent or overlapping dendrites, at least 40 μm long, and extending at an angle from the imaging
plane less than 45 degrees were chosen for tracing in Imaris. Each dendrite and all its dendritic
protrusions less than 10 μm in length were manually traced using the Filaments function. Data from
each dendrite were exported in a text file and imported into Matlab for plotting. For spine density, the
Imaris property "Filament No. Spine Terminal Pts" was used (meaning that branched spines were
counted by the number of spine heads rather than attachment points). For spine head volume, the
Imaris property "Spine Part Volume Head" was used. For the classification of filopodial and mushroom
spines (Figure 5—figure supplement 1, we used the Imaris properties "HeadMaxDiameter" and
"NeckMeanDiameter" and classified spines as filopodial if they had maximum head diameter less than
1.5 times the neck mean diameter. This threshold was chosen because it led to a similar ratio of
mushroom to filopodial spines in controls as previously observed (Breton-Provencher et al., 2014).
637

Electrophysiology (Figures 6 and 7)
Two experiments with two littermate pairs each (control and PLX-treated) were performed for
electrophysiology. Mice were deeply anesthetized with a mixture of ketamine (100 mg/kg) and xylazine
(10 mg/kg) and perfused with ice-cold modified ACSF solution (in mM: 120 choline chloride, 25 glucose,
25 NaHCO3, 2.5 KCl, 0.5 CaCl2, 7 MgSO4, 11.6 ascorbic acid, 3.1 pyruvic acid, 1.25 NaH2PO4). Brains
were removed and placed in the same ice-cold modified ACSF. Horizontal slices (300 μm thick) of
olfactory bulbs were cut using a vibratome (VT1000S; Leica, Germany). Slices were incubated in
oxygenated holding solution (in mM: 119 NaCl, 26.2 NaHCO3, 1 NaH2PO4*H2O, 2.5 KCl, 22 glucose, 1.3
CaCl2, 2.5 MgSO4) at 33°C for at least 30 minutes before being transferred to oxygenated ACSF (in mM:
119 NaCl, 26.2 NaHCO3, 1 NaH2PO4*H2O, 2.5 KCl, 22 glucose, 2.5 CaCl2, 1.3 MgSO4). Extra slices were
maintained in holding solution at room temperature. Whole-cell recordings (Bessel filtered at 2.2 kHz
and acquired at 10 kHz) were performed using patch pipettes filled with internal solution (in mM: 130
cesium gluconate, 5 NaCl, 10 HEPES, 12 phosphocreatine di(tris) salt, 3 Mg*ATP, 0.2 Na*GTP, 1 EGTA,
2.0 Na2-ATP, 0.5 Na3-GTP, pH 7.3) using a Multiclamp 700B amplifier (Molecular Devices, Palo Alto, CA)
at 36 °C. Cells were visualized with dTomato and DIC with custom-built optics on a BX51WI microscope
(Olympus Optical, Tokyo, Japan) and recorded with pClamp 10.3 (Molecular Devices). Cell identity was
confirmed by the presence of fluorescence material in the patch pipet after membrane rupture and/or
cell fill with Alexa Fluor 488, and only cells that had a proximal dendrite that extended from the soma in
the direction of the EPL while remaining beneath the surface of the slice (i.e. cells that had dendrites
that did not appear to have been cut during slicing) were targeted for patching. Patch pipets had 8-11

MΩ open tip resistance. Series resistance was not compensated. Cells were recorded in continuous 10-
second sweeps for five minutes with a test pulse at the beginning of every sweep, which was used to
calculate series resistance and holding current in Matlab for initial quality control. sEPSCs were recorded
at -70 mV and sIPSCs were recorded at 0 mV. We waited at least one minute after breaking into the cell
before beginning recording for sEPSCs and at least 30 seconds after switching the holding potential for
sIPSCs; sEPSCs were always recorded first. Cells were recorded within 8 hours of slicing, and there did
not appear to be any relationship between the time of recording and the frequency of synaptic events
(data not shown). Experimenters were not blind to experimental group during recording.
Analysis
Experimenters were blind to experimental group during analysis. Cells with an initial series resistance of
<50 MΩ were used for analysis. Sweeps that deviated from the average series resistance across the first
three sweeps by more than 25% or had a holding current of more than 100 pA at -70 mV were excluded.
The cell was excluded entirely if less than half of the recording sweeps remained after quality control.
For each cell, separate test pulses (-10 mV, 20 ms) with 50 repetitions at 20 kHz were recorded before
and after each set of recordings, and these files were used to calculate series resistance (based on the
maximum current recorded at the beginning of the pulse), membrane resistance (based on the steady
state current during the last 20% of the pulse), and cell capacitance (based on the time constant of an
exponential fit between 20% and 80% of the current decay). Reported values in Figures 6 and 7
supplements are the means of these parameters from before and after the 5 minutes of sEPSC
recordings. For sEPSCs and sIPSCs, all sweeps were concatenated for each cell (excluding 0.5 s around
the test pulse), filtered with a 60Hz band-stop filter with five harmonics in Matlab, and exported into
Mini Analysis v. 6.0.7 (Synaptosoft). sEPSCs were detected with the following parameters: threshold 5,
period to search a local minimum 10000, time before a peak to baseline 15000, period to search a decay
time 20000, fraction of peak to find a decay time 0.37, period to average a baseline 1000, area threshold
20, number of points to average for peak 1, direction of peak "negative." After the initial detection step,
"Scan and detect double peaks" was selected. sIPSCs were detected with the following parameters:
threshold 8, period to search a local minimum 10000, time before a peak to baseline 6000, period to
search a decay time 20000, fraction of peak to find a decay time 0.37, period to average a baseline 1000,
area threshold 80, number of points to average for peak 1, direction of peak "positive." In both cases,
the detection was manually inspected, and the timepoints spanning any sections of the trace that
exhibited increased noise (typically due to fluctuations in seal quality that caused many obviously false
positive events) were noted. Event data was exported to a text file and imported into Matlab and noisy
sections were excluded before further analysis. Cells that had many noisy sections were excluded. This
included 3 control cells and 2 PLX cells for EPSCs and 2 control cells and 4 PLX cells for IPSCs (Figure 6) as
well as 3 control and 3 PLX cells for EPSCs and 8 control cells and 9 PLX cells for IPSCs (Figure 7). Events
that were less than 3 ms apart were considered doubly detected and excluded.
694
Statistical analysis
The number of mice to be used in each experiment was determined in advance, and all data that could
be obtained from these mice under the experimental conditions and quality standards described in each
section above were included in the analyses. To determine the number of mice for two photon imaging,
we took into account our previous success rate with combined cranial windows/virus injections (about
66%) and aimed to image a similar number of mice as in our previous work (4-7 mice for different
experiments, (Wallace et al., 2017)). For electrophysiology, we aimed to obtain data from at least 20

cells per group from 3-4 mice, consistent with other electrophysiological investigations in the field that
were able to reveal effects of various manipulations (Pallotto et al., 2012)(Quast et al., 2017).
Littermates were randomly allocated into control or PLX-treated groups for all experiments. Raw data is
shown in every figure, in the form of a scatterplot overlaid on a bar graph when N < 40 or as a
cumulative distribution otherwise. Non-parametric statistical tests were used for all comparisons, so no
assumption of normality of data was made. Details for each statistical test can be found in the
corresponding figure legend, and the rationale behind each is described here. The two-tailedWilcoxon
rank sum test was used to compare the medians of all unpaired non-normal distributions and
correspondingly, the medians are plotted with bars in the figures. A one-tailed permutation test was
used to test whether means in the real data were significantly higher than those in offset imaging data
in Figure 1. A two-tailed permutation test was used to test for differences in principal components for
the analysis of response timecourses in Figure 3—figure supplement 1 (details in section above "Two
photon imaging, Temporal dynamics"). The hierarchical bootstrap (Carpenter et al., 2003)(Saravanan et
al., 2019) was used to compare groups in the case of hierarchical datasets (details below in section
"Hierarchical bootstrap"). The statistical test and p value for each test is found in the legend
corresponding to each figure letter. All n values for each statistical test can be found at the end of each
figure legend. In the text, we report mean ± SEM for datasets where means were compared or median
(interquartile range) for datasets where medians were compared. For comparisons with hierarchical
bootstrap, we report sampling distribution mean ± sampling distribution standard deviation.
721
Hierarchical bootstrap
Sampling was performed 1,000 times for each dataset as described(Saravanan et al., 2019). Specifically,
a sample was taken with replacement across the units of the upper level. For example, in the two
photon imaging data, a sample was taken of the individual mice that were imaged (the sample size was
equal to the number of mice). Next, a sample was taken with replacement across the lower level, in this
case the dendrites that were imaged. We chose to sample an equal number of dendrites from each
mouse, regardless of the original number of dendrites imaged, to ensure that each animal would be
represented equally (since the number of dendrites originally imaged for each animal was related to
experimental considerations such as the density of viral labeling in each animal). For this level, we chose
a sample size (100 dendrites) slightly larger than the largest sample per unit in the original data to
ensure a representative sample would be obtained from each mouse. We then report the mean and the
standard error of the bootstrapped parameter (either mean, median, or proportion, which is stated in
each case) across the 1,000 samples. We calculate a p value directly, which represents the probability
that the parameter for the control group is greater than that for the PLX-treated group (for cases where
the parameter for the PLX-treated group is significantly greater, such as the proportion of dendrites that
were unresponsive in the imaging data, we report 1 – p to avoid confusion with standard p values). The
same procedure was performed to analyze datasets for spine morphology (upper level = dendrites,
lower level = spines, sample size = 100 spines) and electrophysiology (upper level = cells, lower level =
EPSCs or IPSCs, sample size = 5000 EPSCs or IPSCs).

References

Abraham NM, Egger V, Shimshek DR, Renden R, Fukunaga I, Sprengel R, Seeburg PH, Klugmann M, Margrie TW, Schaefer AT, Kuner T. 2010. Synaptic inhibition in the olfactory bulb accelerates odor discrimination in mice. *Neuron* **65**:399–411. doi:10.1016/j.neuron.2010.01.009

Allen M, Poggiali D, Whitaker K, Marshall TR, Kievit R. 2018. Raincloud plots: a multi-platform tool for robust data visualization. *PeerJ Prepr* **6**:e27137v1. doi:10.1044/1092-4388(2010/09-0205)

Arganda-Carreras I, Kaynig V, Rueden C, Eliceiri KW, Schindelin J, Cardona A, Seung HS. 2017. Trainable Weka Segmentation: A machine learning tool for microscopy pixel classification. *Bioinformatics* **33**:2424–2426. doi:10.1093/bioinformatics/btx180

Bolós M, Perea JR, Terreros-Roncal J, Pallas-Bazarra N, Jurado-Arjona J, Ávila J, Llorens-Martín M. 2018. Absence of microglial CX3CR1 impairs the synaptic integration of adult-born hippocampal granule neurons. *Brain Behav Immun* **68**:76–89. doi:10.1016/j.bbi.2017.10.002

Breton-Provencher V, Coté D, Saghatelyan A. 2014. Activity of the Principal Cells of the Olfactory Bulb Promotes a Structural Dynamic on the Distal Dendrites of Immature Adult-Born Granule Cells via Activation of NMDA Receptors. *J Neurosci* **34**:1748–59. doi:10.1523/JNEUROSCI.3013-13.2014

Breton-Provencher V, Lemasson M, Peralta MR, Saghatelyan A. 2009. Interneurons produced in adulthood are required for the normal functioning of the olfactory bulb network and for the execution of selected olfactory behaviors. *J Neurosci* **29**:15245–15257. doi:10.1523/JNEUROSCI.3606-09.2009

Bruttger J, Karram K, Wörtge S, Regen T, Marini F, Hoppmann N, Klein M, Blank T, Yona S, Wolf Y, Mack M, Pinteaux E, Müller W, Zipp F, Binder H, Bopp T, Prinz M, Jung S, Waisman A. 2015. Genetic Cell Ablation Reveals Clusters of Local Self-Renewing Microglia in the Mammalian Central Nervous System. *Immunity* 92–106. doi:10.1016/j.immuni.2015.06.012

Carleton A, Petreanu LT, Lansford R, Alvarez-Buylla A, Lledo P-M. 2003. Becoming a new neuron in the adult olfactory bulb. *Nat Neurosci* **6**:507–18. doi:10.1038/nn1048

Carpenter JR, Goldstein H, Rasbash J. 2003. A novel bootstrap procedure for assessing the relationship between class size and achievement. *J R Stat Soc Ser C Appl Stat* **52**:431–443. doi:10.1111/1467-9876.00415

Consiglio A, Gritti A, Dolcetta D, Follenzi A, Bordignon C, Gage FH, Vescovi AL, Naldini L. 2004. Robust in vivo gene transfer into adult mammalian neural stem cells by lentiviral vectors. *Proc Natl Acad Sci U S A* **101**:14835–14840. doi:10.1073/pnas.0404180101

Denizet M, Cotter L, Lledo P-M, Lazarini F. 2016. Sensory deprivation increases phagocytosis of adult-born neurons by activated microglia in the olfactory bulb. *Brain Behav Immun* 6–11. doi:10.1016/j.bbi.2016.09.015

Dombeck D, Tank D. 2014. Two-photon imaging of neural activity in awake mobile mice. *Cold Spring Harb Protoc* **2014**:726–736. doi:10.1101/pdb.top081810

Dubbs A, Guevara J, Yuste R. 2016. moco: Fast Motion Correction for Calcium Imaging. *Front Neuroinform* **10**:1–7. doi:10.3389/fninf.2016.00006

Dunn KW, Kamocka MM, McDonald JH. 2011. A practical guide to evaluating colocalization in biological microscopy. *Am J Physiol - Cell Physiol* **300**:723–742. doi:10.1152/ajpcell.00462.2010

Egger V. 2008. Synaptic sodium spikes trigger long-lasting depolarizations and slow calcium entry in rat
olfactory bulb granule cells. *Eur J Neurosci* **27**:2066–2075. doi:10.1111/j.1460-9568.2008.06170.x

Egger V, Svoboda K, Mainen ZF. 2005. Dendrodendritic synaptic signals in olfactory bulb granule cells:
local spine boost and global low-threshold spike. *J Neurosci* **25**:3521–3530.
doi:10.1523/JNEUROSCI.4746-04.2005

Egger V, Urban NN. 2006. Dynamic connectivity in the mitral cell-granule cell microcircuit. *Semin Cell Dev*
Biol **17**:424–432. doi:10.1016/j.semcdb.2006.04.006

Ekdahl CT. 2012. Microglial activation - tuning and pruning adult neurogenesis. *Front Pharmacol* **3**:41.
doi:10.3389/fphar.2012.00041

Ekdahl CT, Claasen J, Bonde S, Kokaia Z, Lindvall O. 2003. Inflammation is detrimental for neurogenesis
in adult brain **100**.

Elmore MRP, Najafi AR, Koike M a, Dagher NN, Spangenberg EE, Rice R a, Kitazawa M, Matusow B,
Nguyen H, West BL, Green KN. 2014. Colony-stimulating factor 1 receptor signaling is necessary for
microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron* **82**:380–97.
doi:10.1016/j.neuron.2014.02.040

Grimaud J, Murthy VN. 2018. How to monitor breathing in laboratory rodents: a review of the current
methods. *J Neurophysiol* **120**:624–632. doi:10.1152/jn.00708.2017

Gunner G, Cheadle L, Johnson KM, Ayata P, Badimon A, Mondo E, Nagy MA, Liu L, Bemiller SM, Kim K,
Lira SA, Lamb BT, Tapper AR, Ransohoff RM, Greenberg ME, Schaefer A, Schafer DP. 2019. Sensory
lesioning induces microglial synapse elimination via ADAM10 and fractalkine signaling. *Nat*
Neurosci **22**. doi:10.1038/s41593-019-0419-y

Guo Z V., Hires SA, Li N, O'Connor DH, Komiyama T, Ophir E, Huber D, Bonardi C, Morandell K, Gutnisky
D, Peron S, Xu NL, Cox J, Svoboda K. 2014. Procedures for behavioral experiments in head-fixed
mice. *PLoS One* **9**. doi:10.1371/journal.pone.0088678

Hammond TR, Robinton D, Stevens B. 2018. Microglia and the Brain: Complementary Partners in
Development and Disease. *Annu Rev Cell Dev Biol* **34**:523–544. doi:10.1146/annurev-cellbio-
100616-060509

Hioki H, Kuramoto E, Konno M, Kameda H, Takahashi Y, Nakano T, Nakamura KC, Kaneko T. 2009. High-
level transgene expression in neurons by lentivirus with Tet-Off system. *Neurosci Res* **63**:149–54.
doi:10.1016/j.neures.2008.10.010

Hong S, Dissing-Olesen L, Stevens B. 2016. New insights on the role of microglia in synaptic pruning in
health and disease. *Curr Opin Neurobiol* **36**:128–134. doi:10.1016/j.conb.2015.12.004

Hoshiko M, Arnoux I, Avignone E, Yamamoto N, Audinat E. 2012. Deficiency of the microglial receptor
CX3CR1 impairs postnatal functional development of thalamocortical synapses in the barrel cortex.
J Neurosci **32**:15106–11. doi:10.1523/JNEUROSCI.1167-12.2012

Jakubs K, Bonde S, Iosif RE, Ekdahl CT, Kokaia Z, Kokaia M, Lindvall O. 2008. Inflammation regulates
functional integration of neurons born in adult brain. *J Neurosci* **28**:12477–88.
doi:10.1523/JNEUROSCI.3240-08.2008

Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A, Littman DANR. 2000. Analysis of

Fractalkine Receptor CX 3 CR1 Function by Targeted Deletion and Green Fluorescent Protein
Reporter Gene Insertion **20**:4106–4114.

Kato HK, Chu MW, Isaacson JS, Komiyama T. 2012. Dynamic sensory representations in the olfactory
bulb: modulation by wakefulness and experience. *Neuron* **76**:962–75.
doi:10.1016/j.neuron.2012.09.037

Kelsch W, Lin C-W, Lois C. 2008. Sequential development of synapses in dendritic domains during adult
neurogenesis. *Proc Natl Acad Sci U S A* **105**:16803–8. doi:10.1073/pnas.0807970105

Kyle J, Wu M, Gourzi S, Tsirka SE. 2019. Proliferation and Differentiation in the Adult Subventricular Zone
Are Not Affected by CSF1R Inhibition. *Front Cell Neurosci* **13**:1–10. doi:10.3389/fncel.2019.00097

Lazarini F, Gabellec M-M, Torquet N, Lledo P-M. 2012. Early activation of microglia triggers long-lasting
impairment of adult neurogenesis in the olfactory bulb. *J Neurosci* **32**:3652–64.
doi:10.1523/JNEUROSCI.6394-11.2012

Lepousez G, Valley MT, Lledo P-M. 2013. The impact of adult neurogenesis on olfactory bulb circuits and
computations. *Annu Rev Physiol* **75**:339–63. doi:10.1146/annurev-physiol-030212-183731

Li Y, Du X-F, Liu C-S, Wen Z-L, Du J-L. 2012. Reciprocal regulation between resting microglial dynamics
and neuronal activity in vivo. *Dev Cell* **23**:1189–202. doi:10.1016/j.devcel.2012.10.027

Liddelow SA, Barres BA. 2017. Reactive Astrocytes: Production, Function, and Therapeutic Potential.
Immunity **46**:957–967. doi:10.1016/j.immuni.2017.06.006

Lin C-W, Sim S, Ainsworth A, Okada M, Kelsch W, Lois C. 2010. Genetically increased cell-intrinsic
excitability enhances neuronal integration into adult brain circuits. *Neuron* **65**:32–9.
doi:10.1016/j.neuron.2009.12.001

Livneh Y, Mizrahi A. 2012. Experience-dependent plasticity of mature adult-born neurons. *Nat Neurosci*
15:26–8. doi:10.1038/nn.2980

Miyamoto A, Wake H, Ishikawa AW, Eto K, Shibata K, Murakoshi H, Koizumi S, Moorhouse AJ, Yoshimura
Y, Nabekura J. 2016. Microglia contact induces synapse formation in developing somatosensory
cortex. *Nat Commun* **7**:12540. doi:10.1038/ncomms12540

Mizrahi A. 2007. Dendritic development and plasticity of adult-born neurons in the mouse olfactory
bulb. *Nat Neurosci* **10**:444–52. doi:10.1038/nn1875

Monje ML, Toda H, Palmer TD. 2003. Inflammatory blockade restores adult hippocampal neurogenesis.
Science **302**:1760–5. doi:10.1126/science.1088417

Nakayama H, Abe M, Morimoto C, Iida T, Okabe S, Sakimura K, Hashimoto K. 2018. Microglia permit
climbing fiber elimination by promoting GABAergic inhibition in the developing cerebellum. *Nat*
Commun **9**:1–2. doi:10.1038/s41467-018-05100-z

Nimmerjahn A, Kirchhoff F, Helmchen F. 2005. Resting microglial cells are highly dynamic surveillants of
brain parenchyma in vivo. *Science* **308**:1314–8. doi:10.1126/science.1110647

Nissant A, Bardy C, Katagiri H, Murray K, Lledo P-M. 2009. Adult neurogenesis promotes synaptic
plasticity in the olfactory bulb. *Nat Neurosci* **12**:728–30. doi:10.1038/nn.2298

Pallotto M, Nissant A, Fritschy J-M, Rudolph U, Sassoè-Pognetto M, Panzanelli P, Lledo P-M. 2012. Early

formation of GABAergic synapses governs the development of adult-born neurons in the olfactory
bulb. *J Neurosci* **32**:9103–15. doi:10.1523/JNEUROSCI.0214-12.2012

Paolicelli RC, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, Giustetto M, Ferreira TA, Guiducci E,
Dumas L, Ragozzino D, Gross CT. 2011. Synaptic pruning by microglia is necessary for normal brain
development. *Science* **333**:1456–8. doi:10.1126/science.1202529

Peng J, Liu Y, Umpierre AD, Xie M, Tian DS, Richardson JR, Wu LJ. 2019. Microglial P2Y12 receptor
regulates ventral hippocampal CA1 neuronal excitability and innate fear in mice. *Mol Brain* **12**:1–
10. doi:10.1186/s13041-019-0492-x

Quast KB, Ung K, Froudarakis E, Huang L, Herman I, Addison AP, Ortiz-Guzman J, Cordiner K, Saggau P,
Tolias AS, Arenkiel BR. 2017. Developmental broadening of inhibitory sensory maps. *Nat Neurosci*
20:189–199. doi:10.1038/nn.4467

Rall W, Shepherd GM, Reese TS, Brightman MW. 1966. Dendrodendritic synaptic pathway for inhibition
in the olfactory bulb. *Exp Neurol* **14**:44–56. doi:10.1016/0014-4886(66)90023-9

Reshef R, Kudryavitskaya E, Shani-Narkiss H, Isaacson B, Rimmerman N, Mizrahi A, Yirmiya R. 2017. The
role of microglia and their CX3CR1 signaling in adult neurogenesis in the olfactory bulb. *Elife* **6**:1–
30. doi:10.7554/eLife.30809

Ribeiro Xavier a. L, Kress BT, Goldman S a., Lacerda de Menezes JR, Nedergaard M. 2015. A Distinct
Population of Microglia Supports Adult Neurogenesis in the Subventricular Zone. *J Neurosci*
35:11848–11861. doi:10.1523/JNEUROSCI.1217-15.2015

Rodríguez-Iglesias N, Sierra A, Valero J. 2019. Rewiring of Memory Circuits: Connecting Adult Newborn
Neurons With the Help of Microglia. *Front Cell Dev Biol* **7**:1–20. doi:10.3389/fcell.2019.00024

Sailor KA, Valley MT, Wiechert MT, Riecke H, Sun GJ, Adams W, Dennis JC, Sharafi S, Ming G, Song H,
Lledo P-M. 2016. Persistent Structural Plasticity Optimizes Sensory Information Processing in the
Olfactory Bulb. *Neuron* **91**:384–396. doi:10.1016/j.neuron.2016.06.004

Salter MW, Beggs S. 2014. Sublime Microglia: Expanding Roles for the Guardians of the CNS. *Cell*
158:15–24. doi:10.1016/j.cell.2014.06.008

Saravanan V, Berman GJ, Sober SJ. 2019. Application of the hierarchical bootstrap to multi-level data in
neuroscience. *bioRxiv*. doi:10.1101/819334

Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, Ransohoff RM, Greenberg
ME, Barres B a, Stevens B. 2012. Microglia sculpt postnatal neural circuits in an activity and
complement-dependent manner. *Neuron* **74**:691–705. doi:10.1016/j.neuron.2012.03.026

Sierra A, Encinas JM, Deudero JJP, Chancey JH, Enikolopov G, Overstreet-Wadiche LS, Tsirka SE, Maletic-
Savatic M. 2010. Microglia shape adult hippocampal neurogenesis through apoptosis-coupled
phagocytosis. *Cell Stem Cell* **7**:483–95. doi:10.1016/j.stem.2010.08.014

Sipe GO, Lowery, RL, Tremblay M-È, Kelly EA, Lamantia CE, Majewska AK. 2016. Microglial P2Y12 is
necessary for synaptic plasticity in mouse visual cortex. *Nat Commun* **7**:10905.
doi:10.1038/ncomms10905

Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N, Micheva KD, Mehalow AK,
Huberman AD, Stafford B, Sher A, Litke AM, Lambris JD, Smith SJ, John SWM, Barres B a. 2007. The

classical complement cascade mediates CNS synapse elimination. *Cell* **131**:1164–78.
doi:10.1016/j.cell.2007.10.036

Torres L, Danver J, Ji K, Miyauchi JT, Chen D, Anderson ME, West BL, Robinson JK, Tsirka SE. 2016. Brain ,
Behavior , and Immunity Dynamic microglial modulation of spatial learning and social behavior.
Brain Behav Immun **55**:6–16. doi:10.1016/j.bbi.2015.09.001

Tremblay M-È, Lowery RL, Majewska AK. 2010. Microglial interactions with synapses are modulated by
visual experience. *PLoS Biol* **8**:e1000527. doi:10.1371/journal.pbio.1000527

Tremblay M-È, Stevens B, Sierra A, Wake H, Bessis A, Nimmerjahn A. 2011. The role of microglia in the
healthy brain. *J Neurosci* **31**:16064–9. doi:10.1523/JNEUROSCI.4158-11.2011

Wake H, Moorhouse AJ, Jinno S, Kohsaka S, Nabekura J. 2009. Resting microglia directly monitor the
functional state of synapses in vivo and determine the fate of ischemic terminals. *J Neurosci*
29:3974–80. doi:10.1523/JNEUROSCI.4363-08.2009

Wallace JL, Wienisch M, Murthy VN. 2017. Development and Refinement of Functional Properties of
Adult-Born Neurons. *Neuron* **96**:883–896. doi:10.1016/j.neuron.2017.09.039

Weinhard L, Di Bartolomei G, Bolasco G, Machado P, Schieber NL, Neniskyte U, Exiga M, Vadisiute A,
Raggioli A, Schertel A, Schwab Y, Gross CT. 2018. Microglia remodel synapses by presynaptic
trogocytosis and spine head filopodia induction. *Nat Commun* **9**. doi:10.1038/s41467-018-03566-5

Whitman MC, Greer C a. 2007. Synaptic integration of adult-generated olfactory bulb granule cells: basal
axodendritic centrifugal input precedes apical dendrodendritic local circuits. *J Neurosci* **27**:9951–
61. doi:10.1523/JNEUROSCI.1633-07.2007

Wienisch M, Murthy VN. 2016. Population imaging at subcellular resolution supports specific and local
inhibition by granule cells in the olfactory bulb. *Sci Rep* **6**:29308. doi:10.1038/srep29308

Willmore B, Tolhurst DJ. 2001. Characterizing the sparseness of neural codes. *Network* **12**:255–270.
doi:10.1088/0954-898X/12/3/302

Wu Y, Dissing-Olesen L, MacVicar BA, Stevens B. 2015. Microglia: Dynamic Mediators of Synapse
Development and Plasticity. *Trends Immunol* **36**:605–613. doi:10.1016/j.it.2015.08.008

Zhan Y, Paolicelli RC, Sforazzini F, Weinhard L, Bolasco G, Pagani F, Vyssotski AL, Bifone A, Gozzi A,
Ragozzino D, Gross CT. 2014. Deficient neuron-microglia signaling results in impaired functional
brain connectivity and social behavior. *Nat Neurosci* **17**:400–6. doi:10.1038/nn.3641

Ziv Y, Ron N, Butovsky O, Landa G, Sudai E, Greenberg N, Cohen H, Kipnis J, Schwartz M. 2006. Immune
cells contribute to the maintenance of neurogenesis and spatial learning abilities in adulthood. *Nat*
Neurosci **9**:268–275. doi:10.1038/nn1629

Figure 1

Figure 1: Microglia preferentially interact with mushroom spines and filopodial spines on developing abGCs

(A) Maximum intensity projection (10 μm volume at the first imaging timepoint) showing dTomato-labeled abGCs in a CX3CR1-GFP heterozygous mouse imaged 4 weeks after lentivirus injection. Brightness and contrast adjusted for display only.

(B) Single plane time series showing the region marked in (A). Inset shows the calculated percent microglial coverage for the spine marked with the arrowhead (images shown for 6 frames, 7th frame not shown but the value is plotted, showing the end of the interaction) with the larger circle marking the value for the corresponding frame. Brightness and contrast adjusted with the same parameters for each timepoint for display only.

(C) Single plane image showing an example of a spine classified as a mushroom spine because the spine head is wider and brighter than the spine neck.

(D) Probability distribution showing the proportion of mushroom spines with at least one microglial interaction for mushroom spines in the real data (black line, "Data") compared to values calculated from iteratively translating the microglial channel relative to the dendritic imaging channel (gray histogram, "Offsets"). The proportion of spines with interactions was not significantly higher than chance (one-tailed permutation test, $p = 0.13$).

(E) Probability distribution showing the number of interactions (normalized to 10 minutes) for mushroom spines. The mean number of interactions (value for each dendrite is the mean number across all mushroom spines on that dendrite) in the real data was significantly higher than chance (one-tailed permutation test, $p = 0.048$).

(F) Probability distribution showing interaction length for mushroom spines (for spines that had at least one frame that met the criteria for an interaction, see Methods). The mean interaction length across all dendrites (value for each dendrite is the mean interaction length across all interactions for all mushroom spines) was not higher than chance (one-tailed permutation test, $p = 0.96$).

(G) Probability distribution showing maximum percent coverage (mean across all interactions for a given spine for spines that had at least one frame that met the criteria for an interaction). The mean maximum percent coverage across all dendrites (value for each dendrite is the mean interaction length across all interactions for all mushroom spines) was not higher in the real data (one-tailed permutation test, $p = 0.72$).

(H) Single plane image of a spine classified as a filopodial spine because it has no clear spine head.

(I) Probability distribution showing the proportion of spines with at least one microglial interaction for filopodial spines. The proportion of filopodial spines with interactions was not significantly higher than chance (one-tailed permutation test, $p = 0.39$).

(J) Probability distribution showing the number of interactions (normalized to 10 minutes) for filopodial spines. The mean number of interactions (value for each dendrite is the mean number across all filopodial spines on that dendrite) in the real data was not significantly higher than chance (one-tailed permutation test, $p = 0.72$).

(K) Probability distribution showing interaction length for filopodial spines (for spines that had at least one frame that met the criteria for an interaction, see Methods). The mean interaction length across all dendrites (value for each dendrite is the mean interaction length across all interactions for all filopodial spines) was not significantly higher than chance (one-tailed permutation test, $p = 0.23$).

(L) Probability distribution showing maximum percent coverage (mean across all interactions for a given spine for spines that had at least one frame that met the criteria for an interaction). The mean maximum percent coverage across all dendrites (value for each dendrite is the mean interaction length across all interactions for all filopodial spines) was not higher in the real data (one-tailed permutation test, $p = 0.84$).

n = 726 spines (271 mushroom spines and 455 filopodial spines) from 48 dendrites combined at 1, 2, 3, and 4 weeks post injection in 3 mice

ns, not significant; *$p<0.05$

Figure 2

Figure 2: Microglia ablation during development reduces odor-evoked responses of abGCs in anesthetized mice.

(A) Experimental timeline for microglia ablation during development of abGCs. A cranial window was implanted and 3 weeks later mice were given control or PLX5622-containing chow for the remainder of the experiment. After 3 weeks of chow consumption, a lentivirus was injected into the RMS to label abGCs, which were imaged 5-6 weeks later.

(B) Left, images of Iba1 staining in the olfactory bulb of control (top) and PLX-treated (bottom) mice. White squares show the locations of the enlarged insets. Dotted lines mark the upper edge of the glomerular and granule cell layers. Right, schematic showing injection of a lentivirus encoding dTomato and GCaMP6s and microglia ablation.

(C) Example fields of view showing an average intensity projection of dTomato structural images of abGC dendrites (left) and overlaid heatmaps of GCaMP6s-recorded activity (right) in response to ethyl valerate in control (top) and PLX5622-treated (bottom) mice.

(D) GCaMP6s traces showing odor responses of example ROIs from control (top) and PLX-treated (bottom) mice (chosen to have the same ranked response to the first odor). Gray traces represent responses on individual trials and colored trace is the mean across trials. Individual trial traces were median filtered over three frames before averaging for presentation. *, odor responses for which the mean response was above threshold

(E) Heatmap traces from the 100 ROIs with the largest odor-evoked Ca^{2+} signals across all mice ranked separately for each of 15 odors (molecular structures shown above). Black bar denotes odor time. Bottom, mean response time course for each odor across all ROIs.

(F) Cumulative distribution showing that the distribution of responses (averaged across odors for each dendrite) is shifted to the left in PLX-treated mice (Two sample Kolmogorov–Smirnov test for probability distributions, $D = 0.25$, $p = 2.56e-08$) while the noise distributions constructed from blank trials are not different ($D = 0.042$, $p = 0.96$).

(G) Cumulative distribution showing the number of effective odors (odors that evoked responses above the ROC threshold 0.39, which was calculated across all dendrites from both groups). The median number of effective odors was significantly lower in the PLX-treated group (Wilcoxon rank sum test, $z = 3.86$, $p = 1.15e-04$).

(H) Raincloud plot showing the distribution of lifetime sparseness across all dendrites. Above, kernel density estimate. Below, boxplot showing the median, interquartile range (box), and 1.5 times the interquartile range (whiskers) superimposed on a dot plot of all the data (one dot per dendrite). Median lifetime sparseness was significantly lower in the PLX-treated group (Wilcoxon rank sum test, $z = 3.53$, $p = 4.18e-04$).

n = 287 dendrites from 5 control mice and 277 dendrites from 7 PLX-treated mice

*$p<0.05$, **$p<0.01$, ***$p <0.001$

Figure 2.1

Figure 2—figure supplement 1

(A) Microglia cells were gated on CD45^{intermediate} and CD11b^{high} using the following gating strategy: Debris and dead cells were excluded based on the fluorescent intensity of a dead cell stain. Next, single cells where determined based on of their light scattering properties (first by side over forward scatter, SSC-A/FSC-A, and second by forward scatter height over area, FSC-H/FSC-A). The counting beads were identified based on their fluorescent emission at 594 nm.

(B) Plot showing the percent of live microglia remaining (normalized to the mean of the controls) in mice treated with PLX compared to controls (number of microglia in each sample normalized to

counting beads) in whole brain samples after 1 week of control or PLX diet demonstrating 97.4% ablation.

(C) Plot showing the percent of live microglia remaining in mice treated with PLX (normalized to the mean of the controls) compared to controls (number of microglia in each sample normalized to counting beads) in whole brain samples after 1 week of control or PLX diet, demonstrating 99.0% ablation.

(D) Plot showing the percent of live microglia remaining in mice treated with PLX (normalized to the mean of the controls) compared to controls measured via flow cytometry in olfactory bulb samples after 2 weeks of control or PLX diet, demonstrating 96.4% ablation.

(E) Interrogation of light scattering properties (SSC-A/FSC-A) independent of surface marker labeling revealed that hardly any cells remained in the microglia enriched samples after treatment with PLX compared to control samples. This becomes visually more apparent when microglia cells (CD45^{intermediate}, CD11b^{high}, control and PLX-treated plots on left) are backgated to SSC-A/FSC-A plots on right (in dark blue). Note that counting beads can be observed in red.

Bars represent the mean across mice (circles).

Whole brain: n = 5 control and 5 PLX mice

OB: n = 2 control and 2 PLX samples (each sample contained both OBs from 2 littermates which were combined after dissection)

Figure 2.2

Figure 2—figure supplement 2
- (A) Maximum intensity projection showing microglia stained with anti-Iba1 in the granule cell layer of the olfactory bulb in control (left) and PLX-treated (right) mice after 4 weeks of treatment.
- (B) Plot showing the percent of microglia remaining in mice treated with PLX compared to control littermates based on counting of cell bodies stained with Iba1 after 1 week of treatment, demonstrating 84.7% ablation.
- (C) Percent microglia remaining after 4 weeks of treatment (same mice used for spine quantification in Figure 5), demonstrating 91.7% ablation.
- (D) Percent microglia remaining after 9 weeks of treatment (PLX mice are the same mice used for imaging at the 9 weeks timepoint in Figure 4 Supplement 1 and control mice are age-matched controls), demonstrating 86.2% ablation.

Bars represent the mean across mice (circles).

n = 3 mice (1 week), 4 mice (4 weeks), and 2 mice (9 weeks)

Figure 2.3

Figure 2—figure supplement 3

    (A) Maximum intensity projection astrocytes stained with anti-GFAP in the granule cell layer of the olfactory bulb in control mice 1 week (left) and 4 weeks (right) after beginning treatment.

(B) Maximum intensity projection astrocytes stained with anti-GFAP in the granule cell layer of the olfactory bulb in PLX-treated mice 1 week (left) and 4 weeks (right) after beginning treatment.

(C) Plot showing the number of astrocytes counted in control versus PLX-treated mice 1 week after the start of treatment. The median numbers were not significantly different between groups (Wilcoxon rank sum test, rank sum = 12, p = 0.70).

(D) Plot showing the number of astrocytes counted in control versus PLX-treated mice 4 weeks after the start of treatment. The median numbers were not significantly different between groups (Wilcoxon rank sum test, rank sum = 11, p = 0.86).

Bars represent the median across mice (circles).

n = 3 control versus 3 PLX-treated mice (1 week) and 3 control versus 4 PLX-treated mice (4 weeks)

ns, not significant

Figure 2.4

Figure 2—figure supplement 4

(A) Maximum intensity projection showing NeuN staining (left) and BrdU staining (right) in the olfactory bulb. Dotted line indicates boundary of the granule cell layer, where positive cells were quantified.

(B) Plot showing the density of BrdU+/NeuN+ cells per mm^3 in the granule cell layer of the olfactory bulb in control mice and mice treated with PLX for 4 weeks, beginning 3 days after BrdU injection. The density was not different between groups (Wilcoxon rank sum test, rank sum = 35, p = 0.15).

Bars represent the median across mice (circles).

n = 5 control and 5 PLX-treated mice

ns, not significant

Figure 2.5

Figure 2—figure supplement 5

(A) Sampling distributions of the mean response amplitude obtained with the hierarchical bootstrap. The mean amplitude (estimated by the mean of the sampling distribution, dark line) were not significantly different between dendrites in control versus PLX-treated mice (hierarchical bootstrap, p = 0.21).

(B) Sampling distributions of the mean number of effective odors (of the 15-odor panel) evoking a significant response. The mean number of effective odors was significantly greater in control than PLX-treated mice (hierarchical bootstrap, p = 0.050).

(C) Sampling distributions of the mean lifetime sparseness. The mean lifetime sparseness was significantly greater in control than PLX-treated mice (hierarchical bootstrap, p = 0.023).

(D) Sampling distributions of the median response amplitude. The median amplitude was significantly higher in control than PLX-treated mice (hierarchical bootstrap, p = 0.0011).

(E) Sampling distributions of the proportion of dendrites that did not respond to any of the odors. The mean proportion of unresponsive dendrites was significantly greater in PLX-treated mice (hierarchical bootstrap, p = 0.0013).

(F) Sampling distributions of the median response amplitude, only considering responses that were above the threshold. The median response amplitude above threshold was not different between groups (hierarchical bootstrap, p = 0.77).

The means of the sampling distributions are shown as dark lines and were used to estimate the value of the parameter (x-axis label) for each group.

n = 287 dendrites from 5 control mice and 277 dendrites from 7 PLX-treated mice

ns, not significant; *p<0.05, **p<0.01

Figure 3

Figure 3: Microglia ablation during development reduces odor-evoked responses of abGCs in awake mice.

(A) Above, Heatmap traces from the 100 ROIs with the largest odor-evoked Ca2+ signals across all mice ranked for each of 15 odors (molecular structures shown above). Below, Mean response time course for each odor across all ROIs. Black bar denotes odor time.

(B) Cumulative distribution showing that the distribution of responses (averaged across odors for each dendrite) is shifted to the left in PLX-treated mice (Two sample Kolmogorov–Smirnov test for probability distributions, D = 0.18, p = 0.037) while the noise distributions constructed from blank trials are not different (D = 0.11, p = 0.45).

(C) Cumulative distribution showing the number of effective odors (odors that evoked responses above the ROC threshold 0.52, which was calculated across all dendrites from both groups). There was a trend toward a lower median number of effective odors in the PLX-treated group (Wilcoxon rank sum test, z = 1.95, p = 0.052).

(D) Raincloud plot showing the distribution of lifetime sparseness across all dendrite. Above, kernel density estimate. Below, boxplot showing the median, interquartile range (box), and 1.5 times the interquartile range (whiskers) superimposed on a dot plot of all the data (one dot per dendrite). There was a trend toward lower median lifetime sparseness in the PLX-treated group (Wilcoxon rank sum test, z = 1.91, p = 0.056).

n = 105 dendrites from 3 control mice and 132 dendrites from 4 PLX-treated mice

*p<0.05

Figure 3.1

Figure 3—figure supplement 1

(A) Top, the first three principal components (PCs) obtained from PCA of the temporal profiles of responses of all abGC dendrites ($\Delta F/F_\sigma$ traces, GCaMP6s) in anesthetized control mice to all odors. The x-axis represents time, and the y-axis represents the magnitude of $\Delta F/F_\sigma$ for each PC. Percent variance explained: 36.3%, 4.19%, 1.85% for the first three PCs, respectively. Bottom, The first three PCs for odor responses in anesthetized PLX-treated mice. Percent variance explained: 33.7%, 3.82%, 1.96%. Shading indicates the odor presentation period (2 s). The timecourses of responses for the odor analysis period (5 s after odor onset) for the first three PCs was not different between control and PLX-treated mice (permutation test, p = 0.41).

(B) Top, the first three PCs for odor responses in awake control mice. Percent variance explained: 43.7%, 12.6%, 3.36%. Bottom, the first three PCs for odor responses in awake PLX-treated mice. Percent variance explained: 44.9%, 10.2%, 2.86%. The timecourses of responses for the odor

analysis period for the first three PCs were significantly different between control and PLX-treated mice (permutation test, p = 0.011).

(C) Raincloud plot showing the distribution of coefficient of variation values for all dendrites in anesthetized mice averaged across all odors for which the dendrite had a mean response above threshold (ROC thresholds calculated separately for control (threshold = 0.37) and PLX (threshold = 0.5) to ensure the same proportion of true responses). Above, kernel density estimate. Below, boxplot showing the median, interquartile range (box), and 1.5 times the interquartile range (whiskers) superimposed on a dot plot of all the data (one dot per cell, unless the cell had no responses above threshold in which case it is not includeD). The median CVs were not significantly different (Control: 0.59, PLX: 0.62, Wilcoxon rank sum test, z = 0.58, p = 0.56).

(D) Raincloud plot showing the distribution of coefficient of variation values for all dendrites in awake mice averaged across all odors for which the dendrite had a response above threshold (ROC thresholds calculated separately for control (threshold = 0.46) and PLX (threshold = 0.85) to ensure the same proportion of true responses). The median CVs were not significantly different (Control: 0.55, PLX: 0.48, Wilcoxon rank sum test, z = 1.47, p = 0.14).

(E) Cumulative distribution (responses averaged across odors for each dendrite) showing that the distribution of responses detected with event analysis (ROC threshold calculated across both groups combined, event detected during the odor analysis period if 6 or more frames were at least 2.6 standard deviations above baseline) is not significantly different in PLX-treated mice (Two sample Kolmogorov–Smirnov test for probability distributions, D = 0.18, p = 0.23).

(F) Cumulative distribution showing the number of effective odors (odors for which an event was detected in at least one repetition). The median number of effective odors was lower in the PLX-treated group (Wilcoxon rank sum test, z = 2.09, p = 0.037).

(G) Cumulative distribution showing that the latency from odor onset to event detection (averaged across odors for each dendrite) is not significantly different in PLX-treated mice (Two sample Kolmogorov–Smirnov test for probability distributions, D = 0.17, p = 0.31).

ns, not significant; *p<0.05

Figure 3.2

Figure 3—figure supplement 2

 (A) Example trace showing breathing recordings during one trial from a control mouse. The gray bar indicates the time the odor is on and the black circles indicate sniffs that were detected.

 (B) Example trace showing breathing recordings during one trial from a PLX-treated mouse.

 (C) Mean breathing rate during baseline periods (10 s before the odor comes on) and during the odor period (2 s) calculated from all sniffs recorded in all trials from each mouse. The breathing rates in the two groups were not different during the baseline period (Two sample t test, t = - 0.11, p = 0.91) or the odor period (Two sample t test, t = 0.25, p = 0.81).

Bars represent the mean across mice (circles).

n = 3 control mice (Baseline: 15,086 detected sniffs, Odor: 2,895 detected sniffs) and 4 PLX-treated mice (Baseline: 19,248 detected sniffs, Odor: 3,468 detected sniffs)

ns, not significant

Figure 4

Figure 4: Microglia ablation after development has no effect on odor-evoked responses of abGCs

(A) Experimental timeline for microglia ablation after development of abGCs. AbGCs were labeled via lentivirus injection and allowed to mature for 3 months. A control imaging session was performed immediately before administration of PLX chow and the second imaging session occurred 3 weeks later.

(B) Heatmap traces from the 100 ROIs with the largest odor-evoked Ca2+ signals across all mice ranked for each of 15 odors (molecular structures shown above). Black bar denotes odor time. Bottom, mean response time course for each odor across all ROIs.

(C) Cumulative distribution showing that the distribution of responses (averaged across odors for each dendrite) is not different before and after PLX diet administration (Two sample Kolmogorov–Smirnov test, D = 0.087, p = 0.45) and the noise distributions constructed from blank trials are also not different (D = 0.11, p = 0.16).

(D) Cumulative distribution showing the number of effective odors (odors that evoked responses above the ROC threshold 0.53, which was calculated across all dendrites from both groups). The median number of effective odors was not different between groups (Wilcoxon rank sum test, z = 0.0038, p = 1.00).

(E) Raincloud plot showing the distribution of lifetime sparseness across all dendrite. Above, kernel density estimate. Below, boxplot showing the median, interquartile range (box), and 1.5 times the interquartile range (whiskers) superimposed on a dot plot of all the data (one dot per dendrite). Median lifetime sparseness was not different between groups (Wilcoxon rank sum test, z = 0.18, p = 0.87).

n = 198 dendrites before and 185 dendrites after 3 weeks of PLX administration from 3 mice

ns, not significant

Figure 4.1

Figure 4—figure supplement 1

   (A) Experimental timeline for microglia ablation after development of abGCs. AbGCs were labeled via lentivirus injection and allowed to mature for 3 months. One imaging session was performed and a second was performed 3 weeks later in which the same dendrites were imaged again (the second is the same "Control" imaging session from the main figure).

   (B) Scatterplot showing the responses of all dendrite-odor pairs for the first and second imaging sessions. Dotted line is unity. Linear correlation coefficient $R^2 = 0.73$, p = 0.

   (C) Cumulative distribution showing that the distribution of responses (averaged across odors for each dendrite) is not different between the two imaging sessions (Two sample Kolmogorov–Smirnov test for probability distributions, D = 0.097, p = 0.61) and the noise distributions constructed from blank trials are also not different (D = 0.11, p = 0.48).

   (D) Cumulative distribution showing the number of effective odors (odors that evoked responses above the ROC threshold 0.39, which was calculated across all dendrites from both groups). The median number of effective odors was not different between groups (Wilcoxon rank sum test, z = 0.20, p = 0.84).

   (E) Raincloud plot showing the distribution of lifetime sparseness for all dendrites. Above, kernel density estimate. Below, boxplot showing the median, interquartile range (box), and 1.5 times the interquartile range (whiskers) superimposed on a dot plot of all the data (one dot per dendrite). The median lifetime sparseness values were not significantly different (Wilcoxon rank sum test, z = 0.30, p = 0.76).

n = 121 dendrites from 3 mice (same mice from the main figure)

ns, not significant

Figure 4.2

Figure 4—figure supplement 2

- (A) Experimental timeline for microglia ablation after development of abGCs. AbGCs were labeled via lentivirus injection and allowed to mature for 3 months. A control imaging session was performed immediately before administration of PLX chow and the second imaging session occurred 9 weeks later.
- (B) Heatmap traces from the 100 ROIs with the largest odor-evoked Ca2+ signals across all mice ranked for each of 15 odors (molecular structures shown above). Black bar denotes odor time. Bottom, mean response time course for each odor across all ROIs.
- (C) Cumulative distribution showing that the distribution of responses (averaged across odors for each dendrite) is not different before and after PLX diet administration (Two sample Kolmogorov–Smirnov test for probability distributions, D = 0.089, p = 0.61) and the noise distributions constructed from blank trials are also not different (D = 0.14, p = 0.095).
- (D) Cumulative distribution showing the number of effective odors (odors that evoked responses above the ROC threshold 0.78, which was calculated across all dendrites from both groups). The

median number of effective odors was not different between groups (Wilcoxon rank sum test, z = -0.17, p = 0.87).

(E) Raincloud plot showing the distribution of lifetime sparseness for all dendrites in control and PLX-treated mice. Above, kernel density estimate. Below, boxplot showing the median, interquartile range (box), and 1.5 times the interquartile range (whiskers) superimposed on a dot plot of all the data (one dot per dendrite). Median lifetime sparseness was not different between groups (Wilcoxon rank sum test z = -0.26, p = 0.79).

n = 168 dendrites before and 122 dendrites after 9 weeks of PLX administration from 3 mice (1 mouse was also included in the 3 weeks group that is shown in the main figure)

ns, not significant

Figure 5

Figure 5: Microglia ablation during development reduces spine head volume in abGCs

    (A) Experimental timeline for microglia ablation during development of abGCs. Mice were given control or PLX5622-containing chow on the same day that a lentivirus was injected into the RMS to label abGCs. Spine numbers and morphology were quantified 4 weeks later.

    (B) Above, sample images showing two apical dendrites from one cell that were analyzed in a control mouse (left) and PLX-treated mouse (right). Below, insets from the images shown above, showing spine morphology in more detail.

    (C) Spine density averaged across 1-5 apical dendrites from each abGC. Spine density was higher in PLX-treated mice (Wilcoxon rank sum test, z = -2.07, p = 0.039). Bars indicate medians across cells (circles).

(D) Cumulative distribution showing the volume of all spines analyzed. Inset, sampling distributions of the mean head volume obtained with the hierarchical bootstrap. The mean head volume (dark lines) was greater in control versus PLX-treated mice (hierarchical bootstrap, p = 0.057).

n = 810 spines from 17 abGCs from 3 control mice and 1551 spines from 17 abGCs from 4 PLX mice

*p<0.05

Figure 5.1

Figure 5—figure supplement 1

- (A) Spine density for filopodial spines (defined as spines with spine head width less than 1.5 times the mean width of the spine neck). Spine density was higher in PLX-treated mice (Wilcoxon rank sum test, z = -2.69, p = 0.0072). Bars indicate medians across cells (circles).
- (B) Spine density for mushroom spines (defined as spines with the width of the spine head greater than or equal to 1.5 times greater than the mean width of the spine neck). Spine density was higher in PLX-treated mice (Wilcoxon rank sum test, z = -1.96, p = 0.050). Bars indicate medians across cells (circles).
- (C) Cumulative distribution showing filopodial spine volume. Inset, sampling distributions of the mean filopodial head volume obtained with the hierarchical bootstrap. The mean head volume (dark lines) was not different between groups (hierarchical bootstrap, p = 0.72)
- (D) Cumulative distribution showing mushroom spine volume. Inset, sampling distributions of the mean mushroom head volume obtained with the hierarchical bootstrap. The mean head volume (dark lines) was significantly larger in control mice (hierarchical bootstrap, p = 0.036).

n = 810 spines from 17 abGCs from 3 control mice and 1551 spines from 17 abGCs from 4 PLX mice

ns, not significant; *p<0.05; **p<0.01

Figure 6

Figure 6: Microglia ablation during abGC development reduces the amplitude of excitatory synaptic currents but does not affect inhibitory synaptic currents

- (A) Experimental timeline for electrophysiological recording in abGCs
- (B) Left, sample sections from raw traces recorded from abGCs in control (top) and PLX-treated (bottom) mice. Right, median EPSCs across all EPSCs detected from all mice.
- (C) Cumulative distribution showing the inter-event intervals from all recorded EPSCs. Inset, sampling distributions of the median frequency obtained with the hierarchical bootstrap. The median frequencies (dark lines) were not significantly different (hierarchical bootstrap, p = 0.48).
- (D) Cumulative distribution showing the amplitudes from all recorded EPSCs. Inset, sampling distributions of the median amplitude obtained with the hierarchical bootstrap. The median amplitude (dark line) was significantly higher in control cells (hierarchical bootstrap, p = 0.0068).
- (E) Left, Sample sections from raw traces recorded from abGCs in control (top) and PLX-treated (bottom) mice. Right, median IPSCs across all IPSCs detected from all mice.
- (F) Cumulative distribution showing the inter-event intervals from all recorded IPSCs. Inset, sampling distributions of the median frequency obtained with the hierarchical bootstrap. The median frequencies (dark lines) were not significantly different (hierarchical bootstrap, p = 0.77).

(G) Cumulative distribution showing the amplitudes from all recorded IPSCs. Inset, sampling distributions of the median amplitude obtained with the hierarchical bootstrap. The median amplitudes (dark lines) were not significantly different (hierarchical bootstrap, $p = 0.79$).

For EPSCs: n = 30 abGCs from 4 control mice and 33 abGCs from 4 PLX mice

For IPSCs: n = 29 abGCs from 4 control mice and 30 abGCs from 4 PLX mice (same mice in both cases and cells used for both EPSCs and IPSCs if the recordings met criteria stated in Methods)

ns, not significant; **$p<0.01$

Figure 6.1

Figure 6—figure supplement 1

(A) Membrane resistance (mean of measurements taken before and after EPSC recordings) in abGCs. There was no difference between the two groups (Wilcoxon rank sum test, z = 1.01, p = 0.31).

(B) Membrane capacitance (mean of measurements taken before and after EPSC recordings) in abGCs. There was no difference between the two groups (Wilcoxon rank sum test, z = 0.52, p = 0.61).

(C) Series resistance (mean of measurements taken before and after EPSC recordings) during abGC recordings. There was no difference between the two groups (Wilcoxon rank sum test, z = 0.048, p = 0.96).

(D) Distance of the cell body from the mitral cell layer (measured in 2D, not taking into account depth from the surface of the slice) for all cells recorded. Both superficial and deep abGCs were included in the dataset and there was no difference in the mean distance to the mitral cell layer between the two groups (Wilcoxon rank sum test, z = -0.52, p = 0.60).

Bars indicate medians across cells (circles).

n = 30 abGCs from 4 control mice and 33 abGCs from 4 PLX mice

ns, not significant

Figure 7

Figure 7: Microglia ablation after abGC development has no effect on excitatory synaptic currents

    (A) Experimental timeline for electrophysiological recordings in abGCs after their development

    (B) Left, Sample sections from raw traces recorded from abGCs in control (top) and PLX-treated (bottom) mice. Right, median EPSCs detected from all mice.

    (C) Cumulative distribution showing the inter-event intervals from all recorded EPSCs. Inset, sampling distributions of the mean frequency obtained with the hierarchical bootstrap. The mean frequencies (dark lines) were not significantly different (hierarchical bootstrap, p = 0.76).

    (D) Cumulative distribution showing the amplitude from all recorded EPSCs. Inset, sampling distributions of the mean frequency obtained with the hierarchical bootstrap. The mean amplitudes (dark lines) were not significantly different (hierarchical bootstrap, p = 0.093).

n = 23 abGCs from 3 control mice and 27 abGCs from 3 PLX mice

ns, not significant

Figure 7.1

Figure 7—figure supplement 1
- (A) Left, Sample sections from raw traces recorded from abGCs in control (top) and PLX-treated (bottom) mice. Right, median IPSCs calculated across all IPSCs detected from all mice.
- (B) Cumulative distribution showing the inter-event intervals from all recorded IPSCs. Inset, sampling distributions of the mean frequency obtained with the hierarchical bootstrap. The mean frequencies (dark lines) were not significantly different (hierarchical bootstrap, p = 0.73).
- (E) Cumulative distribution showing the amplitudes from all recorded IPSCs. Inset, sampling distributions of the mean amplitude obtained with the hierarchical bootstrap. The mean amplitudes (dark lines) were not significantly different (hierarchical bootstrap, p = 0.14).

n = 11 abGCs from 3 control mice and 13 abGCs from 3 PLX mice (subset of the cells from the main figure that met criteria stated in Methods)

ns, not significant

Figure 8

Figure 8: Summary of results and model for the role of microglia in abGC maturation
Young abGCs enter existing circuits and begin to make synapses, a process that involves the extension of filopodial spines that sample potential synaptic partners (Breton-Provencher et al., 2014). In control mice, abGCs undergo a coordinated process of synaptic formation and elimination (Mizrahi, 2007)(Sailor et al., 2016), which leads to an increasing number of mushroom spines with mature synapses and higher frequency and amplitude excitatory synaptic currents as they mature (Whitman and Greer, 2007)(Kelsch et al., 2008)(Breton-Provencher et al., 2014). Maturation likely also involves a decrease in dendritic excitability (Carleton et al., 2003)(Nissant et al., 2009)(Wallace et al., 2017), such that stronger, coordinated synaptic inputs with particular odor tuning are necessary to activate mature abGCs, leading to more selective odor responses. In mice treated with PLX5622 to ablate microglia throughout the

timecourse of abGC development, we hypothesize that abGCs undergo some parts of the maturation process, but not others. We show that aspects of excitability, such as input resistance, mature normally, and overall spine density increases to levels even above controls. However, synapses fail to mature, leading to overall smaller spines and lower amplitude sEPSCs. This may contribute to lower odor responsiveness in abGCs that have matured in the absence of microglia.

Video 1. Z stack showing microglia and abGC dendrites. Movie showing the entire field of view at the beginning of the imaging session for the example shown in Figure 1. The z stack volume spans 29 µm and was taken at 32 dpi. The analysis in Figure 1 was performed on individual planes from such z stacks.

Video 2. Time series showing interactions between microglia and dendritic spines of abGCs. Movie showing a single plane taken from the z stack in Video 1 across 48 minutes of imaging (images taken 3 minutes apart). The time course of the interaction between a microglial process and the mushroom spine shown in the example in Figure 1 can be observed. The analysis in Figure 1 was performed on individual planes from such time series.

Figure 1-source data 1

This spreadsheet contains the values used to create the histograms in Figure 1D-G and I-L.

Figure 2-source data 1

This spreadsheet contains values from each dendrite from each mouse for Figure 2F, G, and H.

Figure 3-source data 1

This spreadsheet contains values from each dendrite for Figure 3B, C, and D.

Figure 4-source data 1

This spreadsheet contains values from each dendrite for Figure 3C, D, and E.

Figure 5-source data 1

This spreadsheet contains spine density values for each dendrite for Figure 5C and values for the head volume of each spine from each dendrite for Figure 5D.

Figure 6-source data 1

This spreadsheet contains the frequency and amplitude values for all detected events from each dendrite for Figure 6C, D, F, and G.

Figure 7-source data 1

This spreadsheet contains the frequency and amplitude values for all sEPSCs from each dendrite for Figure 7C and D.