Dietary vs non-dietary fatty acid profiles of lake trout morphs from Lake Superior and Great Bear Lake: Are fish really what they eat?

Chavarie, L[1] *, J. Hoffmann[2], A.M. Muir[3], C.C. Krueger[4], C.R. Bronte[5], K.L. Howland[6,7], C. Gallagher[6], S.P. Sitar[8], M.J. Hansen[9], M.R. Vinson[9], L.L. Loseto[6], W. Tonn[7], and H. Swanson[2]

[1]University of British Columbia, Biodiversity Center, 2212 Main Mall, Vancouver, BC, Canada, V6T 1Z4

[2] University of Waterloo, Department of Biology, 200 University Avenue West, Waterloo, Ontario, Canada, N2L 3G

[3]Great Lakes Fishery Commission, 2100 Commonwealth Blvd., Suite 100, Ann Arbor, Michigan, USA, 48105

[4]Michigan State University, Center for Systems Integration and Sustainability, 1405 South Harrison Road, 115 Manly Miles Building, East Lansing, Michigan USA 48823

[5]U.S. Fish and Wildlife Service, Green Bay Fish and Wildlife Conservation Office, New Franken, WI, USA

[6]Fisheries and Oceans Canada, 501 University Crescent, Winnipeg, Canada

[7]University of Alberta, Department of Biological Sciences, CW 405 BioSciences Building, Edmonton, Canada

[8]Michigan Department of Natural Resources, Marquette, MI, USA

[9]U.S. Geological Survey Great Lakes Science Center, Ashland, WI, USA

*Corresponding author: chavarie@zoology.ubc.ca

Key words: Intraspecific diversity, salmonid, aquatic ecology, trophic ecology, Great Lakes, *Salvelinus namaycush*, fatty acids

Abstract:

Fatty acids are well-established biomarkers used to characterize trophic ecology, food-web linkages, and organismal niche for a diversity of taxa. Most often, fatty acids examined include only those previously identified as "dietary" or "extended dietary". Consequently, non-dietary fatty acids are commonly discarded, representing approximately half of the fatty acids that can be extracted from fish muscle. In this study, we explored whether non-dietary fatty acids can provide worthwhile information by assessing their ability to discriminate intraspecific diversity within and between lakes. Non-dietary fatty acids delineated variation geographically (between lakes), between locations within a lake, and among morphs within a species. Non-dietary fatty acids were useful in delineating morphs of lake trout in both lakes and, for Lake Superior, performed better than dietary fatty acids. Physiological differences that arise from dynamic differences in energy processing can be adaptive and linked to habitat use (i.e., shallow vs. deep depths) by species morphs, and likely explains why non-dietary fatty acids can be a relevant tool to delineate intraspecific diversity in lake trout, which are known to inhabit a large range of habitats. The level of differences in non-dietary fatty acids were as pronounced among morphs within Great Bear Lake as observed in Lake Superior. Little is known what causes non-dietary-mediated differences in fatty acid composition, but our results showed that assessing non-dietary fatty acids can be useful in delineating intraspecific diversity and spatial variation.

Introduction:

Introduction:

Constraints on traditional methods to investigate diets of organisms within aquatic systems

have led to the use of biochemical tracers in contemporary investigations (Vinson & Budy, 2010).

Among these, fatty acids have gained popularity as both qualitative and quantitative trophic

markers that reflect foraging patterns and food-web dynamics (Iverson, 2009; Galloway *et al.*,

2014). As such, fatty acids have been used to characterize variation within populations and species

(e.g., evolutionary units linked to trophic polymorphism; Logan *et al.*, 2008; Scharnweber *et al.*,

2016), and to explore geographical (Pomerleau *et al.*, 2014; Hiltunen *et al.*, 2016; Quérouil *et al.*,

2013) and temporal variation in trophic ecology (e.g., seasonal and annual variations; Eloranta *et*

al., 2013; Hartwich, Martin-Creuzburg & Wacker, 2012).

Fatty acids are the "building blocks" of lipids and represent the largest constituent of

neutral lipids (e.g., triacylglycerols and wax esters) and polar phospholipids (Iverson, 2009). The

array of fatty acids present in nature is exceptionally complex, routinely ~70 fatty acids can be

identified within an organism (Iverson, 2009; Budge, Iverson & Koopman, 2006). The utility of

fatty acid analyses to reflect foraging patterns and food web dynamics relies on the assumption

that lipids are broken down into their constituent fatty acids and incorporated relatively unchanged

into consumer tissues (Howell *et al.*, 2003; Iverson *et al.*, 2004; Iverson, 2009). The storage

patterns of fatty acids depend on the biochemical limitations of organisms to biosynthesize, modify

chain-length, and introduce double bonds in fatty acids (e.g., with complexity culminating in

vertebrates) (Iverson 2009). A critical analytical assumption is that storage characteristics of fatty

acids within lipids are not being degraded during digestion, i.e., that they are stored in tissues in

their basic form (Iverson, 2009). Fatty acids labelled as "dietary" or "extended dietary" tracers

(Iverson, Frost & Lowry, 1997, 2004; Budge, Iverson & Koopman, 2006) have been the targets

for analyses in ecological studies. Consequently, approximately half of the fatty acids extracted

from tissue samples are commonly omitted, with only those recognized to be typical markers of

diet being analyzed.

The exclusion of these "non-dietary" fatty acids from analyses is not misguided, as the

purpose of most studies is to accurately describe diet patterns, and this practice has resulted in

reliable information being produced across taxa (Iverson *et al.*, 2004; Galloway *et al.*, 2014;

Iverson, 2009). However, it is unknown whether valid information for other research questions

may be lost when investigators discard a portion of the fatty acids in their samples and the potential

discriminatory data that could have resulted if analysed. Biological (e.g., phenotypic and genetic)

and environmental variables have been shown to affect lipid and fatty acid composition in fishes

(Olsen & Skjervold, 1995). For example, temperature (Olsen, 1999; Farkas *et al.*, 1980), salinity

(Borlongan & Benitez, 1992), and light (Ota & Yamada, 1971) have induced variation in lipid

composition of fish tissue. Thus, fatty acids not labelled as dietary markers could be useful when

the aim of a study is to delineate or better understand intraspecific diversity. To investigate these

questions, we compared "non-dietary" fatty acids among morphs of lake trout (*Salvelinus*

namaycush) in Lake Superior to those among lake trout morphs in Great Bear Lake, as these

morphs represent important intraspecific diversity in these lakes.

Salmonids, such as lake trout, inhabit evolutionarily young ecosystems believed to be

10,000 to 15,000 years old, e.g., post-glacial lakes colonized from non-glaciated refugia. The

depauperate communities of post-glacial lakes are commonly characterized by reduced

competition and predation, which allows colonizers access to a relatively wide array of resources,

conditions that favour development of intraspecific diversity (Smith & Skulason, 1996; McPhail,

1993; Robinson & Wilson, 1994). This high level of ecological opportunity, together with an

increase in intraspecific competition after colonization, can promote individual specialization

within a population, and the development of groups of individuals with similar patterns of resource

use (Svanbäck *et al.*, 2007). Observable characteristics of individuals in such groups can evolve

as niches expand to incorporate novel foraging resources, referred to as resource polymorphism

(Skulason & Smith, 1995; Robinson & Parsons, 2002). Morphs (or ecotypes) can differ from each

other in morphology, genetics, physiology, life-history, and/or behaviour, and these differences

are typically associated with niche use (Schluter & McPhail, 1992; Smith & Skulason, 1996;

Bolnick *et al.*, 2007).

Intraspecific diversity in lake trout has been mostly linked to differences in depth

distribution and, not surprisingly, is best known from large (> 500km^2), deep lakes, such as Lake

Superior (Muir *et al.*, 2014; Moore & Bronte, 2001), Lake Mistassini (Zimmerman, Krueger &

Eshenroder, 2007), Great Slave Lake (Zimmerman, Krueger & Eshenroder, 2006), and Great Bear

Lake (Chavarie, Howland & Tonn, 2013). Although lake trout diversification has often focused on

isolation-by-depth (without excluding isolation-by-adaptation) in large, deep lakes (e.g., Lake

Superior), diversification also occurs in small lakes or within shallow-water habitats (e.g., Great

Bear Lake; Chavarie, Howland & Tonn, 2013; Bernatchez *et al.*, 2016; Chavarie *et al.*, 2016c;

Morissette *et al.*, 2018).

A number of studies have assessed lake trout diets with fatty acids, either qualitatively

(Chavarie *et al.*, 2016b; Hoffmann, 2017; Happel *et al.*, 2017b; Happel *et al.*, 2017a) or

quantitatively (Happel *et al.*, 2016b; Happel *et al.*, 2016a), generating reliable information about

dietary patterns. However, pronounced and systematic non-dietary-mediated differences in fatty

acids composition have been reported in lake trout, supporting the rationale that non-dietary fatty

acids could be important in delineating intraspecific diversity within this species. Goetz *et al.*

(2013) found physiological differences between lean and siscowet morphs of lake trout that

reflected genetically based differences in lipid synthesis, metabolism, and transport (Goetz *et al.*,

2010), which suggests information from non-dietary fatty acids might be important, especially if

they have genetic-based mechanisms. Consequently, where intraspecific diversity is manifested as

physiological differences among morphs, non-dietary fatty acids could assist in identifying

variation. Yet, little is known about the non-dietary-mediated differences in fatty acid composition

of fish (or other taxa) in an ecological context.

To investigate the use of non-dietary fatty acids as an ecological tool, we analyzed a

comprehensive set of non-dietary and dietary fatty acids from lake trout morphs collected from

two lakes that sustain intraspecific diversity along different axes of intraspecific diversification

(e.g., depth-dependent and depth-independent). Specifically, we compared: 1) non-dietary and

dietary fatty acids among four morphs of lake trout from each of Great Bear Lake Lake Superior,

and 2) how well non-dietary and dietary delineated intraspecific diversity within and between

lakes. Answers to these questions will help determine if non-dietary fatty acids can discriminate

fish geographically and among groups within a species, and if they offer different perspectives

than dietary fatty acids.

Material and methods

Study systems and intraspecific diversity of lake trout

Located in the northeast corner of Northwest Territories (Canada), 65°92′ N, 120°82′ W,

Great Bear Lake is the most northerly lake of its size (~31 000 km^2), has a maximum depth of 446

m, and is the fifteenth deepest freshwater lake in the world (Fig. 1; Johnson, 1975). A UNESCO

biosphere reserve, Great Bear Lake is 250 km south of the Arctic Ocean and has characteristics

typical of an Arctic lake. The lake is ultra-oligotrophic and, despite its size, has a simple food web,

supporting only 15 fish species (Johnson, 1975; MacDonald *et al.*, 2004). The lake and its biota

have remained relatively isolated and unexploited, and is one of the most pristine large lakes in

North America. Samples of lake trout were collected from multiple locations, Great Bear Lake has

five semi-isolated arms, but due to sample size, data were pooled across multiple sites (see

Chavarie *et al.*, 2016b for details).

Great Bear Lake sustains a noteworthy example of lake trout divergence (Fig. 2). With its

intraspecific diversity independent of depth-based segregation, the lake also presents an unusual

ecological framework for lake trout differentiation (Chavarie *et al.*, 2016a). Currently, four lake

trout morphs are described in Great Bear Lake; three are common (Morphs 1-3), and one is rare

(Morph 4). Morph 1 has the smallest head and jaws, intermediate fin lengths and body shape

intermediate between Morphs 2 and 3. Morph 2 has the largest head and jaws but the smallest fins

(lean-like), and a streamlined body shape. Morph 3 has the longest fins, a robust body shape, and

a sub-terminal mouth. Morph 4 has a thick and curved lower jaw, a streamlined body shape, and

the smallest caudal peduncle among the morphs. Although Morph 4 is a specialist in the pelagic

habitat, whereas Morphs 1-3 have more general feeding habits, with varying degrees of omnivory

along a weak benthic-pelagic gradient in shallow-water (Chavarie *et al.*, 2016a; Chavarie *et al.*,

2016b).

Lake Superior is a post-glacial, oligotrophic lake shared between Canada and the USA

(47°43′ N, 86°56′ W) (Fig. 1). Lake Superior is the largest lake in the world by surface area (82

100 km^2). Most of the waters of Lake Superior can be classified as offshore; 77% of the total area

is greater than 80 m deep (maximum depth = 406 m) (see Horns *et al.*, 2003; Gorman, Yule &

Stockwell, 2012; Hoffmann, 2017). Lake Superior supports 87 fish species, with lake trout as a

main target of the commercial fishery since the 1800s. Lake trout were sampled from two sites,

Superior Shoal (48° 3'43.54" N, 87° 8'52.57" W) and Stannard Rock (47°12'26.26" N, 87°12'3.82"

W), in Lake Superior (Fig. 1). Superior Shoal and Stannard Rock were known to support humper,

lean, and siscowet Lake Trout morphs, whereas redfins, described by Muir *et al.* (2014), were not

yet known to be present at these two sites.

Lake Superior supports one of the highest levels of sympatric diversity expressed within

lake trout (Fig. 2). Four morphs are currently recognized; the siscowet, humper, and redfin morphs

inhabit deep-water (> 70 m), whereas the lean morph occupies shallow-water habitats (< 70 m)

(Moore & Bronte, 2001; Bronte & Moore, 2007; Muir *et al.*, 2014). The siscowet is characterized

by a large head, short snout, long maxilla, large eye, short and deep caudle peduncle, and

moderately long paired fins (Muir *et al.*, 2015; Bronte & Moore, 2007; Moore & Bronte, 2001).

The humper lake trout has a small head, short snout, short maxillae, large eyes, and short and

narrow caudal peduncle. (Muir *et al.*, 2015; Moore & Bronte, 2001). The redfin has the largest

head, snout, and eyes, the longest and deepest caudal peduncle, and much longer pelvic and

pectoral fins than the other morphs (Muir *et al.*, 2014). Finally, the lean has a large, narrow, and

pointed head, long snout, small eyes, long and narrow caudal peduncle, short paired fins, and low

body lipid content. (Endler, 1978; Muir *et al.*, 2015; Khan & Qadri, 1970).

Fatty acids

Dorsal muscle samples (Budge, Penney & Lall, 2011) from lake trout in both lakes were

stored at -20 °C (Budge, Iverson & Koopman, 2006; Loseto *et al.*, 2009; Kavanagh *et al.*, 2010;

Chavarie *et al.*, 2016b). Lipids were extracted from 1 g of the homogenate material; after passive

overnight extraction (at -20) in 2:1 chloroform:methanol containing 0.01% BHT (v/v/w) (Folch,

Lees & Sloane-Stanley, 1957), samples were filtered through Whatman Grade 1 Qualitative filter

paper and the filter paper sample was rinsed twice with 2 mL of 2:1 chloroform:methanol. Sample

extract was collected in a test tube and 7 mL 0.88 NaCl solution were added because NaCl

encourages fatty acids to move into the organic (chloroform) layer. The aqueous layer was

discarded after which the chloroform was dried with sodium sulfate prior to total lipid

determination. The extracted lipid was used to prepare fatty acid methyl esters (FAME) by

transesterification with Hilditch reagent (0.5 N H_2SO_4 in methanol) (Morrison & Smith, 1964).

Samples were heated for 1 h at 100 °C. Gas chromatographic (GC) analysis was performed on an

Agilent Technologies 7890N GC equipped with a 30 m J&W DB-23 column (0.25 mm I.D; 0.15

μm film thickness). The GC was coupled to a Flame Ionization Detector operating at 350 °C.

Hydrogen was used as carrier gas flowing at 1.25 mL/min for 14 minutes, increasing to 2.5 mL/min

for 5 minutes. The split/splitless injector was heated to 260 °C and run in splitless mode. The oven

program was as follows: 60 °C for 0.66 min, increasing by 22.82 °C/min to 165 °C with a 1.97 min

hold; increasing by 4.56 °C/min to 174 °C and by 7.61 °C/min to 200 °C with a 6 min hold. Peaks

were quantified using Agilent Technologies ChemStation software. FA standards were obtained

from Supelco (37 component FAME mix) and Nuchek (54 component mix GLC-463). FAMEs

were identified via retention time and known standard mixtures and are reported as percentages of

total fatty acids. All FA values were converted to a mass percentage of the total array, and were

named according the IUPAC nomenclature as X:Y n-z, where X is the number of carbons in the

FA, Y is the number of methylene-interrupted double bonds present in the chain, and n-z denotes

the position of the last double bond relative to the methyl terminus (Ronconi *et al.*, 2010). All

laboratory analyses were conducted at the Freshwater Institute, Fisheries and Oceans Canada,

Winnipeg MB.

For both lakes, tissue samples were taken from indivual lake trout previously identified to

corresponding morphotypes: 126 samples were collected from Great Bear Lake (Morph 1 = 32,

Morph 2 = 35, Morph 3 = 38, and Morph 4 = 21, see Chavarie *et al.*, 2016b for more details) and

210 samples were collected from Lake Superior (60 siscowet, 60 humper, 30 redfin, and 60 lean,

see Hoffmann, 2017 for more details). Overall, FA analysis procedures were divided in two steps,

using non-dietary and dietary fatty acids (see Appendix 1), following the methods of Iverson, Frost

and Lowry (1997, 2004) and Budge, Iverson and Koopman (2006) to identify dietary fatty acids.

Thirty-eight dietary and 24 non-dietary fatty acids were found to be shared between the two lakes,

and these were selected for further analyses (Table A1 and A2).

Statistical Analyses

Unless noted otherwise, statistical analyses were conducted using R software version 3.5.3

(R Core Team, 2017). Prior to analysis, fatty acid concentrations were logit transformed (log(p/1-

p) to normalize the data, and then scaled and centered using a z-score transformation ($z=x\mu/\sigma$).

(Witten & Tibshirani, 2011; Clemmensen *et al.*, 2011). Principal Canonical Analysis (PCA) was

performed on all dietary and non-dietary fatty acids to provide inference about patterns of variation

among locations and morphs (Chavarie *et al.*, 2016b). For PCAs, 12 outlier individuals from

Superior Shoal were excluded because PCAs are sensitive to outlier variation (but see Fig. A1 for

PCAs with their inclusion) (Filzmoser, Hron & Reimann, 2009; Kriegel *et al.*, 2008).

To test for differences in fatty acid composition among morphs within each lake (dietary,

non-dietary, and overall), Permutational Multivariate Analysis of Variance (PERMANOVA) was

used (a non-parametric analog of Multivariate Analysis of Variance), followed by post-hoc

comparisons with Bonferroni corrections. PERMANOVAs were performed in PAST 3 (Hammer,

Harper & Ryan, 2001) using 9999 permutations. A similarity percentage routine (SIMPER) using

Bray-Curtis was used to determine which fatty acids (dietary, non-dietary) were primarily

responsible for observed differences among morphs for each lake (King & Jackson, 1999). We

also performed linear discriminant analysis with a jacknife validation procedure using 20% of our

data as unknown on fatty acids (dietary, and non-dietary) from each location to evaluate their

contribution to delineating intraspecific diversity in lake trout. Finally, PERMANOVA (using

9999 permutations), and linear discriminant analyses and jacknife validation procedures were

performed on the ten most discriminating dietary and non-dietary fatty acids (20 fatty acids in total

to reduce the number of vectors versus number of individuals tested for linear discriminant

analysis); these fatty acids were identified from previous SIMPER analyses for each location.

Results

Combined lakes analyses

The first two axes of non-dietary fatty acids PCA explained 42.4 % of the variation among

lake trout from Great Bear Lake and Lake Superior (Fig. 3). Lake trout from Great Bear Lake

clustered together with 37 lake trout overlapped with those from Lake Superior, whereas those

from Stannard Rock and Superior shoal overlapped. The separation between the two lakes

appeared to be driven by the following fatty acids: 13:1, 14:1n-7, 15:0, 15:1n8, 15:1n6, 15:0 iso,

16:1n11, 17:0 iso, 20:0, 22:0, 20:2n9, and 24:1n9. Similar to the non-dietary fatty acids, the first

two axes of the dietary fatty acids PCA explained 41.5% of the variation among lake trout from

Great Bear Lake and Lake Superior. Lake trout from Great Bear Lake grouped together, with 32

individuals overlapped with Lake Superior lake trout, whereas lake trout from Stannard Rock

totally overlapped with Superior Shoal individuals (Fig. 3). Separation between the two lakes

appeared to be driven by fatty acids associated with pelagic habitat (14:0, 20:1n-9, 20:1n-7, 20:1n-

11 and 22:1n-9; toward Lake Superior) versus one fatty acid associated with cannibalism or/and

carnivory (20:5n3; toward Great Bear Lake) (Appendix, Table 1). Finally, the first two axes of the

PCA based on all fatty acids combined explained 39.0% of the variation among lake trout morphs

from Great Bear Lake and Lake Superior. Lake trout from Great Bear Lake clustered together but

with 15 lake trout overlapped with those from Lake Superior. Lake trout from Stannard Rock and

Superior Shoal in Lake Superior overlapped completely.

Lake Trout intraspecific diversity of Great Bear Lake

Composition of both non-dietary and dietary fatty acids were able to discriminate among

the four lake trout morphs from Bear Lake. Non-dietary fatty acids differed among the four Great

Bear Lake morphs (one-way PERMANOVA, $F_{3,122} = 3.6$, $P < 0.01$). Similarly, comparison of non-

dietary fatty acids showed that all pairs of morphs differed from one another ($P < 0.01$) except for

morphs 1 and 3. The ten most discriminating non-dietary fatty acids from SIMPER explained

~63.8% of the dissimilarity among groups (Table 1). The first two axes of the linear discriminant

analysis explained 55.7 % and 27.8% of the variation, and 57.9% of all individuals were correctly

classified to morph based on non-dietary fatty acids (Fig. 4). Dietary fatty acids also differed

among the four lake trout morphs from Great Bear Lake (one-way PERMANOVA, $F_{3,122} = 2.95$,

$P < 0.01$), and most pairs of morphs differed significantly from one another (all $P < 0.01$ except

for morphs 1 vs. 3). The ten most discriminating dietary fatty acids from SIMPER explained

~48.8% of the dissimilarity among groups (Table 2). The first two axes of the linear discriminant

analysis explained 46.7% and 39.0% of the variation, and 68.3% of all individuals were correctly

classified to morph based on dietary fatty acids (Fig. 4). Finally, when the 10 dietary and 10 non-

dietary most discriminating fatty acids were considered together, morphs differed from one another

(PERMANOVA, $F_{3,122} = 2.8$, $P < 0.01$). All pairs of morphs differed ($P < 0.01$) except for morphs

1 and 3. The first two axes of the linear discriminant analysis explained 63.6% and 24.9% of the

variation, and 54.8% of all individuals were correctly classified (Fig. 4).

Lake Trout intraspecific diversity of Lake Superior, Stannard Rock

Non-dietary fatty acids differed among the three morphs of lake trout from Stannard Rock

(one-way PERMANOVA, $F_{2,87} = 3.9$, $P < 0.01$), and leans differed significantly from both

siscowets and humpers ($P < 0.01$). The ten most discriminating non-dietary fatty acids from

SIMPER explained 70.9% of the dissimilarity among morphs (Table 1). The first two axes of the

linear discriminant analysis explained 60.2% and 39.8% of the variation, respectively, and 52.2%

of all individuals were correctly classified to morph based on non-dietary fatty acids (Fig. 4). In

contrast to results from non-dietary fatty acid, no differences in dietary fatty acid composition

occurred among the three lake trout morphs from Stannard Rock, Lake Superior (one-way

PERMANOVA, $F_{2,87} = 1.12$, $P = 0.3$). The ten most discriminating fatty acids from SIMPER

explained ~55.2 % of the dissimilarity among groups (Table 2), and the first two axes of the linear

discriminant analysis explained 60.9% and 39.1% of the variation. Fifty percent of individuals

were correctly classified to morph based on dietary fatty acids (Fig. 4). Finally, the three morphs

also differed when the 10 most discriminating non-dietary and dietary fatty acids were considered

together (one-way PERMANOVA, $F_{2,87} = 2.1$, $P < 0.02$), and leans differed significantly from

siscowets ($P < 0.01$). The first two axes of the linear discriminant analysis explained 73.3% and

26.8% of the variation, respectively, and 48.9% of all individuals were correctly classified to

morph based on their dietary fatty acid profile (Fig. 4).

Lake Trout intraspecific diversity of Lake Superior, Superior Shoal

Like Stannard Rock, composition of non-dietary fatty acid differed significantly among the

four morphs (one-way PERMANOVA, $F_{3,116}$ = 2.5, P < 0.01), and leans differed significantly

from redfins (P < 0.01) and were marginally significantly different between redfin and siscowet

and between redfin and humper (P < 0.06). The ten most discriminating non-dietary fatty acids

from SIMPER explained ~69.9% of the dissimilarity among groups (Table 1). The first two axes

of the linear discriminant analysis explained 64.3% and 25.6% of the variation, and 45.0%

individuals were correctly classified to morph using non-dietary fatty acids (Fig. 4). Based on

dietary fatty acid composition, we found no differences among the four lake trout morphs from

Superior Shoal (one-way PERMANOVA, $F_{3,116}$ = 0.8, P = 0.3), similarly to what was found with

fish from Stannard Rock. The ten most discriminating fatty acids from SIMPER explained ~55.8

% of the dissimilarity among groups (Table 2). The first two axes of the linear discriminant

analysis explained 38.6% and 35.6% of the variation, respectively, and 31.7% individuals were

correctly classified to morph based on their dietary fatty acid profile (Fig. 4). In contrast to the

results for Great Bear Lake and Stannard Rock, when the 10 non-dietary and 10 dietary most

discriminating fatty acids were combined for lake trout from Superior Shoal, morphs did not differ

significantly from one another (one-way PERMANOVA, $F_{3,116}$ = 1.3, P = 0.16). The first two

axes of the linear discriminant analysis explained 72.9% and 19.7% of the variation, respectively,

and 42.5% of all individuals were correctly classified (Fig. 4).

Discussion

While characterizing trophic divergence, food-web linkages, and time-integrated niche use

across a large array of taxa, most investigators use a common set of fatty acids that are selected

from lists that classify fatty acids as "dietary" or "extended dietary" (Budge, Iverson & Koopman,

2006; Iverson *et al.*, 2004). However, including fatty acids associated with supplementary (non-

dietary) information might be worth considering for some studies. In this study, non-dietary fatty

acids showed variation geographically (between lakes), between locations within a lake (i.e.,

Stannard Rock versus Superior Shoal), and among morphs within a species. Although some

overlap was found, our results showed that when investigating intraspecific diversity, non-dietary

fatty acids can be a useful tool to delineate groups, and that sometimes, such as at sites in Lake

Superior, were more discriminatory than dietary fatty acids. Accordingly, our results suggested

that discarding non-dietary fatty acid data as a matter of course may result in inadvertent loss of

information.

Not all fatty acids provide equivalent information about diet due to metabolism and *de novo*

synthesis (Iverson *et al.*, 2004; Iverson *et al.*, 1993). Metabolism can differ within species, as

physiological differences often exists among sets of individuals, i.e., "morphs" (Pryke *et al.*, 2007;

Miles *et al.*, 2007). Such physiological differences within a species may result in non-dietary fatty

acids being a relevant tool to delineate intraspecific diversity, as we show here for lake trout. In

Lake Superior, fat content has long been recognized as an important characteristic that

distinguishes lake trout morphs along a depth axis (Eschmeyer & Phillips, 1965; Eshenroder,

2008), and these lipid differences have been linked to genetic differences among morphs

(Eschmeyer & Phillips, 1965; Goetz *et al.*, 2013; Goetz *et al.*, 2010). Thus, fatty acid deposition

and metabolism appears to have undergone selection along a depth gradient likely in part to assist

with buoyancy compensation, with differences reported between siscowet (deep-water morph) and

lean (shallow-water morph) lake trout (Goetz *et al.*, 2010; Eschmeyer & Phillips, 1965). Until

now, no information has been available for redfin and humper morphs.

In addition to the variation in lipid accumulation, Goetz *et al.* (2013) also found differences

in lipid composition between siscowet and lean morphs. Because a common garden design was

used in the Goetz *et al.* (2013) experiment, the higher proportion of polyunsaturated fatty acids

(PUFAs) found in the muscle lipid profile of siscowet than in lean lake trout could not be attributed

to differences in diet. The differences in non-dietary fatty acids we observed among the four

morphs of lake trout from Lake Superior was consistent with the concept of metabolotypes (Goetz

et al., 2013). Altogether, differences in energy processing and storage between lean and siscowet

lake trout in Lake Superior are adaptive to their respective habitats, deep- vs. shallow-water, and

to their life-histories (Goetz *et al.*, 2013). Lipid content, intertwined with buoyancy variations,

have been linked to differences in depth distributions and swimming tactics among lake trout

morphs (Zimmerman, Krueger & Eshenroder, 2006; Zimmerman, Krueger & Eshenroder, 2007;

Zimmerman *et al.*, 2009). While deep-water morphs use hydrostatic lift related to lipid content to

enhance vertical migration while foraging for *Mysis* and cisco (Henderson & Anderson, 2002), the

shallow-water lean morph likely relies more on hydrodynamic lift, linked to the cruising

movements of pelagic predators (Webb, 1984). Consistent with the findings of Goetz et al. (2013),

we found that non-dietary fatty acids differed between shallow- and deep-water morphs at

Stannard Rock. However, at Superior Shoal, non-dietary fatty acids differed only between the

shallow-water lean morph and one deep-water morph – the redfin. This observation, along with

some more subtle differences in non-dietary fatty acids among deep-water morphs, requires further

study. At both Stannard Rock and Superior Shoal, diet overlap was found among morphs, and

present and future diet overlap among morphs may be linked to ongoing changes in the prey base

(Bronte *et al.*, 2003; Gorman *et al.*, 2013). Changes in the prey base occurred in Lake Superior

prior to sampling, including loss of diversity of some coregonine forage fishes (Bronte *et al.*, 2003;

Gorman *et al.*, 2013). Because conserving diversity of lake trout morphs is a priority for managers

(Zimmerman & Krueger, 2009), a deeper understanding how diet overlap among morphs responds

to changes in the prey base also deserves more consideration.

The concept that dynamics of energy processing and storage are adaptive along a gradient

that is at least partially driven by depth does not apply to the intraspecific diversity of lake trout in

Great Bear Lake. This diversity is limited to shallow-water habitat, and appears to be independent

of major habitat and resource partitioning (Chavarie *et al.*, 2016a; Chavarie *et al.*, 2018). The

similarity of results between Great Bear Lake and Lake Superior is thus perplexing, as we were

expected greater differences in non-dietary fatty acids among morphs in Lake Superior than in

Great Bear Lake, due to known buoyancy variation associated with a depth gradient in Lake

Superior. If lake trout morphs from Great Bear Lake are also under selection (Harris *et al.*, 2014),

differences in energy processing and storage may be as pronounced as those that have been

observed in Lake Superior. Another question raised by our results is the extent to which morphs

are independent of major habitat or resource partitioning (e.g., same question is pertinent for Lake

Superior), especially because dietary fatty acids in Great Bear Lake were slightly better at

delineating intraspecific groups than non-dietary fatty acids.

Ecological differences in allopatry are often found to be more pronounced than those in

sympatry, due to disparate environments and isolation (Rundle & Nosil, 2005; Yoder *et al.*, 2010;

Heggenes, 2002; Fraser *et al.*, 2011). In this study, differences between lakes in both dietary and

non-dietary fatty acids appear to be greater than variation among morphs within a lake. For dietary

fatty acids, differences between the two lakes appeared to be due to fatty acids associated with a

pelagic environment, such as C20 and C22 monounsaturates, that can be used as biomarkers of

food webs based on pelagic copepods (Ahlgren, Vrede & Goedkoop, 2009; Budge, Iverson &

Koopman, 2006; Dalsgaard *et al.*, 2003). Although benthic productivity generally dominates in

Arctic lakes (Chavarie *et al.*, 2018; Johnson, 1975), some lake trout from Great Bear Lake

specialize on pelagic resources, including some individuals of Morph 2 and Morph 4, which have

fatty acids signatures that overlap with lake trout from Lake Superior (Chavarie et al., submitted;

Chavarie *et al.*, 2016a; Chavarie *et al.*, 2016b). With lake productivity in Arctic regions known to

be benthic-oriented, combined with the known depth distributions in the deep-water habitat of lake

trout morphs from Lake Superior (50-150 m; Muir *et al.*, 2014), our results reflected the expected

ecology of this species from these two systems. Despite differences between lake trout inhabiting

Great Bear Lake and Lake Superior, considerable overlap occurred among dietary (e.g., five out

of 10) and non-dietary (e.g., 7 out of 10) fatty acids that were important in identifying intraspecific

diversity in both lakes. The greater number of shared non-dietary fatty acids discriminating lake

trout intraspecific diversity in the two lakes supports the idea of similar physiological differences

(e.g., energy processing and storage) among morphs from both lakes.

A few caveats should be noted that could alter our interpretations. First, no spatial

component was defined for the Great Bear Lake dataset, due to small sample sizes from each

location, which probably introduced variation into the results. Spatial variation can become

important when a large, complex system is studied (Chavarie *et al.*, 2015; Hoffmann, 2017) and

environmental variables (e.g., temperature, light) are known to alter lipid composition in fish tissue

(Olsen, 1999). Second, in Lake Superior, multiple life-stages, e.g., juvenile, mature, and resting

individuals were included together in the analysis, and different life-stages can vary in lipid

metabolism (Sheridan, 1989). In addition, lake trout length can also influence fatty acid

composition. For example, in lakes Huron and Michigan, large lake trout relied more on nearshore-

benthic food web resources than small lake trout (Happel *et al.*, 2017a). Finally, some fatty acids

exist at very low amounts ($\leq 2\%$), which can introduce error in interpreting differences among

fatty acids that are found in only trace amounts (e.g., peak shouldering). Despite these limitations,

we found some consistent patterns with regards to intraspecific diversity between lakes and among

morphs within lakes.

Conclusion

This study demonstrated the benefits of using both dietary and non-dietary fatty acids for

delineating variation and diversity within a species. In some instances, non-dietary fatty acids were

better for discriminating morphs than dietary fatty acids, in contrast to the popular maxim

associated with trophic markers, "you are what you eat". Dietary fatty acids can determine the

occurrence (or absence) of discrete niche use among sets of individuals (i.e., morphs) within a

species (Chavarie *et al.*, 2016b). The fatty-acid composition of individuals reflecting their diet has

been validated for lake trout (Happel *et al.*, 2016a; Happel *et al.*, 2016b). However, physiological

differences in the dynamics of energy processing (e.g., metabolism) and storage can be adaptive

among morphs (Goetz *et al.*, 2013; Goetz *et al.*, 2010; Eschmeyer & Phillips, 1965), and thus

non-dietary fatty acids may be useful molecular biomarkers. Ultimately, the relative importance

of dietary vs. non-dietary fatty acids depends on the question being asked. In our study, non-dietary

fatty acids were valuable in delineating intraspecific variation within a lake, but also in examining

differences between lakes. Non-dietary fatty acids can provide reliable information; therefore, one

should carefully consider if such information is superfluous or not before data from these fatty

acids are discarded.

Acknowledgement

We thank Déline Renewable Resources Council, Déline Lands and Finance Corporation,

the community of Déline, DFO in Hay River, and the Department of Environment and Natural

Data Availability Statement:

All data presented are available by request via e-mail to the first author.

Conflict Of Interest

The authors declare that they have no conflict of interest.

Literature:
Ahlgren G., Vrede T. & Goedkoop W. (2009) Fatty Acid Ratios in Freshwater Fish, Zooplankton and
Zoobenthos – Are There Specific Optima? In: *Lipids in Aquatic Ecosystems.* (Eds M. Kainz & M.T.
Brett & M.T. Arts), pp. 147-178. Springer New York, New York, NY.
Bernatchez S., Laporte M., Perrier C., Sirois P. & Bernatchez L. (2016) Investigating genomic and
phenotypic parallelism between piscivorous and planktivorous lake trout (*Salvelinus namaycush*)
ecotypes by means of RADseq and morphometrics analyses. *Molecular Ecology,* **25,** 4773-4792.
Bolnick D.I., Svanbäck R., Araújo M.S. & Persson L. (2007) Comparative support for the niche variation
hypothesis that more generalized populations also are more heterogeneous. *Proceedings of the*
National Academy of Sciences, **104,** 10075-10079.
Borlongan I.G. & Benitez L.V. (1992) Lipid and fatty acid composition of milkfish (Chanos chanos Forsskal)
grown in freshwater and seawater. *Aquaculture,* **104,** 79-89.
Bronte C.R., Ebener M.P., Schreiner D.R., Devault D.S., Petzold M.M., Jensen D.A., Richards C. & Lozano
S.J. (2003) Fish community change in Lake Superior, 1970-2000. *Canadian Journal of Fisheries and*
Aquatic Sciences, **60,** 1552-1574.

Bronte C.R. & Moore S.A. (2007) Morphological variation of siscowet lake trout in Lake Superior.
Transactions of the American Fisheries Society, **136,** 509-517.

Budge S.M., Iverson S.J. & Koopman H.N. (2006) Studying trophic ecology in marine ecosystems using fatty
acids: a primer on analysis and interpretation. *Marine Mammal Science,* **22,** 759-801.

Budge S.M., Penney S.N. & Lall S.P. (2011) Response of tissue lipids to diet variation in Atlantic salmon
(Salmo salar): Implications for estimating diets with fatty acid analysis. *Journal of Experimental*
Marine Biology and Ecology, **409,** 267-274.

Chavarie L., Harford W.J., Howland K.L., Fitzsimons J., Muir A.M., Krueger C.C. & Tonn W.M. (2016a)
Multiple generalist morphs of Lake Trout: Avoiding constraints on the evolution of intraspecific
divergence? *Ecology and Evolution,* **6,** 7727–7741.

Chavarie L., Howland K., Gallagher C. & Tonn W. (2016b) Fatty acid signatures and stomach contents of
four sympatric Lake Trout: assessment of trophic patterns among morphotypes in Great Bear
Lake. *Ecology of Freshwater Fish,* **25,** 109-124.

Chavarie L., Howland K., Harris L. & Tonn W. (2015) Polymorphism in lake trout in Great Bear Lake: intra-
lake morphological diversification at two spatial scales. *Biological Journal of the Linnean Society,*
114, 109-125.

Chavarie L., Howland K.L., Harris L.N., Hansen M.J., Harford W.J., Gallagher C.P., Baillie S.M., Malley B.,
Tonn W.M., Muir A.M. & Krueger C.C. (2018) From top to bottom: Do Lake Trout diversify along a
depth gradient in Great Bear Lake, NT, Canada? *PloS one,* **13,** e0193925.

Chavarie L., Howland K.L. & Tonn W.M. (2013) Sympatric polymorphism in lake trout: the coexistence of
multiple shallow-water morphotypes in Great Bear Lake. *Transactions of the American Fisheries*
Society, **142,** 814-823.

Chavarie L., Muir A.M., Zimmerman M.S., Baillie S.M., Hansen M.J., Nate N.A., Yule D.L., Middel T., Bentzen
P. & Krueger C.C. (2016c) Challenge to the model of lake charr evolution: shallow- and deep-water
morphs exist within a small postglacial lake. *Biological Journal of the Linnean Society***,** n/a-n/a.

Clemmensen L., Witten D., Hastie T. & Ersbøll B. (2011) Sparse Discriminant Analysis. *Technometrics,* **53,**
406-413.

Dalsgaard J., St. John M., Kattner G., Müller-Navarra D. & Hagen W. (2003) Fatty acid trophic markers in
the pelagic marine environment. In: *Advances in Marine Biology* pp. 225-340. Academic Press.

Eloranta A.P., Mariash H.L., Rautio M. & Power M. (2013) Lipid-rich zooplankton subsidise the winter diet
of benthivorous Arctic charr (Salvelinus alpinus) in a subarctic lake. *Freshwater Biology,* **58,** 2541-
2554.

Endler J.A. (1978) A predator's view of animal color patterns. In: *Evolutionary Biology.* (Eds M.K. Hecht &
W.C. Steere & B. Wallace). Plenum Press, New York and London.

Eschmeyer P.H. & Phillips A.M., Jr. (1965) Fat content of the flesh of siscowets and lake trout from Lake
Superior. *Transactions of the American Fisheries Society,* **1965,** 62-74.

Eshenroder R.L. (2008) Differentiation of deep-water lake charr *Salvelinus namaycush* in North American
lakes. *Environmental Biology of Fishes,* **83,** 77-90.

Farkas T., Csengeri I., Majoros F. & Oláh J. (1980) Metabolism of fatty acids in fish: III. Combined effect of
environmental temperature and diet on formation and deposition of fatty acids in the carp,
Cyprinus carpio Linnaeus 1758. *Aquaculture,* **20,** 29-40.

Filzmoser P., Hron K. & Reimann C. (2009) Principal component analysis for compositional data with
outliers. *Environmetrics,* **20,** 621-632.

Folch J., Lees M. & Sloane-Stanley G. (1957) A simple method for the isolation and purification of total
lipids from animal tissues. *Journal of Biological Chemistry,* **226,** 497-509.

Fraser D.J., Weir L.K., Bernatchez L., Hansen M.M. & Taylor E.B. (2011) Extent and scale of local adaptation
in salmonid fishes: review and meta-analysis. *Heredity,* **106,** 404.

Galloway A.W.E., Eisenlord M.E., Dethier M.N., Holtgrieve G.W. & Brett M.T. (2014) Quantitative estimates of isopod resource utilization using a Bayesian fatty acid mixing model. *Marine Ecology Progress Series,* **507,** 219-232.

Goetz F., Jasonowicz A., Johnson R., Biga P., Fischer G. & Sitar S. (2013) Physiological differences between lean and siscowet lake trout morphotypes: Are these metabolotypes? *Canadian Journal of Fisheries and Aquatic Sciences,* **71,** 427-435.

Goetz F., Rosauer D., Sitar S.P., Goetz G., Simchick C., Roberts S., Johnson R., Murphy C., Bronte C.R. & Mackenzie S. (2010) A genetic basis for the phenotypic differentiation between siscowet and lean lake trout (*Salvelinus namaycush*). *Molecular Ecology,* **19 (Suppl. 1),** 176-196.

Gorman O.T., Evrard L.M., Cholwek G.A. & Vinson M.R. (2013) Status and trends in the fish community of LAke Superior, 2012. (Ed U.G. Survey), Washington, DC.

Gorman O.T., Yule D.L. & Stockwell J.D. (2012) Habitat use by fishes of Lake Superior. II. Consequences of diel habitat use for habitat linkages and habitat coupling in nearshore and offshore waters. *Aquatic Ecosystem Health & Management,* **15,** 355-368.

Hammer Ø., Harper D. & Ryan P. (2001) Past: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica,* **4,** 1-9.

Happel A., Jonas J.L., Mckenna P., Rinchard J., He J.X. & Czesny S.J. (2017a) Spatial Variability of Lake Trout Diets in Lakes Huron and Michigan Revealed by Stomach Content and Fatty Acid Profiles. *Canadian Journal of Fisheries and Aquatic Sciences*.

Happel A., Pattridge R., Walsh M. & Rinchard J. (2017b) Assessing diet compositions of Lake Ontario predators using fatty acid profiles of prey fishes. *Journal of Great Lakes Research*.

Happel A., Stratton L., Kolb C., Hays C., Rinchard J. & Czesny S. (2016a) Evaluating quantitative fatty acid signature analysis (QFASA) in fish using controlled feeding experiments. *Canadian Journal of Fisheries and Aquatic Sciences,* **73,** 1222-1229.

Happel A., Stratton L., Pattridge R., Rinchard J. & Czesny S. (2016b) Fatty-acid profiles of juvenile lake trout reflect experimental diets consisting of natural prey. *Freshwater Biology,* **61,** 1466-1476.

Harris L.N., Chavarie L., Bajno R., Howland K.L., Wiley S.H., Tonn W.M. & Taylor E.B. (2014) Evolution and origin of sympatric shallow-water morphotypes of Lake Trout, Salvelinus namaycush, in Canada's Great Bear Lake. *Heredity,* **114,** 94.

Hartwich M., Martin-Creuzburg D. & Wacker A. (2012) Seasonal changes in the accumulation of polyunsaturated fatty acids in zooplankton. *Journal of Plankton Research,* **35,** 121-134.

Heggenes J. (2002) Flexible Summer Habitat Selection by Wild, Allopatric Brown Trout in Lotic Environments. *Transactions of the American Fisheries Society,* **131,** 287-298.

Henderson B.A. & Anderson D.M. (2002) Phenotypic differences in buoyancy and energetics of lean and siscowet lake charr in Lake Superior. *Environmental Biology of Fishes,* **64,** 203-209.

Hiltunen M., Taipale S.J., Strandberg U., Kahilainen K.K. & Kankaala P. (2016) High intraspecific variation in fatty acids of Eudiaptomus in boreal and subarctic lakes. *Journal of Plankton Research,* **38,** 468-477.

Hoffmann J.M. (2017) *Investigating trophic ecology and dietary niche overlap among morphs of Lake Trout in Lake Superior.* MS thesis, University of Waterloo.

Horns W., Bronte C.R., Busiahn T.R., Ebener M.P., Eshenroder R.L., Gorenflo T., Kmieck N., Mattes W., Peck J.W., Petzold M. & Schreiner D.R. (2003) Fish-community objectives for Lake Superior. Great Lakes Fishery Commission Special Publication 03-01 Ann Arbor, Michigan.

Howell K.L., Pond D.W., Billett D.S. & Tyler P.A. (2003) Feeding ecology of deep-sea seastars (Echinodermata: Asteroidea): a fatty-acid biomarker approach. *Marine Ecology Progress Series,* **255,** 193-206.

Iverson S.J. (2009) Tracing aquatic food webs using fatty acids: from qualitative indicators to quantitative determination. In: *Lipids in Aquatic Ecosystems* pp. 281-308. Springer.

Iverson S.J., Bowen W.D., Boness D.J. & Oftedal O.T. (1993) The Effect of Maternal Size and Milk Energy
Output on Pup Growth in Grey Seals (Halichoerus grypus). *Physiological Zoology,* **66,** 61-88.
Iverson S.J., Field C., Bowen W.D. & Blanchard W. (2004) Quantitative fatty acid signature analysis: a new
method of estimating predator diets. *EcologicalMonographs,* **74,** 211–235.
Iverson S.J., Frost K.J. & Lowry L.F. (1997) Fatty acid signatures reveal fine scale structure of foraging
distribution of harbor seals and their prey in Prince William Sound, Alaska. *Marine Ecology
Progress Series,* **151,** 255-271.
Johnson L. (1975) Distribution of fish species in Great Bear Lake, Northwest Territories, with reference to
zooplankton, benthic invertebrates, and environmental conditions. *Journal of the Fisheries
Research Board of Canada,* **32,** 1989-2004.
Kavanagh A.S., Cronin M.A., Walton M. & Rogan E. (2010) Diet of the harbour seal (*Phoca vitulina vitulina*)
in the west and south-west of Ireland. *Journal of the Marine Biological Association of the United
Kingdom,* **90,** 1517-1527.
Khan N.Y. & Qadri S.U. (1970) Morphological differences in Lake Superior lake char. *Journal of the Fisheries
Research Board of Canada,* **27,** 161-167.
King J.R. & Jackson D.A. (1999) Variable selection in large environmental data sets using principal
components analysis. *Environmetrics,* **10,** 67-77.
Kriegel H.-P., Kröger P., Schubert E. & Zimek A. A General Framework for Increasing the Robustness of
PCA-Based Correlation Clustering Algorithms. In: *Scientific and Statistical Database Management.*
(Eds B. Ludäscher & N. Mamoulis), pp. 418-435. 2008// 2008. Springer Berlin Heidelberg.
Logan M.S., Walde S.J., Iverson S.J., Ruzzante D.E., Alonso M.F., Macchi P.J. & Cussac V.E. (2008) Long term
thet differences between morphs in trophically polymorphic Percichthys trucha (Pisces:
Percichthyidae) populations from the southern Andes. *Biological Journal of the Linnean Society,*
69, 599-616.
Loseto L., Stern G., Connelly T., Deibel D., Gemmill B., Prokopowicz A., Fortier L. & Ferguson S. (2009)
Summer diet of beluga whales inferred by fatty acid analysis of the eastern Beaufort Sea food
web. *Journal of Experimental Marine Biology and Ecology,* **374,** 12-18.
Macdonald D., Levy D., Czarnecki A., Low G. & Richea N. (2004) State of the aquatic knowledge of Great
Bear Lake watershed. Report to Indian and Northern Affairs Canada. *Water Resources Division,
MacDonald Environmental Sciences, Nanaimo, British Columbia*.
Mcphail J. (1993) Ecology and evolution of sympatric sticklebacks (Gasterosteus): origin of the species
pairs. *Canadian Journal of Zoology,* **71,** 515-523.
Miles D.B., Sinervo B., Hazard L.C., Svensson E.I. & Costa D. (2007) Relating endocrinology, physiology and
behaviour using species with alternative mating strategies. *Functional Ecology,* **21,** 653-665.
Moore S.A. & Bronte C.R. (2001) Delineation of sympatric morphotypes of lake trout in Lake Superior.
Transactions of the American Fisheries Society, **130,** 1233-1240.
Morissette O., Sirois P., Lester N.P., Wilson C.C. & Bernatchez L. (2018) Supplementation stocking of Lake
Trout (Salvelinus namaycush) in small boreal lakes: Ecotypes influence on growth and condition.
PloS one, **13,** e0200599.
Morrison W.R. & Smith L.M. (1964) Preparation of fatty acid methyl esters and dimethylacetals from lipids
with boron fluoride–methanol. *Journal of Lipid Research,* **5,** 600-608.
Muir A.M., Bronte C.R., Zimmerman M.S., Quinlan H.R., Glase J.D. & Krueger C.C. (2014) Ecomorphological
diversity of Lake Charr *Salvelinus namaycush* at Isle Royale, Lake Superior. *Transactions of the
American Fisheries Society,* **143,** 972-987.
Muir A.M., Hansen M.J., Bronte C.R. & Krueger C.C. (2015) If Arctic charr *Salvelinus alpinus* is 'the most
diverse vertebrate', what is the lake charr *Salvelinus namaycush*? *Fish and Fisheries***,**
DOI:10.1111/faf.12114.

Olsen Y. (1999) Lipids and essential fatty acids in aquatic food webs: what can freshwater ecologists learn
from mariculture? In: *Lipids in freshwater ecosystems.* (Eds M.T. Arts & B.C. Wainman), pp. 161-
202. Springer-Verlag, New York.

Olsen Y. & Skjervold H. (1995) Variation in content of Ω3 fatty acids in farmed Atlantic salmon, with special
emphasis on effects of non-dietary factors. *Aquaculture International,* **3,** 22-35.

Ota T. & Yamada M. (1971) Lipids of Masu salmon Oncorhynchus masou. I. Variations of the lipid content
and the fatty acid composition of juvenile Masu salmon during the period of smolt-
transformation, and on the influence of light upon those variations. *Hokkaido Univ Fac Fish Bull*.

Pomerleau C., Lesage V., Winkler G., Rosenberg B. & Ferguson S.H. (2014) Contemporary Diet of Bowhead
Whales (Balaena mysticetus) from the Eastern Canadian Arctic Inferred from Fatty Acid
Biomarkers. *Arctic,* **67,** 84-92.

Pryke S.R., Astheimer Lee B., Buttemer William A. & Griffith Simon C. (2007) Frequency-dependent
physiological trade-offs between competing colour morphs. *Biology Letters,* **3,** 494-497.

Quérouil S., Kiszka J., Cordeiro A.R., Cascão I., Freitas L., Dinis A., Alves F., Santos R.S. & Bandarra N.M.
(2013) Investigating stock structure and trophic relationships among island-associated dolphins
in the oceanic waters of the North Atlantic using fatty acid and stable isotope analyses. *Marine
Biology,* **160,** 1325-1337.

R Core Team. (2017) R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria.

Robinson B.W. & Parsons K.J. (2002) Changing times, spaces, and faces: tests and implications of adaptive
morphological plasticity in the fishes of northern postglacial lakes. *Canadian Journal of Fisheries
and Aquatic Sciences,* **59,** 1819-1833.

Robinson B.W. & Wilson D.S. (1994) Character release and displacement in fishes: a neglected literature.
American Naturalist, **144,** 596-627.

Ronconi R., Koopman H.N., Mckinstry C., Wong S. & Westgate A.J. (2010) Inter-annual variability in diet of
non-breeding pelagic seabirds *Puffinus* spp. at migratory staging areas: evidence from stable
isotopes and fatty acids. *Marine Ecology Progress Series,* **419,** 267-282.

Rundle H.D. & Nosil P. (2005) Ecological speciation. *Ecology Letters,* **8,** 336-352.

Scharnweber K., Strandberg U., Marklund M.H.K. & Eklöv P. (2016) Combining resource use assessment
techniques reveals trade-offs in trophic specialization of polymorphic perch. *Ecosphere,* **7,**
e01387.

Schluter D. & Mcphail J.D. (1992) Ecological character displacement and speciation in sticklebacks.
American Naturalist, **140,** 85-108.

Sheridan M.A. (1989) Alterations in lipid metabolism accompanying smoltification and seawater
adaptation of salmonid fish. *Aquaculture,* **82,** 191-203.

Skulason S. & Smith T.B. (1995) Resource polymorphisms in vertebrates. *Trends in Ecology & Evolution,*
10, 366-370.

Smith T.B. & Skulason S. (1996) Evoloutionary significance of resource polymorphims in fishes,
amphibians, and birds. *Annual Review of Ecology Evolution and Systematics,* **27,** 111-133.

Svanbäck R., Eklöv P., Fransson R. & Holmgren K. (2007) Intraspecific competition drives multiple species
resource polymorphism in fish communities. *Oikos,* **117,** 114-124.

Vinson M.R. & Budy P. (2010) Sources of variability and comparability between salmonid stomach
contents and isotopic analyses: study design lessons and recommendations. *Canadian Journal of
Fisheries and Aquatic Sciences,* **68,** 137-151.

Webb P.W. (1984) Body form, locomotion and foraging in aquatic vertebrates. *American Zoologist,* **24,**
107-120.

Witten D.M. & Tibshirani R. (2011) Penalized classification using Fisher's linear discriminant. *Journal of the
Royal Statistical Society. Series B, Statistical methodology,* **73,** 753-772.

Yoder J.B., Clancey E., Des Roches S., Eastman J.M., Gentry L., Godsoe W., Hagey T.J., Jochimsen D., Oswald
B.P., Robertson J., Sarver B.a.J., Schenk J.J., Spear S.F. & Harmon L.J. (2010) Ecological opportunity
and the origin of adaptive radiations. *Journal of Evolutionary Biology,* **23,** 1581-1596.
Zimmerman M.S. & Krueger C.C. (2009) An ecosystem perspective on re-establishing native deepwater
fishes in the Laurentian Great Lakes. *North American Jounal of Fisheries Management,* **29,** 1352-
1371.
Zimmerman M.S., Krueger C.C. & Eshenroder R.L. (2006) Phenotypic diversity of lake trout in Great Slave
Lake: differences in morphology, buoyancy, and habitat depth. *Transactions of the American
Fisheries Society,* **135,** 1056-1067.
Zimmerman M.S., Krueger C.C. & Eshenroder R.L. (2007) Morphological and ecological differences
between shallow- and deep-water lake trout in Lake Mistassini, Quebec. *Journal of Great Lakes
Research,* **33,** 156-169.
Zimmerman M.S., Schmidt S.N., Krueger C.C., Vander Zanden M.J. & Eshenroder R.L. (2009) Ontogenetic
niche shifts and resource partitioning of lake trout morphotypes. *Canadian Journal of Fisheries
and Aquatic Sciences,* **66,** 1007-1018.

Table 1. The ten most discriminating non-dietary fatty acids from SIMPER analyses. Results are presented for each region, including percentage contribution to overall fatty acid dissimilarity among lake trout morphs, and their order of importance (1-10, in parentheses). Fatty acids are listed in order of elution; those highlighted in grey are shared among the study three regions. The total percentage of the ten most discriminating fatty acids are given for each regions.

	Fatty acids	Great Bear Lake (63.8%)	Stannard Rock (70.9%)	Superior Shoal (69.9%)
Non-dietary Fatty Acids	12:0		3.6% (9)	6.1% (3)
	12:1	8.3% (3)	8.9% (2)	14.7% (2)
	13:1			
	14:1n-9	5.4% (6)	3.0% (10)	4.5% (6)
	14:1n-7	3.8% (9)		3.3% (10)
	14:1n-5			4.2% (8)
	14:0 iso		6.4% (4)	
	14:0 ante			
	15:0			
	15:1n-8	8.5 % (2)	25.9% (1)	18.8% (1)
	15:1n-6	8.2 % (4)	3.7% (7)	4.2% (7)
	15:0 iso	3.8% (10)	3.6% (8)	3.6% (9)
	16:1n-11			
	16:1n-9			
	16:1n-5			
	16:0 iso	7.1% (5)		
	7Methyl16:0	4.4% (7)		
	17:1			
	17:0 iso			
	18:1n-5			
	20:0		4.1% (6)	
	20:2n-9	10.2% (1)	4.5% (5)	5.2% (5)
	22:0	4.1% (8)	7.2% (3)	5.3% (4)
	24:1n-9			

698 Table 2. The ten most discriminating dietary fatty acids from SIMPER analyses. Results are
699 presented for each region, including percentage contribution to overall fatty acid
700 dissimilarity among lake trout morphs, and their order of importance (1-10, in parentheses). Fatty
701 acids are listed in order of elution; those highlighted in grey are shared among the study three
702 regions. The total percentage of the ten most discriminating fatty acids are given for each regions.

	Fatty acids	Great Bear Lake (48.8%)	Stannard Rock (55.2%)	Superior Shoal (55.8%)
Dietary and Extended-Dietary Fatty Acids	14:0			
	16:0			
	16:1n-7			
	16:2n-6			
	16:2n-4			
	17:0			
	16:3n-4	4.0% (6)	2.5 % (9)	2.9% (10)
	16:4n-3	3.0% (10)	2.6% (8)	
	16:4n-1		7.3% (3)	6.3% (3)
	18:0			
	18:1n-9		2.4% (10)	
	18:1n-7			
	18:2n-6			
	18:2n-4			
	18:3n-6			
	18:3n-4			
	18:3n-3			
	18:3n-1	3.3% (9)		3.5% (7)
	18:4n-3			
	18:4n-1	3.5% (8)	10.5% (2)	11.1% (2)
	20:1n-11	4.0% (5)		
	20:1n-9			
	20:1n-7	3.6% (7)		
	20:2n-6			
	20:3n-6			
	20:4n-6			
	20:3n-3			
	20:4n-3			
	20:5n-3			
	22:1n-11	8.6% (1)	11.9% (1)	12.1 % (1)
	22:1n-9			
	22:1n-7		2.7% (6)	4.6% (5)
	22:2n-6	6.3% (3)		
	21:5n-3	7.5% (2)		4.7 % (4)
	22:5n-6		7.2% (4)	3.0% (9)
	22:4n-3	5.0% (4)	2.6% (7)	4.2% (6)
	22:5n-3			
	22:6n-3		5.5% (5)	3.4% (8)

728   Fig. 1.

747   A)   B)

748   Fig 2.

760   Fig. 3.

763   Fig. 4.

 Appendix:

Table A1. List of 38 of 41 fatty acids shared by the two datasets, considered as either "dietary"

fatty acids or "extended-dietary" fatty acids and used in this study (see Iverson *et al*., 2004), and

the dietary component they are associated with, based on literature: Sargent *et al*., 1995; Brett &

Müller-Navarra, 1997; Kattner *et al*., 1998; Virtue *et al*., 2000; Budge et al., 2002; Dalsgaard *et*

al., 2003; Iverson *et al*., 2004; Käkelä *et al*., 2005; Alfaro *et al*., 2006; Tucker *et al*., 2008;

Ahlgren *et al*., 2009; Gladyshev *et al*., 2009; Loseto *et al*., 2009; Stowasser *et al*., 2009; ; Piché

et al., 2010; Mariash *et al*., 2011. The fatty acids highlighted are the one discarded because they

were not quantified for both lakes.

Dietary Fatty acids	Dietary component
14:0	Pelagic (Zooplankton) + diatom
16:0	Cannibalism or/and carnivorous
16:1n7	Benthic (bacterial synthesis + diatoms) Cannibalism or/and carnivorous
16:2n6	
16:2n4	Diatom
17:0	
16:3n-4	
16:4n3	
16:4n1	Diatom
18:0	Cannibalism or/and carnivorous
18:1n9	Pelagic (Zooplankton)
18:1n7	Benthic (bacterial synthesis + algal)
18:2n6	Terrestrial
18:2n4	
18:3n6	
18:3n4	
18:3n3	Terrestrial
18:3n1	
18:4n3	Pelagic (Zooplankton)
18:4n1	
20:0	*not used
20:1n11	Copepod (Iverson, 2009)
20:1n9	Pelagic (Calanoid copepods diet based)
20:1n7	Pelagic (Zooplankton)
20:2n9	*not used
20:2n6	
20:3n6	
20:4n6	Benthic (diatom)
20:3n3	
20:4n3	

20:5n3	Cannibalism or/and carnivorous
22:1n11	Copepod
22:1n9	Pelagic (Zooplankton)
22:1n7	
22:2n6	
21:5n3	
22:4n6	*not used
22:5n6	Pelagic (Calanoid copepods diet based)
22:4n3	
22:5n3	
22:6n3	Benthic (Pennate diatoms + dinoflagellates + bivalves) Cannibalism

780   Table A2. List of 24 fatty acids considered as "non-dietary" in this study.

Non-Dietary Fatty acids
12:0
12:1
13:1
14:1n9
14:1n7
14:1n5
14:0 iso
14:0 ante
15:0
15:1n8
15:1n6
15:0 iso
16:1n11
16:1n9
16:1n5
16:0 iso
7Methyl 16:0
17:1
17:0 iso
18:1n5
20:0
20:2n9
22:0
24:1n9

Fig. A1. PCA of all fatty acids (top panels), dietary fatty acids (middle panels), and non-dietary fatty acids (bottom panels) of each morph of Lake Trout from Great Bear Lake, Stannard Rock (Lake Superior), and Superior Shoal (Lake Superior), including the 12 outlier individuals. Vectors of individual fatty acids important to the positioning of lake trout are represented to the right of each PCA. Angles and lengths of vectors represent the direction and the strength, respectively, of relationships between variables and the principal components.