Dietary vs non-dietary fatty acid profiles of lake trout ecotypes from Lake Superior and

Great Bear Lake: Are fish really what they eat?

Chavarie, L[1,2] *, J. Hoffmann[3], A.M. Muir[4], C.C. Krueger[5], C.R. Bronte[6], K.L. Howland[7,8], C.P.

Gallagher[7], S.P. Sitar[9], M.J. Hansen[10], M.R. Vinson[11], L.F. Baker[3], L.L. Loseto[7], W. Tonn[8], and

H. Swanson[3]

[1]University of British Columbia, Biodiversity Center, 2212 Main Mall, Vancouver, BC, Canada,

V6T 1Z4

[2] Scottish Centre for Ecology and the Natural Environment, Institute of Biodiversity, Animal

Health & Comparative Medicine, University of Glasgow, Glasgow, UK

[3]University of Waterloo, Department of Biology, 200 University Avenue West, Waterloo,

Ontario, Canada, N2L 3G

[4]Great Lakes Fishery Commission, 2100 Commonwealth Blvd., Suite 100, Ann Arbor,

Michigan, USA, 48105

[5]Michigan State University, Center for Systems Integration and Sustainability, 1405 South

Harrison Road, 115 Manly Miles Building, East Lansing, Michigan USA 48823

[6]U.S. Fish and Wildlife Service, Green Bay Fish and Wildlife Conservation Office, New

Franken, WI, USA

[7]Fisheries and Oceans Canada, 501 University Crescent, Winnipeg, Canada

[8]University of Alberta, Department of Biological Sciences, CW 405 BioSciences Building,

Edmonton, Canada

[9]Michigan Department of Natural Resources, Marquette, MI, USA

[10]U.S. Geological Survey Great Lakes Science Center, Hammond Bay Biological Station, USA

(retired)

[11]U.S. Geological Survey Great Lakes Science Center, Ashland, WI, USA

*Corresponding author: chavarie@ualberta.ca

Abstract:

Fatty acids are well-established biomarkers used to characterize trophic ecology, food-web linkages, and the ecological niche of many different taxa. Most often, fatty acids that are examined include only those previously identified as "dietary" or "extended dietary" biomarkers. Fatty acids considered as non-dietary biomarkers, however, represent numerous fatty acids that can be extracted. Some studies may include non-dietary fatty acids (i.e., combined with dietary fatty acids), but do not specifically assess them, whereas in other studies, these data are discarded. In this study, we explored whether non-dietary biomarkers fatty acids can provide worthwhile information by assessing their ability to discriminate intraspecific diversity within and between lakes. Non-dietary fatty acids used as biomarkers delineated variation among regions, among locations within a lake, and among ecotypes within a species. Physiological differences that arise from differences in energy processing can be adaptive and linked to habitat use by a species' ecotypes, and likely explains why non-dietary fatty acids biomarkers can be a relevant tool to delineate intraspecific diversity. Little is known about the non-dietary-mediated differences in fatty acid composition, but our results showed that non-dietary fatty acids biomarkers can be useful tool in identifying variation.

Key words: Intraspecific diversity, salmonid, aquatic ecology, trophic ecology, Great Lakes, *Salvelinus namaycush*

Introduction:

Constraints on traditional methods to investigate diets of organisms within aquatic systems

have led to the development and use of biochemical tracers (Vinson and Budy 2010). Among

these, fatty acids have gained popularity as both qualitative and quantitative trophic markers that

reflect foraging patterns and food-web dynamics (Galloway et al. 2014, Iverson 2009). As such,

fatty acids have been used to characterize variation within populations and species (e.g.,

evolutionary units linked to trophic polymorphism; Logan et al. 2000, Scharnweber et al. 2016),

and to explore geographical (Hiltunen et al. 2016, Pomerleau et al. 2014, Quérouil et al. 2013) and

temporal variation in trophic ecology (e.g., seasonal and annual variations; Eloranta et al. 2013,

Hartwich et al. 2012).

Fatty acids are the "building blocks" of lipids and represent the largest constituent of

neutral lipids (e.g., triacylglycerols and wax esters) and polar phospholipids (Iverson 2009). The

array of fatty acids present in nature is exceptionally complex, routinely ~70 fatty acids can be

identified within an organism (Budge et al. 2006, Iverson 2009). The utility of fatty acid analyses

to reflect foraging patterns and food web dynamics relies on the assumption that lipids are broken

down into their constituent fatty acids and incorporated relatively unchanged into consumer tissues

(Howell et al. 2003, Iverson 2009, Iverson et al. 2004). The storage patterns of fatty acids depend

on the biochemical limitations of organisms to biosynthesize, modify chain-length, and introduce

double bonds into fatty acids, which culminate in vertebrates (Iverson 2009). Fatty acids that are

stored in predator tissues, with no or little modification from their prey, have been labelled as

"dietary" or "extended dietary" tracers (Budge et al. 2006, Iverson et al. 1997, 2004) and have

been the target of analyses in ecological studies. Accordingly, for purposes of this study, we use

the terminology of "dietary" versus "non-dietary" derived fatty acids based on the classification

of Iverson et al. (2004), who identified typical dietary fatty acid markers that are now used across

a variety taxa. Often, "non-dietary" fatty acids extracted from tissue samples are not examined,

and only those recognized as dietary are analyzed. In other studies, dietary and non-dietary fatty

acids may be combined to characterize food-web relationships, but information gained from

inclusion of non-dietary fatty acids is not specifically addressed (Hiltunen et al. 2019, Mariash et

al. 2017, McMeans et al. 2015, Taipale et al. 2019).

The exclusion or ignoring non-dietary fatty acids from analyses is not misguided, as the

purpose of most studies is to describe diet patterns, and this practice has resulted in reliable

information being produced across taxa (Galloway et al. 2014, Grosbois et al. 2017, Iverson 2009,

Iverson et al. 2004). However, it is unknown whether valid information for other research

questions is lost when investigators discard non-dietary fatty acids. Biological (e.g., phenotypic

and genetic) and environmental variables can affect lipid and fatty acid composition in fishes

(Olsen and Skjervold 1995), including temperature (Farkas et al. 1980, Olsen 1999), salinity

(Borlongan and Benitez 1992), and light (Ota and Yamada 1971). Thus, fatty acids not labelled as

"dietary" markers could be useful when the aim of a study is to delineate or better understand

intraspecific diversity. To investigate this, we compared non-dietary to dietary fatty acids

biomarkers among ecotypes of lake trout (*Salvelinus namaycush*) in Lake Superior and Great Bear

Lake, as these ecotypes represent important intraspecific diversity in these lakes.

Salmonids, such as lake trout, inhabit young ecosystems believed to be 10,000 to 15,000

years old, e.g., post-glacial lakes colonized from non-glaciated refugia. The depauperate

communities of post-glacial lakes are commonly characterized by reduced interspecific

competition and predation, which allows colonizers access to a relatively wide array of resources,

conditions that favour development of intraspecific diversity (McPhail 1993, Robinson and Wilson

1994, Smith and Skulason 1996). This high level of ecological opportunity, together with an

increase in intraspecific competition after colonization, can promote specialization and divergence

within a population, e.g., the development of groups of individuals with similar patterns of

resource use (Svanbäck et al. 2007). Phenotypic characteristics of individuals in such groups may

evolve as niches incorporate novel foraging resources, referred to as resource divergence

(Robinson and Parsons 2002, Skulason and Smith 1995). As niches diverge, ecotypes can develop

differences in morpholology, genetics, physiology, life-history, and/or behaviour (Bolnick et al.

2007, Schluter and McPhail 1992, Smith and Skulason 1996).

Intraspecific diversity in lake trout has been mostly linked to differences in depth

distribution and, not surprisingly, is best known from large (> 500km^2), deep lakes, such as Lake

Superior (Moore and Bronte 2001, Muir et al. 2014), Lake Mistassini (Zimmerman et al. 2007),

Great Slave Lake (Zimmerman et al. 2006), and Great Bear Lake (Chavarie et al. 2013). Although

lake trout diversification has often focused on isolation-by-depth (without excluding isolation-by-

adaptation) in large, deep lakes (e.g., Lake Superior), diversification also occurs in small lakes or

within shallow-water habitats (Bernatchez et al. 2016, Chavarie et al. 2013, Chavarie et al. 2016c,

Morissette et al. 2018).

A number of studies have assessed lake trout diets with fatty acids, either qualitatively

(Chavarie et al. 2016b, Happel et al. 2017a, Happel et al. 2017b, Hoffmann 2017) or quantitatively

(Happel et al. 2016a, Happel et al. 2016b), generating reliable information about dietary patterns.

However, pronounced and systematic non-dietary-mediated differences in composition of fatty

acids have been reported in lake trout, supporting the rationale that non-dietary fatty acids

biomarkers could be important in delineating intraspecific diversity within this species. Goetz et

al. (2013) found physiological differences between lean and siscowet ecotypes of lake trout that

reflected genetically based differences in lipid synthesis, metabolism, and transport (Goetz et al.

2010), which suggests information from non-dietary fatty acids biomarkers might be important,

especially if they have genetic-based mechanisms. Consequently, where intraspecific diversity is

manifested as physiological differences among ecotypes, non-dietary fatty acids biomarkers could

assist in identifying variation. Yet, little is known about the non-dietary-mediated differences in

fatty acid composition of fish (or other taxa) in an ecological context.

To investigate the use of non-dietary fatty acids biomarkers as an ecological tool, we

analyzed a set of non-dietary and dietary fatty acid biomarkers (as classified by the literature),

from lake trout ecotypes collected from two lakes that sustain intraspecific diversity along different

axes of intraspecific diversification (e.g., depth-dependent and depth-independent). Specifically,

we compared: 1) non-dietary and dietary fatty acids biomarkers among four ecotypes of lake trout

from each of Great Bear Lake and Lake Superior, and 2) how well non-dietary and dietary

biomarkers delineated intraspecific diversity within and between lakes. Answers to these

comparisons will help determine if non-dietary fatty acid biomarkers can discriminate fish

geographically and among groups within a species, and if they offer different perspectives than

dietary fatty acids biomarkers.

Material and methods

Study systems and intraspecific diversity of lake trout

Located in the northeast corner of Northwest Territories (Canada; 65°92′ N, 120°82′ W),

Great Bear Lake is the most northerly lake of its size (~31 000 km^2) and is the fifteenth deepest

freshwater lake in the world (Fig. 1; Johnson 1975). A UNESCO biosphere reserve, Great Bear

Lake is 250 km south of the Arctic Ocean and has characteristics typical of an Arctic lake. The

lake is ultra-oligotrophic and, despite its size, has a simple food web, supporting only 15 fish

species (Johnson 1975, MacDonald et al. 2004). The lake and its biota have remained relatively

isolated and unexploited and is one of the most pristine large lakes in North America. Great Bear

Lake has five semi-isolated arms, but due to sample sizes, data were pooled across multiple sites

(see Chavarie et al. 2016b for details).

Great Bear Lake sustains a noteworthy example of lake trout divergence (Fig. 2). With its

intraspecific diversity independent of depth-based segregation, the lake also presents an unusual

ecological framework for lake trout differentiation (Chavarie et al. 2016a). Currently, four

shallow-water ecotypes are described in Great Bear Lake; three are common (Ecotypes 1-3), and

one is rare (Ecotype 4). Ecotype 1 has the smallest head and jaws, intermediate fin lengths and

body shape intermediate between Ecotypes 2 and 3. The lean-like Ecotype 2 has the largest head

and jaws but the smallest fins, and a streamlined body shape. Ecotype 3 has the longest fins, a

robust body shape, and a sub-terminal mouth. Ecotype 4 has a thick and curved lower jaw, a

streamlined body shape, and the smallest caudal peduncle among the ecotypes. Although Ecotype

4 is a pelagic specialist, Ecotypes 1-3 have more general feeding habits, with varying degrees of

omnivory along a weak benthic-pelagic gradient (Chavarie et al. 2016a, Chavarie et al. 2016b).

Lake Superior is a post-glacial, oligotrophic lake between Canada and the USA (47°43′ N,

86°56′ W) (Fig. 1). Lake Superior is the largest lake in the world by surface area (82 100 km^2).

Most of the waters of Lake Superior can be classified as offshore; 77% of the total area is greater

than 80 m deep (maximum depth = 406 m) (see Gorman et al. 2012, Hoffmann 2017, Horns et al.

2003). Lake Superior supports 87 fish species, with lake trout as a main target of the commercial

fishery since the 1800s. Lake trout were sampled from two sites, Superior Shoal (48° 3'43.54" N,

87° 8'52.57" W) and Stannard Rock (47°12'26.26" N, 87°12'3.82" W) (Fig. 1).

Lake Superior supports one of the highest levels of sympatric diversity expressed within

lake trout (Fig. 2). Four ecotypes are currently recognized; the siscowet, humper, and redfin

ecotypes inhabit deep-water (> 70 m), whereas the lean ecotype occupies shallow-water habitats

(< 70 m) (Bronte et al. 2003, Bronte and Moore 2007, Muir et al. 2014). The siscowet is

characterized by a large head, short snout, long maxilla, large eye, short and deep caudle peduncle,

and moderately long paired fins (Muir et al. 2015). Humpers have a small head, short snout, short

maxillae, large eyes, and short and narrow caudal peduncle. (Moore and Bronte 2001, Muir et al.

2015). The redfin has the largest head, snout, and eyes, the longest and deepest caudal peduncle,

and much longer pelvic and pectoral fins than the other ecotypes (Muir et al. 2014). Finally, lean

lake trout have a large, narrow, and pointed head, long snout, small eyes, long and narrow caudal

peduncle, short paired fins, and low body lipid content. (Endler 1978, Khan and Qadri 1970, Muir

et al. 2015). As vertical migrating visual predators, the three deep-water ecotypes are likely feeding

mostly on *Mysis* and deep-water ciscoes (*Coregonus artedi* complex) (Hoffmann 2017, Hrabik et

al. 2006, Muir et al. 2014), whereas the lean ecotype is adapted for daytime predation on pelagic

fishes in shallow-water habitats (piscivorous feeding strategy; Harvey and Kitchell 2000, Harvey

et al. 2003, Janhunen et al. 2009).

Fatty acids

Dorsal muscle samples (Budge et al. 2011) from lake trout in both lakes were stored at -20

°C (Budge et al. 2006, Chavarie et al. 2016b, Kavanagh et al. 2010, Loseto et al. 2009). Lipids

were extracted from 1 g of the homogenate material; after passive overnight extraction (at -20) in

2:1 chloroform:methanol containing 0.01% BHT (v/v/w) (Folch et al. 1957), samples were filtered

through Whatman Grade 1 Qualitative filter paper and the filter paper sample was rinsed twice

with 2 mL of 2:1 chloroform:methanol. Sample extract was collected in a test tube and 7 mL of

0.88 NaCl solution were added to encourage fatty acids to move into the organic (chloroform)

layer. The aqueous layer was discarded, after which the chloroform was dried with sodium sulfate

prior to total lipid determination. The extracted lipid was used to prepare fatty acid methyl esters

(FAME) by transesterification with Hilditch reagent (0.5 N H_2SO_4 in methanol) (Morrison and

Smith 1964). Samples were heated for 1 h at 100 °C. Gas chromatographic (GC) analysis was

performed on an Agilent Technologies 7890N GC equipped with a 30 m J&W DB-23 column

(0.25 mm I.D; 0.15 μm film thickness). The GC was coupled to a Flame Ionization Detector

operating at 350 °C. Hydrogen was used as carrier gas flowing at 1.25 mL/min for 14 minutes,

increasing to 2.5 mL/min for 5 minutes. The split/splitless injector was heated to 260 °C and run

in splitless mode. The oven program was as follows: 60 °C for 0.66 min, increasing by 22.82

°C/min to 165 °C with a 1.97 min hold; increasing by 4.56 °C/min to 174 °C and by 7.61 °C/min

to 200 °C with a 6 min hold. Peaks were quantified using Agilent Technologies ChemStation

software. Fatty acids standards were obtained from Supelco (37 component FAME mix) and

Nuchek (54 component mix GLC-463). FAMEs were identified via retention time and known

standard mixtures and are reported as percentages of total fatty acids. Fatty acid standards were

obtained from Supelco - Oakville Ontario, Canada (37 component FAME mix) and Nuchek -

Elysian Minnesota, USA (54 component mix GLC-463). Standards were run to allow creation of

a 4 level calibration curve for each set of samples at the start of each sample set. A standard was

repeated every 10 samples thereafter. Every 10 th sample was injected in duplicate. All fatty acids

values were converted to a mass percentage of the total array, and were named according the

IUPAC nomenclature as X:Y n-z, where X is the number of carbons in the fatty acid, Y is the

number of methylene-interrupted double bonds present in the chain, and n-z denotes the position

of the last double bond relative to the methyl terminus (Ronconi et al. 2010). All laboratory

analyses were conducted at the Freshwater Institute, Fisheries and Oceans Canada, Winnipeg MB.

For both lakes, tissue samples were taken from indivual lake trout previously identified to

corresponding ecotypes: 126 samples were collected from Great Bear Lake (Ecotype 1 = 32,

Ecotype 2 = 35, Ecotype 3 = 38, and Ecotype 4 = 21, see Chavarie et al. 2016b for more details)

and 210 samples were collected from Lake Superior (60 siscowet, 60 humper, 30 redfin, and 60

lean, see Hoffmann 2017 for more details). Overall, fatty acid analysis procedures were divided

into two steps, using non-dietary and dietary fatty acid biomarkers (see Appendix 1), following

the methods of Iverson et al. (1997, 2004) and Budge et al. (2006) to identify dietary fatty acids

biomarkers. Dietary biomarkers were defined as fatty acids $\geq$ 14 carbons that are generally

incorporated into animal tissue from the diet with no or little modification (e.g., rather than come

from biosynthesis) (Iverson 2009). Thirty-eight dietary and 24 non-dietary fatty acids biomarkers

were found to be shared between the two lakes, and these were selected for further analyses (Table

A1 and A2).

Statistical Analyses

Unless noted otherwise, statistical analyses were conducted using R software version 3.5.3

(R Core Team 2017). Prior to analysis, fatty acid concentrations were logit transformed (log(p/(1-

p)) to normalize the data, and then scaled and centered using a z-score transformation ($z=x\mu/\sigma$).

(Clemmensen et al. 2011, Witten and Tibshirani 2011). Principal Component Analysis (PCA) was

performed on all dietary and non-dietary fatty acid biomarkers to provide inference about patterns

of variation among locations and ecotypes (Chavarie et al. 2016b). PCA summarizes similarities

and differences among individuals, based on their fatty acid profiles, independent of ecotype and

location (Chavarie et al. 2016b). For PCAs, 12 outlier individuals from Superior Shoal were

excluded because PCAs are sensitive to outlier variation (but see Fig. A1 for PCAs with outliers

included) (Filzmoser et al. 2009, Kriegel et al. 2008).

To test for differences in fatty acid composition among ecotypes within each lake (dietary

and non-dietary), we used Permutational Multivariate Analysis of Variance (PERMANOVA; a

non-parametric analog of Multivariate Analysis of Variance), followed by post-hoc comparisons

with Bonferroni corrections. PERMANOVAs were performed in PAST 3 (Hammer et al. 2001)

using 9999 permutations. A similarity percentage routine (SIMPER) using Bray-Curtis was used

to determine which fatty acids (dietary and non-dietary) were primarily responsible for observed

differences among ecotypes for each lake (King and Jackson 1999). We also performed linear

discriminant analysis on fatty acids (dietary and non-dietary) to delineate differences among

ecotypes at each location. A jacknife validation procedure, using 20% of our data as unknown,

provided a classification success metric to assess how distinct ecotypes appeared in each fatty

acids dataset (dietary vs non-dietary).

Results

Combined lakes analyses

The first two axes of the PCAs explained 42.4 % and 41.5%, respectively, of the variation

among individuals in their non-dietary and dietary fatty acid biomarkers (Fig. 3). In both PCAs,

lake trout from Great Bear Lake were largely distinct from Lake Superior trout (only ~30

individuals from Great Bear Lake overlapped with individuals from Lake Superior), but trout from

the two Lake Superior sites, Stannard Rock and Superior Shoal, overlapped. The non-dietary fatty

acids 13:1, 14:1n-7, 15:0, 15:1n8, 15:1n6, 15:0 iso, 16:1n11, 17:0 iso, 20:0, 22:0, 20:2n9, and

24:1n9 contributed to the separation between the two lakes. Separation between the two lakes in

the dietary PCA appeared to be driven by fatty acids associated with pelagic habitat (14:0, 20:1n-

9, 20:1n-7, 20:1n-11, and 22:1n-9; toward Lake Superior) versus one dietary fatty acid associated

with cannibalism or/and carnivory (20:5n-3; toward Great Bear Lake) (Appendix, Table 1).

Finally, the first two axes of the PCA based on all fatty acids combined explained 39.0% of the

variation among lake trout ecotypes from Great Bear Lake and Lake Superior. As before, lake trout

from Great Bear Lake were largely separated from the Lake Superior trout, whereas lake trout

from Stannard Rock and Superior Shoal in Lake Superior overlapped completely.

Lake Trout intraspecific diversity of Great Bear Lake

Composition of both non-dietary and dietary fatty acid biomarkers were able to

discriminate among the four lake trout ecotypes from Great Bear Lake. Non-dietary fatty acid

biomarkers differed among the four Great Bear Lake ecotypes (one-way PERMANOVA, $F_{3,122} =$

3.6, $P < 0.01$). Similarly, comparison of non-dietary fatty acids biomarkers showed that all pairs

of ecotypes differed from one another ($P < 0.01$) except for ecotypes 1 and 3. The ten most

discriminating non-dietary fatty acid biomarkers from SIMPER explained 63.8% of the

dissimilarity among groups (Table 1). The first two axes of the linear discriminant analysis

explained 55.7 % and 27.8% of the variation, and 57.9% of all individuals were correctly classified

to ecotype based on non-dietary fatty acids biomarkers (Fig. 4). Dietary fatty acid biomarkers also

differed among the four ecotypes from Great Bear Lake (one-way PERMANOVA, $F_{3,122} = 2.95$, P

< 0.01), and most ecotypes differed from each other (all pairwise $P < 0.01$ except for ecotypes 1

vs. 3). The ten most discriminating dietary fatty acid biomarkers from SIMPER explained 48.8%

of the dissimilarity among groups (Table 2). The first two axes of the linear discriminant analysis

explained 46.7% and 39.0% of the variation, and 68.3% of all individuals were correctly classified

to ecotype based on dietary fatty acids biomarkers (Fig. 4).

Lake Trout intraspecific diversity of Lake Superior, Stannard Rock

Non-dietary fatty acid biomarkers differed among the three ecotypes of lake trout from

Stannard Rock (one-way PERMANOVA, $F_{2,87} = 3.9$, $P < 0.01$), and leans differed significantly

from both siscowets and humpers ($P < 0.01$). The ten most discriminating non-dietary fatty acids

biomarkers from SIMPER explained 70.9% of the dissimilarity among ecotypes (Table 1). The

first two axes of the linear discriminant analysis explained 60.2% and 39.8% of the variation,

respectively, and 52.2% of all individuals were correctly classified to ecotype based on non-dietary

fatty acids biomarkers (Fig. 4). In contrast to results from non-dietary fatty acid biomarkers, no

differences in dietary fatty acid composition occurred among the three lake trout ecotypes from

Stannard Rock, Lake Superior (one-way PERMANOVA, $F_{2,87} = 1.12$, $P = 0.3$). The ten most

discriminating fatty acids from SIMPER explained 55.2 % of the dissimilarity among groups

(Table 2), and the first two axes of the linear discriminant analysis explained 60.9% and 39.1% of

the variation. Fifty percent of individuals were correctly classified to ecotype based on dietary

fatty acids biomarkers (Fig. 4).

Lake Trout intraspecific diversity of Lake Superior, Superior Shoal

Like Stannard Rock, composition of non-dietary fatty acid biomarkers differed

significantly among the four ecotypes (one-way PERMANOVA, $F_{3,116} = 2.5$, $P < 0.01$); leans

differed from redfins ($P < 0.01$) and differences were marginally between redfin and siscowet and

between redfin and humper ($P < 0.06$). The ten most discriminating non-dietary fatty acids

biomarkers from SIMPER explained ~69.9% of the dissimilarity among groups (Table 1). The

first two axes of the linear discriminant analysis explained 64.3% and 25.6% of the variation, and

45.0% individuals were correctly classified to ecotype using non-dietary fatty acids biomarkers

(Fig. 4). Similar to what was found at Stannard Rock, we found no differences among the four

ecotypes from Superior Shoal based on dietary fatty acid composition (one-way PERMANOVA,

$F_{3,116}$ = 0.8, P = 0.3). The ten most discriminating fatty acids from SIMPER explained 55.8 % of

the dissimilarity among groups (Table 2). The first two axes of the linear discriminant analysis

explained 38.6% and 35.6% of the variation, respectively, and 31.7% of fish were correctly

classified to ecotype based on their dietary fatty acids (Fig. 4).

Discussion

In this study, non-dietary fatty acids showed variation geographically (between lakes),

between locations within a lake (i.e., Stannard Rock versus Superior Shoal), and among ecotypes

within a lake or location. Although some overlap existed (which reduced the power to

discriminate), our results showed that when investigating intraspecific diversity, non-dietary fatty

acids biomarkers can be a useful tool to delineate groups, and that sometimes, such as at sites in

Lake Superior, were more discriminatory than dietary fatty acids. While characterizing trophic

divergence, food-web linkages, and time-integrated niche use across a large array of taxa, most

investigators use a common set of fatty acids that are selected from lists that classify fatty acids as

"dietary" or "extended dietary" biomarkers (Budge et al. 2006, Iverson et al. 2004). However, our

results suggest that discarding non-dietary fatty acid data as a matter of course may result in

inadvertent loss of information.

Not all fatty acids provide equivalent information about diet due to metabolism and *de novo*

synthesis (Iverson et al. 1993, Iverson et al. 2004). Metabolism, however, can differ within species,

as physiological differences often exists among sets of individuals, i.e., "ecotypes" (Miles et al.

2007, Pryke et al. 2007). Such physiological differences within a species may result in non-dietary

fatty acids biomarkers being a relevant tool to delineate intraspecific diversity, as we show here

for lake trout. In Lake Superior, fat content has long been recognized as an important characteristic

that distinguishes lake trout ecotypes along a depth axis (Eschmeyer and Phillips 1965, Eshenroder

2008), and these lipid differences have been linked to genetic differences among ecotypes

(Eschmeyer and Phillips 1965, Goetz et al. 2013, Goetz et al. 2010). Thus, fatty acid deposition

and metabolism appear to have undergone selection along a depth gradient likely in part to

contribute to buoyancy compensation, with differences reported between siscowet (deep-water

ecotype) and lean (shallow-water ecotype) lake trout (Eschmeyer and Phillips 1965, Goetz et al.

2010). Until now, no information has been available for redfin and humper ecotypes.

In addition to the variation in lipid accumulation, Goetz et al. (2013) also found differences

in lipid composition between siscowet and lean ecotypes. Because a common garden design was

used in their experiment, the higher proportion of polyunsaturated fatty acids (PUFAs) found in

the muscle lipid profile of siscowet than in lean lake trout could not be attributed to differences in

diet. The differences in non-dietary fatty acids biomarkers we observed among the four ecotypes

of lake trout from Lake Superior is consistent with the concept of metabolotypes (Goetz et al.

2013). Altogether, differences in energy processing and storage between lean and siscowet lake

trout in Lake Superior are adaptive to their respective habitats, deep- vs. shallow-water, and to

their life-histories (Goetz et al. 2013). Lipid content, intertwined with buoyancy variations, have

been linked to differences in depth distributions and swimming tactics among lake trout ecotypes

(Zimmerman et al. 2006, Zimmerman et al. 2007, Zimmerman et al. 2009). While deep-water

ecotypes use hydrostatic lift related to lipid content to enhance vertical migration while foraging

for *Mysis* and cisco (Henderson and Anderson 2002), the shallow-water lean ecotype likely relies

more on hydrodynamic lift, linked to the cruising movements of pelagic predators (Webb 1984).

Consistent with the findings of Goetz et al. (2013), we found that non-dietary fatty acid biomarkers

differed between shallow- and deep-water ecotypes at Stannard Rock. However, at Superior Shoal,

non-dietary fatty acids biomarkers differed only between the shallow-water lean ecotype and one

deep-water ecotype – the redfin. This observation, along with some more subtle differences in non-

dietary fatty acid biomarkers among deep-water ecotypes, requires further study as our current

knowledge is limited. It is presently unclear if these lake trout ecotypes have adapted

physiologically to their habitat leading to metabolotypes and how diets differ among all ecotypes

temporally and spatially within Lake Superior. As such, feeding on the same item by different

ecotypes might produce different fatty acid accumulations resulting not only in a physiologically

interesting questions but also questioning the accuracy of predictions about diet composition.

The concept that dynamics of energy processing and storage are adaptive along a gradient

associated with depth does not apply to the intraspecific diversity of lake trout in Great Bear Lake.

This diversity is limited to shallow-water habitat, and appears to be independent of major habitat

and resource partitioning (Chavarie et al. 2016a, Chavarie et al. 2018). The similarity of results

between Great Bear Lake and Lake Superior is thus perplexing, as we were expecting greater

differences in non-dietary fatty acid biomarkers among ecotypes in Lake Superior than in Great

Bear Lake, due to known buoyancy variation associated with a depth gradient in Lake Superior. If

lake trout ecotypes from Great Bear Lake are also under selection (Harris et al. 2014), differences

in energy processing and storage may be as pronounced as those that have been observed in Lake

Superior. Another question raised by our results is the extent to which ecotypes are independent

of major habitat or resource axes (the same question is pertinent for Lake Superior), especially

because dietary fatty acid biomarkers in Great Bear Lake were slightly better at delineating

intraspecific groups than non-dietary biomarkers than in Lake Superior.

Differences between morphs and sites may be influenced by the total content of fatty acids

(μg FA/m) in muscle, rather than by their composition (%). Tissue-specific storage of fatty

acids can vary across space or time, and some fish store lipids as modified adipose or lipid pockets

in their muscle (e.g., salmonids; Iverson 2009, Sasaki et al. 1989). In a comparison of fatty acid

content between dorsal muscle vs. belly tissue, (Happel et al. 2019) found a threshold response;

the tissues became increasingly dissimilar when lipid content of muscle was >

~10% . Nevertheless, Happel et al. (2019) found that fatty acid profiles were specific to each of

the five lakes they examined, i.e., lake trout displayed broad variation among locations.

Ecological differences in allopatry are often more pronounced than those in sympatry, due

to disparate environments and isolation (Fraser et al. 2011, Heggenes 2002, Rundle and Nosil

2005, Yoder et al. 2010). Our results were consistent with this trend, i.e., differences between lakes

were greater than differences among ecotypes within a lake for both dietary and non-dietary fatty

acid biomarkers. For dietary fatty acids, differences between the two lakes appeared to be due to

fatty acids associated with a pelagic environment, such as C20 and C22 monounsaturates, that can

be used as biomarkers of food webs based on pelagic copepods (Ahlgren et al. 2009, Budge et al.

2006, Dalsgaard et al. 2003). Although benthic productivity generally dominates in Arctic lakes

(Chavarie et al. 2018, Johnson 1975), few lake trout in Great Bear Lake specialized on pelagic

resources (i.e., few ecotypes 2 and 4 had fatty acid signatures that overlapped with lake trout from

Lake Superior; Chavarie et al., submitted; Chavarie et al. 2016a, Chavarie et al. 2016b).

With the general benthic orientation of lake productivity in Arctic regions, combined with

the known distributions in deep-water habitats of lake trout ecotypes from Lake Superior (50-150

m; Muir et al. 2014), our overall results reflected the expected ecological differences of this species

from these two systems. Despite these inter-lake differences, similar dietary (e.g., five out of 10)

and non-dietary (e.g., 7 out of 10) fatty acid biomarkers were important in identifying intraspecific

diversity within each lakes. The greater number of shared non-dietary fatty acids biomarkers

discriminating lake trout intraspecific diversity in the two lakes supports the idea of similar

physiological differences (e.g., energy processing and storage) among ecotypes from both lakes.

A few caveats should be noted that could alter interpretations of this study. First, we cannot

ensure the fatty acid biomarkers defined based on Iverson et al. (2004) truly reflect dietary vs non-

dietary origins, due to the lack of taxa-specific studies on the integration of prey fatty acids. Despite

this uncertainty, the aim of this paper was to examine loss of information from discarding fatty

acids generally considered as non-dietary biomarkers by the literature. Second, no spatial

component was defined for the Great Bear Lake dataset, due to small sample sizes from each

location, which may have introduced variation into the results. The importance of spatial variation

in large, complex systems was shown here for Lake Superior (Chavarie et al. 2015, Hoffmann

2017) and environmental variables (e.g., temperature, light), which may vary spatially, are known

to alter lipid composition in fish tissue (Olsen 1999). Third, multiple sizes and life-stages, e.g.,

juvenile, mature, and resting individuals, were included in the Lake Superior analysis; different

life-stages can vary in lipid metabolism (Sheridan 1989) and large lake trout can rely more on

nearshore-benthic food web resources than small lake trout (Happel et al. 2017a). Finally, some

fatty acids exist at very low amounts ($\leq$ 2%), which can introduce error when interpreting

differences among fatty acids that are found in only trace amounts (e.g., peak shouldering)

(Christie 1998). Despite these limitations, we found some consistent patterns with regards to

intraspecific diversity between lakes and among ecotypes within lakes from two distinct datasets.

Conclusion

Our study demonstrated the potential benefits of using both dietary and non-dietary fatty

acid biomarkers for delineating variation within a species. In some instances, non-dietary fatty

acids were better for discriminating ecotypes than dietary ones, in contrast to the popular maxim

associated with trophic markers, "you are what you eat". Dietary fatty acid biomarkers can

document the occurrence of discrete niche use among sets of individuals (i.e., ecotypes) within a

species (Chavarie et al. 2016b). The fatty-acid composition of individuals that reflects their diet

has been validated for lake trout (Happel et al. 2016a, Happel et al. 2016b). However,

physiological differences in the dynamics of energy processing (e.g., metabolism) and storage

can also be adaptive among ecotypes (Eschmeyer and Phillips 1965, Goetz et al. 2013, Goetz et

al. 2010), which may result differences in non-dietary fatty acids (e.g., greater than dietary fatty

acids) and thus, be useful molecular tools. The lack of information on non-dietary fatty acids in

lake trout (and other taxa), however, raises questions about their physiological role and if

consumers acquire them or not from their diet (e.g., synthesis de novo, elongation). This would

be a fruitful area for future research.

Ultimately, the relative importance of dietary vs. non-dietary fatty acid biomarkers depends

on the question being asked. In our study, non-dietary fatty acids were valuable in delineating

intraspecific variation within a lake, but also in examining differences between lakes. Non-dietary

fatty acid biomarkers can provide useful information; therefore, one should carefully consider if

such information is superfluous or not before data from these fatty acids are discarded.

Acknowledgement

We thank Déline Renewable Resources Council, Déline Lands and Finance Corporation,

the community of Déline, DFO in Hay River, and the Department of Environment and Natural

Resources in Déline, which provided valuable help with field planning and logistics. Financial

support was provided by Fisheries and Oceans Canada (DFO), Natural Sciences and Engineering

Research Council of Canada, Sahtu Renewable Resource Board, Association of Canadian

Universities for Northern Studies, Canadian Circumpolar Institute's Circumpolar/Boreal Alberta

Research and Northern Scientific Training Program, D. Alan Birdsall Memorial Scholarship Fund,

Aboriginal Affairs and Northern Development Canada Northwest Territories Cumulative Impacts

Monitoring Program grants, and the Great Lakes Fishery Commission. Logistical and in-kind

support were provided by the Polar Continental Shelf Program and USGS. The findings and

conclusions in this article are those of the authors and do not necessarily represent those of the

U.S. Geological Survey or the U.S. Fish and Wildlife Service.

Data Availability Statement:

All data presented are available by request via e-mail to the first author.

Conflict Of Interest

The authors declare that they have no conflict of interest.

Literature:
Ahlgren, G., Vrede, T., and Goedkoop, W. 2009. Fatty Acid Ratios in Freshwater Fish, Zooplankton and
Zoobenthos – Are There Specific Optima? *In* Lipids in Aquatic Ecosystems. *Edited by* M. Kainz, M.T. Brett
and M.T. Arts. Springer New York, New York, NY. pp. 147-178.
Alfaro, A.C., Thomas, F., Sergent, L., and Duxbury, M. 2006. Identification of trophic interactions within an
estuarine food web (northern New Zealand) using fatty acid biomarkers and stable isotopes. Estuarine,
Coastal and Shelf Science **70**(1): 271-286.
Bernatchez, S., Laporte, M., Perrier, C., Sirois, P., and Bernatchez, L. 2016. Investigating genomic and
phenotypic parallelism between piscivorous and planktivorous lake trout (*Salvelinus namaycush*) ecotypes
by means of RADseq and morphometrics analyses. Molecular Ecology **25**(19): 4773-4792.
Bolnick, D.I., Svanbäck, R., Araújo, M.S., and Persson, L. 2007. Comparative support for the niche variation
hypothesis that more generalized populations also are more heterogeneous. Proceedings of the National
Academy of Sciences **104**(24): 10075-10079.
Borlongan, I.G., and Benitez, L.V. 1992. Lipid and fatty acid composition of milkfish (Chanos chanos
Forsskal) grown in freshwater and seawater. Aquaculture **104**(1): 79-89.
Bronte, C.R., Ebener, M.P., Schreiner, D.R., DeVault, D.S., Petzold, M.M., Jensen, D.A., Richards, C., and
Lozano, S.J. 2003. Fish community change in Lake Superior, 1970-2000. Canadian Journal of Fisheries and
Aquatic Sciences **60**(12): 1552-1574.
Bronte, C.R., and Moore, S.A. 2007. Morphological variation of siscowet lake trout in Lake Superior. Trans
Am Fish Soc **136**(2): 509-517.
Budge, S.M., Iverson, S.J., Bowen, W.D., and Ackman, R.G. 2002. Among- and within-species variability in
fatty acid signatures of marine fish and invertebrates on the Scotian Shelf, Georges Bank, and southern
Gulf of St. Lawrence. Canadian Journal of Fisheries and Aquatic Sciences **59**(5): 886-898.

Budge, S.M., Iverson, S.J., and Koopman, H.N. 2006. Studying trophic ecology in marine ecosystems using
fatty acids: a primer on analysis and interpretation. Marine Mammal Science **22**(4): 759-801.

Budge, S.M., Penney, S.N., and Lall, S.P. 2011. Response of tissue lipids to diet variation in Atlantic salmon
(Salmo salar): Implications for estimating diets with fatty acid analysis. J Exp Mar Biol Ecol **409**(1): 267-
274.

Chavarie, L., Harford, W.J., Howland, K.L., Fitzsimons, J., Muir, A.M., Krueger, C.C., and Tonn, W.M. 2016a.
Multiple generalist morphs of Lake Trout: Avoiding constraints on the evolution of intraspecific
divergence? Ecology and Evolution **6**: 7727–7741.

Chavarie, L., Howland, K., Gallagher, C., and Tonn, W. 2016b. Fatty acid signatures and stomach contents
of four sympatric Lake Trout: assessment of trophic patterns among morphotypes in Great Bear Lake.
Ecology of Freshwater Fish **25**: 109-124.

Chavarie, L., Howland, K., Harris, L., and Tonn, W. 2015. Polymorphism in lake trout in Great Bear Lake:
intra-lake morphological diversification at two spatial scales. Biol J Linn Soc **114**(1): 109-125.

Chavarie, L., Howland, K.L., Harris, L.N., Hansen, M.J., Harford, W.J., Gallagher, C.P., Baillie, S.M., Malley,
B., Tonn, W.M., Muir, A.M., and Krueger, C.C. 2018. From top to bottom: Do Lake Trout diversify along a
depth gradient in Great Bear Lake, NT, Canada? PLoS One **13**(3): e0193925.

Chavarie, L., Howland, K.L., and Tonn, W.M. 2013. Sympatric polymorphism in lake trout: the coexistence
of multiple shallow-water morphotypes in Great Bear Lake. Transactions of the American Fisheries Society
142(3): 814-823.

Chavarie, L., Muir, A.M., Zimmerman, M.S., Baillie, S.M., Hansen, M.J., Nate, N.A., Yule, D.L., Middel, T.,
Bentzen, P., and Krueger, C.C. 2016c. Challenge to the model of lake charr evolution: shallow- and deep-
water morphs exist within a small postglacial lake. Biological Journal of the Linnean Society: n/a-n/a.

Clemmensen, L., Witten, D., Hastie, T., and Ersbøll, B. 2011. Sparse Discriminant Analysis. Technometrics
53(4): 406-413.

Dalsgaard, J., St. John, M., Kattner, G., Müller-Navarra, D., and Hagen, W. 2003. Fatty acid trophic markers
in the pelagic marine environment. *In* Adv Mar Biol. Academic Press. pp. 225-340.

Eloranta, A.P., Mariash, H.L., Rautio, M., and Power, M. 2013. Lipid-rich zooplankton subsidise the winter
diet of benthivorous Arctic charr (Salvelinus alpinus) in a subarctic lake. Freshw Biol **58**(12): 2541-2554.

Endler, J.A. 1978. A predator's view of animal color patterns. *In* Evolutionary Biology. *Edited by* M.K. Hecht,
W.C. Steere and B. Wallace. Plenum Press, New York and London.

Eschmeyer, P.H., and Phillips, A.M., Jr. 1965. Fat content of the flesh of siscowets and lake trout from Lake
Superior. Trans Am Fish Soc **1965**: 62-74.

Eshenroder, R.L. 2008. Differentiation of deep-water lake charr *Salvelinus namaycush* in North American
lakes. Environ Biol Fishes **83**: 77-90.

Farkas, T., Csengeri, I., Majoros, F., and Oláh, J. 1980. Metabolism of fatty acids in fish: III. Combined effect
of environmental temperature and diet on formation and deposition of fatty acids in the carp, Cyprinus
carpio Linnaeus 1758. Aquaculture **20**(1): 29-40.

Filzmoser, P., Hron, K., and Reimann, C. 2009. Principal component analysis for compositional data with
outliers. Environmetrics **20**(6): 621-632.

Folch, J., Lees, M., and Sloane-Stanley, G. 1957. A simple method for the isolation and purification of total
lipids from animal tissues. J biol Chem **226**(1): 497-509.

Fraser, D.J., Weir, L.K., Bernatchez, L., Hansen, M.M., and Taylor, E.B. 2011. Extent and scale of local
adaptation in salmonid fishes: review and meta-analysis. Heredity **106**: 404.

Galloway, A.W.E., Eisenlord, M.E., Dethier, M.N., Holtgrieve, G.W., and Brett, M.T. 2014. Quantitative
estimates of isopod resource utilization using a Bayesian fatty acid mixing model. Marine Ecology Progress
Series **507**: 219-232.

Gladyshev, M.I., Sushchik, N.N., Anishchenko, O.V., Makhutova, O.N., Kalachova, G.S., and Gribovskaya, I.V. 2009. Benefit-risk ratio of food fish intake as the source of essential fatty acids vs. heavy metals: A case study of Siberian grayling from the Yenisei River. Food Chemistry **115**(2): 545-550.

Goetz, F., Jasonowicz, A., Johnson, R., Biga, P., Fischer, G., and Sitar, S. 2013. Physiological differences between lean and siscowet lake trout morphotypes: Are these metabolotypes? Canadian Journal of Fisheries and Aquatic Sciences **71**(3): 427-435.

Goetz, F., Rosauer, D., Sitar, S.P., Goetz, G., Simchick, C., Roberts, S., Johnson, R., Murphy, C., Bronte, C.R., and Mackenzie, S. 2010. A genetic basis for the phenotypic differentiation between siscowet and lean lake trout (*Salvelinus namaycush*). Mol Ecol **19 (Suppl. 1)**: 176-196.

Gorman, O.T., Yule, D.L., and Stockwell, J.D. 2012. Habitat use by fishes of Lake Superior. II. Consequences of diel habitat use for habitat linkages and habitat coupling in nearshore and offshore waters. Aquat Ecosyst Health Manag **15**(3): 355-368.

Grosbois, G., Mariash, H., Schneider, T., and Rautio, M. 2017. Under-ice availability of phytoplankton lipids is key to freshwater zooplankton winter survival. Scientific Reports **7**(1): 11543.

Hammer, Ø., Harper, D., and Ryan, P. 2001. Past: paleontological statistics software package for education and data analysis. Palaeontologia Electronica **4**: 1-9.

Happel, A., Jonas, J.L., McKenna, P., Rinchard, J., He, J.X., and Czesny, S.J. 2017a. Spatial Variability of Lake Trout Diets in Lakes Huron and Michigan Revealed by Stomach Content and Fatty Acid Profiles. Canadian Journal of Fisheries and Aquatic Sciences.

Happel, A., Pattridge, R., Walsh, M., and Rinchard, J. 2017b. Assessing diet compositions of Lake Ontario predators using fatty acid profiles of prey fishes. Journal of Great Lakes Research.

Happel, A., Stafford, C.P., Rinchard, J., and Czesny, S. 2019. Fatty acid profiles of lake trout reveal the importance of lipid content for interpreting trophic relationships within and across lakes. Journal of Great Lakes Research.

Happel, A., Stratton, L., Kolb, C., Hays, C., Rinchard, J., and Czesny, S. 2016a. Evaluating quantitative fatty acid signature analysis (QFASA) in fish using controlled feeding experiments. Canadian Journal of Fisheries and Aquatic Sciences **73**(8): 1222-1229.

Happel, A., Stratton, L., Pattridge, R., Rinchard, J., and Czesny, S. 2016b. Fatty-acid profiles of juvenile lake trout reflect experimental diets consisting of natural prey. Freshwater Biology **61**(9): 1466-1476.

Harris, L.N., Chavarie, L., Bajno, R., Howland, K.L., Wiley, S.H., Tonn, W.M., and Taylor, E.B. 2014. Evolution and origin of sympatric shallow-water morphotypes of Lake Trout, Salvelinus namaycush, in Canada's Great Bear Lake. Heredity **114**: 94.

Hartwich, M., Martin-Creuzburg, D., and Wacker, A. 2012. Seasonal changes in the accumulation of polyunsaturated fatty acids in zooplankton. J Plankton Res **35**(1): 121-134.

Harvey, C.J., and Kitchell, J.F. 2000. A stable isotope evaluation of the structure and spatial heterogeneity of a Lake Superior food web. Can J Fish Aquat Sci **57**: 1395-1403.

Harvey, C.J., Schram, S.T., and Kitchell, J.F. 2003. Trophic relationships among lean and siscowet lake trout in Lake Superior. Transactions of the American Fisheries Society **132**(2): 219-228.

Heggenes, J. 2002. Flexible Summer Habitat Selection by Wild, Allopatric Brown Trout in Lotic Environments. Trans Am Fish Soc **131**(2): 287-298.

Henderson, B.A., and Anderson, D.M. 2002. Phenotypic differences in buoyancy and energetics of lean and siscowet lake charr in Lake Superior. Environ Biol Fishes **64**: 203-209.

Hiltunen, M., Peltomaa, E., Brett, M.T., Aalto, S.L., Strandberg, U., Oudenampsen, J., Burgwal, L.M., and Taipale, S.J. 2019. Terrestrial organic matter quantity or decomposition state does not compensate for its poor nutritional quality for Daphnia. Freshwater Biology **64**(10): 1769-1786.

Hiltunen, M., Taipale, S.J., Strandberg, U., Kahilainen, K.K., and Kankaala, P. 2016. High intraspecific variation in fatty acids of Eudiaptomus in boreal and subarctic lakes. J Plankton Res **38**(3): 468-477.

Hoffmann, J.M. 2017. Investigating trophic ecology and dietary niche overlap among morphs of Lake Trout
in Lake Superior, University of Waterloo.

Horns, W., Bronte, C.R., Busiahn, T.R., Ebener, M.P., Eshenroder, R.L., Gorenflo, T., Kmieck, N., Mattes,
W., Peck, J.W., Petzold, M., and Schreiner, D.R. 2003. Fish-community objectives for Lake Superior. Great
Lakes Fishery Commission Special Publication 03-01 Ann Arbor, Michigan.

Howell, K.L., Pond, D.W., Billett, D.S., and Tyler, P.A. 2003. Feeding ecology of deep-sea seastars
(Echinodermata: Asteroidea): a fatty-acid biomarker approach. Marine Ecology Progress Series **255**: 193-
206.

Hrabik, T.R., Jensen, O.P., Martell, S.J.D., Walters, C.J., and Kitchell, J.F. 2006. Diel vertical migration in the
Lake Superior pelagic community. I. Changes in vertical migration of coregonids in response to varying
perdation risk. Can J Fish Aquat Sci **63**: 2286-2295.

Iverson, S.J. 2009. Tracing aquatic food webs using fatty acids: from qualitative indicators to quantitative
determination. *In* Lipids in Aquatic Ecosystems. Springer. pp. 281-308.

Iverson, S.J., Bowen, W.D., Boness, D.J., and Oftedal, O.T. 1993. The Effect of Maternal Size and Milk
Energy Output on Pup Growth in Grey Seals (Halichoerus grypus). Physiol Zool **66**(1): 61-88.

Iverson, S.J., Field, C., Bowen, W.D., and Blanchard, W. 2004. Quantitative fatty acid signature analysis: a
new method of estimating predator diets. EcologicalMonographs **74**: 211–235.

Iverson, S.J., Frost, K.J., and Lowry, L.F. 1997. Fatty acid signatures reveal fine scale structure of foraging
distribution of harbor seals and their prey in Prince William Sound, Alaska. Marine Ecology Progress Series
151: 255-271.

Janhunen, M., Peuhkuri, N., and Piironen, J. 2009. Morphological variability among three geographically
distinct Arctic charr (Salvelinus alpinus L.) populations reared in a common hatchery environment. Ecol
Freshw Fish **18**(1): 106-116.

Johnson, L. 1975. Distribution of fish species in Great Bear Lake, Northwest Territories, with reference to
zooplankton, benthic invertebrates, and environmental conditions. J Fish Res Board Can **32**: 1989-2004.

Käkelä, R., Käkelä, A., Kahle, S., Becker, P.H., Kelly, A., and Furness, R.W. 2005. Fatty acid signatures in
plasma of captive herring gulls as indicators of demersal or pelagic fish diet. Marine Ecology Progress
Series **293**: 191-200.

Kattner, G., Hagen, W., Graeve, M., and Albers, C. 1998. Exceptional lipids and fatty acids in the pteropod
Clione limacina (Gastropoda) from both polar oceans. Marine Chemistry **61**(3): 219-228.

Kavanagh, A.S., Cronin, M.A., Walton, M., and Rogan, E. 2010. Diet of the harbour seal (*Phoca vitulina
vitulina*) in the west and south-west of Ireland. Journal of the Marine Biological Association of the United
Kingdom **90**(08): 1517-1527.

Khan, N.Y., and Qadri, S.U. 1970. Morphological differences in Lake Superior lake char. J Fish Res Board
Can **27**: 161-167.

King, J.R., and Jackson, D.A. 1999. Variable selection in large environmental data sets using principal
components analysis. Environmetrics **10**(1): 67-77.

Kriegel, H.-P., Kröger, P., Schubert, E., and Zimek, A. 2008. A General Framework for Increasing the
Robustness of PCA-Based Correlation Clustering Algorithms, Berlin, Heidelberg, 2008//, Springer Berlin
Heidelberg, pp. 418-435.

Logan, M.S., Iverson, S.J., Ruzzante, D.E., Walde, S.J., Macchi, P.J., Alonso, M.F., and Cussac, V.E. 2000.
Long term thet differences between morphs in trophically polymorphic Percichthys trucha (Pisces:
Percichthyidae) populations from the southern Andes. Biological Journal of the Linnean Society **69**(4): 599-
616.

Loseto, L., Stern, G., Connelly, T., Deibel, D., Gemmill, B., Prokopowicz, A., Fortier, L., and Ferguson, S.
2009. Summer diet of beluga whales inferred by fatty acid analysis of the eastern Beaufort Sea food web.
Journal of Experimental Marine Biology and Ecology **374**(1): 12-18.

MacDonald, D., Levy, D., Czarnecki, A., Low, G., and Richea, N. 2004. State of the aquatic knowledge of
Great Bear Lake watershed. Report to Indian and Northern Affairs Canada. Water Resources Division,
MacDonald Environmental Sciences, Nanaimo, British Columbia.

Mariash, H.L., Cazzanelli, M., Kainz, M.J., and Rautio, M. 2011. Food sources and lipid retention of
zooplankton in subarctic ponds. Freshwater Biology **56**(9): 1850-1862.

Mariash, H.L., Cusson, M., and Rautio, M. 2017. Fall Composition of Storage Lipids is Associated with the
Overwintering Strategy of Daphnia. Lipids **52**(1): 83-91.

McMeans, B.C., McCann, K.S., Humphries, M., Rooney, N., and Fisk, A.T. 2015. Food Web Structure in
Temporally-Forced Ecosystems. Trends in Ecology & Evolution **30**(11): 662-672.

McPhail, J. 1993. Ecology and evolution of sympatric sticklebacks (Gasterosteus): origin of the species
pairs. Canadian Journal of Zoology **71**(3): 515-523.

Miles, D.B., Sinervo, B., Hazard, L.C., Svensson, E.I., and Costa, D. 2007. Relating endocrinology, physiology
and behaviour using species with alternative mating strategies. Funct Ecol **21**(4): 653-665.

Moore, S.A., and Bronte, C.R. 2001. Delineation of sympatric morphotypes of lake trout in Lake Superior.
Transactions of the American Fisheries Society **130**(6): 1233-1240.

Morissette, O., Sirois, P., Lester, N.P., Wilson, C.C., and Bernatchez, L. 2018. Supplementation stocking of
Lake Trout (Salvelinus namaycush) in small boreal lakes: Ecotypes influence on growth and condition. PLoS
One **13**(7): e0200599.

Morrison, W.R., and Smith, L.M. 1964. Preparation of fatty acid methyl esters and dimethylacetals from
lipids with boron fluoride–methanol. Journal of lipid research **5**(4): 600-608.

Muir, A.M., Bronte, C.R., Zimmerman, M.S., Quinlan, H.R., Glase, J.D., and Krueger, C.C. 2014.
Ecomorphological diversity of Lake Charr *Salvelinus namaycush* at Isle Royale, Lake Superior. Trans Am
Fish Soc **143**(4): 972-987.

Muir, A.M., Hansen, M.J., Bronte, C.R., and Krueger, C.C. 2015. If Arctic charr *Salvelinus alpinus* is 'the
most diverse vertebrate', what is the lake charr *Salvelinus namaycush*? Fish and Fisheries:
DOI:10.1111/faf.12114.

Olsen, Y. 1999. Lipids and essential fatty acids in aquatic food webs: what can freshwater ecologists learn
from mariculture? *In* Lipids in freshwater ecosystems. *Edited by* M.T. Arts and B.C. Wainman. Springer-
Verlag, New York. pp. 161-202.

Olsen, Y., and Skjervold, H. 1995. Variation in content of Ω3 fatty acids in farmed Atlantic salmon, with
special emphasis on effects of non-dietary factors. Aquaculture International **3**(1): 22-35.

Ota, T., and Yamada, M. 1971. Lipids of Masu salmon Oncorhynchus masou. I. Variations of the lipid
content and the fatty acid composition of juvenile Masu salmon during the period of smolt-
transformation, and on the influence of light upon those variations. Hokkaido Univ Fac Fish Bull.

Piché, J., Iverson, S.J., Parrish, F.A., and Dollar, R. 2010. Characterization of forage fish and invertebrates
in the northwestern Hawaiian Islands using fatty acid signatures: species and ecological groups. Mar Ecol
Prog Ser **418**: 1-15.

Pomerleau, C., Lesage, V., Winkler, G., Rosenberg, B., and Ferguson, S.H. 2014. Contemporary Diet of
Bowhead Whales (Balaena mysticetus) from the Eastern Canadian Arctic Inferred from Fatty Acid
Biomarkers. Arctic **67**(1): 84-92.

Pryke, S.R., Astheimer Lee, B., Buttemer William, A., and Griffith Simon, C. 2007. Frequency-dependent
physiological trade-offs between competing colour morphs. Biology Letters **3**(5): 494-497.

Quérouil, S., Kiszka, J., Cordeiro, A.R., Cascão, I., Freitas, L., Dinis, A., Alves, F., Santos, R.S., and Bandarra,
N.M. 2013. Investigating stock structure and trophic relationships among island-associated dolphins in the
oceanic waters of the North Atlantic using fatty acid and stable isotope analyses. Mar Biol **160**(6): 1325-
1337.

R Core Team. 2017. R: A language and environment for statistical computing. Available from
http://www.R-project.org/.

Robinson, B.W., and Parsons, K.J. 2002. Changing times, spaces, and faces: tests and implications of
adaptive morphological plasticity in the fishes of northern postglacial lakes. Canadian Journal of Fisheries
and Aquatic Sciences **59**(11): 1819-1833.
Robinson, B.W., and Wilson, D.S. 1994. Character release and displacement in fishes: a neglected
literature. Am Nat **144**: 596-627.
Ronconi, R., Koopman, H.N., McKinstry, C., Wong, S., and Westgate, A.J. 2010. Inter-annual variability in
diet of non-breeding pelagic seabirds *Puffinus* spp. at migratory staging areas: evidence from stable
isotopes and fatty acids. Marine Ecology Progress Series **419**: 267-282.
Rundle, H.D., and Nosil, P. 2005. Ecological speciation. Ecol Lett **8**(3): 336-352.
Sasaki, S., Ota, T., and Takagi, T. 1989. Compositions of Fatty Acids in the Lipids of Chum Salmon during
Spawning Migration. NIPPON SUISAN GAKKAISHI **55**(12): 2191-2197.
Scharnweber, K., Strandberg, U., Marklund, M.H.K., and Eklöv, P. 2016. Combining resource use
assessment techniques reveals trade-offs in trophic specialization of polymorphic perch. Ecosphere **7**(8):
e01387.
Schluter, D., and McPhail, J.D. 1992. Ecological character displacement and speciation in sticklebacks.
American Naturalist **140**: 85-108.
Sheridan, M.A. 1989. Alterations in lipid metabolism accompanying smoltification and seawater
adaptation of salmonid fish. Aquacult **82**(1): 191-203.
Skulason, S., and Smith, T.B. 1995. Resource polymorphisms in vertebrates. Trends Ecol Evol **10**: 366-370.
Smith, T.B., and Skulason, S. 1996. Evoloutionary significance of resource polymorphims in fishes,
amphibians, and birds. Annual Reviews in Ecology and Systematics **27**: 111-133.
Svanbäck, R., Eklöv, P., Fransson, R., and Holmgren, K. 2007. Intraspecific competition drives multiple
species resource polymorphism in fish communities. Oikos **117**(1): 114-124.
Taipale, S.J., Vuorio, K., Aalto, S.L., Peltomaa, E., and Tiirola, M. 2019. Eutrophication reduces the
nutritional value of phytoplankton in boreal lakes. Environmental Research **179**: 108836.
Tucker, S., Bowen, W.D., and Iverson, S.J. 2008. Convergence of diet estimates derived from fatty acids
and stable isotopes within individual grey seals. Marine Ecology Progress Series **354**: 267.
Vinson, M.R., and Budy, P. 2010. Sources of variability and comparability between salmonid stomach
contents and isotopic analyses: study design lessons and recommendations. Canadian Journal of Fisheries
and Aquatic Sciences **68**(1): 137-151.
Webb, P.W. 1984. Body form, locomotion and foraging in aquatic vertebrates. Am Zool **24**: 107-120.
Witten, D.M., and Tibshirani, R. 2011. Penalized classification using Fisher's linear discriminant. Journal of
the Royal Statistical Society. Series B, Statistical methodology **73**(5): 753-772.
Yoder, J.B., Clancey, E., Des Roches, S., Eastman, J.M., Gentry, L., Godsoe, W., Hagey, T.J., Jochimsen, D.,
Oswald, B.P., Robertson, J., Sarver, B.A.J., Schenk, J.J., Spear, S.F., and Harmon, L.J. 2010. Ecological
opportunity and the origin of adaptive radiations. J Evol Biol **23**(8): 1581-1596.
Zimmerman, M.S., Krueger, C.C., and Eshenroder, R.L. 2006. Phenotypic diversity of lake trout in Great
Slave Lake: differences in morphology, buoyancy, and habitat depth. Trans Am Fish Soc **135**: 1056-1067.
Zimmerman, M.S., Krueger, C.C., and Eshenroder, R.L. 2007. Morphological and ecological differences
between shallow- and deep-water lake trout in Lake Mistassini, Quebec. J Great Lakes Res **33**: 156-169.
Zimmerman, M.S., Schmidt, S.N., Krueger, C.C., Vander Zanden, M.J., and Eshenroder, R.L. 2009.
Ontogenetic niche shifts and resource partitioning of lake trout morphotypes. Can J Fish Aquat Sci **66**:
1007-1018.

Table 1. The ten most discriminating non-dietary fatty acids biomarkers from SIMPER analyses
to determine which fatty acids were primarily responsible for observed differences. Results are
presented for each region, including percentage contribution to overall fatty acid dissimilarity
among lake trout morphs. Fatty acids are listed in order of elution; those highlighted in grey are
shared among the study three regions. The total percentage of the ten most discriminating fatty
acids are given for each region.

	Fatty acids	Great Bear Lake (63.8%)	Stannard Rock (70.9%)	Superior Shoal (69.9%)
Non-dietary Fatty Acids	12:0		3.6%	6.1%
	12:1	8.3%	8.9%	14.7%
	13:1			
	14:1n-9	5.4%	3.0%	4.5%
	14:1n-7	3.8%		3.3%
	14:1n-5			4.2%
	14:0 iso		6.4%	
	14:0 ante			
	15:0			
	15:1n-8	8.5 %	25.9%	18.8%
	15:1n-6	8.2 %	3.7%	4.2%
	15:0 iso	3.8%	3.6%	3.6%
	16:1n-11			
	16:1n-9			
	16:1n-5			
	16:0 iso	7.1%		
	7Methyl16:0	4.4%		
	17:1			
	17:0 iso			
	18:1n-5			
	20:0		4.1%	
	20:2n-9	10.2%	4.5%	5.2%
	22:0	4.1%	7.2%	5.3%
	24:1n-9			

Table 2. The ten most discriminating dietary fatty acids biomarkers from SIMPER analyses to determine which fatty acids were primarily responsible for observed differences. Results are presented for each region, including percentage contribution to overall fatty acid dissimilarity among lake trout morphs. Fatty acids are listed in order of elution; those highlighted in grey are shared among the study three regions. The total percentage of the ten most discriminating fatty acids are given for each regions.

	Fatty acids	Great Bear Lake (48.8%)	Stannard Rock (55.2%)	Superior Shoal (55.8%)
Dietary and Extended-Dietary Fatty Acids	14:0			
	16:0			
	16:1n-7			
	16:2n-6			
	16:2n-4			
	17:0			
	16:3n-4	4.0%	2.5 %	2.9%
	16:4n-3	3.0%	2.6%	
	16:4n-1		7.3%	6.3%
	18:0			
	18:1n-9		2.4%	
	18:1n-7			
	18:2n-6			
	18:2n-4			
	18:3n-6			
	18:3n-4			
	18:3n-3			
	18:3n-1	3.3%		3.5%
	18:4n-3			
	18:4n-1	3.5%	10.5%	11.1%
	20:1n-11	4.0%		
	20:1n-9			
	20:1n-7	3.6%		
	20:2n-6			
	20:3n-6			
	20:4n-6			
	20:3n-3			
	20:4n-3			
	20:5n-3			
	22:1n-11	8.6%	11.9%	12.1 %
	22:1n-9			
	22:1n-7		2.7%	4.6%
	22:2n-6	6.3%		
	21:5n-3	7.5%		4.7 %
	22:5n-6		7.2%	3.0%
	22:4n-3	5.0%	2.6%	4.2%
	22:5n-3			
	22:6n-3		5.5%	3.4%

List of Figures:

Fig. 1. Location of (1) Great Bear Lake (Canada) and (2) Lake Superior (Canada-USA). Two

sampling sites were defined in Lake Superior to account for spatial variation: Superior Shoal

and Stannard Rock (QGIS 3.0, Canada Basemap; Hoffmann 2017).

Fig 2. A) The four morphotypes of Lake Trout in Lake Superior, the lean, humper, siscowet,

and redfin (defined in Muir et al. 2014). B) The four shallow-water morphotypes of Lake

Trout from Great Bear Lake, Morphs1-4 (defined in Chavarie et al. 2013).

Fig. 3. PCA of non-dietary fatty acids biomarkers (top panels), dietary fatty acids biomarkers

(middle panels), and all fatty acids (bottom panels) for individual Lake Trout collected from

Great Bear Lake, Stannard Rock (Lake Superior), and Superior Shoal (Lake Superior). Vectors

of individual fatty acids important to the positioning of lake trout are represented to the right

of each PCA. Angles and lengths of vectors represent the direction and strength of

relationships, respectively, between variables and the principal components.

Fig. 4. Results of linear discriminant function analyses of non-dietary fatty acids biomarkers

(top panel) and dietary fatty acids biomarkers (bottom panel) for Lake Trout collected from

Great Bear Lake, Stannard Rock (Lake Superior), and Superior Shoal (Lake Superior). The

95% ellipse of each morph is also provided.

750   Fig. 1.

770 A)

771 Fig 2.

Fig. 3.

Fig. 4.

Appendix:

Table A1. List of 38 of 41 fatty acids shared by the two datasets, considered as either "dietary"
fatty acids or "extended-dietary" fatty acids biomarkers and used in this study (see Iverson *et al*.,
2004), and the dietary component they are associated with, based on literature: Sargent *et al*.,
1995; Brett & Müller-Navarra, 1997; Kattner *et al*., 1998; Virtue *et al*., 2000; Budge et al., 2002;
Dalsgaard *et al*., 2003; Iverson *et al*., 2004; Käkelä *et al*., 2005; Alfaro *et al*., 2006; Tucker *et al*.,
2008; Ahlgren *et al*., 2009; Gladyshev *et al*., 2009; Loseto *et al*., 2009; Stowasser *et al*., 2009;
Piché *et al*., 2010; Mariash *et al*., 2011. The fatty acids highlighted are the one discarded because
they were not quantified for both lakes.

Dietary Fatty acids	Dietary component
14:0	Pelagic (Zooplankton) + diatom
16:0	Cannibalism or/and carnivorous
16:1n7	Benthic (bacterial synthesis + diatoms) Cannibalism or/and carnivorous
16:2n6	
16:2n4	Diatom
17:0	
16:3n-4	
16:4n3	
16:4n1	Diatom
18:0	Cannibalism or/and carnivorous
18:1n9	Pelagic (Zooplankton)
18:1n7	Benthic (bacterial synthesis + algal)
18:2n6	Terrestrial
18:2n4	
18:3n6	
18:3n4	
18:3n3	Terrestrial
18:3n1	
18:4n3	Pelagic (Zooplankton)
18:4n1	
20:0	*not used
20:1n11	Copepod (Iverson, 2009)
20:1n9	Pelagic (Calanoid copepods diet based)
20:1n7	Pelagic (Zooplankton)
20:2n9	*not used
20:2n6	
20:3n6	
20:4n6	Benthic (diatom)
20:3n3	
20:4n3	

20:5n3	Cannibalism or/and carnivorous
22:1n11	Copepod
22:1n9	Pelagic (Zooplankton)
22:1n7	
22:2n6	
21:5n3	
22:4n6	*not used
22:5n6	Pelagic (Calanoid copepods diet based)
22:4n3	
22:5n3	
22:6n3	Benthic (Pennate diatoms + dinoflagellates + bivalves) Cannibalism

797   Table A2. List of 24 fatty acids considered as "non-dietary" biomarkers in this study.

Non-Dietary Fatty acids
12:0
12:1
13:1
14:1n9
14:1n7
14:1n5
14:0 iso
14:0 ante
15:0
15:1n8
15:1n6
15:0 iso
16:1n11
16:1n9
16:1n5
16:0 iso
7Methyl 16:0
17:1
17:0 iso
18:1n5
20:0
20:2n9
22:0
24:1n9

Fig. A1. PCA of all fatty acids (top panels), dietary fatty acids biomarkers (middle panels), and
non-dietary fatty acids biomarkers (bottom panels) of each morph of Lake Trout from Great Bear
Lake, Stannard Rock (Lake Superior), and Superior Shoal (Lake Superior), including the 12 outlier
individuals. Vectors of individual fatty acids important to the positioning of lake trout are

804   represented to the right of each PCA. Angles and lengths of vectors represent the direction and the

805   strength, respectively, of relationships between variables and the principal components.